



THE UNIVERSITY
OF ILLINOIS
LIBRARY

581.942
L92a2
v. 1

CPLA

Return this book on or before the
Latest Date stamped below.

University of Illinois Library

LL1-H3

8 17/18 1/2
D. A. -

ARBORETUM ET FRUTICETUM BRITANNICUM;

OR,

THE TREES AND SHRUBS OF BRITAIN,

Native and Foreign, Hardy and Half-Hardy,

PICTORIALLY AND BOTANICALLY DELINEATED,
AND SCIENTIFICALLY AND POPULARLY DESCRIBED;

WITH

THEIR PROPAGATION, CULTURE, MANAGEMENT,

AND USES IN THE ARTS, IN USEFUL AND ORNAMENTAL PLANTATIONS, AND IN

LANDSCAPE-GARDENING;

PRECEDED BY A HISTORICAL AND GEOGRAPHICAL OUTLINE
OF THE TREES AND SHRUBS OF TEMPERATE CLIMATES
THROUGHOUT THE WORLD.

BY J. C. LOUDON, F.L. & H.S. &c.

AUTHOR OF THE ENCYCLOPEDIAS OF GARDENING AND OF AGRICULTURE.

IN EIGHT VOLUMES:

FOUR OF LETTERPRESS, ILLUSTRATED BY ABOVE 2500 ENGRAVINGS;
AND FOUR OF OCTAVO AND QUARTO PLATES.

VOL. I.

HISTORY, GEOGRAPHY, AND SCIENCE; AND DESCRIPTIONS, FROM
RANUNCULACEÆ TO STAPHYLEACEÆ, P. 494., INCLUSIVE.

SECOND EDITION.

LONDON:

HENRY G. BOHN, YORK STREET, COVENT GARDEN.

1854.

151
L92a2
v.1

TO HIS GRACE,

HUGH, DUKE OF NORTHUMBERLAND, K.G.

Sc. Sc.

MY LORD DUKE,

IN dedicating to Your Grace the accompanying Volumes, I am anxious to show how fully I appreciate the encouragement which your ancestors and yourself have always given to gardening pursuits, and more especially to the introduction and cultivation of foreign trees and shrubs. How much the British Arboretum is indebted to the noble family of Northumberland, for the introduction of trees and shrubs from America during the last century, is evinced by the *Hortus Kewensis*, *Miller's Dictionary*, and other works which record the names of the first introducers of foreign plants; and how various and magnificent are the specimens of foreign trees which exist in the grounds at Syon, the numerous portraits of them which are given in the Volumes now submitted to the public bear ample testimony.

For the kindness which Your Grace has evinced, in having had these portraits made expressly for my work, I am desirous that this dedication should be considered as a public memorial of my lasting gratitude.

I have the honour to subscribe myself

Your Grace's

Very obedient humble Servant,

THE AUTHOR.

Bayswater, May 20. 1838.

P R E F A C E.

THE main object which induced the author to undertake this Work was, the hope of diffusing more generally, among gentlemen of landed property, a taste for introducing a greater variety of trees and shrubs in their plantations and pleasure-grounds. He had observed, for a number of years, that, though many new and beautiful trees and shrubs were annually introduced from foreign countries into our botanic gardens and nurseries, yet the spread of these plants in the grounds of country residences was comparatively slow; and that not only the new sorts were neglected, but many of the fine old species and varieties, which had been in British nurseries for upwards of a century, were forgotten by planters, and had ceased to be propagated by commercial gardeners. In short, it appeared to the author, that the general taste of the country for trees and shrubs bore no just proportion to the taste which prevailed in it for fruits, culinary productions, and flowers. It also appeared to him, that, while the numerous horticultural societies now established in the British Islands had powerfully promoted the general taste for horticultural and floricultural productions, they had rather neglected arboriculture and landscape-gardening.

Viewing trees and shrubs as, next to buildings, the most important ornaments which can be introduced into a country; and considering them, in this respect, greatly superior to herbaceous plants, from the little care that trees and shrubs require when once properly planted, and their magnitude, and permanent influence when grown up, on the general scenery of the country; the author felt desirous of pointing out the great importance of their more general distribution and culture. In order to impress this on the minds of proprietors and their families, and especially on the rising generation among them, he thought it best to adopt, as the main feature of his plan, the description and portraiture of such species and varieties of trees and shrubs as are actually in cultivation in the country, and as grow vigorously in it; referring to gardens or grounds within a limited distance of London, where these species or varieties may be seen in a living state, and to nurseries where they are propagated for sale, and stating the price for which they might be purchased in England, in France and Germany, and in North America. He has thought it advisable to give, not only botanical specimens, but portraits of the greater number of species of trees; in order, by a palpable representation of their forms and magnitudes, to make a stronger impression on the mind of the reader. These pictorial illustrations are of two kinds: first, portraits of trees of ten or twelve years' growth, taken from specimens growing in 1834, 1835, or 1836, within ten miles of London, and all drawn to the same scale of 1 in. to 4 ft.; and, secondly, of full-grown trees, also all drawn to one scale, viz. 1 in. to 12 ft., and for the most part growing within the same distance of London.

The use of the first class of portraits is, to give a palpable idea of the general magnitude, form, and character, which different species and varieties assume when growing in the same soil and climate, even in so short a period as ten or twelve years after planting. A slight comparative view of these portraits shows that the growth of some trees is much more rapid than that of others; and that while the species of some genera when young are comparatively monotonous in their general form, those of the species of other genera of the same age have marked and characteristic features. Hence the valuable assistance afforded by such portraits in the choice of trees for the purposes of landscape-gardening. To solve the problem of ascertaining the species capable of producing any desired effect of wood, or of trees, in a given locality, and in a given time, it is only necessary to turn over the portraits which are contained in the last four volumes of this Work, and to select those species, the portraits of which exhibit trees of such forms and magnitudes as will produce the effect desired.

The second class of portraits represents full-grown trees of the same species as those of which portraits in their young state are given; and these, for the greater part, are drawn from trees within ten miles of London, and all, with one or two exceptions, are to one scale. The object in giving these portraits of full-grown trees is, to show the magnitude and character which particular species attain when they arrive at maturity, and to be a guide to the planter, not only in many particulars having reference to pictorial effect, but also with respect to the uses of trees as productive of timber, shelter, and shade.

These portraits of trees in their young and mature state, which are contained in the last four volumes, together with the engraved botanical specimens, and the scientific and popular descriptions contained in the four volumes of letter-press, are calculated, as the author thinks, to create that interest in trees and shrubs in those who have not previously paid much attention to the subject, to produce which, as already observed, has been his grand object in undertaking this Work.

In addition to the trees and shrubs which have been ten or twelve years in the country, and which are purchasable in British nurseries, those of more recent introduction, which are comparatively scarce, are also described or noticed; as are some which were introduced at former periods, and have been lost, and others which are known to botanists, but which have not yet been introduced.

Besides notices of new and little known hardy trees and shrubs, or of such as are described by botanists but not yet introduced, some attention has been paid throughout the Work to trees and shrubs which, though they will stand through the winter in mild seasons, without protection in favourable situations, or with protection in situations and seasons less favourable, in the climate of London, yet cannot be recommended for general purposes, and are therefore generally described by gardeners as only half-hardy. These half-hardy species have been noticed, because there is perhaps no scene in a British garden more interesting, than one in which the plants of warm

countries, usually seen in green-houses or conservatories, appear in a flourishing state in the open air; and also because the culture and management of such plants call forth a higher degree of scientific knowledge and attention on the part of the gardener, and therefore contribute to his improvement and consequent usefulness. Half-hardy trees and shrubs of the more tender kinds are generally cultivated against what is in this Work called a conservative wall; that is, a wall which may be flued or not according to circumstances, but which admits of putting up a temporary projecting roof, or some other means of protection, during the winter season. Such walls are at present not very generally in use for ornamental exotics; but so great are the interest and beauty which they are calculated to display when properly designed, planted, and managed, that in a few years they will probably be as general as fruit walls; and, as ornaments to a country residence, connected with the flower-garden and shrubbery, will be considered as ranking next to the conservatory and the green-house.

Having given this notice of the objects which have led to the undertaking of this Work, and a general outline of the plan pursued in it, the reader is referred to the Introduction, and to the Explanatory References which precede the Table of Contents, for further details.

The engraving and printing of this Work, for which collections had been making for several years (see the *Gardener's Magazine*, vol. vi., for 1830, p. 582. and p. 718.), was begun in August, 1834; and it has been published in Numbers, the first of which appeared in January, 1835, and the 63d and last on the 1st of July, 1838. It was originally intended to include in it a generalisation of the whole subject of the trees and shrubs of temperate climates; but, finding the Work had increased to more than double the extent originally contemplated, it has been thought advisable to publish the generalisation alluded to separately; and it will accordingly appear at some future time, in one volume, under the title of an *Encyclopædia of Arboriculture*.

It now remains for the author to acknowledge his obligations to the numerous persons who have given him information for this Work. He refers to the List of Contributors, p. xv., and to every individual there named he begs to return his most sincere thanks. He thanks, in a more particular manner, the Council of the Horticultural Society of London, for having granted him permission to take portraits of the trees in the Society's Garden, and to examine and make drawings from the botanical specimens sent home by Douglas and others, in their herbarium; as well as for the information communicated by their head gardener, Mr. Munro, and the foreman of their arboretum, Mr. Gordon. The author is deeply indebted to Messrs. Loddiges, with reference to their arboretum at Hackney, and for the kind and liberal manner in which they have at all times, during the last ten years, since he began to prepare for this Work, not only allowed him to send artists to make drawings, and supplied him with specimens, but permitted him to select these himself; and, in short, to use their unrivalled collection of hardy trees and shrubs as if it had been his own. To the Linnean Society, and their librarian, Professor Don, the author is much indebted for the loan of books, and for permission to examine

the specimens of trees and shrubs in the Linnean herbarium; as he is to A. B. Lambert, Esq., V.P.L.S., &c., for the use of his magnificent library and unique herbarium, and for the living specimens of the Coniferæ from Boyton; and to W. T. Aiton, Esq., F.L.S., H.S., &c., Her Majesty's Garden-Director at Kew, for specimens of the trees and shrubs in the arboretum there, for the loan of manuscripts, and for other valuable information and assistance, communicated either directly by himself, or through that scientific and assiduous botanist and cultivator, Mr. Smith, foreman of the Kew Botanic Garden.

To His Grace the Duke of Northumberland the author is indebted, not only for access for himself and artists to examine and take portraits of the splendid exotic trees at Syon, but for drawings of upwards of a hundred of the largest and rarest of those trees, made for this Work by G. R. Lewis, Esq., at His Grace's expense, and also for engravings from several of the largest of these drawings. To Mrs. Lawrence of Studley Royal he is obliged for the portraits of many of the noble trees in the park at Studley, that lady having employed a London artist, H. W. Jukes, Esq., for several months for that purpose; and to the Rev. J. Charnock, for directing the measurements, and supplying interesting information respecting those trees. To the Countess of Bridgewater he is indebted for portraits of the gigantic beeches at Ashridge: and to Lady Grenville, for permission to take drawings, and for numerous specimens of the *Abietinæ* in the magnificent pinctum at Dropmore; and to Her Ladyship's intelligent and most industrious gardener, Mr. Frost, for supplying valuable information respecting their culture and management, and for lending every assistance in his power. Portraits of trees, and many specimens, have also been sent by various other persons in different parts of Great Britain and Ireland, whose names, lest any individual should be by chance omitted, it is not endeavoured here to enumerate.

The author is also indebted, for portraits, specimens, and information, to various amateurs and gardeners on the Continent, as well as to correspondents in North America and Australia; and more especially to M. Alphonse De Candolle of Geneva; to M. André Michaux, M. Vilmorin, M. Loiseleur Deslongchamps, and M. Soulange-Bodin, of Paris; to Baron Jacquin, and M. Charles Rauch, of Vienna; to Professor Reinwardt, Leyden; Professor Kops, Utrecht; M. Otto, Berlin; Messrs. Booth, Hamburg; Professor Schouw and M. Petersen, Copenhagen; Bishop Agardh, Carlstadt, Sweden; Dr. Fischer, Petersburg; M. Fintelman, Moscow; M. Descemet, Odessa; Sr. G. Manetti, Monza, near Milan; the Honourable Keppel Craven, Naples; Dr. Mease, Philadelphia; Col. Carr, of Bartram's Botanic Garden; and John Thompson, Esq., Surveyor General's office, Sydney.

In the literary department the author received the assistance of the Rev. M. J. Berkeley, M.A., F.L.S., for the lists of Fungi inhabiting different species of trees; of J. O. Westwood, Esq., F.L.S., Secretary to the Entomological Society, for descriptions and drawings of the Insects infesting different species; of Wm. Spence, Esq., F.L.S., and of M. Audouin, Member of the Institute of France, and Professor of Entomology, for information respecting the species of *Scôlytus* injurious to the elm; of Mr. John Denson, A.L.S., in drawing up

the characters of the Orders and Genera, and generally for what may be considered the botanical department of this Work, from its commencement to the end of the genus *Sâlix*. He is, also, indebted to W. Withers, Esq., of Holt, Norfolk, for the loan of his unpublished work on the *Robinia Pseud-Acacia*; to W. Borrer, Esq., F.L.S., for the arrangement into groups of the numerous species of the difficult genus *Sâlix*, and for looking over the proof sheets of the article on that genus; to Professor Alphonse De Candolle, for looking over the proof sheets of the article on *Salisbùria*, as well as for examining the proofs of the history of trees in France and Switzerland; to M. Vilmorin, for looking over the proof sheets of the history of trees and shrubs in France; to the late M. Fischer, of the Botanic Garden, Göttingen, and to M. Otto, of Berlin, for looking over the proof sheets of the history of trees in Germany; to Professor Don, for arranging and looking over the whole of the generic characters, and some of the proof sheets, of the *Coniferæ*; and to Mr. Gorrie, for examining the proof sheets of the article on the Larch.

To His Grace the Duke of Bedford the author is indebted for much valuable information on the subject of trees, and more especially respecting the Oak, the Scotch Pine, and the Cedar of Lebanon, communicated either directly by His Grace, or through his forester at Woburn Abbey, Mr. Ireland; as well as for permission (of which he has availed himself) to make whatever use he chose of that magnificent work the *Salictum Woburnense*. To the Duke of Richmond, the Earl of Aberdeen, Macpherson Grant, Esq., and Mr. Grigor, of Forres, he is indebted for important communications respecting the Scotch Pine in Scotland; to the Duke of Portland for various researches respecting the Oak in Nottinghamshire; and to the Earl of Wicklow, Earl Roden, Lord Viscount Ferrard, Sir Robert Bateson, Mr. Mackay, and Mr. Niven, for information respecting the trees and shrubs of Ireland.

The author, in conclusion, has only to request that the readers of this Work will send to him whatever corrections, additions, or suggestions may occur to them on its perusal, or at any future period. It is his intention to publish whatever additional matter he may receive from correspondents, or procure himself, in an Annual Report in the *Gardener's Magazine*; and to include in this report notices of all the trees and shrubs which may, during the year, have been introduced from foreign countries, or originated in this country from intentional hybridisation or accident. This report will also include such improvements as may from time to time be made in arboricultural nomenclature, such as the re-arrangement of the species of a genus, &c.; and, in short, every thing that may be found requisite to keep up the information contained in the *Arboretum et Fruticetum Britannicum*, with the actual progress of knowledge on the subject. The essence of these annual reports will be collected from time to time, and published in pamphlets, as supplements to the *Arboretum*; so that the possessor of this first edition may have it in his power, at any future time, to bring the Work down to the latest date, without being under the necessity of purchasing a new edition.

J. C. L.

Bayswater, May 20. 1838.

DIRECTIONS TO THE BINDER.

THE Work is to be done up in Eight Volumes: the first Four of Letter-press, and the last Four of Plates.

Vol. I. is to contain the Dedication, Preface, Contents, &c., of the eight volumes, and the text as far as p. 494. inclusive. A Table of the Contents of Vol. I. is to be placed immediately before the Introduction; and the first part of this table being systematic, and the second part alphabetical, an index at the end of the volume is unnecessary. The Signatures at the beginning of the volume run thus:—A, [A], a, a to d d, * B, c, &c., in the regular series.

Vol. II. commences with p. 495., and ends with p. 1256. There is a Title, with a Table of Contents on the same plan as in Vol. I.

Vol. III. commences with 1257., and ends with p. 2030.; having a Table of Contents, &c., as in Vol. I.

Vol. IV. commences with p. 2031., and contains the remaining part of the text; with a Table of Contents, &c., as in Vol. I., at the beginning, and three Alphabetical Indexes at the end.

Vol. V. commences with *Magnòlia grandiflòra*, and ends with *Acàcia dealbàta*. There is a Title, with a Table of Contents arranged on the same plan as that given in the preceding volumes.

Vol. VI. commences with *Amýgdalus commùnis*, and ends with *O'rnus europæ'a*, full-grown tree. Title, Contents, &c., as in preceding volumes.

Vol. VII. commences with *Catálpa syringæfòlia*, and ends with *Quércus álba*. Title, Contents, &c., as before.

Vol. VIII. commences with *Quércus macrocárpa*, and ends with *Juníperus excélsa*; and has Title, Contents, &c., as before. At the end of this volume there is an Alphabetical Index to the Four Volumes of Plates.

* * * A List of the Plates, in the order in which they are to be bound up, is given in the Table of Contents, p. cliv. to clxi. in Vol. I.

The *Cancels* given in order to introduce corrections, &c., are the following:—

TEXT.

Vol. I., pages 1 to 15. 21, 22. { 27, 28. { 29, 30. 73, 74. 157, 158. { 159, 160. { 173, 174. { 175, 176. { 177, 178. 229, 230. 493, 494.		Vol. II., pages 495, 496. 921, 922. 1229, 1230. 1255, 1256. Vol. III., Titlepage. pages 1257 to 1262. { 1273, 1274. { 1275, 1276. 1295, 1296. 1987, 1988. 2029, 2030. Vol. IV., pages 2031, 2032.
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

PLATES.

Vol. VII. *Pterocàrya caucásica*, to be substituted for *Jùglans fraxinifòlia*.

Certain superfluous Plates given with some copies, and which are to be cancelled, are enumerated at the end of the Supplement, Vol. IV. p. 2608.

ARRANGEMENT OF THE CONTENTS.

Explanatory References	-	-	-	-	Page xii
List of Contributors	-	-	-	-	xv
Epitome of Contents	-	-	-	-	[xxiii]
Contents in full	-	-	-	-	*xiii
List of Portraits of Trees forming the Plates in the last Four Volumes	-	-	-	-	cliv
List of Portraits of Trees given along with the Text in the first Four Volumes	-	-	-	-	clxii
List of Landscapes given along with the Text, showing the Effect of particular Species of Trees in Landscape Scenery	-	-	-	-	clxiv
List of Engravings of Botanical Specimens given along with the Text	-	-	-	-	clxv
List of Engravings of Fungi, Lichens, &c., given along with the Text	-	-	-	-	clxxxii
List of Engravings of Insects given along with the Text	-	-	-	-	clxxxiii
List of Engravings of Diagrams and miscellaneous Subjects given along with the Text	-	-	-	-	clxxxiii
List of Authorities for scientific Names	-	-	-	-	clxxxiv
List of Books quoted or referred to	-	-	-	-	exc

Immediately after the Titlepage of each Volume (except Vol. I.), a Table of the Contents is given, first systematically arranged, and next alphabetically.

A Table of the Contents of Vol. I. will be found from p. ccxxvii. to p. ccxxx.

 ARRANGEMENT OF THE ALPHABETICAL INDEXES.

- Alphabetical Index of Genera is given at the end of Vol. IV. p. 2655.
- Alphabetical Index of miscellaneous Subjects, at the end of Vol. IV. p. 2667.
- Alphabetical Index of Persons and Places, at the end of Vol. IV. p. 2672.
- Alphabetical Index of the Portraits of Trees given in the last Four Volumes, at the end of Vol. VIII.

EXPLANATORY REFERENCES.

THE greater part of the letterpress of this Work consists of the description, history, geography, uses, propagation, culture, &c., of the species and varieties of the trees and shrubs cultivated in the British Islands; and this is always printed in type corresponding in size with that used in this sentence.

The short descriptive notices of species and varieties which it would be desirable to introduce, of such as have been introduced and lost, of such as have not been seen by the author, even though mentioned in British catalogues, and of such as are half-hardy in the climate of London, or supposed to be so, are uniformly printed in a smaller type.

The statistics, or accounts of the dimensions of trees, which we have received from different parts of the British Islands or the Continent, are also in small type, in order to save room.

The portraits of trees which form the last four volumes are sometimes, referred to as in the second volume, and sometimes as in the last volume. These references were made under the idea of binding up all the letterpress in one volume, and all the plates in another, which, now that the Work is finished, is found to be impracticable. The readiest way of finding the portrait of any particular species of tree is, to refer to the Alphabetical Index of Portraits of Trees, given at the end of the eighth volume. The readiest way of finding the description and history of any particular species or variety, and of ascertaining whether or not a figure is given of it, is by referring to the general Alphabetical Index, at the end of the fourth volume.

In various parts of the Work reference will be found to Part IV., and to the *Encyclopædia of Arboriculture*. These two references refer to one and the same work, viz. the *Encyclopædia of Arboriculture*, which it was originally intended to include in the *Arboretum Britannicum*, but which, for the reasons stated in the Preface, will now be published separately in one volume; and in which the subject of trees and their culture will be generalised, and their mode of treatment given *en masse*, whether as seedlings in the nursery, as useful and ornamental plantations, as yielding timber and other useful products, or as ornaments in the lawn and shrubbery.

The engravings of the botanical specimens, whether printed along with the text, or along with the portraits of the trees in the four last volumes, are invariably to one and the same scale of 2 in. to 1 ft. Where any portion of the plant is given of the natural size, it is distinguished by a cross, thus, +. Where dissections are given, *m.* signifies male, *f.* female, and *mag.* magnified.

The portraits of the entire trees and shrubs given along with the text are to different scales, which are always indicated in the descriptions: they are chiefly 1 in. to 12 ft., 1 in. to 24 ft., and 1 in. to 50 ft.

The portraits of the trees in the last four volumes are nearly all from individuals that were growing within ten miles of London, in the years 1834, 1835, 1836, or 1837. These portraits are chiefly of young trees, of ten or twelve years' growth, and are drawn to a scale of 1 in. to 4 ft.: but there are also portraits of full-grown trees, of all the principal kinds of which full-grown specimens are to be found within ten miles of London (taken chiefly from Syon), and these are drawn to a scale of 1 in. to 12 ft.

Of some species of trees good full-grown examples could not be found within ten miles of London; and of these portraits have been taken from trees growing in different parts of Great Britain (particularly from Studley Park and Dropmore) and Ireland, and, in one or two instances, from trees on the Continent.

The botanical specimens of the young trees exhibit a branch in flower, the winter's wood when the tree is deciduous, and a branch with ripe fruit, and with

the autumnal leaves; besides dissections of the flowers and fruit. These were all drawn on purpose for this Work, and, with a few exceptions, by J. D. C. Sowerby, Esq., F.L.S., &c. See this subject further explained in p. 223.

The specimens of foliage placed at the bottom of the plates of the full-grown trees may be called "artist's foliage," rather than "botanical specimens." They are, for the most part, drawn by the artist who took the portrait of the tree (generally no botanist, and who only knew the artistical differences in the aspect of trees), in order to get a more correct idea of what is called the "touch."

The portraits of all the trees, both young and full grown, were taken chiefly during the months of August, September, and October; but a number were also taken during the winter season, in order to show the skeleton tree without its foliage; several species being as readily known, even to a general observer, when they are naked, as when they are clothed with leaves.

All the engravings of trees and shrubs given in this Work, whether along with the text, or in the last four volumes, have been drawn from nature, on purpose for it, by competent artists, whose names, as well as the names of the places, where the trees are now growing, or grew when their portraits were taken, are given in the List of Trees in the Table of Contents, p. cliv.; and the greater number of the original drawings may still be seen in the possession of the author.

In the descriptive part of this Work, under the titles of the chapters, and sometimes under those of the sections, are given signs, intended to show at a glance the general habit of the trees or shrubs described in that chapter or section. These signs represent large, small, and middle-sized plants, and are as follows; the first sign in each row indicating a deciduous tree or shrub, the next an evergreen, and so on alternately:—



1. Round-headed trees; such as the oak, ash, elm, beech, &c.



2. & 3. Spiry-topped trees; such as the spruce fir, larch, pine, deciduous cypress, &c.



4. Fastigate trees; such as the Lombardy poplar, cypress, &c.



5. Drooping trees; such as the weeping willow, weeping elm, &c.



6. Shrubs.




7. Under-shrubs



8. Twining shrubs; such as the honeysuckle, menispermum, aristolochia, &c.





9. Climbing shrubs; such as the clematis, ampelopsis, vine, &c.

10. Trailing shrubs, the branches of which lie prostrate on the ground, but do not root into it; such as many species of willow, *Cistus*, &c.

11. Creeping shrubs, or such as send up shoots from their creeping roots; as many species of *Spiræa*, &c.

The signs put before each individual species and variety which is described as enduring the open air in the climate of London, and in cultivation in British gardens, are the same as those used in the *Gardener's Magazine*, and in the *Hortus Britannicus*, viz.:—

☞ Deciduous tree.

♁ Evergreen tree.

☞ Deciduous shrub.

♁ Evergreen shrub.

☞ Deciduous under-shrub.

♁ Evergreen under-shrub.

☞ Deciduous twiner.

♁ Evergreen twiner.

☞ Deciduous climber.

♁ Evergreen climber.

☞ Deciduous trailer.

♁ Evergreen trailer.

☞ Deciduous creeper.

♁ Evergreen creeper.

The sign \lrcorner (or \lrcorner , indicating a greater degree of tenderness), added to any of the above signs in the Table of Contents, indicates that the tree or shrub, in the climate of London, requires protection during winter, but is considered likely to live against a conservative wall. Throughout the Work, wherever the dimensions of any tree or shrub are given, and the year when these dimensions were taken is not stated, the autumn of the year 1834, when the Work was commenced, is to be understood.

All the botanic names throughout the Work are accented, and have their origin indicated, as in the *Hortus Britannicus* and the *Gardener's Magazine*. The vowels which are sounded short are marked with an acute accent, thus (´), as *A´ceras*; and those which are sounded long are marked with a grave accent, thus (˘), as *A˘brus*. The origin of each name is indicated thus: where the name has been applied to a plant by the ancients, the first letter is in Italic, as *Pinus*; where it is commemorative of some individual, the letters additional to the name are in Italic, as *Banksia*, *Lambertiana*, *Douglasii*; and where an aboriginal name has been adopted, or where the name is of uncertain derivation, the whole word is in Italic as, *Ailantus*, *Caragana*, &c. When the name would otherwise be in Italic, as in the case of synonymes, headings to paragraphs, &c., these distinctions are of course reversed, as *Pinus*, *Banksia*, *Ailantus*. All the other scientific names, generic or specific, are composed from the Greek or Latin, with the exception of a very few, which are taken from places: as *Araucaria*, from the country of the Araucanians; *Quercus gramuntia*, from the estate of Grammont; *Acer monspeliensis*, &c. Further details respecting the particulars entered into in classing, describing, and recording the trees and shrubs included in this Work, will be found in the Introduction, p. 1. to p. 14., and in Part II. Chap. IV. p. 222. to p. 230., which we recommend the reader to peruse with attention.

In the course of the Work, a few exceptions will be found to what is stated in these explanatory pages as general; but they are so very few as to be hardly worth notice; and the reasons for them will either be found given where they occur, or they are considered to be sufficiently obvious.

LIST OF CONTRIBUTORS.

IN 1834, before the *Arboretum* was commenced, about three thousand printed lists of trees and shrubs, agreeable to the form shown in Appendix No. I., were put in circulation; and answers to these, or other information relating to trees and shrubs for this *Arboretum*, were received from the following persons:—

A.

- Aberdeen, George Earl of, F.R.S.,
L.S., H.S., &c.
Ackland, Sir T. D., Bart., M.P., F.
H.S., &c.
Agardh, D. C. A., Bishop of Carlstadt,
F.L.S., &c.
Agardh, jun., Professor of Botany,
Lund.
Aglionby, H. A., M.P., F.H.S., &c.
Aiken, H.
Aikin, A., M.D., F.G.S., &c.
Ailsa, A. Marquess of, F.R.S., H.
S., &c.
Airlie and Linrathen, D. O. Earl of.
Aiton, W. T., F.L.S., H.S., &c.
Albett, Josh.
Alexander, J.
Alsop, J.
Amherst, W. P. A. Earl, P.C.
Anderson, A.
Anderson, G., F.H.S.
Anderson, J., F.H.S., &c.
Anderson, W., F.L.S., H.S.
Andrews, M., F.H.S., &c.
Argyll, G. W. Duke of.
Arundel, H. B. Lord.
Ashbury, W.
Ashby, W. Ashby.
Atkins, J.
Atty, Geo.
Audibert, M. Urban, C.M.H.S., To-
nelle, near Tarascon.
- B.
- Babington, C. C., M.A., F.L.S., &c.
Backhouse, J., C.M.H.S., Sydney.
Backhouse, Thos., C.M.H.S.
Bagnold, Captain.
Bagot, W. Lord, F.L.S., H.S., &c.
Bailly de Merlieux, M.
Baker, A. J. Creswell, F.H.S.
- Baker, W. R., F.H.S.
Balfour, W., C.M.H.S.
Banks, W. J., M.P.
Barnes, Thomas.
Barnet, James.
Barron, W.
Bateman, J., F.L.S., H.S., &c.
Bateson, Sir Robert, M.P.
Bath, Marquess of.
Bauer, Francis, F.R.S., L.S.
Baumann, N., Bollwyller.
Baxter, W., A.L.S., F.H.S., &c.
Baxter, W. H.
Beaton, D.
Bedford, John Duke of, F.L.S., A.
S., H.S., &c.
Belfield, F., jun.
Bell, R.
Bella, M., Grignon, near Versailles.
Bennett, S., C.M.H.S.
Bentinck, Lieut.-Col. H.
Bentley, A. C.
Bentley, M. C.
Berkeley, Rev. M. J., M.A., F.L.S.
Bethell, R., M.P., F.H.S., &c.
Biggs, Arthur, F.L.S., F.H.S., &c.
Bingham, M. C.
Bishop, Thomas, C.M.H.S., C.H.S.,
&c.
Black, Dr.
Black, James.
Blackadder, W.
Blaikie, T., C.M.C.H.S.
Blair, Sir D. H.
Blore, G.
Bolton, Lady.
Bond, John, F.H.S., &c.
Booth, John, A.L.S., C.M.H.S., &c.
Borrer, W., F.R.S., L.S., H.S.
Bosch, J. W., C.M.H.S., Stuttgart.
Bossin, M.
Boulton, M.
Bourgeois, M., Paris.
Bourne, Frederick, F.H.S., &c.

- Bourne, W. Sturges, M.P., M.A., F.
 R.S., Z.S., H.S., &c.
 Bowens, J., F.H.S.
 Bowes, J., F.H.S., &c.
 Bowman, J. E., F.L.S.
 Braithwaite, Edmund.
 Braybroke, R. G. Lord.
 Breadalbane, J. C. Marquess of.
 Bree, Rev. W. T., A.M., C.M.H.S.,
 &c.
 Breese, J., C.M.H.S.
 Brennan, D., C.M.H.S.
 Britton, J., F.A.S., &c.
 Bromfield, W. A., M.D., F.L.S., &c.
 Bromley, William, F.L.S., H.S., &c.
 Brooks, Major J. T.
 Brown, George.
 Brown, John, F.H.S.
 Brown, M., C.M.H.S.
 Brown, Richard.
 Browne, Hay.
 Brown, Henry Augustus.
 Bruce, James.
 Bruen, J. H.
 Buccleugh, W. F. Duke of, K.G.,
 F.R.S., L.S., &c.
 Buchan, Daniel.
 Buchan, W., F.H.S.
 Buchanan, W., F.L.S., H.S., &c.
 Buchanan, W. J.
 Buel, Jesse, C.M.H.S., Albany,
 United States.
 Bullock, J.
 Bunbury, Lieut.-Gen. Sir H. E.,
 Bart., F.L.S., H.S., &c.
 Burchell, W. J., F.L.S., &c.
 Burdin, F., C.M.H.S., Turin.
 Burgess, W.
 Burlington, W. Earl of, F.H.S., &c.
 Busch, J., C.M.H.S., St. Peters-
 burg.
 Bushnan, J., F.L.S., &c.
 Butcher, Robert.
 Byers, R. W., F.L.S., &c.
- C.
- Cady, P.
 Call, M. M., C.M.H.S., St. Peters-
 burg.
 Calthorpe, G. F. Lord, F.Z.S., H.S.,
 &c.
 Cameron, D., A.L.S.
 Campbell, Lord A.
 Campbell, Lord John.
 Campbell, Alexander, A.L.S.
 Campbell, J. D.
 Canterbury, W. Lord Archbishop of.
 D.D., F.R.S., H.S., &c.
 Carberry, J. E. Lord, F.H.S., &c.
- Cardigan, Earl of.
 Carew, Lady H.
 Carnarvon, H. J. G. H. Earl of.
 Carrick, Countess of.
 Carroll, Edward.
 Carter, J.
 Case, P. J.
 Castle, G.
 Castles, Robert.
 Cathie, Nichol.
 Cato, John.
 Cawdor, J. F. Earl of, F.R.S., H.S.,
 &c.
 Chalmers, W., C.M.H.S.
 Chandler, R., F.H.S.
 Chapman, R., C.M.H.S.
 Charlwood, G., F.L.S., H.S. &c.
 Charnock, Rev. J.
 Cholmondeley, H. C. Marquess of.
 Christie, W., jun., F.L.S., &c.
 Christy, W. F.
 Churchill, Lord C. S., F.L.S., &c.
 Clanwilliam, R. M. Earl of.
 Clarke, Fingal.
 Clarke, John.
 Clarke, N.
 Clarke, Thomas.
 Clarke, William.
 Claus, C.
 Clive, E. Viscount, M.P., F.H.S.
 Clive, Hon. R. P., F.H.S., &c.
 Cockburn, J., F.H.S.
 Coghlan, A.
 Colchester, C. Lord, F.H.S.
 Collier, J., M.P.
 Collins, Peter.
 Collinson, John.
 Combermere, S. S. C. Viscount, P.
 C., &c.
 Cook, Captain, S. E.
 Cooper, J., F.H.S.
 Cooper, W.
 Copeland, W. T., M.P., F.H.S., &c.
 Cornwallis, J. M. Earl of.
 Corson, T.
 Coventry, Earl of.
 Cowan, Peter.
 Cowens, John.
 Craggs, W.
 Craven, Hon. Keppel.
 Cree, John, F.H.S.
 Crichton, Daniel.
 Cullis, John, F.H.S.
 Cullum, Rev. Sir T. G., Bart., A.M.,
 F.H.S.
 Cuming, G.
 Cumming, W. C.
 Cunningham, Allan, F.L.S., &c.
 Currer, Miss.

Curtis, S., F.L.S.
 Curzon, Hon. and Rev. Frederick,
 F.H.S., &c.

D.

Dalhousie, G. R. Earl of.
 Dall, James.
 Damer, Hon.—
 Darnley, Earl of.
 Dartmouth, W. Earl of, F.R.S., H.
 S., &c.
 Daubeny, Professor, M.D., F.R.S.,
 H. S., &c.
 Davidson, John.
 Davis, T., F.H.S.
 Dawson, T.
 Deady, M.
 De Candolle, Professor Alphonse.
 De Candolle, A. P., F.R.S., L.S.,
 H.S., &c.
 Dees, William.
 De Lamarre, M.
 De Lançon, M. Durand, Contance,
 Normandy.
 De la Warr, G. J. Earl of, F.H.S., &c.
 De l'Isle, P. C. Lord, F.H.S., &c.
 Delile, M., C.M.H.S., Montpellier.
 Denbigh, W. B. P. F. Earl of.
 Denison, J., A.L.S.
 Derum, M.
 Descemet, M.
 Deslongchamps, M. Loiseleur, Paris.
 De Vere, Sir Aubrey.
 Devon, W. Earl of.
 Devonshire, W. S. C. Duke of, K.
 G., F.H.S., &c.
 De Wael, Emilius, Antwerp.
 Diack, A., C.M.H.S.
 Dickson, Archd., M.C.H.S.
 Dickson, F., C.M.H.S.
 Dickson, Robert., M.D., F.L.S., &c.
 Dickson and Turnbull, Messrs.
 Dillwyn, L. W., M.P., F.R.S., L.S.
 Dinorben, W. L. Lord, F.H.S., &c.
 Divett, Edward, M.P.
 Dobson, —
 Dolphin, Marianne.
 Don, Professor.
 Don, G., F.L.S., &c.
 Don, J. E. S.
 Donald, Robert, F.H.S., &c.
 Donaldson, J.
 Dovaston, J. F. M.
 Downshire, A. B. S. T. H. Mar-
 quess of.
 Drummond, Dr.
 Drummond, H. Home.
 Drummond, Messrs.
 Drummond, J., C.M.H.S.

Dublin, J. Archbishop of.
 Dubreuil, M., C.M.H.S., Rouen.
 Ducie, T. R. Lord, F.R.S., H.S., &c.
 Duffie, Thomas.
 Duffies, Alexander.
 Duncan, George.
 Duncan, J.
 Duncannon, J. W. Viscount, F.H.S.
 Dunmore, G. Earl of.
 Dymock, W. G.
 Dymoke, Hon. Champion H.

E.

Eaton, James
 Ebbotton, D.
 Ebrington, Lord, M.P.
 Edmunds, C.
 Elford, Sir William, F.R.S., L.S.
 Elliott, G. M.
 Elworthy, J.
 Emslie, F.
 Esdaile, E. J.
 Essex, G. C. Earl of.
 Euston, Earl.
 Evans, —
 Ewing, John, M.P.
 Exmouth, Viscountess.

F.

Faldermann, C., C.M.H.S., St. Pe-
 tersburg.
 Farmandy, A.
 Fennell, J. F.
 Fennessey, W.
 Ferguson, Robert, M.P., F.H.S., &c.
 Ferrard, T. H. Lord Viscount, F.H.
 S., &c.
 Fintelmann, M., Potsdam.
 Fischer, Professor, Moscow.
 Fischer, C. A., C.M.H.S.
 Fischer, F. E. L., M.D., F.H.S., &c.
 St. Petersburg.
 Fitzwilliam, C. W. Earl, F.H.S.
 Fleming, Captain, R.N.
 Floud, J.
 Floyer, M.
 Folkes, Sir W. Browne, M.P.
 Forbes, James, A.L.S., &c.
 Forbes, Hon. S. R., Dresden.
 Forbes, W., M.P.
 Ford, Daniel.
 Forester, M.
 Forrest, James.
 Forrest, Richard, F.L.S., H.S.
 Forrest, Thomas, C.M.H.S.
 Forster, E., F.L.S. &c.
 Fox, H.
 Fox, J.

- Fox, Samuel.
 Fraser, John, F.H.S.
 Frederick, Sir Richard, F.H.S.
 Freeling, Sir Francis.
 Frost, Philip.
- Gamball, J.
 Gardiner, Sir R.
 Gardiner, Robert.
 Garraway, J.
 Garritt, James.
 Gathell, William.
 Gendall, J.
 Gentz, C.
 George, W.
 Gibbs, Lady.
 Gibbs, T., F.H.S., &c.
 Gibson, T.
 Gilpin, Henry.
 Glazebrook, T. K., F.L.S., &c.
 Glendinning, R.
 Goldie, John.
 Goldsmid, J. L., F.L.S., &c.
 Golton, J. Howard.
 Gordon, G.
 Gordon, W.
 Gorrie, Archibald, F.H.S., C.S., &c.
 Gosier, L'Abbé.
 Gostling, J.
 Gowan, J. R., F.H.S.
 Gower, Peter.
 Gower, Hon. William Leveson, F.
 H.S., &c.
 Grafton, G. H. Duke of, F.H.S., &c.
 Graham, Robert, M.D., F.R.S., L.S.
 Grainger, J.
 Grant, Col.
 Grant, T. Macpherson.
 Gray, T., C.M.H.S., Ropsha, St.
 Petersburg.
 Green, S.
 Greenshields, W., F.H.S.
 Gregory, Richard.
 Grey of Groby, G. H. G. Lord.
 Grierson, Wilham.
 Grigor, J.
 Groom, H., F.H.S., &c.
 Guest, T., M.P.
 Guildford, Earl of, F.H.S.
 Guthrie, Charles.
- H.
- Hailes, N.
 Halkett, C. C.
 Hamilton, Dr.
 Hammond, W.
 Harbeson James
- Harding, J. D.
 Hare, J., jun.
 Hare, W.
 Harewood, H. Earl of, F.H.S., &c.
 Harkness, H.
 Harrington, Earl of.
 Harris, John.
 Harrison, J., F.L.S., H.S., &c.
 Harrison, William, F.R.S., L.S., H.S.
 Hartland, Lord.
 Harvey, John.
 Haseley, S.
 Haslewood, William.
 Hathwaite, F.
 Haythorn, J., C.M.H.S.
 Haythorn, Z.
 Headly, J.
 Healy, F.
 Hebbert, C.
 Headerson, A., M.D., F.H.S., &c.
 Henderson, James.
 Henderson, T.
 Henslow, Professor, J. S., M.A., &c.
 Herbert, Hon. and Rev. W., F.H.S.
 Hereford, H. F. D. Lord Viscount,
 C.P.
 Heron, Captain Basil, R.
 Hervy, M.
 Higgins, Joshua.
 Hill, Captain.
 Hill, Lord, G.L.B., F.H.S., &c.
 Hislop, J., C.M.H.S..
 Hoare, Sir A. C., F.L.S., &c.
 Hodgins, Edward, C.M.H.S.
 Hodgson, Rev. G.
 Hodgson, Rev. John.
 Hodgson, N., C.M.H.S.
 Holford, R. S., F.L.S., H.S., &c.
 Holland, H. R. V. Lord, P.C., &c.
 Holt, W.
 Honner, R., C.M.H.S., Hobart
 Town, Van Diemen's Land.
 Hood, C. H. Lord Viscount.
 Hoogvorsts, Baron.
 Hooker, H.
 Hooker, Sir W. J., F.L.S., H.S., &c.
 Hope, H. J., M.P.
 Hope, Professor, T. C., F.R.S.
 Hopgood, Thomas.
 Hosack, D., M.D., F.H.S.
 Hoscic, R., C.M.H.S.
 Howard, Robert.
 Howden, John.
 Howden, Gen., Lord.
 Hughes, Robert.
 Hume, Sir Abraham, Bart., F.H.S.
 Hunneman, J., A.L.S.
 Hunt, T. K.
 Hunt, T. L.

Hutchinson, William
Hutton, R., F.H.S., &c.

I.

Ireland, R.
Irvine, Robert.
Irving, T.

J.

Jackson, Thomas.
Jacquin, Baron, J. F., F.H.S., &c.,
Vienna
James, J.
Jardine, Sir William Bart., F.R.S.E.,
L.S., &c.
Jennings, John.
Johnson, J.
Johnston, Sir Alexander, F.R.S.
Johnston, R.
Johnstone, Edmund, F.H.S.
Johnstone, Henry, M.P.
Jones, B.
Judd, D., F.H.S.
Jukes, H. W.

K.

Karwinsky, Baron.
Kelly, Rev. Walter.
Ker, W.
King, W. Lord, F.H.S., &c.
Klouweitz, Hans.
Knight, T. A., F.R.S., L.S., P.H.S.
Knight, J., F.H.S., &c.
Knipe, J. B.
Knowles, G. B., F.L.S., &c.
Kops, Professor, Utrecht.

L.

Labouchère, P. C., F.H.S., &c.
La Croix, L. L., Paris.
La Gasca, Professor, F.L.S., &c.,
Madrid.
Laidlaw, W., C.M.H.S.
Lainell, J.
Lamb, John.
Lamb, Sir Charles M., F.H.S., &c.
Lambert, A. B., F.R.S., V.P.L.S.,
Langelier, P., C.M.H.S., Jersey.
Lansdowne, Henry Marquess of,
F.R.S., L.S., H.S., &c.
Latham, R., F.L.S., H.S., &c.
Lauder, Sir Thomas Dick, Bart.
Law, J.
Lawrence, Charles, F.H.S., &c.
Lawson, Charles.

Lawson, John.
Lawson, P. and Son.
Le Blanc, Thomas.
Le Candele, J., Humbeque Nursery,
near Antwerp.
Lees, E., F.L.S., &c.
Lemon, Sir C., Bart., M.P., F.H.S.
Lenné, M., Potsdam.
Le Roy, M. Angers.
Lewin, F. E.
Lewis, G. R.
Lichfield and Coventry, Hon. H.
Ryder, Bishop of.
Lindsay, T. M.
Lines, J.
Llewelyn, J. D., F.H.S., &c.
Loch, J., M.P., F.G.S., &c.
Loddiges, Conrad, and Sons.
London, C. J. Bishop of, D.D., &c.
Long, H. L.
Longman, T. N., F.H.S., &c.
Lorimer, Robt.
Lothian, J. W. R. Kerr Marquess of.
Loudon, James, Warsaw.
Loudon, James, Ham House.
Low, H., F.H.S.
Lowndes, H.
Lowndes, Richard.
Lucombe and Pince, Messrs.
Luke, Richard.
Lye, W. L.
Lyll, Charles, jun., F.R.S., L.S.,
G.S., &c.
Lyon, Geo.
Lyttleton, W. H. Lord.

M.

M^cArthur, P., F.H.S.
Mackay, F.
Mackay, J. T., A.L.S., M.R.I.A., &c.
Mackey, Constantine.
Mackenzie, Sir Jas. Stuart, Bart.,
M.P., F.H.S., &c.
Mackenzie, P., Philadelphia.
Mackie, J.
MacLeay, A., F.R.S., L.S., &c.
M^cNab, J., Edinburgh.
M^cNab, T., C.M.H.S., Dollar.
M^cNab, W., A.L.S., Edinburgh.
Maddison, J., Ghent.
Main, James, A.L.S., &c.
Major, J.
Malcolm, William, F.L.S., H.S., &c.
Mallet, R., A.M., &c.
Malpas, B.
Manetti, Giuseppe, C.M.H.S., Monza.
Mangles, Capt. F.
Mangles, R., F.H.S.

- Manners, Mrs. L.
 Marnock J.
 Marnock, R.
 Marryatt, Mrs., F.H.S.
 Marsham, R.
 Martin, John.
 Martin, M. J.
 Martius, Dr., Munich.
 Mason, Rev. J. H.
 Masters, W., F.H.S.
 Mathews, Maria Ruthven.
 Mathies, J.
 Matthews, A., A.L.S., Lima.
 Maund, B., F.L.S., &c.
 Maunder, W.
 Maxwell, J., M.P., F.R.S., H.S., &c.
 Maxwell, Lady Heron.
 May, E.
 May, G.
 M'Donald, J., C.M.H.S.
 M'Duff, J.
 M'Diarmaid, William.
 Meara, J.
 Mearns, J., F.A.S.
 Mease, Dr., Philadelphia.
 Melbourne, Lord Viscount, P.C., &c.
 Melville, Lord Viscount, K.T., P.C., &c.
 Menteth, James Stuart.
 Menzies, A., F.L.S., &c.
 Metzger, M., C.M.H.S., Heidelberg.
 Michaux, André, Paris.
 Middlemiss, R.
 Miller, J., A.L.S., F.H.S.
 Mills, J., F.H.S.
 Mills, Philip John, M.P.
 Mills, Rev. Thomas.
 M'Intosh, Alexander.
 M'Intos' C., F.H.S.
 M'Intyre, James.
 Milne, T., F.H.S., A.L.S., &c.
 Mirbel, Professor, F.L.S., &c., Paris.
 Mitchenson, J.
 Mitford, Rev. J.
 M'Launaehan, A.
 M'Leish, George.
 M'Leod, Donald.
 M'Murtrie, W., F.H.S.
 Moffat, J.
 Moffat, Thomas.
 Monck, Sir C., F.H.S., &c.
 Monckton, Lieut.-General H., F.H.S.
 Montagu, Lord.
 Moray, Earl of, K.T., &c.
 Morgan, Octavius.
 Morley, Earl of, F.R.S., &c.
 Mosley, Sir O., Bart., F.H.S.
 Moss, E.
 Moss, William.
 Mount Cashell, Earl of.
 Mountrorris, G., Earl of, F.R.S., H.S., &c.
 M'Pherson, A.
 Muirhead, Alexander, C.M.H.S.
 Munchausen, Baron, Hanover.
 Munro, James.
 Murdoch, A., C.M.H.S.
 Mure, William.
 Murphy, Edward.
 Murray, Mungo.
 Murray, Stewart, F.H.S., &c.
 Murray, W., F.H.S.
- N.
- Napier, Major-General, Charles Jas.
 Nash, C.
 Nasmyth, Sir J., Bart.
 Neill, Dr. P., F.L.S., &c.
 Nérières, M., Nantes.
 Nesfield, W. A.
 Nevill, Rev. W. H.
 Newberry, F.
 Newburgh, Earl of.
 Newell, J.
 Nickson, J.
 Nicol, W.
 Niven N., C.M.H.S.
 Normann, Baron.
 Northampton, Marquess of, F.H.S.
 Northumberland, Hugh Duke of, K.G., F.R.S., L.S., &c.
 Nott, W.
- O.
- Ogle, H. C.
 Oldaker, J., F.H.S.
 Oliver, J., C.M.H.S.
 Osborn, R., F.H.S.
 Osborn, William.
 Otto, F., C.M.H.S., Berlin.
- P.
- Page, W. B., C.M.H.S.
 Pauplin, W., jun., A.L.S.
 Pappenheim, M. le Baron, Paris.
 Parks, J., F.H.S.
 Parker, T. W.
 Parkins, Thomas.
 Paterson, W.
 Patterson, M.
 Paxton, Josh., F.L.S., H.S., &c.
 Pearson, B.
 Pearson, J.
 Pease, J., M.P.
 Penny, G., A.L.S.

Perkins, H.
 Petersen, J. P.
 Phipps, Hon. and Rev. Augustus, J.
 Pince, R. T.
 Plumtre, J. P., M.P.
 Plumtree, Rev. H.
 Polhill, G.
 Poore, Sir Edward, Bart.
 Pope, J., and Sons.
 Porter, G.
 Portland, Duke of.
 Pound, M.
 Powis, Edward Earl of, F.H.S.
 Poyntz, W. S., M.P., F.H.S., &c.
 Pratt, Henry.
 Prescott, A.
 Press, Thomas.
 Preston, C.
 Prevot, fils.
 Priest, Myles.
 Price, Sir Robert, Bart., M.P., F.H.S.
 Pringle, A., M.P.

R.

Racine, M.
 Radcliffe, Anna.
 Radnor, Earl of, F.H.S.
 Ramsey, H.
 Rauch, Charles.
 Rauch, Francis.
 Rauch, Stephen.
 Reeves, J., F.R.S., H.S., &c.
 Reid, J., C.M.H.S.
 Reinhardt, Professor C. G. C.,
 F.L.S., Leyden.
 Rhodes, Rev. J. A.
 Richardson, A.
 Richardson, R. H.
 Richardson, W.
 Richmond, Duke of.
 Riddell, George.
 Riegler, M.
 Ripon, F. J., Earl of, F.R.S., H.S.
 Rivers, T., jun.
 Robert, M.
 Roberts, Job.
 Roberts, T.
 Robertson, F.
 Robertson, John, F.H.S.
 Robins, G.
 Roek, D.
 Roden, Earl of, K.P., P.C., &c.
 Rodney, Hon. W.
 Rodney, Lord.
 Rogers, J., C.M.H.S.
 Rogers, Rev. J., F.H.S.
 Rolle, J. Lord, F.H.S., &c.
 Rollison, G., F.H.S.

Ronalds, J., F.H.S.
 Ross, J., C.M.H.S.
 Roy, J., Aberdeen.
 Royle, J. F., M.D., F.R.S., L.S., &c.
 Rutger, T.
 Rutland, Duke of, H.C., &c.

S.

Salisbury, Marquess of, F.H.S., &c.
 Salmon, Rev. T. W.
 Sanders, G.
 Sanderson, Robert.
 Sang, E., M.C.H.S.
 Saunders, B., of Jersey.
 Saunders, R.
 Schneevoght, G. V., C.M.H.S.,
 Haarlem.
 Schoch, M.
 Schouw, Professor, Copenhagen.
 Schubert, M., C.M.H.S., Warsaw.
 Sekell, C., C.M.H.S., Munich
 Scott, J.
 Seitz, C. J.
 Shaftesbury, Countess of.
 Shepherd, H., F.L.S., &c.
 Shields, George.
 Shirley, W. P.
 Shrewsbury, J. Earl of, F.H.S., &c.
 Shuttleworth, Charles.
 Sidmouth, Lord Viscount, P.C., &c.
 Silverlock, G., F.H.S., &c.
 Simmons, T.
 Simpson, Henry.
 Simpson, Joseph.
 Sinclair, James.
 Sinnot, Rev. Dr.
 Skelmersdale, Edward B. Lord,
 F.H.S.
 Skinner, J., C.M.H.S.
 Smallman, John, Architect.
 Smith, Alexander.
 Smith, D.
 Smith, James, F.H.S.
 Smith, John, A.L.S., F.H.S., &c.
 Smith, R.
 Smith, Rev. Sydney.
 Smith, W., F.H.S.
 Somerset, Duke of, K.G., F.R.S.,
 P.L.S.
 Soulange-Bodin, Le Chevalier E.,
 C.M.H.S., Fromont.
 Sowerby, J. D. C., F.L.S., &c.
 Sowerby, C. E., A.L.S.
 Sparrow, John.
 Speirs, Alexander, M.P.
 Spence, W., F.R.S., L.S., &c.
 Spencer, J.
 Spiker, Dr., Berlin.

Spong, Charles, C.M.H.S.
 Spring, C.
 St. Aubyn, Sir John, Bart., F.L.S.
 Stamford, Earl of.
 Stanhope, P. H. Earl, F.R.S., H.S.
 Stanley, E., M.P.
 Stanley, Sir T. S. M., Bart., F.H.S.
 Staunton, Sir George Thomas, Bart.,
 F.R.S., L.S., &c.
 Stephens, H.
 Steven, C., C.M.H.S., Nikitka, Cri-
 mea.
 Stewart, A.
 Stewart, D., F.R.S., L.S., &c.
 Stewart, Robert Bruce.
 Stewart, Sir M. Shaw, M.P.
 St. Germain, Earl.
 St. Marie, M.
 Stock, Daniel, A.L.S.
 Stone, George.
 Storey, Rev. —
 Stowell, Lord.
 Strachan, G.
 Strangways, Hon. W. T. H. F.,
 F.H.S., &c.
 Street, J., C.M.H.S.
 Strickland, Miss.
 Strom, M.
 Strong, Charles.
 Stuart, W.
 Surrey, C. H. Earl of, M.P., F.H.S.,
 Sutherland, G. G., Duke of, F.H.S.,
 Sydney, M. J. F.

T.

Tait, John.
 Taunton, W. P.
 Taylor, F., C.M.H.S.
 Taylor, S.
 Taylor, Thomas, M.D., F.L.S., &c.
 Taylor, William.
 Templeton, Mrs.
 Thorburn, G., C.M.H.S., New York.
 Thorn, William.
 Tighe, William F.
 Tongard, M.
 Toward, Andrew.
 Trail, Miss, F.H.S.
 Traill, James, A.L.S., Cairo.
 Trevelyan, W. C., F.H.S., &c.
 Turnbull, A., C.M.H.S.
 Turner, G. and T.
 Turner, H., F.H.S.
 Turner, T.
 Tyler, W.
 Tyso, Rev. J.

V.

Vandamme, M.

Vandergelt, M.
 Vandermaelen, M., F.H.S., &c.
 Van Eden, A. C., C.M.H.S., Haar-
 lem.
 Van Wael, M., Antwerp.
 Vaughan, Sir R. W., Bart., F.H.S.
 Varden, R.
 Veitch, J., F.H.S.
 Veltheim, Graf, Hanover.
 Verney, H.
 Vernon, Lord.
 Verulam, Earl of.
 Vilmorin, P. P. A., C.M.H.S., Paris.
 Vulliamy, T. S.
 Vymer, R.

W.

Wagstaff, E.
 Wakeford, Robert.
 Walker, C.
 Walker, J., F.H.S.
 Walker, T. E.
 Wall, C. B., M.P., F.R.S., H.S., &c.
 Wallace, J.
 Walloch, N., M.D., F.R.S., L.S., &c.
 Walsingham, Rev. T. Lord, F.L.S.
 Walsh, Rev. R., L.L.D., C.M.H.S.
 Walter, J., M.P.
 Wank, Franz, Briick on the Leytha.
 Ward, John, F.H.S., Z.S., &c.
 Ward, Lady.
 Warmesley, G., M.P.
 Warren De Tably, Lord.
 Waterer, M., F.H.S., &c.
 Watkin, A.
 Watson, H. C., F.L.S., &c.
 Watson, J.
 Watt, James.
 Watts, W.
 Webster, J., F.H.S.
 Webster, William.
 Weddell, H.
 Wells, W., F.H.S., &c.
 Wemyss, Earl of.
 Westminster, Marquess of, P.C.
 Westwood, J. O., F.L.S.
 Whalley, J., F.H.S.
 White, J., F.H.S.
 Whittit, G., M.C., H.S.
 Wicklow, Earl of.
 Wilkes, R. S.
 Wilkins, W., C.M.H.S.
 Williams, G. W., M.P.
 Wilmore, Mrs.
 Wilmot, Sir Eardley.
 Wilson, D.
 Wilson, J., C.M.H.S.
 Wilson, John S.

Wilson, Neill.
 Wilson, Sir Griffin.
 Wilson, Sir T. M., F.H.S.
 Winchester, Bishop of, F.H.S., &c.
 Winch, N. J., A.L.S.
 Winchester, Henry, F.H.S., &c.
 Wodzicki, Count Stanislaus, Cracow.
 Wood, G.
 Wood, J. F. F.
 Wood, Neville.
 Woods, H., A.L.S.
 Woolf, M.
 Wooluck, J.
 Wraight, W.
 Wright, Henry.

Wrightson, Mrs.
 Wrottesley, Sir J., Bart., M.P.

Y.

Youell, J., A.L.S.
 Young, A.
 Young, Messrs. C. J. and P.
 Young, J., C.M.H.S.
 Young and Penny, Messrs.
 Young, R., F.L.S.
 Young, William, M.D., F.L.S.

Z.

Zettersteck, M.

EPITOME OF THE CONTENTS.

The Roman numerals refer to the Contents in full; the Arabic figures, to the body of the work.

VOL. I.

	Contents.	Text.		Contents.	Text.
INTRODUCTION	-	-	xiii.	I	
<hr/>					
Part I. GEOGRAPHY AND HISTORY	-	-	xiii.	15	
CHAP. I.					
Among the Nations of Antiquity	-	-	xiii.	15	
CHAP. II.					
In the British Islands	-	-	xiii.	20	
SECT. I. Native Trees and Shrubs	-	-	xiii.	20	
SECT. II. Foreign Trees and Shrubs introduced	-	-	xiii.	31	
SUBSECT. 1. Trees and Shrubs introduced by the Romans	-	-	xiii.	32	
SUBSECT. 2. Trees and Shrubs introduced in the 16th Century	-	-	xiii.	35	
SUBSECT. 3. Trees and Shrubs introduced in the 17th Century	-	-	xiii.	40	
SUBSECT. 4. Trees and Shrubs introduced in the 18th Century	-	-	xiii.	44	
SUBSECT. 5. Trees and Shrubs introduced in the 19th Century	-	-	xiv.	117	
CHAP. III.					
History and Geography of the Trees and Shrubs of the Continent of Europe	-	-	xiv.	132	
SECT. I. Trees and Shrubs of France	-	-	xiv.	132	
SECT. II. Trees and Shrubs of Holland and the Netherlands	-	-	xiv.	143	
SECT. III. Trees and Shrubs of Germany and Hungary	-	-	xiv.	145	
SECT. IV. Trees and Shrubs of Denmark, Holstein, Sweden, Lapland, Finland, Iceland, Greenland, and the Faroe Islands	-	-	xv.	153	
SECT. V. Trees and Shrubs of Russia and Poland	-	-	xv.	155	
SECT. VI. Trees and Shrubs of Switzerland	-	-	xv.	161	
SECT. VII. Trees and Shrubs of Italy, Greece, Spain, Portugal, and the Mediterranean Islands	-	-	xv.	164	
SUBSECT. 1. Trees and Shrubs of Italy	-	-	xv.	168	
SUBSECT. 2. Trees and Shrubs of Spain and Portugal	-	-	xv.	170	

	Contents. Text.		Contents. Text.
SUBSECT. 3. Trees and Shrubs of Turkey and Modern Greece - - -	xv. 171	SECT. I. In Uncultivated Nature - - -	xvi. 220
CHAP. IV.		SECT. II. With Reference to Man - - -	xvi. 221
Trees and Shrubs of Asia, Africa, America, and Australia, suitable for the Climate of Britain - - -	xv. 172	CHAP. IV.	
SECT. I. Trees and Shrubs of Asia suitable for Britain - - -	xv. 172	Summary of the Description and Natural and Economical History of Trees and Shrubs - - -	xvi. 222
SECT. II. Trees and Shrubs of Africa suitable for Britain - - -	xv. 177		
SECT. III. Trees and Shrubs of America suitable for Britain - - -	xv. 173	Part III. ARBORETUM AND FRUTICETUM. - - -	xvii. 231
SUBSECT. 1. Trees and Shrubs of North America - - -	xv. 178	<i>Ranunculàcææ</i> Dec. - - -	xvii. 231
SUBSECT. 2. Trees and Shrubs of South America - - -	xv. 185	<i>Clématis L.</i> - - -	xvii. 232
SECT. IV. Trees and Shrubs of Australia and Polynesia - - -	xv. 185	<i>Atrágene L.</i> - - -	xviii. 246
CHAP. V.		<i>Pæoniàcææ</i> Dec. - - -	xviii. 249
Literature of the Trees and Shrubs of Temperate Climates - - -	xv. 187	<i>Pæonia L.</i> - - -	xviii. 249
CHAP. VI.		<i>Xanthorhiza L.</i> - - -	xviii. 255
Conclusion - - -	xvi. 190	<i>Winteràcææ R. Br.</i> - - -	xviii. 256
		<i>Illicium L.</i> - - -	xviii. 256
Part II. SCIENCE OF THE STUDY OF TREES - - -	xvi. 192	<i>Magnoliàcææ</i> Dec. - - -	xix. 259
CHAP. I.		<i>Magnolia L.</i> - - -	xix. 260
Study of Trees Pictorially - - -	xvi. 193	<i>Liriodéndron L.</i> - - -	xix. 284
SECT. I. Study of Forms - - -	xvi. 193	<i>Mangliétia</i> - - -	xix. 291
SECT. II. Study of Expression and Character - - -	xvi. 198	<i>Michelia</i> - - -	xix. 291
SECT. III. Mode of drawing Trees from Nature, so as to give the Pictorial Expression of the Species - - -	xvi. 202	<i>Dilleniàcææ</i> Dec. - - -	xix. 292
CHAP. II.		<i>Ilibértia</i> - - -	xix. 292
Trees and Shrubs considered Botanically - - -	xvi. 211	<i>Anonàcææ</i> Rich. - - -	xx. 292
SECT. I. Classification - - -	xvi. 211	<i>Asimina</i> Adans. - - -	xx. 292
SECT. II. Distinction between Species and Varieties - - -	xvi. 212	<i>Annóna L.</i> - - -	xx. 292
SECT. III. Mode of describing Trees and Shrubs - - -	xvi. 212	<i>Orchidocírcupum</i> Mx. - - -	xx. 292
CHAP. III.		<i>Porcélia</i> Pers. - - -	xx. 292
Uses of Trees and Shrubs - - -	xvi. 219	<i>Schizandrácææ</i> Blume. - - -	xx. 295
		<i>Schizándra</i> Mx - - -	xx. 295
		<i>Sphaerostéma</i> Bl. - - -	xx. 295
		<i>Kadsura J.</i> - - -	xx. 295
		<i>Sarcocodryon</i> Bl. - - -	xx. 295
		<i>Menispermàcææ</i> Dec. - - -	xx. 296
		<i>Menispermum L.</i> - - -	xx. 296
		<i>Cócculus Bauh.</i> - - -	xx. 297
		<i>Wendlandia Willd.</i> - - -	xx. 297
		<i>Andróphilax</i> Wench. - - -	xx. 297
		<i>Baumgártia Mench.</i> - - -	xx. 298
		<i>Berberàcææ</i> Lindl. - - -	xx. 298
		<i>Berberidææ</i> Vent. - - -	xx. 298
		<i>Bérberis L.</i> - - -	xx. 298
		<i>Mahónia</i> Nutt - - -	xxi. 308
		<i>Odosstémon</i> Raf. - - -	xxi. 308
		<i>Nandina</i> - - -	xxvi. 257
		<i>Crucifèrææ</i> Lindl. - - -	xxi. 312
		<i>Crucifèrææ</i> Juss. - - -	xxi. 312
		<i>Vélla L.</i> - - -	xxi. 312
		<i>Cheiránthus</i> - - -	xxi. 313
		<i>Vesciaria</i> - - -	xxi. 313
		<i>Ay-sum</i> - - -	xxi. 313
		<i>Ibèris</i> - - -	xxi. 313
		<i>Lepidium</i> - - -	xxi. 313
		<i>Sisymbrium</i> - - -	xxi. 313

	Contents.	Text.		Contents.	Text.
<i>Capparidææ</i> Lindl.	xxi.	313	<i>Hypéricum L.</i>	xxix.	397
<i>Capparidææ</i> Juss.			<i>Trágium</i> Clus.	xxix.	397
<i>Cápparis L.</i>	xxi.	313	<i>Androsæmum Chois.</i>	xxx.	403
<i>Cistææ</i> Lindl.	xxi.	316	<i>Clymenon</i> L'Obel.	xxx.	403
<i>Cistineæ</i> Dec.			<i>Acerææ</i> Lindl.	xxx.	404
<i>Cístus L.</i>	xxi.	317	<i>Acerineæ</i> Dec.		
<i>eliánthemum Tourn.</i>	xxii.	328	<i>A'cer</i>	xxx.	405
<i>Hudsonia L.</i>	xxv.	354	<i>Negúndo Mœnch</i>	xxxii.	460
<i>Polygalææ</i> Lindl.	xxvi.	355	<i>Negúndium Rafin.</i>		
<i>Polygónææ</i> Juss.			<i>Dobinea</i> Hamilt.	xxxii.	462
<i>Po'ýgala L.</i>	xxvi.	356	<i>Æsculææ</i> Lindl.	xxxii.	462
<i>Pittosporææ</i> Lindl.	xxvi.	356	<i>Castaneææ</i> Lk.		
<i>Pittosporææ</i> R. Br.			<i>Hippocástaneæ</i> Dec.		
<i>Billardiæra</i> Sm.	xxvi.	356	<i>Æ'sculus L.</i>	xxxii.	462
<i>Sólyra</i> Lindl.	xxvi.	357	<i>Hippocástanum</i> Tourn.		
<i>Senécia</i> Dec.	xxvi.	359	<i>Pavía</i> Boer.	xxxii.	469
<i>Bursaria</i> Cav.	xxvi.	359	<i>Sapindææ</i> Juss.	xxxiii.	475
<i>Cheiranthæra</i> Lindl.	xxvi.	359	<i>Költreutéria</i> Laxm.	xxxiii.	475
<i>Caryophyllææ</i> Lindl.	xxvi.	359	<i>Sapíndus L.</i> fil.		
<i>Caryophýllææ</i> Juss.			<i>Dodone'a L.</i>	xxxiii.	476
<i>Diánthus L.</i>	xxvi.	359	<i>Meliææ</i> Juss.	xxxiii.	476
<i>Siléne L.</i>	xxvi.	359	<i>Mélia L.</i>	xxxiii.	477
<i>Arenária W.</i>	xxvi.	359	<i>Vitææ.</i> Lindl.	xxxiii.	477
<i>Drypis L.</i>	xxvi.	359	<i>Vítis L.</i>	xxxiii.	477
<i>Linææ</i> Lindl.	xxvi.	360	<i>Ampelópsis Michx.</i>	xxxiii.	481
<i>Lineæ</i> Dec.			<i>Cissus L.</i>	xxxiii.	483
<i>Linum L.</i>	xxvi.	360	<i>Geraniææ</i> Dec.	xxxiv.	483
<i>Malvææ</i> Juss.	xxvi.	360	<i>Pelargónium L'Herit.</i>	xxxiv.	483
<i>Lavatera L.</i>	xxvi.	360	<i>Zygophyllææ</i> Lindl.	xxxiv.	484
<i>Hibiscus L.</i>	xxvi.	361	<i>Zygophýllææ</i> R. Br.		
<i>Sida Bonp.</i>	xxvi.	363	<i>Meliánthus L.</i>	xxxiv.	484
<i>Abúttion Hook.</i>	xxvi.	363	<i>Zygophýllum L.</i>	xxxiv.	484
<i>Sterculiææ</i> Vent.	xxvi.	363	<i>Rutææ</i> Lindl.	xxxiv.	484
<i>Sterculia L.</i>	xxvi.	363	<i>Rútææ</i> Juss.		
<i>Tiliææ</i> Juss.	xxvii.	364	<i>Rúta L.</i>	xxxiv.	484
<i>Tília L.</i>	xxvii.	364	<i>Apophýllum And. Juss.</i>	xxxiv.	487
<i>Gréwia L.</i>	xxvii.	366	<i>Xanthoxylææ</i> Lindl.	xxxiv.	487
<i>Ternströmiææ</i> Dec.	xxvii.	376	<i>Xanthoxýllææ</i> Nees.		
<i>Malachodéndron Cav.</i>	xxvii.	377	<i>Pteleææ</i> Kunth.		
<i>Stewartia L.</i>			<i>Xanthóxyllum L.</i>	xxxiv.	488
<i>Stuártia Cav.</i>	xxviii.	378	<i>Kampmánnia Rafin.</i>		
<i>Gordónia Ellis</i>	xxviii.	378	<i>Fagára Lam.</i>		
<i>Lacathæa</i> Sal.			<i>Ptèlea L.</i>	xxxiv.	489
<i>Franklínia Marsh.</i>			<i>Bellúcia Adams.</i>		
<i>Caméllia L.</i>	xxviii.	381	<i>Ailántus Desf.</i>	xxxiv.	490
<i>Théa</i>	xxix.	392	<i>Coriææ</i> Lindl.	xxxiv.	493
<i>Eúrya Royle</i>	xxix.	395	<i>Coriæææ</i> Dec.		
<i>Aurantîææ</i> Corr.	xxix.	395	<i>Coriária Niss.</i>	xxxiv.	492
<i>Citrus L.</i>	xxix.	395	<i>Staphyleææ</i> Dec.	xxxiv.	492
<i>Límónia Dec.</i>	xxix.	396	<i>Staphyléa L.</i>	xxxiv.	493
<i>Hypericææ</i> Lindl.	xxix.	396	<i>Staphylodéndron</i> Tourn.		
<i>Hypericîneæ</i> Dec.					
<i>Celastrææ</i> Dec.	xxxiv.	495	<i>Máytenus</i> Fenill	xxxv.	503
<i>Euónymus Tourn.</i>	xxxiv.	496	<i>Senécia Lam.</i>		
<i>Celástrus L.</i>	xxxv.	502	<i>Cussine L.</i>	xxxv.	503
<i>Euonymóides</i> Mœnch.			<i>Maurocénia Mill.</i>		
<i>Nemopánthes Raf.</i>	xxxv.	503	<i>Hartóglia Dec.</i>	xxxv.	504
<i>Ilícioídes Dum. Cours.</i>			<i>Schrebera Thunb.</i>		
			<i>Elæocéndron Spreng.</i>		

VOL. II.

	Contents.	Text.		Contents.	Text.
<i>Aquifoliaceæ</i> Dec.	xxxvi.	505	<i>Podalyria</i> R. Br.	xli.	568
<i>Myginda</i> Jacq.	xxxvi.	505	<i>Chorizanthe</i> R. Br.	xli.	568
<i>Plex</i> L.	xxxvi.	505	<i>Podolobium</i> R. Br.	xli.	568
<i>Aquifolium</i> Tourn.			<i>Oxylobium</i> R. Br.	xli.	568
<i>Prinos</i> L.	xxxvi.	520	<i>Callistachys Sims</i>	xli.	568
<i>Agéria</i> Adans.			<i>Brachysema</i> R. Br.	xli.	569
<i>Rhamnaceæ</i> Lindl.	xxxvii.	523	<i>Gompholobium Smith</i>	xli.	569
<i>Rhamnæ</i> Dec.			<i>Burtonia</i> R. Br.	xli.	569
<i>Zizyphus Tura.</i>	xxxviii.	521	<i>Jacksonia</i> R. Br.	xli.	569
<i>Nabea Arn.</i>			<i>Viminaria Sm.</i>	xli.	569
<i>Ænophia Bauh.</i>			<i>Sphaerolobium Sm.</i>	xli.	569
<i>Paliurus</i> L.	xxxvii.	527	<i>Aotus G. Don</i>	xli.	569
<i>Berberis</i> Neck.	xxxvii.	528	<i>Dillwynia Sm.</i>	xli.	569
<i>Rhamnus</i> Lam.	xxxvii.	529	<i>Eutasia R. Br.</i>	xli.	569
<i>Atalernus</i> Mill.			<i>Sclerolobium R. Br.</i>	xli.	569
<i>Ceanothus</i> L.	xxxviii.	539	<i>Gastrolobium R. Br.</i>	xli.	569
<i>Spharocarya</i> Wall.	xxxix.	541	<i>Euchitus R. Br.</i>	xli.	569
<i>Condalia Gay.</i>	xxxix.	541	<i>Pultenæa Bot. Mag.</i>	xli.	569
<i>Sageretia Brongn.</i>	xxxix.	541	<i>Daviesia R. Br.</i>	xli.	569
<i>Scutia Brongn.</i>	xxxix.	541	<i>Mirbesia Smith</i>	xli.	570
<i>Retanilla Brongn.</i>	xxxix.	541	<i>Ulex</i> L.	xli.	571
<i>Colletia Kunth</i>	xxxix.	541	<i>Stauracanthus</i> Lk.	xli.	576
<i>Trevoa Mey.</i>	xxxix.	541	<i>Spartium</i> Dec.	xli.	576
<i>Piscaria Hook.</i>	xxxix.	541	<i>Spartianthus</i> Lk.		
<i>Hovenia Thunb.</i>	xxxix.	541	<i>Genista</i> Lam.	xli.	577
<i>Colubrina Brongn.</i>	xxxix.	542	<i>Genistoides Mæneh.</i>		
<i>Willemetia Brongn.</i>	xxxix.	542	<i>Cytisus</i> L.	xliii.	588
<i>Pomaderris Lab.</i>	xxxix.	542	<i>Adenocarpus</i> Dec.	xliv.	603
<i>Cryptandra Smith</i>	xxxix.	542	<i>Ononis</i> L.	xliv.	604
<i>Hartungia Brongn.</i>	xxxix.	542	<i>Anonis Mæneh.</i>		
<i>Solenanthe G. Don</i>	xxxix.	542	<i>Natrix Mæneh.</i>		
<i>Trichoccephalus Brongn.</i>	xxxix.	542	<i>Amorpha</i>	xlv.	606
<i>Phytica</i> L.	xxxix.	542	<i>Bonafidia</i> Neck.		
<i>Soulingia Brongn.</i>	xxxix.	542	<i>Robinia</i> L.	xlv.	609
<i>Gouania Jacq.</i>	xxxix.	542	<i>Pseudacacia</i> Tourn.		
<i>Carpodetus Forst.</i>	xxxix.	542	<i>Æschynomene</i> Roxb.		
<i>Oleua</i> Thunb.	xxxix.	542	<i>Caragana</i> Lam.	xlv.	629
<i>Bruniaceæ</i> R. Br.	xxxix.	542	<i>Halimodendron</i> Fisch.	xlv.	634
<i>Homalinaceæ</i> Lindl.	xxxix.	542	<i>Halodendron</i> Dec.		
<i>Homalinæ</i> R. Br.			<i>Calophaca</i> Fisch.	xlvi.	635
<i>Aristotelia</i> L'Hérit.	xxxix.	543	<i>Cytisus</i> Pall.		
<i>Azara</i> R. et P.	xxxix.	544	<i>Colutea</i> R. Br.	xlvi.	637
<i>Blackwellia</i> Dec.	xxxix.	544	<i>Astragalus</i> Dec.	xlvi.	637
<i>Astranthus Lour.</i>	xxxix.	544	<i>Dorcnium Tourn.</i>	xlvii.	639
<i>Neillia</i> D. Don.	xxxix.	544	<i>Plagiolobium Sret.</i>	xlvii.	639
<i>Anacardiaceæ</i> Lindl.	xxxix.	545	<i>Platylobium Sm.</i>	xlvii.	639
<i>Anacardiæ</i> Brown.			<i>Hovea</i> R. Br.	xlvii.	639
<i>Pistacia</i> L.	xxxix.	545	<i>Boschia Vent.</i>	xlvii.	640
<i>Terebinthus</i> Juss.			<i>Goeldia Sal.</i>	xlvii.	640
<i>Rhus</i> L.	xxxix.	548	<i>Scutia R. Br.</i>	xlvii.	640
<i>Côtinus</i> Scop.			<i>Templetonia Sims</i>	xlvii.	640
<i>Toxicodendron</i> Mill.			<i>Rafnia</i> Thunb.	xlvii.	640
<i>Myrica</i> Hort.			<i>Vascoa</i> Dec.	xlvii.	640
<i>Duvaâa</i> Kunth	xl.	558	<i>Borbonia</i> L.	xlvii.	640
<i>Sabia</i> Wall.	xl.	560	<i>Achyronia Wendt.</i>	xlvii.	640
<i>Schinus</i> L.	xl.	560	<i>Epiparia</i> L.	xlvii.	640
<i>Fricetos Lour.</i>	xl.	560	<i>Proctéra</i> Dec.	xlvii.	640
<i>Heptadendron Desf.</i>	xl.	560	<i>Crotalaria</i> L.	xlvii.	640
<i>Stylobasium Desf.</i>	xl.	560	<i>Hypocalyptus Thunb.</i>	xlvii.	640
<i>Cneorum</i> L.	xl.	560	<i>Viborgia Spreng.</i>	xlvii.	640
<i>Burseraceæ</i> Kunth.	xl.	561	<i>Loddigesia Sims</i>	xlvii.	641
<i>Halimodendron Kunth</i>	xl.	561	<i>Dieltus</i> Dec.	xlvii.	641
<i>Pharium R. Br.</i>	xl.	561	<i>Leleckia</i> Thunb.	xlvii.	641
<i>Fagostrium G. Don.</i>	xl.	561	<i>Sarcophyllum Thunb.</i>	xlvii.	641
<i>Amyridaceæ</i> Lindl.	xl.	561	<i>Aspidanthus</i> L.	xlvii.	641
<i>Amyrideæ</i> R. Br.			<i>Requienia</i> Dec.	xlvii.	641
<i>Amyris</i> L.	xl.	561	<i>Anthyllis</i> L.	xlvii.	641
<i>Leguminosæ</i> Juss.	xl.	561	<i>Fulneraria</i> Lam.		
<i>Sophora</i> R. Br.	xl.	563	<i>Medicago</i> L.	xlvii.	642
<i>Virgilia</i> L.	xli.	565	<i>Mellilotus Cust.</i>	xlvii.	642
<i>Piptanthus Swt.</i>	xli.	566	<i>Lotus</i> L.	xlvii.	642
<i>Thermopsis</i> D. Dop.			<i>Carmichaelia</i> R. Br.	xlvii.	642
<i>Anagyris</i> Wall.			<i>Psoralea</i> L.	xlvii.	642
<i>Baptisia</i> Hook.			<i>Indigofera</i> L.	xlvii.	642
<i>Edwardia Miers</i>	xli.	567	<i>Swainsonia</i> Sal.	xlvii.	643
<i>Cyclopia</i> R. Br.	xli.	567	<i>Tessaria</i> Dec.	xlvii.	643
			<i>Sutherlandia</i> R. Br.	xlvii.	643
			<i>Coronilla</i> Neck.	xlvii.	643
			<i>Emerus</i> Mill.		
			<i>Hedysarum</i> L.	xlvii.	643
			<i>Hippocrepis</i> Jacq.	xlvii.	643
			<i>Adesmia</i> Dec.	xlvii.	643
			<i>Urtica</i> Desf.	xlvii.	643
			<i>Hesmodium Desf.</i>	xlvii.	643
			<i>Dicentra</i> Dec.	xlvii.	646
			<i>Zornia Pers.</i>		
			<i>Tavernera</i> Dec.	xlvii.	646
			<i>Lespedeza</i> Mx.	xlvii.	646
			<i>Flemingia</i> Roxb.	xlviii.	646
			<i>Ehretia</i> L.	xlviii.	646
			<i>Anthyllis</i> Lam.		
			<i>Albogi</i> Tourn.	xlviii.	646
			<i>Máuna</i> D. Don.		

	Contents	Text.
<i>Clíanthus Soland.</i>	xlviij.	646
<i>Dona G. and. D. Don.</i>		
Wistária Nutt.	xlviij.	647
<i>Glýcine L.</i>	xlviij.	649
<i>Thýrsínthus Elliott.</i>	xlviij.	649
<i>Kýralúhia Rafin.</i>	xlviij.	649
<i>A'pios Pursh.</i>	xlviij.	649
<i>Anonymus Walt.</i>	xlviij.	649
<i>Phascólóides H. Angl.</i>	xlviij.	649
<i>Lupinus L.</i>	xlviij.	649
<i>Dólichos L.</i>	xlviij.	649
<i>Pachýrhizus Dec.</i>	xlviij.	649
<i>Mucóna Wall.</i>	xlviij.	649
<i>Erythrina L.</i>	xlviij.	649
Gleditschia L.	xlviij.	650
Gymnócladus Lam.	xlviij.	656
<i>Guilandina L.</i>		
<i>Hyperanthera Vahl.</i>		
Céreis L.	xlviij.	657
<i>Siliquastrum Tourn.</i>		
<i>Cassalpinia Ait.</i>	xlix.	660
<i>Cádia L'Hérit.</i>	xlix.	660
<i>Zuccagnia Cav.</i>	xlix.	660
<i>Cerantonia L.</i>	xlix.	660
<i>Castanospermum Cunn.</i>	xlix.	660
<i>Cassia L.</i>	xlix.	660
<i>Chamaefistula Don's Mill.</i>	xlix.	661
<i>Schottia Ait.</i>	xlix.	661
<i>Derris G. Don.</i>	xlix.	661
<i>Pterocarpus Dec.</i>	xlix.	661
<i>Prosopis L.</i>	xlix.	661
<i>Lagonichium Bieb.</i>	xlix.	661
<i>Acacia Neck.</i>	xlix.	662
Rosaceæ Dec.	xlix.	670
Amýgdalus Tourn.	xlix.	673
<i>Amygdalophora Neck.</i>		
Pérsica Tourn.	l.	679
<i>Trichocarpus Neck.</i>		
Armeniaca Tourn.	l.	681
Prúnus Tourn.	l.	684
<i>Prunophora Neck.</i>		
Cérasus Juss.	li.	692
<i>Laurocérasus Tourn.</i>		
Púrshia Dec.	liij.	721
<i>Tigarea Pursh.</i>		
Kérria Dec.	liij.	722
<i>Córchorus Thunb.</i>		
Spiræa L.	liij.	722
Rúbus L.	liv.	733
Potentilla L.	lvi.	747
Ròsa L.	lvi.	748
<i>Rhodophora Neck.</i>		
Lòwea Lindl.	lxi.	812
Cratægus Lindl.	lxi.	813
Photinia Lindl.	lxix.	868
Cotonæster Med.	lxv.	869
Amelánchier Med.	lxv.	874
Méspilus Lindl.	lxv.	877
<i>Mespilophora Neck.</i>		
Pýrus Lindl.	lxv.	879
<i>Morus Tourn.</i>		
<i>Sórbus Tourn.</i>		
<i>Pyróphorum Neck.</i>		
<i>Apyróphorum Neck.</i>		
<i>Pyrúster Ray.</i>		
<i>Aria L'Obel.</i>		
<i>Arónia Pers.</i>		
Cydonia Tourn.	lxviii.	929
Chænoméles	lxviii.	931
<i>Raphidolepis Lindl.</i>	lxix.	952
<i>Eriobotrya Lindl.</i>	lxix.	953
<i>Kagoneckia R. et P.</i>	lxix.	954
<i>Margyricarpus R. et P.</i>	lxix.	954
<i>Eriopetrum Lam.</i>		
<i>Ancistrum Lam.</i>		
<i>Cercocarpus H. B. et K.</i>	lxix.	955
<i>Eretolonia Moc. et Sess.</i>		
<i>Acæna Vahl.</i>	lxix.	955
<i>Patérum L.</i>	lxix.	955
<i>Cliffortia L.</i>	lxix.	955

	Contents.	Text.
Calycanthaceæ Lindl.	lxix.	935
Calycánthus Lindl.	lxix.	936
<i>Buttnèria Du Ham.</i>		
<i>Beuryèria Ehrh.</i>		
<i>Bastèria Adans.</i>		
<i>Pompadoúra Buchoz.</i>		
Chimonánthus Lindl.	lxix.	937
<i>Merátia Nees.</i>		
Granatæcæ D. Don.	lxix.	939
Púnica Tourn.	lxix.	939
Onagræcæ Lindl.	lxix.	942
<i>Fúchia L.</i>	lxix.	944
<i>Skimra Mench.</i>		
<i>Nahúsia Schneegrogt.</i>		
Lythracæ Lindl.	lxx.	944
Hèimia	lxx.	945
Lagerstræ'mia L.	lxx.	945
Tamaricæcæ.	lxx.	946
Támarix Desv.	lxx.	947
Tamariscus All.	lxx.	949
<i>Myricaria Desv.</i>		
Philadelphæcæ.	lxx.	950
Philadélphus L.	lxx.	951
<i>Syringa Tourn.</i>		
Decumária L.	lxxi.	955
<i>Forsýthia Walt.</i>		
<i>Deutzia Thunb.</i>	lxxi.	956
Myrtæcæ.	lxxi.	956
<i>Tristánia R. Br.</i>	lxxi.	956
<i>Beaufórtia R. Br.</i>	lxxi.	957
<i>Calobádmanns R. Br.</i>	lxxi.	957
<i>Melaleuca L.</i>	lxxi.	957
<i>Eudesmia R. Br.</i>	lxxi.	958
<i>Eucalyptus L'Hérit.</i>	lxxi.	958
<i>Angophora Cav.</i>	lxxi.	960
<i>Callistemon Dec.</i>	lxxi.	960
<i>Metrosideros Gaertn.</i>	lxxi.	961
<i>Leptospermum Forst.</i>	lxxi.	961
<i>Bilbatia R. Br.</i>	lxxi.	961
<i>Fabricia Gaertn.</i>	lxxi.	961
<i>Be'ckia Andr.</i>	lxxi.	961
<i>Psidium L.</i>	lxxi.	961
<i>Myrtus L.</i>	lxxi.	961
<i>Chama-laucium Desf.</i>	lxxii.	964
<i>Cálythrix R. Br.</i>	lxxii.	964
<i>Darwinia Rudge.</i>	lxxii.	964
Passifloræcæ Lindl.	lxxii.	964
<i>Passifloræcæ Juss.</i>		
<i>Passiflora L.</i>	lxxii.	964
<i>Disemma Dec.</i>	lxxii.	965
<i>Tucaná Juss.</i>	lxxii.	965
Crassulæcæ Dec.	lxxii.	965
<i>Sédum L.</i>	lxxii.	965
<i>Sempervivum L.</i>	lxxii.	965
<i>Anacampseros Haw.</i>	lxxii.	965
Mesembryæcæ Lindl.	lxxii.	966
<i>Mesembryánthemum</i>	lxxii.	966
Nitrariæcæ Lindl.	lxxii.	966
Nitrària L.	lxxii.	966
Cactæcæ Lindl.	lxxii.	967
<i>Cácti Juss.</i>		
<i>Opúntia Mill.</i>	lxxii.	967
<i>Cactus L.</i>		

	Contents.	Text.		Contents.	Text.
<i>Grossulacæ</i> Dec.	lxxii.	967	<i>Leycesteria</i> Wall	lxxxii.	1060
<i>Ribes</i> L.	lxxii.	968	<i>Hamelia</i> Puer.		
<i>Grossularia</i> Tourn.			<i>Rubiaceæ</i> Juss.	lxxxii.	1061
<i>Chrysobothrya</i> Spach.			<i>Cephalanthus</i> L.	lxxxii.	1061
<i>Calibothrya</i> Spach.			Pinckneya Michx.	lxxxii.	1062
<i>Coreosma</i> Spach.			<i>Cinchona</i> Poir.	lxxxii.	1062
<i>Ribes</i> Spach.			<i>Serissa</i> Conium.	lxxxii.	1062
<i>Escalloniacæ</i> R. Br.	lxxiv.	992	<i>Dysoela</i> Lour.	lxxxii.	1062
<i>Itea</i> L.	lxxiv.	992	<i>Buchasia</i> L'Herit.	lxxxii.	1062
<i>Cedrela</i> Lour.			<i>Spermacoe</i> Desf.	lxxxii.	1062
<i>Dicondangia</i> Mx.			<i>Placania</i> Ait.	lxxxii.	1062
<i>Escallonia</i> Mutis	lxxiv.	993	<i>Bortlagia</i> Rehb.		
<i>Stereoxylon</i> R. et P.			<i>Phyllis</i> L.	lxxxii.	1062
<i>Saxifragæ</i> Dec.	lxxv.	994	<i>Anthospermum</i> L.	lxxxii.	1062
<i>Hydrangea</i> L.	lxxv.	994	<i>Ambraria</i> Walsh.	lxxxii.	1062
<i>Hortensia</i> Juss.			<i>Rubia</i> L.	lxxxii.	1062
<i>Umbellifera</i> Juss.	lxxv.	997	<i>Bouvardia</i> H. B. et K.	lxxxii.	1062
<i>Bupleurum</i> Tourn.	lxxv.	997	<i>Houstonia</i> L.	lxxxii.	1062
<i>Tenoria</i> Spreng.			<i>Manettia</i> Cham. et Schl.	lxxxii.	1062
<i>Buprestis</i> Spreng.			<i>Lobeliacæ</i> Juss.	lxxxii.	1063
<i>Seseli</i> Bauh.			<i>Trapa</i> G. Don	lxxxii.	1063
<i>Araliacæ</i> Juss.	lxxv.	998	<i>Lobelia</i> Forst.		
<i>Aralia</i> L.	lxxv.	998	<i>Campanulacæ</i> R. Br.	lxxxii.	1063
<i>Hedera</i> Swartz	lxxv.	999	<i>Mussaëia</i> Dumort.	lxxxii.	1063
<i>Hamamelidæ</i> Lindl.	lxxv.	1006	<i>Campanula</i> L.		
<i>Hamamelidæ</i> R. Br.			<i>Compositæ</i> Adans.	lxxxii.	1063
<i>Hamamelis</i> L.	lxxv.	1007	<i>Stachelia</i> Lessing	lxxxii.	1064
<i>Trilopus</i> Mith.			<i>Baccharis</i> R. Br.	lxxxii.	1065
<i>Fothergilla</i> L.	lxxvi.	100	<i>Iva</i> L.	lxxxii.	1066
<i>Cornucæ</i> Lindl.	lxxvi.	1009	<i>Santolina</i> L.	lxxxii.	1066
<i>Corneæ</i> Dec.			<i>Artemisia</i> Cass.	lxxxii.	1068
<i>Cornus</i> L.	lxxvi.	1009	<i>Abrótanus</i> Dod.		
<i>Virga</i> Matth.			<i>Absinthium</i> Lob.		
<i>Benthämia</i> Lindl.	lxxvii.	1019	<i>Helicrysum</i> Lessing	lxxxii.	1070
<i>Loranthæ</i> Lindl.	lxxvii.	1020	<i>Gnaphalium</i> L.		
<i>Loranthæ</i> Juss.			<i>Stachas</i> Dod.		
<i>Viscum</i> L.	lxxvii.	1021	<i>Astelma</i> R. Br.	lxxxii.	1070
<i>Aucuba</i> Thunb.	lxxvii.	1026	<i>Cineraria</i> Lessing	lxxxii.	1070
<i>Eubasis</i> Sal.			<i>Jacobæa</i> Bonp.		
<i>Loranthus</i> L.	lxxvii.	1026	<i>Agatha</i> Cas.		
<i>Caprifoliacæ</i> Richd.	lxxvii.	1027	<i>Carlowizia</i> Mench.	lxxxii.	1072
<i>Sambucus</i> Tourn.	lxxvii.	1027	<i>Onobryma</i> Lk.		
<i>Phytolima</i> Lour.			<i>Didelta</i> L'Herit.	lxxxii.	1072
<i>Fibrium</i> L.	lxxvii.	1032	<i>Bercheya</i> Willd.	lxxxii.	1072
<i>Opytus</i> Tourn.			<i>Cullamia</i> R. Br.	lxxxii.	1072
<i>Tinus</i> Tourn.			<i>Othónna</i> L.	lxxxii.	1072
<i>Diervilla</i>	lxxix.	1042	<i>Osteospermum</i> L.	lxxxii.	1072
<i>Wiegela</i> Thunb.			<i>Calendula</i> Vent.	lxxxii.	1072
<i>Wiegelia</i> Pers.			<i>Mutisia</i> D. Don	lxxxii.	1072
<i>Lonicera</i> Desf.	lxxix.	1042	<i>Dahlia</i> Cav.	lxxxii.	1075
<i>Caprifolium</i> Juss.			<i>Atalaphus</i> D. Don	lxxxii.	1075
<i>Xylosteum</i> Juss.			<i>Prenanthes</i> L.		
<i>Chamaecerasus</i> Tourn.			<i>Sonchus</i> Jacq.	lxxxii.	1075
<i>Periclymenum</i> Tourn.			<i>Vernonia</i> Hook.	lxxxii.	1075
<i>Xylosteon</i> Tourn.			<i>Aster</i> Lab.	lxxxii.	1075
<i>Sintonia</i> Swt.			<i>Houstonia</i> Caley.		
<i>Iska</i> Borek.			<i>Chrysocoma</i> L.	lxxxii.	1075
<i>Isca</i> Mench.			<i>Brachyactina</i> Swt.	lxxxii.	1075
<i>Symphoricarpos</i> Dill.	lxxxii.	1058	<i>Conyza</i> L.	lxxxii.	1075
<i>Symphoricarpa</i> Neck.			<i>Podanthus</i> Lindl.	lxxxii.	1075
<i>Symphoria</i> Pers.			<i>Culectium</i> Spreng.	lxxxii.	1075
<i>Anisanthus</i> Wild.			<i>Cacalia</i> L.	lxxxii.	1075
<i>Lonicera</i> L.			<i>Franzeria</i> Willd.	lxxxii.	1075
			<i>Gleditsia</i> Thunb.	lxxxii.	1075
			<i>Pyrethrum</i> Willd.	lxxxii.	1075
			<i>Imelia</i> Cass.	lxxxii.	1075
			<i>Athanasia</i> L.	lxxxii.	1075
			<i>Balsamita</i> Desf.	lxxxii.	1075
			<i>Picntia</i> Willd.	lxxxii.	1075
			<i>Tanacetum</i> L'Herit.		
			<i>Eriocapillus</i> L.	lxxxii.	1075
			<i>Senecio</i> L.	lxxxii.	1075
			<i>Tarcondanthus</i> Lam.	lxxxii.	1075
			<i>Eriocoma</i> D. Don	lxxxii.	1075
			<i>Epaeridæ</i> Lindl.	lxxxiii.	1075
			<i>Epaeridæ</i> Br.		
			<i>Styphelia</i> R. Br.	lxxxiii.	1075
			<i>Sienanthera</i> R. Br.	lxxxiii.	1075
			<i>Cyathodes</i> Lab.	lxxxiii.	1075
			<i>Lissanthe</i> R. Br.	lxxxiii.	1075
			<i>Leucopogon</i> R. Br.	lxxxiii.	1075
			<i>Monotoca</i> R. Br.	lxxxiii.	1075
			<i>Trochocarpus</i> R. Br.	lxxxiii.	1075
			<i>Pyrethrum</i> Forst.	lxxxiii.	1075
			<i>Lysanema</i> R. Br.	lxxxiii.	1075
			<i>Androselia</i> R. Br.	lxxxiii.	1075

	Contents.	Text.		Contents.	Text.
Sprengelia Sm.	lxxxiii.	1075	Virèya	xciv.	1173
Sphenotoma Sm.	lxxxiii.	1075	Befaria H. et B.	xciv.	1173
Dracophyllum R. Br.			Hymenánthes Blom.	xciv.	1173
			Gavliu-acéia H. B. et K.	xciv.	1173
Ericàcæ Juss.	lxxxiii.	1076	Thibaudia H. B. et K.	xciv.	1173
Erica D. Don	lxxxiii.	1079	Cavendishia Lindl.	xciv.	1173
Gypsocállis Sal.	lxxxiii.	1082	Agapetes D. Don	xciv.	1173
Calluna Sal.	lxxxv.	1084	Ceratostema R. et P.	xciv.	1173
Callista D. Don	lxxxiv.	1089	Symplocàcæ D. Don.	xcv.	1186
Ceramia D. Don	lxxxiv.	1089	Symplocos L.	xciv.	1186
Dasyánthes D. Don	lxxxiv.	1089	Styràcæ Rich.	xciv.	1187
Désmia D. Don	lxxxiv.	1089	Styrax L.	xciv.	1187
Eurylepis D. Don	lxxxiv.	1090	Halesiàcæ D. Don		
Páchysa D. Don	lxxxiv.	1091	and Link.	xciv.	1189
Syringodea D. Don	lxxxiv.	1091	Halèsia Ellis	xciv.	1189
Bætria L.	lxxxiv.	1091	Sapotàcæ Lindl.	xciv.	1191
Ectasia D. Don	lxxxiv.	1092	Sapotæ R. Br.		
Eurystégia D. Don	lxxxiv.	1095	Argània Rœm. et Schult.	xciv.	1191
Lophándra D. Don	lxxxiv.	1095	Elæodéndron Retz.		
Lámprotis D. Don	lxxxiv.	1095	Sideróxyton L.		
Andróméda L.	lxxxv.	1105	Bumélia Swartz.	xcv.	1192
Potifolia Buxb.			Achéras L.		
Cassiøpe D. Don	lxxxvi.	1107	Chrysophýllum Aubl.		
Cassándra D. Don	lxxxvi.	1108	Ebenàcæ Dec.	xcv.	1194
Zenòbia D. Don	lxxxvi.	1108	Diospýros L.	xcv.	1194
Lyònia Nutt.	lxxxvi.	1109	Eúbenus Comm.		
Leucóthøe D. Don	lxxxvi.	1113	Guaiaçaina Tourn.		
Pieris D. Don	lxxxvii.	1114	Pseudólötus Matth.		
Phyllódoce Sal.	lxxxvii.	1115	Embryópterus L.	xcv.	1197
Dabæ'cia D. Don	lxxxvii.	1116	Oleàcæ Lindl.	xcv.	1198
Menzièsia Sm.			Oleína Hoff.		
Arbutus Camer.	lxxxvii.	1117	Ligústrum Tourn.	xcv.	1198
Arctostáphylos Adans.	lxxxviii.	1123	Phillyrea Tourn.	xcv.	1203
Pernéttya Gaud.	lxxxviii.	1124	Chionánthus L.	xcvi.	1205
Gaulthéria L.	lxxxviii.	1125	O'lea L.	xcvi.	1207
Epigæ'ia L.	lxxxviii.	1126	Syringa L.	xcvi.	1208
Mémécylum Mx.			Fontanèsia Lab.	xcvii.	1213
Phaleroçárpus G. Don	lxxxviii.	1127	Fráxinus Tourn.	xcvii.	1213
Cléthra L.	lxxxviii.	1127	Ornus Pers.	xcviii.	1241
Rhododéndron L.	lxxxix.	1130	Jasminàcæ Lindl.	xcix.	1248
Azàlea L.			Jasminæ Juss.		
Rhòdra L.			Jasminum Forsk.	xcix.	1248
Chamærhododéndros Tourn.			Mongórium Lam.		
Kálmia L.	xcii.	1151	Apocynàcæ Lindl.	xcix.	1254
Menzièsia Sm.	xcii.	1152	Apocynæ Juss.		
Azàlea D. Don	xcii.	1153	Vinca L.	xcix.	1254
Loisèleuria Desf.			Pervinca Tourn.		
Chamælèdon Lk.			Gelæmium Mx.	xcix.	1256
Leiophýllum Pers.	xcii.	1154	Nerium L.	xcix.	1256
Ammýrsinc Pursh.					
Fischera Swartz.					
Lèdum L.	xcii.	1155			
Vaccinium L.	xcii.	1156			
Vitis idæ'a	xcii.	1156			
Oxycóccus Pers.	xciii.	1168			
Bryánthus Gmel.	xciv.	1171			
Enkiánthus Lour.	xciv.	1172			
Meladòra Sal.					

VOL. III.

Asclepiàdæ Lindl.	xcix.	1257	Cobæàcæ D. Don.	c.	1264
Asclepiðæ R. Br.			Cobæ'a Cav.	c.	1264
Períploca L.	xcix.	1257	Convulvulàcæ R. Br.	c.	1264
Bignoniàcæ R. Br.	c.	1258	Convólulus L.	c.	1264
Bignònia Tourn.	c.	1258	Boraginàcæ Lindl.	c.	1265
Tècoma Juss.	c.	1259	Boraginæ Juss.		
Catálpa Juss.	c.	1261	Lithospermum L.	c.	1265
Eccremocárpus Humb.	c.	1263	Echium L.	c.	1265
Calámpelis D. Don	c.	1263	Heliotrópium L.	c.	1265

	Contents.	Text.		Contents.	Text.
<i>Cordiàcæ R. Br.</i>	c.	1265	<i>Chenopodiàcæ Lindl.</i>	cii.	1287
<i>Ehretia Roxb.</i>	c.	1265	<i>Chenopodeæ Vent.</i>		
<i>Solanàcæ Lindl.</i>	c.	1266	<i>Chenopodium L.</i>	cii.	1288
<i>Solanææ Juss.</i>			<i>Salsola L.</i>		
<i>Solanum L.</i>	c.	1266	<i>Suaeda Forsk.</i>		
<i>Melongèna Tourn.</i>			<i>Atriplex L.</i>	ciii.	1289
<i>Pseudo-Capsicum Mœnch.</i>			<i>Halmis Bauh.</i>		
<i>Nyctèrium Vent.</i>			<i>Diôtis Schreb.</i>	ciii.	1290
<i>Aquária Jacq.</i>			<i>Ceratoides Tourn.</i>		
<i>Lýcium L.</i>	ci.	1269	<i>Aryris L.</i>		
<i>Jasminoides Niss.</i>			<i>Ceratospèrum Persk.</i>		
<i>Grabówskiä Schlecht</i>	ci.	1273	<i>Achyranthes Forsk.</i>		
<i>Nicotiana Grab.</i>	ei.	1274	<i>Krascheninnikovia Guild.</i>		
<i>Brugmansia R. et P.</i>	ei.	1274	<i>Urtica Roy</i>		
<i>Solantra L.</i>	ei.	1274	<i>Andræsis L.</i>	clii.	1291
<i>Cestrum L.</i>	ei.	1274	<i>Kuchia Schr.</i>	ciii.	1291
<i>Vestia Walld.</i>	ei.	1274	<i>Boson L.</i>	ciii.	1291
<i>Cantua Juss.</i>			<i>Camphorosma Schk.</i>	ciii.	1291
<i>Scrophulariàcæ Lindl.</i>	ci.	1276	<i>Polygonàcæ Lindl.</i>	ciii.	1292
<i>Scrophularinæ R. Br.</i>			<i>Tragopyrum Bieb.</i>	ciii.	1292
<i>Búddlea L.</i>	ci.	1276	<i>Polygonum L.</i>		
<i>Halleria L.</i>	ci.	1277	<i>Atraphaxis L.</i>	ciii.	1294
<i>Maurandia Jacq.</i>	ci.	1277	<i>Calligonum L.</i>	ciii.	1295
<i>Mianulus Willd.</i>	ci.	1277	<i>Pallasia L.</i>		
<i>Anthocercis R. Br.</i>	ci.	1277	<i>Pterococcus Pall.</i>		
<i>Calceolaria R. Br.</i>	ci.	1277	<i>Brunnichia Giertn.</i>	ciii.	1296
<i>Feronia Ait.</i>	ci.	1277	<i>Rumex L.</i>	ciii.	1296
<i>Celsia Jacq.</i>	ei.	1277	<i>Polygonum R. Br.</i>	ciii.	1296
<i>Capraria L.</i>	ei.	1277			
<i>Fragaria Bot. Mag.</i>			<i>Lauràcæ Lindl.</i>	ciii.	1296
<i>Alonsoa R. et P.</i>	ci.	1277	<i>Laurinæ Vent.</i>		
<i>Angelonia H. B. et K.</i>	ci.	1277	<i>Laurus Plin.</i>	ciii.	1296
<i>Lophospèrum Don</i>	ci.	1277	<i>Sassafras Esenb.</i>		
<i>Rhodochiton Zucc.</i>	ei.	1277	<i>Benzoin Esenb.</i>		
<i>Nycteria D. Don</i>	ei.	1277	<i>Cinnamomum R. Br.</i>	civ.	1305
			<i>Percua Spr.</i>		
<i>Labiàtæ Juss.</i>	ci.	1278	<i>Proteàcæ Juss.</i>	civ.	1306
<i>Satureja L.</i>	ci.	1278	<i>Banksia R. Br.</i>	civ.	1306
<i>Thymus L.</i>	ci.	1278	<i>Grevillea Gunn.</i>	civ.	1306
<i>Hyssopus L.</i>	cii.	1278	<i>Hakea R. Br.</i>	civ.	1306
<i>Teucrium Schreb.</i>	cii.	1279	<i>Thymelàcæ Juss.</i>	civ.	1306
<i>Phlomis L.</i>	cii.	1279	<i>Dáphne L.</i>	civ.	1307
<i>Rosmarinus L.</i>	cii.	1279	<i>Thymelæa Tourn.</i>		
<i>Stachys L.</i>	cii.	1281	<i>Daphnoides Gesn.</i>		
<i>Lavandula L.</i>	cii.	1281	<i>Lauræola Roy.</i>		
<i>Acyros Lk.</i>	cii.	1282	<i>Sanamunda Bauh.</i>		
<i>Gardoquia Hook.</i>	cii.	1282	<i>Chamaedaphnoides Alpin.</i>		
<i>Westringia Sm.</i>	cii.	1282	<i>Dírea L.</i>	cv.	1314
<i>Sálvia L.</i>	cii.	1282	<i>Gnidia L.</i>	cv.	1315
<i>Audibertia Benth.</i>	cii.	1283	<i>Passerina L.</i>	cv.	1315
<i>Plectranthus L'Herit</i>	cii.	1283	<i>Pimelea</i>	cv.	1315
<i>Sideritis Ait.</i>	cii.	1283	<i>Santalàcæ Brown.</i>	cv.	1316
<i>Leonotis R. Br.</i>	cii.	1283	<i>Nýssa L.</i>	cv.	1315
<i>Sphæcele Benth.</i>	cii.	1283	<i>Osyris L.</i>	cvi.	1320
<i>Dracæcephalum Com.</i>	cii.	1283	<i>Casia Camer.</i>		
<i>Præsum L.</i>	cii.	1283	<i>Elæagnàcæ Lindl.</i>	cvi.	1321
<i>Prostanthera Lab.</i>	cii.	1283	<i>Elæagnus Tourn.</i>	cvi.	1321
<i>Verbenàcæ Juss.</i>	cii.	1285	<i>Hippophæc L.</i>	cvi.	1324
<i>Vitex L.</i>	cii.	1285	<i>Rhamnoides Tourn.</i>		
<i>Cerodéndron R. Br.</i>	cii.	1286	<i>Shepherdia Nutt.</i>	cvi.	1327
<i>Volkammeria L.</i>	cii.	1286	<i>Aristolochiàcæ R. Br.</i>	cvi.	1328
<i>Duranta Hook.</i>	cii.	1286	<i>Aristolochia L.</i>	cvi.	1328
<i>Aloysia Or.</i>	cii.	1286	<i>Euphorbiàcæ Juss.</i>	cvi.	1330
<i>Verbena L'Herit.</i>			<i>Euphorbia</i>	cvi.	1331
<i>Lippia Kunth.</i>			<i>Tithymalus</i>		
<i>Myopórinæ R. Br.</i>	cii.	1287	<i>Stillingia Garden</i>	cvii.	1332
<i>Myoporum</i>	cii.	1287	<i>Búxus Tourn.</i>	cvii.	1332
<i>Globulariàcæ Lindl.</i>	cii.	1287	<i>Plagiântus Vent.</i>	cvii.	1341
<i>Globulária Dec.</i>			<i>Cluytia Bot. Mag.</i>	cvii.	1341
<i>Globulária L.</i>	cii.	1287			
<i>Plumbaginàcæ R. Br.</i>	cii.	1287			
<i>Statice L.</i>	cii.	1287			
<i>Plumbago L.</i>	cii.	1287			

	Contents.	Text.
<i>Urticacæ Lindl.</i>	cvii.	1342
<i>Urticæ Juss.</i>		
<i>Morus Tourn.</i> -	cvii.	1343
<i>Broussonètia Vent.</i> -	cviii.	1361
<i>Papÿrus Encyc. Bot.</i>		
<i>Maclura Nutt.</i> -	cviii.	1362
<i>Tôzilou Raf.</i>		
<i>Ficus Tourn.</i> -	cviii.	1365
<i>Bôrya Willd.</i> -	cviii.	1370
<i>Adelia Michx.</i>		
<i>Bigelôvia Sm.</i>		
<i>Ulmacæ Mirb.</i>	cviii.	1371
<i>Ulmus L.</i> -	cviii.	1373
<i>Plânera Gmel.</i> -	cx.	1409
<i>Celtis Tourn.</i> -	cx.	1413
<i>Juglandacæ Lindl.</i>	cx.	1420
<i>Juglandæ Juss.</i>		
<i>Juglans L.</i> -	cx.	1421
<i>Carya Nutt.</i> -	cx.	1421
<i>Hicôrius Raf.</i>		

	Contents.	Text.
<i>Pterocarya Kunth</i> -	cx.	1451
<i>Salicacæ Lindl.</i>	cx.	1453
<i>Saliciflora Rich.</i>		
<i>Salix L.</i> -	cx.	1453
<i>Pôpulus Tourn.</i> -	cxxi.	1636
<i>Betulacæ Dum.</i>	cxxiii.	1677
<i>Alnus Tourn.</i> -	cxxiii.	1677
<i>Bétula Tourn.</i> -	cxxiii.	1690
<i>Corylacæ Lindl.</i>	cxxiv.	1715
<i>Cupuliferæ Rich.</i>		
<i>Quercus L.</i> -	cxxiv.	1717
<i>Flex Tourn.</i>		
<i>Siber Tourn.</i>		
<i>Fâgus L.</i> -	cxxix.	1949
<i>Castanea Tourn.</i>		
<i>Castanea Tourn.</i> -	cxxx.	1983
<i>Cârpinus L.</i> -	cxxx.	2004
<i>O'strya Willd.</i> -	cxxx.	2015
<i>Cârpinus L.</i>		
<i>Côrylus L.</i> -	cxxx.	2016

VOL. IV.

<i>Garryacæ Lindl.</i>	cxxx.	2031
<i>Garrya Doug.</i> -	cxxx.	2031
<i>Platanacæ Lindl.</i>	cxxx.	2032
<i>Platanus L.</i> -	cxxx.	2033
<i>Balsamacæ Lindl.</i>	cxxxii.	2048
<i>Liquidambar L.</i> -	cxxxii.	2049
<i>Altingia Nov.</i>		
<i>Myricacæ Lindl.</i>	cxxxii.	2055
<i>Myrica L.</i> -	cxxxii.	2055
<i>Comptonia Banks.</i>	cxxxii.	2059
<i>Liquidambar L.</i>		
<i>Myrica L.</i>		
<i>Casuaracæ. R. Br.</i>	cxxxii.	2060
<i>Casuarina Ait.</i> -	cxxxii.	2060
<i>Gnetacæ Lindl.</i>	cxxxiii.	2062
<i>E'phedra L.</i> -	cxxxiii.	2062
<i>Taxacæ Lindl.</i>	cxxxiii.	2065
<i>Taxus L.</i> -	cxxxiii.	2066
<i>Salisbûria Sm.</i> -	cxxxiii.	2094
<i>Ginkgo Kæmpf.</i>		
<i>Podocarpus L'Herit.</i> -	cxxxiii.	2100
<i>Dacrydium Soland.</i> -	cxxxiii.	2101
<i>Phylloctadus Rich.</i> -	cxxxiii.	2102
<i>Coniferæ Juss.</i>	cxxxiii.	2103
<i>Abietinæ.</i>	cxxxiii.	2106
<i>Pinus L.</i> -	cxxxiii.	2152
<i>Abies D. Don</i> -	cxxxvii.	2293
<i>Picea Lk.</i>		
<i>Picea D. Don</i> -	cxxxviii.	2329
<i>Abies Lk.</i>		
<i>Lârix Tourn.</i> -	cxxxix.	2350
<i>Cèdrus Barrel.</i> -	cxl.	2402
<i>Araucaria R. et P.</i> -	cxl.	2432
<i>Eutassa Sal.</i>		
<i>Colymbèa Sal.</i>		
<i>Dombèya Lam.</i>		

<i>Cunninghâmia R. Br.</i> -	cxl.	2445
<i>Dammara Rumph.</i> -	cxl.	2447
<i>§ Cupressinæ.</i>	cxl.	2453
<i>Thûja L.</i> -	cxl.	2454
<i>Câllitris Vent.</i> -	cxli.	2462
<i>Fresnèlia Mirb.</i>		
<i>Cupressus</i> -	cxli.	2464
<i>Taxodium Rich.</i> -	cxlii.	2480
<i>Schubèrtia Mirb.</i>		
<i>Condylôcirpus Sal.</i>		
<i>Juniperus L.</i> -	cxlii.	2487
<i>Empetracæ Nutt.</i>	cxliii.	2506
<i>Empetrum L.</i> -	cxliii.	2506
<i>Corèma D. Don</i> -	cxliii.	2508
<i>Ceratiola Mx.</i> -	cxliii.	2508
<i>Smilacæ Brown.</i>	cxliii.	2509
<i>Smilax L.</i> -	cxliii.	2510
<i>Liliacæ Dec.</i>	cxliv.	2215
<i>Aspâragus L.</i> -	cxliv.	2516
<i>Ruscus L.</i> -	cxliv.	2517
<i>Yûca L.</i> -	cxlv.	2521
<i>Amaryllidacæ Lindl.</i>		
<i>Fourcroya Vent.</i> -	cxlv.	2527
<i>Littæa Brig.</i> -	cxlv.	2528
<i>Agave Ker.</i>		
<i>Bonapârtèa Haw.</i>		
<i>Agave L.</i> -	cxlv.	2529
<i>Phôrmium Thun.</i> -	cxlv.	2529
<i>Pâlmæ Juss.</i>		
<i>Chamæ'rops L.</i> -	cxlv.	2530
<i>Graminæ R. Br.</i>		
<i>Bambûsa L.</i> -	cxlv.	2532
<i>Arûndo Beauv.</i> -	cxlv.	2532

EPITOME OF THE PLATES.

VOL. V.

Magnoliaceæ.

Magnòlia, 9 sp. and 2 var.;
12 plates.

Liriodéndron, 1 sp.; 2 pl.

Tiliaceæ.

Tilia, 8 sp. and 1 var.; 10 pl.

Aceraceæ.

Acer, 15 sp. and 1 var.; 21 pl.

Negúndo, 1 sp. and 1 var.; 2 pl.

Æsculaceæ.

Æ'sculus, 2 sp.; 3 pl.

Pàvia, 3 sp. and 1 var.; 6 pl.

Sapindaceæ.

Kölreutèria, 1 sp.; 1 pl.

Xanthoxylaceæ.

Xanthóxyllum, 1 sp.; 1 pl.

Ptèlea, 1 sp.; 1 pl.

Ailántus, 1 sp.; 2 pl.

Celastraceæ.

Euónymus, 2 sp.; 2 pl.

Aquifoliaceæ.

Flex, 2 sp.; 4 pl.

Rhamnaceæ.

Paliùrus, 1 sp.; 2 pl.

Rhámnus, 4 sp.; 4 pl.

Homalinaceæ.

Aristotèlia, 1 sp.; 1 pl.

Leguminosæ.

Sophòra, 1 sp. and 1 var.; 3 pl.

Virgíliã, 1 sp.; 1 pl.

Cýtisus, 2 sp. and 2 var.; 4 pl.

Ròbinia, 3 sp. and 2 var.; 6 pl.

Caragàna, 1 sp.; 1 pl.

Gleditschia, 7 sp.; 9 pl.

Gymnócladus, 1 sp.; 2 pl.

Céreis, 2 sp.; 3 pl.

Acácia, 1 sp.; 1 pl.

VOL. VI.

Rosaceæ.

Amýgdalus, 1 sp.; 1 pl.

Pérsica, 1 sp.; 1 pl.

Armeniaca, 2 sp.; 2 pl.

Prúnus, 3 sp.; 3 pl.

Cérasus, 7 sp.; 9 pl.

Cratægus, 21 spec. and 13
var.; 36 pl.

Photínia, 1 sp.; 1 pl.

Cotoneáster, 4 sp.; 4 pl.

Amelánchier, 1 sp.; 2 pl.

Méspilus, 2 sp.; 2 pl.

Pýrus, 18 sp. and 2 var.;
22 pl.

Cydònia, 2 sp.; 2 pl.

Myrtaceæ.

Eucalyptus, 1 sp.; 1 pl.

Cornaceæ.

Córnu, 1 sp.; 1 pl.

Caprifoliaceæ.

Sambucus, 1 sp.; 1 pl.

Vibúrnum, 1 sp.; 1 pl.

Ericaceæ.

Arbutus, 2 sp.; 2 pl.

Halesiaceæ.

Halèsia, 1 sp.; 2 pl.

Ebenaceæ.

Diospýros, 2 sp.; 4 pl.

Oleaceæ.

Fráxinus, 8 sp. and 1 var.;
11 pl.

Órnus, 1 sp.; 2 pl.

VOL. VII.

Bignoniaceæ.

Catálpa, 1 sp.; 2 pl.

Lauraceæ.

Laúrus, 2 sp.; 3 pl.

Santalaceæ.

Nýssa, 1 sp.; 1 pl.

Elæagnaceæ.

Elæágnus, 1 sp.; 1 pl.

Hippóphæe, 1 sp.; 1 pl.

Urticaceæ.

Mòrus, 3 sp.; 5 pl.

Broussonètia, 1 sp.; 1 pl.

Ficus, 1 sp.; 1 pl.

Ulmaceæ.

Ulmus, 7 sp. and 5 var.; 15 pl.

Plánera, 1 sp.; 2 pl.

Céltis, 2 sp.; 3 pl.

Juglandaceæ.

Juglans, 3 sp.; 6 pl.

Càrya, 1 sp.; 1 pl.

Pterocàrya, 1 sp.; 1 pl.

Salicaceæ.

Sàlix, 4 sp. and 1 var.; 6 pl.

Pópulus, 10 sp. and 1 var.;
17 pl.

Betulaceæ.

A'lnus, 2 sp. and 1 var.; 4 pl.

Bétula, 4 sp.; 5 pl.

Corylaceæ, or Cupuliferaæ.

Quérens, 6 sp. and 4 var.;
20 pl.

VOL. VIII.

Corylaceæ, or Cupuliferaæ.

Quérens, 13 sp. and 6 var.;
24 pl.

Fágu, 1 sp. and 2 var.; 4 pl.

Castànea, 1 sp.; 2 pl.

Cárpinus, 1 sp.; 2 pl.

O'strya, 2 sp.; 3 pl.

Córylus, 1 sp.; 2 pl.

Platanaceæ.

Plátanus, 2 sp. and 2 var.; 6 pl.

Balsamaceæ.

Liquidámbar, 1 sp.; 2 pl.

Taxaceæ.

Táxus, 1 sp. and 1 var.; 4 pl.

Salisbùria, 1 sp.; 2 pl.

Coniferaæ, § Abiétinaæ.

Pínus, 11 sp. and 1 var.; 17 pl.

Abies, 5 sp. and 1 var.; 10 pl.

Picea, 2 sp.; 4 pl.

Làrix, 2 sp.; 4 pl.

Cèdrus, 1 sp. and 1 var.; 5 pl.

Abiétinaæ, § Cuprèssinaæ.

Thùja, 2 sp.; 2 pl.

Cuprèssus, 2 sp.; 1 pl.

Taxòdium, 1 sp.; 2 pl.

Juniperus, 3 sp.; 5 pl.

CONTENTS.

INTRODUCTION.

Uses of Trees, 1. Ornaments, 2. Interchange of Trees, 3. Arrangement of the Work, 4.

PART I.

GENERAL OUTLINE OF THE HISTORY AND GEOGRAPHY OF THE TREES AND SHRUBS OF TEMPERATE CLIMATES.

CHAP. I.

Of the Knowledge of Trees and Shrubs which existed among the Nations of Antiquity, 15.

Sacred Trees, 15. Trees of the Egyptians and Persians, 16. Enumeration of the Species of Trees indigenous to Greece as given by Theophrastus, 17. Trees of the Romans, 19.

CHAP. II.

Of the History and Geography of the Trees and Shrubs now in the British Islands, 20.

SECT. I. Of the Native Trees and Shrubs of the British Islands, 20.

Trees found by Cæsar, 20. Trees, the indigeneness of which is considered doubtful, 21. Beech, Sweet Chestnut, English Elm, &c., 22. The Box, 25. Trees known to our Saxon Ancestors, 26. Enumeration of the Species and Varieties of Ligneous Plants included in the British Flora, from Smith and Hooker, 27. Classification of the Ligneous British Flora, according to height, habit, and popular character, 30.

SECT. II. Of the Foreign Trees and Shrubs introduced into the British Isles, 31.

SUBJECT. 1. Of the Foreign Trees and Shrubs introduced into Britain by the Romans, and during the Middle Ages, till the end of the 15th Century, 32.

Trees and Shrubs introduced by the Romans 32.; by the Occupiers of Monasteries and other religious Houses, and by the Crusaders, 33. Introduction of the cultivated Rose, 33. York and Lancaster Rose, 34. Foreign Trees and Shrubs of Scotland and Ireland, 34.

SUBJECT. 2. Of the Foreign Trees and Shrubs introduced into Britain in the 16th Century, 35.

Trees and Shrubs introduced in the time of Henry VIII., 35. Fitzherbert, Turner, Goodge, L'Obel, Grindal, Hampden, Gerard, &c., 35. Trees and Shrubs in the Earl of Essex's Garden at Barn Elms, 36. Corbett's Nursery at Twickenham, 36. Raleigh's Park in Dorsetshire, 36. Enumeration of the Species of Trees and Shrubs introduced into England during the 16th Century, by Hugh Morgan, Gray, L'Obel, Grindal, Gerard, &c., 36. 39.

SUBJECT. 3. Of the Foreign Trees and Shrubs introduced into Britain in the 17th Century, 40.

Tradesant's Garden at Lambeth, Bishop Compton's Garden at Fulham, 41. Ray's Historia Plantarum, 41. Dr. Mitchel, Sir Wm. Watson, Mr. Bentick, Dr. Plukenet, Sir Hans Sloane, Furber of Kensington, Gray of Fulham, &c., 41. Enumeration of Species of Foreign Trees and Shrubs found in the Episcopal Garden at Fulham in 1751, 42.; in 1793, and again in 1809, by Lyson, 43.; in 1835, 44. Banister, Evelyn, Hermann, Gibson, Sir Stephen Fox, London and Wise, Cooke, Lukar, London and Field, Dr. Uvedale, Miller, &c., 42. 46. Enumeration of the Trees and Shrubs introduced into England during the 17th Century, according to the Hortus Kewensis, 49. Harefield Park, 52. Edinburgh Botanic Garden, Oxford Botanic Garden, 53.

SUBJECT. 4. Of the Foreign Trees and Shrubs introduced into Britain during the 18th Century, 54.

Trees and Shrubs introduced at Whitton, Goodwood, Syon, &c., 54. Miller, Bradley, Switzer, Linnæus, Peter Collinson, the London Nurserymen, &c., 54. Ridgway House Garden, 54. London Gardens in 1712, 55. Lord Petre's Collection, 55.

John Clark, a Huteher at Barnes, famous for raising Cedars from Seed, &c., 55. The Great Cedar at Hendon Place, the Cedars at Whitton, 55. Fine Gardens noticed by Collinson on a blank Leaf of his Copy of Miller's Dictionary, 56. Enumeration of Trees and Shrubs, with their Dimensions, found in the Garden at Mill Hill (which formerly belonged to Collinson) in January, 1835, 56. Peterborough House, Whitton, 57. Twickenham Botanic Garden, Thornodon Hall, Essex, 58. Cedar House, Cashobury, 59. Catalogue of Trees and Shrubs propagated for Sale in the London Nurseries in 1730, 60. Celebrated Amateurs and Gardeners of that Time: Dr. Compton, S. Reynardson, Esq., Dr. Uvedale, Duchess of Beaufort, Earl of Pembroke, 61. Dubois, 62. List of Species and Varieties of Trees and Shrubs given in the Catalogue of the Society of Gardeners, arranged according to the Natural System, 64. Catesby's Travels and Publications, 68. Plan for importing Acorns and other Seeds of American Trees, 70. Dr. Garden of Charlestown, 70. Planting of Stowe, Blenheim, Corby Castle, Pain's Hall, &c., 70, 71. Upton House, and its fine Specimens of Foreign Trees and Shrubs, 71. Trees and Shrubs at Purser's Cross in 1756, their comparative Dimensions taken at different periods between 1793 and 1835, 72. Trees and Shrubs at Syon, at Croome, and at various other Places, 73. Collection in the Chelsea Botanic Garden, 74. Botanic Garden and Arboretum at Kew, Sherard's Botanic Garden at Eltham, Twickenham Botanic Garden, 75. Brompton Park Nursery, 76. Kensington Nursery, Hoxton Nursery, Mile End Nursery, 77. Putney Nursery, Hammersmith Nursery, 78. Hackney Botanic Garden and Nursery, and those of Brentford, Lewisham, New Cross, &c., 79. Botanical and Horticultural Authors and Artists of the 18th Century, 79. Enumeration of the Trees and Shrubs introduced during each Decade of the 18th Century, 80-84. Biographical Notices of John and William Bartram, 85. Trees and Shrubs introduced into Scotland during the 18th Century, 86. Dr. Walker's Exertions, 87. Mr. Sang's *Planter's Calendar*, 89. Lists of the Species of Foreign Trees and Shrubs introduced into Scotland during the 17th and 18th Centuries according to Dr. Walker, 90, 91. Taymouth, Inverary Castle, 91. Prestonfield, 92. Hamilton Palace, 92. Panmure, New Posso, 93. Holyrood House, 94. Bargally, the Property of Andrew Heron, one of the greatest Botanists and Horticulturists that has ever appeared in Scotland, 95, 99. Dunkeld, 99. New Hailes, Arbigland, Loudon Castle, Dalmahoy, 101. Hopetoun House, Carmichael, Tynningham, 102. Scottish Nurseries, 104. Indigenous Trees of Ireland, 105. Irish Yew in a Garden at Mayland near Antrim, 106. Antrim Castle, 106. Mitchelstown, 107. Moira, and other Places celebrated for Foreign Trees; Lords Oriel and Clanbrasil introduced by far the greater number of Trees into Ireland in the 18th Century, 108. Dimensions of many fine Specimens of Trees and Shrubs at Oriel Temple, 109.; at Antrim Castle, 109.; at Tollymore Park, at Dundalk, 110.; at Cypress Grove, at Moira, and at Cranmore, 111.; at Castle Ward, 112.; at Howth Castle, Charleville Forest, and Shelton Abley, 113.; at Castle Freke, Florence Court, Killrudey House, Mount Anville Hill, Castletown, 114. The finest Cedars in Ireland, 114. The largest Oak in Ireland, 115. Foreign Trees and Shrubs at Hillsborough, 115. Irish Nurseries, 115. Kilkenny Nursery, &c., 116. Glasnevin and Trinity College Botanic Gardens, 116, 117.

SUBJECT. 5. Of the Foreign Trees and Shrubs introduced into Britain during the 19th Century, 117.

The number of Trees and Shrubs introduced in the three first Decades, 117. Biographical Notice of John Fraser, the Botanical Collector, 119.; of John Lyon, 122.; of David Douglas, 123. List of the Species and Varieties of Trees and Shrubs introduced into Europe by Douglas, 125. Number of Species of Foreign Trees and Shrubs introduced in Britain during every 10 Years, between 1548 and 1830; Number of Species furnished to Britain by each particular Foreign Country, 126. Planters of British Arboretums and Fruticetums since the Commencement of the 19th Century, 129. Public Bodies that have planted Arboretums, 129. British Nurserymen who have formed Arboretums, 130.

CHAP. III.

Of the History and Geography of the Trees and Shrubs of the Continent of Europe, 132.

SECT. I. Of the Indigenous and Foreign Trees and Shrubs of France, 132.

Enumeration of the Species of Trees and Shrubs indigenous to France, according to the *Botanicon Gallicum*, 132. Comparison between the Ligneous Flora of France, and that of Britain, 134. Classification of the Indigenous and Foreign Trees of France by Professor Thouin, 135. Comparison between the Number of Ligneous Plants in the Jardin des Plantes at Paris, and the London Horticultural Society's Garden, in 1834, 135. Robin, Duhamel, Jansen, Lemonnier, &c., 137. Remains of Collections of Foreign Trees and Shrubs in France, existing in 1834, 138. Existing Collectors of Foreign Trees and Shrubs in France, 139. Notice of Duhamel du Monceau, 140.; of André Michaux, 140.; of F. A. Michaux, 142.; of Dumont de Courset, and of the Gardens at Courset, 142.

SECT. II. Of the Indigenous and Foreign Trees and Shrubs of Holland and the Netherlands, 143.

Botanic Garden at Antwerp, 144. Country Seats near Rotterdam, 144. Garden at Læken, with Dimensions of some of the Exotic Trees there, 145. Brussels, Ghent, and other Botanic Gardens, 145.

SECT. III. Of the Indigenous and Foreign Trees and Shrubs of Germany, including Hungary, 145.

Enumeration of the Ligneous Plants indigenous to Germany and Hungary, 146. Historical Notices of the Places most celebrated for their Collection of Exotic Trees and Shrubs, 147. Heidelberg, Baden, Durlach, Schwäbber, Harbke, Wörlitz, Schönbrunn, 145. Botanic Garden of the University of Vienna, Kopenzel, Brünn on the Leytha, Hadersdorf, Eisenstadt, 150. Enumeration of the Species of Evergreen Trees and Shrubs, foreign and indigenous, which stand the Winter at Vienna without Protection, 150. Remarkable Specimens of Trees and Shrubs in the Botanic Garden at Berlin, 151. Foreign Trees and

Shrubs in Bavaria, in Saxony, in Hanover, 151.; in Wirtemberg, in Baden, in Hesse Cassel, in Nassau, 152.; in Mecklenburg, Anhalt, Frankfort, &c., 153. Principal German Nurseries, 153.

SECT. IV. Of the Indigenous and Foreign Trees and Shrubs of Scandinavia, including Denmark, Holstein, Sweden, Lapland, Finland, Iceland, Greenland, and the Faroe Islands, 153.

Enumeration of the Species of Ligneous Plants indigenous in the Scandinavian Peninsula, 153.; in the Faroe Islands, 154. Arboreal Flora of Sweden, of Denmark, 154. Swedish Noblemen most conspicuous for introducing Foreign Trees and Shrubs, 155.

SECT. V. Of the Indigenous and Foreign Trees and Shrubs of the Russian Empire, 155.

Enumeration of the Ligneous Species indigenous to Russia, from Pallas's *Flora Rossica*, 156. Trees and Shrubs foreign and indigenous, which stand the Winter in the Petersburg Gardens, 157. Trees and Shrubs, foreign and indigenous, which endure the Winter in Moscow, 158. Trees and Shrubs which grow in the open Air in the Crimea, 159.

SECT. VI. Of the Indigenous and Foreign Trees and Shrubs of Switzerland, 161.

Enumeration of the Indigenous Species from Gaudin's *Flora Helvetica*, 160. Exotic Trees cultivated in Switzerland, Swiss Nurseries, 161. Places celebrated for Collections of Trees, 162. Bourdigny, 163. Botanic Gardens of Geneva, 164.

SECT. VII. Of the Indigenous and Foreign Trees and Shrubs of Italy, Greece, Spain, Portugal, and the Mediterranean Islands, 164.

Enumeration of Species from various Authors, 164—168.

SUBJECT. 1. Of the Trees and Shrubs of Italy, 168.

First Introduction, 168.; into Lombardy, 168. Remarkable Specimens in different Italian Gardens, Park and Gardens of Monza, &c., Isola Bella, 169.

SUBJECT. 2. Of the Trees and Shrubs of Spain and Portugal, 170.

La Gasea, Capt. S. C. Cook, 170. Discovery by Capt. Cook and Mr. Drummond that the *Alerce* is the *Thuja articulata*, 171.

SUBJECT. 3. Of the Trees and Shrubs of Turkey and Modern Greece, 171.

CHAP. IV.

Of the Trees and Shrubs of Asia, Africa, America, and Australia, which are suitable for temperate Climates, 172.

SECT. I. Of the Trees and Shrubs suitable for Temperate Climates, indigenous or introduced, in Asia, 172.

Enumeration of the Species inhabiting Regions temperate from their Elevation, and which, it is believed, would for the greater part endure the open air in the Climate of London, from Royle's *Illustrations* and other Sources, 173. Enumeration of the Species composing the Ligneous Flora of China, and which, it is believed, would for the greater part endure the Climate of London, 176.

SECT. II. Of the Indigenous and Foreign Trees and Shrubs of Africa, 177.

SECT. III. Of the Indigenous and Foreign Trees and Shrubs of America, 178.

SUBJECT. 1. Of the Indigenous and Foreign Trees and Shrubs of North America, 178.

Enumeration of the Indigenous Species, the greater part of which have been already introduced into Britain, 178. European Trees and Shrubs introduced into America, 181. General Aspect of the Ligneous Vegetation of America, 182.; about New York, Lake Champlain, St. Lawrence, Montreal, Lake Ontario, Toronto, Niagara, Hamilton, New London, Goderich, 183.; St. Clair, Lake Huron, Lake Erie, Pittsburg, Alleghany Mountains, 184.; Mexico, 185.

SUBJECT. 2. Of the Indigenous and Foreign Trees and Shrubs of South America, 185.

SECT. IV. Of the Indigenous and Foreign Trees and Shrubs of Australia and Polynesia.

Trees and Shrubs of New Zealand, 185.; of Van Diemen's Land; of New Holland, &c., 186.

CHAP. V.

Of the Literature of the Trees and Shrubs of Temperate Climates, 187.

Aristotle, Theophrastus, Vitruvius, Crescentius, Belon, Meursius, Johnston, Aldrovandus, Evelyn, 187. Hanbury, Miller, Earl of Haddington, Mead, Morel, Boutecher, Dr. Anderson, Marshal, Nichol, Sang, Pontey, Lambert, Dr. Wade, Phillips, Watson, Cobbett, George Don, 188. In France; Dubamel, Loiseleur Deslongchamps, Desfontaines, Jaume St. Hilaire, Thouin, André Michaux, F. A. Michaux, Dumont de Courset, 189. In Holland; Krause, 189. In Germany; Du Roi, Schmidt, C. L. Willdenow. F. J. Hayne, F. Otto, 190. In America, D. J. Brown, 190.

CHAP. VI.

Conclusion, 190.

Free and universal Exchange and Intercourse, 191. Equalisation of the Plants of different Regions of the Temperate Zone; Establishment of Botanic Gardens and Arboretums all over the World, 192. Cooperative System applied to the collecting, distributing, and cultivating of Trees and Shrubs, 192.

PART II.

OF THE SCIENCE OF THE STUDY OF TREES.

CHAP. I.

Of the Study of Trees pictorially, or as component Parts of General Scenery, 193.

SECT. I. Of the Study of the Forms of Trees and Shrubs, 193.

Height and Breadth, Form and Outline, 194. Light and Shade, Colours, Trunks of Trees, 195. Branches, 196. Mode of Growth, Mode of Tufting, Leaves, Spray, Buds, 197.

SECT. II. Of the Expression and Character of Trees and Shrubs considered pictorially, 198.

Character, Regularity, 198. Symmetry, Organic Beauty, Moral and Historical Expression, 199. Picturesque Beauty, Gardenesque Beauty, Architectural and Sculpturesque Trees, Viewing Trees with Reference to their Beauty as Organic Forms, 200. Trees with Local Associations, 201.

SECT. III. Of the Mode of drawing Trees from Nature, in such a Manner as to give the general Pictorial Expression of the Species of Tree delineated, 202.

Choice of Specimens, 202. ; Details of the Process, 203. Drawing to a Scale, 204. Drawing the Leaves in order to exhibit the "Touch," 207. Botanical Specimens, 209.

CHAP. II.

Trees and Shrubs considered botanically, 211.

SECT. I. Of the Classification of Trees and Shrubs, 211.

Alphabetical Arrangement, 211. Linnæan Arrangement; Jussieuan, or Natural, Arrangement, 212.

SECT. II. Of the Distinction between Species and Varieties in Trees and Shrubs, 212.

Definition of a Species; Cultivated Varieties, Accidental Varieties, 213. Botanical Species, 214. Species according to De Candolle, 215.; Races according to the same Author, 215. Varieties, Variations, Mules and Hybrids, Botanical Species according to De Candolle 216.

SECT. III. Of the Mode of describing Trees and Shrubs, 217.

Unavoidable Evils of describing Plants from dried Specimens, 218.

CHAP. III.

Trees and Shrubs considered with Reference to their Uses in the Economy of Nature, and to Man, 219.

SECT. I. Trees and Shrubs considered with Reference to uncultivated Nature, 220.

Influence of a Predominance of Forest on the indigenous Animals; on the herbaceous Plants; on the Waters, Rivers, and Lakes; in increasing the Moisture of the Atmosphere; in moderating the Heat of Summer, and the Cold of Winter, 220. Forests of Britain, of France, of Germany, of Italy, 220.; of Holland, Belgium, &c., 221.

SECT. II. Trees and Shrubs considered with Reference to Man, 221.

Uses of Timber in Arts and Manufactures, 221. Of the Fruit, Flowers, and other Products of Trees in Domestic Economy, 222.

CHAP. IV.

Summary of Particulars to be taken into Consideration, in preparing the Description, and Natural and Economical History, of Trees and Shrubs, 222.

Classification: Genera, Distinctive Character, Identifications, Synonymes, Derivations, 222. Engravings: Engravings of Trees, Pictorial Signs, Descriptions, Descriptive Details, 223. General Descriptions, 224. Casualties: Insects and Vermin, Parasitical Plants, Diseases, Accidents, 225. Geographical Distribution, 226. History: Retrospective History, Prospective History, 226. Use: Arts of Construction, 226.; Manufacture of Machinery, &c., the Arts of Fabrication, Chemical Arts, Arts of Domestic Economy, the Arts of Rural Economy, Medicine, the Use of Trees by the Priests of particular Religions; Poetical, Mythological, and Legendary Associations; the Picturesque and Decorative Uses of Trees, 227. Propagation: Natural Propagation, Artificial Propagation, 227. Culture: the Soil, Situation, and Exposure, Culture in the Nursery, 228.; Choice of Plants, and planting out; Culture after final removal; Species adapted for Succession, 229. Statistics: Geographical Statistics, 229.; Commercial Statistics, 230.

PART III.

THE DESCRIPTION, GEOGRAPHY, HISTORY, PROPERTIES AND USES, SOIL AND SITUATION, PROPAGATION AND CULTURE, ACCIDENTS, DISEASES, INSECTS, AND STATISTICS, OF THE HARDY TREES AND SHRUBS OF BRITAIN, INDIGENOUS AND FOREIGN; WITH NOTICES OF THE HALF-HARDY SPECIES.

The Synonymes are printed in Italics. The letters *pl.* refer to the portraits of entire trees; and the letters *fig.* and *f.* to the botanical figures incorporated in the text.

	Page		Page
<i>Ranunculàcæ</i> Dec. $\text{f. } \text{f. } \text{f. } \text{f.}$	231	<i>Clematite brutante, Clematite des Haycs, L'Herbe aux Gueux, La Fiorne des Pauvres, Fr. Gemeine Waldrebe, Ger.</i>	
Sect. I. CLEMATIDÆ. $\text{f. } \text{f.}$	232	Variety - - -	236
I. CLEMATIS L. $\text{f. } \text{f.}$ - -	232	2 integrata f.	
THE CLEMATIS, or <i>Virgin's Bower. Ladies' Bower, Gerard. Clematite, Fr. Waldrebe, Ger. Clematide, Ital.</i>		7. virginiana L. $\text{f. } \text{N. America f. 13.}$	237
§ i. <i>Flammula</i> Dec. $\text{f. } \text{—}$	233	The Virginian Clematis.	
1. <i>Flammula</i> L. $\text{f. } \text{France fig. 9.}$	233	<i>C. canadensis</i> Tourn.	
The inflammatory-juiced Clematis.		<i>C. cordifolia</i> Mench.	
<i>C. òrens</i> Gerard.		Variety - - -	237
<i>C. maritima</i> All. Ped.		2 bracteata Dec. f.	
<i>C. suaveolens</i> Sal. Prod.		<i>C. bracteata</i> Mench.	
The sweet-scented <i>Virgin's Bower. Clematite odorante, Fr. Scharfe Waldrebe, Ger.</i>		8. triternata Dec. $\text{f. } \text{E. Indies}$	238
Varieties - - -	233	The triternate-leaved Clematis.	
2 rotundifolia Dec. f.		<i>Atrágene triternata</i> Desf.	
<i>C. frágans</i> Zea.		9. Viórna L. $\text{f. } \text{N. America fig. 14.}$	238
3 maritima Dec. f.		The Road-ornamenting Clematis.	
4 rubella Dec. f.		<i>C. purpúrea ripens</i> Ray.	
5 cæspitosa Dec. f.		<i>Flammula scindens</i> Dill.	
<i>C. cæspitosa</i> Scop.		<i>Leathery-flowered Clematis. Glocken-blüthige Waldrebe, Ger.</i>	
<i>C. Flammula</i> Bertol.		Variety - - -	238
2. orientalis L. $\text{f. } \text{Levant fig. 10.}$	234	2 Símsii f.	
The Oriental Clematis.		10. cylíndrica Sims. $\text{f. } \text{N. Am. fig. 15.}$	239
<i>Flammula scindens</i> Dill. Elt.		The cylindrical-flowered Clematis.	
<i>Clematis flava</i> Mench Meth.		<i>C. crispa</i> Lam.	
3. glauca Willd. $\text{f. } \text{Siberia fig. 11.}$	235	<i>C. Viórna</i> Andr.	
The glaucous-leaved Clematis.		<i>C. divaricata</i> Jacq.	
4. chinensis Retz. $\text{f. } \text{China}$	235	<i>Long-flowered Virgin's Bower.</i>	
The Chinese Clematis.		Variety - - -	238
5. paniculata Thunb. $\text{f. } \text{Japan}$	235	11. Símsii Swt. $\text{f. } \text{N. America f. 16.}$	240
The panicled Clematis.		Sims's Clematis.	
6. Vitálba L. $\text{f. } \text{England fig. 12.}$	235	<i>C. cordata</i> Sims.	
The White Vine Clematis.		12. reticulata Walt. $\text{f. } \text{N. Am. f. 17.}$	240
<i>Atrágene</i> Theoph.		The net-veined-leaved Clematis.	
<i>Vitis sylvestris</i> Diosc.		<i>C. róscæ</i> Abbott.	
<i>Clematis latifolia, seu Atrágene, Ray. C. altera</i> Matth.		§ ii. <i>Viticella</i> Dec. f.	240
<i>C. tertia</i> Comm.		13. flórida Thunb. $\text{f. } \text{Japan fig. 18.}$	240
<i>Viórna</i> Ger. and Lob.		The florid Clematis.	
<i>Vitis nigra</i> Fuchs.		<i>Atrágene indica</i> Desf.	
<i>Vitalba</i> Dod.		<i>A. flórida</i> Pers.	
<i>Traveler's Joy, Old Man's Beard, Bind-with, Common Virgin's Bower, Wild Climber, Great Wild Climber.</i>		<i>Clematite à grandes Fleurs, Fr. Grosblüthige Waldrebe, Ger.</i>	
		Variety - - -	241
		flóre pléno f.	

	Page		Page
14. <i>Viticella L.</i> § Spain - fig. 19.	241	3. <i>ochotensis Pall.</i> § Siberia -	248
The Vine-bower Clematis.		The Ochot-koi Atragene.	
<i>Viticella deltoidea</i> Moench.		A. <i>violacea</i> Pall.	
The red-flowered Lady's Bower, Gerard.		<i>Clematis ochotensis</i> Poir.	
Italianische Waldrebe, Ger.			
<i>Varieties</i> - - -	241	4. <i>americana Sims.</i> § N. Am. fig. 27.	248
1 <i>caerulea</i> §		The American Atragene.	
2 <i>purpurea</i> §		<i>Clematis verticillaris</i> Dec.	
3 <i>multiplex G. Don.</i> §		<i>Variety</i> -	248
<i>C. pulchella</i> Pers. §		2 <i>obliqua Dougl.</i> §	
4 <i>tenuifolia</i> Dec. §		5. <i>occidentalis Horn.</i> § N. America	249
5 <i>baccata</i> Dec. §		The Western Atragene.	
15. <i>campaniflora Brot.</i> § Sp. f. 20.	242	<i>Clematis occidentalis</i> Dec.	
The Bell-flowered Clematis.			
<i>C. cionoides</i> Schrad.		Sect. II. <i>PEONIA</i> CEE Dec. § 22	249
<i>C. parviflora</i> Dec.		I. <i>PEONIA</i> L. § -	249
16. <i>crispa L.</i> § N. America f. 21.	243	THE PEONY.	
The curled-sepal'd Clematis.		<i>Piony.</i>	
<i>C. flore crispa</i> Dill. Eth.		<i>Pivonne, Fr.</i>	
§ iii. <i>Choirópsis</i> Dec. § -	243	<i>Gichterrase, Pönnie, Ger.</i>	
17. <i>cirrhosa L.</i> § - fig. 22.	243	<i>Rosa del Monte, Span.</i>	
The tendriled Clematis.		<i>Peonia, Ital.</i>	
<i>Atragene cirrhosa</i> Pers.		1. <i>Moutan Sims.</i> § China -	250
<i>Traveller's Joy of Candia, Spanish Travel-</i>		The Moutan Peony.	
<i>ler's Joy, Johnson's Gerard.</i>		<i>Varieties</i> 250	
<i>Spanish Wild Cucumber, Parkinson.</i>		1 <i>papaveracea</i> <i>And.</i> § f. 28.	
<i>Evergreen Clematis.</i>		2 <i>variegata D. Don.</i> §	
<i>Einfachblüttrige Waldrebe, Ger.</i>		3 <i>Banksii</i> Andrews § fig. 29.	
<i>Variety</i> - - -	243	4 <i>Humei</i> Ker. §	
2 <i>pedicellata</i> Dec. §		5 <i>rosea</i> Dec. §	
<i>C. baccata</i> Pers.		6 <i>rosea semiplena</i> §	
<i>C. pedicellata</i> Swt.		7 <i>rosea plena</i> §	
<i>C. cirrhosa</i> Sims.		8 <i>Rawesii</i> Hort. Trans. §	
18. <i>baccata Rich.</i> § Minorca -	244	9 <i>carnea plena Hort. Trans.</i> §	
The Minorca Clematis.		10 <i>albida plena Hort. Trans.</i> §	
<i>C. calycina</i> Ait.		11 <i>Anneslei</i> Hort. Trans. §	
<i>Clematis de Mahon, Fr.</i>		12 <i>lucera Lindl.</i> §	
§ iv. <i>Anemoniflora.</i> §		Expected Varieties.	
19. <i>montana Ham.</i> § Nepal f. 23, 24.	245	II. <i>XANTHORIZA L.</i> § -	255
The Mountain Clematis.		THE YELLOW ROOT.	
<i>C. anemoniflora</i> D. Don.		<i>Zanthorize, Fr.</i>	
App. i. <i>Doubtful Species of Clematis.</i> 246		<i>Gelbwurz, Ger.</i>	
<i>C. tenuiflora</i> Dec., <i>biterata</i> Dec., <i>dahurica</i> Dec.,		1. <i>apiifolia L'Herit.</i> § N. Am. f. 31.	255
<i>diversifolia</i> Dec., <i>japonica</i> Dec., <i>semitriloba</i> Dec.,		The Parsley-leaved Yellow Root.	
<i>campaniflora</i> Dec.			
App. ii. <i>Anticipated Introductions.</i> 246		—	
<i>C. nepalensis, pallenscens, nitifolia, Buchananiana,</i>		<i>Winteracea R. Br.</i> § -	256
<i>intricata, holosericea, Walteri, Catesbyana.</i>		I. <i>ILLIUM L.</i> § -	256
II. <i>ATRA'GENE L.</i> § -	246	THE ILLIUM, or Aniseed Tree.	
THE ATRAGENE.		<i>Badiane, Anis étoilé, Fr.</i>	
<i>Clematis</i> Lam. and Dec.		<i>Sternanis, Ger.</i>	
<i>Atragene, Fr. and Ger.</i>		1. <i>floridamm Ellis.</i> § N. Am. fig. 32.	256
1. <i>alpina L.</i> § Austria fig. 25.	247	The Florida Illium.	
The Alpine Atragene.		<i>Red-flowered Aniseed Tree, Mor. Hist.</i>	
<i>Clematis caerulea</i> Bauh.		<i>Unächter Sternanis, Ger.</i>	
<i>Atragene austriaca</i> Scop.		2. <i>anisatum L.</i> § China -	257
<i>A. clematodes</i> Crantz.		The Anise Illium.	
<i>Clematis alpina</i> Mill. Diet.		<i>Chinese Aniseed Tree.</i>	
2. <i>sibirica L.</i> § Siberia fig. 26.	247	<i>Badiane de la Chine, Anis étoilé, Fr.</i>	
The Siberian Atragene.		<i>Leichter Sternanis, Ger.</i>	
A. <i>alpina</i> Gmel. Sib.		3. <i>parviflorum Michx.</i> § N. Amer.	258
<i>Clematis sibirica</i> Mill. Diet.		The small-flowered Illium.	
<i>Variety</i> - - -	248	1. <i>anisatum</i> Bartram.	

	Page		Page
<i>Magnoliaceæ Dec.</i> † † †	259	8. pyramidata <i>Bartr.</i> † <i>Carol. pl.</i> 10.	277
1. MAGNOLIA <i>L.</i> † † †	260	The pyramidal-headed Magnolia.	
THE MAGNOLIA.			
<i>Magnolie</i> , Fr.			
<i>Bieberbaum</i> , Ger.			
§ i. <i>Magnoliástrum.</i>	261	§ ii. <i>Gwillimia Rott.</i>	278
1. grandiflora <i>L.</i> † <i>Carolina pl.</i> 1.	261	9. conspicua <i>Salisb.</i> † <i>China</i>	278
The large-flowered Magnolia.		<i>pl.</i> 11. fig. 34, 35.	
<i>Laurel-leaved Magnolia, Laurel Bay, Big Laurel, Large Magnolia, Laurier Tulipier</i> , Fr.		The <i>Yulan</i> , or conspicuous-flowered, Magnolia.	
<i>Varieties</i>	261	<i>M. præcia</i> Correa.	
2 obovata <i>Ait.</i> †		<i>M. Yulan</i> Desf.	
3 rotundifolia <i>Sut.</i> †		<i>Yu lan</i> , Chinese.	
4 exoniensis <i>Hort.</i> † <i>pl.</i> 2.		<i>Lily-flowered Magnolia.</i>	
5 ferruginea <i>Sims.</i> †		<i>Magnolier Yulans</i> , Fr.	
6 lanceolata <i>Ait.</i> †		<i>Yulans Bieberbaum</i> , Ger.	
7 elliptica <i>Ait.</i> †		<i>Varieties</i>	272
8 angustifolia <i>Hort.</i> †		2 <i>Soulangeana</i>	
9 præcox <i>Hort.</i> †		<i>M. Soulangeana Ann. Hort. Soc. Par.</i> †	
Other Varieties.			
2. glauca <i>L.</i> † <i>N. America</i>	266	3 <i>Alexandrina Hort.</i> †	
The glaucous-leaved Magnolia.		4 speciosa <i>Hort.</i> †	
<i>M. frigrans</i> Salisb.		5 citriodora <i>Hort.</i> †	
<i>Swamp Sassafras, Beaver-wood, White Bay, Small Magnolia, Swamp Magnolia, Arbre de Castor</i> , Fr.		Other Varieties.	
<i>Varieties</i>	267	10. purpurea <i>Bol. Mag.</i> † <i>China</i>	282
2 <i>Thompsoniana</i> <i>Thomp.</i> †		The purple-flowered Magnolia.	
<i>M. glauca major</i> Bot. Mag.		<i>M. obovata</i> Thunb.	
3 longifolia <i>Hort.</i> †		<i>M. discolor</i> Vent.	
Other Varieties.			
3. tripétala <i>L.</i> † <i>N. America</i>	269	<i>M. denudata</i> Lam.	
The three-petaled Magnolia.		11. grácilis † <i>China</i>	283
<i>M. umbrellá</i> Lam.		The slender-growing Magnolia.	
<i>M. frondosa</i> Salisb.		<i>M. Kobus Kämpf.</i>	
<i>The Umbrella Tree, Umbrella Magnolia, Elkwood.</i>		<i>M. glauca</i> Thunb.	
<i>Magnolie Parasol, Arbre Parasol</i> , Fr.		<i>M. tomentosa</i> Thunb. in Lin. Soc.	
		<i>Sidi Kobusi</i> , Jap.	
4. macrophýlla <i>Michx.</i> † <i>Carolina</i>	271	App. i. <i>Half-hardy Magnolias.</i>	
The large-leaved Magnolia.		<i>M. fuscata Andr., anonaefolia Sal., pumila Andr.</i>	
<i>Large-leaved Umbrella Tree, Amer. M. Michauxii Hort.</i>		App. ii. <i>Additional Magnolias.</i>	
<i>Magnolier Bannanier</i> , Fr.		II. LIRIODENDRON <i>L.</i> †	
5. acuminata <i>L.</i> † <i>North America</i>	273	THE TULIP TREE.	
The pointed-leaved Magnolia.		<i>Tulipier</i> , Fr.	
<i>M. rustica</i> and <i>M. pennsylvanica</i> of some.		<i>Bieberbaum</i> , Ger.	
<i>The blue Magnolia, Eng. The Cucumber Tree, U. S.</i>		1. Tulipifera <i>L.</i> † <i>N. Am.</i>	284
<i>Varieties</i>	273	The Tulip-bearing Liriodendron, or Tulip Tree.	
2 <i>Candollei</i> <i>Savi.</i> †		<i>The Poplar, White Wood, Canoe Wood, the Tulip Tree, Amer., Virginian Poplar, Tulip-bearing Lily Tree, Saddle Tree</i> , Eng.	
3 <i>máxima</i> <i>Lodd.</i> †		<i>Tulipier de Virginie</i> , Fr.	
<i>Virginischer Tulipcerbaum</i> , Ger.			
6. cordata <i>Michx.</i> † <i>North America</i>	275	<i>Varieties</i>	285
The heart-leaved Magnolia.		2 obtusiloba <i>Michx.</i> †	
<i>The heart-leaved Cucumber Tree, Amer.</i>		<i>L. T. integrifolia Hort.</i>	
7. auriculata <i>Lam.</i> † <i>Carolina</i>	276	<i>Yellow Wood, Yellow Poplar.</i>	
The auricled-leaved Magnolia.		3 acutifolia <i>Michx.</i> †	
<i>M. Fræseri</i> <i>Walt.</i>		4 flava <i>Hort.</i> †	
<i>M. auriculatis</i> Salisb.		App. I. <i>Expected Additions to the Order Magnoliaceæ.</i>	
<i>Indian Physic, Long-leaved Cucumber Tree, Amer.</i>		<i>Magnolia</i> - - - 291	
		<i>Manglietia (Magnolia) insignis; Michelia lanuginosa, excelsa, Kisopa, Doltsopa.</i>	
—————			
<i>Dilleniaceæ Dec.</i> † † - 292			
1. <i>Hibbertia volubilis</i> † Cape of Good Hope		- fig. 38, 292	
2. <i>H. dentata</i> † New Holland		- - - fig. 37, 292	
5. <i>H. grossulariifolia</i> † New Holland		- - - 292	

	Page		Page
Anonæacæ Rich.			
I. ASIMINA Adams. ☼ -	292	II. COCCULUS Bauh. ☼ -	297
THE ASIMINA.		THE COCCULUS.	
<i>Annona L.</i>		<i>Menispermum L.</i>	
<i>Orchidocarpum</i> Mx.		<i>Wendlandia Willd.</i>	
<i>Porcchia Pers.</i>		<i>Androphilar Wendl.</i>	
<i>Custard Apple.</i>		<i>Menispermæ, Fr.</i>	
<i>Asiminier, Fr.</i>		<i>Mondsaauc, Ger.</i>	
<i>Flaschenbaum, Ger.</i>			
1. <i>parviflora</i> Dun. ☼ N. America -	293	1. <i>carolinus</i> Dec. ☼ — N. A. f. 45.	298
The small-flowered Asimina.		The Carolina Cocculus.	
<i>Porcchia parviflora Pers.</i>		<i>Menispermum carolinum L.</i>	
<i>Orchidocarpum parviflorum Mx.</i>		<i>Wendlandia populifolia Willd.</i>	
		<i>Androphilar scandens Wendl.</i>	
		<i>Baumgärtia scandens Mench.</i>	
2. <i>triloba</i> Dun. ☼ N. America f. 39.	293	App. I. <i>Anticipated Menispermæacæ.</i>	298
The three-lobed-calyced Asimina.			
<i>Annona triloba L.</i>		Berberæcæ Lindl. ☼ ■ 298	
<i>Porcchia triloba Pers.</i>		I. BERBERIS L. ☼ ■ -	298
<i>Orchidocarpum arcticum Mx.</i>		THE BERBERIS.	
The Papaw, Amer.		<i>Piperidge Bush.</i>	
<i>Asiminier de Virginie, Fr.</i>		<i>Épine vinette, Fr.</i>	
		<i>Berberitze, Ger.</i>	
3. <i>pygmæa</i> Dun. ☼ N. America f. 40.	294	1. <i>sibirica</i> Pall. ☼ Siberia f. 46.	301
The pygmy Asimina.		The Siberian Berberry.	
<i>Annona pygmæa, Bartr.</i>		<i>B. altaica</i> Pall.	
<i>Orchidocarpum pygmæum Mx.</i>		2. <i>vulgàris</i> L. ☼ England -	301
<i>Porcchia pygmæa Pers.</i>		The common Berberry.	
4. <i>grandiflora</i> Dun. ☼ N. America	295	<i>Varieties - - 301</i>	
The large-flowered Asimina.		2 <i>lutea</i> ☼	
<i>Annona grandiflora Bartr.</i>		3 <i>alba</i> ☼	
<i>A. oborata Willd.</i>		4 <i>violæca</i> ☼	
<i>Orchidocarpum grandiflorum Mx.</i>		5 <i>purpurea</i> ☼	
<i>Porcchia grandiflora Pers.</i>		6 <i>nigra</i> ☼	
		7 <i>dulcis</i> ☼	
		8 <i>asperma</i> ☼	
		Other Varieties: <i>B. daurica</i> and <i>altaica</i> .	
Schizandræcæ. ☼ - 295		3. <i>canadensis</i> Mill. ☼ N. Am. f. 48.	303
I. SCHIZANDRA Michx. ☼ —	295	The Canadian Berberry.	
THE SCHIZANDRA.		<i>B. vulgàris</i> Michx.	
1. <i>coccinea</i> Michx. ☼ — N. Amer. f. 41.	295	<i>B. v. canadensis</i> Mart. Mill.	
The scarlet-flowered Schizandra.		4. <i>emarginata</i> Willd. ☼ Siberia f. 49.	303
App. I. <i>Anticipated Additions to Schizandrae.</i>	295	The emarginate-pectalcd Berberry.	
<i>Sphaerostema grandiflorum, Kotsura japonica</i>	29	<i>Ausgeraudete Berberitze, Ger.</i>	
Menispermæcæ Dec. ☼ ■ 296		5. <i>ibérica</i> Stev. ☼ Iberia - f. 50.	304
I. MENISPERMUM L. ☼ ☼ -	296	The Iberian Berberry.	
THE MOONSEED.		<i>B. vulgàris f. n. ibérica</i> Dec. Syst.	
<i>Menispermæ, Fr.</i>		<i>B. sinensis</i> Wal.	
<i>Mondsaauc, Ger.</i>		6. <i>sinensis</i> Desf. ☼ China -	304
1. <i>canadense</i> L. ☼ N. Amer. f. 42.	296	The Chinese Berberry.	
The Canadian Moonseed.		<i>B. vulgàris</i> Thunb.	
<i>M. canadense var. α</i> Lam.		7. <i>crética</i> L. ☼ Candia -	304
<i>M. angulatum</i> Mench.		The Cretan Berberry.	
<i>Variety - - 296</i>		<i>B. crética buxifolia</i> Tourn.	
2 <i>lobatum</i> Dec. ☼		<i>Variety - - 305</i>	
<i>M. virginicum</i> L.		2 <i>serratifolia</i> Poir.	
2. <i>dauricum</i> Dec. ☼ Dauria f. 43.	297	8. <i>cratægina</i> Dec. ☼ Asia Minor	305
The Daurian Moonseed.		The Cratægus-like Berberry.	
<i>Tetlophus impellisægia</i> Fisch.		9. <i>dulcis</i> Swt. ☼ S. America f. 51.	305
<i>M. canadense var. β</i> Lam.		The sweet-fruited Berberry.	
3. <i>smilacinum</i> Dec. ☼ -	fig. 44. 297		
The Smilax like Moonseed.			
<i>Cissampelos sulcatissima</i> L.			

	Page
10. heterophýlla <i>Juss.</i> S. Am. f. 52.	305
The various-leaved Berberry.	
<i>B. ilicifolia</i> Forst.	
<i>B. cuspidata</i> Smith.	
11. empetrifolia <i>Lam.</i> Chili -	306
The Empetrum-leaved Berberry.	
12. floribunda <i>Wall.</i> Nepal -	305
The many-flowered Berberry.	
<i>B. affinis.</i>	
<i>B. ceratophýlla.</i>	
13. asiática <i>Rox.</i> Nepal -	306
The Asiatic Berberry.	
<i>B. tinctoria</i> Lech.	
The Raisin Berberry.	
14. dealbata <i>Lindl.</i> Mexico f. 35.	307
The whitened-leaved Berberry.	
<i>B. glauca</i> Hort.	
15. aristata <i>Dec.</i> Nepal fig. 54.	307
The bristle-toothed-leaved Berberry.	
<i>B. Chitria</i> Buch.	
<i>B. angustifolia</i> Roxb.	
<i>B. sinensis</i> Delf.	

App. i. *Additional Species of Bérberis.* 308
B. Wallichiana Dec., syn. *atro-viridis*, *B. kunawurensis*, *B. ac-*
tinacantha, *B. buxifolia* (f. 55.), *B. ilicifolia.*

II. MAHONIA *Nutt.* - 308
 THE MAHONIA.

<i>Ashberry.</i>	
<i>Bérberis of Authors.</i>	
<i>Oostemon</i> Raf.	
1. fascicularis <i>Dec.</i> N. Am. f. 56.	309
The crowded-racemed Mahonia.	
<i>Bérberis pinnata</i> Lag.	
<i>B. fascicularis</i> Pen. Cyc.	
2. Aquifolium <i>Nutt.</i> N. Am. f. 57.	309
The Holly-leaved Mahonia.	
<i>B. Aquifolium</i> Pursh.	
<i>Variety</i> - - 309	
2 nutkana <i>Dec.</i>	
3. nervosa <i>Nutt.</i> N. Amer. f. 58.	310
The nerved-leaved Mahonia.	
<i>Bérberis nervosa</i> Pursh.	
<i>Mahonia glutinosa</i> Dec.	
<i>Bérberis glutinosa</i> Pen. Cyc.	
4. repens <i>G. Don.</i> N. Amer. f. 59.	311
The creeping-rooted Mahonia.	
<i>Bérberis Aquifolium</i> Lindl.	
<i>B. repens</i> Pen. Cyc.	

App. i. *Additional Species of Mahonia.* 311
M. nepalensis, *acanthifolia*, *tragacanthoides*, and
caraganaefolia.

Cruciacea. - 312

I. VELLA *L.* - 312
 THE VELLA.

1. Pseudo-Cytisus <i>L.</i> Spain f. 60.	312
The False-Cytisus Cross-Rocket.	
<i>V. integrifolia</i> Sal.	
<i>Faux-Cytise.</i>	
<i>Strauchartige Velle</i> , Ger.	
App. I. <i>Other Cruciacea.</i> - 313	
<i>Cheiranthus Cheri</i> , <i>C. C. fruticulosus</i> ; <i>Vesicaria</i>	
<i>utriculata</i> ; <i>Alyssum argenteum</i> , saxatile (f. 61.),	
<i>gemonense</i> ; <i>Iberis sempervirens</i> (f. 62.), sax-	
<i>atilis</i> , <i>corifolia</i> ; <i>Lepidium subulatum</i> , sulfruti-	
<i>cosum</i> ; <i>Sisymbrium Millefolium.</i>	


Capparidacea *Lindl.* - 313

I. CAPPARIS *L.* -
 THE CAPER BUSH.

<i>Caprier</i> , <i>Tapenier</i> , Fr.	
<i>Capernstrauch</i> , Ger.	
<i>Capriolo</i> , <i>Cappero</i> , Ital.	
1. spinosa <i>L.</i> Levant - fig. 63.	314
The spined, or common, Caper Bush.	
2. Fontanèsi Dec. Barbary -	316
<i>Desfontaine's Caper Bush.</i>	
<i>C. ovata</i> Desf.	
<i>Caprier oval</i> , Fr.	

C. ægypta *Lam.*, *nepalensis* *Dec.*, *nummularia* *Dec.*, *quini-*
flora *Dec.*, *umbellata* *R. Br.*, *canescens* *Banks*, *heteracanth-*
tha *Dec.*, *leucophylla* *Dec.*, *Volkameria* *Dec.*, *citrifolia* *Lam.*,
cluytiofolia *Burch.*, *oleoides* *Burch.*, *coriacea* *Burch.*, *albu-*
trunca *Burch.*, *punctata* *Burch.*, *racemosa* *Dec.*, *saligna*
Vahl.

Cista *Dec.* *Lindl.* - 316



I. CISTUS *L.* - 317
 THE ROCK ROSE.

<i>Holly Rose</i> , Gerard.	
<i>Gum Cistus.</i>	
<i>Ciste</i> , Fr.	
<i>Cisten Rose</i> , Ger.	

§ i. *Erythrocistus* *Dec.* - 318

1. purpureus <i>Lam.</i> Levant. f. 64.	318
The purple-flowered Cistus.	
<i>C. creticus</i> Hort. Kew.	
2. heterophyllus <i>Desf.</i> Alg. f. 65.	318
The various-leaved Cistus.	
The <i>Cistus of Algiers.</i>	
3. parviflorus <i>Lam.</i> Crete -	319
The small-flowered Cistus.	
4. complicatus <i>Lam.</i> Spain -	319
The complicated Cistus.	
5. villosus <i>Lam.</i> S. Europe -	319
The villous Cistus.	
<i>C. salvifolius</i> Hort.	
<i>C. undulatus</i> Mœnch.	
<i>C. más major</i> , &c., Duh.	
<i>C. creticus</i> Hort.	
The <i>shrubby Cistus</i> , Mart. Mill.	
<i>Variety</i> - - 319	
2 rotundifolius	
<i>C. rotundifolius</i> Swt.	
<i>C. villosus</i> β <i>virescens</i> <i>Dec.</i>	

6. creticus <i>L.</i> Levant fig. 66.	319
The Cretan Cistus.	
<i>Lèdon</i> Diosc.	

7. inèanus <i>L.</i> S. Europe fig. 67.	320
The hoary Cistus.	

<i>C. albidus</i> Hort.	
<i>C. cymosus</i> <i>Dec.</i>	
<i>Ciste colonnecur</i> , Fr.	
<i>Bestaubte Cisten Rose</i> , Ger.	

Variety - - 321

2 canescens	
<i>C. canescens</i> Swt.	
<i>C. más</i> Clus.	
<i>C. inèanus</i> var. β <i>Dec.</i>	
8. undulatus <i>Dec.</i> -	321
The waved-leaved Cistus.	
<i>C. creticus</i> Swt., afterwards <i>C. undulatus</i>	
<i>Swt.</i>	
? <i>C. crispus</i> var. <i>Don.</i>	

	Page		Page
9. <i>crispus</i> L. \underline{n} France -	321	28. <i>asperifolius</i> Swt. \underline{n} -	326
The curled-leaved Cistus.		The rough-leaved Cistus.	
10. <i>albidus</i> L. \underline{n} France -	321	29. <i>psilosépalus</i> Swt. \underline{n} -	326
The white-leaved Cistus.		The glabrous-sepaled Cistus.	
11. <i>candidissimus</i> Dum. \underline{n} Canaries	322	30. <i>longifolius</i> Lam. \underline{n} S. Europe	326
The whitest-leaved Cistus.		The long-leaved Cistus.	
12. <i>vaginatus</i> Ait. \underline{n} Teneriffé -	322	<i>C. ulgricosa</i> Pourr.	
The sheathed-petioled Cistus.		<i>C. populifolius</i> var. <i>longifolius</i> Dumont.	
<i>C. symphytifolius</i> Lam.		31. <i>populifolius</i> L. \underline{n} Spain -	326
<i>Ciste à Feuilles de Consoude</i> , Fr.		The Poplar-leaved Cistus.	
<i>Scheidnartige Cisten</i> Rose, Ger.		Variety \underline{n} -	326
13. <i>sericeus</i> Vahl. \underline{n} Spain -	322	2 minor Dec. \underline{n}	
The silky-leaved Cistus.		<i>Lédon latifolium</i> , ii., Clus.	
14. <i>hybridus</i> Vahl. \underline{n} Spain -	322	<i>C. populifolius</i> Cav.	
The hybrid Cistus.		32. <i>latifolius</i> Swt. \underline{n} Barbary -	327
15. <i>cymosus</i> Dum. \underline{n} Spain -	322	The broad-leaved Cistus.	
The cymose-flowered Cistus.		<i>C. populifolius</i> var. <i>major</i> Dec.	
<i>C. incanus</i> Sib.		33. <i>laurifolius</i> Lin. \underline{n} Spain -	327
§ ii. <i>Ladonia</i> . \underline{n} \underline{n} \underline{n} \underline{n} \underline{n}	323	The Laurel-leaved Cistus.	
16. <i>salviaefolius</i> L. \underline{n} S. Europe	323	34. <i>cyprius</i> Lam. \underline{n} Greece -	327
The Sage-leaved Cistus.		The Gum Cistus.	
<i>C. foemina</i> Clus.		<i>C. ladaniferus</i> Bot. Mag.	
<i>Ciste à Feuilles de Sauge</i> , Fr.		<i>C. stenophyllus</i> Link.	
<i>Salbei-blättrige Cisten</i> Rose, Ger.		<i>C. salicifolius</i> of some.	
Varieties \underline{n} -	323	35. <i>ladaniferus</i> L. \underline{n} Spain -	328
2 erectiúsculus Dec. \underline{n}		The Ladanum-bearing Gum Cistus.	
3 ochroleucus Dec. \underline{n}		Varieties \underline{n} -	328
17. <i>obtusifolius</i> Swt. \underline{n} Spain -	323	1 albiflorus Dec. \underline{n}	
The obtuse-leaved Cistus.		2 maculatus Dec. \underline{n}	
18. <i>Cupaniæus</i> Presl. \underline{n} Sicily -	324	3 plenifolius Dec. \underline{n}	
Cupani's Cistus.		36. <i>Clusii</i> Dunal. \underline{n} Spain -	328
Variety \underline{n} -	324	Clusius's Cistus.	
2 acutifolius \underline{n}		<i>C. Libanotis</i> β Lam.	
The acute-leaved Cupani's Cistus.		<i>C. undulatus</i> Link.	
<i>C. acutifolius</i> Swt.		<i>Lédon</i> , vii., Clus.	
<i>C. salviaefolius</i> β <i>humifusus</i> Dec.		II. HELIANTHEMUM Tourn. \underline{n} \underline{n}	317. 328
19. <i>corbáriensis</i> Pourr. \underline{n} Spain	324	THE HELIANTHEMUM, or Sun Rose.	
The Corbières Cistus.		<i>Cisti</i> sp. L.	
<i>C. salviaefolius</i> β Dec.		<i>Helianthemum</i> Sonnen Gurtel, Ger.	
<i>C. populifolius</i> minor, in some nurseries.		§ i. <i>Halimium</i> Dec. \underline{n} \underline{n} \underline{n} \underline{n}	329
<i>C. hybridus</i> Pourr., not of Vahl.		1. <i>Libanotis</i> Willd. \underline{n} \underline{n} Spain	329
20. <i>florentinus</i> Lam. \underline{n} Spain -	324	The Rosemary-leaved Helianthemum.	
The Florentine Cistus.		<i>Cistus Libanotis</i> L.	
21. <i>monspeliensis</i> L. \underline{n} S. Europe	324	<i>H. rosmarinifolium</i> Lag.	
The Montpellier Cistus.		<i>Lédon</i> , viii., Clus.	
22. <i>platysépalus</i> Swt. \underline{n} Crete -	325	<i>Ciste à Feuilles de Rosmarin</i> , Fr.	
The broad-sepaled Cistus.		2. <i>umbellatum</i> Mill. \underline{n} \underline{n} S. Eur.	329
23. <i>Lédon</i> Lam. \underline{n} France -	325	The umbellate-flowered Helianthemum.	
The Lédon Cistus.		<i>Cistus umbellatum</i> , L.	
<i>C. undulatus</i> Link. ?		<i>Cistus Libanotis</i> of some.	
<i>C. ladaniferus</i> <i>monspeliensis</i> Haub.		Varieties \underline{n} \underline{n}	330
<i>Ciste d' Montpellier qui donne du ladanum</i> , <i>Ciste de Lédon</i> , Fr.		2 erectum Dec. \underline{n} \underline{n}	
24. <i>hirsutus</i> Lam. \underline{n} Spain -	325	3 subdecumbens Dec. \underline{n} \underline{n}	
The hairy Cistus.		3. <i>ocymoides</i> Pers. \underline{n} \underline{n} Spain	330
25. <i>Sideritis</i> Presl. \underline{n} Sicily -	325	The Basil-like Helianthemum.	
The Ironwort-like Cistus.		<i>Cistus ocymoides</i> Lam.	
26. <i>lævis</i> Ait. \underline{n} Spain -	325	<i>Cistus sampsucifolius</i> Cav.	
The loose flowering Cistus.		4. <i>alysoides</i> Vent. \underline{n} \underline{n} Spain	330
<i>C. aspasia</i> L.		The Alyssum-like Helianthemum.	
<i>S. Miffle Cisten</i> Rose, Ger.		<i>Cistus alyssoides</i> a Lam.	
27. <i>oblongifolius</i> Swt. \underline{n} Spain -	326	5. <i>rugosus</i> Dum. \underline{n} \underline{n} Portugal	330
The oblong-leaved Cistus.		The wrinkled-leaved Helianthemum.	

- | | Page | | Page |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----------------------------------------------------------------------------------|------|
| 6. <i>microphyllum</i> <i>Swt.</i> n. France | 330 | § v. <i>Fumàna</i> <i>Dec.</i> n. — — — — | 335 |
| The small-leaved Helianthemum. | | 23. <i>Fumàna</i> <i>Mill.</i> n. France f. 72. | 336 |
| <i>H. rugosum</i> β <i>microphyllum</i> <i>Dec. Prod.</i> | | The Fumana Helianthemum. | |
| <i>H. alyssoides</i> β <i>microphyllum</i> <i>Dec. Fl. Fr.</i> | | <i>Cistus Fumàna</i> <i>L.</i> | |
| 7. <i>scabrosum</i> <i>Pers.</i> n. Port. f. 68. | 331 | <i>Cistus humilis, seu Chamæcistus Erica julio luteus ex ætior.</i> <i>Bauh.</i> | |
| The rough Helianthemum. | | <i>Varieties</i> n. - - - | 336 |
| <i>Cistus scabrosus</i> <i>Ait.</i> | | 2 major <i>Dec.</i> n. | |
| 8. <i>algarvænsis</i> <i>Dun.</i> n. — Portugal | 331 | 2 minor <i>Dec.</i> n. | |
| The Algarve Helianthemum. | | 3 virgata <i>Dec.</i> n. | |
| <i>Cistus algarvænsis</i> <i>Bot. Mag.</i> | | 24. <i>procumbens</i> <i>Dun.</i> n. — S. Europe | 336 |
| 9. <i>formosum</i> <i>Dun.</i> n. — Portugal | 331 | The procumbent Heath-like Helianthemum. | |
| The beautiful Helianthemum. | | <i>Cistus humilis, sive Chamæcistus Erica julio humilior.</i> <i>Magn. Bot.</i> | |
| <i>Cistus formosus</i> <i>Bot. Mag.</i> | | 25. <i>arabicum</i> <i>Pers.</i> n. — Arabia | 336 |
| 10. <i>atriplicifolium</i> <i>Willd.</i> n. — Spain | 332 | The Arabian Helianthemum. | |
| The Orache-leaved Helianthemum. | | <i>Cistus arabicum</i> <i>L.</i> | |
| <i>Cistus atriplicifolius</i> <i>Lam.</i> | | <i>Cistus ferrugineus</i> <i>Lam.</i> | |
| 11. <i>lasianthum</i> <i>Pers.</i> n. — Spain | 332 | <i>Cistus Savi</i> <i>Berol.</i> | |
| The hairy-flowered Helianthemum. | | <i>H. viscidulum</i> <i>Stev.</i> | |
| <i>Cistus lasianthus</i> <i>Lam.</i> | | 26. <i>lævipes</i> <i>Willd.</i> n. — Spain - | 336 |
| 12. <i>involutratum</i> <i>Pers.</i> n. — Spain | 332 | The smooth-peduncled Helianthemum. | |
| The involuted-flowered Helianthemum. | | <i>Cistus lævipes</i> <i>L.</i> | |
| <i>Cistus involutatus</i> <i>Lam.</i> | | 27. <i>læve</i> <i>Pers.</i> n. — Spain - | 337 |
| 13. <i>cheiranthoides</i> <i>Pers.</i> n. — Port. 332 | | The smooth Helianthemum. | |
| The Wallflower-like Helianthemum. | | <i>Cistus lævis</i> <i>Cav.</i> | |
| <i>Cistus cheiranthoides</i> <i>Lam.</i> | | 28. <i>viride Tenore</i> n. — Sicily - | 337 |
| <i>Cistus elongatus</i> <i>Vahl.</i> | | The green-towered Helianthemum. | |
| <i>Cistus halimifolius</i> , <i>ii.</i> , <i>Clus.</i> | | 29. <i>juniperinum</i> <i>Lag.</i> n. — S. Europe | 337 |
| 14. <i>candidum</i> <i>Swt.</i> n. Spain | 332 | The Juniper-like Helianthemum. | |
| The white-leaved Helianthemum. | | <i>Cistus lævipes</i> <i>Durand.</i> | |
| 15. <i>halimifolium</i> <i>Willd.</i> n. — Spain | 333 | <i>Cistus mauritanicus</i> <i>Thib.</i> | |
| The Sea-Purslane-leaved Helianthemum. | | 30. <i>Barrelieri Tenore.</i> n. — Italy - | 337 |
| <i>Cistus halimifolius</i> <i>L.</i> | | The Barrelier's Helianthemum. | |
| <i>Cistus julio Halimi</i> , <i>i.</i> , <i>Clus.</i> | | 31. <i>thymifolium</i> <i>Pers.</i> n. — Spain | 337 |
| § <i>ii.</i> <i>Lecheoides</i> <i>Dec.</i> n. n. — — — — | 333 | The Thyme-leaved Helianthemum. | |
| 16. <i>corymbosum</i> <i>Michx.</i> n. — Mexico | 333 | <i>Cistus thymifolius</i> <i>L.</i> | |
| The corymbose-flowered Helianthemum. | | <i>H. glutinosum</i> β <i>Dec.</i> | |
| 17. <i>glomeratum</i> <i>Lag.</i> n. — Mex. f. 69. | 333 | 32. <i>glutinösium</i> <i>Pers.</i> n. — Spain | 337 |
| The glomerate-flowered Helianthemum. | | The clammy Helianthemum. | |
| <i>Cistus glomeratus</i> <i>Lag. Gen.</i> | | <i>Cistus glutinosus</i> <i>L.</i> | |
| 18. <i>brasiliense</i> <i>Pers.</i> n. — Brazil f. 70. | 334 | § <i>vi.</i> <i>Pseudo-Cistus</i> <i>Dec.</i> n. — — — — | 338 |
| The Brazilian Helianthemum. | | 33. <i>molle</i> <i>Pers.</i> n. n. — Spain - | 338 |
| <i>Cistus brasiliensis</i> <i>Lam.</i> | | The soft-leaved Helianthemum. | |
| <i>Cistus alternifolius</i> <i>Vahl.</i> | | <i>Cistus mollis</i> <i>Cav.</i> | |
| <i>Other Species belonging to this Division of</i> | | 34. <i>origanifolium</i> <i>Pers.</i> n. — Spain | 338 |
| <i>Lecheoides.</i> | 334 | The Marjoram-leaved Helianthemum. | |
| <i>H. polygalæfolium</i> <i>Swt.</i> n. — <i>ástylum</i> <i>Moc. et Sesse,</i> <i>tripetalum</i> <i>Moc. et Sesse,</i> <i>obcordatum</i> <i>Moc. et Sesse.</i> | | <i>Cistus origanifolius</i> <i>Lam.</i> | |
| § <i>iii.</i> <i>Tuberaria</i> <i>Dec.</i> n. | 334 | 35. <i>dichotomum</i> <i>Dunal.</i> n. — Spain | 338 |
| 19. <i>lignosum</i> <i>Swt.</i> n. S. Europe f. 71. | 334 | The dichotomous-branched Helianthemum. | |
| The woody Helianthemum. | | <i>Cistus dichotomus</i> <i>Cav.</i> | |
| § <i>iv.</i> <i>Eriocarpon</i> <i>Dec.</i> n. n. — — | 335 | 36. <i>celandicum</i> <i>Dec.</i> n. Europe - | 338 |
| 20. <i>Lippii</i> <i>Pers.</i> n. — Egypt | 335 | The Eland Helianthemum. | |
| Lippi's Helianthemum. | | <i>Cistus celandicus</i> <i>L.</i> | |
| <i>Cistus Lippii</i> <i>L.</i> | | <i>Chamæcistus, ii.</i> , <i>Clus.</i> | |
| 21. <i>sessiliflorum</i> <i>Pers.</i> n. — N. Africa | 335 | 37. <i>pulchellum</i> <i>Swt.</i> n. Germany | 339 |
| The sessile-flowered Helianthemum. | | The neat Helianthemum. | |
| <i>Cistus sessiliflorus</i> <i>Desf.</i> | | <i>H. alpestre</i> <i>Spreng.</i> | |
| 22. <i>kahiricum</i> <i>Del.</i> n. — Egypt | 335 | 38. <i>alpestre</i> <i>Dunal.</i> n. S. Europe | 339 |
| The Cairo Helianthemum. | | The Alpine Helianthemum. | |
| <i>Cistus stipulatus</i> β <i>Forsk.</i> | | <i>Cistus alpestris</i> <i>Crantz.</i> | |
| | | <i>Cistus celandicus</i> <i>Jacq.</i> | |
| | | <i>Varieties</i> n. - - - | 339 |
| | | <i>glabratum</i> <i>Dec.</i> n. | |
| | | <i>elongatum</i> <i>Dec.</i> n. | |
| | | <i>canescens</i> <i>Dec.</i> n. | |

	Page
39. penicillatum <i>Thib.</i> 2 France - 339 The pencilled Helianthemum. <i>Cistus echinoides</i> Lam. <i>Cistus anglicus</i> L.	339
40. obovatum <i>Dunal.</i> 2 Spain - 339 The obovate-leaved Helianthemum. <i>Cistus italicus</i> L.	339
41. italicum <i>Pers.</i> 2 Italy - 339 The Italian Helianthemum. <i>Cistus italicus</i> L. <i>Cistus marifolius</i> Bieb.	339
Varieties 2 - 339	
strigosum <i>Dec.</i> 2 - 339	
candidissimum <i>Dec.</i> 2 - 339	
albium <i>Dec.</i> 2 - 339	
42. vineale <i>Pers.</i> 2 Germany - 340 The Vineyard Helianthemum. <i>Cistus vinealis</i> Willd.	340
43. canum <i>Dunal.</i> 2 S. Europe 340 The hoary Helianthemum <i>Cistus canus</i> L. <i>Chamaecistus</i> , iii., Clus.	340
44. marifolium <i>Dec.</i> 2 France - 340 The Marum-leaved Helianthemum. <i>Cistus marifolius</i> L.	340
45. crassifolium <i>Pers.</i> 2 Spain - 340 The thick-leaved Helianthemum. <i>Cistus glaucus</i> Desf. <i>H. Sixte</i> Lag.	340
46. paniculatum <i>Dunal.</i> 2 Spain 340 The panicked-flowered Helianthemum. <i>Cistus marifolius</i> Thib. <i>Cistus nummularius</i> var. Lag. <i>H. sp. nova</i> Schouw.	340
§ vii. <i>Euhelianthemum</i> <i>Dec.</i> 2 2 340	340
47. lavandulæfolium <i>Dec.</i> 2 France 341 The Lavender-leaved Helianthemum. <i>Cistus lavandulæfolius</i> Lat.	341
Varieties 2 - 341	
syriacum <i>Dec.</i> 2 - 341	
<i>Cistus syriacus</i> Jacq.	
Thibaudi <i>Pers.</i> 2 - 341	
<i>Cistus racemosus</i> Cav.	
48. stœchadifolium <i>Pers.</i> 2 Spain 341 The French-Lavender-leaved Helianthemum. <i>Cistus stœchadifolius</i> Brot.	341
49. crœcum <i>Pers.</i> 2 Spain - 341 The Saffron-coloured-flowered Helianthemum. <i>Cistus crœcus</i> Desf.	341
Varieties - - 341	
50. Andersoni <i>Swt.</i> 2 Hybrid - 342 Anderson's Helianthemum.	342
51. nudicaule <i>Dunal.</i> 2 Spain - 342 The naked-stemmed Helianthemum.	342
52. glaucum <i>Pers.</i> 2 Spain - 342 The glaucous Helianthemum. <i>Cistus glaucus</i> Cav.	342
Varieties 2 - 342	
acutiusculum <i>Dec.</i> 2 - 342	
obtusiusculum <i>Dec.</i> 2 - 342	
53. tomentosum <i>Dunal.</i> 2 France 342 The tomentose Helianthemum.	342
54. barbatum <i>Pers.</i> 2 S. Europe 343 The bearded-racemed Helianthemum. <i>Cistus barbatus</i> Lam.	343
55. leptophyllum <i>Dunal.</i> 2 Spain 343 The slender-leaved Helianthemum. <i>Cistus angustifolius</i> Lag. <i>Cistus stœchadifolius</i> Hort.	343
56. acuminatum <i>Pers.</i> 2 France - 343 The acuminated Helianthemum. <i>Cistus serpyllifolius</i> Mill. <i>Cistus acuminatus</i> Niv.	343
57. serpyllifolium <i>Mill.</i> 2 S. Europe 343 The Wild-Thyme-leaved Helianthemum. <i>Cistus serpyllifolius</i> L.	343
58. vulgare <i>Gart.</i> 2 Britain - 343 The common Helianthemum. <i>Cistus Helianthemum</i> L.	343
Varieties - - 344	
Double-flowered.	
Lee's new double yellow.	
Tomentose pubescent branches,	
<i>Dec.</i>	
Branches glabrous at the base,	
<i>Dec.</i>	
59. surrejanum <i>Mill.</i> 2 England 344 The Surrey Helianthemum. <i>Cistus surrejanus</i> L.	344
60. ovatum <i>Dunal.</i> 2 Geneva - 344 The ovate-leaved Helianthemum. <i>Cistus ovatum</i> Viv.	344
61. grandiflorum <i>Dec.</i> 2 Pyrenees 344 The large-flowered Helianthemum. <i>Cistus grandiflorus</i> Scop.	344
62. obscurum <i>Pers.</i> 2 S. Europe - 344 The obscure Helianthemum. <i>H. obscurum</i> a <i>Dec.</i>	344
63. tauricum <i>Fisch.</i> 2 Tauria - 345 The Taurian Helianthemum.	345
64. lucidum <i>Horn.</i> 2 - 345 The shining-leaved Helianthemum.	345
65. nummularium <i>Mill.</i> 2 France 345 The Moneywort-leaved Helianthemum. <i>Cistus nummularius</i> a L. <i>H. obscurum</i> β <i>nummularium</i> <i>Dec.</i> <i>H. angustifolium</i> Hort.	345
66. angustifolium <i>Pers.</i> 2 Cyprus 345 The narrow-leaved Helianthemum. <i>Cistus angustifolius</i> Jacq. ? <i>H. nummularium</i> <i>Dec.</i>	345
67. hirtum <i>Pers.</i> 2 Spain - 345 The hairy Helianthemum. <i>Cistus hirtus</i> L.	345
Varieties 2 - 346	
hæticum <i>Dec.</i> 2 - 346	
aureum <i>Dec.</i> 2 - 346	
teretifolium <i>Dec.</i> 2 - 346	
68. Lagasœa <i>Dunal.</i> 2 Spain 346 <i>La Gasca's</i> Helianthemum. <i>H. hirtum</i> var. Lag.	346
69. violaceum <i>Pers.</i> 2 Spain - 346 The Violet-coloured-colored Helianthemum. <i>Cistus violaceus</i> Cav.	346
70. racemosum <i>Dunal.</i> 2 Spain - 346 The racemose-flowered Helianthemum. <i>Cistus racemosus</i> L.	346
71. farinosum <i>Scot.</i> 2 Spain - 346 The mealy-leaved Helianthemum. <i>H. racemosum</i> β <i>farinosum</i> <i>Dec.</i>	346
72. strictum <i>Pers.</i> 2 Spain - 346 The straight-branched Helianthemum. <i>Cistus strictus</i> Cav.	346

	Page		Page
73. pilòsum Pers. 2 France	- 346	86. variegatum Swt. 2 Hybrid	350
The pilose Helianthemum.		The variegated Helianthemum.	
<i>Cistus pilòsus</i> L.			
<i>Chamaecistus</i> , iv., Clus.			
<i>Varieties</i> 2	- 346	87. versicolor Swt. 2 S. Europe	350
With linear leaves, hoary, Dec. 2		The various-coloured Helianthemum.	
With linear leaves, oblong and shining,			
Dec. 2		88. sulphureum Willd. 2 Spain	351
		The sulphur-colour-flowered Helianthemum.	
74. lineare Pers. 2 Spain	- 347	89. stramineum Swt. 2 Hybrid	351
The linear-leaved Helianthemum.		The straw-colour-flowered Helianthemum.	
<i>Cistus linearis</i> Cav.		<i>Variety</i> 2	- 351
<i>Cistus pilòsus</i> Dec.		2 moltiplex Swt. 2	
75. virgatum Pers. 2 Barbary	- 347	90. diversifolium Swt 2	- 351
The twiggly Helianthemum.		The various-leaved Helianthemum.	
<i>Varieties</i> 2	- 347	<i>Variety</i> 2	- 351
albiflorum Dec. 2		2 moltiplex 2	
roseum Dec. 2			
76. apenninum Dec. 2 S. Europe	347	91. eriosépalon Swt. 2 S. Europe	351
The Apennine Helianthemum.		The woolly-sepaled Helianthemum.	
<i>Cistus apenninus</i> L.			
<i>Cistus hispídus</i> β Lam.			
<i>Varieties</i> 2	- 347	92. roseum Dec. 2 S. Europe	352
Leaves broad and flattish, Dec. 2		The Rose-colour-flowered Helianthemum.	
Leaves linear and narrow, Dec. 2		<i>Cistus roseus</i> All.	
		<i>Variety</i> 2	- 352
77. hispídum Dunal. 2 France	- 347	2 moltiplex Swt. 2	
The hispíd-herbage Helianthemum.			
<i>Cistus hispídus</i> Lam.			
<i>Cistus pilòsus</i> β Gouan.			
<i>H. majoranæfolium</i> β Dec.			
78. pulverulentum Dec. 2 France	348	93. foetidum Pers. 2 Hybrid	- 352
The powdered-leaved Helianthemum.		The fetid Helianthemum.	
<i>Cistus pulverulentus</i> Poir.		<i>Cistus foetidus</i> Jacq.	
<i>Cistus polifolius</i> Lam.			
79. macranthum Swt. 2	- 348	94. hyssopifolium Tenore. 2 Italy	352
The large-flowered Helianthemum.		The Hyssop-leaved Helianthemum.	
<i>Variety</i> 2	- 348	<i>Varieties</i> 2	- 352
2 moltiplex Swt. 2		1 crocæum Swt. 2	
		2 cùpreum Swt. 2	
		3 moltiplex Swt. 2 fig. 75.	
80. rhodanthum Dunal. 2 Spain	348	95. cùpreum Swt. 2 Hybrid	- 353
The red-flowered Helianthemum.		The copper-colour-flowered Helianthemum.	
<i>Cistus roseus</i> Jacq.			
<i>Cistus angustifolius</i> , formerly in the Royal			
Botanic Garden, Paris.			
<i>Cistus pituliferus</i> Thib.			
<i>Varieties</i> 2	- 348	96. venustum Swt. 2	- 353
oblongifolium Dec. 2		The handsome Helianthemum.	
subhirsutum Dec. 2			
càrneum Dec. 2		97. Milleri Swt. 2 Hybrid	- 353
		Miller's Helianthemum.	
81. canescens Swt. 2 Spain	- 349	98. majoranæfolium Dec. 2 S. Europe	353
The canescent-leaved Helianthemum.		The Marjoram-leaved Helianthemum.	
		<i>H. m. var. a</i> Dec.	
		<i>Cistus marjoranæfolius</i> Gouan.	
82. confusum Swt. 2 France	349	99. hirsutum Dec. 2 Pyrenees	- 353
The confused Helianthemum.		The hairy Helianthemum.	
<i>H. polifolium</i> Dec.		<i>Cistus hirsutus</i> Lapeyr., not of Lam.	
83. lanceolatatum Swt. 2 Hybrid	349	III. HUDSONIA L. 2	317. 354
The lanceolate-leaved Helianthemum.		THE HUDSONIA.	
84. polifolium Pers. 2 England	349	1. ericoides L. 2 N. Am. fig. 76.	354
The Polium-leaved Helianthemum.		The Heath-like Hudsonia.	
<i>Cistus polifolius</i> L.			
85. mutabile Pers. 2 Spain	- 350	2. [? e.] Nuttalli Swt. 2 N. America	354
The changeable-coloured-flowered Helianthemum.		Nuttall's Hudsonia.	
<i>Cistus mutabilis</i> Jacq.		<i>H. ericoides</i> Nutt.	
<i>Varieties</i> 2	- 350	3. [? e.] tomentosa Nutt. 2 N. America	354
With white flowers. 2		The tomento-e-leaved Helianthemum	
With smaller flowers of a rose red. 2		b	
With double rose-coloured flowers. 2			

Polygalàcææ. ■ □ ■ - 355

1. *Polygala Chamabáxus* L. ■ Europe fig. 77. 356
The Dwarf Box Polygala, or Box-leaved Milkwort.

Half-hardy Polygalas. ■ □ - 356

- P. oppositifolia* L. ■ □ major ■ (P.) *grandiflora* of some fig. 78.
- P. latifolia* Ker. ■ □
- P. myrtifolia* ■ □
- P. grandiflora* Lodd. ■ □
- P. bracteolata* L. ■ □
- P. speciosa* Bot. Mag. ■ □
- P. attenuata* Lodd. ■ □

Pittosporàcææ. ■ □ ■ □ 356

I. **BILLARDIÈ'RA** Sm. ■ □ - 356
THE BILLARDIERIA, or Apple Berry.

- 1. *longiflora* Labill. ■ □ V. D. L. f. 79. 357
The long-flowered Billardiera.
- 2. *ovàlis* Lindl. ■ □ V. Diemen's L. 357
The oval-leaved Billardiera.
- 3. *mutàbilis* H. K. ■ □ N. S. W. f. 80. 357
The changeable-coloured-flowered Billardiera.

II. **SO'LLYA** Lindl. ■ □ - 357
THE SOLLYA.
Billardièra, in one species.

- 1. *heterophýlla* Lindl. ■ □ N. II. f. 81. 357
The various-leaved Sollya.
- 2. *angustifolia* Lindl. ■ □ V. D. L. 358
The narrow-leaved Sollya.
Billardièra fusiformis Lab.

III. **PITTO'SPORUM** Banks. ■ □ 538
THE PITTO'SPORUM.

- 1. *Tobira* Ait. ■ □ China fig. 82. 358
The Tobira Pittosporum.
Euclypnus Tobira Thunb.
P. chinensis Donn.
Tobira Japan. Fr.
Chinesischer Klebsaame, Ger.
- 2. *undulàtum* ■ □ China fig. 83. 358
The undulated-leaved Pittosporum.

App. i. *Other Species of Pittosporum.* ■ □ 359

- P. revolútum* Ait. ■ □
- P. tomentosum* Bomp. ■ □
- P. fulvum* Rudge. ■ □
- P. hirtum* Willd. ■ □
- P. eriocarpum*. ■ □

App. I. *Pittosporàcææ probably half-hardy.*
■ □ - 359
Senecia nepalensis Dec. ■ □, *Bursaria spinosa* Cav. ■ □,
Cheiranthra linearis Lindl.

Caryophýllacææ. ■ 359

1. *Dianthus Caryophýllus* var. *fruticòsus* Hort. ■ - - 359
The shrubby Clove Pink.

Other suffruticosæ evergreen Caryophýllacææ.
■ - - 359

- Dianthus arbuscula* Bot. Reg. ■, *arboreus* L. ■ fig. 81.
fruticòsus L. ■, *suffruticosus* W. ■
- Silene fruticosa* L. ■
- Arenaria verticillata* W. ■
- Drypis spinosa* L. ■

Lanàcææ. ■ ■ □ - 360

- Lanum arbdreum* L. ■ □ fig. 86.
- L. taúricum* W. ■
- L. salsòfolius* Lam. ■
- L. suffruticosum* ■ □

Malvàcææ. ■ ? ■ ■ □ 362

I. **LAVATERA** L. ? ■ ■ □ - 360
THE LAVATERA, or Tree Mallow.
Lavatiere, Fr and Ger.

- 1. *marítima* Gouan ■ □ France f. 87. 360
The sea-side-inhabiting Lavatera.
L. hispanica Mill.
L. rotundifolia Lam.
- 2. *triloba* L. □ ■ Spain - 361
The three-lobed-leaved Lavatera.
- 3. *subovàta* Dec. ■ Morocco - 361
The subovate-leaved Lavatera.
- 4. *africàna* Cav. ■ □ N. Africa 361
The African Lavatera.
L. hispida var. Willd.

Other Species of Lavatera, likely to prove half-hardy. ■ □ - 361

- L. Pseudo-Olibia* Vair. ■ □
- L. phœnicea* Vent. ■ □
- L. Olibia* L. ■ □
- L. unguiculata* Desf. ■ □
- L. hispida* Desf. ■ □

II. **HIBISCUS** L. ■ - 360, 361
THE HIBISCUS.

Kelmie, Fr.
Eibisch, Ger.

1. *syriacus* L. ■ Syria - fig. 88. 362
THE SYRIAN HIBISCUS.

Althæa Frutic.
Kelmie des Jardins, Fr.
Syrischer Eibisch, Ger.

Varieties ■ - 362

- 2 *foliis variegatis* ■
- 3 *flòre variegato* ■
- 4 *flòre purpureo* ■
- 5 *flòre purpureo-plèno* ■
- 6 *flòre rubro* ■
- 7 *flòre albo* ■
- 8 *flòre albo-plèno* ■

App. I. *Other ligneous Plants of the Order Malvàcææ, which will probably be found hardy or half-hardy.* ■ □ - 363

- Sida pulchella* Bomp. ■ □ fig. 89.
- Abutilon pulchellum* Bot. Mag.

Stereuliàcææ. ■ □ - 633

Stereulia plantanifolia L. ■ □ fig. 90.

- S. lanceolata* Cav.
- S. diversifolia* G. Don. ■ □
- S. tomentosa* Thunb.
- S. peltata* G. Don.

Tiliàcea Lindl. † - Page 364

1. *TYLIA* L. † - - 364

THE LIME TREE.

- Lime Tree*, Gerard.
- Lind*, Anglo-Sax.
- Tilleul*, Fr.
- Lindc*, Ger. and Dutch.
- Tiglio*, Ital.
- Tilo*, Spain.
- Lipa*, Russ.

1. *europæa* L. † Europe pl. 14, 15. 364

The European, or common, Lime Tree.
T. intermedia Dec.
T. vulgaris Hayne.
T. europæa borealis Wahl.

Varieties. †

2 *microphýlla* - - 364

The small-leaved European Lime Tree.

- T. microphýlla* Vent.
- T. c. var γ* L.
- T. ulmifolia* Scop.
- T. sylvestris* Desf.
- T. parvifolia* Ehrh.
- T. cordata* Mill.
- Tilleul à petites Feuilles*, Fr.
- Kleinblättrige Linde*, *Winterlinde*, Ger.

? *Subvariety.* † - - 365
T. parvifolia glauca Hort. †

3 *platyphýlla* † - pl. 16. 365

The broad-leaved European Lime Tree.

- T. platyphýlla* Scop.
- T. cordifolia* Bess.
- T. europæa* Desf.
- T. grandifolia* Ehrh.
- Broad-leaved downy Lime Tree*.
- Tilleul à grandes Feuilles*, *Tilleul de Hollande*, Fr.

4 *rùbra* † - - 365

The red-twigged European Lime Tree.

- T. triflora* Puer.
- T. cornuthaca* Bosc.
- T. corallina* Hort. Kew.
- T. europæa β rùbra* Sibthorp.
- T. europæa γ* Smith Fl. Brit.
- T. grandiflora β* Smith Eng. Flor.

5 *laciniàta* - pl. 17. 366

The cut-leaved European Lime Tree.

T. platyphýlla laciniàta Hort.

6 *aúrea* † - pl. 18. 366

The golden-twigged European Lime Tree.

7 *platyphýlla aúrea* † - 366

The golden-twigged broad-leaved European Lime Tree.

8 *dasýstyla* † - - 366

The hairy-styled European Lime Tree.

T. dasýstyla Steven.

Other Varieties - 366

With variegated Leaves.

- T. a spleniifolia nová* Boll. Cat., a subvar. of *e. laciniàta*.
- e. aúrea*, a new var. of, *Bonn.*
- e. rubicaulis* Bot. Gard. Ant.

2. (cur.) *álba* Waldst. et Kit. † Hungary pl. 19. 372

The white-leaved European Lime Tree.

- T. americana* Du Roi.
- T. argéntea* Desf.
- T. rotundifolia* Vent.
- T. tomentosa* Mench.

? *Variety* † - - - 373

petiolàris Dec. †

The long-petioled-leaved Lime Tree.

3. *americana* L. † North America pl. 20. 372

The American Lime Tree.

- T. glàbra* Vent.
- T. caroliniana* Wangenh.
- T. canadensis* Michx.
- The smooth-leaved, or black, Lime Tree, and Brass Wood, Amer.

Varieties.

2 *laxiflòra* † - 374

The loose-cymed-flowered American Lime Tree.

T. laxiflòra Michx.

3 *pubescens* † - pl. 22. 374

The pubescent-leaved American Lime Tree.

- T. pubescens* Hort. Kew.
- T. caroliniana* Mill.
- T. americana* Walt.

4 *pubescens leptophýlla* † - 375

The thin-leaved pubescent American Lime Tree.

- T. pubescens leptophýlla* Vent.
- T. mississippiensis* Desf.

5 *heterophýlla* † - 375

The various-leaved American Lime-Tree.

- T. heterophýlla* Vent.
- T. álba* Michx.
- The White Lime.

App. i. *Doubtful Varieties of Tilia europæa and americana.* - - 176

T. platyphýlla vèra, *T. p. minor*, *T. præcox*, *T. vitifolia*, *T. laxiflòra microphýlla*, and *T. pubescens rugosa*.

App. I. *Other Species belonging to the Order Tiliàcea.* ■ □ ■ ▭ - 376

- Grèwia* L. ■ □ ■ ▭
- oppositifolia* Rozb. ■ □
- didyma*. ■
- biloba* G. Don. ■
- occidentàlis* L. ■ ▭
- populifolia* Vahl. ■
- pumila* Ham. ■
- velutina* Vahl. ■
- echinulata* Delile. ■

Ternströmiàcea. † † ■ ▭ 376

Sect. 1. *Hardy and half-hardy lignous Plants belonging to the Tribe Gordoniàcea.* 376

1. *MALACHODE'NDRON* Cav. † 377

THE MALACHODENDRON.
Stuàrta L'Hérit.
Steuàrta L.

1. *ovàtum* Cav. † N. Am. fig. 91. 377

The ovate-leaved Malachodendron.
Stuàrta pentagýnia L'Hérit.
Steuàrta Malachodéndron.
Stewartia à cinq Styles, Fr.

II. STUARTIA Cav. * - 377, 378
THE STUARTIA.

1. virginica Cav. * N. America f.92. 378
The Virginian Stuartia.
Stewartia Malachodendron L.
Stewartia virginiana Bot. Rep.
Stewartia à un Style, Fr.
Eingriffliche (one-styled) Stuartie, Ger.

III. GORDONIA Ellis. † 377, 378
THE GORDONIA.

1. Lasiánthus L. † N. Am. fig. 93. 379
The woolly-flowered Gordonia, or Loblolly Bay.
Hypéricium Lasiánthus L.
Gordonia à Feuilles glabres, Alécé de la Floride, Fr.
Langstielige Gordonie, Ger.

2. pubescens Ph. † Georgia f.94. 380
The pubescent Gordonia.
Lacathia florida Sal.
Franklinia americana Marsh.
The Franklinia, Amer.
Bchaarte Gordonie, Ger.

Varieties - - - 380

- velutina Dec.
G. pubescens L'Hér.
subglabra Dec.
G. Franklini L'Hér.
Franklinia Alhatamaha Marsh.

Sect. II. Hardy and half-hardy ligneous
Plants belonging to the Tribe Camelliæ.
* - - - 381

I. CAMELLIA L. * - 381
THE CAMELLIA.

The Japan Rose.
Camellier, Rose du Japon, et de la
Chine, Fr.
Camellie, Ger.

1. japonica L. * Japan fig.95. 382
The Japanese Camellia, or Red single-flowered
Camellia.

Varieties.

Chinese in general Cultivation.

- * - - - 383
- 2 variegata Bot. Rep. * - - -
3 incarnata Bot. Rep. * - - -
4 alba plena Bot. Rep. * - - -
5 fimbriata Lodd. * - - -
6 rubra plena Bot. Rep. * - - -
7 anemoneflora * - - - fig. 96.
8 crassinervis Lodd. * - - -
9 myrtifolia Bot. Mag. * - - -
10 involuta Bot. Reg. * - - -
11 variabilis * - - -
12 Pomponia Bot. Reg. * - - -
Subvarieties * - - -
 pawoniceflora rosea * - - -
 alba * - - -
 pallida * - - -
13 semiduplex Bot. Rep. * - - -
14 atro-rubens Bot. Cab. * - - -
15 Welbánkii Chandl. Ill. * - - -
 lateo-albicans Bot. Reg.
 flavescens.
16 rosea Don's Mill. * - - -

- 17 speciosa Hort. Trans. * - - -
 C. Rawesiana Hort.

- 18 carnea Bot. Reg. * - - -
19 imbricata Hort. Trans. * - - -
20 Parksii Hort. Trans. * - - -
21 Sabiniána Hort. Trans. * - - -

Chinese and other Foreign Varieties
not in general Cultivation, but in
all probability as hardy as the
others. * - - - 3-6

- 22 candidissima * - - -
23 Donklæri * - - -
24 francofurtensis * - - -
25 hybrida Makoy. * - - -

Other Names of Foreign Varieties.
* - - - 386

argentea, Gunnelli, Pronayana, violacea su-
perba.
The semidouble white, the rose-coloured
waratah, hexangularis.





Varieties originated in Britain.

- * - - - 386
- 26 corallina Chandl. * - - -
27 eximia Chandl. * - - -
28 Wiltoni Chandl. * - - -
29 Chándleri Chandl. * - - -
 versicolor Bot. Reg.
30 flore-albo Chandl. * - - -
31 althææflora Chandl. * - - -
32 anemoneflora alba Chandl. * - - -
33 Woodsii Chandl. * - - -
34 punctata Chandl. * - - -
35 elegans Chandl. * - - -
36 florida Chandl. * - - -
37 rosea Chandl. * - - -
38 celipsis Chandl. * - - -
39 insigne Chandl. * - - -
40 alba semiduplex Chandl. * - - -
41 concinna Chandl. * - - -
42 splendens Chandl. * - - -
 coccinea Hort. Brit.
43 Røsa sinensis Lodd. * - - -
44 Røssii Chandl. * - - -
45 Aitonii Chandl. * - - -
46 epsomensis Chandl. * - - -
47 Colvilli Swt. * - - -
48 Sweetiana Swt. * - - -
49 Knightii Lodd. * - - -
50 Røsa mundi Don's Mill. * - - -
51 Præssii Don's Mill. * - - -
52 rubro-punctata * - - -
53 Elphinstonia * - - -
54 single-striped and dotted * - - -
55 acubæfolia Hort. Brit. * - - -
56 expansa Hort. Brit. * - - -
57 Susanna Gard. Mag. * - - -
58 Martha Gard. Mag. * - - -
59 Waheana Gard. Mag. * - - -

Varieties mentioned in Gard. Mag.
without any Description being
given. - - - 3-9



Cliveana, conspecta, decora, princeps, ro-
tundifolia, Palmieri, Reevesii, longifolia,
Dorsetii, pendula, Allmuta alba, superba.



2. reticulata Lindl. * - - - China - 389
The reticulate-leaved Camellia, or Capt. Rawes's
Camellia.
3. maliflora Lindl. * - - - China f.97. 390
The Apple-blossom-flowered Camellia.
C. Sasánqua Bot. Mag.
C. Sasánqua rosea Hort.
Palmer's double Sasanqua.
4. Sasánqua Thunb. * - - - China f.98. 390
Sasanqua, or Lady Banks's, Camellia.
Sasánqua Kämpf.
Cha-tcha, Chinese.
5. Kíssi Wall. * - - - Nepal fig.99. 390
The Kissi Camellia.
C. Keina Hamil. MSS.


6. *oleifera* *Abel.*   Cochin-China Page
fig. 100. 391
The oil-bearing Camellia.
7. *euryoides* *Lindl.*   China fig. 101. 391
The Eurya-like Camellia.
Thèa euryoides Booth.



II. *THEA* *L.*  - - 381. 392
THE TEA TREE.


1. *viridis* *L.*   China - fig. 102. 393
The common, or green, Tea.
T. Bohèa stricta Ait.
T. sinensis Sims.
T. chinensis var. α *viridis* Dec.
Camèllia viridis Link.
Thèa cantonensis Lam.

2. *Bohèa* *L.*   China fig. 103. 393
The Bohea, or black, Tea.
T. chinensis β *Bohèa* Sims.


3. *cochinchinensis* *Lour.*   Cochin-China - 395
The Cochin-China Tea.

4. *oleosa* *Lour.*   China - 395
The oily Tea.







- Anticipated Ternströmiaceæ.* ?  - 395
Eurya acuminata *Royl.* ? 







Aurantiaceæ.  - 395

- I. *CITRUS*  - fig. 104. 395



- II. *LIMONIA*  - 396
L. Laurèla *Dec.*
L. citrifolia *Willd.*
L. parviflora *Bot. Mag.*
L. australis.



Hypericaceæ *Lindl.* 362


         






- I. *HYPERICUM* *L.*                  - 397
THE ST. JOHN'S WORT.
Fuga Dæmonum.
Mille Pertuis, Fr.
Johanniskraut, Ger.

§ i. *Ascyrella* *Chois.* - 398



1. *elatum* *Ait.*   N. America 398
The tall St. John's Wort.



2. *hircinum* *L.*   Mediterranean. fig. 105 398



The Goat-scented St. John's Wort.
Tràgium Clus.
Androsæmum scetidum Bauh.
Mille Pertuis à Odeur de Bouc, Fr.
Varieties  - - 398



- 2 obtusifolium *Dec.*     



- 3 minus *Wats.* 







































3. *grandiflorum* *Chois.*   Teneriffe Page
298
The large-flowered St. John's Wort.
H. canariense Willd., not of L.



4. *foliosum* *Ait.*   Azores - 398
The leafy St. John's Wort.
Shiàng St. John's Wort.



5. *floribundum* *Ait.*   Canaries 398
The abundant-flowered St. John's Wort.
H. frutescens Comm.
Many-flowered St. John's Wort.



6. *olympicum* *L.*   Mount Olympus fig. 106. 399
The Olympian St. John's Wort.
H. montis Olympi Wheel.
H. orientalis flore majus Tourm.



7. *canariense* *L.*   Canaries - 399
The Canary Island St. John's Wort.



Varieties   - - 399
triphýllum *Dec.*                                    



8. *chinense* *L.*   China - 399
The Chinese St. John's Wort.
H. monogynum Mill.
H. aureum Lour.



9. *cordifolium* *Chois.*   Nepal 399
The heart-leaved St. John's Wort.
H. bracteatum.
H. Lungisium Ham. MSS.

10. *pátulum* *Thun.*   Japan - 399
The spreading St. John's Wort.



11. *Kalmianum* *Lam.*   N. America 400
Kalm's St. John's Wort.
H. Bartramium Mill.
Virginia St. John's Wort.



12. *Urðum* *Ham.*   Nepal fig. 107. 400
The Urala St. John's Wort.



13. *calycinum* *L.*   Britain - 400
The large-calyxed St. John's Wort.
Androsæmum constantinopolitanum flore miximo Wheeler.
The large-flowered St. John's Wort, The large-flowering Tutsan, The terrestrial Sun, Aaron's Beard.
Mille Pertuis à grands Fleurs, Fr.
Grossblumiger Johanniskraut, Ger.



14. *balæricum* *L.*   Majorca - 400
The Majorca St. John's Wort.



§ ii. *Perforaria* *Chois.* - - 401

15. *prolificum* *L.*   N. America 401
The prolific St. John's Wort.
H. foliosum Jacq.
H. Kalmianum Du Roi.



16. *heterophyllum* *Vent.*   Persia 401
The various-leaved St. John's Wort.

17. *ægyptiacum* *L.*   Egypt 401
The Egyptian St. John's Wort.

18. *galioides* *Lam.*   N. America 401
The Galium-like-leaved St. John's Wort.

19. *axillare* *Lam.*   N. America 401
The axillary-flowered St. John's Wort.
H. fasciculatum Willd.
H. Còris Walt.

20. *glandulosum* *Ait.*   Madeira 402
The glandular St. John's Wort.

21. *serpyllifolium* *Lam.*   Levant 402
The Wild-Thyme-leaved St. John's Wort.

22. *empetrifolium* Willd. α — S. Europe ^{Page} 402
fig. 108. The Empetrum-leaved St. John's Wort.
23. *Còris L.* α — Levant. 402
The *Còris*-leaved St. John's Wort.
24. *ericoides L.* α — Spain 402
The Heath-like St. John's Wort.
§ iii. *Bràthya* Chois. α — 402
25. *fasciculatum Lam.* α — Carolina 402
The fascicled-leaved St. John's Wort.
II. *asplathoides* Pursh.
- App. i. *Other Species of Hypéricum.* 403
II. *repens*, *H. linearifolium*, *H. oblongifolium*, and
H. japonicum Dec.
- II. **ANDROSÆMUM** Chois. α 397. 403
THE ANDROSÆMUM, or *Tutsan*.
Hypéricum L.
Androsæmum, Fr.
Johanniskraut, Ger.
1. *officinàlis Allioni.* Levant α f. 109. 403
The officinal Androsæmum, or common *Tutsan*.
Clyconum Italorum L'Obel.
Hypéricum Androsæmum L.
Park Leaves.
Androsæmum officinale, Fr.
Breit-blättriges (broad-leaved) Johanniskraut, Ger.
-
- Acerdææ* Lindl. γ δ ζ — η — 404
- I. **ACER** L. γ δ ζ — η — 405
THE MAPLE.
E'rablè, Fr.
Ahorn, Ger.
Accro, Ital.
Arec, Spanish.
1. *oblòngum Will.* ζ — Nepal fig. 113.
in p. 433. - - - 405
The oblong-leaved Maple.
A. laurifolium D. Don.
A. Buzimpala Hamilt.
2. *tatàricum L.* ζ Tartary pl. 23. f. 114.
in p. 134. - - - 406
The Tartarian Maple.
E'rablè de Tartarie, Fr.
Tartarische Ahorn, Ger.
Zarza-modon, or *Locust Tree*, Russ.
3. *spicatum L.* ζ North America pl. 24.
fig. 115. in p. 435. - - - 406
The spike-flowered Maple.
A. montanum Ait.
A. pennsylvanicum Du Roi.
A. parryflorum Ehrh.
Mountain Maple.
E'rablè de Montagne, Fr.
Berge Ahorn, Ger.
4. *striatum L.* ζ North America pl. 25.
fig. 116. in p. 436, 437. - - - 407
The striped-barked Maple.
A. pennsylvanicum L.
A. canadense Marsh.
Snake-barked Maple, *Moose Wood*, *Striped Maple*, *E'rablè jaspé*, Fr.
Gestreifter Ahorn, Ger.
5. *macrophyllum Pursh.* ζ North America ^{Page} 408
pl. 26. fig. 117. in p. 438, 439., and
118. in p. 440, 441. - - - 408
The long, or large, leaved Maple.
6. *platanoides L.* ζ Europe pl. 27.
fig. 119. in p. 442, 443. - - - 408
The Platanus-like, or *Norway Maple*,
E'rablè blanc, *E'rablè de Norvège*, Fr.
Spitz-Ahorn, *Spitz-blättriger Ahorn*, Ger.
- Varieties.* ζ
- 2 *Lobèlii* ζ f. 120. in p. 444. 409
Lobel's Platanus-like Maple.
A. Lobèlii Tenore.
A. platanoides Don's Mill.
- 3 *pubescens Hayne.* ζ - - - 409
The downy-leaved Platanus-like Maple.
- 4 *variegatum Hort.* ζ - - - 409
The silver variegated-leaved Platanus-like Maple.
albo variegatum Hayne.
- 5 *aureo variegatum* ζ - - - 409
The golden variegated-leaved Platanus-like Maple.
- 6 *laciniatum Dec.* ζ pl. 28. fig.
121. in p. 445. - - - 409
The cut-leaved Platanus-like Maple.
? A. p. crispum Lanth.
The Eagle's Claw, or *Hawk's Foot Maple*.
7. *saccharinum L.* ζ North America pl.
29. fig. 122. in p. 446, 447. 411
The Sugar Maple.
Rock Maple, *Hard Maple*, *Bird's-Eye Maple*.
- Variety* ζ - - - 411
- 2 *nigrum* ζ
The black Sugar Tree, or *Rock Maple*.
A. nigrum Michx.
8. *Pseudo-Platanus L.* ζ Europe pl. 30,
31. fig. 123. in p. 448, 449. 414
The Mock Plane Tree, the *Sycamore*, or *Great Maple*.
Plane Tree, Scotch.
E'rablè Sycamore, *E'rablè blanc de Montagne*, *Fausse Platane*, *Grand E'rablè*, Fr.
Ehrenbaum, *Weisser Ahorn*, *Gemeine Ahorn*, Ger.
- Varieties.*
- 2 *flava variegata* ζ - - - 414
The yellow variegated Sycamore, or *Cotswoldine Plane*, with leaves variegated with yellow.
- 3 *albo variegata Hayne* ζ - - - 414
The white variegated-leaved Sycamore.
- 4 *purpurea Hort.* ζ - - - 415
The purple-leaved Sycamore.
- 5 *subobtusà Dec.* ζ - - - 415
The half-obtusè-leaved Sycamore.
A. opulifolium Thuil.
A. vitifolium Opiz.

	Page		Page
<i>Other Varieties</i> - - -	415	<i>Varieties</i> † - - -	424
Hodgkins's Seedling, Hort.		2 coccineum †	
Leslie's Seedling, Hort.		3 intermedium †	
stenóptera Hayne.		17. monspessulanum L. † South Europe	
macroptera Hayne.		pl. 39, 40. fig. 131. in p. 458.	427
microptera Hayne.		The Montpellier Maple.	
		A. trilobum Mench.	
9. obtusatum Kit. † Hungary pl. 32.	420	A. trifolium Duh.	
fig. 124. in p. 450, 451.		A. trilobatum Lam.	
The obtuse-lobed-leaved Maple.		E'rablc de Montpellier, Fr.	
A. neapolitanum Tenore.		Französischer Ahorn, Ger.	
A. hybridum Hort.		18. campestre L. † Europe pl. 41, 42.	
The Neapolitan Maple.		fig. 132. in 458. - - -	428
10. barbatum Michx. † North America		The common, or Field, Maple.	
pl. 33. fig. 125. in p. 452.	420	E'rablc champêtre, Fr.	
The bearded-calyced Maple.		Kleiner Ahorn, Feld Ahorn, Ger.	
A. carolinianum Bolt.		<i>Varieties.</i> †	
A. trilobatum Hort.		1 hebecarpum Dec. † - - -	428
11. Opalus Dec. † Italy pl. 34. fig. 126.		The downy-fruited Field Maple.	
in p. 453. - - -	421	A. campestre Wallr.	
The Opal, or Italian, Maple.		A. mollis Opiz.	
A. Opalus L.		2 foliis variegatis † - - -	428
A. rotundifolium Lam.		The variegated-leaved Field Maple.	
A. italicum Lanth.		A. Pseudo-Platanus.	
A. villosum Pres.		3 collinum Wallr. † - - -	428
L'E'rablc Opale, E'rablc à Feuilles rondes,		The hill-inhabiting Field Maple.	
E'rablc d'Italie, Fr.		A. affine and A. macrocarpum	
		Opiz.	
<i>Variety</i> † - - -	421	4 austriacum Tratt. Arch. † 428	
? A. coriaceum Lodd. †		The Austrian Field Maple.	
12. opulifolium Vill. † Pyrenees - - -	421	19. creticum L. † Crete pl. 43. fig.	
The Guelder-Rose-leaved Maple.		132. in p. 459. - - -	430
A. hispanicum Pour.		The Cretan Maple.	
A. vernum Reyn.		A. heterophyllum Willd.	
A. montanum C. Bauhin.		A. sempervirens L.	
E'rablc duret, E'rablc à Feuilles d'Obier,		E'rablc de Crète, Fr.	
Ayart in Dauphiné, Fr.		Cretischer Ahorn, Ger.	
Schnecboll-blättriger Ahorn, Ger.		App. i. Doubtful Species of Acer. - 430	
13. circinatum Pursh. † North America		A. Opalus, A. opulifolium, A. barbatum, A. cori-	
fig. 112., and fig. 127. in p. 454. 422		aceum, A. lobëlii, A. ulgrum, A. platanoides, A.	
The round-leaved Maple.		sacharinum, A. hybridum, A. obtusifolium, A.	
		obtusatum, A. lobatum.	
14. palmatum Thunb. ㄩ Japan fig.		App. ii. Anticipated Species of Acer. † 431	
128. in p. 455. - - -	422	A. ibericum Bieb. †	
The palmate-leaved Maple.		A. obtusifolium Sibth. †	
		A. parvifolium Tausch. †	
		A. parvifolium of some,	
		A. monspessulanum of others.	
15. eriocarpum Michx. † N. Amer. pl.		A. glabrum Torrey. †	
35, 36. fig. 129. in p. 456. - - -	423	A. laevigatum Wall. †	
The hairy-fruited, or white, Maple.		A. acuminatum Wall. †	
A. dasycarpum Willd.		A. cultratum Wall. †	
A. tomentosum Hort. Par.		A. caudatum Wall. †	
A. glaucum Marsh.		A. villosum Wall. †	
A. virginianum Duh.		A. sterculiaceum Wall. †	
A. rubrum Wagenh.		A. dissectum Thun. †	
White, or soft, Maple, U. S.		A. japonicum Thun. †	
Sir Charles Wager's Maple.		A. septenlobum Thun. †	
E'rablc à Fruits cotoneux, E'rablc blanc,		A. pictum Thun. †	
Fr.		A. trifidum Thun. †	
Rauher Ahorn, Ger.		A. truncatum Bunge. †	
16. rubrum L. † N. America pl. 37, 38.		App. iii. Half-hardy Species of Acer. 432	
fig. 130. in p. 457. - - -	424	† ㄩ ㄩ ㄩ	
The red-flowering, or scarlet, Maple.		A. palmatum Thun. ㄩ	
A. virginianum Herm.		A. oblongum Wall. † ㄩ	
A. coccineum Ait.		II. NEGUNDO Mench. † 405. 460	
Soft Maple, Swamp Maple, Red Maple.		THE NEGUNDO, or Box Elder.	
E'rablc rouge, Fr.		A'cer L.	
Rother Ahorn, Ger.		Negundium Rafin.	

- I. fraxinifolium Nutt.** † North America Page
460
pl. 44. 460
- The Ash-leaved Negundo.
Acer Negundo L.
N. aceroides Muench.
Negundium americanum Rafin.
The Ash-leaved Maple, The black Ash.
L'érable à Feuilles de Frêne, Fr.
L'érable à Giguères, Illinois.
Eschenblättriger Ahorn, Ger.
- Variety † 460
2 *crispum G. Don.* † - pl. 45.
The curled-leaved Ash-leaved Negundo, or
Box Elder
- App. i. *Anticipated Species of Negundo.* † 462
- N. mexicanum Dec.* †
A. ternatum Moc.
? *N. fraxinifolium var. Dec.*
N. cochinchinense Dec. †
- App. I. *Other Aceræcæ.* † - 462
Dobinæa vulgaris Hamilt. †
- Æsculæcæ Lindl.** † † - 462
Castanæcæ Link.
Hippocastanæ Dec.
- I. ÆSCULUS L.** † - 462
THE HORSECHESTNUT.
Hippocastanum Tourn.
Marronnier d'Inde, Fr.
Rosskastanie, Ger.
- J. *Hippocastanum L.* †
pl. 46, 47, 463
- The common Horsechestnut.
Hippocastanum vulgare Tourn.
Marronnier d'Inde, Fr.
Gemeine Rosskastanie, Ger.
- Varieties. †
2 *flore pleno* † - 463
The double-flowered Horsechestnut.
3 *variegata* † - 463
The variegated-leaved Horsechestnut.
2. (H.) *ohioënsis Michx.* † N. Am. 467
The Ohio Æsculus, or Horsechestnut.
3. (H.) *rubicunda Lois.* † N. America
pl. 48, 467
The reddish-flowered Æsculus, or Horsechestnut.
Æ. carnea Hort.
Æ. rosea Hort.
Æ. coccinea Hort.
Æ. Hippocastanum var. rubicundum Schu-
hart
Whitley's fine scarlet.
4. (H.) *glabra Willd.* † North America
fig. 133. 467
The smooth-leaved Æsculus, or Horsechestnut.
5. (H.) *pallida Willd.* † N. Am. f. 134. 468
The pale-flowered Æsculus, or Horsechestnut.
Gebliche Rosskastanie, Ger.
6. (H.) *Lyoni Hort.* † 468
Lyon's Æsculus.
- Whitley's fine scarlet
Æ. americana
Æ. H. incisum
Æ. H. pubescens
Æ. H. tortuosum
Æ. H. nigrum
- II. PAVIA Boerh.** † † 462, 469
THE PAVIA, or Smooth-fruited Horsechestnut
Tree.
Pavier, Fr.
- I. rubra Lam.** † N. Am. pl. 49, 50. 469
The red-flowered Pavia.
Æsculus Pavia L.
Æsculus Pavia var. α rubra Hayne.
P. parviflora Hort.
Small Buckeye, Amer.
Marronnier Pavie, Pavie à Fleurs rouges,
Fr.
Rothe Rosskastanie, Ger.
- Varieties. †
2 *arguta G. Don* † fig. 135. 469
The sharp-toothed-leaved red-flowered
Pavia.
Æsculus Pavia parviflora Hort.
- 3 *sublaciniata Wats.* † - 470
The slightly cut-leaved red-flowered
Pavia.
- 4 *humilis* † - pl. 51. 470
The dwarf red-flowered Pavia.
P. humilis G. Don.
Æsculus humilis Lodd.
- 5 *humilis pendula* † - 470
The pendulous-branched dwarf
red-flowered Pavia.
2. *flava Dec.* † Carolina pl. 52, 53. 471
The yellow-flowered Pavia.
Æsculus flava Ait.
Æ. litca Wangh.
P. litca Poir.
The large Buckeye, Big Buckeye, Amer.
The yellow Pavia.
3. *discolor Swt.* † N. America - 472
The two-coloured-flowered Pavia.
Æsculus discolor Ph.
4. *hybrida Dec.* † Hybrid - 472
The hybrid Pavia.
Æsculus hybrida Dec.
5. *neglecta G. Don.* † fig. 136. 472
The neglected Pavia.
Æsculus neglecta Lindl.
6. *macrocarpa Hort.* † pl. 54. 473
The long-fruited Pavia.
Æsculus Pavia macrocarpa Lodd.
Pavia macrocarpa Hort.
7. *macrostachya Lois.* † North America
fig. 137. 473
The long-racemed Pavia.
Æsculus parviflora Walt.
Æ. macrostachya Mx.
Pavia alba Poir.
Pavia ciliata Voit.
Pavier à longs Épis, Pavier nain, Fr.
Langährige Rosskastanie, Ger.
- App. i. *Other Varieties of Pavia.* - 474
P. serrata, P. erecta, Æsculus Pavia flava var.
Hort.

	Page
<i>Sapindàcæa.</i> ☿ ■ □ ■ ⊔	474
I. KÖLREUTERIA Laxm. ☿ -	475
THE KÖLREUTERIA.	
<i>Sapindus</i> sp. L. fil.	
1. paniculata Laxm. ☿ China pl. 55.	475
The panicked-flowering Kolreuteria.	
<i>Sapindus chinensis</i> L. fil.	
<i>K. paullinoides</i> L'Herit.	
<i>Savonnier paniculé</i> , Fr.	
<i>Rispentragende Kolreuterie</i> , Ger.	
App. I. Half-hardy ligneous Species of Sa-	
pidàcæa. ■ □ ■ ⊔	- 476

- Dodonea viscosa* L. ■ ⊔
- D. alternata* Cunning.
- D. jamaicensis* Dec. ■ □
- D. angustifolia* Swz.
- D. viscosa* Cav.
- D. salicifolia* Dec. ■ ⊔
- D. angustifolia* Lam.
- Bols de reinette*, Fr.
- D. laurina* Sieb. ■ ⊔
- D. triquetra* Bot. Rep. ■ ⊔
- D. cuneata* Smith. ■ ⊔
- D. asplenifolia* Rudge. ■ ⊔

Meliàcæa. ☿ ⊔ - 476

- Melia** 1. *Azederach* L. ☿ ⊔ - fig. 138. 476
- The Bead Tree, or Indian Lilac.
- 2. *sempervirens* Swz. ☿ ⊔ - 476
- The evergreen Melia, or Bead Tree.
- 3. *australis* Smt. ☿ ⊔ - 477
- 4. *japonica* G. Don. ☿ ⊔ - 477

Vitacæa Lindl. ■ ■ □ - 477

- I. VITIS** L. ■ ■ □ - 477
- THE GRAPE VINE.
- Giud.* Celtic.
- Vid.* Span.
- Vigne.* Fr.
- Wein.* Ger.
- 1. vinifera** L. ■ Persia - fig. 139. 477
- The wine-bearing Vine.
- Vigne.* Fr.
- Gemeiner Weinstock.* Ger.
- Varieties* ■ - 478
- 2 foliis incanis** ■ - 478
- The hoary-leaved Grape Vine.
- Miller's Grape.* or *Miller's Black Cluster Grape.*
- 3 foliis rubescéntibus** ■ 478
- The rubescent-leaved Grape Vine.
- The Claret Grape.*
- Tenturier.* Fr., not *Clairette*, *Duham.*
- 4 apiifolia** Hort. ■ fig. 140. 478
- The Parsley-leaved Grape Vine.
- Crotal.* Fr.
- V. laciniata* L.
- 2. Labrusca** L. ■ N. Am. fig. 141. 479
- The wild Vine, or *Fox Grape.*
- V. taurina* Walt.
- Filsiger Wein.* Ger.

- 3. æstivalis** Michx. ■ N. Am. fig. 142. 479
- The Summer Vine, or *Grape Vine.*
- V. Labrusca* Walt.
- V. vulpina* Willd.
- 4. sinuata** G. Don. ■ N. America - 479
- The scallop-leaved Vine, or *Summer Grape Vine.*
- V. æstivalis* var. *sinuata* Ph.
- ? *V. labruscoides* Muhl.
- 5. cordifolia** Michx. ■ N. Am. f. 143. 480
- The heart-shape-leaved Vine, or *Chicken Grape.*
- V. incisa* Jacq.
- V. vulpina* L.
- 6. riparia** Michx. ■ N. Am. f. 144. 480
- The river-side, or *sweet-scented*, Vine.
- V. odoratissima* Donn.
- Vigne de Baturcs.* Amer.
- 7. rotundifolia** Michx. ■ N. Am. 480
- The round-leaved Vine, or *Bullet Grape.*
- 8. caribæa** Dec. ■ □ Jamaica - 480
- The Caribbean Vine.
- V. indica* Swz.
- App. i. Other hardy or half-hardy Species of**
- Vitis.** ■ - 481
- V. Wällichii* Dec. ■
- V. glabrata* Roth. ■ □
- V. indica* L. ■ □ fig. 145.
- V. caesia* Hort. Trans.

II. AMPELOPSIS Michx. ■ 477. 481

- THE AMPELOPSIS.
- Vitis* sp.
- Cissus* sp.
- 1. cordata** Michx. ■ N. America 481
- The cordate-leaved Ampelopsis.
- Cissus Ampelopsis* Pers.
- Vitis indivisa* Willd.
- 2. hederacea** Michx. ■ North America
- fig. 146. 482
- The Ivy-like Ampelopsis, or *Five-leaved Ivy.*
- Hedera quinquefolia* L.
- Vitis quinquefolia* Lam.
- Cissus hederacea* Ph.
- Cissus quinquefolia* Hort.
- Vitis hederacea* Willd.
- A. quinquefolia* Hook.
- Vigne Vierge.* Fr.
- Jungfern Reben.* Wilder Wein, Ger.
- 3. (H.) hirsuta** Donn. ■ N. America 482
- The hairy-leaved Ampelopsis.
- Cissus hederacea* var. *hirsuta* Ph.
- 4. bipinnata** Michx. ■ N. America 482
- The bipinnate-leaved Ampelopsis.
- Vitis arborea* Willd.
- Cissus stans* Pers.
- App. i. Anticipated hardy Species of Ampe-**
- lopsis.** ■ - 482
- A. hötuya* Dec. ■
- A. heterophylla* Blume. ■
- Vitis javanica* Spreng.
- A. capreolata* G. Don. ■
- Vitis capreolata* D. Don.

III. CISSUS L. ■ ⊔ - 477. 483

- THE CISSUS.
- 1. orientalis** Lam. ■ ⊔ Levant - 485
- 2. quinata** Ait. ■ ⊔ Cape of Good Hope - 485
- 3. antarctica** Vent. ■ ⊔ New Holland - fig. 147. 485
- 1. capensis** Willd. ■ ⊔ Cape of Good Hope - 485
- 5. ritiginea** ■ ⊔ - fig. 148. 485
- 6. quinquefolia** ■ ⊔ - fig. 149. 485

Page

Geraniaceæ. ̳ ̳ - 483

Pelargonium Harringtonii ̳ ̳, cucullatum ̳ ̳ fig. 150., macranthou ̳ ̳, megalanthon ̳ ̳, calaministratum, quercifolium ̳ ̳, peltatum ̳ ̳, zonale ̳ ̳ fig. 151., Benticianum ̳ ̳, inquinans ̳ ̳ fig. 152.

Zygophyllaceæ. ̳ ̳ ̳ 484

Melanthus major L. ̳ - - - fig. 155. 484
Zygophyllum sessilifolium L. ̳ ̳ - - - fig. 151. 484
Morgiana L. ̳ ̳ - - - - - 484

Rutaceæ. ̳ ̳ ̳ ̳ 481

I. RUTA L. ̳ ̳ ̳ ̳ - 484

THE RUE.
Ruc, Fr.
Raute, Ger.

1. graveolens L. ̳ S. Europe f. 155. 485
The heavy-scented, or common, Rue.
R. hartensis Mill. Diet.
Ruc, Fr.
Gartenraute, Ger.

App. i. Half-hardy Species of Ruta. 487

R. pinnata L. ̳ ̳, R. bracteosa Dec. ̳ ̳, R. angustifolia Pers. ̳ ̳ fig. 156., R. macrophylla Sol. ̳ ̳, R. montana Clus. ̳ ̳, R. divaricata Tenore ̳ ̳, R. corsica Dec. ̳ ̳, R. albiflora Hook. ̳ ̳

II. APLOPHYLLUM Andr. Juss. ̳ ̳ - 484. 487

THE APOPHYLLUM, or Simple-leaved Rue.

1. linifolium G. Don. ̳ ̳ Spain fig. 157. 487
Ruta linifolia L.

2. suaveolens G. Don. ̳ ̳ Tauria 487
Ruta suaveolens Dec.

3. fruticosum G. Don. ̳ ̳ Syria 487
Ruta fruticulosa Lab.

Xanthoxylaceæ. ̳ ̳ - 487

I. XANTHOXYLUM L. ̳ ̳ 487, 488
THE XANTHOXYLUM, or Toothache Tree.

Xanthoxylum of many.
Kaupmannia Rafin.
Clavaler, Fr.
Zahnwechholz, Ger.

1. fraxineum Willd. ̳ North America pl. 6. fig. 158. 488

The Ash-leaved Xanthoxylum, or Common Toothache Tree.

Xanthoxylum ramiflorum Michx.
Z. Clava Herculis var. I.
Z. americanum Mill.
Z. caribæum Gært.
Clavaler à Feuilles de Frêne, Fr.
Eschen-blüttrages Zahnwechholz, Ger.

Variety ̳ - - - 488

2 virginicum ̳
X. virginicum Lodd

2. tricarpum Michx. ̳ N. America 488
The three fruited Xanthoxylum, or Toothache Tree
Fagara fraxinifolia Lam.

3. mite Willd. ̳ North America - 489
The smooth, or thornless, Xanthoxylum, or Toothache Tree.

App. i. Half-hardy Species. 489

II. PTELEA L. ̳ ̳? ̳ - 487. 489

THE PTELEA, or Shrubby Trefoil.
Bellucia Adans.
Orme de Samarie, Fr.
Lederblume, Ger.

1. trifoliata L. ̳ N. Amer. pl. 57. 489
The three-leafed-leaved Ptelea, or Shrubby Trefoil.

Orme de Samarie à trois Feuilles, Fr.
Dreyblättrige Lederblume, Ger.

Varieties ̳ - - 489
2 pentaphylla Munchh. ̳
3 pubescens Pursh. ̳

App. i. Other ligneous Species of Ptelea, hardy or half-hardy. ̳ - 490

P. monophylla Lam. ̳? ̳
P. pentandra Moc. ̳? ̳
P. ovata Lour. ̳? ̳

III. AILANTHUS Desf. ̳ 487. 490

THE AILANTO.
Rhus Ehrh
Verne du Japon, Fr.
Götterbaum, Ger.

1. glandulosa Desf. ̳ China pl. 58, 59. fig. 159. 490

The glandulous-leaved Ailanto.
A. procera Sal.
Rhus hypoleudendron Mœneh.
R. cacodendron Ehrh.
R. sinense Ellis.
Aylante glanduleux, Fr.
Drüsiger Gotterbaum, Ger.

Coriaceæ. ̳ - 492

I. CORIARIA Niss. ̳ - - 492

THE CORIARIA.
Redoul, Fr.
Gerberstrauch, Ger.

1. myrtifolia L. ̳ Europe f. 160. 492
The Myrtle-leaved Coriaria.

Fustet des Corroyeurs, Redoul à Feuilles de Myrte, Fr.
Myrtenblüttriger Gerberstrauch, Ger.

2. nepalensis Wall. ̳ Nepal - 493
The Nepal Coriaria.

3. microphylla Poir. N. Zealand ̳? ̳ 493
The small-leaved Coriaria.
C. armentosa Forst.

Staphyleaceæ ̳ - 493

I. STAPHYLEA L. ̳ - 493

THE STAPHYLEA, or Bladder-nut Tree.
Staphylodendron Tourn.
Staphilier, Faux Pistachier, Fr.
Pompennuss, Ger.

1. trifolia L. * N. Am. fig. 161, 162. 493
The three-lobed-leaved Staphylea.
Staphilier à Feuilles ternées, Fr.
Virginsche Pimpernuss, Ger.
2. pinnata L. * Europe - fig. 163. 494
The pinnated-leaved Staphylea.
Staphylodendron pinnatum Ray.
Staphilier à Feuilles ailées, Fr.
Gemeine Pimpernuss, Ger.

App. i. Anticipated Species of *Staphylæa*.
 * ? ┘ - 495

- S. Bumalda Dec. * ? ┘
S. heterophylla R. et P. * ? ┘

Celastrææ Dec. † * * - 495

1. EUONYMUS Tourn. † * * 495, 496

THE EUONYMUS, or Spindle Tree.
Fusain, Bonnet de Prêtre, Bois à Lardoire, Fr.
Spindelbaum, Ger.

1. europæus L. † E. pl. 60. f. 164. 496
The European Euonymus, or Spindle Tree.

E. vulgaris Mill.
Prick-timber, Gerard.
Louse Berry, Dogwood, Gatteridge Tree.
Fusain d'Europe, Bonnet de Prêtre commun, Fr.
Gemeine Spindelbaum, Ger.

Varieties † * - - 496

- 2 foliis variegatis Lodd. †
3 latifolius Lodd. †
4 nanus Lodd. *
5 fructu albo Lodd. †

2. verrucosus Scop. * Europe f. 165. 498
The warted-barked Euonymus, or Spindle Tree.

E. europæus leprosus L. fil.
Fusain galeux, ou verrucoux, Fr.
Warziger Spindelbaum, Ger.

3. latifolius C. Bauh. † Europe
pl. 61. fig. 661. 498

The broad-leaved Euonymus, or Spindle Tree.
E. europæus var. 2. L.
Fusain à larges Feuilles, Fr.
Breitblättriger Spindelbaum, Ger.

4. nanus Bieb. * Caucasus - 499
The dwarf Euonymus, or Spindle Tree.

5. atropurpureus Jacq. * North America
fig. 167. 499

The dark-purple-flowered Euonymus.
E. carolinensis Marsh.
E. latifolius Marsh.

6. americanus L. * North America
fig. 168. 169. 499

The American Euonymus, or Spindle Tree.
E. sempervirens Marsh.
E. alternifolius Mench.
The Burning Bush, Amer.

7. sarmentosus Nutt. * N. America 500
The trailing-stemmed Euonymus, or Spindle Tree.

E. scandens Hort.
E. americanus var. sarmentosus Dec.

8. obovatus Nutt. * N. America - 500
The obovate-leaved Euonymus, or Spindle Tree.

9. angustifolius Ph. * N. America 500
The narrow-leaved Euonymus, or Spindle Tree.

10. Hamiltonianus Wall. † Nepal 500
Hamilton's Euonymus, or Spindle Tree.
E. atropurpureus Wall.

11. garciniaefolius Roxb. † Nepal - 501
The Garcinia-leaved Euonymus.
E. laevis Ham.

12. grandiflorus Wall. * Nepal - 501
The large-flowered Euonymus.

App. i. Half-hardy Species of *Euonymus*. 501

- E. grossus* Wall. †
E. micranthus D. Don *
E. laevis D. Don *
E. japonicus Thunb. †
E. echinatus Wall. †
E. tingens Wall. †
E. glaber Roxb. †
E. imbricatus Wall. †
E. medicus Heyne *
E. vagans Wall. †
E. subtriflorus Blume *
E. Thunbergianus Blume *
E. pendulus Wall. †
E. frigidus Wall. †

II. CELASTRUS L. † † - 502

THE STAFF TREE.
Euonymoides Mench.
Celastre, Fr.
Celaster, Ger.

1. scandens L. † N. America f. 171. 502
The climbing Staff Tree.

Bourreau des Arbres, Fr.
Baumwörder, Ger.

2. bullatus L. † Virginia - - 502
The studded-capsuled Staff Tree.

App. i. Half-hardy Species of *Celastrus*. 503

III. NEMOPANTHES Rafin. * 503

THE NEMOPANTHES.
Iticiöldes Dum. Cours.

1. canadensis Dec. * Canada f. 172. 503
The Canadian Nemopanthes.

Flex canadensis Mx.
N. fasciculata Rafin.
Flex delicatula Bart.
Prinos lucidus Ait.
Houx du Canada, Fr.

IV. MAYTENUS Feuill. * - 503

THE MAYTENUS.
Senécia Lam.
Celastrus Willd.

1. chilensis Dec. * Chili f. 173. 503
The Chili Maytenus.

Senécia Maytenus Lam.
Celastrus Maytenus Willd.

V. CASSINE L. * - - 503

THE CASSINE.

1. Maurocœnia L. * Ethiopia - 501
Maurocœni's Cassine.

The Hottentot Cherry.
Maurocœnia frangularia Mill.

2. capensis L. * Cape of Good Hope 174. 504
The Cape Cassine.

Cape Phillyrea.

C. excelsa Wall., discolor Wall., Colpoia Thunb. 501

- VI. **HARTOGIA** Dec. - - - 504
THE HARTOGIA.
Schrœbera Thunb.
Elaeodendron Spreng.
1. *capensis* L. ♀ C. of Good Hope 504
The Cape Hartogia.
Schrœbera schinoides Thunb.
Elaeodendron schinoides Spreng.
-
- Aquifoliaceæ* Dec.**
- I. **MYGINDA** Jacq. - - - 505
THE MYGINDA.
Plex Pursh.
1. *myrtifolia* Nutt. ♀ N. Amer. 175. 505
The Myrtle-leaved Myginda. †
Plex Aquifolium Pursh.
- II. **FLEX** L. - - - 505
THE HOLLY.
Aquifolium Tourn. †
Houx, Fr.
Heiße, Stechpalme, Ger.
1. *Aquifolium* L. † Eur. pl. 62, 63. 505
The prickly-leaved Holly.
Hulcer, Hulferer, Holme, Eng.
Stechpalme, Stecheriche, Stechbaum, Stech-
laub, Hulse, Hulsenhbaum, Hulsenstrauß,
Hulst, Hulchs, Holst, Habze, Hüllgenol,
Myrtendorn, Christdorn, Maistorn,
Zwieseldorn, Kleczebusch, Stechapsel,
Stechweide, Waldistel, Ger.
Stikpalme, Maretorn, Christorn, Skoutisdet,
Dan.
Jernck, Christorn, Swed.
Le Houx, le grand Housson, l'Agron grand
Pardon, Bois Franc, Fr.
Agrifoglio, Aloro spinoso, Ital.
Acbo, Agrifolio, Span.
Azerinhó, Agrifolio, Acrifolio, Aginfolio,
Port.
Wagföschel, Ostrokof, Padub, Russ.
Schubbig hardkelt, Dutch.
- Varieties - - - 506
- 2 *heterophyllum* Hort. †
3 *angustifolium* Hort. †
4 *latifolium* Hort. †
5 *altaicænsis* Hort. †
6 *marginatum* Hort. † f. 176.
7 *laurifolium* Hort. † f. 177.
8 *ciliatum* Hort. † f. 179.
9 *ciliatum minus* Hort. †
10 *recurvum* Hort. † f. 181.
11 *serratifolium* Hort. † f. 182.
12 *crispum* Hort. †
13 *ferox* Hort. † f. 180.
14 *crassifolium* Hort. † f. 178.
15 *senescens* Sweet †
16 *albo-marginatum* Hort. †
17 *aureo-marginatum* Hort. †
18 *albo-pictum* Hort. †
19 *aureo-pictum* Hort. †
20 *ferox argenteum* Hort. †
21 *ferox aureum* Hort. †
22 *fructu luteo* Hort. †
23 *fructu albo* Hort. †
2. *halearica* Desf. † Minorca f. 183. 516
The Minorca Holly.
1. *Aquifolium* var. † Lam.
3. *opaca* Ait. N. Amer. pl. 61, 65. 516
The opaque-leaved Holly.
Agrifolium vulgare Clayt.
1. *Aquifolium* Gron.
The American Holly.
Varieties - - - 517
- 2 *macrodon*.
3 *latifolia*.
4 *acuminata*.
5 *globosa*.
4. *laxiflora* Lam. ♀ Carolina - - - 517
The loose-flowered Holly
1. *Aquifolium* *baccia flava*.
5. *Cassine* Ait. ♀ Carolina - - - 517
The Cassine-like Holly.
Aquifolium carolinense Cat.
1. *caroliniana* Mill.
1. *cassinoides* Link.
1. *Dahoon* Wall.
Cassena, Amer. Indians.
Dahoon Holly.
Variety - - - 517
- 2 *angustifolia* † f. 184.
6. *angustifolia* Willd. ♀ N. Amer. f. 185. 517
The narrow-leaved Holly.
1. *myrtifolia* Walt.
1. *rosmarinaifolia* Lam.
Variety - - - 518
- 2 *ligustrifolia* Pursh † f. 186.
7. *vomitaria* Ait. † N. Am. f. 186. 518
The Emetic Holly.
1. *Cassine vera* Walt.
1. *ligustrina* Jacq.
Cassine Peragna Mill.
1. *Cassena Michx.*
1. *religiosa* Bart.
1. *floridana* Lam.
Houx apalachine, Fr.
True Cassene, Cassene, Floridan,
The Yapon, Virginian,
The evergreen Cassena, Cassiberry-bush,
South Sea Tea, Eng.
8. *canariensis* Poir. † Canaries - - - 519
The Canary Holly.
9. *Dahoon* Wall. ♀ Carolina - - - 519
The Dahoon Holly.
1. *Cassine* Willd.
Variety - - - 519
- 2 *laurifolia* Nutt.
- App. i. *Species of Flex not yet introduced.* 519
E. colorata Ham., *cuneifolia* L., *C. bonariensis*, *ligustrifolia* G. Don
(*angustifolia* Nutt.), *neapolensis* Spreng.
- App. ii. *Half-hardy Species of Flex.* 519
E. dipyrrena Wall., *excelsa* Wall., *serrata* Boyle, *Perado* Ait. (*ma-*
derensis Lam., and fig. 187.), *chinensis* Sims (f. 188.), *hetero-*
phylla G. Don, *macrophylla*, *elliptica* B. B. et Kunth,
scopulorum B. B. et Kunth, *rapicola* H. B. et Kunth, *Pallsoni*
Pers., *emarginata* Thunb., *crenata*, *Thunb.*, *serrata* Thunb.,
latifolia Thunb., *in ricoides* Thunb., *asiatica* L., *inter-*
media, *rotunda* Thunb., *lanelloides* H. B. et Kunth, *paragua-*
riensis (f. 189.).
- III. **PRINOS** L. - - - 520
THE WINTER BERRY.
Agéria Adans.
Apalanche, Fr.
Winterbeere, Ger.

	Page		Page
§ i. <i>Prinoides</i> Dec. ☼ - -	520	4. <i>incúrva</i> Roxb. ☼ Nepal - -	526
1. <i>decíduus</i> Dec. ☼ N. America -	520	The incurved-spined Jujube. <i>Z. paniculata</i> Hamilt.	
The deciduous Winter Berry. <i>Plex prinoides</i> Ait. <i>Plex decídua</i> Walt.		5. <i>flexuòsa</i> Wall. ☼ Nepal - -	526
<i>Variety</i> - - -		The flexible Jujube.	
2. <i>æstivàlis</i> Dec.	521	App. i. <i>Half-hardy Species of Zizyphus</i>	526
<i>Plex æstivàlis</i> Lam.		<i>Z. Lótus</i> Lam. (<i>Rhámnus Lótus</i> L.) fig. 194.	
2. <i>ambíguus</i> Michx. ☼ N. Am. f. 190.	521	<i>Z. nitida</i> Roxb., <i>parvifólia</i> Del., <i>mucronata</i> Willd., <i>glábra</i> Roxb., (<i>Enóplia Mill.</i> , <i>tomentósa</i> Roxb., <i>Z. álbens</i> Roxb., <i>agrésis</i> Schult., <i>soporíferus</i> Schult., <i>capénsis</i> , <i>Jújuba</i> Lamb. (<i>Rhámnus Jújuba</i> L.))	
The ambiguous Winter Berry. <i>Cassine caroliniana</i> Walt.		II. <i>PALIURUS</i> L. ☼ - -	528
§ ii. <i>Agèria</i> Dec. ☼ -	521	CHRIST'S THORN. <i>Paliure</i> , <i>Porte-chapeau</i> , Fr.	
3. <i>verticillátus</i> L. ☼ N. Amer. f. 191.	522	1. <i>aculeátus</i> Lam. ☼ Asia	527
The whorled Winter Berry. <i>P. padifolius</i> Willd. <i>P. Gronóvi</i> Michx. <i>P. confértus</i> Mench. <i>P. pruifólius</i> Lodl.		pl. 66, 67. fig. 195.	
4. <i>dúbius</i> G. Don. ☼ N. America -	521	The prickly Christ's Thorn. <i>P. pélasus</i> Dum. <i>P. austràlis</i> Gærtn. <i>P. vulgaris</i> D. Don. <i>Rhámnus Paliurus</i> L. <i>Zizyphus Paliurus</i> Willd. <i>Christ's Thorn</i> , or <i>Lamb of Libya</i> , Gerard. <i>E'pine de Christ</i> , Argalon, <i>Porte-chapeau</i> , Fr. <i>Geflügelter Judendorn</i> , Ger.	
5. <i>lævigátus</i> Pursh. ☼ - f. 191.	522	III. <i>BERCHEMIA</i> Neck. ☼ -	528
6. <i>lanceolátus</i> Pursh. ☼ N. Amer.	522	THE BERCHEMIA. <i>Enóplia</i> Hedw.	
The lanceolate-leaved Winter Berry.		1. <i>volúbilis</i> Dec. ☼ N. Am. f. 196.	528
§ iii. <i>Wintèrlia</i> Mæneh. ☼ -	522	The twining Berchemia. <i>Rhámnus volúbilis</i> L. <i>Zizyphus volúbilis</i> Willd. <i>Enoplia volúbilis</i> Schult. <i>Supple Jack</i> , Virginian.	
7. <i>gláber</i> L. ☼ North America -	522	App. i. <i>Other Species of Berchemia.</i>	529
The glabrous Winter Berry.		<i>B. flavescens</i> Brong. (<i>Zizyphus flavescens</i> Willd.) <i>B. lineáta</i> Dec. (<i>Rhámnus lineatus</i> L. <i>B. Loureirána</i> Dec. (<i>Rhámnus lineatus</i> Lam.))	
8. <i>atomárius</i> Nutt. ☼ N. America -	522	IV. <i>RHA'MNUS</i> Lam. ☼ ☼ ☼ ☼	529
The atom-bearing Winter Berry.		THE BUCKTHORN. <i>The Ram</i> , or <i>Hart's Thorne</i> , Gerard. <i>Box Thorn</i> . <i>Neyprun</i> , Fr. <i>Wegdorn</i> , Ger.	
9. <i>coriáceus</i> Pursh. ☼ N. America	523	§ i. <i>Marcorélla</i> Neck.	529
The coriaceous-leaved Winter Berry. <i>P. gláber</i> Wats.		1. <i>Alatérnus</i> L. ☼ S. Europe f. 197.	529
App. i. <i>Other Species of Prinós.</i>	523	The Alaternus. <i>Alatérnus Phyllýra</i> Mill. <i>Varieties</i> - - -	
<i>P. dioicus</i> Vahl, <i>nitidus</i> Vahl.		2. <i>baleárica</i> Hort. Par. ☼	
		3. <i>hispánica</i> Hort. Par. ☼	
		4. <i>fóliis maculátis</i> ☼	
		5. <i>fóliis áureis</i> ☼	
		6. <i>fóliis argénteis</i> ☼	
		7. <i>angustifólia</i> ☼	
		<i>R. Clusii</i> Willd.	
<i>Rhamnáceæ</i> Lindl. ☼ ☼ ☼ ☼ ☼ ☼	523	2. <i>hýbridus</i> L' <i>Hérit.</i> ☼ Hybrid	531
1. <i>ZIZYPHUS</i> Tourn. ☼ -	524	The hybrid Alaternus. <i>R. burgundiacus</i> Hort. Par. <i>R. sempervirens</i> Hortulan.	
<i>Jujubier</i> , Fr. <i>Judendorn</i> , Ger.		3. <i>longifólius</i> Link. ☼ - -	531
1. <i>vulgàris</i> Lam. ☼ Syria - f. 193.	524	The long-leaved Buckthorn. <i>R. Willdenowianus</i> Rom. et Schult.	
The common Jujube. <i>Rhámnus Zizyphus</i> L. <i>Z. sativa</i> Desl. <i>Z. Jújuba</i> Mill. <i>Jujubier cultivé</i> , Fr. <i>Brustbeeren</i> , Ger. <i>Giuggiol</i> , Ital.			
2. <i>sinénsis</i> Lam. ☼ China - -	525		
The Chinese Jujube. <i>Rhámnus Zizyphus</i> Lour.			
3. <i>spína-Christi</i> Lam. ☼ Africa -	526		
Christ's Thorn Jujube. <i>Rhámnus spína-Christi</i> L. <i>R. Nábeca</i> Forsk. <i>Z. africana</i> Mill. <i>Z. Nábeca</i> Lam. <i>Nábeca Alp.</i> <i>Enóplia spinosa</i> Benth. <i>Christskronen Judendorn</i> , Ger.			
<i>Variety</i> - - -			
2. <i>inérmis</i> Dec. ☼	526		

- | | Page | | Page |
|-------------------------------------------------|------|----------------------------------------------------------------|------|
| 1. catharticus L. ♀ Europe; | 531 | 20. franguloides Michx. ♂ North America | 536 |
| pl. 68. fig. 198. | | fig. 207. | |
| The purging Buckthorn. | | The Frangula-like Buckthorn. | |
| <i>The White Thorn</i> of the modern Greeks. | | <i>R. alnifolia</i> var. <i>franguloides</i> Dec. | |
| Variety - - - | 531 | 21. pumilus L. ♂ South Europe | 536 |
| 2 hydruntina Jac. ♀ | | pl. 69. fig. 208. | |
| 5. virgatus Roxb. ♂ Himalaya - - | 532 | The Alpine Buckthorn. | |
| The twiggy Buckthorn. | | 22. pumilus L. ♂ Austria - - | 536 |
| <i>R. catharticus</i> Hamilt. | | The dwarf Buckthorn. | |
| 6. tinctorius Waldst. ♂ Hung. f. 199. | 532 | <i>R. rupetris</i> Scop. | |
| The Dyer's Buckthorn. | | § ii. <i>Frangula</i> Tourn. ♀ ♂ | 537 |
| <i>R. cardiosepnus</i> Willd. | | 23. carolinianus Wall. ♂ Carolina | 537 |
| 7. infectorius L. ♂ S. Europe f. 200. | 533 | The Carolina Buckthorn. | |
| The staining Buckthorn. | | 24. Frangula L. ♀ Europe | 539 |
| <i>Rhamnus Lycium</i> Scop. | | pl. 70. fig. 209. | |
| <i>Dwarf, or yellow-berryed, Buckthorn;</i> | | The breaking Buckthorn. | |
| <i>Avignon Berry.</i> | | <i>Berry-bearing Alder.</i> | |
| <i>Nerprun des Teinturiers, Grainc d'Avi-</i> | | <i>Nerprun Bourgeois, Aune noir, Fr.</i> | |
| <i>Farbender Wegdorn, Ger.</i> | | <i>Glatter Wegdorn, Ger.</i> | |
| 8. saxatilis L. ♂ S. Europe f. 201. | 533 | Variety - - - | 537 |
| The Stone Buckthorn. | | 2 angustifolia Hort. ♀ | |
| <i>R. longifolia</i> Mill. | | 25. latifolia L'Hérit. ♀ Azores | 538 |
| <i>Stein Wegdorn, Ger.</i> | | pl. 71. fig. 210. | |
| 9. oleoides L. ♂ Sicily - fig. 202. | 534 | The broad-leaved Buckthorn. | |
| The Olive-like Buckthorn. | | App. i. <i>Hardy Species of Rhamnus not yet</i> | 538 |
| <i>R. oleifolia</i> Hort. | | <i>introduced.</i> - - - | |
| 10. luxifolia Poir. ♂ Africa f. 203. | 534 | <i>R. amygdalinus</i> Desf. | |
| The Box-leaved Buckthorn. | | <i>R. persicifolia</i> Moris. | |
| 11. pubescens Poir. ♂ Levant - - | 534 | <i>R. prunifolia</i> Smith. | |
| The pubescent Buckthorn. | | <i>R. Sibthorpianus</i> Schult. (<i>R. pumilus</i> Sibth.) | |
| <i>R. oleoides</i> Lam. | | <i>R. Purshianus</i> Dec. (<i>R. alnifolia</i> Pursh) f. 211. | |
| 12. lycioides L. ♂ Spain - - | 534 | <i>R. sanguineus</i> Pers. | |
| The Lycium-like Buckthorn. | | <i>R. minutiflorus</i> Pursh. | |
| Variety - - - | 534 | <i>R. carpinifolia</i> Pall. f. 212. | |
| 2 arragonensis Asso. | | App. ii. <i>Half-hardy Species.</i> - - - | 538 |
| 13. Erythroxylon Pall. ♂ Siberia | 534 | <i>R. integrifolia</i> Dec. (<i>R. coriaceus</i> Nees.) | |
| fig. 204. | | <i>R. prinoides</i> L'Hérit. (<i>Zizyphus lucidus</i> Mench.) | |
| The red-wooded Buckthorn. | | <i>R. celtidifolia</i> Thunb. | |
| Variety - - - | 295 | <i>R. crenulata</i> Ait. | |
| 2. angustissimum ♂ fig. 205. | | <i>R. serrulata</i> H. B. et Kunth. | |
| <i>R. lycioides</i> Pall. | | <i>R. microphyllus</i> Willd. | |
| 14. rupetris Will. ♂ France | 555 | <i>R. umbellata</i> Car. | |
| The Rock Buckthorn. | | <i>R. tenuifolia</i> Moc. | |
| <i>R. pumilus</i> & <i>rupetris</i> Dec. | | V. CEANOTHUS L. ♂ ♂ - - | 539 |
| 15. valentinus Willd. ♂ Spain - | 553 | THE CEANOTHUS, or Red Root. | |
| The Valencia Buckthorn. | | <i>Rhamnus</i> L. | |
| <i>R. pumilus</i> Cav. | | <i>Ceanothe</i> , Fr. | |
| <i>R. pumilus</i> var. <i>valentinus</i> Dec. | | <i>Säkebaum</i> , Ger. | |
| 16. Wulfenii Spreng. ♂ Austria | 535 | 1. azureus Desf. ♂ Mexico f. 213. | 539 |
| Wulfen's Buckthorn. | | The azure-flowered Red Root. | |
| <i>R. pumilus</i> Wulf. | | <i>C. cæruleus</i> Lag. | |
| <i>R. pumilus</i> var. <i>Wulfenii</i> Dec. | | <i>C. bicolor</i> Willd. | |
| 17. pusillus Ten. ♂ Naples - | 535 | 2. americanus L. ♂ N. Amer. 211. | 539 |
| The small Buckthorn. | | The American Red Root, or New Jersey Tea. | |
| <i>R. pumilus</i> var. <i>neapolitanus</i> Dec. | | 3. tardiflorus Horn. N. America | 510 |
| 18. dahuricus Pall. ♂ Dahuria - | 535 | The late-flowering Red Root. | |
| The Dahurian Buckthorn. | | 4. ovatus Desf. ♂ N. America - | 510 |
| 19. alnifolius L'Hérit. ♂ North America | 536 | The ovate-leaved Red Root. | |
| fig. 206. | | 5. intermedius Pursh. ♂ N. America | 510 |
| The Alder-leaved Buckthorn | | The intermediate Red Root. | |
| | | 6. sanguineus Pursh ♂ N. America | 510 |
| | | The bloody-branched Red Root. | |

7. *microphyllus* Michx. n. N. America 540
 The small-leaved Red Root.
C. hypericoides L'Hérit.

App. i. *Other Species of Ceanothus.* 540

C. velutinus Hook. fig. 215.
C. levigatus Hook.
C. thyrsoïdorus Esch.

App. I. *Half-hardy Rhamnaceæ.* 541

Sphaerocarya edulis Wall.
Condalia microphylla Cav. fig. 216.
Sageretia Theezans Brongn.
Rhamnus Theezans L.
R. Thun Osb.
S. oppositifolia Brongn.
Rhamnus oppositifolia L.
S. hamosa Brongn.
Rhamnus hamosa L.
Scutia capensis Brongn.
Rhamnus capensis Thunb.
Ceanothus capensis Dec.
Retanilla obcordata Brongn.
Rhamnus Retanilla Domb.
Collitia Retanilla Vent.
R. E'phedra Brongn.
R. E'phedra Brongn.
Collitia E'phedra Vent.
Collitia spinosa Kunth. fig. 217.
C. polyacantha Willd.
C. serratifolia Vent.
Rhamnus Spartium Domb.
C. cruciata Gill.
C. foex Gill.
C. ulicina Gill.
C. Chacayé G. Don.
Rhamnus Chacayé Domb.
C. tetrasida Brongn.
Troxou quinquerivis Mey.
T. trinervis
Discaria americana Hook.
D. australis Hook.
Hovenia dulcis Thunb.
Columbinia trilobata Brongn.
Rhamnus trilobata Moc.
C. Mociuana G. Don.
Ceanothus Mociuanus Dec.
Willemctia africana Brongn.
Ceanothus africanus Lin.
Pomaderris elliptica Lab. fig. 218.
Cryp'tandra Sm., *Bartlingia* Brongn., *Solenanthes* G. Don, *Tri-chocephalus* Brongn., *Physica* L., *Soulangea* Brongn., *Gouania* Jacq., *Carpodetus* Forst., *Oleua* Thunb.
Physica ericoides L. fig. 219.

Bruniæcæ R. Br. - 542

Homalinæcæ Lindl. 541

I. *ARISTOTELIA* L'Hérit. 543
 THE ARISTOTELIA.

1. *Macqui* L'Hérit. Chili pl. 72. 543
 The Macqui Aristotelia.
A. glandulosa R. et P.
A. Macqui Dec.
Variety - 543
 2 *foliis variegatis* 543

II. *AZARA* R. et P. 544
 THE AZARA.

1. *dentata* R. et P. Chili f. 220. 544
 The tooth-leaved Azara.
 2. *integrifolia* R. et P. Chili - 544
 The entire-leaved Azara.

App. I. *Other Species of Homalinæcæ.* 541

Blackwellia nepalensis Dec.
Astránthus cochinchinensis Lour.
Neillia thyrsoïdora D. Don.
N. rubiflora D. Don.

Anacardiæcæ Lindl. 545

I. *PISTA'CIA* L. 545

THE PISTACIA TREE.
Terebinthus Juss.
Pistachier, Fr.
Pistacie, Ger.
Pistachio, Ital.

1. *vèra* L. Syria - fig. 221. 545
 The true Pistacia Nut Tree.
P. officinarum Hort. Kew.

Varieties - 545

2 *trifolia* L. 545

3 *narbonensis* Bocc. 545

P. reticulata Willd.

2. *Terebinthus* L. S. Europe 546

The Turpentine Pistacia.
T. vulgaris Tourn.
P. vèra Mill.
Venetian, or *Chian*, *Turpentine Tree*.
Pistachier Terebinthe, Fr.
Turpentin Pistacie, Ger.
Terebinto, Ital.

Variety - 546

2 *sphaerocarpa* Dec. 546

3. *Lentiscus* L. Levant - 547

The Mastich Tree.

Varieties - 547

2 *angustifolia* Dec. 547

P. massiliensis Mill.

3 *chia* N. Du Ham. 547

P. chia Desf.

4. *atlantica* Desf. Africa - 548

The Mount Altas Mastich, or *Turpentine Tree*.

II. *RHU'S* L. 548

THE SUMACH.

§ i. *Cótinus* Tourn. 548

1. *Cótinus* L. S. Europe f. 223. 549

The *Cótinus* Rhus.
Cótinus Coggágria Scop.
Cótinus coriácca Duh.
Venus Sumach, *venice Sumach*, *Wild Olive*.
Sumach Fustet, *Arbre aux Péruques*,
 Fr.
Perücken Sumach, Ger.
Scotino, Ital.

§ ii. *Sùmach* Dec. 550

2. *typhina* L. N. America - 550

The Fever Rhus.
R. virginiana Baulh.
Virginian Sumach, *Stag's Horn Stumach*.

Varieties - 550

2 *arboræscens* Willd. 550

3 *frutæscens* 550

3. *viridiflora* Poir. N. America 551

The green-flowered Sumach.
R. canadense Mill.

	Page		Page
4. glabra L. ♀ N. America f. 225.	551	§ iv. Lobodium Dec. ♀	557
The glabrous Sumach.		14. suaveolens Ait. ♀ Carolina	557
<i>Varieties</i> -	551	The sweet-scented Sumach.	
1 hermaphrodita ♀		<i>Myrica trifoliata</i> Hortul.	
R. glabra Willd.		<i>Toxicodendron erectum</i> Mill.	
2 dioica Lam. ♀		15. aromatica Ait. ♀ N. America	557
3 coccinea ♀		The aromatic Sumach.	
R. carolinianum Mill.		App. i. Other Species of Rhus. -	557
R. elegans Ait.		R. lobata Hook.	
5. pumila Michx. ♀ N. America	552	R. acuminata Dec.	
The dwarf Sumach.		R. Anala D. Don. (R. Bucku-Amela H. B.)	
6. vernicifera Dec. ♀ Japan	552	R. bahianensis G. Don.	
The Varnish-yielding Sumach.		R. oxyacantha H. B.	
Sitz, Urus, Japanese.		R. oxyacanthoides H. B.	
R. vernix L.		R. dioica G. Don.	
R. juglandifolium Wall.		R. heterophylla	
7. venenata Dec. ♀ N. Am. f. 226.	552	III. DUVAUVA Kunth. † ♀ -	558
The poisonous Sumach.		THE DUVAUVA.	
R. vernix L.		Schinus Andr.	
Toxicodendron pinnatum Mill.		Amirris Cav.	
Poison Sumach, Swamp Sumach, Poison Elder.		1. dependens Dec. † Chili fig. 232.	559
8. Coriaria L. ♀ South Europe	fig. 227, 228. 553	The drooping-branched Duvaava.	
The hide-tanning Sumach, or Elm-leaved Sumach.		Amirris polygama Cav.	
9. copallina L. ♀ N. Amer. f. 229.	554	Schinus dependens Ort.	
The Gum Copal Rhus, or Mastich-tree-leaved Sumach.		Duvaava dependens a Hook.	
<i>Variety</i> -	554	2. ovata Lindl. ♀ Chili	559
2 leucantha Jac. ♀		The ovate-leaved Duvaava.	
10. radicans L. † ♂ North America	fig. 230. 555	3. latifolia Gill. ♀ Chili - fig. 243.	559
The rooting-branched Sumach.		The broad-leaved Duvaava.	
R. Toxicodendron var. α Michx.		D. dependens γ Hook.	
R. Toxicodendron var. β Torrey.		4. dentata Dec. ♀ Chili	559
<i>Varieties.</i>		The toothed-leaved Duvaava.	
1 vulgaris †	555	Schinus dentata Andr.	
R. Toxicodendron vulgare Pursh.		App. i. Other Species of Duvaava.	560
Toxicodendron vulgare, and T. volubile Mill.		App. I. Other Species of Anacardiaceæ. 560	
2 volubile †	555	Sibia parviflora Wall.	
Toxicodendron volubile Mill.		S. campanulata Wall.	
3 microcarpa †	555	Schinus Mülli L. ♀ Brazil fig. 234.	
R. Toxicodendron microcarpon Ph.		S. M. 2 Azeva L., S. Huizua Mol., and S. virgata Sret.	
11. Toxicodendron L. ♂ † N. America	fig. 231. 556	Triceros cochinchinensis.	
The Poison-tree Sumach.		Heterodendron olenifolium Desf.	
R. Toxicodendron quercifolium Michx.		Stylopladium spatulatum Desf.	
Toxicodendron pubescens Mill.		Cinorum triocceum L.	
R. T. serratum Mill.		C. pulverulentum.	
Poison Oak, Poison Nut, Poison Vine.		<hr/>	
§ iii. Thezera Dec. ♀	556	Burseraceæ Kunth. -	561
12. pentaphylla Desf. ♀ Sicily	556	Balsamodendron gilcadense - - - - -	fig. 255. 561
The five-leaved Sumach.		Amirris glandulosa L.	
Rhamnus pentaphyllus Jac.		Canarium Pimenta Kunig.	561
R. Thezera Pers.		Fagastrium G. Don.	561
13. zizyphina Ait. ♀ Sicily	557	<hr/>	
The Zizyphus like Sumach		Amirridaceæ Lindl. -	561
Rhamnus zizyphifolia Pers.		Amirris toxicifera Willd.	561
		A. balsamifera L.	561
		A. floridana Nutt.	561
		<hr/>	
		Leguminaceæ, ♂ ♀ ♂ † ♂ -	561
		Sect. I. SOPHOREÆ. † -	563
		I. SOPHORA R. Br. †	562, 563
		THE SOPHORA.	

	Page		Page
I. japonica L. Japan pl. 73, 74.	563	2. nana Forst. □ Britain	
The Japan Sophora.		fig. 264, 265.	575
S. sinica Ros.		The dwarf Furze.	
Varieties - - -	563	U. minor Roth.	
2 variegata Hort.		U. europæus ß Lin.	
3 pendula Hort. ¶ pl. 75.		3. provinciâlis Lois. ■ France -	575
II. VIRGILII L. ¶ - - -	565	The Provence Furze.	
THE VIRGILIA.		4. stricta Mackay. ■ Ireland -	575
1. lutea Michx. ¶ N. Amer. pl. 76.	565	The upright-growing Furze.	
The yellow-wooded Virgilia, or Yellow Wood.		U. hibernica Don.	
III. PIPTANTHUS Swt. ■ - - -	566	U. fastigiata Hort.	
THE PIPTANTHUS.		V. STAURACANTHUS Link. ■ 576	
Thermopsis D. Don.		THE STAURACANTHUS.	
Anagyris Wall.		Ulex Brot.	
Baptisia Hook.		Leafless Furze.	
1. nepalensis Swt. ■ Nepal 237, 238.	566	1. aphyllus Link. ■ Spain -	576
The Nepal Piptanthus.		The leafless Stauracanthus.	
Thermopsis habruifolia D. Don.		Ulex genistoides Brot.	
Anagyris indica Wall.		U. mitis Hort.	
Baptisia nepalensis Hook.		VI. SPARTIUM Dec. ■ - - -	576
App. I. Half-hardy Species of Sophoræ. 567		THE SPANISH BROOM.	
Sophora velutina Lindl.		Spartianthus Link.	
S. tormentosa Hook. (S. occidentalis Lindl.)		Genista Lam.	
Edwardsia chilensis Miers (Sophora macrocarpa Smith), f. 259.		Genêt d'Espagne, Fr.	
E. grandiflora Sal. (Sophora tetraptera Ait.) f. 240.		Biuscartige Pfirricmen, Ger.	
E. microphylla Sal. (Sophora microphylla Ait.) f. 241.		1. junceum L. ■ South Europe -	575
E. myriophylla Wand. (E. minima Lodd. Cat.)		The Rush-like Spanish Broom.	
E. chrysophylla Sal.		Genista juncea, Lam.	
Cyclopia genistoides R. Br. (Gompholobium maculatum Bot. Rep.)		G. odorata Mench.	
Podalyria sericea R. Br. (Sophora sericea Bot. Rep.) f. 242.		Spartianthus junceus Mench.	
Chorozeina Henchmannii R. Br. f. 243.		Variety - - -	576
Podolobium trilobatum R. Br. f. 244, 245.		flôre pleno ■	
Oxylobium arborescens R. Br. f. 246.		VII. GENISTA Lam. ■ ■ □ ■ □ ■	
Callistachys ovata Sims.		■ □ ■ ■ ■ ■ ■ □ ■ ■	577
Brachysema latifolium R. Br. f. 247.		THE GENISTA.	
Gompholobium grandiflorum Smith.		Genista and Spartium, spec., L.	
Burtonia R. Br.		Genêt, Fr.	
Jacksonia scoparia R. Br.		Giuster, Ger.	
Viminaria denudata Smith, f. 249.		1. parviflora Dec. ■ Levant -	578
Sphaerolobium vimineum Smith, f. 251, 252.		The small-flowered Genista.	
S. medium.		Spartium parviflorum Vent.	
Aotus ericoides G. Don. f. 255.		2. clavata Poir. ■ □ Spain -	578
Dillwynia glaberrima Smith, f. 255, 256.		The club-shaped-calyxed Genista.	
Eutaxia myrtifolia R. Br. f. 254.		Spartium sericeum Vent., not of Ait.	
E. pinguis Swt.		3. cándicans L. ■ Levant fig. 267.	578
Sclerothamnus microphyllus B. Br.		The whitish-surfaced Genista.	
G. asrolobium bilobum Ker.		Cytisus cándicans L.	
Euchilus obovatus R. Br. f. 258, 257.		Cytisus pubescens Mench.	
Pultenaea stricta Bot. Mag. f. 259.		4. triquetra Ait. ✱ Spain fig. 268.	578
Daviesia latifolia R. Br. f. 261.		The triangular-stemmed Genista.	
Mirbelia reticulata Smith, f. 260.		G. triquetra Lam. ?	
Sect. II. LOTEÆ. - - -	571	5. bracteolata Lk. ■ □ - - -	579
¶ ■ ■ ■ ■ ■ ■ ■		The bracteolated Genista.	
IV. ULEX L. ■ ■ - - -	571	6. umbellata Poir. ■ □ Barbary -	579
THE FURZE.		The umbellate-flowered Genista.	
Ajone, Fr.		Spartium umbellatum Desf.	
Hecksuame, Ger.		Variety ■ □ - - -	579
1. europæa L. ■ Europe f. 262, 263.	571	2 capitata Dec. ■ □	
The European Furze		7. lusitânica L. ■ Portugal -	579
Genista spinosa L'Obel.		The Portugal Genista.	
U. grandiflorus Pour.			
U. vernalis Thore.			
Whin, Gorsc, Prickly Broome.			
Ajone commun, jone marin, Jomarin.			
Genêt épineux Fr.			
Varieties - - -	571		
2 flôre pleno ■			
provinciâlis.			
stricta.			

	Page		Page
8. (l.) radiata Scop. ♀ Italy f. 269.	519	26. tinctoria L. ♂ Europe f. 276.	583
The rayed-branched Genista.		The Dyer's Broom, or Green Weed.	
<i>Spartium radiatum</i> L.		Base Broom, Grecu Weed, Dyer's Weed,	
<i>G. ibensis</i> Dalech.		Wood-waxen.	
9. ephedroides Dec. ♀ Sardinia	- 580	Genêt des Teinturiers, Genêt de Sibirie,	
The Ephedra-like Genista.		Fr.	
10. triacanthos Brox. ♀ Portugal	580	Farbender Ginster, Ger.	
The three-spined Genista.		Varieties ♂ - - - 583	
<i>G. rostrata</i> Poir.		2 latifolia Dec. ♂	
Variety ♀ - - - 580		3 hirsuta Dec. ♂	
2 interrupta Dec. ♀		4 pratensis Poll. ♂	
11. horrida Dec. ♀ Pyrenees	- 580	27. (t.) sibirica L. ♀ Siberia	- 584
The horrid Genista.		The Siberian Genista.	
<i>Spartium horridum</i> Vahl.		Genistoides elata Mench.	
<i>G. eriuca</i> Gilib.		<i>G. tinctoria</i> var. N. Du Ham.	
12. sylvestris Scop. ♀ Austria	- 580	28. (t.) tetragona Besser. ♂ Podolia	584
The Wood Genista.		The quadrangular-horned Genista.	
<i>G. hispanica</i> Jacq.		29. (t.) polygalifolia Dec. ♀ Port.	584
13. Scorpins Dec. ♀ Europe	- 580	The Milkwort-leaved Genista.	
The Scorpion Genista.		<i>G. polygalifolia</i> Brot.	
<i>Spartium Scorpins</i> L.		<i>G. erubata</i> Link.	
<i>G. spiniflora</i> Lam.		<i>G. tinctoria lusitanica maxima</i> Tourm.	
Scorpion Furze, Gerard.		30. (t.) florida L. ♀ Spain	- 584
14. hispanica L. ♂ Spain	- 581	The florid Genista.	
The Spanish Genista.		31. mantica Poll. ♂ Italy	- 584
Spanish Furze, Hort.		The Mantuan Genista.	
15. anglica L. ♂ Europe f. 270.	581	32. ovata Waldst. ♀ Hungary	- 584
The English Genista, or Petty Whin.		The ovate-leaved Genista.	
<i>G. minor</i> Lam.		<i>G. nevata</i> Kit. in Litt.	
16. germanica L. ♀ Europe f. 271.	581	33. patula Bieb. ♀ Tauria	- 584
The German Genista.		The spreading Genista.	
<i>Scorpius spinosus</i> Mench.		34. triangularis Willd. ♂ Hungary	- 584
<i>Foglera spinosa</i> Fl. Wett.		The triangular-stemmed Genista.	
Variety ♀ - - - 581		<i>G. triquetra</i> Waldst., not of Ait.	
2 inermis Dec. ♀		35. sagittalis L. ♂ Europe, fig. 277.	585
17. purgans L. ♀ France	- 581	The arrow-jointed Genista.	
The purgans Genista.		<i>G. herbacea</i> Lam.	
<i>Spartium purgans</i> L.		Genistella racemosa Mench.	
18. sericea Wolf. ♂ Austria	- 581	Saltzweidelia sagittalis Fl. Wett.	
The silky Genista.		Variety ♂ - - - 585	
19. humifusa L. ♂ Levant	- 582	2 minor Dec. ♂	
The trailing Genista		36. diffusa Willd. ♂ Italy	- 584
20. aphylla Dec. ♀ Siberia	- 528	The diffuse Genista.	
The leafless Genista.		<i>G. humifusa</i> Wulf.	
<i>Spartium aphyllum</i> L.		<i>Spartium procumbens</i> Jacq., not of Ait.	
<i>G. virgata</i> Lam.		37. prostrata Lam. ♂ France fig. 278.	585
21. monosperma Lam. ♀ Mediterranean	fig. 272. 582	The prostrate Genista.	
The one-seeded Genista.		<i>G. pendiculata</i> L'Hérit.	
<i>Spartium monospermum</i> L.		<i>G. decumbens</i> Dur. Bourg.	
<i>G. Raetam</i> Forsk.		<i>G. Halleri</i> Reyn.	
22. sphaerocarpa Lam. ♀ S. Europe	582	38. procumbens Waldst. et Kil. ♂	Hungary - - - 585
The round fruited Genista.		The procumbent Genista.	
<i>Spartium sphaerocarpon</i> L.		39. pilosa L. ♂ Europe - fig. 279.	586
23. aethiœnsis Dec. ♀ Etna	fig. 273. 582	The hairy Genista.	
The Mount Etna Genista.		<i>G. rœpans</i> Lam.	
<i>Spartium aethiœnsis</i> Biv.		Genistoides tuberculata Mench.	
<i>Spartium trisperrum</i> Smith.		40. pilocarpa Link. ♂	- 586
24. scariosa Vir. ♀ Naples	- 583	The hairy-fruited Genista.	
The scarious-margined-leaved Genista.		App. i. Hardy Species of Genista not yet in-	
<i>G. janensis</i> Viv. Cat.		roduced. - - - - - 586	
<i>G. genuana</i> Pers.		<i>G. mollis</i> Dec.	
25. anxantiœca Ten. ♂ Nap. f. 274, 275.	583	<i>Spartium molle</i> Cav.	
The Anxantiœca Genista.			
<i>G. ansantiœca</i> Tenore.			

	Page
G. patens Dec.	
<i>Spartium patens</i> Cav.	
G. sessilifolia Dec.	
G. casparrina Guss.	
G. acanthoclada Dec.	
<i>Lobelia</i> Dec.	
<i>Spartium erinacoides</i> .	
G. parvifolia G. Dou.	
<i>G. microphylla</i> Moris.	
O. Salzmanni Dec.	
<i>G. umbellata</i> Salzm.	
G. aspalathoides Lam.	
<i>Spartium aspalathoides</i> Desf.	
G. ferox Poir.	
<i>Spartium heterophyllum</i> L'Hérit.	
<i>Spartium ferox</i> Desf.	
G. Cupani Guss.	
<i>Acacia triphylla</i> Cup.	
G. infesta G. Don.	
<i>Spartium infestum</i> Guss.	
G. hirsuta Vahl.	
var. 2 <i>cuspidata</i> Dec.	
3 <i>orientalis</i> Dec.	
G. algarbiensis Brot.	
<i>G. hirsuta</i> var. 8? <i>algarbiensis</i> Dec.	
G. truncispinata Desf.	
<i>Spartium truncispinatum</i> Cav.	
G. gibraltaria Dec.	
G. falcata Brot.	
G. corsica Dec.	
<i>Spartium corsicum</i> Lois.	
G. ramosissima Poir.	
<i>Spartium ramosissimum</i> Desf.	
G. cinerea Dec.	
<i>Spartium cinereum</i> Vill. Prop.	
<i>G. scoparia</i> Vill. Dauph.	
<i>G. florida</i> Asso.	
G. stylosa Spreng.	
<i>G. bracteolata</i> Willd.	
G. multicaulis Lam.	
G. tenuifolia Lois.	
G. depressa Bieb.	
G. Perreymondii Lois.	
G. pulchella Pistani.	
G. tridentata L.	
G. albida Willd.	
G. micrantha Ort.	
G. ? heterophylla Dec.	
<i>Cytisus heterophyllus</i> La Peyr.	
G. angulata G. Don.	

App. ii. *Half-hardy Species of Genista.*

G. canariensis L.	588
<i>Spartium albicans</i> Cav.	
<i>Cytisus paniculatus</i> Lois.	
<i>Cytisus ramosissimus</i> Poir.	
G. linifolia L.	f. 280.
<i>Spartium linifolium</i> Desf.	
<i>Cytisus linifolius</i> Lam.	
<i>Genista linifolia</i> Moench.	
G. biflora Dec.	
<i>Spartium biflorum</i> Desf.	
G. microphylla Dec.	
<i>Spartium microphyllum</i> Cav.	
G. tridens Cav.	
G. aegyptiaca Spreng.	
G. virgata Dec. fig. 281.	
<i>Spartium virgatum</i> Ait.	
<i>G. gracilis</i> Poir.	
<i>Cytisus tener</i> Jacq.	
G. cuspidata Dec.	
<i>Spartium cuspidatum</i> Burch.	
G. congesta Dec.	
<i>Spartium congestum</i> Willd.	
G. desiderata Dec.	
G. scandens Lois.	

VIII. CYTISUS Dec.

THE CYTISUS.
Cytisus and *Spartium*, spec., L.
 and Lam., &c.
Cytise, Fr.
Bohnenbaum, Ger.

§ i. *Albumoides* Dec. - 589

1. <i>albus</i> Link. Levant fig. 282. 589	
The white Cytisus, or Portugal Broom.	
<i>Genista alba</i> Lam.	
<i>Spartium album</i> Desf.	
<i>Spartium multiflorum</i> Ait.	
<i>Spartium dispersum</i> Moench.	
<i>Genista multiflora</i> N. Du Ham.	
<i>Spartium a Fleurs blanches</i> , Fr.	
<i>Weisse Pflümen</i> , Ger.	
Variety - - 590	
2 <i>incarnatus</i>	

	Page
§ ii. <i>Laburnum</i> Dec. - 590	
2. <i>Laburnum</i> L. Europe f. 77. 590	
The common Laburnum.	
<i>C. alpinus</i> Lam.	
<i>Bean-trefoile Tree</i> , and <i>Peascod Tree</i> , Gerard.	
<i>Pea Tree</i> , Scotch.	
<i>Golden Chain</i> .	
<i>L'Aubours</i> , <i>Faux E'bénier</i> , <i>Arbois</i> or <i>Arc Bois</i> , Fr.	
<i>Geneine Bohnenbaum</i> , Ger.	
Varieties - - 590	
2 <i>quercifolium</i> Hort. pl. 78.	
<i>C. L. incisum</i> .	
3 <i>pendulum</i> Hort.	
4 <i>foliis variegatis</i>	
5 <i>purpurascens</i> Hort.	
<i>C. L. purpureum</i> Hort.	
<i>C. Adami</i> Poir.	
<i>C. L. coccineum</i> Baum.	
3. (<i>L.</i>) <i>alpinus</i> Mill. Alps pl. 79. 591	
The Alpine, or Scotch, Laburnum.	
<i>C. Laburnum</i> β Ait.	
<i>C. angustifolium</i> Moench.	
<i>C. Laburnum</i> var. <i>latifolium</i> Pers.	
<i>Cytise des Alps</i> , <i>L'Aubours</i> , Fr.	
<i>Alpen Bohnenbaum</i> , Ger.	
<i>Maggio pendolino</i> , Ital.	
Variety - - 591	
2 <i>pendulus</i> pl. 80.	
4. <i>nigricans</i> L. France f. 283. 593	
The black Cytisus.	
5. <i>sessilifolius</i> L. Fr. f. 284, 285. 594	
The sessile-leaved Cytisus.	
6. <i>triflorus</i> L'Hérit. South Europe fig. 286. 594	
The three-flowered Cytisus.	
<i>C. villosus</i> Pour.	
7. <i>mollis</i> Willd. - 595	
The soft Cytisus.	
8. <i>patens</i> L. Portugal - 595	
The spreading Cytisus.	
<i>C. pendulum</i> L. Fil.	
<i>Genista tomentosa</i> Poir.	
<i>Spartium patens</i> L., not of Cav.	
9. <i>grandiflorus</i> Dec. Portugal - 595	
The great-flowered Cytisus.	
<i>Spartium grandiflorum</i> Brot.	
10. <i>scoparium</i> Link. Europe f. 287. 595	
The common Broom.	
<i>Spartium scoparium</i> L.	
<i>Genista scoparia</i> Lam., not of Vill.	
<i>Genista hirsuta</i> Moench.	
<i>Genêt à Balais</i> , <i>Genêt commun</i> , Fr.	
<i>Geneine Pflümen</i> , Ger.	
Varieties - - 595	
2 <i>albus</i> Hort.	
3 <i>flore pleno</i> Hort.	
§ iii. <i>Calyeotome</i> Link. 597	
11. <i>spinosus</i> Lam. S. Europe - 597	
The spiny Cytisus.	
<i>Spartium spinosum</i> L.	
d 2	

	Page
12. lanigerus Dec. ☼ ☐ S. Europe	597
The wool-bearing Cytisus.	
<i>Spartium lanigerum</i> Desf.	
<i>Calycotome villosa</i> Link.	
<i>Spartium villosum</i> Brot.	
Variety ☼ ☐ - - - 597	
2 rigidus Dec. ☼ ☐ - - - 598	
§ iv. Tubercytisus Dec. ☼ ☼ ☼	598
13. leucanthus Walst. et Kit. Austria	598
fig. 288.	
The white-flowered Cytisus.	
14. purpureus Scop. ✕ Austria	598
fig. 289, 290.	
The purple-flowered Cytisus.	
Variety ✕ - - - 598	
2 flore albo Hart. ✕	
15. elongatus Walst. et Kit. ☼ Hun.	598
The elongated Cytisus.	
16. multiflorus Lindl. ☼ Europe	599
The many-flowered Cytisus.	
<i>C. elongatus</i> Hort., not of Kit.	
<i>C. elongatus</i> ♂ multiflorus Dec.	
17. falcatus Walst. et Kit. ☼ Austria	599
The sickle-like-podded Cytisus.	
18. austriacus L. ☼ Austria f. 291.	599
The Austrian Cytisus.	
19. supinus Jacq. ✕ Europe f. 292.	599
The supine Cytisus.	
<i>C. lotoides</i> Pour.	
20. hirsutus L. ☼ Hungary -	599
The hairy Cytisus.	
<i>C. supinus</i> Bertol.	
<i>C. triflorus</i> Lam., not of L'Hérit.	
<i>C. Tournefortianus</i> Lois.	
21. capitatus Jacq. ☼ Italy -	600
The headed-flowered Cytisus.	
<i>C. hirsutus</i> Lam.	
<i>C. supinus</i> L.	
22. ciliatus Wahl. ☼ - - -	600
The ciliated-podded Cytisus.	
23. polýtrichus Bieb. ✕ Tauria -	600
The many-haired Cytisus.	
§ v. Lotoides Dec. - - - 600	
24. argentens L. ✕ France -	600
The silvery Cytisus.	
<i>Lotus argentens</i> Brot.	
25. calycinus Bieb. ✕ Caucasus -	601
The large-calyced Cytisus.	
<i>C. pauciflorus</i> Willd.	
26. nanus Willd. ✕ Levant fig. 293.	601
The dwarf Cytisus.	
§ vi. Chronanthus Dec. ☼ - - - 601	
27. orientalis Lois. ☼ Levant -	601
The Oriental Cytisus.	
<i>C. orientalis</i> , &c. - Greek and Vaid. Herb.	

	Page
App. i. Hardy Species of Cytisus not yet introduced.	601
<i>C. arboreus</i> Dec.	
<i>Spartium arboreum</i> Desf.	
<i>Gentia pendula</i> Poir.	
<i>C. Wédeni</i> Visiani.	
<i>C. albidus</i> Dec.	
<i>C. alpinus</i> L'Hérit.	
<i>C. supinus</i> Jacq.	
<i>C. hirsutus</i> Griseb.	
<i>C. hirsutus</i> and <i>C. supinus</i> Bieb.	
<i>C. macrospermum</i> Bégis.	
<i>var.</i> 2 glaber L.	
5 subspinescens.	
<i>C. serotinus</i> Kit. in Litt.	
<i>C. pygmaeus</i> Willd.	
<i>C. ponticus</i> Willd.	
<i>C. ponticus humifusus vulgno flore.</i>	
<i>C. caucasicus</i> Lois.	
<i>C. africanus</i> Lois.	
<i>C. africanus hirsutus angustifolius.</i>	
<i>C. procerus</i> Link.	
<i>Spartium procerum.</i>	
<i>C. 2 persicus</i> Burm.	
<i>Spartium persicum.</i>	
App. ii. Half-hardy Species of Cytisus.	602
☼ ☐ - - - 602	
<i>C. proliferus</i> L. ☼ ☐ fig. 204.	
<i>C. pallidus</i> Poir.	
<i>C. nubigenus</i> Link. ☼ ☐	
<i>Spartium nubigenum</i> Ait.	
<i>C. supranubium</i> L. Fil.	
<i>C. fedgrana</i> Lam.	
<i>C. bracteolatus</i> Hort.	
<i>C. tetragonocladius</i> Hort.	
<i>C. racemosus</i> Hort.	
App. iii. Anticipated hardy and half-hardy Species of Cytisus.	602
IX. ADENOCARPUS Dec. ☼ ☼ ☐ 603	
THE ADENOCARPUS.	
<i>Cytisus.</i>	
<i>Spartium.</i>	
<i>Genista.</i>	
1. hispanicus Dec. ☼ Spain -	603
The Spanish Adenocarpus.	
<i>Cytisus hispanicus</i> Lam.	
<i>Cytisus Anagyris</i> L'Hérit.	
2. intermedius Dec. ☼ Port. f. 295.	603
The intermediate Adenocarpus.	
<i>Cytisus complicatus</i> Brot.	
3. parvifolius Dec. ☼ France f. 296.	603
The small-leaved Adenocarpus.	
<i>Cytisus parvifolius</i> N. Du Ham.	
<i>Cytisus divaricatus</i> L'Hérit.	
<i>Cytisus complicatus</i> Dec.	
<i>Spartium complicatum</i> Lois.	
4. toulonensis Dec. ☼ Pyren. f. 297.	604
The Toulon Adenocarpus.	
<i>Cytisus toulonensis</i> Lois.	
<i>Spartium complicatum</i> Gouan.	
App. i. Half-hardy Species of Adenocarpus.	604
☼ ☐ - - - 604	
<i>A. frankenioides</i> Choix. ☼ ☐ - - - 604	
<i>Genista viscosa</i> Willd.	
<i>A. foliolosus</i> Dec. ☼ ☐ - - - 604	
<i>Cytisus foliolosus</i> Ait.	
X. ONONIS L. ☼ ☼ ☼ ☐ ☐ ☐ 604	604
THE RESTHARROW.	
<i>Ononis</i> and <i>Natrix</i> Monch.	
<i>Arche-bœuf</i> , <i>Bugrane</i> , Fr.	
<i>Hauwechel</i> , Ger.	

	Page
1. fruticòsa L. ☼ Europe fig. 298.	604
The shrubby Restharrow.	
Variety ☼ - -	605
2 microphýlla Dec. ☼	
O. fruticòsa Asso.	
2. rotundifòlia L. ☼ Alps fig. 299.	605
The round-leaved Restharrow.	
O. latifòlia Asso.	
Nátrix rotundifòlia Mærch.	
Variety - -	605
2 aristàta Dec.	
3. (r.) tribractèata Dec. ☼ Carinthia	605
The three-bracted-coloured Restharrow.	
O. rotundifòlia L.	
4. Nátrix Dec. ☼ — Europe fig. 300.	605
The Goat-root Restharrow.	
Nátrix plúguis Mærch.	
5. arenària Dec. ☼ — France -	606
The Sand Restharrow.	
Anònis spinis cárens latea minor Magn. Bot.	
6. cenisia L. ☼ France - -	606
The Mount Cenis Restharrow.	
O. cristata Mill. Dict.	
Variety ☼ - -	606
2 subaristàta Dec. ☼	
O. cenisia Asso.	
7. aragonénsis Asso. ☼ Spain - -	606
The Aragon Restharrow.	
E. dumosa Lapeyr.	
App. i. Other suffruticose Species of Ondònis.	
☼ — — - -	606
O. pedunculàris Lindl. fig. 301.	
O. crispa L.	
O. hispànica	
O. vaginàlis L. ☼ —	
O. arachnoidea Lapeyr.	
O. longifòlia Willd. ☼ —	
O. falcàta Willd. ☼ —	
O. ramosíssima Desf. ☼ —	
O. tridentàta L. ☼ —	
O. angustifòlia Lam. ☼ —	
O. foetida Schousb.	
XI. AMORPHA L. ☼ ☼ — ☼ —	606
THE AMORPHA, or Bastard Indigo.	
Bonafidia Neck.	
1. fruticòsa L. ☼ N. America f. 302.	607
The shrubby Amorpha.	
Wild Indigo.	
Faux Indigo. Fr.	
Strauchartiger Unform, Ger.	
Varieties ☼ - -	607
2 angustifòlia Pursh. ☼	
3 emarginàta Pursh. ☼	
4 Lewisii Lodd. Cat. ☼	
5 cærùlea Lodd. Cat. ☼	
2. (f.) glàbra Desf. ☼ — N. America	607
The glabrous Amorpha.	
3. (f.) nàna Nutt. ☼ — N. America	607
The dwarf Amorpha.	
A. microphýlla Pursh.	
4. (f.) fràgrans Swt. ☼ North America	fig. 303. 608
The fragrant Amorpha.	
A. nàna Sims.	

5. (f.) cròceo-lanàta Wats. ☼ —	Page
North America - fig. 304.	608
The Saffron-coloured woolly Amorpha.	
6. (f.) canéscens Nutt. ☼ — N. Am.	608
The canescent Amorpha.	
? A. pubéscens Pursh.	
XII. ROBINIA L. ☼ ☼ ☼	609
THE ROBINIA, or Locust Tree.	
Pseudacàcia.	
Robinier. Fr.	
Robinie, Ger.	
1. Pseud-Acàcia Lin. North America	pl. 81, 82. fig. 305. 609.
The common Robinia, or False Acacia	
Eschynómene Pseudacacia Roxb.	
Pseudacàcia odorata Mærch.	
Locust Tree, Amer.	
The Bastard Acacia.	
Acacia blanc, Carouge des Américains, Fr.	
Gemeine Acacia, Schotendorn, Ger.	
Varieties ☼ - -	609
2 flòre lùtea Dumont. ☼	
3 inermis Dec. ☼	
4 crispa Dec. ☼	
5 umbraculifera Dec. ☼ pl. 83.	
R. inermis Dum.	
6 tortuosa Dec. ☼ pl. 84.	
7 sophorafòlia Lodd. Cat. ☼	
8 amorphæfòlia Lk. ☼	
9 stricta Lk. ☼	
10 procèra Lodd. Cat. ☼	
11 péndula Ort. ☼	
12 monstròsa Lodd. Cat. ☼	
13 macrophýlla Lodd. Cat. ☼	
14 microphýlla Lodd. Cat. ☼	
R. angustifòlia Hort.	
15 spectàbilis Dum. ☼	
16 latisiliqua Prince's Cat. ☼	
2. [P.] viscòsa Vent. ☼ North America	pl. 85. fig. 336. 626
The clammy-barked Robinia.	
R. glutinosa Curt.	
R. montana Bartram.	
The Rose-flowering Locust.	
3. dùbia Fouc. ☼ Hybrid. - -	627
The doubtful Robinia.	
R. hybrida Audin.	
R. ambigua Poir.	
? R. chinata Mill. Diet.	
R. intermedia Soulange-Bodin.	
4. hispida L. ☼ N. Amer. f. 307.	627
The hispid Robinia, or Rose Acacia.	
R. rosea N. Du Ham.	
R. montana Bartr. Vor.	
Eschynómene hispida Roxb.	
Varieties ☼ ☼ - -	628
2 nàna Dec. ☼	
3 rosea Pursh. ☼	
4 macrophýlla Dec. ☼ pl. 86.	
R. grandiflora Hort.	
XIII. CARAGANA Lam. ☼ ☼ ☼	629
THE CARAGANA, or Siberian Pea Tree	
Robinia sp. L.	

- | | Page | | Page |
|-----------------------------------------------------|------|---------------------------------------------------------|------|
| 1. arborëscens Lam. γ Siberia pl. 87. | 629 | XIV. HALIMODE'NDRON Fisch. ♂ | 634 |
| The arborescent Caragana. | | THE HALIMODENDRON, or Salt Tree. | |
| <i>Robinia Caragana</i> L. | | <i>Halodendron</i> Dec., not of Petit | |
| <i>C. sibirica</i> Ray. | | Thouars. | |
| <i>Acacie de Sibérie, Robinie de Sibérie, Arbre</i> | | | |
| <i>aux Pois des Russes, Fr.</i> | | | |
| <i>Sibirische Erbsenbaum, Ger.</i> | | | |
| <i>Gorochnik, Russ.</i> | | | |
| Varieties ♀ - - - 629 | | | |
| 2 inermis Hort. ♀ | | 1. argenteum Dec. ♂ Siberia f. 315. | 634 |
| | | The silvery-leaved Halimodendron. | |
| | | <i>Robinia Halodendron</i> L. | |
| | | <i>Caragana argentea</i> Lam. | |
| | | Varieties ♂ - - - 634 | |
| 2. (a.) Altagana Poir. ♂ Siber. f. 308. | 630 | 1 vulgare Dec. ♂ | |
| The Altagana Caragana. | | 2 brachyscema Dec. ♂ | |
| <i>Robinia</i> Altagana Pall. | | | |
| <i>Caragana microphylla</i> Lam | | | |
| | | | |
| 3. (a.) microphylla Dec. ♂ Siberia | 630 | | |
| The small-leaved Caragana. | | 2. (a.) subvirëscens Dou. ♂ Siberia | 634 |
| <i>Robinia microphylla</i> Pall. | | The greenish Halimodendron. | |
| <i>Caragana Altagana var. Poir.</i> | | <i>Robinia triflora</i> L'Herit. | |
| | | <i>H. argenteum</i> β <i>subvirëscens</i> Dec. | |
| | | | |
| 4. (a.) Redowski Dec. ♂ Siberia | 630 | XV. CALOPHACA Fisch. ♂ - 635 | |
| Redowski's Caragana. | | THE CALOPHACA. | |
| Variety ♂ - - - 631 | | <i>Cytisus</i> Pall. and L. | |
| 2 præcox Fisch. ♂ | | <i>Colutea</i> Lam. | |
| | | <i>Adenocarpus</i> Spreng. | |
| | | | |
| 5. (a.) arenaria Donn and Sims. ♂ | | 1. wolgarica Fisch. ♂ Russia f. 316. | 635 |
| Siberia - - - fig. 309. | 631 | The Wolga Calophaca. | |
| The Sand Caragana. | | <i>Cytisus nigricans</i> Pall. Itin. | |
| | | <i>Cytisus pinnatus</i> Pall. Fl. Ross. | |
| | | <i>Cytisus wolgaricus</i> L. | |
| | | <i>Colutea wolgarica</i> Lam. | |
| | | <i>Adenocarpus wolgensis</i> Spreng. | |
| | | | |
| 6. frutescens Dec. ♂ Russia f. 310. | 631 | XVI. COLUTEA R. Br. ♂ - 635 | |
| The shrubby Caragana. | | THE COLUTEA, or Bladder Scana. | |
| <i>Robinia frutescens</i> L. | | | |
| <i>C. digitata</i> Lam. | | | |
| Varieties ♂ - - - 631 | | 1. arborëscens L. ♂ Europe f. 317. | 635 |
| 1 latifolia ♂ | | The arborescent Colutea. | |
| 2 angustifolia ♂ | | <i>C. hirsuta</i> Roth. | |
| | | | |
| 7. (f.) mollis Bess. ♂ Tauria - - - 631 | | 2. (a.) eruënta Lit. ♂ Levant f. 318. | 636 |
| The soft Caragana. | | The bloody-flowered Colutea, or Oriental Bladder Scana. | |
| <i>Robinia mollis</i> Bieb. | | <i>C. orientalis</i> Lam. | |
| <i>Robinia tomentosa</i> Fisch. | | <i>C. sanguinea</i> Pall. | |
| <i>C. f. var. mollis</i> Dec. | | <i>C. aptera</i> Schmidt. | |
| | | <i>C. humilis</i> Scop. | |
| | | | |
| 8. pygmaea Dec. ♂ Siberia f. 311. | 631 | 3. (a.) media Willd. ♂ Hybrid - - 635 | |
| The pygmy Caragana. | | The intermediate Colutea. | |
| <i>Robinia pygmaea</i> L. | | | |
| Variety ♂ - - - 632 | | | |
| 2 arenaria Fisch. ♂ | | | |
| | | | |
| 9. spinosa Dec. ♂ China - - - 632 | | 4. (a.) halëppica Lam. ♂ Syria - 637 | |
| The spiny Caragana. | | The Aleppo Colutea. | |
| <i>Robinia spinosa</i> L. | | <i>C. Pouchkii</i> Lit. | |
| <i>Robinia ferax</i> Pall. | | <i>C. istria</i> Mill. Diet. | |
| <i>Robinia spinosissima</i> Laxm. | | <i>C. procumbens</i> L'Herit. | |
| <i>C. ferax</i> Lam. | | | |
| | | | |
| 10. tragacanthoides Poir. ♂ Siberia | 632 | 5. nepalensis Hook. ♂ Nepal f. 319. | 637 |
| The Goat's-thorn-like Caragana. | | The Nepal Colutea. | |
| <i>Robinia tragacanthoides</i> Pall. | | | |
| <i>Robinia macracantha</i> Lodd. Cat. | | | |
| | | | |
| 11. jubata Poir. ♂ Siberia - - - 633 | | XVII. ASTRA'GALUS Dec. ♂ 637 | |
| The crested Caragana. | | THE MILK VETCH. | |
| <i>Robinia jubata</i> Pall. | | | |
| | | 1. Tragacantha L. ♂ S. Europe f. 320. | 637 |
| | | The Goat's Thorn Milk Vetch, or Great Goat's | |
| | | Thorn. | |
| | | <i>A. massiliensis</i> Lam. | |
| | | <i>A. creticus</i> - - - fig. 321. | 638 |
| | | | |
| 12. grandiflora Dec. ♂ Georgia - 633 | | App. i. Other ligneous Species of Astragalus | 638 |
| The great-flowered Caragana. | | in Cultivation. - - - 638 | |
| <i>Robinia grandiflora</i> Bieb. | | | |
| | | | |
| 13. Chamliqua Lam. ♂ China f. 311. | 633 | | |
| The Chamliqua, or Chinese, Caragana. | | <i>A. altaicus</i> Lodd. | |
| <i>Robinia Chamliqua</i> L'Herit. | | <i>A. aristatus</i> L'Herit. ♂ fig. 322. | |
| | | <i>A. involucratus</i> Lodd. fig. 325. | |
| | | <i>A. massiliensis</i> Lam. | |
| | | 2 <i>A. Tragacantha</i> var. with white flowers. | |

App. ii. *Hardy Species of Astragalus not yet introduced.* - - 638

A. aboriginorum Richards.

App. I. *Suffruticose hardy Species belonging to the Tribe Loteæ.* 2 2 2 2 - 639

Dorjencium Tourn. 2 2 2 2
 suffruticosum Mill. 2 2
Lotus Dorjencium L.
 rectum Ser. 2
Lotus rectus L.
 latifolium Willd. 2
 hirsutum Ser. 2
Lotus hirsutus L.
 tomentosum G. Don. 2
D. hirsutum var. *incanum* Ser.

App. II. *Half-hardy ligneous Species of Loteæ.* 2 2 2 2 2 2 - 639

Hovea R. Br. 2 - - - 639
Celsi Bonp. 2 fig. 524.
latifolia Lodd. 2 fig. 525.
lancofolata Sims. 2
Plagiobolium Smt. 2 - - - 639
flicifolium Sm. 2
chorozema-folium Smt. 2
Platylobium Sm. 2 - - - 639
formosum Sm. 2 fig. 526.
triangulare R. Br. 2
Bossiaea Vent. 2 - - - 640
ensata Sieb. 2
microphylla Smith. 2
Goddia latifolia Salisb. 2 - fig. 527. 640
pubescens Bot. Mag. 2
polysperma Bot. Rep. 2
Scottia dentata R. Br. 2 - - - 640
angu-stifolia Bot. Reg. 2
Templetonia glauca Sims. 2 - - - 640
Rafines Thunb. 2
triflora Bot. Mag. fig. 551, 552.
Vascosa Dec. 2 - - - 640
amplexicaulis Dec. 2
perfoliata Dec. 2
Borbonia L. 2 - - - 640
crenata L. 2
Achyronia villosa Wendl. 2 - - - 640
Liparia sphaerica L. 2 - - - 640
Priestleya Dec. 2 - - - 640
vestita Dec. 2
Crotalaria 2 - - - 640
pulchella Andr. 2
Hypocleptes obovatus Thunb. 2 - - - 640
Viborgia Spreng. 2 - - - 640
obcordata Thunb. 2 fig. 553.
Crotalaria floribunda Bot. Cab.
Loddigesia Sims. 2 - - - 641
oxalidifolia Bot. Mag. fig. 554.
Dichilus Dec. 2 - - - 641
Lebeckia Thunb. 2 - - - 641
sepaliata Thunb.
Sarcophyllum carnosum Thunb. 2 - - - 641
Aspalathus L. 2 - - - 641
callosa L. 2 fig. 556.
Requienia Dec. 2 - - - 641
sphaerosperma
Anthyllis L. 2 2 2 2 - - - 641
Burba Jovis L. 2 fig. 557.
Vulvararia *Burba Jovis* Lam.
Burba Jovis *argyrophylla* Moench.
cytisoides L. 2
Hermannia L. 2
Aspalathi Dec. 2
Spartium *criticum*.
erinacea L. 2
Gemista DuRoi.
splendens Willd.
indica Lour.
sericea Lag.
heterophylla L.
Medicago L. 2 - - - 642
arbores L. 2 fig. 558.
Melilotus *arbores* *Custogae* in Litt. 2 - - - 642

Lotus L. - - - - - 642
ereticus L. 2
anthylloides Vent. 2
atropurpureus Dec. 2
jacobinus L. 2 fig. 539.
Brous one. ii L.
spectabilis
Carmichaelia *australis* R. Br. 2 - - - 642
Psoralea L. 2 2 2
bituminosa L. 2
glandulosa L. 2 fig. 540, 541.
pubescens Balb. 2
picata Delil.
obtusifolia Dec.
Indigofera L. 2 2 2 - - - 642
denudata Jacq. 2 2 fig. 542.
anureba Ait. 2
atropurpurea Hamilt. 2
australis Willd. 2 fig. 545.
sylvatica Bieb. 2
I. angulata Bot. Reg.
Swainsonia *salish*. 2 - - - 645
galeifolia R. Br. 2 fig. 544.
coronilla-folia Salisb. 2
lessertia-folia Dec. 2
Lessertia Dec. 2 - - - 645
fruticosa Lind. 2 fig. 545.
Sutherlandia *frutescens* R. Br. 2 - - - 645
microphylla Burch. 2

Sect. III. HEDYSA'REÆ.

XVIII. CORONILLA Neck. 2 2 - 643

THE CORONILLA.

1. *E'merus* L. 2 S. Europe f. 347. 644
 The Scorpion Senna Coronilla.
E'merus major Mill. Icon., t. 132. f. 1.
E'merus minor Mill. Icon., t. 132. f. 2.
C. pauciflora Lam.
2. *Jancea* L. 2 France - f. 348. 644
 The rusby-branched Coronilla.

App. i. *Half-hardy ligneous Species of Coronilla.* 2 - - - 644





C. stipularis Lam. 2 - - - 649.
C. valentina L.
C. hispanica Mill.
C. pentaphylla Desf. 2
C. glauca L. 2 fig. 550.
C. argentea L. 2
C. multiflora Dec. 2





App. I. *Hardy suffruticose Species of Hedy-sarææ.* 2 - - - 645



Hedysarum *fruticosum* L. 2 - - - 645

App. II. *Half-hardy ligneous Species of Hedy-sarææ.* 2 ? 2 2 2 2 2 - 645

Hippocrepis *baleatica* Jacq. 2 - - - 645
Adesmia Dec. 2 - - - 645
microphylla Hook. fig. 555, 554.
Loudouia Hook. fig. 555, 556.
vistosa Gill. et Hook. fig. 557.
us-pallatensis Gill.
Uraria *Desr.* 2 - - - 645
arbores G. Don. 2
Hedysarum *arboresum* Hamilt.
Desmodium Dec. 2 - - - 645
retusum G. Don. 2
Hedysarum *retusum* Hamilt.
Dicerna *elegans* Dec. 2 - - - 646
Hedysarum *elegans* Lour.
Zornia *elegans* Pers.
Taverniera Dec. 2 - - - 646
nummularia Dec. 2 2
Hedysarum *oliværi* Spreng.
Lepedeza *Michx.* 2 - - - 646
frutescens Pers. 2
Flemingia *semialata* Karst. 2 - - - 646



<i>Ebenus cratica</i> L. 	Page 646
<i>Anthyllus cratica</i> Lam.	
<i>Alhagi Maurorum</i> Pourn. 	646
<i>Hedysarum Alhagi</i> L.	
<i>Alhagi rotundifera</i> Desf.	
<i>Ononis spinosa</i> Hasskq.	
<i>Manna hebraica</i> D. Don.	
camelorum 	
<i>Cliaanthus puniceus</i> Soland. ? 	fig. 358, 646
<i>Dona punicea</i> of G. and D. Don.	


Sect. IV. PHASEOLEÆ.     647


XIX. WISTARIA Nutt.   - 647


THE WISTARIA.





Glycine sp. L.
Thyrsanthus Elliot.
Kraumbia Rafin.



1. <i>frutescens</i> Dec.  N. Amer. f. 359. 647	
The shrubby Wistaria.	
<i>Glycine frutescens</i> L.	
<i>Apios frutescens</i> Pursh.	
<i>Anonymus frutescens</i> Walt.	
<i>Wistaria speciosa</i> Nutt.	
<i>Thyrsanthus frutescens</i> Elliot.	
<i>Phaseoloides</i> Hort. Angl.	
The Kidneybean Tree.	
2. <i>chinensis</i> Dec.  China f. 360. 648	
The Chinese Wistaria.	
<i>Glycine chinensis</i> Sims.	
<i>Glycine sinensis</i> Ker.	
<i>Wistaria Consequana</i> Loudon.	






App. i. Other Species of Wistaria.  648



W. floribunda Dec. 	648
<i>Dolichos polystachyus</i> Thunb.	
<i>Leijocle floribunda</i> Willd.	
<i>Dolichos japonicus</i> Spreng.	
<i>Findia</i> Kumpf.	

App. I. Suffruticose hardly or half-hardy Species of Phaseolæ.  - 649

<i>Lupinus arboreus</i> Sims. 	fig. 361.
<i>L. multiflorus</i> Desrous. 	
<i>L. albitrons</i> Benth.	
<i>L. Marshalliana</i> Sert. fig. 562.	
<i>L. canaliculatus</i> Sert. 	
<i>L. versicolor</i> Sert. 	
<i>L. pulchellus</i> Sert.	

App. II. Half-hardy Species of Phaseolæ.   - 649




<i>Dolichos lignosus</i> L. 	
<i>Pachyrhizus trifolius</i> Dec. 	
<i>Dolichos trifolius</i> Lour.	
<i>Mucuna macrocarpa</i> Wall. 	fig. 363.
<i>Erythrina Crista-galli</i> L. 	
<i>laurofolia</i> Jacq. 	
<i>E. Crista-galli</i> Bot. Reg.	


Sect. V. CASSIÆÆ   - 660


XX. GLEDITSCHIA L.  - 650







THE GLEDITSCHIA.


Acacia sp. Pluk.
Févier, Fr.
Gleditschie, Ger.


1. <i>triacanthos</i> L.  N. Am. pl. 88, 89. 650	
The three-thorned Gleditschia, or Honey Locust.	
<i>G. triacanthos</i> var. α <i>polysperma</i> Mart.	
<i>G. melloba</i> Walt.	
<i>G. spinosa</i> Du Ham.	
<i>Acacia triacanthos</i> Hort.	
<i>Acacia americana</i> Pluk.	
<i>Févier d'Amérique</i> , Fr.	
Thorny Acacia, Sweet Locust, United States.	
<i>Carouge à Mel</i> , Canada.	
Variety  - 650	
2 <i>inermis</i> Dec.  pl. 90, 91.	
<i>G. lævis</i> Hort.	




2. (t.) <i>monosperma</i> Wall.  N. America	Page fig. 364. 653
The one-seeded Gleditschia, or Water Locust.	
<i>G. carolinensis</i> Lam.	
<i>G. aquatica</i> Marsh.	
<i>G. triacantha</i> Gaert.	


3. (t.) <i>brachycarpa</i> Pursh.  North America	- 653
The short-fruited Gleditschia.	
<i>G. triacanthos</i> β Michx.	




4. <i>sinensis</i> Lam.  China pl. 92. 654	
The Chinese Gleditschia.	
<i>G. horrida</i> Willd.	
Varieties. 	
2 <i>inermis</i> N. Du Ham.  pl. 93.	
<i>G. japonica</i> Lodd.	
<i>G. javanica</i> Lam.	
3 <i>majior</i> Hort.  - 654	
<i>G. horrida majior</i> Lodd.	
4 <i>nana</i> Hort.  - pl. 94. 654	
<i>G. h. nana</i> Hort.	
5 <i>purpurea</i> Hort.  pl. 95. 654	
<i>G. h. purpurea</i> Lodd.	
Other Varieties - 654	

5. (s.) <i>macracantha</i> Desf.  China pl. 96. 654	
The long-spined Gleditschia.	
<i>G. ferox</i> Baudr.	
<i>Févier à grosses Epines</i> , Fr.	

6. (s.) <i>ferox</i> Desf.  China - 655	
The ferocious-prickled Gleditschia.	
<i>G. orientalis</i> Bosc.	
<i>Févier hérissé</i> , Fr.	


7. <i>caspia</i> Desf.  Caspian - 655	
The Caspian Gleditschia.	
<i>G. caspiana</i> Bosc.	
Variety  - 655	
2 <i>subvirescens</i> Hort. 	


App. i. Other Sorts of Gleditschia.  655

<i>G. micracantha</i> Hort. Soc. Gard. 	
<i>G. Boqui</i> Hort. Soc. Gard.	
<i>G. pumilox</i> Hort. Soc. Gard.	
<i>G. aquatica</i> Lodd.	
<i>G. monosperma</i> .	
<i>G. orientalis</i> Lodd. 	
<i>G. ferox</i> .	
<i>G. indica</i> Pers. 	

XXI. GYMNOCLADUS Lam.  656

THE GYMNOCLADUS.

1. <i>canadensis</i> Lam.  Canada pl. 97, 98. 656	
The Canada Gymnocladus, or Kentucky Coffee Tree.	
<i>Guitandou dioica</i> L.	
<i>Hyperanthura dioica</i> Vahl.	
Nicker Tree, and Stump Tree, United States.	
<i>Bonduc</i> , Chiquier, Fr.	
<i>Clécol</i> , Canadian.	
<i>Canadischer Schusserbaum</i> , Ger.	

XXII. CERCIS L.  - 657

THE JUDAS TREE.

Siliquastrum Tourn.
Gaïnier, Fr.

1. *Siliquástrum L.* † Europe pl. 99, 100. 657
 The common Judas Tree.
Siliquástrum orbiculátum Moench.
 Love Tree.
Gañier commun, Arbre de Judée, Fr.
Arbol d' Amor, Span.
Judasbaum, Ger.
Varieties † 𠄎 - 658
 2 parviflorum Dec. 𠄎
 3 flóre álbido Hort. †
 4 rósea Hort. †

2. canadensis L. † Canada pl. 101. 659
 The Canada Judas Tree.
Siliquástrum cordátum Moench.
 Red Bird Tree, Amer.
Bouton Rouge, Fr.
Varieties † - 659
 2 pubescens Ph. †
 3 Foreman's new variety. †

App. I. *Half-hardy Species of the Tribe Cassiæ.* 𠄎 𠄎 - 660

Cassápinia Ait. 𠄎
leibeckii Dec. 𠄎
Culia varia L'Herit. 𠄎
Zuccagnia Cav. 𠄎
punctata Cav. 𠄎
Ceratonia Siliqua L. † 𠄎 fig. 365, 366.
Casinospermum australe Cunningham.
Cassiá L. 𠄎
Barclayana Swt. 𠄎 fig. 367.
australis Hook. 𠄎 fig. 368.
lanecolata 𠄎
obovata 𠄎
Chamaefistula Don's Mill. 𠄎
corymbosa G. Don. 𠄎 fig. 370.
Cassia corymbosa Lam. 𠄎
Schötiá stipulata Ait. 𠄎
alata Thunb. 𠄎
latifolia Jacq. 𠄎 fig. 369.

App. I. *Other half-hardy ligneous Species of the Order Leguminæ.* - 661

† i. *Dalbergiæ.* - 661
Détris sericea G. Don. 𠄎
Pterocárus peitárius Dec. 𠄎
 † ii. *Mimósæ.* - 661
 𠄎 𠄎 𠄎 𠄎 𠄎 𠄎 𠄎 𠄎 𠄎

Prosopis L. 𠄎
glandulosa Torrey
Siliquástrum Dec. 𠄎
Lagoníchium Stephanianum Bieb. Supp. 𠄎
Acacia Stephaniana Bieb. Fl. Taur. 𠄎
Mimosa micrantha Vahl. 𠄎
Acácia Neck 𠄎 𠄎 𠄎 𠄎 𠄎

1. *Phyllodineæ.* 𠄎 𠄎 - 662
A. alata R. Br. 𠄎 fig. 371.
A. armata R. Br. 𠄎 fig. 372.
A. juniperina Willd. 𠄎 fig. 373.
Mimosa juniperina Vent.
Mimosa ulicifolia Wendl.
A. verticillata Sieb.
A. diffusa Ker. 𠄎 fig. 374, 375.
A. prostrata Lodd
A. stricta Willd. 𠄎 fig. 376, 377.
A. laurifolia Willd. 𠄎
Mimosa simplicifolia L.
A. melanoxylon R. Br. 𠄎 fig. 378, 379.
A. heterophylla Willd. 𠄎
Mimosa heterophylla Lam.
A. myrtifolia Willd. 𠄎 fig. 380.
Mimosa myrtifolia Sm.
A. lanata Lodd.
A. suaveolens Willd. 𠄎 fig. 381.
Mimosa suaveolens Sm.
A. oxycedrus Sibb. 𠄎 fig. 382, 385.
A. tarifolia Lodd.
 i. *verticillata Willd.* 𠄎 fig. 384.
var. 5 latifolia Dec. 𠄎

2. *Conjugato-pinnatæ.* 𠄎 𠄎 - 664
A. gummifera Willd. 𠄎
A. coronillifolia Desf. 𠄎
A. pulchella R. Br. 𠄎 fig. 385, 386.
A. detinens Burch. 𠄎
A. viridiramis Burch.
 3. *Spicifloræ.* 𠄎 𠄎 - 664
A. lophántha Willd. 𠄎 fig. 387.
A. lophántha Willd. 𠄎 fig. 387.
Mimosa Algiana Bot. Rep.
A. cæfra Willd. 𠄎
Mimosa cæfra Thunb.
A. álbida Delil. ? 𠄎

4. *Globifloræ.* 𠄎 𠄎 - 665
A. farnesiána Willd. † fig. 388.
Mimosa farnesiána L.
Mimosa scorpioides Forsk.
Galzia, Ital.
A. Cavénia Hook.
Mimosa Cavénia Moll.
A. nigricans R. Br. 𠄎 fig. 589, 590.
Mimosa nigricans Labill.
A. strigosa Link. 𠄎
A. ciliata R. Br.
A. glauca Willd. 𠄎
Mimosa glauca L.
A. Lamletiana D. Don. 𠄎
A. discolor Willd. 𠄎
A. angulata Desf. 𠄎
A. pubescens R. Br. 𠄎

A. Julibrissin Willd. † fig. 391.
Mimosa Julibrissin Scop.
Mimosa arbutæca Forsk.
A. decurrens Willd. 𠄎
Mimosa decurrens Vent.
A. mollissima Willd. 𠄎 fig. 592, 595.
A. decurrens var. ß mollis Bot. Reg.
A. mollis Swt.

A. dealbata Link. 𠄎 pl. 102.
A. affinis Hor.
The Black Wattle Mimosa, Van Diemen's Land.
A. mollis Wall. 𠄎

App. II. *Remarks on cultivating the half-hardy Leguminæ in British Gardens.* 667

Rosæcæ Dec. - 670
 † † † † † † † † † †
 † † † † † † † † † †

Sect. I. *AMYGDALÆ Juss.* - 671, 673
 I. *AMYGDALUS Tourn.* † 671, 673
 THE ALMOND TREE.

Amygdalophora Neck.
Amándier, Fr.
Mandcbaum, Ger.
 1. *nàna L.* 𠄎 Russia fig. 394, 395. 673
 The dwarf, or shrubby, Almond.
Prúnus inermis Gmel.
A. nana var. æ vulgaris Dec.
Zuerchenmandel, Ger.

Varieties. 𠄎
 2 *geórgica Dec.* 𠄎 - 673
 The Georgian dwarf Almond.
A. geórgica Desf.
 3 *campéstris Ser.* - 674
 The field dwarf Almond.
A. campéstris Bess.
A. Besseriana Schott.
 4 *incána Pall.* 𠄎 - 674
 The hoary-leaved dwarf Almond.
 ? 5 *sibirica Lodd.* 𠄎 - 674
 The Siberian Almond.

2. *communis* L. ♀ Mauritania pl. 103. 674
The common Almond Tree.

Varieties. ♂

1 *amara* Dec. ♀ - - 674
The bitter-kerneled common Almond Tree.

Amandier amer, Fr.
Gemeine Mandelbaum, Ger.

2 *dulcis* Dec. ♀ - - 675
The sweet-kerneled common Almond Tree.

Amandier à petits Fruits, *Amande douce*, Fr.

Süsse Mandel, Ger.

3 *âtre plèno* Baum. Cat. ♀ - 675

4 *follis variegatis* Baum. Cat. ♀ 675

5 *fragilis* Ser. ♀ - 675

The brittle-shell common Almond Tree.

A. fragilis Bell.

Amandier des Dames, N. Du Ham.

Coque molle, Fr.

Abellan, Provence.

6 *macrocarpa* Ser. ♀ - 675

The long-fruited common Almond Tree.

Amandier à gros Fruits, N. Du Ham.

Amandier Sultan, *Amandier des Dames*,

Amandier Pistache, Fr.

7 *persicoides* Ser. ♀ - 675

The Peach-like leaved common Almond Tree.

Other Varieties - - 675

3. *orientalis* Ait. ♂ Levant - - 679
The Eastern Almond Tree.

A. argentea Lam.

App. i. Other Species of *Amýgdalus*. ♀ ♂ 679

A. Tournefortii Bosc. ♂

A. cochinchinensis Lour. ♀

A. microphylla H. B. et Kunth. ♂

II. *PE'RSICA* Tourn. ♀ ♂ 671. 679

THE PEACH TREE.

Amýgdalus sp. L.

Trichocárpus Neck.

Pécher, Fr.

Pfirschenbaum, Ger.

1. *vulgáris* Mill. ♀ Persia pl. 104. 680
The common Peach Tree.

Amýgdalus Pérsica L.

Pêche duveteuse, Fr.

Pfirsche, Ger.

Varieties. ♀ ♂

1 The freestone common Peach Tree. ♀

Pêche, Fr.

2 The clingstone common Peach Tree. ♀

Pavie, Fr.

3 *flòre plèno* Hort. ♀
The double-flowering common Peach.

4 *álba* Lindl. ♂

The white-flowering common Peach.

5 *follis variegatis* Hort. ♀

The variegated-leaved Peach Tree.

6 *comprèssa* Hort. ♂ f. 397.

The flat Peach of China.

2. (v.) *la'vis* Dec. ♀ Persia - 680

The smooth-skinned Peach, or Nectarine Tree.

Amýgdalus Pérsica Lam.

Amýgdalus Pérsica Nectarina Ait.

Brugnon, *Pêche lisse*, Fr.

Varieties ♀ - - 680

1 The freestone Nectarine. ♀

Pêche lisse, Fr.

2 The clingstone Nectarine. ♀
Brugnon, Fr.

III. *ARMENIACA* Tourn. ♀ 671. 681

THE APRICOT.

Prúnus sp. L. and others.

Abricotier, Fr.

Aprikosenbaum, Ger.

1. *vulgáris* Lam. ♀ Asia pl. 105. 682
The common Apricot Tree.

Prúnus Armeniaca L.

Varieties ♀ - - 682

1 *ovalifolia* Ser. ♀ fig. 398.

The oval-leaved common Apricot Tree.

2 *cordifolia* Ser. ♀ fig. 399.

The heart-shaped-leaved common Apricot Tree.

3 *follis variegatis* Hort. ♀

The variegated-leaved common Apricot Tree.

4 *flòre plèno* Hort. ♀

The double-blossomed common Apricot Tree.

2. *dasy'carpa* Pers. ♀ f. 400, 401. 683

The thick-fruited Apricot Tree.

A. atropurpúrea Lois.

Prúnus dasy'carpa Ehrh.

Prúnus Armeniaca niger Desf.

The black Apricot.

Variety ♀ - - 683

2 *persicifolia* Lois. ♀ f. 402

The Peach-leaved thick-fruited Apricot Tree.

3. (v.) *sib'rica* Pers. ♀ Siberia pl. 106. 683

The Siberian Apricot Tree.

Prúnus sib'rica L.

4. (v.) *brigantiaca* Pers. ♀ Fr. f. 403. 684

The Briangon Apricot Tree.

Prúnus brigantiaca Vill.

IV. *PRUNUS* Tourn. ♀ ♂ 671. 684

THE PLUM.

Prunóphora Neck.

1. *spin'osa* L. ♀ Europe pl. 107. 684

The spiny Plum Tree, or common Sloe Thorn.

P. sylv'istris Fuch.

Blackthorn.

Prunier épineux, *Prunellier*, *L'pine noire*,

Mêre-du-Bois, Fr.

Schleodorn, *Schlen Pflaum*, Ger.

Varieties ♀ - - 685

1 *vulgáris* Ser. ♀

The common Sloe Thorn.

P. spin'osa Lois.

2 *follis variegatis* Ser. ♀

The variegated-leaved Sloe Thorn.

3 *microcarpa* Wallr. ♀

The small-fruited Sloe Thorn.

4 *macrocarpa* Wallr. ♀

The large-fruited Sloe Thorn.

5 *ovata* Ser. ♀

The ovate-leaved Sloe Thorn.

6 *flòre plèno* ♀

The double-flowered Sloe Thorn.

2. *insititia* L. † Barbary pl. 108. 687
 The engrafted Plum Tree, or Bullace Plum.
P. sylvestris præcox Attior Tourn.
P. sylvestris major Ray.
Prunier sauvage, Fr.
Kirschen Pflaume, Ger.
 Varieties † - - 687
 1 fructu nigro Hort. †
 The black-fruited, or common, Bullace.
 2 fructu luteo- albo Hort. †
 The yellowish-white-fruited Bullace.
 3 fructu rubro Hort. †
 The red-fruited Bullace.
 4 flore pleno Hort. †
 The double-flowered Bullace.
3. *domestica* L. † S. Europe - 687
 The domestic cultivated Plum Tree.
P. sativa Fuchs and Ray.
Gemeine Pflaume, Ger.
 Varieties † - - 688
 2 flore pleno Hort. †
 The double-blossomed Plum.
 3 foliis variegatis Hort. †
 The variegated-leaved Plum.
 4 myrobálana L. † pl. 109.
 The Myrobalan, or Cherry, Plum.
P. Myrobalan Du Ham.
P. myrobálana Lois.
P. cerasifera Ehrh.
Prunier Myrobalan, Cerisette, Fr.
Kirschpflaume, Ger.
 Subvariety. †
 5 m. foliis variegatis N. Du Ham.
 The variegated-leaved Myrobalan, or Cherry, Plum.
 6 armenioides Ser. †
 The Apricot-like Plum, or *Drap d'Or*.
4. *cándicans* Balb. ✽ f. 404, 405. 690
 The whitish-leaved Plum Tree.
5. *Cocomilla* Tenore. ✽ Calabria - 691
 The Cocomilla Plum Tree.
6. *marítima* Wangenheim. ✽ North America - 691
 The sea-side-inhabiting Plum Tree.
7. *pubescens* Poir. ✽ - 691
 The pubescent-leaved Plum Tree.
8. ? *divaricata* Led. ✽ Caucasus - 691
 The divaricate (? branched) Plum Tree.
- App. i. *Other Species of Prunus.* † ✽ 691
P. microcarpa Meyer. ✽
P. tomentosa Thunb. †
P. chinensis Blum. †
 Other Species.
- V. *CE'RASUS* Juss. † † † † ✽ ✽ ✽ ✽
 672. 692
- THE CHERRY.
Cerasus and *Laurocerasus* Tourn.
Prunus sp. L.
Cerisier, Fr.
Kirsche, Ger.
- § i. *Cerasophora* Dec. † ✽ ✽ - 692
 The Cherries cultivated in Gardens.
1. *sylvestris* Bauh. and Ray. † Europe pl. 110, 111. 693
 The wild black-fruited Cherry Tree.
C. avium Moench.
C. nigra Mill. Diet., not of Alt.
 † *Prunus avium* L.
Prunus avium var. *α ct β* Willd.
 † *Prunus nigricans* and *Prunus varia* Ehrh.
Gean, Bigarreau, Corone, Coroon, Small Black, Black Hertfordshire, Black Heart, Black Mazzard,
The Merry Tree of the Cheshire peasants, The Merries in Suffolk.
Mérisier, Mérisse grosse noire, Guignier, Bigarreautier, Heaumiér, Fr.
Süss Kirsche, Ger.
 Varieties - - - 693
 1 Mérisiers, or Merries. †
 2 Guigniers, or Geans. †
C. Juliána Dec.
C. decumána DeLaunay.
 3 Heaumiérs. †
 The Helmet-shaped Cherries.
C. Juliána var. *heaumiána* Dec.
 Variety of this race used for ornamental purposes:
durácina 2 flore pleno Hort. †
 The double-flowered wild black Cherry.
Mérisier Reunculier, Fr.
 4 Bigarreautiers. †
 The Bigarreau, or hard-fleshed Cherries.
C. durácina Dec.
2. *vulgáris* Mill. † Europe pl. 112. 693
 The Common Cherry Tree.
Prunus Cerasus L.
C. horténsis Pers.
C. caproniána Dec.
P. austera and *P. ácida* Ehrh.
Cherry, Kentish or Flemish Cherry, Morello, May Duke.
Cerise de Montmorency, Cerise de Paris, Cerise á Fruits ronds, Cerise du Nord, Cerisier, Griottier, Fr.
Saurc Kirsche, Ger.
 Varieties † - - 694
 2 flore semipleno Hort. †
 The semidouble-flowered common Cherry.
 3 flore pleno Hort. †
 The double-flowered common Cherry.
 4 persiciflora Hort. †
 The Peach-blossomed common Cherry.
 5 foliis variegatis Hort. †
 The variegated-leaved common Cherry.
3. (v.) *semperflórens* Dec. †
 pl. 113. 701
 The ever-flowering Cherry Tree.
Prunus semperflórens Ehrh.
Prunus serótina Roth.
The weeping Cherry, The Allsaints Cherry, Cerise de la Toussaint, Cerise de St. Martin, Cerise tardive, Fr.
4. *serruláta* G. Don. † China f. 406. 701
 The serrulated-leaved Cherry Tree.
Prunus serruláta Lindl.
The double Chinese Cherry, Yung-To, Chinese.

- | | Page |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 5. <i>Pseudo-Cerasus Lindl.</i> ♀ China
fig. 407. 701 | 706 |
| The False Cherry Tree.
<i>Prunus Pseudo-Cerasus Lindl.</i>
<i>Prunus paniculata</i> Ker. not of Thunb. | |
| 6. <i>Chamaecerasus Lois.</i> ♂ Sib. f. 408. 702 | 707 |
| The Ground Cherry Tree, or Siberian Cherry.
<i>C. intermedia</i> Lois.
<i>Prunus intermedia</i> Poir.
<i>Prunus fruticosus</i> Pall.
<i>C. pumila</i> C. Bauh.
<i>Chamaecerasus fruticosus</i> Pers. | |
| 7. <i>prostrata Ser.</i> ♂ Levant f. 409. 702 | 707 |
| The prostrate Cherry Tree.
<i>Prunus prostrata</i> Lab.
<i>Amygdalus incana</i> Pall.
<i>Prunus incana</i> Steven. | |
| 8. <i>persicifolia Lois.</i> ♀ N. America 702 | 707 |
| The Peach-tree-leaved Cherry Tree.
<i>Prunus persicifolia</i> Desf. | |
| 9. <i>borealis Michx.</i> ♀ N. Amer. f. 410. 703 | 707 |
| The North American Cherry Tree.
<i>Prunus borealis</i> Poir.
The Northern Choke Cherry, Amer. | |
| 10. <i>pumila Michx.</i> ♂ N. America 703 | 707 |
| The dwarf Cherry Tree.
<i>Prunus pumila</i> L.
<i>C. glauca</i> Moench.
Ragouminier, Nega, Mencl du Canada, Fr. | |
| 11. <i>depressa Ph.</i> ♂ North America 704 | 707 |
| The depressed, or prostrate, Cherry Tree.
<i>C. pumila</i> Michx., not the <i>Prunus pumila</i> L.
<i>Prunus Susquehænæ</i> Willd. | |
| 12. <i>pygmæa Lois.</i> ♂ North America 704 | 707 |
| The Pygmy Cherry Tree.
<i>Prunus pygmæa</i> Willd. | |
| 13. <i>nigra Lois.</i> ♀ Canada f. 411, 412. 704 | 707 |
| The black Cherry Tree.
<i>Prunus nigra</i> Ait.
<i>Prunus americana</i> Darlington. | |
| 14. <i>hymæalis Michx.</i> ♂ N. America 704 | 707 |
| The winter Cherry Tree.
<i>Prunus hymæalis</i> Michx.
The Black Choke Cherry. | |
| 15. <i>chicasa Michx.</i> ♂ N. America 705 | 707 |
| The Chicasaw Cherry Tree.
<i>Prunus chicasa</i> Pursh.
<i>Prunus insillita</i> Walt.
Chicasaw Plum, in Carolina. | |
| 16. <i>pubescens Ser.</i> ♂ N. America 705 | 707 |
| The pubescent Cherry Tree.
<i>Prunus pubescens</i> Pursh.
<i>Prunus sphaerocarpa</i> Michx., not of Swartz. | |
| 17. <i>pennsylvanica Lois.</i> ♀ N. America 705 | 707 |
| The Pennsylvanian Cherry Tree.
<i>Prunus pennsylvanica</i> L.
<i>Prunus lanceolata</i> Willd. | |
| 18. <i>japonica Lois.</i> ♂ Japan f. 413, 414. 705 | 707 |
| The Japan Cherry Tree.
<i>Prunus japonica</i> Thunb.
<i>Prunus sinensis</i> Pers.

Variety ♂ - - 706
2 multiplex Ser. ♂ figs. 415, 416.
<i>Amygdalus pumila</i> L. | |
| 19. <i>sinensis G. Don.</i> ♂ China f. 417. 706 | 707 |
| The Chinese Cherry.
<i>Prunus japonica</i> Ker. | |
| 20. <i>salicina G. Don.</i> ♂ China - 707 | 707 |
| The Willow-leaved Cherry Tree.
<i>Prunus salicina</i> Lindl.
Ching-Cho-Lee, or Tung-Choh-Lee, Chinese. | |
| Species belonging to the preceding Subdivision,
not yet introduced. ♀ ♂ ♂ 707 | |
| <i>C. Phòshia</i> Hamilt. ♀
<i>Prunus cerasoides</i> D. Don. | |
| <i>C. Páddum</i> Roxb. ♀
<i>C. glandulosa</i> Lois. ♂
<i>C. aspera</i> Lois. ♂
<i>C. incisa</i> Lois. ♂
<i>C. humilis</i> Moris. ♂ | |
| § ii. <i>Pádi reri</i> Ser. ♀ ♀ — - 707 | |
| 21. <i>Mahaleb Mill.</i> ♀ South Europe
pl. 114. 707 | 707 |
| The Mahaleb, or perfumed, Cherry Tree.
<i>Prunus Mahaleb</i> L.
<i>Bois de Sainte Lucie, Prunier odorant,</i>
Fr.

Varieties ♀ - - 707
1 <i>foliis variegatis</i> Hort. ♀
2 <i>fructu flavo</i> Hort. ♀
3 <i>latifolium</i> Hort. ♀ | |
| 22. <i>Párus Dec.</i> ♀ Europe pl. 115. 709 | 709 |
| The Bird Cherry Tree.
<i>Prunus Párus</i> L.
<i>Bird Cherry, Fowl Cherry,</i>
<i>Hag-berry,</i> Scot.
<i>Cerisier à Grappes, Mérisier à Grappes,</i>
<i>Laurier-Putier or Putier, Faux Bois de</i>
<i>Ste. Lucie,</i> Fr.
<i>Hag-bier,</i> Swedish.
<i>Traubeden Kirsche,</i> Ger.

Varieties ♀ - - 709
1 <i>vulgáris</i> Ser. ♀
<i>C. Párus</i> Dec.
2 <i>parviflora</i> Ser. ♀
3 <i>rúbra</i> Ser. ♀
<i>C. Párus fructu rubro</i> Dec.
4 <i>bracteosa</i> Ser. ♀ - - 702 | |
| 23. <i>virginiana Michx.</i> ♀ Virg. f. 418. 710 | 710 |
| The Virginian Bird Cherry Tree.
<i>Prunus rubra</i> Ait.
<i>Prunus arguta</i> Bigelow.
Wild Cherry Tree, Amer. | |
| 24. (v.) <i>serótina Lois.</i> ♀ North America
pl. 116. fig. 419. 712 | 712 |
| The late-flowering, or American, Bird Cherry
Tree.
<i>Prunus serótina</i> Willd.
<i>Prunus virginiana</i> Mill.

Variety ♀ - - 712
2 <i>retusa</i> Ser. ♀ | |
| 25. (v.) <i>Capóllin Dec.</i> ♀ Mexico
fig. 420. 713 | 713 |
| The Capóllin Bird Cherry Tree.
<i>Prunus virginiana</i> Flora Mexic. | |

26. (v.) canadensis Lois. ʎ	Canada	- 713	Page
The Canadian Bird Cherry Tree. <i>Prunus canadensis</i> Willd.			
27. nepalensis Ser. ʎ —	Nepal	- 713	
The Nepal Bird Cherry Tree.			
Species of Bird Cherry Trees which have not yet been introduced. ʎ — - 714			
C. paniculata Lois. ʎ	<i>Prunus paniculata</i> Thunb.		
C. acuminata Wall. ʎ	<i>C. mollis</i> Dougl. ʎ		
C. emarginata Dougl. ʎ	<i>C. capricida</i> G. Don. ʎ		
C. capricida G. Don. ʎ	The Goat-killing Bird Cherry Tree. <i>Prunus capricida</i> Wall. <i>Prunus undulata</i> Hamilt. <i>C. undulata</i> Dec.		
C. elliptica Lois. ʎ	<i>Prunus elliptica</i> Thunb.		
§ iii. Laurocerasi. ʎ —	- 714		
28. lusitânica Lois. ʎ	Portugal	pl. 117, 118. fig. 421.	714
The Portugal Laurel Cherry, or common Portugal Laurel. <i>Prunus lusitânica</i> L. The Cherry Bay. Azarciro, Portuguese.			
Variety ʎ	- 714		
2 Hixa Ser. ʎ	- 714		
<i>Prunus Hixa</i> Broussonet. <i>Prunus multiglandulosa</i> Cav.			
29. Laurocérusus Lois. ʎ	Asia Minor	fig. 422.	716
The Laurel Cherry, or common Laurel. <i>Prunus Laurocérusus</i> L. Cherry Bay, Cherry Laurel. Laurier au Lait, Laurier Amandier, Fr. Kirsche Lorbeer, Ger.			
Varieties	- 716		
2 variegata Hort. ʎ	- 716		
3 angustifolia Hort. ʎ	<i>Hortgia capensis</i> Hort.		
30. caroliniana Michx. ʎ	Carolina	fig. 423.	720
The Carolina Bird Cherry Tree. <i>Prunus caroliniana</i> Ait. <i>Prunus scampervirens</i> Willd. <i>Párus caroliniana</i> Mill. Dict. Wild Orange, Amer.			
App. i. Other Species of Cérusus.	- 721		
C. spherocarpa Lois.	<i>Prunus spherocarpa</i> Swartz.		
C. occidentalis Loitz.	<i>Prunus occidentalis</i> Swartz.		
Sect. II. SPIRÆÆ Dec.	672. 721		
VI. PURSHIA Dec. ʎ - -	672. 721		
THE PURSHIA. Tigarea Pursh, not of Aublet.			
1. tridentata Dec. ʎ	North America	fig. 424, 425.	721
The three-toothed-leaved Purshia. Tigarea tridentata Pursh.			

VII. KE'RRIA Dec. ʎ	- 672. 722	Page
THE KERRIA. <i>Ribus</i> L. <i>Corychorus</i> Thunb. <i>Spiræa</i> Camb.		
1. japónica Dec. ʎ	Japan	fig. 426. 722
The Japan Kerria. <i>Ribus japonicus</i> L. <i>Corychorus japonicus</i> Thunb. <i>Spiræa japonica</i> Camb.		
VIII. SPIRÆA L. ʎ —	- 672. 722	
THE SPIRÆA. <i>Spiræe</i> , Fr. <i>Spiræstaude</i> , Ger.		
§ i. Physocérpos Camb. ʎ	- 723	
1. opulifolia L. ʎ	North America	fig. 427, 428. 723
The Guelder-Rose-leaved Spiræa, or Virginian Guelder Rose. Nine Bark, Amer. Variety ʎ - - 723		
2 tomentella Ser. ʎ		
2. capitata Ph. ʎ	North America	- 723
The capitata-corymboid Spiræa. S. opulifolia var. Hook.		
3. monógyna Torrey. ʎ	N. America	723
The monogynous Spiræa.		
§ ii. Chamædryon Ser. ʎ —	- 724	
4. chamædrifolia L. ʎ	Siberia	f. 429. 724
The Germander-leaved Spiræa. S. caucasiensis Lour. Varieties ʎ - - 724		
1 vulgaris Camb. ʎ		
2 média Pursh. ʎ		
3 oblongifolia Camb. ʎ		
S. oblongifolia Waldst.		
4 subracemosa Ser. ʎ		
5 incisa Hort. ʎ		
? S. incisa Thunb.		
5. (c.) ulmifolia Scop. ʎ	Carinthia	fig. 430. 724
The Elm-leaved Spiræa. S. chamædrifolia Jacq. Variety ʎ - - 725		
2 phyllantha Ser. ʎ		
6. (c.) flexuosa Fisch. ʎ	- 725
The flexible-branched Spiræa. S. alpina Hort.		
7. (c.) eratægifolia Lk. ʎ	- 725
The Crataegus-leaved Spiræa.		
8. (c.) betulæfolia Pall. ʎ	N. Amer.	725
The Birch-leaved Spiræa. ? S. corymbosa Raf. ? S. eratægifolia Lk.		
9. cæna Waldst. et Kit. ʎ	Austria	- 725
The hoary-leaved Spiræa.		
10. trilobata L. ʎ - -	fig. 431.	725
The three-lobed-leaved Spiræa. S. triloba Don's Mill.		

- | | Page | | Page |
|--------------------------------------------------------------------------------------------------------------------------------------------------------|------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 11. alpina <i>Pall.</i> ❄ Siberia fig. 432. | 726 | 20. tomentosa <i>L.</i> ❄ Canada f. 444. | 730 |
| The Siberian alpine Spiræa. | | The downy Spiræa. | |
| 12. hypericifolia <i>Dec.</i> ❄ Europe and
America - - - fig. 433. | 726 | 21. lævigata <i>L.</i> ❄ Siberia fig. 445. | 731 |
| The Hypericum-leaved Spiræa. | | The smooth-leaved Spiræa. | |
| <i>Hypericum frutex</i> Hort. | | <i>S. altaicensis</i> Laxm. | |
| <i>Italian May.</i> | | <i>S. altaica</i> Pall. | |
| <i>Varieties.</i> ❄ | | 22. ariæfolia <i>Smith.</i> ❄ North America
fig. 446, 447. | 731 |
| 1 uralensis <i>Ser.</i> ❄ - - - | 726 | The White-Beam-tree-leaved Spiræa. | |
| <i>S. crenata</i> L. | | § iv. Sorbària <i>Ser.</i> ❄ - - - | 731 |
| <i>S. hypericifolia</i> Camb. | | 23. sorbifolia <i>L.</i> ❄ Siberia fig. 448. | 731 |
| 2 Plukenetiana <i>Ser.</i> ❄ - - - | 726 | The Sorbus-leaved Spiræa. | |
| <i>S. hypericifolia</i> L. | | <i>S. pinnata</i> Meuch. | |
| <i>S. h. var. β</i> Dec. | | <i>Variety</i> - - - | 732 |
| 3 acuta <i>Ser.</i> ❄ - - - fig. 434. | 727 | 2 alpina <i>Pall.</i> ❄ | |
| <i>S. acutifolia</i> Willd. | | <i>S. grandiflora</i> Hort. Brit. | |
| <i>S. sibirica</i> Hort. | | <i>S. Pallasi</i> Don's Mill. | |
| <i>S. ambigua</i> Pall. | | App. i. <i>Species or Varieties of Spiræa not
yet introduced.</i> ❄ - - - | 732 |
| 4 crenata <i>Ser.</i> ❄ - - - fig. 435. | 727 | <i>S. Blumei</i> G. Don. ❄ | |
| <i>S. obovata</i> Waldst. et Kit. | | <i>S. chamærifolia japonica</i> Blume. | |
| <i>S. h. γ</i> Dec. | | <i>S. lanceolata</i> Poir. ❄ | |
| <i>S. crenata</i> L. | | <i>S. argentea</i> Mutis. ❄ | |
| 5 savranica <i>Ser.</i> ❄ fig. 536. | 727 | <i>S. Thunbergii</i> Blume ❄ | |
| <i>S. savranica</i> Besser. | | <i>S. magellanica</i> Poir. ❄ | |
| <i>S. crenata</i> Pall. | | <i>S. japonica</i> Sieb. ❄ | |
| <i>S. h. var. β longifolia</i> Led. | | <i>S. Menziesii</i> Hook. ❄ | |
| 6 Besseriàna <i>Ser.</i> ❄ - - - | 725 | <i>S. Douglasii</i> Hook. ❄ | |
| <i>S. crenata</i> Besser. | | <i>S. callosa</i> Thunb. ❄ | |
| <i>S. savranica β Besseriàna</i> Don's
Mill. | | <i>S. capansa</i> Wall. | |
| 13. (<i>h.</i>) thaliectroides <i>Pall.</i> ❄ Dahuria
fig. 437. | 727. | <i>S. œrulescens</i> Poir. ❄ | |
| The Meadow-Rue-leaved Spiræa. | | <i>S. discolor</i> Pursh. ❄ | |
| <i>S. aquilegifolia</i> Pall. | | Sect. III. POTENTILLÆ <i>Juss.</i> - 733
(<i>Dryadac</i> Vent.) | |
| 14. pikowiensis <i>Besser.</i> ❄ Podolia - 728
The Pikow Spiræa. | | IX. RUBUS <i>L.</i> ❄ ❄ ❄ ❄ ❄ ❄ ❄
672. 733 | |
| 15. ecanothifolia <i>Horn.</i> ❄ - 728
The Ceanothus-leaved Spiræa. | | THE BRAMBLE.
<i>Ronce, Framboisier, Fr.</i>
<i>Himbeere, Brombeerstrauch, Ger.</i> | |
| 16. corymbosa <i>Raf.</i> ❄ Virg. f. 438. 728
The corymbose-flowering Spiræa. | | § i. <i>Leaves pinnate, of 3—7 Leaflets.</i> ❄ ❄ 735 | |
| <i>Variety</i> - - - - - 728 | | 1. suberectus <i>Anders.</i> ❄ Britain 735. 746
The sub-erect Bramble.
<i>R. nessensis</i> Hall.
<i>R. plicatus</i> W. et N.
<i>R. corylifolius</i> Wahlenb. | |
| 2 sororia - - - - -
<i>S. sororia</i> Penny. | | 2. affinis <i>Weibe et Nees.</i> ❄ Germany 735
The related Bramble.
<i>R. collinus</i> Dec. } according to Lindley.
<i>R. nitidus</i> Smith }
<i>R. plicatus</i> Borrer } | |
| 17. vacciniifolia <i>D. Don.</i> ❄ Nepal
fig. 439. 728
The Vaccinium-leaved Spiræa. | | <i>Variety</i> - - - - - 735
2 bracteatus <i>Ser.</i>
<i>R. α, γ, et δ</i> W. et N. | |
| 18. bella <i>Sims.</i> ❄ Nepal fig. 440. 729
The beautiful Spiræa. | | 3. fissus <i>Lindl.</i> ❄ Britain - - - 735
The cleft Bramble.
<i>R. fastigiatus</i> Lindl. Synops., ed. 1., not of Weibe
and Nees. | |
| § iii. <i>Spiræria</i> <i>Ser.</i> ❄ - - - 729 | | 4. micranthus <i>D. Don.</i> ❄ Nepal
fig. 449, 450, and 450 a. 735, 736
The small-flowered Bramble.
<i>R. pauciflorus</i> Lindl. | |
| 19. salicifolia <i>L.</i> ❄ Siberia fig. 443. 729
The Willow-leaved Spiræa.
<i>Spiræa frutex</i> Hort.
<i>Bridewort, Queen's Needle-work.</i> | | | |
| <i>Varieties</i> ❄ - - - - - 729 | | | |
| 1 carnea <i>Ait.</i> ❄ f. 443. | | | |
| 2 alpestris <i>Pall.</i> ❄ | | | |
| 3 paniculata <i>Willd.</i> ❄
<i>L. alba</i> Ehrh. | | | |
| 4 latifolia <i>Willd.</i> ❄ f. 441.
<i>S. obovata</i> Raf.
<i>S. carpiniifolia</i> Willd. | | | |
| 5 grandiflora ❄ f. 442.
<i>S. grandiflora</i> Lodd. | | | |

	Page		Page
5. <i>dístans</i> <i>D. Don.</i> ☼ Nepal - - 736 The distant-leafleted Bramble. <i>R. pinnátus</i> <i>D. Don.</i>	736	17. <i>corylifólius</i> <i>Smith.</i> ✱ Europe fig. 457. 740. 746 The Hazel-leaved Bramble. <i>R. vulgáris</i> <i>W. & N.</i> <i>R. nemorósus</i> <i>Heyne.</i>	740
6. <i>strigósus</i> <i>Michx.</i> ☼ North America 736 The strigose Bramble. <i>R. pennsylvánicus</i> <i>Poir.</i>	736	<i>Varieties</i> ✱ - - - 740 2 <i>cánus</i> <i>Wallr.</i> ✱ 3 <i>glandulósus</i> <i>Wallr.</i> ✱ <i>R. glandulósus</i> <i>Spreng.</i>	740
7. <i>occidentális</i> <i>L.</i> ☼ North America fig. 451. 736. 746 The Western, or American, Bramble. <i>R. virginianus</i> <i>Hort.</i> <i>R. idæus fructu nigro</i> <i>Dill.</i>	746	British kinds of <i>Rubus</i> which, according to Dr. Lindley, may be associated with <i>R. corylifólius</i> <i>Sm.</i> , either as related species, or as varieties:— - - 741 <i>R. macrophýllus</i> <i>W. et N.</i> The large-leaved Bramble. <i>R. carpinifólius</i> <i>W. et N.</i> The Horn beam-leaved Bramble. <i>R. fúscó-ater</i> <i>W. et N.</i> The brownish-black Bramble. <i>R. Köhleri</i> <i>W. et N.</i> Köhler's Bramble. <i>R. glandulósus</i> <i>Smith.</i> The glandulous-bristled Bramble. <i>R. rúdis</i> <i>W. et N.</i> The rough Bramble. <i>R. celnátus</i> <i>Lindl.</i> <i>R. diversifólius</i> <i>Lindl. Synops. ed. 1.</i> The diverse-leaved Bramble. <i>R. diversifólius</i> <i>W. et N.</i>	741
8. <i>ásper</i> <i>D. Don.</i> ☼ Nepal - - 737 The rough-branched and petioled Bramble.	737	18. (c.) <i>agréstis</i> <i>Wallst. et Kit.</i> ✱ Hung. 741 The Field Bramble.	741
9. <i>idæus</i> <i>L.</i> ☼ Europe, Asia, Africa, and America - - fig. 452. 737. 746 The Mount Ida Bramble, or common Raspberry. <i>R. frambésianus</i> <i>Lam.</i> <i>Gemeine Brombeere</i> , Ger. <i>Batos idæia</i> , Greek. <i>Raspis, Framboise, Hinde-berry</i> , Gerard. <i>Varieties</i> ☼ - - - 737 With red fruit. ☼ With yellow fruit. ☼ With white fruit. ☼ 2 <i>microphýllus</i> <i>Wallr.</i> ☼ Varieties cultivated in British Gardens.	737	19. <i>spectábilis</i> <i>Ph.</i> ☼ North America fig. 458. 741. 746 The showy-flowered Bramble. <i>R. ribifólius</i> <i>Willd.</i>	741
§ ii. <i>Leaves digitate, of 3—5 Leaflets.</i> ☼ ✱ 2 - - - 738	738	20. <i>ulmifólius</i> <i>Schott.</i> ✱ Gibraltar - 741 The Elm-leaved Bramble.	741
10. <i>laciniátus</i> <i>W.</i> ✱ . . . f. 453. 738. 746 The cut-leaved Bramble.	746	21. <i>Linkianus</i> <i>Ser.</i> ✱ . . . - - 742 Link's Bramble. <i>R. paniculátus</i> <i>Schlecht.</i>	742
11. <i>cæ'sius</i> <i>L.</i> ✱ North-eastern Asia fig. 454. 739. 746 The grey Bramble, or Dewberry. <i>Varieties</i> ✱ - - - 739 2 <i>arvénis</i> <i>Wallr.</i> ✱ <i>R. pseudo-cæ'sius</i> <i>Weihe.</i> 3 <i>grandiflórus</i> <i>Ser.</i> ✱ 4 <i>parvifólius</i> <i>Wallr.</i> ✱ f. 455. 5 <i>fóliis variegátis</i> <i>Hort.</i> ✱	739	22. <i>fruticósus</i> <i>L.</i> ✱ Europe fig. 459. 742. 746 The shrubby Bramble, or common Blackberry. <i>R. discolor</i> and <i>R. abruptus</i> <i>Lindl. Synops., ed. 1.</i> <i>Varieties</i> ✱ - - - 742 2 <i>pompónius</i> <i>Ser.</i> ✱ f. 460. <i>R. fruticósus</i> <i>W. et N.</i> 3 <i>tauricus</i> <i>Hort.</i> ✱ 4 <i>flóre rósea plèno Baum. Cat.</i> ✱ The double pink-flowered Bramble. 5 <i>fóliis variegátis</i> ✱ The variegated-leaved Bramble. 6 <i>leucocárpus</i> <i>Ser.</i> ✱ 7 <i>inermis</i> <i>Ser.</i> ✱ 8 <i>dalmáticus</i> <i>Tratt. Ros.</i> ✱ 9 <i>cóncolor</i> <i>Wallr.</i> ✱ 10 <i>glandulósus</i> <i>Wallr.</i> ✱ 11 <i>rharnifólius</i> <i>W. & N.</i> ✱ The Buckthorn-leaved Bramble. <i>R. cordifólius</i> <i>Lindl. ? Weih.</i> 12 <i>leucóstachys</i> <i>Schl.</i> ✱ The white-spiked Bramble.	742
12. <i>hírtus</i> <i>W. et K.</i> ✱ Hungary - 739 The hairy Bramble. <i>R. villosus</i> <i>Ait.</i> <i>var. glandulosus</i> <i>Ser.</i> <i>R. glandulosus</i> <i>Bell.</i> <i>R. hybridus</i> <i>Vill.</i>	739	<i>R. tomentosus</i> <i>Weihe.</i> ✱ - - - fig. 461. 743 <i>R. fastigiátus</i> <i>Weihe.</i> ✱ - - - 743 <i>R. filicifólius</i> <i>Weihe.</i> ✱ - - - 743 <i>R. Menkii</i> <i>Weihe.</i> ✱ - - - 743 <i>R. Schlechtendáhlíi</i> <i>Weihe.</i> ✱ - - - 743 <i>R. Schleicheri</i> <i>Weihe.</i> ✱ - - - 743 <i>R. hórridus</i> <i>Weihe.</i> ✱ - - - 748 <i>R. nitidus</i> <i>Weihe.</i> ✱ - - - 745 <i>R. rubricáulis</i> <i>Weihe.</i> ✱ - - - 745	743
13. <i>Sprengèlii</i> <i>Weihe.</i> ✱ Germany - 740 Sprengel's Bramble. <i>R. vulpinus</i> <i>Desf.</i> <i>R. villosus</i> <i>β vulpinus</i> <i>Ser.</i>	740		
14. <i>dumetórum</i> <i>W. et N.</i> ✱ Britain 740 The Bramble of Thickets.	740		
15. <i>foliolósus</i> <i>Don.</i> ✱ Nepal - - 740 The leafy Bramble. <i>R. microphýllus</i> <i>Don.</i>	740		
16. <i>flagelláris</i> <i>Willd.</i> ✱ North America 740 The Rod-like, or Runner, Bramble. <i>Variety</i> ✱ - - - 740 2 <i>inermis</i> <i>Ser.</i> ✱ <i>R. inermis</i> <i>Willd.</i>	740		

- | | Page |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|
| 23. <i>sánctus</i> Schreb. ✱ East - - 744
The holy Bramble.
R. <i>obtusifolius</i> Willd. | 744 |
| 24. <i>canéscens</i> Dec. ✱ Italy - - 744
The grey Bramble. | 744 |
| 25. <i>setósus</i> Big. ✱ North America - 744
The bristly-stemmed Bramble. | 744 |
| 26. <i>argútus</i> Lk. ✱ North America - 744
The sharp-toothed-toothed Bramble. | 744 |
| 27. <i>cuneifólius</i> Ph. ✱ N. America 745
The wedge-shaped-leafted Bramble.
R. <i>parvifolius</i> Walt. | 745 |
| 28. <i>hispídus</i> L. ✱ Canada fig. 462. 745
The hispid-stemmed Bramble.
R. <i>trivialis</i> Michx.
R. <i>poenambens</i> Mühl.
R. <i>flagellaria</i> Willd. | 745 |
| 29. <i>lanuginósus</i> Steven. ? ✱ Caucasus 745
The woolly Bramble. | 745 |
| 30. <i>canadáensis</i> L. ✱ Canada - - 745
The Canadian Bramble. | 745 |
| § iii. <i>Leaves lobed, not pinnate or digitate.</i>
- - - 745 | 745 |
| 31. <i>odorátus</i> L. ✱ North America
fig. 464, 745. 746
The sweet-scented Bramble.
R. <i>occidentális</i> Hort.
The Virginian Raspberry, The flowering
Raspberry. | 746 |
| 32. <i>nutkáus</i> Moc. ✱ California
fig. 464. 745, 746
The Nootka Sound Bramble.
R. <i>odorátus</i> Hort., not of L. | 746 |
| App. i. <i>Species and Varieties of Rubus best
deserving of Cultivation in British Gardens,
as ornamental Shrubs</i> - - - 746 | 746 |
| App. ii. <i>Other Sorts of shrubby Rubuses.</i>
- - - 746 | 746 |
| R. <i>macroptálus</i> Dougl. MS. fig. 165.
R. <i>deliciosus</i> Torrey.
R. <i>áilíceus</i> Smith ✱
? R. <i>cordifólius</i> D. Don
R. <i>acuminátus</i> Smith
R. <i>betulinus</i> D. Don
R. <i>reflexus</i> Ker. ✱ fig. 466.
R. <i>obovátus</i> Ait.
R. <i>rosafólius</i> Smith ✱
R. <i>r. coronáris</i> Sims ✱ fig. 467. | |
| X. POTENTILLA L. ✱ ✱ - 747
THE POTENTILLA, or Shrubby Cinqufoil. | 747 |
| 1. <i>fruticósa</i> L. ✱ Europe fig. 468. 747
The Shrubby Potentilla, or Cinqufoil.
<i>Varieties.</i> ✱ ✱
2 <i>dahúrica</i> Ser. ✱ - 747
P. <i>dahurica</i> Nestl.
P. <i>fruticosa</i> B. Lelom.
3 <i>tenuiloba</i> ✱ fig. 469. 748
P. <i>fruticosa</i> B. Nestl.
P. <i>floribunda</i> Pursh.
P. <i>tenuifolia</i> Sch'chtend. | 747
747
748 |
| 2. <i>glabra</i> Lodd. ✱ Siberia fig. 470. 748
The glabrous Potentilla.
P. <i>fruticosa</i> álba Busch. | 748 |
| 3. <i>Salesóvii</i> Steph. ✱ Siberia - 748
Salesovius's Potentilla.
Sect. IV. ROSEÆ Dec. - 672. 748 | 748 |
| XI. RO'SA Tourn. ✱ ✱ ✱ ✱ ✱ ✱ ✱
- - - 672. 748 | 748 |
| THE ROSE TREE.
<i>Rhodóphora</i> Neck.
<i>Rosier</i> , Fr.
<i>Rosenstock</i> , Ger.
<i>Roozboom</i> , Dutch.
<i>Rozajó</i> , Ital.
<i>Rosal</i> , Spanish.
<i>Roseira</i> , Portuguese.
§ i. <i>Feróces</i> Lindl. ✱ - - 750 | 750 |
| 1. <i>férox</i> Laur. ✱ Caucasus fig. 471. 750
The fiercely-prickled Rose.
R. <i>kamtschática</i> Med.
R. <i>kamtschática</i> B. <i>férox</i> Ser.
R. <i>echináta</i> Dupont.
<i>Variety</i> ✱ - - 750
2 <i>nítens</i> Lindl. ✱ | 750 |
| 2. (f.) <i>kamtschática</i> Vent. ✱ Kamtschatka
fig. 472. 750
The Kamtschatka Rose.
§ ii. <i>Bracteáta</i> . ✱ - - 750 | 750 |
| 3. <i>bracteáta</i> Wendl. ✱ China f. 473. 750
The large-bracted Rose.
<i>Lord Macartney's Rose.</i>
<i>Varieties</i> ✱ - - 751
2 <i>scabriceális</i> Lindl. ✱
R. <i>b. véra</i> Lodd. ✱
R. <i>b. flore pléno</i> Lodd. ✱ | 751 |
| 4. (b.) <i>microphýlla</i> Roxb. ✱ China
fig. 474. 751
The small-leafted Rose.
<i>Hoi-tong-hong</i> , Chinese. | 751 |
| 5. (b.) <i>involveráta</i> Roxb. ✱ Nep. f. 475. 751
The involucred-corymbed Rose.
R. <i>Lindleyana</i> Tratt. Ros.
R. <i>palustris</i> Buchan.
§ iii. <i>Cinnamómea</i> Lindl. ✱ ✱ - 751 | 751 |
| 6. <i>lúcida</i> Ehrh. ✱ N. America f. 476. 752
The shining-leaved Rose.
R. <i>rúbra lúcida</i> Rossig.
<i>Rose Turneps.</i>
<i>Rosier a Feuilles de Frénc</i> , Fr. | 752 |
| 7. (l.) <i>nítida</i> W. ✱ Newfoundl. f. 477. 752
The glossy-leaved Rose.
R. <i>Rubra vulcánica</i> Thory.
<i>The dwarf Labrador Rose.</i> | 752 |
| 8. (l.) <i>Rápa Bose</i> . ✱ N. Amer. f. 478. 752
The Turnip-fruited Rose.
R. <i>turgida</i> Pers.
R. <i>fraxinifolia</i> Dumont. | 752 |
| 9. <i>Woodsii</i> Lindl. ✱ North America 753
Woods's Rose.
R. <i>luta nigra</i> Pronv. | 753 |
| 10. <i>frutetórum</i> Bess. ✱ Volhynia - 753
The Coppice Rose. | 753 |
| 11. <i>carolina</i> L. ✱ Carolina - - 753
The Carolina Rose.
R. <i>virginiana</i> Du Roi.
R. <i>palustris</i> Marsh.
R. <i>corymbosa</i> Ehrh.
R. <i>pennsylvánica</i> Michx.
R. <i>hudsoniana</i> Bed.
R. <i>cor. liniana</i> Regel. | 753 |

	Page		Page
12. Lindlèyi Spreng. ☼ North America	753	14 turbinàta Desv. ☼	
Lindley's Carolina Rose.		R. inermis Delam., not of Ait.	
R. laxa Lindl.		R. a. flore pleno Godefroy.	
R. carolina e Ait.		R. a. multiplex Ser.	
R. carolina pimpinellifolia Andr.		R. mixta Tratt.	
13. parviflora Ehrh. ☼ N. Amer. f. 479.	753	21. suavis Willd. ☼	fig. 484. 756
The small-flowered, or Pennsylvania, Rose.		The sweet Rose.	
R. humilis Marsh.		22. acicularis Lindl. ☼	- - - 756
R. caroliniana Michx.		The needle-prickled Rose.	
R. carolina y et e Ait.		R. alpina e aculeata Ser.	
Pennsylvania Rose, Lawr.		23. lutescens Pursh. ☼ North America	fig. 485. 756
Variety ☼ - - - - -	753	The yellow American Rose.	
2 flore pleno Red. ☼ fig. 479.		R. hispida Curt.	
14. fraxinifolia Bork. ☼ North America	fig. 480. 754	24. sulphurea Ait. ☼ Levant f. 486.	756
The Ash-leaved Rose.		The sulphur-coloured-flowered Rose.	
R. virginiana Mill. Dict.		R. hemisphærica Herm.	
R. blanda e Sol. MSS.		R. glaucophylla Ehrh.	
R. corymbosa Bosc.		R. lutea flore pleno Ray.	
R. alpina β Ait.		R. lutea Brot.	
R. alpina lævis Red.		The double yellow Rose.	
15. cinnamomea Besl. ☼ Europe	fig. 481. 754	25. spinosissima L. ☼ Europe f. 487.	757
The Cinnamon-scented Rose.		The most spiny, or Scotch, Rose.	
R. fecundissima Munch.		Varieties - - - - -	757
R. majalis Herm.		26. hibernica Smith. ☼ Ireland	- 757
16. majalis Retz. ☼ Sweden - - - - -	754	The Irish Rose.	
The May Rose.		27. oxyacantha Bieb. ☼ Siberia	- 757
R. mutica Fl. Dan.		The sharp-prickled Rose. ☼	
R. spinosissima Gorter.		28. sanguisorbifolia Don. ☼	758
R. collincola Ehrh.		The Burnet-leaved Rose.	
R. cinnamomea Eng. Bot.		R. spinosissima var. sanguisorbifolia Lindl.	
17. Dicksoniana Lindl. ☼ Ireland - 754		R. spinosissima var. macrophylla Ser.	
Dickson's Rose.		29. grandiflora Lindl. ☼ Siberia f. 488.	758
18. taurica Bieb. ☼ Tauria - - - - -	754	The large-flowered Rose.	
The Taurian Rose.		R. pimpinellifolia Bieb.	
19. dahurica Pall. ☼ Dahuria - - - - -	754	30. myriacantha Dec. ☼ France f. 489.	758
The Dahurian Rose.		The myriad prickled Rose.	
§ iv. Pimpinellifoliæ Lindl. ☼ ☼ - 755		R. parvifolia Pall.	
20. alpina L. ☼ S. Europe fig. 482.	755	R. provincialis Bieb.	
The Alpine Rose.		R. spinosissima var. η myriacantha Ser.	
R. rupëstris Crantz.		31. involuta Smith. ☼ Hebrides - 758	
R. monspeliaca Gouan.		The involute-petaled Rose.	
R. inermis Mill. Dict.		R. nivalis Donn.	
R. hybrida Vill. Dauph.		32. reversa Waldst. et Kit. ☼ Hungary	758
R. lagenaria Vill.		The reversed-prickled Rose. . . .	
R. biflora Krok.		33. Sabini Woods. ☼ Britain - - - 758	
Varieties ☼ - - - - -	755	Sabine's Rose.	
2 pyrenæica Lindl. ☼		R. sabini β Lindl.	
R. pyrenæica Gouan.		34. Doniana Woods. ☼ Scotland - 758	
R. alpina Jacq.		Don's Rose.	
R. hispida Krok.		§ v. Centifoliæ Lindl. ☼ ☼ ☼ - 759	
R. turbinata Vill. Dauph.		35. damascena Mill. ☼ Syria f. 490.	759
R. alpina β Dec.		The Damascus, or Damask, Rose.	
R. hispida ?		R. bérigica Mill. Dict.	
R. alpina coronata Desv.		R. calandærum Munch.	
3 pendulina Lindl. ☼		R. biceps Poir.	
R. pendulina Ait.		Rose a quatre Saisons, Fr.	
R. a. latifolia Ser.		Varieties - - - - -	759
4 pimpinellifolia Lindl. ☼		36. centifolia Lin. ☼ Caucasus f. 491.	760
R. glandulosa Bell.		The hundred-petaled, Provence, or Cabbage Rose.	
R. pygmaea Bieb.		R. provincialis Mill. Dict.	
R. pyrenæica β Smith.		R. polyanthos Rossig.	
5 lagenaria Ser. ☼		R. caryophyllæ Poir.	
6 sorbinella Ser. ☼		R. unguiculata Desf.	
7 hispidella Ser. ☼		R. varians Pohl.	
8 lævis Ser., not of Desv. or Red. ☼	fig. 483.	f	
R. Sanguisorba majoris, &c. Dill.			
R. alpina glabra Desv.			
R. a. vulgaris Red.			
9 pyriformis Ser. ☼			
10 setosa Ser. ☼			
R. a. hircina Desv.			
11 globosa Desv. ☼			
R. canina ambigua Desv.			
12 hellebörina Ser. ☼			
13 pilosula Ser. ☼			

- | | Page | | Page |
|--------------------------------------------|------|-------------------------------------------------------------|------|
| <i>Varieties</i> ☼ - - - | 760 | 41. <i>gracilis</i> Woods. ☼ Britain - | 763 |
| 1 provinciális Mill. ☼ | | The slender Rose. | |
| The Province, or Cabbage, Roses. | | R. villosa Smith. | |
| 2 muscosa Mill. ☼ fig. 492. | | 42. tomentosa Smith. ☼ Europe - | 763 |
| The Moss Roses. | | The tomentose, or woolly-leaved, Rose. | |
| 3 pomponia Dec. ☼ | | R. villosa Ehrh. | |
| The Pomponé Roses. | | R. mollissima Börk. | |
| 4 bipinnata Red. ☼ | | R. diibia Wibel. | |
| | | R. villosa β Huds. | |
| | | <i>Variety</i> ☼ - - - | 763 |
| 37. gállica L. ☼ Europe fig. 493. | 760 | 2 scabriuscula Sm. ☼ | |
| The French Rose. | | R. fa'tida Batard. | |
| R. centifolia Mill. Dict. | | 43. Sherárdi Davies. ☼ England - | 764 |
| R. sylvatica Gater. | | Sherard's Rose. | |
| R. rubra Lam. | | R. subglabrosa Sm. | |
| R. holosericea Rossig. | | R. tomentosa var. α et η Woods. | |
| R. belgica Brot. Fl. Lus. | | 44. sylvestris Lindl. ☼ England - | 764 |
| R. blanda Brot. | | The Wood Rose. | |
| Rose de Provins, Fr. | | R. tomentosa sylvestris Woods. | |
| Essig Rose, Ger. | | 45. mollis Led. ☼ Caucasus - | 764 |
| <i>Varieties</i> ☼ ☼ - - - | 761 | The soft-leaved Rose. | |
| 2 pumila Lindl. ☼ | | R. Ledbourii Spreng. | |
| R. pumila L. | | 46. álba L. ☼ Europe fig. 496. | 764 |
| R. repens Munch. Haussv. | | The common white Rose. | |
| R. hispida Munch. | | R. usitatissima Gat. | |
| R. austrálica Crantz. | | <i>Variety</i> - - - | 764 |
| R. olympica Donn. | | § vii. Rubiginosa Lindl. ☼ ☼ ☼ - | 764 |
| 3 arvina Lindl. ☼ | | 47. lutea Dodon. ☼ Germany f. 497. | 765 |
| R. arvina Krok. | | The yellow Eglantine Rose. | |
| 4 inaperta Ser. ☼ | | R. Eglanteria L. | |
| The Vilmorin Rose. | | R. fa'tida Herm. | |
| 5 Agatha Red. ☼ | | R. chlorophylla Ehrh. | |
| The Agatha Rose. | | R. cæca Rossig. | |
| 6 inermis Ser. ☼ | | <i>Varieties</i> ☼ - - - | 765 |
| 7 parvifolia Ser. ☼ f. 491. | | 2 subrubra Red. ☼ | |
| The Burgundy Rose. | | 3 punicea Lindl. ☼ f. 498. | |
| R. parvifolia Ehrh. | | R. punicea Mill. Dict. | |
| R. burgundica Rossig. | | R. cinnamomea Roth. | |
| R. remensis Desf. | | R. lutea bicolor Jacq. | |
| | | R. Eglanteria punicea Red. | |
| | | R. Eglanteria bicolor Dec. | |
| 38. pulchélla Willd. ☼ | 762 | | |
| The neat Rose. | | | |
| § vi. Villosa. ☼ ☼ - - - | 762 | | |
| 39. turbinata Ail. ☼ Germany f. 495. | 762 | | |
| The turbinate-calyced, or Frankfort, Rose. | | | |
| R. campanulata Ehrh. | | | |
| R. francofurtana Munch. | | | |
| R. francofurtensis Rossig. | | | |
| <i>Varieties</i> ☼ - - - | 763 | | |
| 1 francofurtana Ser. ☼ | | | |
| The Frankfort Rose. | | | |
| R. turbinata Red. | | | |
| R. campanulata Ehrh. | | | |
| R. francofurtana Gmel. | | | |
| R. francofurtensis Desf. | | | |
| 2 orbessanea Ser. ☼ | | | |
| The Orbessan Rose. | | | |
| R. orbessanea Red. | | | |
| 10. villosa L. ☼ Europe - | 763 | | |
| The villous-leaved Rose. | | | |
| R. mollis Sm. | | | |
| R. tomentosa β Lindl. | | | |
| R. heterophylla Woods in L. Trans. | | | |
| R. pulchélla Woods. | | | |
| R. pomifera Herm. | | | |
| <i>Varieties</i> ☼ ☼ - - - | 763 | | |
| 2 resinosa Lindl. ☼ | | | |
| 3 pomifera Desr. ☼ | | | |
| R. hispida Poir. | | | |
| | | 49. suavcolens Pursh. ☼ N. America | 766 |
| | | The sweet-scented Rose, American Sweet Briar, or Eglantine. | |
| | | R. rubiginosa and R. Eglanteria of the Americans. | |

- | | Page | | Page |
|--------------------------------------------------------------|------|------------------------------------------------------------------------|------|
| 50. micrantha Sm. ☼ Britain - - | 766 | 16 ambigua Desv. ☼ - - | 768 |
| The small-flowered Rose, or Sweet Briar. | | R. malnandiarænsis Lejeune. | |
| R. rubiginosa β micrantha Lindl. | | 17 squarrosa - - - | 768 |
| 51. sepium Thuil. ☼ Europe - - | 766 | R. canina β Dict. Fl. Taur. | |
| The Hedge Rose, or Briar. | | 18 rubiflora Ser. ☼ - - | 768 |
| R. helvetica and | | 58. Försteri Sm. ☼ Europe - - | 768 |
| R. myrtifolia Hall. | | Forster's Dog Rose. | |
| R. canina β Dec. | | R. collina β et γ Woods. | |
| R. agræstis Savi. | | 59. dumetorum Thuill. ☼ Europe - - | 768 |
| R. biserrata, R. macrocarpa, and R. stipularis Mer. | | The Thicket Dog Rose. | |
| 52. ibérica Stev. ☼ Iberia - - | 766 | R. leucantha β acutifolia Bast. | |
| The Iberian Rose. | | R. sepium Borkh. | |
| 53. glutinosa Sm. ☼ Levant - - | 766 | R. solstitialis Bess. | |
| The clammy Rose, or Briar. | | R. corymbifera Gmel. | |
| R. rubiginosa cretica Red. | | 60. braetescens Woods. ☼ England - - | 769 |
| R. rubiginosa sphaerocarpa Desv. | | The bractescent Dog Rose. | |
| 54. Klükii Begg. ☼ Tauria - - | 766 | 61. sarmentacea Swartz. ☼ Europe - - | 769 |
| Kluk's Rose, or Sweet Briar. | | The sarmentaceous Dog Rose. | |
| R. rubiginosa Bieb. | | R. glaucophylla Winch. | |
| R. floribunda Stev. | | R. canina Roth. | |
| R. balsamea Bess. | | 62. cæsia Sm. ☼ Scotland - - | 769 |
| 55. Montezumæ Humb. ☼ Mexico - - | 766 | The grey Dog Rose. | |
| Montezuma's Rose, or Briar. | | R. canina pubescens Afr. | |
| § viii. Canina Lindl. ☼ * * * 767 | | R. canina ζ cæsia Lindl. | |
| 56. caucásica Pall. ☼ Caucas. f. 500. 767 | | 63. Borreri Woods. ☼ Britain - - | 769 |
| The Caucasian Dog Rose. | | Borrer's Dog Rose. | |
| R. leucantha Bieb. | | R. dianthorum Sm. | |
| 57. canina L. ☼ Europe - - | 767 | R. rubiginosa β Lindl. | |
| The common Dog Rose. | | R. rubiginosa inodora Hook. | |
| R. dumalis Bechst. | | ? R. sepium Borkh., not of Thuill. | |
| R. andegavensis Bat. | | R. affinis Rau. | |
| R. glauca Lois. | | R. uncinata β Besser. | |
| R. arvensis Schrank. | | 64. rubrifolia Willd. ☼ S. Europe f. 503. 769 | |
| R. glaucescens and R. nitens Mer. | | The red-leaved Dog Rose. | |
| R. teneriffensis Donn. | | R. multiflora Reyn. | |
| R. senticosa Achar. | | R. rubicunda Hall. | |
| Varieties. ☼ - - | | R. livida Andr. | |
| 2 sureulosa Woods. ☼ - - | 767 | R. cinnamomica γ rubrifolia Red. | |
| 3 nuda Woods. ☼ - - | 767 | Varieties. ☼ - - | 769 |
| 4 aciphylla Lindl. ☼ f. 501, 502. 767 | | 2 hispida Ser. ☼ | |
| R. aciphylla Rau. | | R. cinnamomica glauca Desf. | |
| 5 ægyptiaca Lindl. ☼ - - | 767 | 3 Redoutèa Ser. ☼ | |
| R. indica Forsk. | | 4 inermis Ser. ☼ | |
| 6 borboniana Desv. ☼ - - | 767 | 5 pinnatifida Ser. ☼ | |
| 7 nitens Desv. - - - | 768 | R. r. germinalibus-matis and R. montana germinalibus glabris Schlecht. | |
| R. nitens Desv. | | R. canina globosa Desv. | |
| 8 obtusifolia Desv. ☼ - - | 768 | 65. indica L. * China fig. 504. 770 | |
| R. obtusifolia Desv. | | The Indian, or common China, Rose. | |
| R. leucantha Lois. | | R. stueca L. Syst. Veg. | |
| 9 glaucescens Desv. ☼ - - | 768 | R. semperflorens cærnea Rossig. | |
| R. glaucescens Desv. | | R. indica chinensis simpliciflora Ser. | |
| R. canina glauca Desv. | | R. reclinata flore submultiplici Red. | |
| 10 Schottiana Ser. ☼ - - | 768 | The monthly Rose, the blush China Rose, the Tea-scented Rose. | |
| R. glauca Schott. | | Varieties. * * * - - | 770 |
| 11 piloscula Desv. ☼ - - | 768 | 2 Noisettiana Ser. * f. 505. | |
| R. humilis Bess. | | The Noisette Rose. | |
| R. nitidula Bess. | | Subvarieties. | |
| R. friedlanderiana Bess. | | purpurea Red. | |
| R. collina Rau. | | nivea Hort. | |
| 12 fastigiata Desv. ☼ - - | 768 | Smithii Hort. * | |
| R. fastigiata Bast. | | 3 odoratissima Lindl. * f. 506. | |
| R. stylosa β Desv. | | The sweetest, or Tea, scented China Rose. | |
| 13 hispida Desv. ☼ - - | 768 | R. odoratissima Swt. | |
| R. canina var. lancolata, grandidentata, and ovoidalis Desv. | | R. indica fragrans Red. | |
| R. andegavensis Bast. | | 4 longifolia Lindl. * f. 507. 771 | |
| R. semperflorens Bast., not of L. | | R. longifolia Willd. | |
| 14 microcarpa Desv. ☼ - - | 768 | R. semperflorens var. 7. N. Du Ham. | |
| 15 Meratiiana Ser. ☼ - - | 768 | R. satifolia Hort. | |
| ? R. biserrata Mer. | | 5 pumila Lindl. * * * | |
| | | 6 caryophyllæ Red. * * * | |

	Page	Page
7 pannòsa Red. ■		
8 eruénta Red. ■		
9 Fraseriána Hort. Brit. ■		
10 rùga Lindl. †		
11 oehroleùca Bot. Reg. ■		
The yellow China Rose.		
66. sempervlorens Curt. ■ China f. 508. 771		
The ever-flowering China Rose.		
R. diversifolia Vent.		
R. bengalensis Pers.		
R. indica Red.		
67. Lawrenceana Swt. ■ China f. 509. 772		
Miss Lawrence's China Rose.		
R. sempervlorens minima Sims.		
R. indica var. a acuminata Red.		
R. indica Lawrenceana Red.		
§ ix. Systyle Lindl. † † † † † † † † † 772		
68. systyla Bat. † † France - - - 772		
The connate-styled Rose.		
R. collina Sm.		
R. stylosa Desv.		
R. brevistyla Dec. Fl. Fr.		
R. bracteata Dec.		
R. systyla a ovata Lindl.		
69. arvënsis Huds. † † Britain f. 510. 772		
The Field Rose.		
R. sylvestris Hem.		
R. scandens Mench.		
R. herpëthodon Ehrh. Beitr.		
R. Halleri Krok.		
R. fusca Mench.		
R. serpens Ehrh. Arbor.		
R. sempervirens Rossig.		
R. repens Gmel.		
R. rampans Reyn.		
Varieties † † † - - - 772		
2 ayreshiëra Ser. † † †		
R. caprolata Neill.		
3 hýbrida Lindl. † † †		
70. (a.) sempervirens L. † † S. Europe		
fig. 511. 773		
The evergreen (Field) Rose.		
R. scandens Mill. Dict.		
R. balcanica Desf.		
R. atrovirens Viv.		
R. sempervirens globosa Red.		
R. sempervirens var. a scandens Dec.		
Varieties † † † † - - - 773		
2 Russelliána † † †		
3 Clàrei Bot. Reg. † † †		
The Rose Clare.		
4 Leschenaultiána Red. † † †		
71. multiflora Thunb. † Japan f. 512. 773		
The many-flowered Rose.		
R. flava Donn.		
R. florida Poir.		
R. diffusa Roxb.		
Varieties - - - 774		
2 Grevillei Hort. † † † f. 513.		
R. Roxburghii Hort.		
R. platyphylla Red.		
3 Russelliána †		
4 Boursaulti Hort. † † †		
72. Brunoniá Lindl. † † Nepal - 775		
Brown's Rose.		
R. Browni Spreng.		
73. moschàta Mill. † † Barbary		
fig. 514. 775		
The Musk Rose.		
R. oppositèmma Ehrh.		
R. glandulifera Roxb.		
Varieties † - - - 775		
2 flore plèno G. Don. †		
3 nívea Lindl. † fig. 515.		
R. nívea Dupont, not of Dec.		
R. m. ? var. rósea Ser.		
74. rubifolia R. Br. ■ North America		
fig. 516. 776		
The Bramble-leaved Rose.		
Variety † - - - 776		
3 fenestralis Lindl. †		
R. fenestrata.		
§ x. Banksiána Lindl. † - - 776		
75. sínica Ait. † China f. 517. 776		
The trifoliate-leaved China Rose.		
R. trifoliata Bosc.		
R. ternata Poir.		
R. cherokeënsis Donn.		
R. nívea Dec.		
76. Bánksie R. Br. † China f. 518. 777		
Lady Banks's Rose.		
R. Banksiána Abel.		
R. inermis Roxb. ?		
Variety † - - - 777		
2 lútea Lindl. † fig. 519.		
77. microcarpa Lindl. † China f. 520. 777		
The small-fruited Rose.		
R. cynusa Tratt.		
App. i. Hardy Species of the Genus Ròsa		
not yet introduced. - - - 778		
§ i. Feròces. - - - 778		
R. rugosa Thunb.		
§ ii. Bracteata. - - - 778		
R. Lyelli Lindl.		
§ iii. Cinnamòmea. - - - 778		
R. líxa Retz.		
R. kosingiana Desv.		
R. songárica Runge.		
R. aristata Lapey.		
R. macrophylla. fig. 521.		
§ iv. Pimpinellifolia. - - - 778		
R. Candolleana Don's Mill.		
R. rubella Lindl.		
R. pendula Roth.		
R. alpina Fall.		
R. poliphylla Windl.		
R. alpina var. rubella Ser.		
R. Candolleana pedunculata Red.		
R. Candolleana elegans Thor.		
R. flava Wicks.		
R. viminea Lindl.		
R. Weibotiana Wall.		
R. reclinata Red.		
§ v. Centifolia. - - - 778		
R. pygmaea Rich.		
R. adenophylla Willd.		
R. verecúnda Waitz.		
R. dimorpha var.		
§ vi. Villósa - - - 778		
R. hispida Forst.		
R. villosa var. pomifera Desv.		
R. tetralobata Bess.		

§ vii. *Rubiginosæ.* - - Page
- 778

- R. montana* Will.
- R. uncinella* Bess.
- R. caryophyllacea* Bess.
- R. inodora* Fries.
- R. cuspidata* Bieb.
- R. agrestis* Swartz.
- R. pseudo-rubiginosa* Lejeune.
- R. Wolfgangiana* Bess.
- R. dimorpha* Bess.
- R. Willdenowii* Spreng.
- R. microphylla* Willd.
- R. arcurata* Bieb.
- R. floribunda* Bess.

§ viii. *Caninae.* - - Page
- 778

- R. ciliato-petala* Bess.
- R. Gmelini* Bunge.
- R. canina* Sevrès.
- R. collina* Jacq.
- R. baltica* Roth.
- R. Jundzillii* Beggs.
- R. turbinella* Swartz.
- R. venosa* Swartz.
- R. corifolia* Fries.
- R. saxatile* Stec.
- R. sericea* Lindl. fig. 522
- R. atropurpurea* Roth.

§ ix. *Sýstyle.* - - Page
- 779

- R. moschata nepalensis.* fig. 523.
- ? *R. Brunonii* var. *nepalensis* Bot. Reg.
- R. evratina* Rosc.

§ x. *Banksianæ.* - - Page
- 779

- R. hystrix* Lindl. fig. 521.

App. ii. *Half-hardy Species of the Genus
Rosa, not yet introduced.* - 779

§ iv. *Pimpinellifoliae.* - - Page
- 779

- R. nankinensis* Lour.

§ viii. *Caninae.* - - Page
- 779

- R. pseudo-indica* Lindl.

§ x. *Banksianæ.* - - Page
- 779

- R. recurva* Rozb.
- R. triphylla* Roxb.
- ? *R. microcarpa.*
- ? *R. sinica* var.
- R. fragariaeflora* Ser.
- R. amygdalifolia* Ser.

App. iii. *Uncertain Species of Rosa.* 779

App. iv. *A practical Arrangement of Roses
in actual Cultivation in the Nursery of
Messrs. Rivers and Son, Sawbridgeworth,
Hertfordshire.* - - - 779

XII. LOWEA Lindl. $\text{æ} \perp$ - - Page
- 812

THE LOWEA.
Rosa sp. Pall.

1. *berberifolia* Lindl. $\text{æ} \perp$ Persia
fig. 594. 813

The Berberry-leaved Lowea.
Rosa simplicifolia Sal.
Rosa berberifolia Pall.

Varieties $\text{æ} \perp$ - - Page
- 813

- 1 *glabra* Ser. $\text{æ} \perp$
- 2 *velutina* Ser. $\text{æ} \perp$
- 3 *Redouteana* Ser. $\text{æ} \perp$
- R. berberifolia* Red.
- R. simplicifolia* Sal

Sect. V. *POMEÆ* Lindl. - 672. 813

XIII. CRATÆGUS Lindl. $\text{æ} \uparrow \uparrow \perp$

$\uparrow \perp \text{æ} \text{æ} \perp$ fig. 672. 813

THE THORN.

Cratægus and *Méspilus* sp. L. and others.

- Néflier, Atisier, Aubépine, Fr.*
- Doorn, Úzbecr, Mispel, Ger.*
- Doorn, Dutch.*
- Spino, Ital.*
- Espino, Span.*

§ i. *Coccinea.* æ - - Page
- 816

1. *coccinea* L. æ North America pl. 119.
fig. 564. in p. 851. - - Page
- 816

The scarlet-fruited Thorn.

- C. æstivalis* Booth.
- Méspilus æstivalis* Walt.
- Méspilus coccinea* Mill.
- Thornless American* Azarote.
- Néflier écartate, Fr.*
- Scharlachrothe Mispel, Ger.*

Varieties æ - - - Page
- 816

- 2 *corallina* æ f. 565. in p. 852.
- C. corallina* Lodd.
- C. pyriformis* of some collections.
- 3 *indentata* æ f. 566. in p. 852.
- C. indentata* Lodd.
- 4 *máxima* Lodd. æ
- C. c. spinosa* Godefroy.
- C. ? flabellata* Hort.

2. *glandulosa* W. æ North America
pl. 120. fig. 567. in p. 853. - 817

The glandular Thorn.

- ? *C. sanguinea* Pall.
- ? *Méspilus rotundifolia* Ehrh.
- Pýrus glandulosa* Moench.
- C. rotundifolia* Booth.

Varieties æ - - - Page
- 818

- 2 *succulenta* Fisch. æ
- Méspilus succulenta* Booth.
- 3 *subvillósa* æ f. 550., and f. 568.
in p. 853.
- C. subvillósa* Fisch.

§ ii. *Punctata.* æ - - Page
- 818

3. *punctata* Ait. æ North America pl. 124.
fig. 569, 570. in p. 854. - 818

The dotted-fruited Thorn.

- C. Crús-gállí* Du Roi.
- Méspilus cuneifolia* Ehrh.
- Méspilus punctata* Link.
- Méspilus cornifolia* Lam.

Varieties æ - - - Page
- 818

- 1 *rùbra* Pursh æ fig. 569. in 854.
- C. edulis* Ronalds.
- 2 *rùbra stricta* Hort. æ
- C. p. stricta* Ronalds.
- 3 *aúrea* Pursh æ fig. 570. in
p. 854.
- C. p. flava* Hort.
- C. dulcis* Ronalds.
- C. edulis* Lodd.
- C. pentágyna flava* Godefroy.

	Page		Page
4. <i>pyrifolia</i> <i>Ait.</i> ♀ North America pl. 122. fig. 571. in p. 854. - - 819	819	§ v. <i>Nigra</i> . ♀ - - - 822	822
The Pear-tree-leaved Thorn. <i>C. leucophlaeos</i> (white-barked) Mench. <i>C. radula</i> Lodd. <i>C. tomentosa</i> Du Roi. <i>Mespilus latifolia</i> Lam. <i>Mespilus Calpod. ulron</i> Ehrh <i>Mespilus pyrifolia</i> Link. <i>Mespilus cornifolia</i> Poir. <i>C. latifolia</i> Ronalds. <i>C. cornifolia</i> Booth.		9. <i>nigra</i> <i>Waldst. et Kil.</i> ♀ Hungary pl. 130. fig. 581. in p. 857. - 822	822
§ iii. <i>Macracantha</i> . ♀ - - 819	819	The black-fruited Thorn. <i>Mespilus nigra</i> Willd. <i>C. carpatica</i> Lodd.	
5. <i>macracantha</i> <i>Lodd.</i> ♀ North America pl. 123. fig. 572. in p. 855. - 819	819	10. <i>purpurea</i> <i>Bosc.</i> ♀ pl. 131. fig. 582. in p. 857. - - 822	822
The long-spined Thorn. <i>Variety</i> ♀ - - - 819 2 minor ♀ fig. 573. in p. 855. 819	819	The purple-branched Thorn. <i>C. sanguinea</i> Hort. <i>Variety</i> ♀ - - - 823 2 <i>altatica</i> ♀ f. 583. in p. 858.	823
§ iv. <i>Crus-galli</i> . ♀ ☼ - - 820	820	§ vi. <i>Douglasii</i> . ♀ - - - 823	823
6. <i>Crus-galli</i> <i>L.</i> ♀ North America pl. 124, 125. fig. 574. in p. 856. 820	820	11. <i>Douglasii</i> <i>Lindl.</i> ♀ North America pl. 132. fig. 584. in p. 858. - 823	823
The Cock's-spur Thorn. <i>C. lucida</i> Wang. <i>C. cuneifolia</i> Lodd. <i>Mespilus lucida</i> Ehrh. <i>Mespilus Crus-galli</i> Poir. <i>Mespilus hymenitis</i> Walt. <i>Mespilus cuneifolia</i> Mench. <i>Glänzende Mispel</i> , Ger. <i>Varieties</i> . ♀ ☼ 2 <i>splendens</i> <i>Dec.</i> ♀ fig. 575. in p. 856. - - 820 <i>C. arbutifolia</i> and <i>C. splendens</i> Lodd. 3 <i>pyracanthifolia</i> <i>Dec.</i> ♀ pl. 126. fig. 580. in p. 856. - 820 <i>C. pyracanthifolia</i> Lodd. <i>Mespilus lucida</i> Dunn. 4 <i>salicifolia</i> <i>Dec.</i> ♀ pl. 127. f. 551, 552, 553., and fig. 578. in p. 856. - - 820 <i>C. salicifolia</i> . 5 <i>linearis</i> <i>Dec.</i> ♀ fig. 577. in p. 856. - - 821 <i>Mespilus linearis</i> Desf. <i>C. linearis</i> Lodd. 6 <i>nana</i> <i>Dec.</i> ☼ - fig. 552. 821 <i>Mespilus nana</i> Dun.	820	Douglas's Thorn. <i>C. punctata</i> var. <i>brèvespina</i> Douglas. § vii. <i>Flava</i> . ♀ - - - 823	823
7. (C.) <i>ovalifolia</i> <i>Horn.</i> ♀ N. America pl. 128. fig. 579. in p. 856. - 821	821	12. <i>flava</i> <i>Ait.</i> ♀ North America pl. 133. fig. 585. in p. 859. - - 823	823
The oval-leaved Thorn. <i>C. elliptica</i> Lodd. <i>C. Crus-galli ovalifolia</i> Bot. Reg.		The yellow-fruited Thorn. <i>C. glandulosa</i> Michx., not of Walt. <i>Mespilus Michauxii</i> Pers. <i>C. caroliniana</i> Poir. <i>C. flavissima</i> Hort.	
8. (C.) <i>prunifolia</i> <i>Bosc.</i> ♀ N. America pl. 129. fig. 576. in p. 856. - 821	821	13. (f.) <i>lobata</i> <i>Bosc.</i> ♀ fig. 544., and fig. 586. in p. 859. - 824	824
The Plum-leaved Thorn. <i>Mespilus prunifolia</i> Poir. <i>C. caroliniana</i> Lodd. <i>Variety</i> ♀ - - - 821 2 <i>ingestria</i> ♀ <i>C. ingestria</i> Lodd.	821	The lobed-leaved Thorn. <i>Mespilus lobata</i> Poir. <i>C. lutea</i> Hort.	824
		14. <i>trilobata</i> <i>Lodd.</i> ♀ Hybrid fig. 587. in p. 860. - - 824	824
		The three-lobed-leaved Thorn. <i>C. spinosissima</i> Lee. § viii. <i>Apiifolia</i> . ♀ - - - 824	824
		15. <i>apiifolia</i> <i>Michx.</i> ♀ North America pl. 134. fig. 589. in p. 860. - 824	824
		The Parsley-leaved Thorn. <i>C. Oryacantha</i> Walt. <i>C. apiifolia major</i> Lodd. <i>Variety</i> ♀ - - - 825 2 minor ♀ f. 588. in p. 860. 825	825
		§ ix. <i>Microcarpa</i> . ♀ - - - 825	825
		16. <i>cordata</i> <i>Mill.</i> ♀ N. America pl. 135. fig. 590. in p. 861. - - 825	825
		The heart-shaped-leaved Thorn. <i>C. populifolia</i> Walt. <i>Mespilus acerifolia</i> Poir.	
		17. <i>spathulata</i> <i>Elliot.</i> ♀ North America pl. 136. fig. 591. in p. 861. - 825	825
		The spathula-shaped-leaved Thorn. <i>C. microcarpa</i> Lindl.	

- | | Page |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| § x. Azaròli. \checkmark - - - | 826 |
| 18. <i>Azaròlus</i> L. \checkmark France pl. 137.
fig. 592. in p. 862. - - - | 826 |
| The Azarole Thorn.
<i>Pyrus Azaròlus Scop.</i>
<i>Méspilus Azaròlus All.</i>
<i>E'pine d'Espagne, Néflier de Naples, Pommettes à deux Clos, Fr.</i> | |
| <i>Varieties</i> \checkmark - - - | 826 |
| 1 With the leaves hairy beneath. \checkmark
<i>Méspilus Arònia.</i> | |
| 2 With large deep red fruit. \checkmark | |
| 3 With yellowish white fruit. \checkmark | |
| 4 With long fruit of a whitish yellow. \checkmark | |
| 5 With double flowers. \checkmark | |
| 6 The white Azarole of Italy. \checkmark | |
| 19. (<i>A.</i>) <i>maroccàna</i> Pers. \checkmark Morocco
pl. 138. fig. 594. in p. 862. - 827 | 827 |
| The Morocco Thorn.
? <i>C. maúra</i> L. | |
| 20. <i>Arònia</i> Bosc. \checkmark Levant pl. 139.
fig. 593. in p. 862. 827 | 827 |
| The Aronia Thorn.
<i>Méspilus Arònia</i> Willd. Enum.
<i>C. Azaròlus</i> β Willd. sp. | |
| 21. <i>orientàlis</i> Bosc. \checkmark South Europe
pl. 140. fig. 595. in p. 863. 827 | 827 |
| The Eastern Thorn.
<i>Méspilus orientàlis</i> Poir.
<i>C. odoratissima</i> Bot. Rep. | |
| <i>Variety</i> \checkmark - - - | 828 |
| 2 <i>sanguinea</i> \checkmark f. 596. in p. 863. 828 | 828 |
| <i>C. tanacetifolia</i> 2 <i>taurica</i> Dec.
<i>C. sanguinea</i> Schrader.
<i>C. orientàlis</i> Lindl. | |
| 22. <i>tanacetifolia</i> Pers. \checkmark Greece pl. 141.
fig. 597. in p. 863. - - - | 828 |
| The Tansy-leaved Thorn.
<i>Méspilus tanacetifolia</i> Poir.
<i>Méspilus pinnata</i> Dum.
? <i>Méspilus Celsiana</i> Dum. | |
| <i>Varieties</i> \checkmark - - - | 828 |
| 2 <i>glàbra</i> Lodd. \checkmark pl. 142. f. 598.
in p. 863. | |
| 3 <i>Leeàna</i> \checkmark pl. 143. f. 599. in
p. 864. | |
| <i>Lee's Seedling, Hort.</i>
<i>C. incisa</i> Lee. | |
| § xi. <i>Heterophýlla</i> . \checkmark - - - | 829 |
| 23. <i>heterophýlla</i> Flugge \checkmark pl. 144.
fig. 600. in p. 864. - - - | 829 |
| The various-leaved Thorn.
<i>C. neapolitana</i> Hort.
<i>Méspilus constantinopolitana</i> Godefroy. | |
| § xii. <i>Oxyacántha</i> . \checkmark - - - | 829 |
| 24. <i>Oxyacántha</i> L. \checkmark Europe pl. 145,
146. fig. 602. in p. 865. - 829 | 829 |
| The sharp-thorned <i>Cratægus</i> , or common Hawthorn.
<i>The Pyracantha</i> of the Greeks. | |
| <i>Méspilus Oxyacántha</i> Gaertn.
<i>E'pine blanche, Noble E'pine, Bois de Mai, Scuellier Aubépine, Néflier Aubépine, Fr. Hagedorn, Gemeiner Weissdorn, Ger. Hagctoun, Dan. Hagctorn, Swed. Bianco-spino, Ital. Espino blanco, Span. White Thorn, May Bush, Quick, Quicksct, May.</i> | |
| <i>Varieties.</i> \checkmark | |
| 2 obtusàta Dec. \checkmark pl. 147. f. 601.
in p. 864. - - - | 830 |
| <i>Méspilus Oxyacántha integrifolia</i> Wallr.
<i>C. Oxyacanthoides</i> Thuill.
<i>C. Oxyacántha</i> Fl. Dan.
<i>The French Hawthorn.</i> | |
| 3 <i>sibírica</i> \checkmark fig. 555. - 830 | 830 |
| <i>C. sibírica</i> Lodd.
<i>C. mongyana</i> L. | |
| 4 <i>transylvànica</i> Hort. \checkmark - 830 | 830 |
| 5 <i>quercifolia</i> Booth. \checkmark f. 603. in
p. 866. - - - | 830 |
| 6 <i>laciniàta</i> \checkmark pl. 148. f. 663. in
p. 865. - - - | 830 |
| <i>C. laciniàta</i> Lodd. | |
| 7 <i>pteridifolia</i> \checkmark fig. 604. in
p. 865. - - - | 831 |
| <i>C. pterifolia</i> Lodd.
<i>C. pectinata</i> Hort. | |
| 8 <i>eriocàrpa</i> Lindl. \checkmark pl. 149.
fig. 607. in p. 865. - 831 | 831 |
| <i>C. eriocàrpa</i> Lodd. | |
| 9 <i>purpùrea</i> Penny \checkmark f. 611. in
p. 866. - - - | 831 |
| 10 <i>Oliveriàna</i> \checkmark pl. 150. f. 606.
in p. 865. - - - | 831 |
| <i>C. Oliveriàna</i> Bosc.
<i>C. Oliveria</i> Lodd.
<i>C. orientalis</i> Lodd. | |
| 11 <i>melanocàrpa</i> \checkmark pl. 151. f. 605.
in p. 865. - - - | 831 |
| <i>C. fissa</i> Lee.
<i>C. Oxyacántha platyphýlla</i> Lodd.
<i>C. platyphýlla</i> Lindl. | |
| 12 <i>aùrea</i> Hort. \checkmark fig. 610. in
p. 866. - - - | 831 |
| <i>C. flava</i> Hort. | |
| 13 <i>aurantiaca</i> Booth. \checkmark - 831 | 831 |
| 14 <i>leucocàrpa</i> \checkmark - - - | 831 |
| 15 <i>múltiplex</i> Hort. \checkmark f. 609. in
p. 866. - - - | 832 |
| <i>C. O. flore pleno</i> Hort. | |
| 16 <i>ròsea</i> Hort. \checkmark fig. 612. in
p. 866. - - - | 832 |
| <i>E'pniec Marron, Fr.</i> | |
| 17 <i>punicèa</i> Lodd. \checkmark - - - | 832 |
| <i>C. O. rùca superba</i> Hort. | |
| 18 <i>punicèa flore pleno</i> Hort. \checkmark 832 | 832 |
| 19 <i>fòliis aùreis</i> Lodd. \checkmark - 832 | 832 |
| 20 <i>fòliis argéteis</i> Hort. \checkmark - 832 | 832 |
| 21 <i>strieta</i> Lodd. \checkmark pl. 152. 832 | 832 |
| <i>C. O. rigida</i> Ronalds. | |
| 22 <i>Celsiana</i> Hort. \checkmark - - - | 832 |
| 23 <i>péndula</i> Lodd. \checkmark - - - | 832 |
| 24 <i>reginæ</i> Hort. \checkmark
pl. 153. fig. 556. 832 | 832 |
| Queen Mary's Thorn. | |

25	<i>praecox Hort.</i> † - - -	Page 833
	The early-flowering, or <i>Glastonbury</i> , Thorn.	
26	<i>monogyne</i> † - - -	834
	<i>C. monogyne</i> Jacq.	
27	<i>apétala</i> <i>Lodd.</i> † - - -	834
28	<i>lúcida</i> † - - -	834
29	<i>capitata</i> <i>Smith of Agr.</i> † - -	834
30	<i>flexuosa</i> <i>Smith of Agr.</i> † - -	835
§ xiii. <i>Parvifolia.</i> † - - - 841		

25. *parvifolia* *Ait.* ✨ North America
f. 557., and f. 614. in p. 867. - 841
The small-leaved Thorn.
Mespilus axillaris Pers.
Mespilus tomentosa Poir.
Mespilus xanthocarpos L.
Mespilus parvifolia Wats.
C. tomentosa L.
C. uniflora Du Roi.
C. viridis, *C. axillaris*, *C. bctulifolia*,
C. florida, and *C. linearis* *Lodd.*
Gooseberry-leaved Thorn.

Varieties ✨ - - - 842
2 *florida* ✨ fig. 558., and fig. 613.
in p. 867.
C. florida *Lodd.*
3 *grossularifolia* ✨ fig. 559., and
fig. 616. in p. 867.

26. *virginica* *Lodd.* ✨ Virginia fig. 560.,
and fig. 615. in p. 867. - - 842
The Virginian Thorn.
C. virginiana *Hort.*

§ xiv. *Mexicana.* † — - - 843

27. *mexicana* *Moc. et Sesse.* † — Mexico
pl. 154. fig. 617. in p. 867. - 843
The Mexican Thorn.
C. stipulacea *Lodd.*

§ xv. *Pyracantha.* ✨ — — - 844

28. *Pyracantha* *Pers.* ✨ South Europe
fig. 561. 844
The fiery Thorn, or *Pyracantha.*
Mespilus Pyracantha L.
Evergreen Thorn.
Buisson ardent. Fr.
Innegrüne *Mispel.* Ger.

Variety ✨ — - - 844
2 *crenulata* ✨ — — —
C. crenulata *Roxb. MSS.*
Mespilus crenulata *D. Don.*

§ xvi. *Glaúca.* † - - - 844

29. *glaúca* *Wall.* † Nepal f. 562, 563. 844
The glaucous-leaved evergreen Thorn.

App. i. *Synopsis of the Species of Crataegus growing, in 1836, in the Horticultural Society's Garden.* - - - 845

App. ii. *Additional Species of Crataegus.* 848

§ i. *Leaves toothed, or nearly entire, never regularly lobed.* † - - - 848

- C. subspinosa* *Dec.*
Mespilus subspinosa *Vent.*
- C. prunifolia* *Bos.* †
- C. latifolia* *Pers.* †
- C. flexuosa* *Poir.*
- C. alpina* *Mill. Dict.* †
- C. lutea* *Poir.*
- C. pauciflora* *Pers.*
- C. unilateralis* *Pers.*
- C. lucida* *Mill. Dict.* †

§ ii. *Leaves variously lobed, or cut.* † - - - 848

- C. turbinata* *Pursh.*
- C. pentagyna* *Waldst. et Kit.* †
- C. kirtostyla* *Fing.*
- C. laciniata* *Dec.* †
- C. laciniata* *Dec.*
Mespilus laciniata *Poir.*
- C. Poir. Mispilus* *Dec.* †
- Mespilus linearis* *Poir.*
- C. pectinata* *Bosc.*
- C. trifoliata* *Bosc.*
- C. quinquelobata* *Bosc.*
- C. odorata* *Bosc.*
- C. obovata* *Bosc.*
- C. flavescens* *Bosc.*
- C. flabellata* *Bosc.* †
- C. lucida latifolia* *Boll. Cat.* †
- C. lucida media* *Boll. Cat.* †

App. iii. *Alphabetical List of Sorts of Crataegus in the Arboretum of Messrs. Loddiges, as given in their Catalogue, 16th ed. 1836; with some Additions, taken from the Names placed against Plants in their Nursery, but not in the Catalogue; referred to the Species and Varieties of Crataegus as given in this Work.* - - - 848

App. iv. *Alphabetical List of the Species and Varieties of Crataegus described in the Arboretum Britannicum, with the Names which are appended to the Specimen Plants of these Sorts in the Arboretum of Messrs. Loddiges* - - - 849

XIV. PHOTINIA *Lindl.* † — † — 673. 868

THE PHOTINIA.
Crataegus sp. L.

1. *serrulata* *Lindl.* † — Japan
pl. 155. 868
The serrulated-leaved Photinia.
Crataegus glabra *Thunb.*
Strausveisia *Lindl.*
2. *arbutifolia* *Lindl.* † — California
fig. 619. 868
The Arbutus-leaved Photinia.
Crataegus arbutifolia *Ait.* - - - 868
3. *integrifolia* *Lindl.* † — Nepal - 869
The entire leaved Photinia.
Pyrus integrifolia *Wall.*
4. *dubia* *Lindl.* † — Nepal - - 869
The doubtful Photinia.
Mespilus bengalensis *Roxb.*
Mespilus tinctoria *D. Don.*
Crataegus Shicola *Ham. MSS.*

App. i. *Species of Photinia not yet introduced.* - - - 869

- P. *bengalensis* Wall.
 P. *Sieboldii* Don's Mill.
 Mespilus Sieboldii Blum.
 P. *laevis* Dec.
 Crataegus laevis Thunb.
 P. *villosa* Dec.
 Crataegus villosa Thunb.

XV. COTONEASTER *Med.* † † † 673. 869

THE COTONEASTER.
Mespilus sp. L.

- § i. *Leaves deciduous. Shrubs.* 673 - 870
1. *vulgàris* Lindl. 620. 870
 The common Cotoneaster.
Mespilus Cotoneaster L.
Néflier cotonneux, Fr.
Quitten Mispel, Ger.
 Varieties 673 - - - 870
 1 *erythrocarpa* Led. 673
 2 *melanocarpa* Led. 673
 3 *depressa* Fries. 673
2. (v.) *tomentosa* Lindl. 870
 The tomentose, or woolly, Cotoneaster.
Mespilus tomentosa Willd., not of Lam
Mespilus cricocarpa Dec.
3. (v.) *laxiflora* Jacq. 870
 The loose-flowered Cotoneaster.
 fig. 621, 622.
- § ii. *Subevergreen or deciduous. Tall Shrubs, or low Trees.* † † - - 871
4. *frigida* Wall. 156. 871
 The frigid Cotoneaster.
Pyrus Nussia Ham.
5. (f.) *affinis* Lindl. † † Nepal pl. 157. 871
 The related (to *C. frigida*) Cotoneaster.
Mespilus integerrima Ham. MSS.
6. *acuminata* Lindl. † † Nepal pl. 158. 872
 The acuminated-leaved Cotoneaster.
Mespilus acuminata Lodd.
7. *nummularia* Lindl. † † Nepal pl. 159. 872
 The money-like-leaved Cotoneaster.
- § iii. *Leaves evergreen, leathery. Low Shrubs, with prostrate Branches; Trailers, but not properly Creepers.* 673 - - - 872
8. *rotundifolia* Wall. 623, 624. 872
 The round-leaved Cotoneaster.
C. microphylla β *Ura-ursi* Lindl.
 The Bear-berry-leaved Nepal Cotoneaster.
9. (r.) *microphylla* Wall. 625. 873
 The small-leaved Cotoneaster.
10. (r.) *buxifolia* Wall. 625. 873
 The Box-leaved Cotoneaster.

App. i. *Species of Cotoneaster not yet introduced.* - - - 873

- C. bacillaris* Wall.
C. obtusa Wall.

XVI. AMELANCHIER *Med.* † 673. 874

THE AMELANCHIER.
Mespilus L.
Pyrus W.
Aronia Pers.
Crataegus Lam.
Sorbus Crantz.

1. *vulgàris* Mœnch. † Europe f. 626. 874
 The common Amelanchier.
Mespilus Amelanchier L.
Pyrus Amelanchier Willd.
Aronia rotundifolia Pers.
Crataegus rotundifolia Lam.
Sorbus Amelanchier Crantz.
Alisier Amelanchier, Amelanchier des Bois, Néflier à Feuilles rondes, Fr.
Felsenbirne, Ger.
2. (v.) *Botryapium* Dec. † N. America pl. 160, 161. fig. 627, 628. 874
 The Grape-Pear, or Snowy-blossomed, Amelanchier.
Mespilus canadensis L.
Mespilus arborea Michx.
Crataegus racemosa Lam.
Pyrus Botryapium Pers.
 The Canadian Medlar, Snowy Mespilus, June Berry, Wild Pear Tree.
Alisier de Choisy, Alisier à Grappes, Fr.
Traubenbirne, Ger.
3. (v.) *sanguinea* Dec. † North America fig. 630, 631. 875
 The blood-coloured Amelanchier.
Pyrus sanguinea Pursh.
Aronia sanguinea Nutt.
Mespilus canadensis γ rotundifolia Michx.
4. (v.) *ovalis* Dec. † North America fig. 632. 876
 The oval-leaved Amelanchier.
Crataegus spicata Lam.
Mespilus Amelanchier Wall.
A. parviflora Dougl. MSS.
Mespilus canadensis var. α ovalis Michx.
Pyrus ovalis Willd.
Aronia ovalis Pers.
Amelanchier du Canada, Alisier à E'pi, Fr.
Rundblättrige Birne, Ger.
 Variety † - - - 876
 2 *subcordata* Dec. †
 Aronia subcordata Raf.
 Mitus microcarpa Raf.
 3 *semi-integrifolia* Hook. †
5. (v.) *florida* Lindl. † North America fig. 633, 634. 876
 The flowery Amelanchier.
 Variety - - - 877
 2 *parvifolia* †
 A. parvifolia Hort. Soc. Gard.

XVII. ME'SPILUS Lindl. † 673. 877

THE MEDLAR.
Mespilus sp. of Lin. and others.
Mespilophora sp. Neck.

- | | Page | | Page |
|---------------------------------------------------------------------------------------------------|-------|--------------------------------------------------------------------------------------------------|--------------|
| 1. <i>germánica L.</i> Y Europe pl. 162. | 877 | 4. (c.) <i>sináica Thoun.</i> Y Mount Sinai | 889 |
| The German, or common, Medlar. | | | pl. 167. |
| <i>Varieties</i> - - - | - 878 | The Mount Sinai Pear Tree. | |
| 1 <i>sylvéstris Mill. Dict.</i> Y | | <i>P. Sinai</i> Desf. | |
| 2 <i>stricta Dec.</i> Y | | <i>P. persica</i> Pers. | |
| 3 <i>diffusa Dec.</i> Y | | The Mount Sinai Medlar. | |
| <i>Cultivated Varieties.</i> | | 5. (c.) <i>salicifolia L.</i> Y Siberia - 889 | |
| 1. Blake's large-fruited Medlar. | | The Willow-leaved Pear Tree. | |
| 2. Dutch Medlar. | | 6. (c.) <i>clavaginifolia Pall.</i> Y A. Minor 889 | |
| 3. Nottingham, or common, Medlar. | | The Oleaster-leaved Pear Tree. | |
| 4. The stonless Medlar. | | <i>P. orientalis</i> Horn. | |
| 2. <i>Smithi Dec.</i> Y pl. 163. | 878 | 7. (c.) <i>amygdaliformis Vil.</i> Y - 889 | |
| Smith's Medlar. | | The Almond-shaped Pear Tree. | |
| <i>M. grandiflora</i> Sm. | | <i>P. sylvéstris</i> Magnol Bot. | |
| <i>M. lobata</i> Poir. | | <i>P. salicifolia</i> Lois. | |
| XVIII. <i>PYRUS Lindl.</i> Y 673. 879 | | 8. <i>sinéensis Lindl.</i> Y China pl. 168. 889 | |
| THE PEAR TREE. | | The Chinese Pear Tree. | |
| <i>Pyrus, Malus, and Sorbus, Tourn.</i> | | <i>Pyrus communis</i> Lois. | |
| <i>Pyrus and Sorbus L.</i> | | <i>P. sinica</i> Royle. | |
| <i>Pyróphorum and Appyróphorum Neck.</i> | | <i>Ri vulgo</i> Nas, Japanese. | |
| | | The sandy Pear, Snow Pear, Sand Pear. | |
| | | <i>Sha lec</i> , Chinese. | |
| § i. <i>Pyróphorum Dec.</i> Y - 880 | | 9. <i>bollwylleriána Dec.</i> Y France | pl. 169. 890 |
| 1. <i>communis L.</i> Y Eur. pl. 164, 165. 880 | | The Bollwyller Pear Tree. | |
| The common Pear Tree. | | <i>P. Poltéria</i> L. | |
| <i>P. A'chras</i> Gært. n. | | <i>P. auricularis</i> Knoop. | |
| <i>P. sylvéstris</i> Dod. | | 10. <i>crenata Don.</i> Y Nepal f. 628, 639. 890 | |
| <i>Pyráster</i> Ray. | | The notched-leaved Pear Tree. | |
| <i>Poirier</i> , Fr. | | 11. <i>variolosa Wall.</i> Y Nepal pl. 170. 891 | |
| <i>Gemeine Birne, Birnebaum</i> , Ger. | | The variable-leaved Pear Tree. | |
| <i>Pera</i> , Ital. | | <i>P. Páshia</i> Ham. | |
| <i>Pera</i> , Span. | | App. i. <i>Species of Pyrus belonging to the Section Pyróphorum, and not yet introduced.</i> 891 | |
| <i>Gruschka</i> , Russian. | | <i>P. euneifolia</i> Gus. | |
| <i>Varieties</i> Y - - - - - 880 | | <i>P. parviflora</i> Desf. | |
| 1 <i>A'chras</i> Wallr. Y | | <i>P. sylvéstris cética</i> C. Bauh. | |
| 2 <i>Pyráster</i> Wallr. Y | | <i>P. Michauxii</i> Bosc. | |
| 3 <i>foliis variegatis</i> Y | | <i>P. indica</i> Colebr. | |
| 4 <i>fructu variegato</i> Y | | § ii. <i>Malus.</i> Y - - - 891 | |
| 5 <i>sanguinolenta</i> Y | | 12. <i>Malus L.</i> Y Eur. pl. 171, 172. 891 | |
| The sanguinole Pear. | | The common, or wild, Apple Tree. | |
| 6 <i>flöre pléno</i> Y | | <i>P. Malus mitis</i> Wallr. | |
| <i>Poire de l'Arménie</i> Bon Jard. | | <i>Malus communis</i> Dec. | |
| 7 <i>jáspida</i> Y | | <i>Pommier commun</i> , Fr. | |
| <i>Bon Chrétien á Bois jaspé</i> Bon Jard. | | <i>Gemeine Apfelbaum</i> , Ger. | |
| 8 <i>sativa Dec.</i> Y | | 13. (<i>M.</i>) <i>acerba Dec.</i> Y Europe - 892 | |
| <i>Varieties most deserving of Cultivation, selected from the Hort. Soc. Cat. of Fruits</i> - 881 | | The sour-fruited Apple, or common Crab Tree. | |
| <i>Beurré Diel.</i> | | <i>Pyrus Malus austera</i> Wallr. | |
| <i>Beurré de Rans.</i> | | <i>Malus acerba</i> Merat. | |
| <i>Bezi de la Motte.</i> | | <i>Malus communis sylvéstris</i> Desf. | |
| <i>Glout Moreau</i> , pl. 166. | | <i>Malus sylvéstris</i> Fl. Dan. | |
| Napoleon. | | <i>P. Malus</i> Sm. | |
| Swan's Egg. | | <i>Pommier sauvagine</i> , Fr. | |
| <i>Scotch Pears recommended by Mr. Gorrie, as Trees adapted for Landscape Secnery</i> - 881 | | <i>Holzapfelbaum</i> , Ger. | |
| The Benvie. | | 14. (<i>M.</i>) <i>prunifolia H.</i> Y Siberia - 892 | |
| The Golden Knap. | | The Plum-tree-leaved Apple Tree, or Siberian Crab. | |
| The Elcho. | | <i>P. Malus</i> β Ait. | |
| The busked Lady. | | <i>Malus hýbrida</i> Desf. | |
| The Pow Meg. | | 15. (<i>M.</i>) <i>baccata L.</i> Y Siberia - 892 | |
| 2. (c.) <i>salvifolia Dec.</i> Y France 888 | | The berry-like-fruited Apple Tree, or Siberian Crab. | |
| The Sage-leaved, or Annetian, Pear Tree. | | <i>Malus baccata</i> Desf. | |
| <i>Poirier Sauger</i> D'Ourch. | | | |
| 3. (c.) <i>nivalis L. fl.</i> Y Austria - 888 | | | |
| The snowy-leaved Pear Tree. | | | |

	Page		Page
16. (<i>M.</i>) dioica <i>W.</i> \checkmark	892	6 eretica <i>Lindl.</i> \checkmark	
The dioecious-seed Apple Tree.		<i>P. A. rotundifolia</i> Hort.	
<i>P. apétala</i> Münch.		<i>P. græca</i> Hort.	
<i>Malus dioica</i> Audub.		7 bullata <i>Lindl.</i> \checkmark	
		<i>P. A. acuminata</i> Hort.	
17. (<i>M.</i>) astracànica <i>Dec.</i> \checkmark Persia 893		22. (<i>A.</i>) intermèdia <i>Ehrh.</i> \checkmark Europe 912	
The Astrachan Apple Tree.		The intermediate White Beam Tree.	
<i>Malus astracànica</i> Dum.		<i>Cratægus A'ria</i> β L.	
<i>Varieties of P. Malus cultivated for their</i>		<i>Cratægus scândica</i> Wahl	
<i>Fruit.</i> - - - 893		<i>Cratægus suécica</i> Ait.	
The Red Astrachan.		<i>Alisier de Fontainebleau</i> , Fr.	
The White Astrachan.		<i>Schwedischer Mehlsbaum</i> , Ger.	
<i>The transparent Crab of Moscow.</i>		<i>Varieties</i> \checkmark - - - 912	
The Black Crab.		1 latifolia <i>Dec.</i> \checkmark	
The Court pendu plat.		<i>Cratægus latifolia</i> Poir.	
The Lincolnshire Holland Pippin.		<i>Sórbus latifolia</i> Pers.	
The Tulip Apple.		<i>Cratægus dentata</i> Thuil.	
The Violet Apple.		2 angustifolia <i>Dec.</i> \checkmark fig. 640.	
The Cherry Crab.		<i>P. edulis</i> Willd.	
<i>A subvar. of P. (M.) baccata.</i>			
The Supreme Crab.		23. (<i>A.</i>) vestita <i>Wall.</i> \checkmark Nepal pl.178. 912	
Bigg's Everlasting Crab.		The clothed White Beam Tree.	
		<i>Pýrus nepalensis</i> Hort.	
18. coronària <i>L.</i> \checkmark North America		<i>Sórbus vestita</i> Lodd.	
pl. 174. 908			
The garland-flowering Apple Tree.		App. i. <i>Additional Species of Pýrus belong-</i>	
<i>Malus coronària</i> Mill.		<i>ing to the Section A'ria.</i> - 913	
<i>Crab Apple, Sweet-scented Crab, Amer.</i>			
19. (c.) angustifolia <i>Ait.</i> \checkmark Carolina		<i>P. kamaonensis</i> Wall.	
pl. 175. 909		<i>P. lanata</i> D. Don.	
The narrow-leaved Apple Tree.		\S iv. <i>Torminària</i> <i>Dec.</i> \checkmark - 913	
<i>P. coronària</i> Wang.			
<i>Malus sempervirens</i> Desf.		24. torminàlis <i>Ehrh.</i> \checkmark Europe	
<i>P. pàmila</i> Hort.		pl. 179. 913	
20. spectàbilis <i>Ait.</i> \checkmark China		The gripping-fruited Service Tree.	
pl. 176. 909		<i>Cratægus torminàlis</i> L.	
The showy-flowering wild Apple Tree, or Chi-		<i>Sórbus torminàlis</i> Crantz.	
nese Crab Tree.		<i>The Maple-leaved Service Tree.</i>	
<i>Malus spectàbilis</i> Desf.		<i>Alisier des Bois</i> , Fr.	
<i>Malus sinënsis</i> Dum.		<i>Elzbeur Baum</i> , Ger.	
App. i. <i>Additional Species of Pýrus belonging</i>		App. i. <i>Other Species of Pýrus belonging to</i>	
<i>to the Section Malus.</i> - 909		<i>the Section Torminària.</i> - 915	
<i>P. quinqueflora</i> Hamilt.		<i>P. rivularis</i> Dougl. fig. 642, 643.	
<i>P. Sieversii</i> Led.		\S v. <i>Eriólobus</i> <i>Dec.</i> \checkmark - 915	
<i>P. nov. sp. Sievers.</i>			
\S iii. <i>A'ria</i> <i>Dec.</i> \checkmark - - 910		25. trilobata <i>Dec.</i> \checkmark Mount Lebanon 915	
21. A'ria <i>Ehrh.</i> \checkmark Europe - - 910		The three-lobed-leaved Pear Tree.	
The White Beam Tree.		<i>Cratægus trilobata</i> Labill.	
<i>Cratægus A'ria</i> var. α L.		\S vi. <i>Sórbus</i> <i>Dec.</i> \checkmark - 915	
<i>Mespilus A'ria</i> Scop.			
<i>Sórbus A'ria</i> Crantz.		26. auriculata <i>Dec.</i> \checkmark Egypt - 915	
<i>A'ria Theophrasti</i> L'Obel.		The auricled Service Tree.	
<i>White Wild Pear, White Leaf Tree, Red</i>		<i>Sórbus auriculata</i> Pers.	
<i>Chess Apple, Sea Ouler, Cumberland</i>			
<i>Hawthorn, Gerard.</i>		27. pinnatifida <i>Ehrh.</i> \checkmark Britain.	
<i>Alisier Allouchier, Alisier blanc, Fr.</i>		pl. 180. 915	
<i>Mehlbeerbaum, Mehlsbaum, Ger.</i>		The pinnatifid-leaved Service Tree.	
<i>Aria, Sorba pilosa, Ital.</i>		<i>Sórbus híbrida</i> L.	
<i>Mostaco, Span.</i>		<i>Pýrus híbrida</i> Sm., not of Willd.	
<i>Axelbeer, Danish.</i>		<i>The Bastard Service Tree.</i>	
<i>Oxilbeer, Swedish.</i>		<i>Varieties</i> \checkmark - - - 916	
<i>Varieties</i> \checkmark - - - 910		2 lanuginosa \checkmark	
1 obtusifolia <i>Dec.</i> \checkmark		3 pendula \checkmark	
<i>P. A. ovalis</i> Hort.		<i>Sórbus híbrida pendula</i> Lodd.	
2 acutifolia <i>Dec.</i> \checkmark		4 arbúscula <i>Dec.</i> \checkmark	
<i>Cratægus longifolia</i> N. Du Ham.			
? <i>P. alpina</i> Willd.		28. aucupària <i>Gærtu.</i> \checkmark Europe	
3 undulata <i>Lindl.</i> \checkmark pl. 177.		pl. 181. 916	
4 angustifolia <i>Lindl.</i> \checkmark		The Fowler's Service Tree, or Mountain Ash.	
5 rugosa <i>Lindl.</i> \checkmark		<i>Sórbus aucupària</i> L.	
		<i>Mespilus aucupària</i> All.	
		<i>Quicken Tree, Quick Beam, Wild Ash,</i>	
		<i>Wild Service, Wicken Tree, Rowan</i>	
		g 2	

- | | Page | | Page |
|-------------------------------------------------------------------------------------------------------------------------|----------------------------------|---------------------------------------------------|------------------|
| <i>Tree, Rowne Tree, Roan Tree, Roddian, Routry, Mountain Service, Wichen, Wild Sorb, Wichen, Whitten, Wigger Tree.</i> | | 3 <i>serótina</i> Lindl. ☼ | |
| <i>Sorbier des Oiseleurs, Sorbier des Oiseaux, Fr.</i> | | 4 <i>pùmila</i> ☼ f. 647, 648. | |
| <i>Fogel Beerbaum, Ger.</i> | | - <i>Méspilus pùmila</i> Lodd. | |
| <i>Sorbo Salvatico, Ital.</i> | | 35. (a.) <i>melanocarpa</i> W. N. America | fig. 649. 926 |
| <i>Varieties</i> ☼ - - - 916 | | The black-fruited Aronia. | |
| 2 <i>fructu luteo</i> ☼ | | <i>P. arbutifolia</i> β Willd. | |
| 3 <i>foliis variegatis</i> ☼ | | <i>Ardunia arbutifolia</i> Pers. | |
| 4 <i>fastigiata</i> ☼ | | <i>Variety</i> ☼ - - - 927 | |
| 29. <i>americana</i> Dec. ☼ North America | pl. 182. 920 | 36. (a.) <i>floribunda</i> Lindl. ☼ North America | - - - 927 |
| The American Service. | | The abundant-flowered Aronia. | |
| <i>Sorbus americana</i> Pursh. | | 37. (a.) <i>depressa</i> Lindl. ☼ North America | 927 |
| <i>Sorbus americana</i> var. β Michx. | | The depressed Aronia. | |
| 30. <i>microcarpa</i> Dec. ☼ North America | 921 | 38. <i>púbens</i> Lindl. ☼ N. America - 927 | |
| The small-fruited Service. | | The downy-branched Aronia. | |
| <i>Sorbus aucuparia</i> α Michx. | | 39. <i>grandifolia</i> Lindl. ☼ North America | fig. 650. 928 |
| <i>Sorbus micrantha</i> Dum. | | The large-leaved Aronia. | |
| <i>Sorbus microcarpa</i> Pursh. | | § viii. <i>Chamaméspilus</i> Dec. ☼ 928 | |
| 31. <i>Sorbus Gærtu.</i> ☼ Barbary pl. 183. 921 | | 40. <i>Chamaméspilus</i> Lindl. ☼ Europe | fig. 651. 928 |
| The True Service. | | The dwarf Medlar. | |
| <i>Sorbus domestica</i> L. | | <i>Crataegus Chamaméspilus</i> Jacq. | |
| <i>Pyrus domestica</i> Sm. | | <i>Méspilus Chamaméspilus</i> L. | |
| The Whitty Pear Tree. | | <i>Sorbus Chamaméspilus</i> Crantz. | |
| <i>Cormier, Sorbier cultivé, Fr.</i> | | <i>The Bastard Quince.</i> | |
| <i>Speyerlingsbaum, Sperberbaum, Ger.</i> | | App. i. <i>Species of Pyrus not sufficiently</i> | known. - - - 923 |
| <i>Sorbo, Ital.</i> | | <i>P. alnifolia</i> Lindl. | |
| <i>Varieties</i> ☼ - - - 921 | | <i>P. tomentosa</i> Dec. | |
| 2 <i>malifórmis</i> Lodd. ☼ | | - <i>Malus tomentosa</i> Dum. | |
| <i>La Corne-Pomme, Fr.</i> | | <i>P. rubicunda</i> Hoffmans. | |
| 3 <i>pyrifórmis</i> Lodd. ☼ | | XIX. CYDONIA Tourne. ☼ - 929 | |
| <i>La Corne-Poire, Fr.</i> | | THE QUINCE TREE. | |
| 32. <i>lanuginosa</i> Dec. ☼ ? Hungary | pl. 184. 924 | <i>Pyrus</i> sp. L. | |
| The woolly-leaved Service Tree. | | 1. <i>vulgàris</i> Pers. ☼ Crete pl. 186. 929 | |
| <i>P. hybrida lanuginosa</i> Hort. | | The common Quince Tree. | |
| <i>Sorbus lanuginosa</i> Kit. | | <i>Pyrus Cydonia</i> L. | |
| 33. <i>spúria</i> Dec. ☼ Kamtschatka | pl. 185. fig. 645. 924 | <i>C. europæa</i> Sav. | |
| The spurious Service Tree. | | <i>Varieties</i> ☼ - - - 929 | |
| <i>P. hybrida</i> Mœnch. | | 1 <i>pyrifórmis</i> Hort. ☼ | |
| <i>Sorbus spúria</i> Pers. | | 2 <i>malifórmis</i> Hort. ☼ | |
| <i>Méspilus sorbifolia</i> Poir. | | 3 <i>lusitánica</i> Du Ham. ☼ | |
| <i>P. sorbifolia</i> Bose, not of Sm. | | 2. <i>sinénsis</i> Thoun. ☼ China pl. 187. 931 | |
| ? <i>P. sambucifolia</i> Cham. | | The China Quince Tree. | |
| <i>Variety</i> ☼ - - - 925 | | <i>Pyrus sinénsis</i> Poir. | |
| 2 <i>péndula</i> Hort. ☼ | | 3. <i>japónica</i> Pers. ☼ Japan fig. 952. 931 | |
| <i>Sorbus hybrida pëndula</i> Lodd. | | The Japan Quince Tree. | |
| <i>P. spúria sambucifolia</i> Hort. Brit. | | ? <i>Pyrus japónica</i> Thunb. | |
| App. i. <i>Additional Species of Pyrus belonging</i> | to the Section Sorbus. - - - 925 | <i>Chœnomiles japónica</i> Lindl. | |
| <i>P. foliolosa</i> Wall. | | <i>Varieties</i> ☼ - - - 932 | |
| <i>P. hircina</i> Wall. | | 2 <i>flòre álbo</i> ☼ | |
| Other Sorts. | | 3 <i>flòre semi-pleno</i> ☼ | |
| § vii. <i>Adenórachis</i> Dec. ☼ - 925 | | App. i. <i>Other Species of Cydonia.</i> 932 | |
| 34. <i>arbutifolia</i> L. <i>fil.</i> ☼ North America | fig. 616. 925 | <i>C. Sumbushia</i> Hamilt. | |
| The Arbutus-leaved Aronia. | | | |
| <i>Crataegus pyrifolia</i> Lam. | | | |
| <i>Ardunia pyrifolia</i> Pers. | | | |
| ? <i>Crataegus serrata</i> Poir. | | | |
| <i>Méspilus arbutifolia</i> Schmiddl. | | | |
| <i>Varieties</i> ☼ - - - 926 | | | |
| 2 <i>intermèdia</i> Lindl. ☼ | | | |

App. I. *Half-hardy Species of Rosacææ*,
 § *Pômææ*, not belonging to any of the Genera
 containing hardy Species. - - 932

Raphiôlepis Lindl. - - 932

indica Lindl. fig. 653.

Cratægus indica L.
salicifolia Lindl. fig. 654.
rùbra Lindl.
Cratægus rubra Lour.
Méspilus sinensis Poir.
phœostemon Lindl.
R. indica Bot. Reg.
 Laureiri Spreng.
Cratægus indica Lour.
spiralis Don's Mill.
Méspilus spiralis Blume.

Eriobôtrya Lindl. - - 933

japónica Lindl. fig. 655, 656.

Méspilus japónica Thunb.
Cratægus Bibas Lour.
 Lou-Kœt, Japan.
elliptica Lindl.
Méspilus Cuiila Ham. MSS.
cordata Lindl.
obtusifolia Dec.
chinensis G. Don.

Kagenéckia Ruiz et Pav. - - 934

cratægoides D. Don. fig. 657.

K. cratægifolia Lindl.
oblônga R. et P.
lanceolata.
glutinosa.

App. II. *Half-hardy Species of Rosacææ*,
 belonging to the Suborder *Sanguisôrbeæ*. 934

Margyriérpus setôsus R. et P. - - 934

E'mpetrum pinuatum Lam. Dict.
Ancistrum barbatum Lam. III.

Cercocárpus fothergillôides H. B. et Kunth. - - 934

E'rtolônia guierôides Moc. et Sesse.

Acæ'na Fahl. - - 934

Potèrium L. - - 935
spinôsum L. - - 935
caudatum Ait. fig. 658.

Cliffortia - - 935

ilicifolia L.
obcordata L.

Calycanthacææ Lindl. 935

I. CALYCA'NTHUS Lindl. 935, 936

THE CALYCANTHUS, or American Allspice.

Calycanthi sp. 1.
Bittèria Du Ham., not of L.
Bcurrèria Ehrh.
Bastèria Adams.
Pompadoûra Buchoz.
Calycante, Fr.
Kelch Blume, Ger.

1. floridus L. N. America f. 659. 936

The flowery Calycanthus, or Carolina Allspice.

C. stérilis Walt.
Sweet-scented Shrub, in Carolina.
Common American Allspice.

Varieties - - - 936

- 1 oblôngus Dec. 936
- 2 ovatus Dec. 936
- 3 asplenifolius Lodd. 936
- 4 fèrax Lodd. 936
- 5 glaucus Lodd. 936
- 6 inodorus Lodd. 936
- 7 longifolius Lodd. 936
- 8 variegatus Lodd. 936

2. (f.) glaucus Willd. Carol. f. 660. 937
 The glaucous-leaved Calycanthus, or fertile-
 flowered American Allspice.

C. fértilis Walt.

Variety - - - 957

2 oblôngifolius Nutt. 957

3. lævigatus Willd. N. Amer. f. 661. 937

The glabrous-leaved Calycanthus.

C. fèrax Michx.
C. pennsylvanicus Lodd.

II. CHIMONA'NTHUS Lindl. 935, 937

THE CHIMONANTHUS, or Winter Flower.

Méràtia Nees.
Calycanthi sp. L.

1. fragrans Lindl. Japan f. 662. 938

The fragrant-flowered Chimonanthus.

Calycanthus præcox L.
Méràtia fragrans Nees.
O'bai, or *Robai*, Kæmpf.
The Winter Flower.
Calycante de Japon, Fr.
Japanische Kelch Blume, Ger.

Varieties - - - 938

2 grandiflorus Lindl. f. 663.

3 luteus Hort. 938

Granatæcæ D. Don. 939

1. PU'NICA Tourn. 939

THE POMEGRANATE TREE.

The Carthaginian Apple.
Grenadier, Gr.
Granate, Ger.
Melograno, Ital.
Granados, Span.

1. Granatum L. Asia - - 939

The common Pomegranate Tree.

Varieties - - - 939

1 rùbrum Dec. f. 664.

2 r. flore pleno Trew. 939

3 albescens Dec. 939

4 a. flore pleno Dec. 939

5 flavum Hort. 939

2. (G.) nana L. North America

fig. 665. 942

The dwarf Pomegranate.

P. americana nana Tourn.
P. Granatum nanum Pers.

Onogræcæ Lindl. 942

§ FUCHSIE'Æ. 942

Fúchia L. 942

microphylla H. B. et Kunth. fig. 666.

thymifolia H. B. et Kunth.

rosea Ruiz et Pav.

F. lycioides Bot. Reg.

parviflora Lindl.

F. ovata Moc. et Sesse.

arborescens Sims.

F. racemosa Moc. et Sesse.

F. amæ'na Hort.
F. humicoides Moc. et Sesse.

- gracilis Lindl.
 - F. decussata* Grah.
 - var. 2 multiflora Lindl.
 - macrostemon Ruiz et Pav.
 - var. 2 tenella Dec.
 - F. gracilis* var. *tenella* Lindl.
 - cónica Lindl.
 - virgata Smt.
 - coccinea Ait. fig. 667.
 - F. pendula* Saussb.
 - F. mozghanica* Lam.
 - Nahusia coccinea* Schneevogt.
 - Skimmia coccinea* Moench.
 - apétala Ruiz et Pav.
 - discolor Lindl.
 - hacillaris Lindl.
 - hybrida Smt.
 - globosa Hort.
 - var. 2 longiflora Hort.
 - longiflora Hort.
 - F. longipedunculata* Beaton.
 - speciosa Hort.
 - grandiflora Hort.
 - praecox Hort.
 - Thompsonii Hort.
 - excorticata L. fl. fig. 668.
 - Skimmia excorticata* Forst.
- Other Varieties

Lythraceæ.

945

- Heimia - 945
- salicifolia* Link et Otto. fig. 669.
- Nesaea salicifolia* H. B. et Kunth.
- Lythrum flavum* Spreng.
- myrtifolia* Hort. Berol.
- Lythrum apetalum* Spreng.
- linearifolia* Hort.
- H. salicifolia* var.
- syphilitica* Dec. - 945
- Lagerstrœmia
- indica* L. fig. 670.
- parvifolia* Roerb.
- reginae* Roerb.

Tamaricææ. - 946

I. TAMARIX Desr. - 946, 947
THE TAMARISK.

The species of *Tamarix* of authors that have 4 stamens and 5 stamens.
Tamarix, Fr.
Tamariscus, Ger.

1. *gállica* L. France - fig. 671. 947
The French Tamarisk.

- T. narbonensis* Lob.
- Tamariscus gállicus* All.
- Tamariscus pentandrus* Lam.

Varieties - 948

- 1 *súbtillis* Ehrenberg
- 2 *narbonensis* Ehrh.
- 3 *milótica* Ehrh.
- 4 *arborea* Sieb.
- 5 *mannifera* Ehrh.
- 6 *heterophýlla* Ehrh.
- 7 *libanótica* Lodd.
- The Rosemary-like Tamarisk.

App. i. Other hardy Species or Varieties of *Támarix*. - 948

- T. tetrándra* Pal.
- T. gállica* Habl.
- T. láta* Willd.
- T. elongata* Led.
- T. gracilis* Willd.
- T. hispida* Willd.
- T. pentándra* var. *Pal.*
- T. gállica* var. *g* Willd.
- T. tomentosa* Smith.
- T. canescens* Desv.

- T. ramosíssima* Led.
- T. gállica* Sievers.
- T. Pallásii* Desv.
- T. pentándra* Pall.
- T. gállica* Bieb.
- T. paniculata* Stev.
- T. cupressiformis* Led.
- T. parviflora* Dec.
- T. tetrágyra* Ehrenb.
- T. etíusa* Ehrenb.

App. ii. Half-hardy Sorts of *Támarix*. 949

- T. africana* Poir.
 - T. gállica* var. *γ* Willd.
 - T. canariensis* Willd.
 - T. pycnocarpa* Dec
 - T. passerinoides* Del.
 - var. 1 *divaricata* Ehrh.
 - 2 *Hammonia* Ehrh.
 - 3 *macrocarpa* Ehrh.
- Other Species or Sorts.

II. MYRICARIA Desr. 946, 949
THE MYRICARIA.

The species of *Támarix* of authors that have monadelphous stamens.

1. *germánica* Desr. Europe f. 672. 949
The German Myricaria, or German Tamarisk.

- Támarix germánica* L.
- Tamariscus germánicus* Lob.
- Tamariscus decándrus* Lam.
- Támarix decándra* Moench.
- Tamaris d'Allemagne*, Fr.
- Deutschen Tamarisken*, Ger.

2. *dahùrica* Dec. Siberia - 949
The Dahurian Myricaria.
Támarix dahùrica Willd.

App. i. Other Sorts of *Myricària* not yet introduced. - - 950

- M. squamòsa* Desr.
- ? *M. dahùrica* var.
- M. longifolia* Dec.
- Támarix germánica* Pall.
- Támarix decándra* Pall.
- Támarix longifolia* Willd.
- M. linearifolia* Desv.
- M. herbàcea* Desf.
- Támarix germánica subherbàcea* Pall.
- M. bracteata* Royle.
- M. elegans* Royle.

Philadelphææ. 950

I. PHILADELPHUS L. - 950, 951
THE PHILADELPHUS, or Mock Orange.

- Syringa Tourn.*, not of L.
- Philadelphus*, Fr.
- Pfeifenstrauch* (Pipe Shrub), Ger.
- Pipe Prinet*, Gerard.
- The *Syringa* of the gardens.

§ i. Stems stiff and straight. Flowers in Racemes. - - 951

1. *coronarius* L. South Europe fig. 673. 951

- The garland Philadelphus.
- Syringa suarèolens* Moench.
- Varieties** - 951

- 1 *vulgàris* Schkuhr.
- 2 *nànus* Mill.
- 3 *flòre plèno* Lodd.
- 4 *variegatus* Lodd

2. (c.) *inodorus* L. ☼ Carolina f. 674. 952
The scentless-flowered Philadelphus.
Syringa inodora Mönch.
P. lãxus in many English gardens.

3. (c.) *Zeyheri* Schrad. ☼ N. Amer. 952
Zeyher's Philadelphus.

4. *verrucosus* Schrad. ☼ North America
f. 675. 952

The warted Philadelphus.
P. grandiflorus Lindl.

5. (v.) *latifolius* Schrad. ☼ N. America
fig. 676. 953

The broad-leaved Philadelphus.
P. pubescens Cels.

6. (v.) *floribundus* Schrad. ☼ N. Amer. 953
The abundant-flowered Philadelphus.

§ ii. *Stems more slender, rambling, twiggy,
and loose. Flowers solitary, or 2 or 3
together.* - - - - - 954

7. *lãxus* Schrad. ☼ N. Amer. f. 677. 954
The loose-growing Philadelphus.
P. humilis Hort.
P. pubescens Lodd.

8. (l.) *grandiflorus* Willd. ☼ N. Amer. 954
The large-flowered Philadelphus.
P. inodorus Hort.
P. lãxus Lodd.

9. *hirsutus* Nutt. ☼ North America.
f. 678, 678a. 954
The hairy-leaved Philadelphus.
P. villosus and *P. gracilis* Lodd.

10. *tomentosus* Wall. ☼ Nepal - 955
The woolly-leaved Philadelphus.
P. nepalensis Lodd.
? *P. triflorus* Royle.

II. DECUMARIA L. ☼ - ☼ 950. 955

THE DECUMARIA.
Forsythia Walt., not of Vahl.

1. *bãrbara* L. ☼ Carolina f. 679, 680. 955
The barbarous Decumaria.
D. radicans Mönch.
D. Forsythia Michx.
D. prostrata Lodd.

Variety ☼ - - - - - 955
2 *sarmentosa* Dec. ☼
D. sarmentosa Bosc.
Forsythia scandens Walt.

App. I. *Half-hardy ligneous Plants of the
Order Philadelphiceæ.* - 956

Deutzia scãbra Thunb. ☼ - f. 681. 950.

D. corymbosa R. Br.
Philadelphus corymbosus Wall.
D. staminea R. Br.
Philadelphus stamineus Wall.
D. Brunonia Wall.
Leptospermum scãbrum Wall.

Myrtaceæ. ☼ - ☼ - ☼ - ☼ 956

Sect. I. MELALEUCÆÆ. ☼ - ☼ - ☼ - 956

Tristania nerifolia R. Br. ☼ - fig. 682. 956
Melaleuca nerifolia Sims.
Melaleuca salicifolia Bot. Rep.
Beaufortia decussata R. Br. ☼ - fig. 633. 957
Calothamnus villosa R. Br. ☼ - fig. 634. 957

C. gracilis R. Br. ☼ - - - - - 957
C. quadrina R. Br. ☼ - - - - - 957
C. clavata Cunningham. ☼ - - - - - 957
Melaleuca squamea Labill. ☼ - - - - - 957
M. linearifolia Sm. ☼ - - - - - fig. 685. 957
Metrosideros hysteroptifolia Cav.
M. pulchella R. Br. ☼ - - - - - 957
M. hypericifolia Sm. ☼ - - - - - fig. 687. 957
M. squarrosa Sm. ☼ - - - - - 957
Eudesia tetragona R. Br. ☼ - - - - - fig. 686. 958

Sect. II. EULEPTOSPERMÆÆ. ☼ - ☼ - ☼ - 958

Eucalyptus resinifera Sm. ☼ - - - - - fig. 688, 689. 958
The Iron Bark Tree.
E. robusta Sm. ☼ - - - - - pl. 188. 959
The Stringy Bark Tree.
E. viminalis Labill. ☼ - - - - - fig. 692, 693. 959
E. amygdalina Labill. ☼ - - - - - fig. 694, 695. 959
E. piperita Sm. ☼ - - - - - fig. 696, 697. 959
The Blue Gum Tree.
E. cordata Labill. ☼ - - - - - fig. 698. 959
E. pulverulenta Sims. ☼ - - - - - fig. 697. 959
E. cordata Hort. Berol.
E. diversifolia Bonpl. ☼ - - - - - 959
E. conata Dum.
E. piperita Hort. Berol.
E. heterophylla Swt.
E. angustifolia Hort.
Several other Species of *Eucalyptus* - - - - - 959
Angophora cordifolia Car. ☼ - - - - - fig. 699. 960
Metrosideros hispidus Sm.
A. lanceolata Cav. ☼ - - - - - fig. 702. 960
The Apple Tree of New Holland.
Metrosideros splendens Gærtn.
Callistemon salignus Dec. ☼ - - - - - 960
Metrosideros salignus Sm.
C. lanceolatus Dec. ☼ - - - - - fig. 700. 960
Metrosideros lanceolatus Sm.
Metrosideros curvata Bot. Mag.
Metrosideros corifolius Vent. ☼ - - - - - fig. 705. 961
The Coris-leaved Iron Wood.
Leptospermum ambiguum Sm.
Leptospermum grandifolium Sm. ☼ - - - - - fig. 701. 961
L. lanigerum Ait. ☼ - - - - - 961
Philadelphus lanigerus Ait.
Billotia marginata R. Br. ☼ - - - - - 961
Leptospermum marginatum Labill.
B. flexuosa D. Don. ☼ - - - - - 961
Fabricia myrtifolia Gærtn. ☼ - - - - - fig. 703. 961
F. stricta Lodd. ☼ - - - - - 961
Bacckia virgata Andr. ☼ - - - - - fig. 704. 961
Leptospermum virgatum Forsk.
Melaleuca virgata L. fil.

Sect. III. MYRTEÆÆ. ☼ - ☼ - ☼ - 961

Psidium Cattle-yauum Sabine. ☼ - - - - - fig. 706. 961
Cattle's Guava.
P. coriaceum Marsh.
P. chinense Lodd.

Myrtus communis L. ☼ - - - - - 961
The common Myrtle.

Varieties ☼ - - - - - 963

§ i. *Melanocarpa* Dec. ☼ - - - - - 963

- 1 *romãna* Dec. ☼ - - - - -
The common broad-leaved, or Roman, Myrtle.
- 2 *tarentina* Mill. Jc. ☼ - - - - -
The Tarentum, or Box-leaved, Myrtle.
- 3 *italica* Mill. Dict. ☼ - - - - -
The Italian, or upright, Myrtle.
- 4 *bac'tica* Mill. Dict. ☼ - - - - -
The Andalusian, or Orange-leaved, Myrtle.
- 5 *lusitanica* L. ☼ - - - - -
The Portuga Myrtle.
M. acuta Mill. Dict.
Subvar.? The Nutmeg Myrtle ☼ - - - - -
- 6 *hãlgica* Mill. Dict. ☼ - - - - -
The broad-leaved Dutch Myrtle.
Subvar. The double-flowered Myrtle.
- 7 *mucronata* L. ☼ - - - - -
The Rosemary, or Thyme-leaved Myrtle.

§ ii. *Leucocarpa* Dec. ☼ - - - - - 963

8 *leucocarpa* Dec. ☼ - - - - -
The white-berried Myrtle.

§ iii. *Garden Varieties.* ☼ - - - - - 963

1. Gold-striped broad-leaved Myrtle ☼ - - - - -

	Page
2. Broad-leaved Jew's Myrtle	961
3. Gold-striped Orange-leaved Myrtle	961
4. Silver-striped Italian Myrtle	961
5. Striped-leaved Myrtle	961
6. Silver-striped Rosemary-leaved Myrtle	961
7. Silver-striped Nutmeg Myrtle	961
8. Cock's-comb, or Bird's-nest Myrtle	961
9. Spotted-leaved Myrtle	961
<i>M. tomentosa</i> Lf.	fig. 707. 964
<i>M. canescens</i> Lour.	
<i>M. tenuifolia</i> Sm.	964

Sect. IV. CHAMELAUCIÆ. 964

<i>Chamaelancium ciliatum</i> Desf.	964
<i>Calythrix glabra</i> R. Br.	fig. 708. 964
<i>C. ericoides</i> Cunningham.	964
<i>Darwinia fascicularis</i> Rudge.	964

Passifloræcæ. 964

<i>Passiflora cærulea</i> L.	fig. 709. 964
var. 2 <i>angustifolia</i> Hort.	
3 <i>glaucophylla</i> Hort.	
4 <i>Colvillii</i> Swt.	
5 <i>racemosa</i> Hort.	
<i>P. incarnata</i> L.	fig. 710. 965
The flesh-coloured Granadilla, or May Apple.	
<i>P. filifolia</i> L.	965
The Lime-tree-leaved Passion Flower.	

Other Species or Varieties. 965

<i>Dioscorea adiantifolia</i> Dec.	fig. 711. 965
<i>Passiflora adiantifolia</i> Bot. Mag.	
<i>Tacsonia pennatisipula</i> Juss.	965
<i>Passiflora pinnatisipula</i> Cav.	

Crassulæcæ. 965

<i>Sedum populifolium</i> L.	965
The Poplar-leaved Sedum, or Stonecrop.	
<i>Anacampseros populifolia</i> Haw.	
<i>Sempervivum arborescens</i> L.	fig. 712. 965
The Tree Houseleek.	

Mesembryæcæ. 966

<i>Mesembryanthemum</i> L.	966
----------------------------	-----

Nitrariæcæ Lindl. 966

1. NITRARIA L.	966
THE NITRARIA.	
1. Schöberi L.	Russia 966
Schober's Nitraria.	
Varieties	966
1 <i>sibirica</i>	fig. 713. N. sibirica Pall.
2 <i>caspica</i>	fig. 714. N. caspica Pall.
2. tridentata Desf.	Barbary 967
The three-toothed-leaved Nitraria.	
<i>Peganum retusum</i> Forsk.	

Cactæcæ. 967

<i>Opuntia vulgaris</i> Mill.	967
The common Indian Fig, or Prickly Pear.	
<i>Cactus Opuntia</i> L.	

Grossulæcæ Dec. 967

I. RIBES L. 967, 968

THE RIBES.	
<i>Grossularia</i> Tourn.	
<i>Chrysobotrya</i> , <i>Calobotrya</i> , <i>Corcosma</i> , and <i>Ribes</i> , Spach.	
<i>Grosseller</i> , Fr.	
<i>Johannisbeere</i> , Ger.	
<i>Kruisbes</i> , Dutch.	
<i>Uva Spina</i> , Ital.	
<i>Grossella</i> , Span.	

§ i. *Grossulariæ* Ach. Rich. 968

A. Flowers greenish white.

1. *oxyacanthoides* L. Canada f. 715. 968
The Hawthorn-leaved Gooseberry.
2. *setosum* Lindl. N. Am. f. 716. 969
The bristly Gooseberry.
3. *triflorum* W. North America
fig. 717. 969
The three-flowered Gooseberry.
R. *stamineum* Horn.
4. (t.) *niveum* Lindl. North America
fig. 718. 969
The snowy-flowered Currant-like Gooseberry.
5. (t.) *Cynosbati* L. Canada f. 719. 970
The Dog-Bramble Gooseberry.
R. ? *triflorum* var.
6. (t.) *divaricatum* Dougl. N. America
fig. 720. 970
The spreading-branched Gooseberry.
R. ? *triflorum* var.
R. ? *Grossularia* var. *triflora subvar.*
7. (t.) *irriguum* Dougl. North America
fig. 721. 971
The well-watered Gooseberry.
R. ? *triflorum* var.
8. *hirtellum* Michx. N. America 971
The slightly hairy-branched Gooseberry.
9. *gracile* Michx. North America 971
The slender-branched Gooseberry.
10. *aciculare* Smith Siberia - - 971
The acicular-spined Gooseberry.
R. *Uva-crispa* Sievers.
11. *Grossularia* L. Europe - 972
The common Gooseberry.
R. *Uva-crispa* (Ed. Fl. Dan. *Grossularia hirsuta* Mill. R. *Uva-crispa* var. 5 *salpa* Dec. *Feaberry*, Cheshire and the N. of England. *Feabs*, Norfolk. *Groszert*, in Scotland. *Grosseller à Maugrean*, Fr. *Griselle*, in Piedmont. *Gemeine Stachelbeere*, Ger. *Uva Spina*, Ital.

	Page		Page
<i>Varieties</i> - - -	972	3 <i>earneum</i> <i>Berl.</i> <i>MSS.</i> ♀	
2 <i>Uva-crispa</i> <i>Sm.</i> ♀		<i>R. rubrum domesticum</i> 2 <i>baccis</i>	
<i>R. Uva-crispa</i> <i>L.</i>		<i>carneis</i> <i>Walfr.</i>	
<i>Uva-crispa</i> <i>Fuch.</i>		4 <i>variegatum</i> <i>Dec.</i> ♀	
<i>Uva-spinna</i> <i>Math.</i>		5 <i>album</i> <i>Desf.</i> ♀	
<i>R. Uva-crispa</i> <i>var. 1 sylvestris</i>		6 <i>foliis lateo variegatis</i> <i>Du Ham.</i> ♀	
<i>Berlandier.</i>		7 <i>foliis albo variegatis</i> <i>Du Ham.</i> ♀	
3 <i>spinosissima</i> <i>Berl.</i> <i>MSS.</i> ♀		18. (r.) <i>alpinum</i> <i>L.</i> ♀ <i>Europe</i> f. 725. 979	
4 <i>reclinata</i> <i>Berl.</i> <i>MSS.</i> ♀		The alpine red Currant.	
<i>R. reclinatum</i> <i>L.</i>		<i>Varieties</i> - - - - 979	
<i>Grossularia reclinata</i> <i>Mill.</i>		1 <i>sterile</i> <i>Walfr.</i> ♀	
5 <i>Besseriana</i> <i>Berl.</i> <i>MSS.</i> ♀		<i>R. dioicum</i> <i>Mench.</i>	
<i>R. hybridum</i> <i>Besser.</i>		2 <i>bacciferum</i> <i>Walfr.</i> ♀	
6 <i>subinermis</i> <i>Berl.</i> <i>MSS.</i> ♀		3 <i>pumilum</i> <i>Lindl.</i> ♀ f. 726.	
? <i>R. G. reclinata</i> <i>subvar.</i>		4 <i>foliis variegatis</i> <i>Hort.</i> ♀	
7 <i>macrocarpa</i> <i>Dec.</i> ♀		19. (r.) <i>petraeum</i> <i>Wulf.</i> ♀ <i>Carinthia</i>	
8 <i>bracteata</i> <i>Berl.</i> <i>MSS.</i> ♀		<i>fig. 727. 979</i>	
<i>Other Varieties.</i> ♀		The rock red Currant.	
The Red Champagne. ♀		<i>R. alpinum</i> <i>Delarb.</i>	
Horseman's Green Gage. ♀		The woolly-leaved Currant.	
The Red Rose. ♀		The red Marsh-mallow-leaved Currant.	
a. <i>Sorts of Gooseberries belonging to this Division not yet introduced.</i> - - - 974		20. (r.) <i>spicatum</i> <i>Robs.</i> ♀ <i>England</i>	
<i>R. saxosum</i> <i>Hook.</i>		<i>fig. 728. 980</i>	
<i>R. saxatile</i> <i>Dougl.</i> <i>MSS.</i>		The spiked-flowered red, or Tree, Currant.	
<i>R. triflorum</i> <i>Büzel.</i>		21. (r.) <i>carpathicum</i> <i>Küt.</i> ♀ <i>Carpathian</i>	
<i>R. rotundifolium</i> <i>Michx.</i>		<i>Momtains</i> - - - - 980	
<i>R. caucasicum</i> <i>Adams.</i>		The Carpathian red Currant.	
? <i>R. G. Uva-crispa</i> <i>subvar.</i>		<i>R. acerinum</i> <i>Rochel.</i>	
<i>R. cuneifolium</i> <i>R. et P.</i>		22. (r.) <i>multiflorum</i> <i>Küt.</i> ♀ <i>Croatia</i>	
<i>R. cucullatum</i> <i>Hook. et Arn.</i>		<i>fig. 729. 980</i>	
<i>B. Flowers red.</i>		The many-flowered red Currant.	
12. <i>speciosum</i> <i>Pursh.</i> ♀ <i>California</i>		<i>R. spicatum</i> <i>Schultes.</i>	
<i>fig. 722. 974</i>		23. (r.) <i>procumbens</i> <i>Pall.</i> ✕ <i>Siberia</i>	
The showy-flowered Gooseberry.		<i>fig. 730. 981</i>	
<i>R. stambucum</i> <i>Sm.</i>		The procumbent red Currant.	
<i>R. fuchsoides</i> <i>Fl. Mex.</i>		<i>R. polycarpum</i> <i>Gmel.</i>	
<i>R. triacanthum</i> <i>Menzies.</i>		24. (r.) <i>prostratum</i> <i>L.</i> ✕ <i>North America</i>	
a. <i>Sorts of Gooseberries not yet introduced belonging to this Subsection.</i> - - - 975		<i>fig. 731. 981</i>	
<i>R. Menziesii</i> <i>Pursh.</i>		The prostrate red Currant.	
<i>R. ferax</i> <i>Sm.</i>		<i>R. glandulosum</i> <i>Ait.</i>	
<i>R. microphyllum</i> <i>H. B. et Kunth.</i>		<i>Variety</i> ✕ - - - - 981	
§ ii. <i>Botrycarpum</i> <i>Dec.</i> ♀ - 975		2 <i>laxiflorum</i> ✕	
13. <i>orientale</i> <i>Poir.</i> ♀ <i>Syria</i> - - 975		<i>R. affine</i> <i>Dougl.</i> <i>MSS.</i>	
The Eastern Currant-like Gooseberry.		<i>R. laxiflorum</i> <i>Pursh.</i>	
14. <i>saxatile</i> <i>Pall.</i> ♀ <i>Dahuria</i> - 976		25. (r.) <i>resinosum</i> <i>Pursh.</i> ♀ <i>N. America</i>	
The rock Currant-like Gooseberry.		<i>fig. 732. 981</i>	
? <i>R. alpinum</i> <i>Sievers.</i>		The resinous red Currant.	
15. <i>Diacantha</i> <i>L. fil.</i> ♀ <i>fig. 723. 976</i>		26. (r.) <i>trifidum</i> <i>Michx.</i> ✕ <i>N. Amer.</i> 981	
The twin-prickled Currant-like Gooseberry.		The trifid-calyced red Currant.	
16. <i>lacustre</i> <i>Poir.</i> ♀ <i>N. Amer.</i> f. 724. 976		27. (r.) <i>albinervum</i> <i>Michx.</i> ♀ <i>Canada</i> 982	
The lake-side Currant-like Gooseberry.		The white-nerved-leaved red Currant.	
? <i>R. oryzacanthoides</i> <i>Michx.</i>		28. <i>rigens</i> <i>Michx.</i> ♀ <i>Canada</i> - 982	
<i>R. echinatum</i> <i>Dougl.</i> <i>M.S.</i>		The stiff racemed red Currant.	
§ iii. <i>Ribesia</i> <i>Dec.</i> ♀ ✕ - 977		29. <i>punctatum</i> <i>R. et P.</i> ♀ <i>Chili</i> f. 733. 982	
A. <i>Flowers greenish, or greenish yellow, or reddish; and Fruit, in a wild State, red.</i> - 977		The dotted-leaved red Currant.	
17. <i>rubrum</i> <i>L.</i> ♀ <i>Europe</i> - - 977		30. (p.) <i>glandulosum</i> <i>R. et P.</i> ♀ <i>Chili</i> 982	
The common red Currant.		The glandular-calyced red Currant.	
<i>R. vulgare</i> <i>N. Du. Ham.</i>		h	
<i>Grosseller</i> <i>common, Fr.</i>			
<i>Gemeine Johannisbeere, Ger.</i>			
<i>Roode Aaltesen Boom, Dutch.</i>			
<i>Varieties</i> - - - - 977			
1 <i>syvestre</i> <i>Dec.</i> ♀			
2 <i>hortense</i> <i>Dec.</i> ♀			
<i>R. rubrum</i> <i>Lois.</i>			

	Page
a. <i>Species or Varieties of Ribes belonging to the Sub-division A of the Section Ribesia, which are not yet introduced.</i>	983
<i>R. fraxans</i> Pall.	
<i>R. heterostachium</i> Meyer.	
<i>R. bracteosum</i> Dougl.	
<i>R. tuberosum</i> Meyer.	
<i>R. tuberosum</i> Fischscholtz	
<i>R. macrobotrys</i> R. et P.	
<i>R. albidum</i> R. et P.	
<i>R. ciliatum</i> Willd.	
<i>R. jordanianum</i> H. B. et Kunth.	
<i>R. hirtum</i> Thunb. et Dougl.	
<i>R. fragilium</i> H. B. et Kunth.	
<i>R. campanulatum</i> Thunb. et Dougl.	
<i>R. affine</i> H. B. et Kunth.	
<i>R. Känthii</i> Bort.	
<i>R. multiflorum</i> H. B. et Kunth, not of Kit.	
<i>R. Takie</i> D. Don.	
<i>R. acuminatum</i> Wall.	
<i>R. villosum</i> Wall.	
B. <i>Flowers greenish yellow, sometimes with the Tips of the Sepals and Petals red. Fruit Black.</i>	983
31. <i>nigrum</i> L. Europe fig. 734. 983	
The black Currant.	
<i>R. ædium</i> Moench	
<i>Capis Poivrier</i> , Fr.	
<i>Schwartz</i> <i>Johannisbeere</i> , Ger.	
<i>Varieties</i>	984
2 <i>baccæ flavidæ</i> Gard. Mag.	
3 <i>baccæ viridæ</i> Hort.	
4 <i>foliis variegatis</i> Vilmorin.	
<i>Garden Varieties.</i>	
32. (n.) <i>triste</i> Pall. Siberia 985	
The sad-coloured, or dark-blossomed, black Currant.	
<i>R. altaicum</i> Lodd.	
33. (n.) <i>floridum</i> L'Herit. N. America fig. 735. 985	
The flowery black Currant.	
<i>R. nigrum</i> 2 L.	
<i>R. pennsylvanicum</i> Lam.	
<i>R. recurvatum</i> Michx.	
<i>Ribesium nigrum</i> , ꝯc. Dill.	
<i>Varieties</i>	986
2 <i>grandiflorum</i> Hort.	
3 <i>parviflorum</i> Hort.	
34. (n.) <i>inebrians</i> Lindl. N. America fig. 736. 986	
The intoxicating black Currant.	
35. <i>cereum</i> Dougl. N. W. America fig. 737. 986	
The waxy-leaved black Currant.	
36. <i>viscosissimum</i> Pursh. N. America fig. 738. 987	
The very clammy black Currant.	
<i>Coreösmia viscosissima</i> Spach.	
37. <i>hudsonianum</i> Richardson. North America 987	
The Hudson's Bay black Currant.	
<i>R. petiolare</i> Dougl.	
38. <i>glaciæ</i> Wall. Nepal - 987	
The icy black Currant.	
a. <i>Species or Varieties of Ribes belonging to the Division B of the Section Ribesia, which have not yet been introduced</i>	987
<i>R. Biebersteinii</i> Berl.	
<i>R. longistachium</i> Bieb.	
<i>R. viscosum</i> R. et P.	

<i>C. Flowers deep red. Fruit black.</i>	Page
39. <i>sanguineum</i> Pursh. N. America fig. 739. 988	
The bloody, or red-flowered, Currant.	
<i>R. malvaceum</i> Sm.	
<i>Calobotrya sanguinea</i> Spach.	
<i>Varieties</i>	988
2 <i>glutinösium</i> Benth. f. 740.	
<i>R. angustum</i> Dougl. MSS.	
3 <i>malvaceum</i> Benth. f. 741.	
4 <i>atro-ribens</i> Hort.	
a. <i>Species or Varieties of Ribes belonging to the Division C of the Section Ribesia, which have not yet been introduced.</i>	989
40. <i>atro-purpureum</i> Meyer. Siberia 989	
The dark purple-flowered Currant.	
<i>Varieties</i>	989
1 Flowers deep purple, &c.	
2 Leaves rather pubescent, &c.	
3 Flowers paler, &c.	
§ iv. <i>Symphocalyx</i> Dec. - 989	
41. <i>aureum</i> Pursh. N. Am. f. 742. 989	
The golden-flowered Currant.	
<i>R. palmata</i> Desf.	
<i>Chrysobotrya revoluta</i> Spach.	
<i>Varieties</i>	989
1 <i>præcox</i> Lindl.	
2 <i>villösium</i> Dec.	
<i>R. longiflorum</i> Fraser's Cat.	
3 <i>serotinum</i> Lindl. f. 743.	
42. (a.) <i>tenniflorum</i> Lindl. N. Amer. fig. 744. 990	
The slender-flowered Currant.	
<i>R. aureum</i> Colla.	
<i>R. flavum</i> Berl.	
<i>R. missouriensis</i> Hort.	
<i>Chrysobotrya Lindleyana</i> Spach.	
<i>Varieties</i>	990
1 <i>fructu nigro</i>	
2 <i>fructu luteo</i>	
43. (a.) <i>flavum</i> Coll. - 990	
The yellow-flowered Currant.	
<i>R. aureum</i> 3 <i>sanguineum</i> Lindl.	
<i>R. palmatum</i> Desf.	
<i>R. aureum</i> Ker, but not of Pursh.	
<i>Chrysobotrya intermedia</i> Spach.	
App. i. <i>A Classification of the Species and Varieties of Ribes in the Horticultural Society's Garden in 1836.</i>	990
—	
<i>Escalloniaceæ.</i>	992
I. <i>ITEA</i> L.	992
THE ITEA.	
<i>Cedrela</i> Lour.	
<i>Dicondangia</i> Michx.	
I. <i>virginica</i> L. N. Amer. f. 745. 992	
The Virginian Itca.	
II. <i>ESCALLONIA</i> Mutis. - 993	
THE ESCALLONIA.	
<i>Stereoxylon</i> R. et P.	

- E. rubra Pers. * — - fig. 746. 993
Varieties * — - - 993
 1 glabrifolia Hook. et Arn. * — —
 2 albiflora Hook. et Arn. * — —
E. glandulosa Bot. Cab.
 3 pubescens Hook. et Arn. * — —

- E. montevidensis Dec. * — fig. 747. 993
E. floribunda var. *β* montevidensis Schlecht.
E. bifida Link et Otto.

- E. floribunda H. B. et Kunth. * — 994

- E. resinosa Pers. * — - - 994

- Stereoxylon resinosum* R. et P.

- E. pulverulenta Pers. * — - - 994

- Stereoxylon pulverulentum* R. et P.

Saxifragæ. * * — — 294

Tribe HYDRANGEÆ. * * — —

- I. HYDRANGEA L. * * — — 994
 THE HYDRANGÆA.

Hydrangæa and *Horténsia* Juss.

1. arborescens L. * North America
 fig. 748. 994

The arborescent Hydrangæa.

Varieties * - - - 995

- 1 vulgaris Ser. Michx.

H. vulgaris Michx. and ? Pursh.

H. arborescens Curt.

H. frutescens Meneh.

- 2 discolor Ser. *

2. (a.) cordata Pursh. * North America
 fig. 749. 995

The cordate-leaved Hydrangæa.

3. nivea Michx. * North America
 fig. 750. 995

The snowy-leaved Hydrangæa.

H. radiata Walt., not of Sm.

Variety - - - 995

- 2 glabella Ser.

4. quercifolia Bartram. * North America
 fig. 751. 995

The Oak-leaved Hydrangæa.

H. radiata Sm., not of Walt.

5. heteromalla D. Don. * Nepal 996
 The diverse-haired-leaved Hydrangæa.

- App. i. *Half-hardy Species of Hydrangæa.* 996

H. horténsia Sieb. * — fig. 752.

The Chinese Guelder Rose.

H. horténsia Sm.

Horténsia opuloides Lam.

Horténsia speciosa Pers.

Prinula mutabilis Lour.

Viburnum serratum and *Viburnum tomentosum* Thunb.

H. vestita Wall. * —

H. altissima Wall. * —

Umbellæcæ. * * — — 997

- I. BUPLEURUM Tourn. * * — — 997
 * — - - - 997

THE BUPLEURUM, or *Hare's Ear.*

Tenoria and *Bupræstis* Spreng.

Séseli Bauh. and Mor.

Buphore, Oreille de Lièvre, Fr.

Hasenöhrllein, Ger.

- I. fruticosum L. * S. Europe f. 753. 997
 The shrubby Bupleurum.

Tenoria fruticosa Spreng.

Bupræstis fruticosa Spreng. Mag.

Séseli æthiopicum Bauh.

Séseli frater Mor.

- App. i. *Half-hardy Species of the Genus Bupleurum.* - - 998

B. gibraltàricum Lam. * — —

B. coriaceum L'Hérit.

B. obliquum Vahl.

B. arborescens Jacq.

Tenoria coriacea Spreng.

B. verticale Ort.

B. plantagineum Desf. * — —

Tenoria plantaginea Spreng.

B. canescens Schoub. * — —

B. frutescens L. * — —

Araliæcæ. * * — - 998

- I. ARALIA L. * - - - 998
 THE ARALIA, or *Angelica Tree.*

Aralia sp. L.

Araliæ vera Blum.

1. spinosa L. * N. America f. 754. 999
 The spiny Aralia.

Aralie, Fr. and Ger.

Spikeward, N. Amer.

Other suffruticose Species of Aralia. - 999

- II. HEDERA Swartz. * - 998, 999
 THE IVY.

Aralia, sect. Gymnopterum Blum.

Hedera and *Aralia* sp. L.

Lierre, Fr.

Ephen, Ger.

1. Hélix L. * Europe - - 1000
 The common Ivy.

Varieties * - - 1000

- 1 vulgaris Dec. * f. 755.

- 2 canariensis Dec. * —

H. canariensis Willd.

- ? 3 chrysocarpa Dec. * —

H. poetica C. Bauh.

H. chrysocarpa Dalech.

H. Dionysias J. Bauh.

H. Hélix Wall.

Additional Varieties in British Gardens.

- 4 foliis argenteis Lodd. * —

The silver-striped Ivy.

- 5 foliis aureis Lodd. * —

The golden-striped Ivy.

- 6 digitata Lodd. * —

The palmate, or hand-shaped, Ivy.

- 7 arborescens Lodd. * —

The arborescent, or Tree, Ivy.

Hamamelidæcæ. * * 1006

- I. HAMAMELIS L. * * 1006, 1007
 THE HAMAMELIS, or *Wych Hazel.*

Trilopus Mith.

1. virginica L. * * North America
 fig. 756, 757. 1007

The Virginian Hamamelis.

Virginische Zaubernuss, Ger.

	Page	Page
<i>Varieties</i> - - -	1005	
2 <i>parviflora</i> Nutt.		
3 <i>macrophylla</i> .		
<i>H. macrophylla</i> Pursh.		
App. i. <i>Other Species, not yet introduced.</i>	1008	
<i>H. persica</i> Dec.		
<i>H. chinensis</i> R. Br.		
II. FOTHERGILLA L. ☼	1007, 1008	
THE FOTHERGILLA.		
<i>Hamanidlis</i> L.		
I. <i>alnifolia</i> L. ☽ North America	1008	
The Alder-leaved Fothergilla.		
<i>F. Gärdeni</i> Michx.		
<i>Hamanidlis monoleuca</i> L.		
<i>Varieties</i> ☽ - - -	1008	
1 <i>obtusata</i> Sims. ☽ f. 759.		
<i>F. major</i> Lodd.		
<i>F. alnifolia</i> L. n.		
2 <i>acuta</i> Sims. ☽		
<i>F. Gärdeni</i> Jacq.		
3 <i>major</i> Sims. ☽ f. 758.		
4 <i>serotina</i> Sims. ☽		
—		
<i>Cornaceæ.</i> ☽ ☼	1009	
I. CORNUS L. ☽ - - -	1009	
THE DOGWOOD.		
<i>Cornouiller</i> , Fr.		
<i>Hartriegel</i> , Ger.		
§ i. <i>Nudiflora</i> Dec. ☽ ☼ -	1010	
1. <i>alternifolia</i> L. ☽ ☽ North America	fig. 760. 1010	
The alternate-leaved Dogwood.		
<i>C. alternata</i> Marsh.		
2. <i>sanguinea</i> L. ☽ North America	fig. 761. 1010	
The blood-red-leaved, or common, Dogwood.		
<i>C. foemina</i> Rafin.		
<i>Virga sanguinea</i> Matth.		
<i>Femle</i> Cornel, Dogberry Tree, Hound		
<i>Tree</i> , <i>Hound's-berry Tree</i> , <i>Prickwood</i> ,		
<i>Gaten</i> or <i>Gatten Tree</i> , <i>Gat'er</i> or <i>Gatter</i>		
<i>Tree</i> , <i>Cateridge Tree</i> , <i>Wild Cornel</i> .		
<i>Cornouiller sauvee</i> , <i>sanguin</i> , or <i>femelle</i> ;		
<i>Puin</i> , or <i>Buis punais</i> , Fr.		
<i>Roth'er Hartriegel</i> , Ger.		
<i>Sanguinello</i> , Ital.		
<i>Varieties</i> ☽ - - -	1011	
2 <i>Pürshii</i> Don. ☽		
<i>C. sanguinea</i> Pursh.		
3 <i>foliis variegatis</i> ☽		
<i>C. canadensis</i> Mill. var. Lodd.		
3. <i>alba</i> L. ☽ N. America fig. 762. 1011		
The white-fruited Dogwood.		
<i>C. solida</i> Jacq. Michx.		
<i>C. latifolia</i> Mill.		
<i>Varieties</i> ☽ - - -	1012	
2 <i>circinata</i> G. Don. ☽		
<i>C. circinata</i> Cham.		
<i>Mispawmawid</i> , <i>M etbquim-pee-</i>		
<i>mawid</i> , <i>and</i> <i>M. asata</i> , of the		
Cree Indians.		
3 <i>sibirica</i> Lodd. ☽		
4. (a.) <i>stricta</i> Lam. ☽ North America	fig. 763, 764. 1012	
The straight-branched Dogwood.		
<i>C. fastigiata</i> Michx.		
<i>C. sanguinea</i> Walt., not of L.		
<i>C. cyanocarpus</i> Gmel.		
<i>C. canadensis</i> Hort. Par.		
<i>C. cerida</i> Meerb., not of Lam.		
<i>Varieties</i> ☽ - - -	1013	
2 <i>asperifolia</i> Lodd. ☽		
3 <i>sempervirens</i> Lodd. ☽		
5. (a.) <i>paniculata</i> L'Hérit. ☽ ☽ North	America - - - fig. 765. 1012	
The panicled-flowering Dogwood.		
<i>C. racemosa</i> Lam.		
<i>C. foemina</i> Mill.		
<i>C. citrifolia</i> Hort. Par.		
<i>Varieties</i> ☽ - - -	1013	
2 <i>albida</i> Ehrh. ☽		
3 <i>radiata</i> Pursh. ☽		
6. (a.) <i>sericea</i> L'Hérit. ☽ N. America	fig. 766. 1013	
The silky Dogwood.		
<i>C. lanuginosa</i> Michx.		
<i>C. alba</i> Walt., not of L.		
<i>C. Anomum</i> Du Roi.		
<i>C. rubiginosa</i> Ehrh.		
<i>C. ferruginea</i> Hort. Par.		
<i>C. candidissima</i> Mill.		
<i>C. cyanocarpus</i> Meench, not of Gmel.		
<i>Varieties</i> ☽ - - -	1013	
2 <i>oblongifolia</i> Dec. ☽		
<i>C. oblongifolia</i> Rafin.		
3 <i>asperifolia</i> Dec. ☽		
<i>C. asperifolia</i> Michx.		
7. (a.) <i>circinata</i> L'Hérit. ☽ N. America	fig. 767. 1014	
The rounded-leaved Dogwood.		
<i>C. tomentosa</i> Michx.		
<i>C. rugosa</i> Lam.		
<i>C. virginiana</i> Hort. Par.		
8. <i>oblonga</i> Wall. ☽ Nepal - - -	1014	
The oblong-leaved Dogwood.		
<i>C. paniculata</i> Hamilt.		
<i>C. macrophylla</i> Wall. ☽ - - -	1011	
<i>C. excelsa</i> H. B. et Kunth. ☽ - - -	1011	
§ ii. <i>Involucrata</i> Dec. ☽ -	1014	
<i>C. disciflora</i> Me. et Sesse. ☽ - - -	1011	
<i>C. grandis</i> Cham.		
<i>C. japonica</i> Thunb. ☽ - - -	1011	
<i>Viburnum japonicum</i> Spreng.		
9. <i>mas</i> L. ☽ Europe pl. 189. f. 768. 1014		
The male Dogwood, the Cornel, or Cornelian		
<i>Cherry Tree</i>		
<i>C. mascula</i> L'Hérit. &c.		
<i>Long Cherry Tree</i> , <i>Cornelia</i> ,		
<i>Cornouiller male</i> , <i>Cornus</i> , <i>Cornettes</i> , Fr.		
<i>Kornel Kirsche</i> , <i>Hartriegel</i> , Ger.		
<i>Varieties</i> ☽ - - -	1015	
2 <i>fructu cære coloris</i> N. Du Ham. ☽		
3 <i>variegatus</i> ☽		
10. <i>florida</i> L. ☽ N. America f. 769. 1017		
The Florida Dogwood.		
<i>Virginian Dogwood</i>		

18. (O.) Oxycóccos Pursh. ☼ North America - 1041
 The Cranberry-fruited Guelder Rose.
V. opuloides Mühl.
V. trilobum Marsh.
V. Opulus americana Ait.
 Variety ☼ - - - 1041
 2 subintegrifolius Hook. ☼

19. (O.) edule Pursh. ☼ North America 1041
 The edible-fruited Guelder Rose.
V. Opulus edulis Michx.

20. (O.) molle Michx. ☼ North America 1041
 The soft-leaved Guelder Rose.
V. adnifolium Marsh.

A. Species of *Viburnum* belonging to the Section *Opulus*, not yet introduced. 1041
V. microcarpum Cham. et Schlecht.
V. polycarpum Wall.
V. cylindricum Huan.
V. grandiflorum Wall.
V. erubescens Wall.
V. coriaceum Blum.
 Several other Species.

Sect. II. LONICEREÆ. ☼ ☼ ☼ ☼ ☼ 1041

II. DIERVYLLA Tourn. ☼ 1027. 1042
 THE DIERVYLLA.

Lonicera sp. L.
Weigela Thunb.
Weigela Pers.

1. canadensis Willd. ☼ North America fig. 794, 795. 1042
 The Canadian Diervilla.
Lonicera Diervilla L.
D. Tournefortii Michx.
D. humilis Pers.
D. lata Pursh.
D. trifida Mönch.
D. arcadiensis Du Ham.

App. i. Species of *Diervilla* not yet introduced. - - 1042

D. japonica Dec.
Weigela japonica Thunb.
D. corænsis Dec.
Weigela corænsis Thunb.
Weigela corænsis Pers.

IV. LONICERA Desf. ☼ ☼ ☼ ☼ ☼ 1027. 1042

THE LONICERA, or Honeysuckle.
Lonicera sp. L. and many authors.
Caprifolium and *Xylostemum* Juss.
Xylostemum, *Caprifolium*, *Chamaecerasus*, and *Periclymenum*, Tourn.
Caprifolium and *Lonicera* Rœm.
Lonicera and *Xylostemum* Torrey.
Chèvrefeuille, Fr.
Geissblättl, *Honightume*, *Lonicere*, Ger.

§ i. *Caprifolium* Dec. ☼ ☼ ☼ ☼ 1043
Caprifolium Juss. and Rœm.
Lonicera Torr., not of Schult.

A. Flowers ringent. — *Caprifolium* Tourn.

1. *Periclymenum* L. ☼ Europe 1043
 The Woodbine, or common Honeysuckle.
Periclymenum, Ger.
Periclymenum germanicum Riv.

Periclymenum hortense Gesn.
Caprifolium Periclymenum Rœm. et Schultes.
Caprifolium sylvaticum Lam.
Caprifolium Itali.
 Woodbind.
Chèvrefeuille des Bois, Fr.
Wildes gemeines Geissblättl, Ger.
Gewoone Kamperfoelie, Dutch.
Lego Bosco, Ital.
Mudre Selva, Span.

Varieties ☼ - - - 1044
 2 serotinum Ait. ☼ fig. 797.
Periclymenum germanicum Mill.
 3 belgicum ☼
Periclymenum germanicum Mill.
 Diet.
 4 quercifolium Ait. ☼

2. *Caprifolium* L. ☼ Europe f. 798. 1045
 The Goat's-leaf, or pale perfoliate, Honeysuckle.
Periclymenum perfoliatum Ger.

3. (C.) etrusca Santi. ☼ fig. 799. 1046
 The Etruscan Honeysuckle.
L. etrusca Hort. Fl. Austr.
Caprifolium etruscum Rœm. et Schult.
Periclymenum Gouan.
Caprifolium italicum perfoliatum præcox Tourn.

4. impléxa Ait. ☼ Sicily fig. 800. 1046
 The interwoven, or Minorca, Honeysuckle.
Caprifolium impléxum Rœm. et Schult.
 Variety ☼ - - - 1047
 2 baleárica Fr. ☼
Caprifolium baleáricum Dum.
L. baleárica Dec.
L. Caprifolium Desf.

5. flava Sims. ☼ North America fig. 801. 1047
 The yellow-flowered Honeysuckle.
Caprifolium flavum Ell.
Caprifolium Frasca Pursh.

6. (f.) pubescens Swt. ☼ North America fig. 802. 1047
 The pubescent Honeysuckle.
Caprifolium pubescens Goldie.
L. hirsuta Eaton.
L. Goldii Spreng.

7. parviflora Lam. ☼ North America fig. 803, 804. 1048
 The small-flowered Honeysuckle.
Caprifolium parviflorum Pursh.
L. dioica L.
L. media Murr.
Caprifolium bracteosum Michx.
Caprifolium dioicum Rœm. et Schult.
Caprifolium glaucum Mönch.
 Glaucous Honeysuckle.
Chèvrefeuille dioique, Fr.
Meevgrünnes Geissblättl, Ger.
Middelboore Kamperfoelie, Dutch.

8. (p.) Douglasii Dec. ☼ N. America 1048
 Douglas's Honeysuckle.
Caprifolium Douglasii Lindl.

9. grata Ait. ☼ N. America f. 805. 1048
 The pleasant, or evergreen, Honeysuckle.
Caprifolium gratum Pursh.
L. virginiana Marsh.
 ? *Periclymenum americanum* Mill.

L. microphylla Hook. - - - 1049

- Page
- B. *Limb of Corolla nearly equal.* — *Periclymenum*
Tourn. - 1049
10. *sempervirens* *Ait.* 2. North America
fig. 806. 1019
The evergreen *Trumpet* Honeysuckle.
Caprifolium sempervirens Michx.
Periclymenum sempervirens Mill.
Alaternus sempervirens Koehl.
Periclymenum virginianum Riv.
Varieties 2. - - - 1049
2 *majör* *Ait.* 2.
The large *Trumpet* Honeysuckle.
3 *minor* 2. —
The small *Trumpet* Honeysuckle.
? *L. comata* Meerb.
11. *ciliösa* *Poir.* 3. North America 1050
The ciliated-*leaved* Honeysuckle.
Caprifolium ciliösum Pursh.
L. ciliata Dietr.
12. *occidentälis* *Hook.* 3. N. Amer. 1050
The Western Honeysuckle.
Caprifolium occidentale Lindl.
Caprifolium ciliösum Dougl. MSS.
L. pilosa Willd. - 1050
Caprifolium villosum H. B. et Kunth.
- § ii. *Xylösteum* Dec. 2. 3. 4. 5. 1050
Xylösteum Juss.
Lonicera *Rum. et Schult.*
Xylösteum and *Chamaecerasus* Tourn.
Xylösteum and *Isika Adams.*
Cöba's Neck.
The *Fly* Honeysuckle.
Hackenkirsche. Ger.
Hondsbezien, Hondsksarsen. Dutch.
- A. *Ovaries and Berries altogether distinct. Stems
scandent. Flowers irregular.* — *Nintöa* Dec.
1050
13. *confusa* *Dec.* 3. Japan fig. 808. 1050
The confused Honeysuckle.
Nintöa confusa Scot.
Lonicera japonica Andr., not of Thunb.
Nintöa, Nintoo, Kampf.
Caprifolium japonicum Lond. Hort. Brit.
14. *longiflora* *Dec.* 3. China - 1051
The long-flowered Honeysuckle.
Caprifolium longiflorum Sabine.
Nintöa longiflora Swt.
Caprifolium japonicum G. Don.
Caprifolium nepalense G. Don.
15. *japönica* *Thunb.* 2. Japan
fig. 809, 810. 1051
The Japan Honeysuckle.
Nintöa japonica Swt.
L. chinensis Hort. Kew.
L. flexuosa Lodd., not of Thunb.
L. glabrata Roxb.
Caprifolium chinense Lond. Hort. Brit.
Caprifolium flexuosum Hort.
- a. *Hardy Species of Lonicera belonging to the Divi-
sion Nintöa of the Section Xylösteum, not yet
introduced.* - - - 1052
L. cochinchinensis Don's Mill.
L. Xylösteum Lour.
L. Tellari Hook. et Arn.
L. Periclymenum Lour.
L. Lechenaultii Wall.
L. glabrata *Hall.*
L. nigra *Thunb.*
L. acuminata *Hall.*
L. diversifolia *Wall.*
L. fistulifera *Wall.*
Xylösteum Agostiniana G. Don.
Xylösteum Nipponica *Thunb.*
- L. lanceolata* *Wall.*
L. canescens *Schousb.*
L. biflora *Desf.*
L. bracteata *Royle.*
Several other species of *Lonicera.*
- b. *Berries distinct, or usually connate together at the
Base, and diverging at the Tip. Corolla hardly
pithous at the Base, or equal.* — *Chamaecerasus*
Dec. - - - 1052
16. *tatarica* *L.* 2. Siberia f. 811, 812. 1052
The Tartarian Honeysuckle.
Xylösteum cordatum Mönch.
Xylösteum tataricum Dum.
Varieties 2. - - - 1053
2 *albiflora* *Dec.* 2.
L. pyrenaica *Willd.*
3 *rubriflora* *Dec.* 2.
L. grandiflorum *Lodd.*
L. sibirica *Hort.*
4 *lutea* *Lodd.* 2.
5 *latifolia* *Lodd.* 2.
17. (t.) *nigra* *L.* 2. Europe - 1053
The black-fruited Honeysuckle.
Caprifolium viscum Lam.
Chamaecerasus nigra *Delarb.*
Variety 2. - - - 1053
2 *campaniflora* 2. fig. 813, 814.
Xylösteum campaniflorum *Lodd.*
18. (t.) *ciliata* *Mühl.* 2. N. Amer. 1053
The ciliated-*leaved* Honeysuckle.
Xylösteum ciliatum Pursh.
L. tatarica *Michx.*, not of *L.*
L. canadensis *Ram. et Schult.*
19. *pyrenaica* *L.* 2. Pyrenees - 1054
The Pyrenean Honeysuckle.
Caprifolium pyrenaicum Lam.
Xylösteum pyrenaicum *Tourn.*
20. *punicea* *Sims.* 2. North America
fig. 815. 1054
The crimson-flowered Honeysuckle.
Symphoricarpos puniceus *Swt.*
21. *Xylösteum* *L.* 2. North America
fig. 816. 1054
The bony-wooded, or upright, *Fly* Honeysuckle.
Caprifolium dumetorum Lam.
Xylösteum dumetorum *Mönch.*
Varieties 2. - - - 1055
2 *leucocarpa* *Dec.* 2.
3 *xanthocarpa* *Dec.* 2.
4 *melanocarpa* *Dec.* 2.
22. *flexuosa* *Thunb.* 2. Japan - - 1055
The flexible-stemmed Honeysuckle.
L. naga *Thunb.*, not of *L.*
L. brachyloba *Dec.*
- a. *Hardy Species of Lonicera, belonging to the Divi-
sion Chamaecerasi of the Section Xylösteum, not
yet introduced.* - - - 1055
L. bispidula *Pall.*
- c. *Berries either distinct or joined together. Corolla
very pithous at the Base. Erect bushy Shrubs.*
— *Cephaantha* *Dec.* - - - 1055
23. *involverata* *Banks.* 2. North America
fig. 817, 818, 819. 1055
The involuerated Honeysuckle.
Xylösteum involueratum *Rich.*

a. *Hardy Species of Lonicera belonging to the Division Caphanthæ of the Section Xylösteum, which arc not yet introduced.* - - - 1056

L. gibbosa Willd.
Xylösteum mexicanum H. B. et Kunth.
 L. Mociniæna Dec.
L. gibbosa Moc. et Sesse.
 L. Ledebourii Eschsch.

D. *Berries two on each Petuncle, joined together in one, which is bi-umbellate at the Apex.* Erect branching Shrubs. — Isikæ Adans. - 1056

24. *alpigena* H. Europe fig. 820, 821. 1056

The alpine Honeysuckle.
Caprifolium alpinum Lam.
Caprifolium alpinum Gærtm.
Isika alpigena Börck.
Isika lucida Mœnch.
Xylösteum alpinum Lodd.
Chamæcerasus alpigena Delarb.
Cherry Woodbine.
Heckenkirsche. Gr.

Variety - - - 1056

2 sibirica Dec. L. sibirica Vest.

25. (a.) *microphýlla* Willd. Siberia 1057
 The small-leaved Honeysuckle.
L. alpigena Sievers.

26. *oblongifolia* Hook. North America fig. 822. 1057

The oblong-leaved Honeysuckle.
Xylösteum oblongifolium Goldie.

27. *cærulea* L. Eur. f. 823, 824. 1057
 The blue-berried Honeysuckle.
L. villosa Mühl.
Xylösteum villosum Michx.
Xylösteum Solomis Eaton.
L. vclutina Dec.
L. altaica Pall.
Xylösteum cæruleum canadense Lam.
Xylösteum canadense Du Ham.
Caprifolium cæruleum Lam.
Chamæcerasus cærulea Delarb.
L. pyrenæica Pall.
L. Pallasiæ Led.

28. (c.) *orientalis* Lam. Asia Minor 1058

The Oriental Honeysuckle.
L. caucásica Pall.
L. cærulea Güld.
Chamæcerasus orientális laurifolia Tourn.

29. *ibérica* Bieb. - - - 1058
 The Georgian Honeysuckle.
Xylösteum ibericum Bieb.

a. *Hardy Species of the Genus Lonicera belonging to the Division Isikæ of the Section Xylösteum, not yet introduced.* - - - 1058

L. Webbiana Wall.
 L. Govaniana Wall.
 L. angustifolia Wall.
 Some other Species.

V. SYMPHORICARPOS Dill. 1027. 1058

THE ST. PETER'S WORT.
Symphoricarpa Neck.
Symphoria Pers.
Anisanthus Willd.
Lonicera sp. L.

1. *vulgáris* Michx. N. Am. f. 825. 1058
 The common St. Peter's Wort.
Lonicera Symphoricarpos L.
S. parviflora Desf.
Symphoria conglomerata Pers.
Symphoria glomerata Pursh.

Variety - - - 1059

2 *fóliis variegáteis* S. *glomerata fóliis variegáteis* Lodd.

2. *racemósus* Michx. fig. 826. 1059
 The racemose-flowered St. Peter's Wort, or Snouberry.
Symphoria racemosa Pursh.
L. lucocarpa Hort.

App. i. *Hardy Species of Symphoricarpos not yet introduced.* - - - 1059

S. occidentális Richards.
 Wolfberry, Amer.

VI. LEYCESTERIA Wall. 1027. 1060

THE LEYCESTERIA.

1. *formosa* Wall. Nepal f. 827. 1060
 The beautiful Leycesteria.
Hamelia connata Puerari MSS.

Rubiaceæ. - 1061

↑ ↓ ☼ ☼ ☼ ☼ ☼ ☼ ☼ ☼

I. CEPHALANTHUS L. - 1061
 THE BUTTON-WOOD.

1. *occidentális* L. North America fig. 828, 829. 1061
 The Western Button-wood.
C. oppositifolius Mœnch.
Swamp Globe Flower. Amer.

Variety - - - 1062

2 *brachypodus* Dec.

Some other Species of Cephalanthus. 1062

App. I. *Half-hardy ligneous Plants belonging to the Order Rubiaceæ.* - 1062

Pinckneya pubens Michx. ↑ ↓ fig. 830.
Pinckneya pubescens Pers.
Cinchona caroliniana Poir.

Sertosa fructida Conn. ↓ ↓ fig. 831.

Lajcium japonicum Thunb.
Lajcium fistulosum L. fil.
Lajcium indicum Kt
Dysodia fasciculata Lamr.
Buchozia coprosmodites L'Herit.
Dysodia fructida Salisb.

Spermacoce fruticosa Desf.

Plócama pendula Ait. ↓ ↓
Bartlingia scoparia Rehb.

Phyllis Nibla L. ↓

Anthospermum æthiopicum L. ↓ ↓
Ambraria Heisteri Walsh.

Rubia fruticosa Ait. ↓ ↓
R. fruticosa canadensis Poir.

Bouvardia Jacquinii H. B. et Kunth. ↓ ↓
B. triphylia Hort.

Honatania coecinea Bot. Rep.
Manettia glabra Cham. et Schlecht. ↓ ↓
M. cordifolia Mart.

Lobeliaceæ. ↓ ↓ 1063

Tupa salicifolia G. Don. ↓ ↓

Lobelia Tupa Ait.
Lobelia gigantea Sims.
Lobelia salicifolia Swt.

Lobelia arborea Forst. ? ↓ ↓

L. superba Cham. ↓ ↓

Campanulaceæ. ↓ ↓ 1063

Muschia aurea Dumort. ↓ ↓
Campanula aurea L.

Compósita. - 1063

I. STEHELINA Lessing. 1063. 1064

THE STEHELINA.

Stacheline, Fr. and Ger.

- 1. dubia L. S. Europe fig. 832. 1064
The doubtful, or Rosemary-leaved, Stachelina.
S. rosmarinifolia Cass.

App. i. Half-hardy Species of Stachelina. 1065

- Stachelina arbor-scens* L.
- S. Chamæpice* L.

II. BACCHARIS R. Br. 1063. 1065

THE BACCHARIS, or Ploughman's Spikenard.
Bacchante, Fr.
Baccharis, Ger.

- 1. halimifolia L. N. Am. f. 833. 1065
The Sea-Purslane-leaved Baccharis, or the
Grundsel Tree.
Senecio arbor-scens Hort. Kew.

- 2. angustifolia Pursh. N. Amer. 1065
The narrow-leaved Baccharis, or Ploughman's
Spikenard.

- B. glomeratiflora* Mich. 1066
- B. Dioscoridis* W. 1066

III. IVA L. 1064. 1066

THE IVA.

- 1. frutescens L. N. Amer. f. 834. 1066
The shrubby Iva.
Agerato affinis peruviana frutescens Pluk.
Bastard Jesuit's Bark Tree.

IV. SANTOLINA L. 1064. 1066

THE SANTOLINA, or Lavender Cotton.
Santoline, Fr.
Heiligenpflanze, Ger.

- 1. Chamæcyparissus L. France
fig. 835. 1067

The Dwarf Cypress Santolina, or common
Lavender Cotton.

- 2. (C.) squarrosa H. Europe 1067
The squarrose (? leaved) Santolina.
Abrotanum junina jolitis Ericæ Moris.

- 3. viridis H. Europe - 1067
The green Santolina.

- 4. rosmarinifolia L. Spain f. 836. 1067
The Rosemary-leaved Santolina.

V. ARTEMISIA Cass. 1064. 1068

THE ARTEMISIA.

- 1. Abrotanum L. Europe fig 837. 1068
The Abrotanum Artemisia, or Southernwood.
Abrotanus mas Dod.
Ohl Man.

*Armoise Aurone, Aurone des Jardins, la
Citronelle, la Garderobe*, Fr.
*Therrancle, Wermuth, Stabwurtz, Garten-
wurtz*, Ger.
Abrotano, Ital., Span., and Port.

- Varieties* - - - 1068
- 2. humile Hort.

- 3. tobolskianum Hort. \blacksquare
A. tobolskiana Lodd.

- 2. Santonica L. Asia fig. 838. 1068
The Santonica Artemisia, Tartarian Southern-
wood, or Wormseed.
Artemisia fruticosa, &c., Gmel.

- 3. arbor-scens L. Levant - 1069
The arborescent Artemisia, or Tree Worm-
wood.
Absinthium arbor-scens Lob.
Absinte, Armoise en Arbre, Fr.

App. i. Other hardy Species of Artemisia. 1069

- A. procera*

App. ii. Half-hardy Species of Artemisia. 1069

- 1. argentea Ait. fig. 851.

VI. HELICHRYSUM Lessing. 1064. 1070

THE HELICHRYSUM, or Everlasting Flower.
Part of *Gnaphalium* L.

- 1. Stechas D. Don. Europe - 1070
The Stechas Helichrysum, or common, Shrubby
Everlasting Flower.
Gnaphalium Stechas L.
Stechas citrina Dod.

App. i. Half-hardy Species of Helichrysum. 1070

- H. frutescens* D. Don. fig. 840.
Astelma frutescens Bot. Reg.
Gnaphalium frutescens L.
Gnaphalium grandiflorum Bot. Rep.

- H. congestum* D. Don. fig. 811.

- Gnaphalium congestum* Lamb.

- H. aequalatum* Lodd.

- H. crassifolium* Lodd.

- H. diversifolium* Lodd.

- H. ericoides* Lodd.

VII. CINERARIA Lessing. 1064. 1071

THE CINERARIA.
Cinéraire, Fr.
Aschenpflanze, Ger.

- 1. maritima L. S. Europe 1071
The sea-side-inhabiting Cineraria, or the Sea
Ragwort.

Cineraria Dod.
Jacobæa maritima Bonp.
Sicilian Ragwort.

App. i. Half-hardy Species of Cineraria. 1071

- C. cruenta* fig. 842.

- C. lactea*

- C. canescens*

- C. hybrida*

- C. populifolia*

- C. bicolor*

- C. lanata* fig. 115.

- C. geoffolia*

- C. amplexoides* fig. 813.

Agnostia calceatis Cass.

App. I. Half-hardy Genera belonging to the Order Compósita. - 1072

- Carlowsia salicifolia* March. \blacksquare
Ombroma salicifolia Link.

- Arctostaphylos aspera* L.

- Dudleya carnosum* H. K.

- D. spmosum* H. K.

- Ber. klavya grandiflora* W.

- Cultania* R. Br.

- Otholma halimifolia* Bot. Cal. fig. 847.

- O. pycnantha* L.




























- O. pinnata* Bot. Mag.


- O. pectinata*





















- O. coronopifolia*

Page


Page

- Osteospermum pistiferum L.  fig. 848., 849., 850.
 Caléndula chrysanthemifolia Ven.  fig. 851.
 Mutisia latifolia D. Don.  fig. 852.
 Dáhlia Cav.
 Atalánthus pinnátus D. Don. 
Prenánthes pinnáta L.
 Sónchus fruticosus Jacq.  fig. 855.
 Vernónia acutifolia Hook. 
Aster argophyllus Lab.  fig. 854.
Hartónia argophylla Caley.
 A. angustifolius Jacq. 
 A. aculeátus Lab.  fig. 855.
 Chrysócoma Comaúrea L.  fig. 856.
 Brachyla'na nereifolia Smt. 
Baccharis nereifolia L.
 Conÿza carolinénsis Jacq. 
 Podánthus *Mithqui* Lindl.  fig. 857.
 Calcitium calcicum Spr.  fig. 858.
Cactilia salicina Lab.
 Cactilia L.
 Franzéria *artemisioides* W. 
 F. ambrosioides Cav. 
 G'edera prolifera Thunb. 
 P. pèthrum *funiculáccum* W. En.  fig. 859.
Jemelia maderénsis 
 Athanasia pubescens L. 
 Balsamita ageratifolia Desf. 
 Pientza flabelliformis W. 
Tanacetum flabelliforme L'Hérit.
 Eriocéphalus africanus L. 
 Senécio élegans plena rábra Bot. Mag. 
 S. lílicianus Link. 
 Tarchonánthus camphorátus Lam. 
 Eriócoma frágrans D. Don. 
 Various other genera of Compositæ.


Epacridàcæa.  1075


- Styphèlia R. Br.
 Stenanthèra pinifolia R. Br. 
Styphèlia pinifolia Spreng.
 Cyathòdes glauca Labil. 
Trochocárpa glauca Spreng.
 C. Oxycèdrus R. Br. 
 C. aceròsa R. Br. 
 Lissánthe sápidá R. Br. 
 L. subuláta 
 L. strigòsa 
 L. daynhüdes 
 L. ciliáta 
 Leucopogon lanceolátus R. Br. 
Styphèlia lanceolata Sm.
Styphèlia purpúflora Andr.
 L. Richei R. Br.  fig. 860.
 L. polystachyus Lodd.
 L. apiculátus Sm.
 L. parviflorus Lindl.
 Monótoea elliptica R. Br. 
 M. álbens 
 M. lineáta 
 M. scopária 
 Trochocárpa laúrina R. Br.  fig. 861.
Styphèlia cornifolia Rudge.
 E'pacrís grandiflora Sm.  fig. 862.
E. longiflora Cav.
 Lysinèma R. Br.
 Andersónia sprengelioides R. Br.  fig. 865.
 Sprengèlia incarnáta Bot. Cab. 
 Sphenótoma gracilis Smt. 
Draconophyllum gerdlei R. Br.
 Various other Genera of Compositæ.


Ericàcæa. - - 1076



Sect. I. ERÍCEÆ. - 1076




§ i. *Ericæa normàles.*  1076

I. ERÍCA D. Don.  1076. 1079

THE HEATH.

Erica sp. of L. and other authors.!





1. Tétralix L.  N. Europe. f. 864. 1079


The four-leaved Heath.

- E. botulifórmis Sal.
- E. barbárica Rail.
- E. pínnula Park.
- E. T. rábra Hort. Eric. Wob.

The cross-leaved Heath.

Varieties  - - - 1079







- 1 rábra Hort. Eric. Wob. 
- 2 cárnea Loudon's II. B. 
- 3 álba Hort. Eric. Wob. 
- 4 Mackaíana 
- E. Mackaíana Bab.

2. cinèrea L.  N. Europe f. 865. 1079

The grey Heath.

- E. mutábilis Salisb.
- E. húmilis Neck.
- E. tenuifolia, Ger.
- E. c. rábra Bedf. Hort. Eric. Wob.

Varieties  - - - 1080




- 2 atropurpúrea Lodd. 
- 3 álba Lodd. 
- 4 pállida Lodd. 
- 5 carnescens Lodd. 
- 6 prolifera Lodd. 
- 7 strícta Lodd. 


3. arbòrea L.  South Europe - 1080

The Tree Heath.

- E. scoparia Thunb., exclusive of the synonym.
- E. cáffra L.
- E. triflóra Berg.
- E. procéra Sal.

Varieties  - - - 1080

- 2 stylòsa Don's Mill. 
- 3 squarròsa Hort. 
- 4 mínima Hort. 

4. (a.) polytrichifolia Sal.  Port. 1080

The Polytrichum-leaved Heath.

- ? E. arbòrea var.
- ? E. c. stylòsa Hort.


5. (a.) codonòdes Lindl.  Europe


fig. 866. 1080

The bell-shaped-flowered Heath.

6. austràlis L.  Spain fig. 867. 1081


The southern Heath.

- E. pistillóris Sal.

7. strícta Donn.  Italy fig. 868. 1081

The upright Heath.


- E. multicaulis Sal.
- E. còrsica Dre.
- E. ramulòsa Viv.

8. ciliàris L.  Portugal fig. 869. 1081


The ciliated-leaved Heath.

9. sícula Schonberg,  Sicily - 1082

The Sicilian Heath.

II. GYPSOCA'LLIS Sal.  1076. 1082

THE GYPSOCALLIS, or Moor Heath.
Ericæ of others.

1. vágans Sal.  England fig. 870. 1082

The wandering Gypsocallis, or Cornish Moor Heath.

- Erica vágans* L.
- Erica voga* Sal.
- Erica multiflóra* Huds.
- Erica didyma* Stokes.
- Erica purpuráscens* Lam.

	Page
Varieties n. - - -	1082
2 pallida Don's Mill. n.	
3 rubescens Brev. n.	
4 purpurascens Brev. n.	
5 alba Don's Mill. n.	
6 tenella Don's Mill. n.	

2. multiflora D. Don. n. South Europe
fig. 871. 1083

The many flowered Gypsocalis.
Erica multiflora L.
Erica juniperifolia, &c., Garidel.
Erica multiflora longi-pedicellata Wendl.
Erica peduncularis Presl.

3. carnea D. Don. n. Germany
fig. 872. 1083

The flesh colour-flowered Gypsocalis.
Erica carnea L. Sp.
Erica herbacea L. Duss.
Erica saxatilis Sal.

4. mediterranea D. Don. n. S. Eur. 1083

The Mediterranean Gypsocalis.
Erica mediterranea L.
Erica lugubris Sal.

III. CALLUNA Sal. n. - 1076. 1084

THE CALLUNA.

Erica sp. L. and others.

1. vulgaris Sal. n. Europe - - 1084

The Ling, or Heather.

Erica vulgaris L.
La Bruyère, Fr.
Heide, Ger.
Lyug, Dan.
Ljung, Swed.
Erica, Ital.
Brezo, Span.
Urzo, Port.
Weresk, Russ.

Varieties n. - - - 1084

- 1 purpurea Don's Mill. n.
- 2 spuria Don's Mill. n.
- 3 decumbens Don's Mill. n.
- 4 tomentosa Don's Mill. n.
- 5 alba Don's Mill. n.
- 6 flore pleno Don's Mill. n.
- 7 foliis variegatis Don's Mill. n.
- 8 aurea Don's Mill. n.
- 9 coccinea Don's Mill. n.
- 10 spicata Don's Mill. n.
- 11 atro-rubens n.
- 12 serotina n.

App. I. List of hardy Species and Varieties of *Ericaceæ* belonging to the Group *Erica* normales, of which Plants are cultivated for Sale in the Tooting Nursery; with some additional Names from the "Hortus Woburnensis." - - - 1086

App. II. Arrangement of the hardy Heaths included in the preceding List; showing which of them are in Flower, in the open Garden, every Month in the Year; and the Colour of the Flower, and the Height of each. - - - 1088

Page
App. III. List of Cape Heaths which will stand in the open Air, in Autumn, or the middle of Winter, without Protection, with Fahrenheit's Thermometer 7 or 8 Degrees below Freezing, without suffering in any way from such a Degree of Cold. n. l. - 1089

Callista acuminata (Erica L.) Bot. Cab. fig. 873.

2 pallida Hort. Brit.
comosa Hort. Kew. Ic.
2 alba Andr. Heath.
3 rubra Wendl. E.
ferruginea Andr. Heath.
hyacinthoides Andr. Heath.
tenuiflora Andr. Heath.
2 alba Hort. Brit.
3 lutea.
tetragona Andr. Heath.
C. pugionifolia Sal.
ventricosa Bot. Mag.

2 coccinea.
3 stellifera.
4 carnea.
5 alba.
6 superba.
7 erecta.
8 nana.
9 hirsuta.

Ceremia (Erica L.) serpyllifolia Lodd. fig. 874.

Dasyanthus (Erica L.) Sparmanni Andr. Heath.

D. dispersa A. H.
D. hystericiflora L. T.

Désmia (Erica L.) conferta Andr. Heath. fig. 875.

Erica aggregata Wendl. Eric. fig. 876.

2 alba Hort. Brit.
campanulata Andr. Heath.

cerinthoides Bot. Mag.

1 glabriuscula Suet.
2 hispida Suet.
3 major H. Wob.
4 minor H. Wob.
5 nana H. Wob.
6 superba Roll.

congesta Wendl. Eric.

corifolia.

E. articulata B. M.
E. calycina A. H.

Lamprotis corifolia Don's Mill

cupressina Roll.
globosa Andr. Heath.
glomerata Andr. Heath.
gracilis Wendl. Eric.
hispidula.
leucanthera.
lucida Andr. Heath.
? Lamprotis lucida Don's Mill.
margaritacea Andr. Heath.
montana.
pendula Lodd.
perlata.
pubescens.

1 major H. Brit.
2 minor Roll.
3 pubescentior H. Brit.
4 verna H. Wob.
raimentacea Andr. Heath.
setacea Andr. Heath.
tenella Andr. Heath.

Eurylepis (Erica L.) triflora Wendl. Eric.

Gypsocalis (Erica L.) intertexta Lodd., fig. 877

longipedunculata Bot. Cal.

nigrita Don's Mill.

Pachysa physodes Bot. Mag., fig. 878, 879.


Syringoblea cruenta Andr. Heath.

2 superba Roll.

curviflora Andr. Heath.

1 aurantia.
2 rubra.
diaphana Don's Mill.
Ewerona Andr. Heath.
2 glabra.
3 speciosa.
4 longiflora.
5 pilosa.
exulans Lodd

grandiflora Bot. Mag.	Page
1 humilis.	
2 superba.	
ignescens Andr. Heath.	
? longiflora Bot. Cal.	
mammosa Andr. Heath.	
2 minor.	
3 pallida.	
4 rosea.	
? tumida Bot. Reg.	
verticillata Andr. Heath.	
2 major.	
viridescens Bot. Cal.	

App. IV. List of Cape Heaths which are tenderer than those mentioned in the preceding List, and which, when exposed to the Degree of Cold there stated, will be injured by it, but will not suffer, although fully exposed to a Temperature 4 or 5 Degrees below Freezing.  - - - - 1091

Blæria cricoides fig. 880.	
Erica Blæria Wendl. Coll.	
Callista (Erica L.) comosa Wendl. Eric.	
1 álba.	
2 rùbra.	
Coventryna Lodd.	
daphneflora.	
mündula Lodd.	
2 major Lodd.	
pellucidata Andr. Heath.	
2 rùbra H. Wob.	
prægnans Bot. Cal.	
2 coccinea H. Wob.	
Cerámia (Erica L.) urceolaris Hort. Kew. Icon.	fig. 881
? articuláris Don's Mill.	

Ectasis Sebæna	
Erica Sebæna aurántia Andr. Heath.	
2 fúscá.	
3 lùtea.	
4 minor.	

Erica assurgens.	
barbata Andr. Heath.	
2 major.	
Bonplandiana Bot. Cab.	
? caltra Bot. Cab.	
2 spicata.	
cerinthoides Bot. Mag.	
1 major H. Wob.	
2 minor H. Wob.	
nana H. Wob.	
decora Andr. Heath.	
depressa Andr. Heath.	
mollissima.	
Persóluta.	
2 álba.	
3 rùbra.	
propendens Andr. Heath.	
pubescens.	
1 major H. Wob.	
2 minor H. Wob.	
3 verna H. Wob.	













quadriflora.	
reflexa.	
2 rùbra.	
rùbens.	
pedunculáris Sal.	
viscaria Hort. Kew. Icon.	
cylindrica.	
divaricata Lodd.	
gélida Bot. Cab.	
incarnata Andr. Heath.	
trivialis M. Nab.	

Eurylepis (Erica L.) álbens Bot. Mag. fig. 882. Ilalicacaba Andr. Heath.	
Eurystegia (Erica L.) triceps Bot. Cab. fig. 883.	
Gypsocalis (Erica L.) nudiflora Sm. Icon.	
Lophandra (Erica L.) cubica Andr. Heath. fig. 884.	
2 minor H. Wob.	
3 major Hort. Brit.	

Lámprotis (Erica L.) calycina Andr. Heath. fig. 885.	
2 major H. Wob.	
Páchysa (Erica L.) báccana Bot. Mag.	

Syringodeca (Erica L.) abiétina.	Page
? claviflora Don's Mill.	
colorans Bot. Reg.	
concinna Andr. Heath.	
discolor Andr. Heath.	
elata Andr. Heath.	
Ewersona Andr. Heath.	
2 glabra.	
3 speciosa.	
4 longiflora.	
5 pilosa.	
Linnaea na.	
S. linnaeoides Andr. Heath.	
S. perspicua Hort. Kew.	
2 superba.	
linnaeoides Andr. Heath.	
perspicua Wendl. Eric.	
2 nana.	
radiata Andr. Heath.	
2 discolor.	
simpliciflora Wendl. Eric.	
spicata Andr. Heath.	
2 pallida H. Wob.	
tubiflora Andr. Heath.	
vestita.	
1 álba.	
2 incarnata.	
3 purpurea.	
4 rosea.	
5 fulgida.	
6 coccinea.	
7 lutea.	
8 mutabilis.	
9 elegans.	

App. V. List of a few of the larger Specimens of exotic Heaths, cultivated in the Royal Botanic Garden, Edinburgh; with their Dimensions, &c., as taken from the Plants, 12th July, 1836. - - - - 1094

App. VI. Culture of the hardy and half-hardy Species and Varieties of Erica belonging to the Group Erica normales. - 1094	
§ II. ANDROMÉDEÆ. - 1105	
       	
   	

IV. ANDROMEDA L. n. 1077. 1105

THE ANDROMEDA.
Polifolia Buxbaum.
Andromeda sp. L.

1. polifolia L. n. Europe fig. 889. 1106
The Poly-leaved-Andromeda, or Moorwort.
Rholoëndron polifolium Scop.
Wild Rosemary, Poly Mountain, Marsh Cistus, Moorwort, Marsh Holy Rose, Andromède, Fr. and Ger.

Varieties n. - - - - 1106	
1 angustifolia Lodd. n. fig. 890.	
2 ericoides n.	
3 grandiflora Lodd. n. fig. 891.	
4 latifolia Lodd. n. fig. 892.	
5 minima n.	
6 revoluta Lodd. n. fig. 893.	
7 scotica n.	
8 stricta n.	

2. rosmarinifolia Pursh. n. Newfoundland 1106
The Rosemary-leaved Andromeda.
A. polyfolia Michx.

	Page		Page
V. CASSIOPE D. Don. α β	1077. 1107	<i>A. Leaves evergreen.</i>	
THE CASSIOPE.		1. ferruginea Nutt. α N. America	1109
<i>Andr6meda sp. L. et Pall.</i>		The rusty-looking Lyonia.	
1. hypnoides D. Don. α β N. Europe	fig. 894. 1107	<i>Andr6meda ferruginea</i> Walt.	
The Hypnum-like Cassiope.		<i>Andr6meda ferruginea</i> β fruticosa Michx.	
<i>Andr6meda hypnoides L. &c.</i>		2. rigida Nutt. α North America	1110
2. tetragona D. Don. α β N. America	fig. 895. 1107	The rigid-leaved Lyonia.	
The four-cornered-branched Cassiope.		<i>Andr6meda ferruginea</i> Willd.	
<i>Andr6meda tetragona L.</i>		<i>Andr6meda ferruginea</i> β arborescens Mx.	
App. i. Hardy Species of Cassiope not yet introduced.	- - 1107	<i>Andr6meda rigida</i> Pursh.	
C. lycopodioides D. Don.		3. marginata D. Don. α North America	fig. 902. 1110
<i>Andr6meda lycopodioides</i> Pall.		The marginated-leaved Lyonia.	
C. ericoides D. Don.		<i>Andr6meda marginata</i> Du Ham.	
<i>Andr6meda ericoides</i> Pall.		<i>Andr6meda coriacea</i> Willd.	
C. Red6wskii G. Don.		<i>Andr6meda lucida</i> Lam.	
<i>Andr6meda Red6wskii</i> Cham. et Schlecht.		<i>Andr6meda mariana</i> Jacq.	
C. Mertensiana G. Don.		Variety α - - - 1110	
<i>Andr6meda Mertensiana</i> Bongard.		2 rubra Lodd. α fig. 900.	
C. fastigiata D. Don.		<i>B. Leaves deciduous.</i>	
<i>Andr6meda fastigiata</i> Wall. Pl. Par.		4. mariana D. Don. α North America	fig. 903. 1110
<i>Andr6meda cupressiformis</i> Wall. MSS.		The Maryland Lyonia.	
VI. CASSANDRA D. Don. α	1077. 1108	<i>Andr6meda mariana L.</i>	
THE CASSANDRA.		Variety α - - - 1111	
<i>Anar6meda sp. L. and others.</i>		2 obl6nga Swt. α	
1. calyculata D. Don. α North America	fig. 896. 1108	5. racemosa D. Don. α N. America	1111
The calyculated Cassandra.		The racemose-flowered Lyonia.	
<i>Andr6meda calyculata L.</i>		<i>Andr6meda racemosa L.</i>	
Varieties α - - - 1108		<i>Andr6meda paniculata</i> Walt.	
1 ventricosa Sims. α		6. arborea D. Don. α N. America	1111
2 latifolia Lodd. α		The Tree Lyonia.	
3 nana Sims. α		<i>Andr6meda arborea L.</i>	
2. (c.) angustifolia G. Don. α N. Amer.	fig. 897. 1108	7. paniculata Nutt. α N. Am. f. 904. 1111	
The narrow-leaved Cassandra.		The panicle-flowered Lyonia.	
<i>Andr6meda calyculata</i> β angustifolia Alt.		<i>Andr6meda paniculata L.</i>	
<i>Andr6meda angustifolia</i> Pursh.		8. salicifolia Wats. α N. Am. f. 905. 1111	
<i>Andr6meda crispata</i> Desf. et Link.		The Willow-leaved Lyonia.	
VII. ZENOBIA D. Don. α β	1077. 1108	9. frondosa Nutt. α N. America	1112
THE ZENOBIA.		The branchy Lyonia.	
<i>Andr6meda sp. Michx.</i>		<i>Andr6meda frondosa</i> Pursh.	
1. speciosa D. Don. α β Carolina	fig. 898. 1109	10. multiflora Wats. α North America	fig. 906. 1112
The snowy flowered Zenobia.		The many-flowered Lyonia.	
<i>Andr6meda speciosa</i> Michx.		11. capreaefolia Wats. α North America	fig. 907. 1112
Varieties α β - - - 1109		The Goat-Willow-leaved Lyonia.	
2 nitida Pursh. α β fig. 899.		App. i. Doubtful Species of Lyonia not yet introduced.	- - 1112
3 pulverulenta Pursh. α β f. 900.		<i>L. rhomboidalis G. Don.</i>	
<i>Andr6meda pulverulenta</i> Bartr.		<i>Andr6meda rhomboidalis N. Du Ham.</i>	
<i>Andr6meda cassinefolia</i> β Vent.		IX. LEUCOTH6E D. Don. α β	1077. 1113
<i>Andr6meda speciosa</i> var. γ glauca Wats.		THE LEUCOTH6E.	
<i>Andr6meda dealbata</i> Lindl.		<i>Andr6meda sp. of authors previously.</i>	
<i>Andr6meda ovata</i> Soland MSS.		1. axillaris D. Don. α North America	fig. 908. 1113
VIII. LYONIA Nutt. α β γ δ ϵ ζ η	1077. 1109	The axillary-racemed Leucoth6e.	
THE LYONIA.		<i>Andr6meda axillaris</i> Solander.	
<i>Andr6meda sp. L. and various others.</i>			

- | | Page |
|------------------------------------------------------------|------------|
| Variety \blacksquare - - - | 1113 |
| 2 longifolia \blacksquare | |
| <i>Andrómèda longifolia</i> Pursh. | |
| <i>Andrómèda Walteri</i> Willd. | |
| 2. spinulosa G. Don. \blacksquare North America | 1113 |
| The spinulose-tooth-leaved Leucothoe. | |
| <i>Andrómèda spinulosa</i> Pursh. | |
| <i>Andrómèda Catesbaei</i> Walt. | |
| 3. acuminata G. Don. \blacksquare North America | 1113 |
| The acuminate-leaved Leucothoe. | |
| <i>Andrómèda acuminata</i> Ait. | |
| <i>Andrómèda lucida</i> Jacq. | |
| <i>Andrómèda populifolia</i> Lam. | |
| <i>Andrómèda reticulata</i> Walt. | |
| <i>Andrómèda formosissima</i> Bartr. | |
| <i>Andrómèda laurina</i> Michx. | |
| Pipe-stem-wood, Amer. | |
| 4. floribunda D. Don. \blacksquare North America | 1114 |
| The numerous-flowered Leucothoe. | |
| <i>Andrómèda floribunda</i> Lyons. | |
| 5. spicata G. Don. \blacksquare N. Am. f. 912. | 1114 |
| The spicate-racemed Leucothoe. | |
| <i>Andrómèda spicata</i> Wats. | |
| X. PIERIS D. Don. ♂ - 1077. | 1114 |
| THE PIERIS. | |
| <i>Andrómèda sp.</i> Wallich. | |
| 1. ovalifolia D. Don. ♂ Nepal f. 913. | 1115 |
| The oval-leaved Pieris. | |
| <i>Andrómèda ovalifolia</i> Wall. | |
| <i>Andrómèda copricida</i> Ham. MSS. | |
| App. i. Half-hardy Species of <i>Pieris</i> not yet | 1115 |
| introduced. - - - | |
| P. formosa D. Don. | |
| <i>Andrómèda formosa</i> Wall. | |
| P. lanceolata D. Don. | |
| <i>Andrómèda lanceolata</i> Wall. | |
| <i>Andrómèda squamulosa</i> D. Don. | |
| P. japonica D. Don. fig. 911. | |
| <i>Andrómèda japonica</i> Thunb. | |
| XI. PHYLLODOCE Sal. \blacksquare ♂ | 1077. 1115 |
| THE PHYLLODOCE. | |
| <i>Andrómèda sp.</i> L. | |
| <i>Menziesia sp.</i> Swartz and Smith. | |
| 1. taxifolia Sal. \blacksquare Europe f. 915. | 1115 |
| The Yew-leaved Phyllodoce. | |
| <i>Menziesia cærulca</i> Swz. | |
| <i>Andrómèda cærulca</i> L. | |
| <i>Andrómèda taxifolia</i> Pall. | |
| <i>Erica cærulca</i> Willd. | |
| 2. empetrifórmis D. Don. ♂ N. America | 1116 |
| The Empetrum-like Phyllodoce. | |
| <i>Menziesia empetrifórmis</i> Sm. | |
| XII. DABŒCIA D. Don. \blacksquare 1077. 1116 | |
| THE DABŒCIA. | |
| <i>Erica sp.</i> Lin. | |
| <i>Andrómèda sp.</i> Lin. | |
| <i>Menziesia sp.</i> Juss. | |
| 1. polifolia D. Don. \blacksquare Ireland | 1116 |
| The Poly-leaved Dabœcia. | |
| <i>Andrómèda Dabœcia</i> Lin. Syst. | |
| <i>Andrómèda Dabœcia</i> Lin. sp. | |
| <i>Menziesia Dabœcia</i> Dec. | |
| <i>Erica hibérnica</i> , &c., Ray. | |
| <i>Menziesia polifolia</i> Juss. | Page |
| <i>Vaccinium cantábriçum</i> Huds. | |
| Irish Whorts, Cantabrian Heath, Saint | |
| Dabocç's Heath. | |
| Variety \blacksquare - - - | 1116 |
| 2 flore albo Swt. \blacksquare | |
| XIII. ARBUTUS Camer. ♂ ♂ ♂ | 1077. 1117 |
| THE ARBUTUS, or Strawberry Tree. | |
| <i>Andràchne</i> Clus. | |
| <i>Arbutus sp.</i> L. | |
| <i>Arbousier</i> , Fr. | |
| <i>Sandbeere</i> , Ger. | |
| <i>Abbatro</i> , Ital. | |
| 1. U'nedo L. ♂ ♂ S. Europe f. 919. | 1117 |
| The Unedo Arbutus, or Strawberry Tree. | |
| <i>L'Arbousier commun</i> , <i>Arbousier des Py-</i> | |
| <i>renées</i> , <i>Fraisier en Arbre</i> , Fr. | |
| <i>Erdbeercartige Sandbeere</i> , Ger. | |
| <i>Komad</i> , Mod. Greek. | |
| Varieties. ♂ ♂ - - - | 1117 |
| 1 álbus Ait. ♂ ♂ | |
| 2 rüber Ait. ♂ ♂ | |
| 3 plenus Ait. ♂ | |
| 4 schizopétalus ♂ | |
| 5 integrifolius Sims. ♂ | |
| 6 crispus ♂ | |
| 7 salicifolius ♂ | |
| 2. híbrida Ker. ♂ ♂ Hybrid | 1119 |
| pl. 192. fig. 920. | |
| The hybrid Arbutus. | |
| A. <i>andrachnoides</i> Link. | |
| Variety ♂ ♂ - - - | 1120 |
| 2 Milleri ♂ ♂ | |
| A. <i>Milleri</i> Mayes. | |
| 3. Andrâchne L. ♂ Levant pl. 193. | 120 |
| The Andrachne Arbutus. | |
| A. <i>integrifolia</i> Lam. | |
| <i>Andràchne Theophrásti</i> Clus. | |
| <i>Andràchne</i> Park. Theatr. | |
| <i>Andrachnè</i> of Theophrastus. | |
| <i>Andrachta</i> , Mod. Greek. | |
| Varieties - - - | 1120 |
| 1. With serrated leaves <i>Tourn.</i> | |
| 2 With large oblong fruit, <i>Tourn.</i> | |
| 3. With large compressed fruit, | |
| <i>Tourn.</i> | |
| 2 serratifolia fig. 821 ♂ ♂ | |
| A. <i>serratifolia</i> Nois. | |
| 4. procera Douglas. ♂ N. America | 1121 |
| The tall Arbutus. | |
| 5. tomentosa Pursh. ♂ N. America | 1122 |
| The downy Arbutus. | |
| <i>Arctostâphylus tomentosa</i> Lindl. | |
| Variety ♂ - - - | 1122 |
| 2 nuda Hook. et Arn. ♂ | |
| 6. densiflora H. B. et Kunth. ♂ ♂ Mexico | 1122 |
| The densely-flowered Arbutus. | |
| App. i. Hardy Species of <i>Arbutus</i> not yet | 1122 |
| introduced. - - - | |
| A. <i>aurifolia</i> L. | |
| A. <i>Menziesii</i> Pursh. | |
| A. <i>cordifolia</i> | |
| <i>Arctostâphylus cordifolia</i> Lindl. | |
| A. <i>glauca</i> | |
| <i>Arctostâphylus glauca</i> Lingl. | |

App. ii. *Half-hardy Species of Arbutus.* 1122
A. canadensis Lam. ■ □ fig. 921.
A. petiolaris H. B. et Kunth.
A. furens Hook. et Arn.
A. xalapensis H. B. et Kunth.
A. mollis H. B. et Kunth.
A. ferruginea L.

XIV. ARCTOSTAPHYLOS Adans.

THE BEARBERRY.
Uva-ursi Dod.
Arbutus sp. L.

1. *Uva-ursi* Spreng. 2 Europe fig. 923. 1123

The common Bearberry.
Arbutus Uva-ursi L.
Arbutus buxifolia Stokes.
Uva-ursi buxifolia Sal.
Bear berries, Bear-orchard-berries, Eng.
Barentraube, Barenberrie, Ger.
Berenduif, Dutch.
La Basserole, Fr.
Uva d'Orzo, Ital.
Uva de Oso, Span.
Uva de Ursa, Port.
Uva-ursi, in works of most old Botanists.

2. *alpina* Spreng. 2 Europe - 1123
 The Alpine Bearberry.
Arbutus alpina L.

App. i. *Half-hardy Species of Arctostaphylos not yet introduced.* - - 1124

A. polifolia H. B. et Kunth.
Andromeda ledifolia Humb.
A. glaucescens H. B. et Kunth.
A. pungens H. B. et Kunth.
A. Hookeri G. Don.
Arbutus pungens Hook.

XV. PERNETTYA Gaud. ■ □ 2 1078. 1124

THE PERNETTYA.

1. *mucronata* Gaud. 2 Terra del Fuego fig. 924. 1124
 The mucronate-leaved Pernettya.
Arbutus mucronata L.

2. *pilosa* G. Don. Mexico - - 1124
 The pilose, or hairy, Pernettya.
Arbutus pilosa Grah.

App. i. *Hardy Species of Pernettya not yet introduced.* - - 1125

P. microphylla Gaud.
Arbutus microphylla Forst.
Arbutus serpyllifolia Lam.
P. Myrsinites G. Don.
Andromeda Myrsinites Lam.

App. ii. *Half-hardy Species of Pernettya.* 1125

P. empetrifolia Gaud.
Arbutus empetrifolia Lindl.
Arbutus punila Willd.
Andromeda empetrifolia Lam.

P. punila Gaud. ■ □
Arbutus punila Forst.
P. Cavaniillesiana V. Don.
Andromeda prostrata Cav.
P. purpurea D. Don.
P. ciliaris D. Don.

XVI. GAULTHERIA L. ■ 2 1078. 1125

THE GAULTHERIA.

1. *procumbens* L. 2 North America fig. 925. 1125

The procumbent Gaultheria.
Partridge Berry, Mountain Tea, Spring Winter Green, Sin.

2. *Shallon* Pursh. 2 N. Am. f. 926. 1126
 The Shallon Gaultheria.

App. i. *Half-hardy Species of Gaultheria.* 1126

G. fragrantissima D. Don. ■ □
G. Fragrans D. Don.
Arbutus laurifolia Hamil.

App. ii. *Hardy and half-hardy Species of Gaultheria, not yet introduced.* - 1126

G. nummularioides D. Don.
G. ciliata Cham. et Schlecht.
 Several other Species

XVII. EPIGÆA L. ■ 2 - 1078. 1126

THE EPIGÆA.

Mimocyrtum Michx.
May Flower, Nova Scotia.

1. *repens* L. 2 North America - 1127
 The creeping Epigæa.

XVIII. PHALEROCA'RPUS G. Don. ■ 2 - - 1078. 1127

THE PHALEROCARPUS.

Vaccinium L.
Gaultheria Pursh.
Oryzococcus Nutt.
Arbutus Lam.

1. *serpyllifolia* G. Don. 2 N. Amer. 1127
 The Wild-Thyme-leaved Phalerocarpus.

Vaccinium hispidulum L.
Gaultheria serpyllifolia Pursh.
Arbutus filiformis Lam.
Oryzococcus hispidulus Pers.

XIX. CLETHRA L. ■ □ ■ 2 1078. 1127

THE CLETHRA.

Cuellaria R. et P.

1. *alnifolia* L. 2 N. Amer. f. 927. 1128
 The Alder-leaved Clethra.

C. alnifolia var. *a. denudata* Ait.

2. (*a.*) *tomentosa* Lam. 2 N. America fig. 928, 929. 1128

The downy Clethra.
C. alnifolia 2 *pubescens* Ait.
C. incana Pers.

3. (*a.*) *paniculata* Ait. 2 N. Amer. 1128
 The panicled-flowered Clethra.

4. (*a.*) *acuminata* Michx. 2 N. America 1128

The acuminated-leaved Clethra.
C. montana Bartr.

5. (*a.*) *scabra* Pers. 2 North America 1128
 The rough-leaved Clethra.

App. i. *Half-hardy Species of Clethra.* 1129

C. arborea Ait. ■ □ fig. 930.
C. ferruginea R. et P. ■ □
C. tinifolia Swartz. ■ □
Tiara occidentalis L.
 other species of Clethra

App. I. *Half-hardy Genera belonging to the Section Erica and § Andromedæ of the Order Ericaceæ.* - - - 1129

Agarista buxifolia G. Don. ■ □
Andromeda buxifolia Lam.

17. nudiflòrum Torr. (*Azàlea nudiflòra*
L.) ♀ N. America fig. 943. 1140

The naked-flowered Azalea.

Azalea nudiflora L.

Azalea periclymenoides Michx.

The American Honey-suckle.

May Flowers; Wild, or upright, Honey-suckle, Amer.

Varieties and Hybrids - 1141

1 coccineum *D. Don.* ♀

Azalea n. coccinea Sims.

2 rutilans *D. Don.* ♀

Azalea n. rutilans Ait.

Azalea periclymenoides rutilans
Pursh.

3 carneum *D. Don.* ♀

Azalea n. carneum Ait.

Azalea p. carneum Pursh.

4 album *D. Don.* ♀

Azalea n. alba Ait.

Azalea p. alba Ait.

5 papilionaceum *D. Don.* ♀

Azalea p. papilionacea Pursh.

6 partitum *D. Don.* ♀

Azalea p. partita Pursh.

7 polyandrum *D. Don.* ♀

Azalea p. polyandra Pursh.

8 Govenianum *D. Don.* ♀ f. 944.

9 rubrum *Lodd.* ♀

10 eximium *D. Don.* ♀

Varieties and Hybrids, according to
Lodd. Cat.

A. v.	2 alba et rubra	- 1142
	3 amara	- 1142
	4 blanda	- 1142
	5 carnea	- 1142
	6 caroliniana	- 1142
	7 Coburgia	- 1142
	8 colorata	- 1142
	9 conspicua	- 1142
	10 crispata	- 1142
	11 cumula	- 1142
	12 discolor	- 1142
	13 fastigiata	- 1142
	14 flore pleno	- 1142
	15 florida	- 1142
	16 globosa	- 1142
	17 grandiflora	- 1142
	18 incana	- 1142
	19 incarnata	- 1142
	20 mirabilis	- 1142
	21 montana	- 1142
	22 ochroleuca	- 1142
	23 pallida	- 1142
	24 pallidosa	- 1142
	25 papilionacea	- 1142
	26 periclymenoides	- 1142
	27 purpurea	- 1142
	28 purpurea	- 1142
	29 rosea	fig. 945. 1142
	30 ruberrima	- 1142
	31 rubicunda	- 1142
	32 rubra	- 1142
	33 rufa	- 1142
	34 rutilans	- 1142
	35 serotina	- 1142
	36 staminea	- 1142
	37 stellata	- 1142
	38 tricolor	- 1142
	39 varia	- 1142
	40 variabilis	- 1142
	41 variegata	- 1142
	42 versicolor	- 1142
	43 violacea	- 1142

18. bicolor *G. Don.* (*A. (n.) bicolor*
Pursh.) ♀ Carolina - 1142

The two-coloured-flowered Azalea.

Azalea bicolor Pursh.

Azalea n. bicolor Ait.

19. calendulaceum Torr. (*A. (n.) calen-*
dulacea Michx.) ♀ Carolina

fig. 946. 1142

The Marigold-flowered Azalea.

Azalea calendulacea Michx.

Azalea nudiflora var. *coccinea* Ait.

Varieties ♀ - 1142

2 Morterii *Swt.* ♀

3 fulgidum *Hook.* ♀

A. c. fulgida Hort.

20. canescens *G. Don.* (*A. (n.) canes-*
cens Michx.) ♀ Carolina - 1143

The canescent Azalea.

Azalea canescens Michx.

21. viscosum Torr. (*A. viscosa* *L.*) ♀

North America - fig. 947. 1143

The clammy-flowered Azalea.

Azalea viscosa L.

Varieties ♀ - 1143

2 ornatum *Swt.* ♀

Varieties and Hybrids, according to

Lodd. Cat.

A. Varieties.

A. v.	2 alba	- 1143
	3 crispata	- 1143
	4 dealbata	- 1143
	5 penicillata	- 1143
	6 praecox	- 1143
	7 pubescens	- 1143
	8 variegata	- 1143
	9 vittata	- 1143
	10 violae odore	- 1143

B. Hybridæ attackerensés. Hybrids
raised at High Clerc.

11 amara	- 1143
12 actinata	- 1143
13 aurora	- 1143
14 basilissa	- 1143
15 calodendron	- 1143
16 calocoryphe	- 1143
17 Caronia	- 1143
18 charissia	- 1143
19 coccinea nobilis	- 1143
20 eudæmon	- 1143
21 euryepes	- 1143
22 Govenia	- 1143
23 Herbertiana	- 1143
24 imperatrix	- 1143
25 inclata	- 1143
26 jasminodora	- 1143
27 lepida	- 1143
28 ochroleuca	- 1143
29 poikila	- 1143
30 poetica <i>Howard</i>	- 1143
hexapla.	
31 pulchella	- 1143
32 regalis	- 1143
33 rugens	- 1143
34 thysaliflora	- 1143

C. Hybridæ belgicæ. Hybrids raised
in Belgium.

35 Agate	- 1143
36 albo pleno	- 1143
37 amabilis	- 1143
38 amarantina	- 1143
39 amœnissima	- 1143
40 ardens	- 1143
41 atro-rubens	- 1143
42 aurantia maxima	- 1143
43 blanda	- 1143
44 calendulacea globosa	- 1143
insignis.	
nova.	
supërba.	
45 cædon	- 1143
46 coccinea maxima	- 1143
speciosa.	
47 concinna	- 1143
48 coruscans	- 1143
49 crocea	- 1143
globosa.	
50 cruenta	- 1143
51 cuprea	- 1143
alba.	
elegans.	
eximia.	
globosa.	
rubra.	
splendens.	
52 decorata	- 1143
53 decus hortorum	- 1143
54 dulcedo	- 1143
55 electa	- 1143
56 elegantissima	- 1143
57 eximula	- 1143
58 Ferrocii	- 1143
59 flamboyante	- 1143

	Page
A. v. 60 fulgida	- 1145
61 fulva	- 1143
62 gloria mundi máxima. minor.	- 1143
65 Gulielmus primus	- 1145
64 hybrida coccifera coccinea. nivea.	- 1145
65 incarnata máxima rubra.	- 1143
66 lépida	- 1145
67 lutea rubicúnda	- 1143
68 magnífica	- 1143
69 naritima incarnata	- 1143
70 miniata	- 1143
71 núbilis	- 1144
72 mixta triúmphans	- 1144
75 miniata	- 1144
74 Morterii	- 1144
75 ne plus últra	- 1144
76 nobilis	- 1144
77 nóritas antilléscens	- 1144
78 nóritata	- 1144
79 óptima	- 1144
80 ornata pállida	- 1144
81 picturata	- 1144
82 póntica glóbosa konink. tricolor var.	- 1144
85 præstantíssima	- 1144
84 pulchella	- 1144
85 punicea	- 1144
86 repui	- 1144
87 regina bélgica	- 1144
88 re-tantíssima	- 1144
89 rigida incarnata	- 1144
90 robústa	- 1144
91 rubra aurántia fulva.	- 1144
92 rubricata	- 1144
93 sanguinea	- 1144
91 Saturni	- 1144
95 severa	- 1144
96 speciosa	- 1144
97 speciosíssima	- 1144
98 splendens	- 1144
99 splendida	- 1144
100 sulphurea	- 1144
101 superba	- 1144
102 tricolor Jacóbi Wolff.	- 1144
103 triúmphans	- 1144
104 variegata	- 1144
105 venusta	- 1144
106 venustíssima	- 1144
107 versicolor	- 1144
22. glaucum G. Don. (A. (v.) glauca Ph.) ☼ North America - - 1144 The glaucous-leaved Azalea. <i>Azalea glauca</i> Pursh. <i>Azalea viscosa</i> var. <i>floribunda</i> Ait.	
23. hispídum Torr. (A. (v.) hispida Ph.) ☼ North America fig. 948. 1144 The hispid Azalea. <i>Azalea hispida</i> Pursh. <i>Azalea viscosa</i> var. <i>glauca</i> Ait.	
24. nítidum Torr. (A. nitida Ph.) ☼ North America - fig. 949. 1144 The glossy-leaved Azalea. <i>Azalea nitida</i> Pursh.	
25. speciósium G. Don. (A. speciosa Willd.) ☼ N. Amer. f. 950. 1145 The showy Azalea. <i>Azalea speciosa</i> Willd. <i>Azalea coccinea</i> Lodd.	
26. arboréscens Torr. (A. arboréscens Pursh.) ☼ North America 1145 The arborescent Azalea. <i>Azalea arboréscens</i> Pursh.	
§ v. Rhodora D. Don. ♀ ♂ ☼ 1145	
27. Rhodora G. Don. (Rhodora canadensis L.) ☼ N. Amer. f. 951. 1145 The Canada Rhodora. <i>Rhodora canadensis</i> L.	

	Page
App. i. Half-hardy Species of <i>Rhododendron</i> (and <i>Azalea</i>). - - 1146	
§ i. Bodram. ♀ ♂ ☼	
A. Species already introduced into British Gardens.	
28. arbóreum Smith. ♀ ♂ Nepal fig. 951. 1146 The scarlet-flowered Tree Rhododendron. <i>R. purpuræum</i> Hamilt. MSS. <i>Varieties and Hybrids</i> ♀ ♂ 1146	
1 sanguineum Lindl. ♀ ♂	
2 roseum Swt. ♀ ♂	
3 niveum Swt. ♀ ♂	
<i>R. album</i> Hamilt. MSS.	
<i>R. a. fl. niveis</i> D. Don.	
<i>R. a. album</i> Wall.	
4 cinnamomeum ♀ ♂	
<i>R. cinnamomeum</i> Wall.	
5 venustum D. Don.	
<i>Other Varieties and Hybrids of R. arboreum and other half-hardy Species</i> - - 1147	
<i>R. a. altaclerense</i> , fig. 952.	
29. campanulatum D. Don. ☼ Nepal fig. 953. 1147 The bell-shaped-flowered Rhododendron.	
<i>R. barbátum</i> Wall. - - - - - 1148	
<i>R. zeylanicum</i> Lodd. - - - - - 1148	
<i>R. strictum</i> Lodd. - - - - - 1148	
B. Species not yet introduced.	
<i>R. formosum</i> Wall. - - - - - 1148	
<i>R. linearifolium</i> Poir. - - - - - 1148	
§ ii. Pogonanthum. ☼	
30. anthopogon D. Don. ☼ Nepal fig. 954. 1148 The bearded-flowered Rhododendron. <i>R. aromatiæum</i> Wall.	
§ iii. Tsutsutsi D. Don. ☼ ♂ ♂ ☼ ☼ ☼ ☼ ☼	
A. Indian half-hardy Azaleas in British Gardens.	
31. índicum Swt. (A. indica L.) ☼ ♂ China fig. 955. 1148 The Indian Azalea. <i>Azalea indica</i> L.	
<i>Varieties and Hybrids</i> ☼ ♂ 1149	
2 phœniceum Don's Mill. ☼ ♂	
<i>Azalea indica phœnicea</i> Swt.	
<i>Azalea ledifolia phœnicea</i> Hook.	
3 flore pleno ☼ ♂	
<i>Azalea indica fl. pleno</i> Hook.	
4 ledifolium fig. 956. ☼ ♂	
<i>R. ledifolium</i> Don's Mill.	
<i>A. i. alba</i> Lindl.	
<i>A. ledifolia</i> Hook.	
<i>R. leucanthum</i> Bunge.	
5 pulchrum Don's Mill. ☼ ♂	
<i>A. i. pulchra</i> Hort.	
<i>R. pulchrum</i> Swt.	
<i>R. i. var. y Smithii</i> Swt.	
6 ignescens Swt. ☼ ♂	
<i>A. i. ignescens</i> Hort.	
7 aurantiacum G. Don ☼ ♂	
<i>A. i. aurantiaca</i> Hort.	
8 luteum Swt. ☼ ♂	
<i>A. i. lutea</i> Hort.	
9 spatulatum Blum. ☼ ♂	
<i>A. i. spatulata</i> Hort.	
10 grandiflorum Blum. ☼ ♂	
<i>A. i. grandiflora</i> Hort.	
11 angustifolium Blum. ☼ ♂	
<i>A. i. angustifolia</i> Hort.	
12 floribundum Blum. ☼ ♂	
<i>A. i. floribunda</i> Hort.	
13 Danielsianum ☼ ♂	
<i>A. i. Danielsiana</i> Paxton.	
14 lateritium ☼ ♂	
<i>A. i. lateritia</i> Lindl.	
15 variegatum Blum. ☼ ♂	
<i>A. i. variegata</i> Lindl.	
16 speciósium D. Don. ☼ ♂	
<i>A. i. speciosa</i> Hort.	

	Page
32. <i>sinéuse</i> Swt. (<i>A. sinénsis</i> Lodd.)	1149
China - - - - - fig. 957.	
The Chinese <i>Azalea</i> .	
<i>Azalea sinensis</i> Lodd.	
<i>Varieties and Hybrids</i>	1149
✱ 2. <i>flavescens</i> Swt.	
<i>A. s. 2. flavescens</i> .	
3. <i>macranthum</i> Don's Mill.	
<i>Azalea macrantha</i> Bunge.	
<i>R. nitens</i> D. Don MSS.	
<i>R. reticulatum</i> D. Don MSS.	1150
<i>Azalea reticulata</i> Hort.	
<i>R. Farrera</i> Tate.	fig. 958. 1150
<i>Azalea Farrera</i> Hort.	
<i>R. decumbens</i> D. Don MSS.	1150
<i>Azalea decumbens</i> Hort.	
B. <i>Indian Azaleas not yet introduced</i> .	
<i>R. scabrum</i> Don's Mill.	1150
<i>Azalea scabra</i> .	
<i>R. maximon</i> Thunb.	
<i>R. mucronatum</i> G. Don.	1150
<i>Azalea mucronata</i> Blum.	
? <i>R. leifolium</i> (A. T. alba Lindl.).	
<i>R. Burmanni</i> G. Don.	1150
<i>Azalea rosmarinifolia</i> Burm.	
<i>R. mollis</i> G. Don.	1150
<i>Azalea mollis</i> Blum.	
<i>R. Loureiriana</i> G. Don.	1150
<i>Azalea punctata</i> Lour.	

§ iv. *Propagation and Culture of the half-hardy Species of Rhododéndron and Azalea.*

XXI. <i>KALMIA</i> L. 1078. 1151	
THE KALMIA.	
<i>American Laurel.</i>	
1. <i>latifolia</i> L. N. Amer. f. 959. 1151	
The broad-leaved Kalmia.	
<i>Mountain Laurel, Calico Bush, Calico Flower, Amer.</i>	
2. <i>angustifolia</i> L. N. Am. f. 960. 1152	
The narrow-leaved Kalmia.	
<i>Sheep Laurel, Amer.</i>	
<i>Variety</i> - - - - - 1152	
2 ovata Pursh.	
3. <i>glauca</i> Ait. N. Amer. f. 961. 1152	
The glaucous-leaved Kalmia.	
<i>K. polifolia</i> Wangh.	
<i>Variety</i> - - - - - 1152	
2 rosmarinifolia Pursh.	
4. <i>cuneata</i> Michx. N. America 1152	
The wedge-shaped-leaved Kalmia.	
5. <i>hirsuta</i> Walt. N. Am. f. 962. 1152	
The hairy Kalmia.	
<i>K. ciliata</i> Bartr.	

XXII. <i>MENZIESIA</i> Smith.	1078. 1152
THE MENZIESIA.	
1. <i>ferruginea</i> Sm. N. Am. f. 963. 1153	
The rusty-flowered Menziesia.	
<i>M. urecoloris</i> Salisb.	
2. <i>globularis</i> Salisb. N. America 1153	
The globular-flowered Menziesia.	
<i>M. Smithii</i> Michx.	
<i>Azalea pilosa</i> Lam.	
<i>M. pilosa</i> Pers.	

XXIII. <i>AZALEA</i> D. Don.	1078. 1153
THE AZALEA.	
<i>Azalea procumbens</i> L. and many authors.	
<i>Lobelia</i> Desf.	
<i>Chamaledon</i> Link.	

1. <i>procumbens</i> L. Britain and North America - - - - - fig. 964. 1154	
The procumbent Azalea.	

XXIV. <i>LEIOPHYLLUM</i> Pers.	1078. 1154
THE LEIOPHYLLUM.	
<i>Ammyrsine</i> Pursh.	
<i>Fischera Swartz.</i>	
<i>Ledum buxifolium</i> Berg.	

1. <i>thymifolium</i> Pers. North America	fig. 965. 1154
The Thyme-leaved Leiophyllum.	
<i>Ledum buxifolium</i> Berg.	
<i>Ledum thymifolium</i> Lam.	
<i>Ledum serpyllifolium</i> L'Herit.	
<i>Ammyrsine buxifolia</i> Pursh.	
<i>Sand Myrtle, New Jersey.</i>	

L. <i>prostratum</i>	1155
<i>Ammyrsine prostrata</i> Swt.	
<i>Ammyrsine Lyonii</i> Swt.	

XXV. <i>LEDUM</i> L. 1078. 1155	
THE LEDUM.	

1. <i>palustre</i> L. N. Amer. f. 966. 1155	
The Marsh Ledum.	
<i>Ledum silesiacum</i> Cms.	
<i>Rosmarinum sylvestre</i> Cam.	
<i>Variety</i> - - - - - 1155	
2 <i>decumbens</i> Ait.	

2. <i>latifolium</i> Ait. N. Am. f. 967. 1155	
The broad-leaved Ledum, or Labrador Tree.	
<i>L. grandlandicum</i> Retz.	
<i>L. palustre</i> Michx.	
<i>Labrador Tea, Amer</i>	

3. <i>canadense</i> Lodd. N. Am. f. 968. 1156	
The Canadian Ledum.	
Sect. III. <i>VACCINIENÆ</i> D. Don. 1156	

XXVI. <i>VACCINIUM</i> L. 1078. 1156	
THE WHORTLEBERRY.	

<i>Vitis idæa</i> Tourn.	
<i>Airelle, Fr.</i>	
<i>Heidelbeere, Ger.</i>	
A. <i>Leaves deciduous.</i>	

1. <i>Myrtillus</i> L. Europe fig. 969. 1156	
The Little-Myrtle-like Whortleberry, or common Bilberry, or Blacberry.	
<i>Variety</i> - - - - - 1157	
2 <i>baccis albis</i>	

2. <i>uliginosum</i> L. Europe f. 970. 1157	
The bog Whortleberry, or great Bilberry.	
<i>Myrtillus grandis</i> Bauh.	

3. <i>angustifolium</i> Ait. N. Amer. 1158	
The narrow-leaved Whortleberry.	
<i>V. Myrtillus</i> Michx.	

4. <i>caespitosum</i> Michx. N. Amer. 1158	
The tufted Whortleberry.	

5. <i>galèzans</i> Michx. N. America 1158	
The Gale-like Whortleberry.	
<i>V. galiformis</i> Smith.	

6. <i>tenullum</i> Ait. N. Amer. f. 971. 1158	
The delicate Whortleberry.	
<i>V. pennsylvanicum</i> Lam.	
<i>Variety</i> - - - - - 1159	

7. <i>ligustrinum</i> Michx. N. Amer. 1159	
The Privet-like Whortleberry.	

	Page		Page
8. pallidum Ait. ✱ North America	1159	22. (?A.) padifolium Sm. ✱ Madeira	1164
The pale-flowered Whortleberry.		fig. 985, 986.	
9. arboreum Marsh. ✱ N. America	1159	The Bird-cherry-leaved Bear's Grape Whortleberry.	
The Tree Whortleberry.		V. Arctostaphylos Andr.	
V. diffusum Ait.		V. madagascensis Link.	
10. stamineum L. ✱ N. Am. f. 972.	1159	V. caucasicum Hort.	
The long-stamened Whortleberry.		V. padifolium caucasicum Hort.	
V. album Pursh.		B. Laves evergreen.	
V. clavatum Banks.		23. caracasenum H. B. et Kunth. ✱ □	1164
Variety ✱ - - -	1160	South America - - -	
2 album H. B. et Kunth. ✱		The Caracas Whortleberry.	
11. dumosum Ait. ✱ N. Am. f. 973.	1160	24. Vitis idæa L. ✱ Europe f. 987.	1164
The bushy Whortleberry.		The Mount Ida Whortleberry, or Cowberry.	
V. frondosum Michx.		Vitis idæa rubra Cam.	
V. hirtellum Ait.		The red Whortleberry.	
Variety ✱ - - -	1160	25. (V.) buxifolium Salisb. ✱ N. America	1165
2 humile Wats. ✱		fig. 988, 989.	
12. corymbosum L. ✱ North America	1160	The Box-leaved Whortleberry.	
fig. 974, 975.		V. brachycerum Michx.	
The corymbose-flowered Whortleberry.		26. myrtifolium Michx. ✱ N. Am.	1166
V. amœnum Ait.		The Myrtle-leaved Whortleberry.	
V. disomorphum Michx.		27. nitidum Andr. ✱ N. Am. f. 990.	1166
V. clavatum Hort.		The glossy-leaved Whortleberry.	
V. album Lam.		28. crassifolium Andr. ✱ N. America	1166
Varieties ✱ - - -	1161	The thick-leaved Whortleberry.	
2 virgatum Ait. ✱ fig. 976.		29. ovatum Pursh. ✱ N. Am. f. 991.	1166
3 fuscatum Ait. ✱ fig. 977.		The ovate-leaved Whortleberry.	
V. formosum Andr.		30. canadense Richards. ✱ N. Am.	1166
V. virgatum Wats., not of Ait.		The Canada Whortleberry.	
4 angustifolium. ✱		31. Myrsinites Michx. ✱ N. America	1167
V. virgatum var. angustifolium Wats.		The Myrsine-like Whortleberry.	
13. albiflorum Hook. ✱ N. Amer.	1161	Varieties ✱ - - -	1166
The white-flowered Whortleberry.		2 lanceolatus Pursh. ✱	
V. album Lam. ?		3 obtusus Pursh. ✱	
14. mariannum Wats. ✱ North America	1162	32. humiflorum Grab. ✱ N. America	1167
fig. 978.		The trailing Whortleberry.	
The Maryland Whortleberry.		App. i. Hardy Species of Vaccinium not yet	
V. marilandicum Lodd.		introduced. - - -	1167
15. grandiflorum Wats. ✱ North America	1162	V. Chamissonis Bongaro.	
fig. 979.		V. Myrtillus Cham.	
The great-flowered Whortleberry.		V. ovalifolium Smith.	
16. elongatum Wats. ✱ North America	1162	V. parvifolium Smith.	
fig. 980.		V. salicinum Cham.	
The elongated Whortleberry.		V. cylindraceum Smith.	
17. minutiflorum Wats. ✱ N. America	1162	V. confertum H. B. et Kunth.	
fig. 981.		V. obtusum Pursh.	
The minute-flowered Whortleberry.		App. ii. Half-hardy Species of Vaccinium	
18. glabrum Wats. ✱ N. Am. f. 982.	1162	not yet introduced. - - -	1167
The glabrous Whortleberry.		V. geminiflorum H. B. et Kunth.	
19. frondosum L. ✱ N. America	1163	V. hirtum Thunb.	
The frondose Whortleberry.		V. calcicum Smith.	
V. glaucum Michx.		V. bracteatum Thunb.	
Blue Tangles, Amer.		V. ciliatum.	
Variety ✱ - - -	1163	V. ledifolium Pohl.	
2 venustum Ait.		V. floribundum H. B. et Kunth.	
V. frondosum var. β lanceolatum		V. leucanthum Cham.	
Pursh.		V. villosum Smith.	
20. resinosum Ait. ✱ N. Am. f. 983.	1163	V. Schlechtensilii G. Don.	
The resinous Whortleberry.		V. scabrum Pohl.	
Andromeda baccata Wang.		V. montanum Pohl.	
Varieties ✱ - - -	1163	V. cereum Forst.	
2 rubescens Pursh. ✱		V. reticulatum Smith.	
3 lutescens Pursh. ✱		V. penduliflorum Goud.	
21. Arctostaphylos L. ✱ Black Sea	1163	V. microphyllum Herb. Reinw.	
fig. 984.		V. dentatum Smith.	
The Bear's Grape Whortleberry.		V. empetrifolium H. B. et Kunth.	
		V. penaeodes H. B. et Kunth.	
		V. acuminatum H. B. et Kunth.	
		V. alaternoides H. B. et Kunth.	
		XXVII. OXYCOCUS Pers. ✱ ✱	1078, 1168
		THE CRANBERRY.	
		Vaccinium sp. of L. and others.	

- Page
1. palustris Pers. ♀ Europe f. 992. 1168
The Marsh, or common, Cranberry.
O. vulgaris Pursh.
O. europæus Nutt.
Vaccinium Oxycoecus L.
Vaccinium Oxycoecus var. *α ovalifolius*
Michx.
Vaccinia palustris Ger. Emac.
Oxycoecum Cord.
Mossberries, Moorberries, Fenberries,
Marshworts, Whortleberries, Corn-
berries, Eng.
Airelle cancheberge, Fr.
Gemeine Moussebeere, Ger.
2. macrocarpus Pursh, ♀ North America
fig. 993. 1170
The large-fruited, or American, Cranberry.
Vaccinium macrocarpum Ait.
Vaccinium hispidulum Wagh.
Vaccinium Oxycoecus β oblongifolius Mx.
Variety ♀ - - - 1170
2 foliis variegatis Hort. ♀
Vaccinium macrocarpum fol. var.
Lodd.
3. erectus Pursh, ♀ N. Am. f. 994. 1171
The erect Cranberry.
Vaccinium erythrocarpum Mx.
- App. I. Genera of Ericaceæ, of which it ap-
pears doubtful if any hardy ligneous Species
have yet been introduced. - - - 1171
1. BRYANTHUS Gmel. ♀ - - - 1171
THE BRYANTHUS.
Andromeda sp. L.
Menziesia Swartz and Pursh.
Erica sp. Thunb.
1. Gmelini D. Don. ♀ Kamtschatka
fig. 995. 1171
Gmelin's Bryanthus.
Menziesia bryanthi Swartz.
Andromeda bryanthi L.
Erica bryanthi Thunb.
Bryanthus repens serpyllifolia flore rubro Gmel.
2. Stelleri D. Don. ♀ North-west America
fig. 996. 1172
Steller's Bryanthus.
Andromeda Stelleriana Pall.
Menziesia empetrifloris Pursh, but not of others.
Cladothamnus pyrolaeflorus Bongard. - 1172
Pyrola frutescens E.-Schischoltz.
- App. II. Half-hardy ligneous Species of
Ericaceæ. - - - 1172
1. ENKIANTHUS Lour. ♀ - - - 1172
THE ENKIANTHUS.
Melodora Sal.
1. quinqueflorus Lour. ♀ - - - 1172
The five-flowered Enkianthus.
E. reticulatus Lindl.
- E. biflorus Lour. ♀ - - - 1173
Virena - 1173
Hefaria Humb. et Bonp. ♀ - - - 1173
Hefaria Mutis.
H. glauca Humb. et Bonp. ♀ - - - 1173
H. racemosa Vent. Cole. ♀ - - - 1173
H. paniculata Michx.
Hymenanthus japonica Blum. - - - 1173
Gastussarcia baxifolia H. B. et Kunth - - - 1173
Thibaudia cordifolia H. B. et Kunth - - - 1173
Cavendishia nobilis Lindl. - - - 1173
Agapetes D. Don - - - 1173
Ceratostema grandiflora R. et P. - - - 1173
- App. III. Of the Cultivation of the Hardy
Ericaceæ, including the Laying out and
Planting of an Ericacetum. - - - 1173

- Page
Symplococææ. 1186
☉ ☐ ☉ ☐
Symplocos sinica Ker. ☉ ☐ China 2 sp. fig. 1007.
S. eratagoides Hamilt. ☉ ☐ Nepal.
- Styracææ.* 1187
1. STYRAX L. ☉ - - - 4 sp. 1187
THE STYRAX.
Alibocifer, Fr.
Storax, Ger.
1. officinale L. ☉ Syria fig. 1008. 1187
The official Storax.
Lagonomia, Modern Greek.
Sturax kulamitès, Ancient Greek.
2. grandifolium Ait. ☉ North America
fig. 1009. 1188
The large-leaved Storax.
S. officinale Walt.
S. grandiflorum Michx.
3. lævigatum Ait. ☉ South Carolina
fig. 1010. 1188
The smooth-leaved Storax.
S. octandrum L'Hérit.
S. globrum Cav.
S. læve Walt.
S. americanum Lam.
4. pulverulentum Michx. ☉ Virginia and
Carolina - - - fig. 1011. 1189
The powdery Storax.
S. lævigatum Curt.
- Halesiæcææ.* ☉ ☉ 1189
1. HALE'SIA Ellis. ☉ ☉ - 3 sp. 1189
THE HALE'SIA, or Snowdrop Tree.
Halesia, Fr. and Ger.
1. tetraptera L. ☉ South Carolina
pl. 194, 195. fig. 1012. 1190
The four-winged-fruited Halesia, or common
Snowdrop Tree.
The Snowdrop Tree, Silver Bell Tree, Amer.
2. (t.) parviflora Michx. ☉ ☉ Florida
fig. 1013. 1190
The small-flowered Halesia.
3. diptera L. ☉ ☉ Georgia and Carolina
fig. 1014. 1191
The two-winged-fruited Halesia.
- Sapotæcææ.* 1191
☉ ☉ ☉ ☐ ☉ ☉
1. ARGANIA Ræm. et Schultes. ☉ ☉
1191
THE ARGANIA.
Sideroxylon spinosum L.
L'Argan, Fr.
Eisenholz, Ger.
1. Sideroxylon Ræm. et Schultes. ☉ ☉
Morocco - - - fig. 1015. 1192
The Iron-wood Argania.
Sideroxylon spinosum L. Sp. &c.
Eleocharon Argan Retz.
Rhæmnus pentaphyllus Jacq.
Rhæmnus siculus L. Syst.

II. *BUMELIA* Swartz. Page
1192

THE BUMELIA.

Achras sp. Lln.
Sideroxylon sp. Lam. and others.
Chrysophyllum sp. Aubl. and others.
Hochstaum, Ger.

1. *lycioides* Gaertn. Carolina
fig. 1016. 1193

The Box-thorn-like Bumelia.
Sideroxylon lycioides Du Ham.
S. laeve Walt.
Lycioides sp. Lln. Hort Cliff.

2. *reclinata* Vent. Georgia - - 1193
The reclinate-branched Bumelia.

Sideroxylon reclinatum Michx.

3. *ténax* Willd. Carolina f. 1017. 1193
The tough-branched Bumelia.

B. chrysophylloides Pursh.
Sideroxylon tenax L.
S. striccum Walt.
S. chrysophyllodes Michx.
Chrysophyllum carolinense Jacq.
C. glabrum Juss.

4. *lanuginosa* Pursh. Carolina and
Georgia - - - 1194
The woolly-leaved Bumelia.

Sideroxylon lanuginosum Michx.
S. tenax Walt.

5. *oblongifolia* Nutt. N. America 1194
The oblong-leaved Bumelia.

B. salicifolia Swz.
Sideroxylon salicifolium Lam.
Achras salicifolia L.

Ebenaceæ. 1194

I. *DIOSPYROS* L. 1194

THE DATE PLUM.

Ebenus Comm.
Guaiaçana Tourn.
Plaqueminier, Fr.
Dattelpflaume, Ger.

1. *Lotus* L. Caucasus pl. 196, 197. 1194
The European Lotus, or common Date Plum.

Pseudolotus Matth.
Guaiaçana patavina Tourn.
Italian Lignum Vitæ, Wood of Life, Pock-
wood, Bastard Menynwood, Gerard.
Date of Trebisonde.
Plaqueminier, *Faux Lotier*, Fr
Italianische Dattelpflaume, Ger.

2. *virginiana* L. North America
pl. 198, 199. 1195

The Virginian Date Plum, or *Persimon*.
Guaiaçana Catsb.

Variety - - - 1196
2 *dulcis* Prince's Cat.
Foreman's sweet Persimon.

3. (v.) *pubescens* Pursh. N. Am. 1196
The downy-leaved Virginian Date Plum.

D. virginiana var. Michx.

App. I. *Other Species of Ebenaceæ.* 1197

Diospyrus angustifolia Lodd. Cat.
D. fertilis Lodd. Cat.
D. lucida Lodd. Cat. North America.
D. Mabotta Roxb. Philip Islands.

Embryopteris Kaki L. Japan. Page
D. chinensis Blume.
Konis, *Kaki*, *Kæmpf.*

Oleaceæ. - 1197

Sect. I. *OLEINEÆ.* 1197, 1198

I. *LIGUSTRUM* Tourn. 1198

THE PRIVET.

Troëne, Fr.
Rainucide, Ger.

1. *vulgare* Trag. Britain
fig. 1019, 1020. 1198

The common Privet.

L. germânicum Bauh.
Prim, *Prim-privet*.
Puine blanc, Fr.
Gemcine Rainucide, Ger.
Ligustrum Olivella, Ital.

Varieties - - 1199

2 *leucocarpum*

The white-berried Privet.

3 *xanthocarpum*

The yellow-berried Privet.

4 *chlorocarpum*

The green-berried Privet.

5 *sempervirens* fig. 1018.

The *Italian*, or evergreen, Privet.
L. italicum Mill.

6 *variegatum*

The variegated-leaved Privet.

7 *angustifolium*

The narrow-leaved Privet.

2. *spicatum* Hamilt. Nepal
fig. 1022. 1201

The spiked-flowered Privet.

L. nepalense Wall.
L. lanceolatum Herb. Lamb.
L. nepalense var. *glabrum* Hook.

3. *lucidum* Ait. fig. 1024, 1025. 1201
The shining-leaved Privet, or Wax Tree.

Variety - - - 1201

2 *floribundum* *Donald's* Cat.

App. i. *Species of Ligustrum not yet introduced.* - - 1202

L. sinense Lour. China.
L. japonicum Thunb. Japan.
L. latifolium Vitm.
L. pubescens Wall. Burmese Empire.
L. bracteatum D. Don. Nepal.
L. japonicum Hamilt.
Phillyrea bracteolata Herb. Lamb.

II. *PHILLYREA* Tourn. 1198, 1203

THE PHILLYREA.

Filaria, Fr.
Steindinde, Ger.

1. *angustifolia* L. Italy and Spain
fig. 1025. 1203

The narrow-leaved Phillyrea.

P. obliqua Tenore Syll.
P. media Tenore Fl. Neap.

Varieties - - - 1204

2 *lancolata* Ait.

3 *rosmarinifolia* Ait. fig. 1026.

4 *brachiata* Ait.

2. *média L.* ■ S. Europe fig. 1027. 1204
 The intermediate, or lance-leaved, Phillyrea.
P. latifolia var. média Lapeyr.
P. ligustrifolia Mill.
P. lævis Tenore Syll.
P. latifolia var. A ligustrifolia Poll.

Varieties ■ - - - 1204
 2 *virgata* Ait. ■
 3 *buxifolia* Ait. ■

3. (m.) *ligustrifolia Ait.* ■ S. Europe 1204
 The Privet-leaved Phillyrea.

4. (m.) *péndula Ait.* ■ S. of Europe 1204
 The drooping-branch'd Phillyrea.
P. média y Willd.

5. (m.) *oleafolia Ait.* ■ S. of Europe 1204
 The Olive-leaved Phillyrea.
P. média & Ait.
P. racemosa Link.

6. *latifolia L.* ■ S. Eur. fig. 1028. 1205
 The broad-leaved Phillyrea.
P. latifolia β serrata Poll.
P. latifolia β Ten. Fl. Neap.
P. spinosa Ten. Syll.
P. latifolia β spinosa Seg.

7. (l.) *lævis Ait.* ■ South of Europe and North of Africa - - 1205
 The smooth Phillyrea.
P. latifolia var. A. Willd. Sp.
P. latifolia Mill. Dict.

8. (l.) *obliqua Ait.* ■ South Europe 1205
 The oblique-leaved Phillyrea.
P. latifolia y Willd. Sp.
P. filarea Link.
Phillyrea v. Clus.

9. (l.) *spinosa Mill.* ■ S. Europe 1205
 The spiny, or Holly-leaved, Phillyrea.
P. ilicifolia Willd. Enum.
P. latifolia β spinosa Willd. Sp.
P. latifolia longifolia Link.
Phillyrea l. Clus.

III. *CHIONANTHUS L.* ☿ ♀ 1098. 1205

THE SNOW-FLOWER, or Fringe Tree.
Chionanthe, Fr.
Schneebäume, Ger.

1. *virginica L.* ☿ North America fig. 1029, 1030. 1206
 The Virginian Snow-Flower.
Snowdrop Tree, Amer.
Arbe de Neige, Fr.
Schneebäume, Ger.

Varieties ■ - - - 1206
 2 *latifolia* Catesh. ♀
C. v. montana Pursh.
 3 *angustifolia Ait.* ♀
C. trifida Mœnch.
 4 *maritima Pursh.* ♀
C. maritima Lodd.

App. I. Half-hardy lignous Species of *Olea*-
cea belonging to the Section *Oleina*. 1207

- Olea europæa L.* ♀
O. Oleaster Hoffm. & Segg.
O. europæa communis Ait.
O. sylvestris Mill.
O. e. sativa ♀ fig. 1051.
O. sativa Hoffm. & Segg.
O. europæa Michx.
Subvarieties.
O. e. s. longifolia Ait. fig. 1052
latifolia.
hispanica Mill.

- oblōnga.*
Picholine, Fr.
Pignolo, Ital.
ferruginea Ait. fig. 1033.
obliqua Ait.
buxifolia Ait.

Subarcticis most esteemed in France.

1. *Olivier pleureur.*
O'lea craniomirpha N. Du Ham.
 2. *Olivier à Fruit arrondi.*
O'lea apiculata N. Du Ham.
 3. *Olivier de Lacques.*
O'lea minor lucida N. Du Ham.
 4. *Olivier à petit Fruit rond, N. Du Ham.*
 5. *Olivier de Salon, N. Du Ham.*
 6. *Olivier amygdalin.*
O'lea amygdalina.
 7. *Olive Picholine.*
O'lea oblonga N. Du Ham.
Pignolo, Ital.

- O. excelsa Ait.* ♀
O. americana L. ♀ fig. 1051. 1208
O. fragrans Thunb. ♀ fig. 1055. 1208
O. capensis L. ♀
O. buxifolia Mill. - 1208

Sect. II. *SYRINGEE* ♀ - 1198. 1208

IV. *SYRINGA L.* ♀ - 1198. 1208
 THE LILAC.

Lilac Tourn.
Lilas, Fr.
Flüder, Ger.

1. *vulgàris L.* ♀ Persia fig. 1036. 1209
 The common Lilac.
Lilac vulgàris Gartn.
Pipe Privet, Pipe Tree.
Lilas commun, Fr.
Gemeiner Flüder, Ger.

Varieties ■ - - - 1209

- 1 *cærulea* Clus. ♀
 The common blue Lilac.
 2 *riolacea Curt.* ♀
 The common purple Lilac.
Scotch Lilac.
 3 *alba* ♀
 The common white Lilac.
 4 *alba major Lodd. Cat.* ♀
 5 *alba plena* ♀
S. plena Lodd. Cat.
 6 *rubra Lodd. Cat.* ♀
 7 *rubra major Lodd.* ♀
 Other Varieties.

2. *Josik'a Jacq.* ♀ Transylvania fig. 1037, 1038. 1201
Josika's Lilac.

3. *pèrsica L.* ♀ Persia fig. 1039. 1211
 The Persian Lilac.
Lilac minor Mœnch.
Lilac pèrsica Lam.
Lilas de Perse, Fr.

Varieties ■ - - - 1211

- 2 *alba Lodd. Cat.* ♀
 The white Persian Lilac.
 3 *laciniata Lodd. Cat.* ♀ fig. 1040.
 The cut-leaved Persian Lilac.
S. capitata Gmel.
Lilas à Feuilles de Persil, Fr.
 4 *salvifolia Lodd. Cat.* ♀


4. *rothomagènsis Renault.* ♀ Rouen fig. 1041. 1212

The Rouen Lilac.
S. dubia Pers.
Lilacum rothomagense Renault
S. média Dum. Cours.
Lilas Varin, N. Du Ham.
S. chinensis Willd.
S. sibirica Hort.
The Siberian Lilac, Hort.

	Page
Varieties ♀ - - -	1212
2 Lilas Royal, <i>Bou. Jard.</i> ♀	
3 saugeàna <i>Hort.</i> ♀	
<i>Lilas saugé</i> , Fr.	
? <i>S. coccinea</i> Lodd. Cat.	
<i>S. chinensis rubra</i> Lodd. Cat.	
App. i. <i>Species of Syringa not yet introduced.</i> - - -	1212
<i>S. Emòdi</i> Wall. fig. 1012.	
<i>S. villosa</i> Vahl.	
? <i>Ligustrum sinense</i> Lour.	
V. FONTANESIA Labill. ♀ †	
THE FONTANESIA. 1198. 1213	
1. phillyreoides Labill. ♀ † Syria	
fig. 1034. 1213	
The Phillyrea-like Fontanesia.	
<i>Forsythia suspensa</i> Vahl. - - -	1213
<i>Syringa suspensa</i> Thunb.	
<i>Lilac perpersè</i> Lam.	
Sect. III. FRAXINIEÆ. † ♀ 1198. 1213	
VI. FRAXINUS Tourn. † ♀ 1198. 1213	
THE ASH.	
<i>Frêne</i> , Fr.	
<i>Esche</i> , Ger.	
A. <i>Leaflets broad, smooth or shining on the upper surface.</i> <i>Natives of Europe.</i>	
1. excelsior L. † Europe	
pl. 200, 201, 202. fig. 1044. 1215	
The taller, or common, Ash.	
<i>F. apétala</i> Lam.	
<i>F. rostrata</i> Guss.	
<i>F. O'rnus</i> Scop.	
<i>F. erosa</i> Pers.	
<i>F. crispa</i> Bosc.	
<i>Le Frêne</i> , Fr.	
<i>Aesche, Esche</i> , Ger. and Dutch.	
<i>Ask</i> , Dan. and Swed.	
<i>Frassino</i> , Ital.	
<i>Fresno</i> , Span.	
<i>Freixo</i> , Port.	
<i>Jas. Jasen. Jasscn.</i> , Russ.	
<i>Æsc</i> , Sax.	
Varieties † - - -	1214
2 pëndula Ait. pl. 203.	
The pendulous, or weeping, Ash.	
pëndula var. †	
<i>The Cowpen Ash.</i>	
3 Kincáirniae †	
The Kincairney Ash.	
4 aúrea Willd. †	
The golden-barked Ash.	
<i>F. aúrea</i> Pers.	
5 aúrea pëndula †	
6 crispa †	
<i>F. crispa</i> Bosc.	
<i>F. atrovirens</i> Desf.	
7 jaspídea Willd. †	
The striped-barked Ash.	
8 purpuráscens Descemet. †	
The purple-barked Ash.	
9 argéntea Desf. †	
The silver-striped-leaved Ash.	
10 lútea †	
The yellow-edge-leafteted Ash.	
11 erosa Pers. †	
12 horizontális Desf. †	
The horizontal-branched Ash.	
13 verrucósa Desf. †	
The warted-barked Ash.	

	Page
14 verrucósa pëndula †	
15 nána Lodd. Cat. †	
The dwarf Ash.	
<i>F. c. hùnilis</i> Hort.	
16 fungósa Lodd. Cat. †	
The fungous-barked Ash.	
17 verticilláta Lodd. Cat. †	
The whorled-leaved Ash.	
18 villósa nóva Descemet. †	
Other Varieties.	
2. (e.) heterophýlla Vahl. † Europe	
pl. 204. 1228	
The various-leaved Ash.	
<i>F. simplicifolia</i> Willd. and Hort.	
<i>F. monophýlla</i> Desf.	
<i>F. c. β diversifolia</i> Ait.	
<i>F. c. var. ε</i> Lam.	
<i>F. c. β heterophýlla</i> Dec.	
<i>F. integrifolia</i> and <i>diversifolia</i> Hort.	
Variety - - -	1229
2 variegáta fig. 1051.	
3. (e.) angustifolia Bauh. † Eur. 1229	
The narrow-leaved Ash.	
Other Sorts of the common European Ash 1229	
B. <i>Leaflets small, smooth or shining above. Natives of the South of Europe, the North of Africa, or the West of Asia.</i>	
4. (e.) parvifolia Willd. † South of Europe, North of Africa, West of Asia	
pl. 205. fig. 1052. 1229	
The small-leaved Ash.	
5. (e. p.) argéntea Lois. † South of Europe, North of Africa, and West of Asia	
- - - - 1230	
The silvery-leaved Ash.	
6. (e. p.) oxycárpa Willd. † South of Europe, North of Africa, and West of Asia	
- - - fig. 1053. 1230	
The sharp-fruited Ash.	
<i>F. oxyphylla</i> Bieb.	
<i>F. O'rnus</i> Pall.	
7. (e.) pállida Bosc. † South of Europe, N. of Africa, and W. of Asia	
1230	
The pale-barked Ash.	
8. lentiscifolia Desf. † South of Europe, North of Africa, and West of Asia	
pl. 206. fig. 1054. 1231	
The Lentiscus-leaved Ash.	
<i>F. tamariscifolia</i> Vahl.	
<i>F. parvifolia</i> Lam.	
<i>F. aleppensis</i> Pluk.	
Variety - - -	1231
2 pëndula	
C. <i>Leaves and Leaflets large, glaucous and downy beneath. Natives exclusively of North America.</i>	
9. americana Willd. † North America	
pl. 207. fig. 1055. 1232	
The American Ash.	
<i>F. acumináta</i> Lam.	
<i>F. discolor</i> Muhl.	
White Ash, Green Ash, Amer.	
Variety † - - -	1232
2 latifolia †	


- | | Page | | Page |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 10. (a.) pubescens <i>Walt.</i> \checkmark N. America
fig. 1056. 1232, 1233 | 1232, 1233 | 22. (a.) viridis <i>Bosc.</i> \checkmark N. Amer.
The green Ash. | 1239 |
| The downy Ash.
<i>F. nigra</i> Du Roi.
<i>F. tomentosa</i> Michx.
<i>Red Ash, Black Ash, Amer.</i>
<i>Varieties</i> \checkmark - - - 1233 | | 23. (a.) cinerea <i>Bosc.</i> N. America
The grey Ash. | 1239 |
| 2 longifolia <i>Willd.</i> \checkmark
<i>F. pennsylvanica</i> Marsh. | | 24. (a.) alba <i>Bosc.</i> \checkmark N. America
The white Ash.
<i>F. caroliniana alba</i> ,
<i>F. americana alba</i> var.,
<i>F. am. alba affinis</i> Hort. Soc. Gard. | 1239 |
| 3 latifolia <i>Willd.</i> \checkmark | | 25. (a.) Richardi <i>Bosc.</i> \checkmark N. Am.
Richard's Ash. | 1239 |
| 4 subpubescens <i>Pers.</i> \checkmark
<i>F. submillosa</i> Bosc. | | 26. (a.) ovata <i>Bosc.</i> \checkmark N. America
The ovate-leaved Ash. | 1239 |
| 11. (a.) sambucifolia <i>Vahl.</i> \checkmark N. America
fig. 1057, 1058. 1234 | 1234 | 27. (a.) nigra <i>Bosc.</i> \checkmark N. America
The black Ash. | 1239 |
| The Elder-leaved Ash.
<i>F. nigra</i> Moench.
<i>F. crispa</i> Hort.
<i>Black Ash, Water Ash, Amer.</i>
<i>Variety</i> \checkmark - - - 1234 | | 28. (a.) elliptica <i>Bosc.</i> \checkmark N. Amer.
The elliptic-leaved Ash. | 1240 |
| 2 crispa <i>Lodd. Cat.</i> \checkmark | | 29. (a.) fusca <i>Bosc.</i> \checkmark N. America
The brown-branched Ash. | 1240 |
| 12. (a.) quadrangulata <i>Michx.</i> \checkmark North
America - fig. 1059, 1060. 1235 | 1235 | 30. (a.) rufa <i>Bosc.</i> \checkmark North
The rufous-haired Ash. | 1240 |
| The quadrangular-branched Ash.
<i>F. tetragona</i> Cels.
<i>F. quadrangularis</i> Lodd. Cat.
<i>Blue Ash, Amer.</i>
<i>Variety</i> \checkmark - - - 1235 | | 31. (a.) pannosa <i>Vent. et Bosc.</i> \checkmark North
America - - - pl. 210. 1240
The cloth-like-leaved Ash. | 1240 |
| 2 nervosa <i>Lodd. Catt.</i> \checkmark | | 32. Boscii <i>G. Don.</i> \checkmark N. America
Bosc's Ash.
<i>F. nana</i> Bosc, not of Willd. | 1240 |
| 13. (a.) juglandifolia <i>Lam.</i> \checkmark N. America
pl. 208, f. 1061, 1062. 1232. 1236 | 1236 | 33. (a.) polemoniifolia <i>Poir.</i> \checkmark North
America - - - 1240
The Greek-Valerian-leaved Ash.
<i>F. nana</i> Desf.
<i>F. nana (appendiculata)</i> Pers. | 1240 |
| The Walnut-leaved Ash.
<i>F. viridis</i> Michx.
<i>F. concolor</i> Mühl.
<i>The green Ash, Amer.</i>
<i>Western black Ash, Pursh.</i>
<i>Variety</i> \checkmark - - - 1236 | | 34. (a.) triptera <i>Nutt.</i> \checkmark N. Amer.
The three-winged-fruited Ash. | 1240 |
| 2 subintegerrima <i>Vahl.</i> \checkmark
<i>F. juglandifolia</i> β <i>subseerrata</i>
<i>Willd.</i>
<i>F. caroliniana</i> Wengenh.
<i>F. Nove-Anglie</i> and <i>F. caroliniana</i> Mill. Dict. | | 35. chinensis <i>Roxb.</i> \checkmark N. America
The China Ash. | 1240 |
| 14. (a.) caroliniana <i>Lam.</i> \checkmark N. America
1232. 1237 | 1237 | 36. Schiedeana <i>Schlecht.</i> \checkmark N. Amer.
Schiede's Ash.
<i>Fagara dubia</i> Benth. et Schult. | 1241 |
| The Carolina Ash.
<i>F. cecelsior</i> Walt.
<i>F. serratifolia</i> Michx.
<i>F. lanceolata</i> Borkh. | | VII. O'RNUS <i>Pers.</i> \checkmark 1198. 1241
THE FLOWERING ASH.
<i>Fraxinus</i> sp. of the older authors.
<i>Le Frêne à Fleurs.</i> Fr.
<i>Die blühende Esche.</i> Ger.
<i>Oren.</i> Hebrew.
<i>Orcinē Mētia.</i> Greek. | |
| 15. (a.) epiptera <i>Vahl.</i> \checkmark N. America
pl. 209. fig. 1232. 1237. | 1237 | 1. europæa <i>Pers.</i> \checkmark South of Europe
pl. 211, 212. 1241 | 1241 |
| The wing-topped, seeded, or two-coloured, Ash.
<i>F. canadensis</i> Gartin.
<i>F. linca</i> Bosc. | | The European Flowering, or <i>Manna</i> , Ash.
<i>F. O'rnus</i> L.
<i>F. O'rnus</i> and <i>F. paniculata</i> Mill. Dict.
<i>F. florifera</i> Scop.
<i>F. hotryoides</i> Mar.
<i>F. vulgarior</i> Segn.
<i>Varieties</i> - - - 1241 | |
| 16. (a.) platycarpa <i>Vahl.</i> \checkmark N. America
fig. 1063, 1064. 1238 | 1238 | rotundifolia.
<i>O. rotundifolia.</i>
americana.
<i>O. americana.</i>
globifera <i>Lodd.</i> | |
| The broad-fruited Ash.
<i>F. caroliniana</i> Catesb.
<i>The Carolina Ash, Amer.</i> | | 2. (c.) rotundifolia <i>Pers.</i> \checkmark Calabria
fig. 1069. 1244 | 1244 |
| 17. (a.) expansa <i>Willd.</i> \checkmark N. Am. 1238
The expanded Ash.
<i>F. aroliniana</i> Hort. Worlitz. | 1238 | The round-leafleted Flowering Ash.
<i>Fraxinus rotundifolia</i> Ait.
<i>F. mannifera</i> Hort. | |
| 18. (a.) mixta <i>Bosc.</i> \checkmark N. America 1238
The mixed Ash. | 1238 | | |
| 19. (a.) pulverulenta <i>Bosc.</i> \checkmark N. Am. 1238
The powdery Ash. | 1238 | | |
| 20. (a.) rubicunda <i>Bosc.</i> \checkmark N. Amer. 1239
The reddish-veined Ash. | 1239 | | |
| 21. (a.) longifolia <i>Bosc.</i> \checkmark N. Amer. 1239
The long-leaved Ash. | 1239 | | |

P. laevigata Ait.   Canary Islands
P. panicifolia Cav. - 1258

Half-hardy Species of Periploca. - 1258

Bignoniaceæ. 1258


I. BIGNONIA Tourn.  - 1258







THE TRUMPET FLOWER.

Bignonia sp. L. and others.

Bignone, Fr.


Trompctenblume, Ger.

1. *capreolata* L.  N. Am. f. 1090. 1259
The tendriled Bignonia.

II. *TECOMA* Juss.    
  - 1258, 1259

THE TECOMA.

Bignonia sp. L. and others.

1. *radicans* Juss.  Carolina f. 1091. 1259
The rooting-branched Tecoma.

Bignonia radicans L.

Bignonia radicans major Hort.

Gelsemium Clematis Barrel.


Bignonia fraxinifolia Catesb.



Jasmin de Virginie, Fr.

Wurzel Bignonia, Ger.

Essenblättrige Bignonia, Dutch.

Variety  - - - 1259

2 major Hort. 

2. *grandiflora* Swt.   China and Japan
fig. 1092. 1260

The great-flowered Tecoma.

Bignonia grandiflora Thunb.


Bignonia chinensis Lam.


Kjotsjo Kämpf.

Incarnilla grandiflora Spreng.


Tung-von-fa, Chinese.

App. i. *Half-hardy Species of Bignonia and Tecoma.* - 1261


Bignonia crucigera Plum.  Virginia.

Tecoma australis R. Br.  New Holland. fig. 1095.

Bignonia Pandoræ Vent.

T. capensis Lindl.  Cape of Good Hope. fig. 1094.

Bignonia capensis Thunb.


III. *CATALPA* Juss.  - 1258, 1261

THE CATALPA.

Bignonia sp. of L. and others.

Bignone Catalpa, Fr.

Gemeine Trompctenblume, Ger.

1. *springata* Sims.  North America
pl. 213, 214. 1261

The Lilac-like leaved Catalpa.

Bignonia Catalpa Lin.

Catalpa bignonioides Walt.

C. cordifolia Nutt.

Bois Shavanon, Catalpa de l'Amérique,

Fr.


Trompctenbaum, Ger.

Catalpa-boom, Dutch.



App. I. *Of the half-hardy ligneous Plants of the Order Bignoniaceæ.* - 1263


Eccremocarpus longiflorus Humb.  Peru. fig. 1095.

E. virolis R. et P. Peru.


Catalpa discolor D. Don.  figs. 1096, 1097.


C. venosa R. et P.

Cobæacææ.   1264

Cobæa scandens Cav.   fig. 1098.

Convolvulacææ. 1364

Convolvulus Dorstenium L.  Levant. fig. 1100.


C. Cneorum L.  Spain, &c. fig. 1099.

C. scoparius L.  Canaries.

C. floridus L.  Canaries.


Boraginacææ. 1265


     

Lithospermum frutescens L.  S. of Europe.


L. frutescens majus Lehmann. Naples. fig. 1101.


L. rosmarinifolium Tenore.


L. prostratum Lois.  France.

Echium giganteum L.  Canaries.



E. cándicans L.  Madeira. fig. 1102.

Heliotropium peruvianum L.  Peru.











H. p. hybridum  Hybrid.







H. corymbosum R. et P.  Peru.

Cordiaceæ. 1265

Ehretia serrata Rozb.   E. Indies and China. fig. 1105.

Solanacææ. 1266

I. SOLANUM Pliny.    
  - - - 1266

THE NIGHTSHADE.

Melongèna Tourn.


Pseudo-Capsicum Mœnch.

Nycteterium Vent.

Aquiritia Jacq.

Morelle, Fr.

Nachtschatcn, Ger.

1. *Dulcamàra* L.  Britain f. 1104. 1266
The Bitter-sweet, or woody, Nightshade.

S. scandens Neck.

Dulcamàra flexuosa Mœnch.

S. scandens seu Dulcamàra Tourn.


Amara dulcis Gerard.


Dulcis Amàra Trag.


Glycypicros seu Dulcamàra Bauh.


La Morelle grimpante Renault.


Varieties  - - - 1267


1 violacea Hort. 


2 álba L. 

3 cárnea Cels. 



4 plèna Tourn. 

5 variegata Munt. 


6 hirsùta Don's Mill. 

7 rupèstris Schmidt. 

2. *suffruticosum* Schousb.  Barbary 1267
The suffruticeo-Nightshade.

3. *crispum* R. & S.   Chiloe
fig. 1105. 1267.

The curled-leaved Solanum.

4. *bonariense* L.  Buenos Ayres
fig. 1106. 1268

The Buenos Ayres Nightshade

App. i. *Half-hardy ligneous or fruticose Species of Solanum.* - 1268

- Solanum Baibisi Donal. S. America.
 S. betaceum Car. S. America.
 S. angulatum R. et S. Lima.
 S. marginatum W. Africa.
 S. Pseudo-Capsicum L. Madeira.
Capsicum Animum Plinij Gerard.
 S. sodomense L. Africa.
 S. ligustrinum Lodd. Chili. fig. 1107.

II. *LY'CIUM L.* 1266. 1269

THE BOX THORN.
Jasminoides Niss.
Matrimony Vine, Amer.
Lycien, Fr.
Bocksdorn, Ger.

I. *europæum L.* South of Europe
 fig. 1108. 1269.

The European Box Thorn.
L. salicifolium Mill.
Jasminoides aculeatum Michx.
 Varieties - 1270
 Fruit yellow
 Fruit roundish

2. (e.) *bárbarum L.* S. of Europe, &c.

The Barbary Box Thorn.
L. latimifolium Mill.
L. bárbarum *et vulgare* Ait.
The Duke of Argyll's Tea Tree.

3. (e.) *chinense Mill.* China
 fig. 1110, 1111. 1271

The Chinese Box Thorn.
L. bárbarum *et chinense* Ait.
L. bárbarum Lour.
L. ovatum N. Du Ham.

4. (e.) *Trewianum G. Don.* China 1271

Trew's Box Thorn.
L. bárbarum Lam.
L. chinense N. Du Ham.

5. (e.) *ruthenicum Murr.* Siberia
 fig. 1112. 1271

The Russian Box Thorn.
L. tataricum Pall.
Lycien de la Russie, Fr.
 Variety - 1272
 2 *caespicum* Pall.

6. (e.) *lanceolatum Poir.* S. Europe 1272

The lanceolate-leaved Box Thorn.
L. europæum *et* Dec.

7. (e.) *turbinatum Du Ham.* China
 fig. 1115. 1272

The turbate-fruited Box Thorn.
L. latimifolium Mill.
L. bárbarum *et* Dec.

8. (?e.) *tetrandrum Thunb.* Cape of
 Good Hope - - 1272

The tetrandrous-flowered Box Thorn.

9. (?e.) *Sháwi Rœm.* Barbary - 1273

Shaw's Box Thorn.
L. europæum Mill.

10. *àfrum L.* Spain f. 1114, 1115. 1273

The African Box Thorn.

III. *CRABOWSKIA Schlecht.*
 1266. 1273

THE CRABOWSKIA.
Lycium sp. L.
Ehretia sp. L'Hérit.

1. *boerhaaviaefolia Schlecht.* South
 of Brazil! - fig. 1116. 1274

The Boerhaavia-leaved Crabowskia.
Lycium boerhaaviaefolium L.
Ehretia halimifolia L'Hérit.
Lycium heterophyllum Murr.
Jasminoides spinosum Du Ham.
Lycium paniculé, Fr.

App. I. *Half-hardy ligneous Plants belonging to the Order Solanaceæ.* - 1274

- Nicotiana glauca* Graham. Buenos Ayres. fig. 1118.
Brugmansia sanguinea R. et P. fig. 1117.
B. bicolor Pers.
 B. suaviscolens Willd. Peru. fig. 1120.
Datura arborea Hort.
Solantra grandiflora L. Jamaica.
Cestrum nocturnum L. E. Indies. fig. 1119.
C. Páruqi L. Chili. fig. 1122.
Vestia lycioides Willd. Chili. fig. 1121.
Cantua ligustrifolia Juss.

Scrophulariæcæ. 1276

I. *BU'DDLEA L.* 1276

THE BUDDLEA.

1. *globosa L.* Chili f. 1123, 1124. 1276

The globe-flowered Buddlea.
B. globiflora N. Du Ham.
B. capitata Jacq.
Pálquin Feuillée H.
Buddleia globuleux, Fr.
Kopftragende Budeje, Ger.

App. i. *Half-hardy Species of Buddlea.* 1277

- B. salvifolia* Lam. C. G. H.
Lantana salvifolia L.
B. paniculata Walt. Nepal.
B. saligna Willd. C. G. H.
B. crispa Royle. Himalayas.

App. I. *Half-hardy ligneous Plants of the Order Scrophulariæcæ.* - 1277

- Hallèria lœuca* L. C. G. H. fig. 1125.
Maurândya semperflorens Jacq. Mexico.
M. Barclayana Bot. Reg. Mexico.
Mimulus glutinosus Willd. California.
Anthoeris viscosa R. Br. New Holland.
Calceolaria integrifolia L. Chili.
C. rugosa Fl. Per. Chili.
C. sessilis Hort. figs. 1127, 1128.
Peróica decussata Ait. Falkland Islands. figs. 1129, 1130.
Celsia lanata Jacq. fig. 1236.
Capraria lanceolata L. C. G. H.
Fraulia salicyfolia Bot. Mag.
Alonsoa R. et P.
Angelonia H. B. et Kunth.
Lophospermum Don.
Rhodochiton Zucc.
Nycteria D. Don.

Labiæcæ. 1278

Satureja montana L. South of Europe
 fig. 1131. 1278

S. capitata Willd. Levant - 1278

Thymus vulgáris L. South of Europe
 fig. 1132. 1278

T. Mastichina L. Spain 1278

T. grandiflorus Hort.

	Page
<i>Hyssopus officinalis</i> L. α S. of Europe	1278
fig. 1133.	
<i>Teucrium angustifolium</i> Schreb. α Spain	1279
<i>T. fruticans</i> L. α Spain	1279
fig. 1135, 1136.	
<i>T. Marum</i> L. α South of Europe	1279
fig. 1134.	
<i>T. flavum</i> L. α South of Europe	1279
<i>T. Pòlium</i> L. α S. of Europe	1279
<i>T. corymbosum</i> R. Br. α Van Diemen's Land	1279
<i>Phlomis fruticosa</i> L. α Spain	1279
f. 1137.	
<i>P. purpurea</i> Sm. α South of Europe	1279
fig. 1138.	
<i>Rosmarinus officinalis</i> L. α South of Europe	1279
fig. 1139.	
<i>Stachys fruticulosa</i> Bieb. α Caucasus	1281
<i>S. stenophylla</i> Spr. α Spain	1281
<i>S. palestina</i> L. α Syria	1281
<i>S. lavandulifolia</i> Pers. α Levant	1281
<i>Lavandula Spica</i> L. α South of Europe	1281
fig. 1140.	
<i>L. latifolia</i> Ehrh. α S. of Europe	1281
<i>Acyros graveolens</i> Link. α Crimea	1282
<i>A. rotundifolia</i> Pers. α Spain	1282
<i>Gardoquia Hookeri</i> Benth. ? α South Carolina	1282
<i>Westringia rosmariniformis</i> Sm. α New South Wales	1282
<i>Salvia officinalis</i> L. α South of Europe	1282
fig. 1141.	
var. Leaves variegated α	
Whole plant of a reddish hue α	
Leaves larger than those of the species α	
fig. 1142	
<i>S. Hablitziana</i> L. α Siberia	1283
f. 1143.	
<i>S. pomifera</i> L. α Candia	1283
fig. 1144.	
<i>S. critica frutescens pomifera</i> Tourn.	
<i>Audibertia incana</i> Benth. α Colombia	1283
fig. 1145.	
App. I. <i>Half-hardy ligneous or suffruticose Species of Labiæcæ.</i>	1283
<i>Lavandula Stoechas</i> L. α S. of Europe	1149.
<i>L. dentata</i> L. α Spain	1146.
<i>L. pinnata</i> B. & M. α Madeira	1147.
<i>L. viridis</i> L'Herit. α Madeira.	
<i>Ph. crotanhus fruticosus</i> L'Herit. α C. G. H.	1148.
<i>Sideritis cædicans</i> Ait. α Mad. Ira.	
<i>Leonotis Leonurus</i> R. Br. α C. G. H.	
<i>Phlomis Leonurus</i> L.	
<i>Sphærole campanulata</i> Benth. α Chili	1151.
<i>S. Lindleyi</i> Benth. Valparaiso.	
<i>Dracocephalum canariense</i> Com. α Canaries.	
<i>Salvia splendens</i> Ker. α S. America.	
<i>S. farinosa</i> Willd. α S. America.	
<i>S. fulgens</i> Cur. α S. America.	
<i>S. Grahami</i> Benth. α S. America.	1151.
<i>S. chamædryoides</i> Cur. α Mexico.	
<i>S. aurea</i> L. α C. G. H.	
<i>Prasium majus</i> L. α Spain.	
<i>Prostanthera lasiantha</i> Tab. α N. S. Wales.	
Other half hardy species.	

	Page
<i>Verbenæcæ.</i>	1285
α α α α α α α α	
I. <i>VITEX</i> L. α α α α - - - 1285	
THE CHASTE TREE.	
<i>Gatilier</i> , Fr.	
<i>Keuschbaum</i> , Ger.	
I. <i>Agnus castus</i> L. α Sicily f. 1152. 1285	
The officinal, or true, Chaste Tree.	
<i>Elicognon Theophrasti</i> Lob.	
<i>Agnus castus</i> Blackw.	
<i>Arbreau Poivre, Poivre sauvage</i> , Fr.	
Variety α - - - 1286	
2 <i>latifolia</i> Mill. α South of France and Italy.	
App. i. <i>Half-hardy Species of Vitæ.</i>	1286
<i>V. incisa</i> Lam. α China.	
<i>V. Negundo</i> Bot. Mag.	
App. I. <i>Half-hardy Plants of the Order Verbenæcæ.</i>	1286
<i>Clerodendrum inerme</i> R. Br. α E. Indies.	1153.
<i>Folkameria incensis</i> L.	
<i>C. speciosissimum</i> Parton ? α Japan.	
<i>Doranta cænea</i> Hort. α S. America.	
<i>Aloysia citriodora</i> Or. α Chili	1286
fig. 1154.	
<i>Verbena triphylla</i> L'Herit.	
<i>Lippia citriodora</i> Kunth.	
—————	
<i>Myopórincæ.</i> α α 1287	
<i>Myoporum parvifolium</i> R. Br. α New Holland.	
<i>M. oppositifolium</i> R. Br. α New Holland.	
<i>M. diffusum</i> R. Br. α New Holland.	
<i>M. ascenden.</i> R. Br. α New Holland.	
—————	
<i>Globulariæcæ.</i> α α 1287	
<i>Globularia longifolia</i> L. α Madeira	1287
fig. 1155.	
<i>G. salicina</i> Lam.	
G. <i>Alypum</i> L. α S. of Europe	1287
var. <i>integrifolium</i> α α	
—————	
<i>Plumbaginæcæ.</i> 1287	
α α α α	
<i>Statice monopétala</i> L. α Sicily.	
<i>S. suffruticosa</i> L. α Siberia.	
<i>Plumbago capensis</i> Thunb. α Cape of Good Hope.	
—————	
<i>Chenopodiæcæ.</i> 1287	
α α α α α α α α α	
I. <i>CHENOPODIUM</i> L. α α 1288	
THE GOOSEFOOT.	
<i>Salsola</i> Sp.	
<i>Inserme</i> , Fr.	
<i>Gause Fuss</i> , Ger.	

1. fruticosum *Schrad.* ■ England Page
 fig. 1156, 1157. 1288
 The shrubby Goosefoot, or *Stoncrop Tree*.
Salsola fruticosa L.
The shrubby Glasswort.
Soude en Arbre, Fr.
Strauchartiges Salzkraut, Ger.
2. parvifolium *R. et S.* ■ Caucasus 1289
 The small-leaved Goosefoot.
C. fruticosum Bieb. Fl. Taur.
C. microphyllum Bieb. Suppl. to Fl. Taur.
Salsola fruticosa Bieb. Casp.
Salsola microphylla Pall.
3. hortense *R. et S.* ■ S. of Europe 1289
 The Garden Goosefoot.
Salsola hortensis Forsk.
Salsola divergens Poir.

II. ATRIPLEX L. ■ ■ 1288, 1289
 THE ORACHE.

1. Halimus L. ■ Spain fig. 1158. 1289
 The Halimus Orache, or *Tree Purslane*.
Halimus latifolius sive fruticosus Bauh.
Halimus i. Clus.
The broad-leaved Sea Purslane Tree.
Arroche, Fr.
Strauchartige Melde, Ger.
2. portulacoides L. ■ Britain f. 1159. 1290
 The Purslane-like, or *shrubby*, Orache, or *Sea Purslane*.
Halimus scandens Clus.
Halimus vulgaris Ger. Emac.
Halimus scu Portulaca marina Bauh.
A. maritima.
Halimus et Portulaca marina dicta angustifolia Ray.
The narrow-leaved Sea Purslane Tree.

III. DIOTIS Schreb. ■ 1288, 1290
 THE DIOTIS.

1. Ceratoides W. ■ Siberia f. 1160. 1291
 The two-horned-calyx'd Diotis.
Axyris Ceratoides L.
Ceratosperrum papposum Pers.
Axyris fruticosa, floribus foemineis lanatis
 Gmel.
Achyronthes papposa Forsk.
Krascheninnikovia Guildenst.
Urtica foliis lanceolatis, foeminis hirsutis,
 Roy.
Ceratoides orientalis fruticosa clavagni
folio Tourn.
Orientalisches Doppellohr, Ger.

App. I. Half-hardy Species of Chenopodiaceæ.
 1291

- Anabasis tamariscifolia L. ■ Spain.
 A. aphylla L. ■ Asia Minor.
Salsola articulata Forst.
 Kochia prostrata Schreb. ■ South of Europe.
Salsola prostrata L.
 Bösen Yeremora L. ■ Canaries.
 Camphorosma monspeliaca Schk. ■ South of Europe.
 Other Genera belonging to Chenopodiaceæ.

Polygonaceæ. 1292

■ ■ ■ ■ ■

I. TRAGOPYRUM Bieb. ■ ■ 1292
 THE GOAT WHEAT.
Polygonum L.

1. lanceolatum Bieb. ■ ■ Siberia Page
 fig. 1161. 1292
 The lanceolate-leaved Goat Wheat.
Polygonum frutescens Willd.
Strauchartiger Knöterig, Ger.
2. buxifolium Bieb. ■ Siberia f. 1162. 1293
 The Box-leaved Goat Wheat.
Polygonum crispulum var. α Sims.
P. caucasicum Hoffmannsegg.
3. polygamum Spr. ■ Carolina f. 1163. 1293
 The polygamous-seed'd Goat Wheat.
Polygonum polygamum Vent.
P. parvifolium Nutt.
- T. pängens Bieb.
 T. glaucum Spr.
 T. grandiflorum Bieb.

II. ATRAPHA'XIS L. ■ 1292. 1294
 THE ATRAPHA'XIS.

1. spinosa L. ■ Levant fig. 1164. 1294
 The spine-branched Atraphaxis.
Atriplex orientalis, frutex aculeatus, flore
pulehro Tourn.
2. undulata L. ■ Cape of Good Hope 1295
 The waved-leaved Atraphaxis.

III. CALLIGONUM L. ■ 1292. 1295
 THE CALLIGONUM.

1. Pallasia L'Hérit ■ Caspian Sea
 figs. 1165, 1166. 1295
 Pallas's Calligonum.
Pterococcus aphyllus Pall. Voy.
Calligonum polygonoides Pall. Itin.
Pallasia caspica L.
Pollisia Pterococcus Pall. Fl. Ross.
Cuspischer Hackenkraut.
- C. comosum L'Hérit. - - - - 1296
 C. Panderi L'Hérit. - - - - 1296

App. I. Half-hardy Species of Polygonaceæ.
 1296

- Brunnichia cirrhosa *Gertn.* ■ Carolina.
 Rumex Lunaria L. ■ Canaries. fig. 1167.
 Polygonum adpressum *R. Br.* ■ Van Diemen's Land.

Lauræce. ■ ■ ■ ■ ■ 1296

I. LAURUS Plin. - - 1296
 THE LAUREL, or Bay, Tree.

- Sassafras and Benzoin, C. G. Von
Esenbeck.
Daphnè, Greek.
 A. Plants evergreen; hardy.
1. nobilis L. ■ ■ S. Europe pl. 215. 1297
 The noble Laurel, or Sweet Bay.
Laurus Camer.
L. vulgaris Bauh.
Laurier commun, Laurier franc, Laurier
d'Apollon, Laurier à sauce, Fr.
Gemeine Lorbeer, Ger.

Varieties ■ - - 1297

- 2 undulata Mill. ■
 3 salicifolia Swt. ■
 L. n. angustifolia Lodd. Cat
 4 variegata Swt. ■
 L. n. fol. var. Lodd. Cat.
 5 latifolia Mill. ■
 6 crispa Lodd. Cat. ■
 7 flore pleno N. du Han. ■

	Page
B. <i>Plants evergreen; half-hardy.</i>	
2. <i>carolinensis</i> <i>Catesh.</i> † North America fig. 1168, 1169. 1299	
The Carolina Laurel, or Red Bay.	
L. <i>Borbônia</i> L. sp.	
L. <i>axillaris</i> Lam.	
L. <i>Borbônia</i> sp. Plum.	
Pérsca <i>Borbônia</i> Spreng.	
The broad-leaved Carolina Bay.	
Laurier rouge, Laurier Bourbon, Laurier de Caroline, Fr.	
Carolinischer Lorbeer, Rother Lorbeer, Ger.	
Varieties † - - 1299	
2 glabra Pursh. †	
3 pubescens Pursh. †	
4 obtusa Pursh. †	
3. <i>Cateshiána</i> Michx. † Georgia - 1300	
Catesby's Laurel.	
4. <i>aggregata</i> Sims. † China f. 1170. 1300	
The grouped-flowered Laurel.	
L. <i>foetens</i> Ait. † Madeira. - - - 1301	
L. <i>maileirensis</i> Lam.	
Pérsca <i>foetens</i> Spreng.	
L. <i>Mýrtila</i> Lour. China - - - 1301	
L. <i>indica</i> L. † India - - - 1301	
C. <i>Leaves deciduous.</i>	
5. <i>Sassafras</i> L. † North America pl. 216, 217. 1301	
The Sassafras Laurel, or Sassafras Tree.	
Cornus más odorata, folio trifido, margine plano, Sassafras dicta. Pluk.	
Sassafras arbor, ex Florida, ficulneo folio, Bauh.	
Sassafras sp. C. G. Nees Von Esenbeck.	
Pérsca Sassafras Spreng.	
Laurier Sassafras, Fr.	
Sassafras Lorbeer, Ger.	
Varieties † - - 1301	
The red †	
L. <i>subgenus Euósmus</i> Nutt.	
Sassafras L.	
The white †	
L. <i>e. álvida</i> Nutt.	
6. <i>Benzóin</i> L. † Virginia f. 1171. 1303	
The Benzoin Laurel, or Benjamin Tree.	
Arbor virginiana citrea vel limoni folio, benzoinum fructens, Comm.	
Laurus <i>æstivális</i> Wangh.	
Pseudo Benzoin Michx.	
Euósmus Benzoin Nutt.	
Benzoin sp. C. G. Nees Von Esenbeck.	
Spice Bush, Spice Wood, Wild Allspice, Amer.	
Laurier faux B'nzoin, Fr.	
Benzoin Lorbeer, Ger.	
7. (<i>B.</i>) <i>Diospýrus</i> Pers. † Virginia fig. 1172. 1304	
The Diospyrus-like Laurel.	
L. <i>Euósmus Diospýrus</i> Nutt.	
L. <i>diospyroides</i> Michx.	
? L. <i>melissafolia</i> Water.	
8. (<i>B.</i>) <i>æstivális</i> L. † Virginia 1304	
The summer Laurel, or Willow-leaved Bay.	
L. <i>cnérvia</i> Mill.	
L. <i>Euósmus æstivális</i> Nutt.	
Pond Bush Amer.	
Sommer Lorbeer, Ger.	
9. <i>geniculata</i> Michx. † Virginia fig. 1173. 1305	
The knee-flexed-branched Laurel.	
L. <i>Euósmus geniculata</i> Nutt.	
L. <i>æstivalis</i> Willd.	

	Page
App. I. <i>Half-hardy Species of Lauræcæ.</i> 1305	
<i>Cinnamótrum Címphora</i> Swt. † □ Japan. fig. 1174.	
The Camphor Tree.	
Laurus <i>Camphora</i> L.	
C. <i>verum</i> Swt. † □ Ceylon. fig. 1175.	
Laurus <i>Cinnamómu</i> L.	
Laurus <i>Cassia</i> Bot. Mag.	
C. <i>Cassia</i> D. Don. † □ Ceylon.	
The wild Cinnamon.	
Laurus <i>Cárta</i> L.	
Laurus <i>Cinnamómu</i> Bot. Rep.	
Pérsca <i>Cassia</i> Spr.	
Other liguosous plants belonging to this order.	

Protæcæ. † □ 1306

<i>Bánkeir litoralis</i> R. Br. † □ New Holland.	
B. <i>oblongifolia</i> Cur. † □ New South Wales.	
<i>Grevillea rosarinifolia</i> Cun. † □ N. South Wales. f. 1176.	
G. <i>acuminata</i> R. Br. † □ New South Wales. f. 1177, 1178.	
<i>Hakea acicularis</i> R. Br. † □ New South Wales.	
H. <i>suaveolens</i> R. Br. † □ New Holland.	
H. <i>pugionifórris</i> R. Br. † □ New South Wales. fig. 1179.	

Thymelæcæ. - 1306

I. DA'PHNE L. † † † † † † 1307	
THE DAPHNE.	
<i>Thymelæa</i> Tourn.	
A. <i>Leaves deciduous.</i>	
1. <i>Mezèreum</i> L. † N. of Eu. f. 1180. 1307	
The Mezereon Daphne, or common Mezereon.	
Spurge Olive, Spurge Flax; Flowering Spurge, Parkinson.	
Dwarf Bay Gerard.	
Lauréole femelle, Bois gentil, Mézèreon, Bois joli, Fr.	
Gemeiner Seidelbast, Kellerbaltz, Ger.	
Peperachtige Daphne, Dutch.	
Lauréola femina, Biondella, Camelia, Ital.	
Lauréola hembra, Span.	
Varieties † - - 1308	
2 flore álbo †	
3 autumnale †	
2. <i>altáica</i> Pall. † Siberia fig. 1181. 1308	
The Altaic Daphne.	
Daphné altáique, Lauréole de Tartarie, Fr.	
Sibirischer Seidelbast, Ger.	
3. <i>alpína</i> L. † Switzerland f. 1182. 1309	
The Alpine Daphne.	
The Alpine Chamelca, Marsh.	
Daphn' des Alpes, Fr.	
Alpen Seidelbast, Ger.	
B. <i>Erect. Leaves persistent. Flowers lateral.</i>	
4. <i>Lauréola</i> L. † Britain f. 1183. 1309	
The Laureola Daphne, or Spurge Laurel.	
Daphnoides verum, vel Lauréola, Gesn.	
Lauréola Ray.	
Thymelæa Lauréola Scop.	
The Evergreen Daphne.	
Lauréole mâle, Lauréole des Anglais, Fr.	
Immergrüner Seidelbast, Ger.	
5. <i>póntica</i> L. † Asia Minor f. 1184. 1310	
The Pontic Daphne, or twin-flowered Spurge Laurel.	
Thymelæa póntica citrei foliis Tourn.	
Lauréole du Levant, Fr.	
Pontischer Seidelbast, Ger.	
Varieties † - - 1310	
2 rubra Hort. †	
3 foliis variegatis Lodd. Cat. †	

6. *Thymelæa L.* ✨ Spain f. 1185. 1310
 The *Thymelæa*, or *Milkwort-like*, Daphne.
Thymelæa foliis polygalæ glabris Bauh.
T. alpina glabra, flosculis subluteis ad foliorum ortum scssilibus, Pluk.
Sanamunda viridis vel glabra Bauh. Prod.
Sanamunda glabra Bauh. Hist.
Passerina Thymelæa Dec.
The Wild Olive.
La Thymelic, Fr.
Astloser Seidelbast, Ger.

7. *Tárton ráira L.* ✨ South of France
 fig. 1186. 1311
 The *Tarton-raira*, or *silvery-leaved*, Daphne.
Thymelæa foliis candidantibus et serici instar mollibus Bauh. Pin.
Tartoo-Raire Gallo-provinciæ Mospeticusium Lob.
Sanamunda arcuata latifolia Barr.
Passerina Tárton-Raira Schrad.
The oval-leaved Daphne.
Lauréole blanche, Fr.
Silberblättriger Seidelbast, Ger.

8. (? *T.*) *pubescens L.* ✨ Austria - 1311
 The pubescent Daphne.
Thymelæa italica, Tarton-raire Gallo-provinciæ similis, sed per omnia major, Micheli.
Bchaarter Seidelbast, Ger.

9. (? *T.*) *tomentosa Lam.* ✨ Asia Minor 1311
 The tomentose Daphne.
Passerina villosa L.
Lauréole cotonneuse, Lam.

C. *Erect. Leaves persistent. Flowers terminal.*

10. *collina Sm.* ✨ S. Italy fig. 1187. 1311
 The hill-inhabiting Daphne, or *Neapolitan Mezereon*.
D. collina æ Bot. Rep.
 ? *D. buxifolia* Vahl.
Daphné des Collines, Lauréole à Feuilles de Santé, Fr.
Stumpfblättriger Seidelbast, Ger.

11. (c.) *neapolitana Lodd.* ✨ S. of Italy fig. 1188. 1312
 The Neapolitan Daphne.
D. collina ß *neapolitana* Lindl.

12. (c.) *oleoides L.* ✨ Crete f. 1189. 1312
 The Olive-like Daphne.
Chamædaphnoides cretica Alpin.
Thymelæa cretica oleæ folio utriusque glabro Tourn.
Daphne salicifolia Lam.
Lauréole à Feuilles d'Olivier, Fr.
Oelbaunblättriger Seidelbast, Ger.

13. (c.) *scricea Vahl.* ✨ Candia - 1312
 The silky-leaved Daphne.
Thymelæa cretica oleæ folio subtus villosa Tourn.
Daphne oleifolia Lam.
Seidenartiger Seidelbast, Ger.

14. *striata Trat.* ✨ Switzerland - 1313
 The striated-calyx'd Daphne.

D. *Erect. Leaves persistent. Flowers in Racemes.*

15. *Gnidium L.* ✨ Spain f. 1190. 1313
 The *Gnidium*, or *Flax-leaved*, Daphne.
Thymelæa foliis lini Bauh.
Spurge Flax, Mountain Widow Wayle.
Daphné Gnidium, Lauréole à Panicule, Fr.
Rispelblättriger Seidelbast, Ger.

E. *Prostrate. Leaves persistent. Flowers terminal aggregate.*

16. *Cneorum L.* ✨ Switzerland fig. 1191. 1313
 The *Garland-flower*, or *trailing*, Daphne.
Cneorum Matth.
Wohlricherender Seidelbast, Ger.
Varieties 2 - - - 1313
 2 *foliis variegatis* 2
 3 *floræ albo* 2

App. i. *Half-hardy Species of Daphne.* 1314

D. odora Thunb. ✨ [] China.
D. sinensis Lam.
The sweet-scented Daphne.
Lauréole de Chine, Daphné odorant, Fr.
Wohlricherender Seidelbast, Ger.
Var. 2 variegata Lodd. Cut. ✨ []
 3 *rubra* D. Don. ✨ [] fig. 1192.

D. hybrida Swt. ✨ [] Hybrid f. 1193.
D. dalphinia of Fr. Gardens.
D. dauphinoi of Eng. Gardens.

D. indica L. ✨ [] India.
D. papiræa Wal. ✨ [] Nepal.
D. canubina Wal.

II. *DIRCA L.* ✨ - - 1307. 1314
 THE *DIRCA*, or *Leather-wood*.
Thymelæa Gron.

1. *palustris L.* ✨ Virginia f. 1194. 1314
 The *Marsh Dirca*.
Moorwood.
Bois de Cuir, Bois de Plomb, Fr.
Sumpf Lederholz, Ger.

App. I. *Half-hardy ligneous Plants belonging to the Order Thymelæaceæ.* - 1315

Gnidia imbricata L. ✨ [] Cape of Good Hope.
G. dendata Bot. Reg.
Passerina filiformis L. ✨ [] Cape of Good Hope.
Pimelea drupacea Lab. ✨ [] New Holland.

Santalàceæ. † ✨ 1315

I. *NYSSA L.* † - - 1315, 1316
 THE *NYSSA*, or *Tupelo Tree*.

1. *biflora Michx.* † Virginia pl. 218. fig. 1195, 1196. 1317
 The twin-flowered Nyssa.
N. aquatica L. sp.
N. caroliniana L.
N. integrifolia Ait.
N. pedunculis unifloris Gron.
Mountain Tupelo Mart.
Gum Tree, Sour Gum Tree, Peperidge, Amer.

2. (b.) *villosa Michx.* † North America fig. 1197, 1198. 1317
 The hairy-leaved Nyssa.
N. sylvatica Michx.
N. multiflora Wanch.
N. montana Hort.
N. pedunculis multifloris Gron.
Sour Gum Tree, Black Gum, Yellow Gum, Amer.
Haariger Tulpeibaum, Ger.
 m

3. cándicans *Michx.* & Carolina
fig. 1199. 1318
The whitish-leaved Nyssa, or Ogechee Lime
Tree.

N. capitata Walt.
N. coccinea Bartr.
Sour Tupelo Tree, Wild Lime.
Weisslicher Tulplobaum, Ger.

4. grandidentata *Michx.* & North America
fig. 1200, 1201. 1319

The deeply toothed-leaved Nyssa, or Large Tupelo Tree.
N. tomentosa and *N. angulizans* Michx.
N. denticulata Ait.
N. angulosa Poir.
N. uniflora, Wangh.
Wild Olive, Amer.
Virginian Tupelo, Mart.

- II. OSYRIS *L.* - - 1316. 1320
THE OSYRIS, or Poet's Cassia.
Casia Camer.

1. álba *L.* - Italy - fig. 1202. 1320
The white-flowered Osyris.

O. foliis linearibus acutis Leeb.
O. frutescens baccifera Bauh.
Casia poetica Montpellieranum Cam.
Casia latiflora Alp.
Casia Montpellier dicta Giesn.
Weisse Osyris, Ger.

Elæagnæcæ. & & & 1320

- I. ELÆAGNUS *Tourn.* & & & 1320, 1321
THE ELÆAGNUS, Oleaster, or Wild Olive Tree.
Chalef, Fr.
Wilde Oelthauw, Ger.

1. horténsis *Bieb.* & South of Europe
pl. 219. fig. 1203. 1321.

The Garden Elæagnus.
E. angustifolia L.
E. inermis Mill.
E. argenteus Mœnch.
E. orientalis Delisle.
? *E. argentea* Wats.
Jerusalem Willow.
Olivier de Bohême, Chalef à Feuilles étroites,
Fr.
Schmalblättriger Oleaster, Ger.

Varieties & - - - 1322

1 angustifolia *Bieb.* &

E. angustifolia L.

2 dactyliformis &

E. orientalis L.

4 spinosa &

E. spinosa L.

2. argentea *Ph.* & Hudson's Bay
fig. 1204. 1323

The silver-leaved Elæagnus.
Missouri Silver Tree, U. S. of N. Amer.

- App. i. Half-hardy Species of Elæagnus. 1324

E. conferta *Rorb.* & Nepal.
E. arborea *Rorb.* & Nepal.
E. latifolia L. & East Indies.
E. salicifolia ? *D. Don.* & fig. 1205.

- II. HIPPOPHAE *L.* & 1321. 1324
THE HIPPOPHAE, Sea Buckthorn, or Sallow-
thorn.

Rhamnoides *Tourn.*
Argoussier, Fr.
Hoffdorn, Sanddorn, Ger.
Espino amarillo, Span.

1. Rhamnoides *L.* & Europe
pl. 220. fig. 1206. 1324
The Buckthorn-like Hippophae.

Rhamnoides florifera sativis folio, *Tourn.*
Rhamnoides fructifera *Ray.*
Argoussier faux Nerprun, Fr.
Weidenblättriger Sanddorn, Ger.
Arve, Saule Épineux, Alps of Switzer-
land.

Varieties & & - - 1325

2 angustifolia & &

3 sibirica & &

11. sibirica *Lodd. Cat.*

2. salicifolia *D. Don.* & Nepal
fig. 1207. 1326
The Willow-leaved Hippophae.
11. conferta *Wall.*

- III. SHEPHERDIA *Nutt.* & &
1321. 1327

THE SHEPHERDIA.
Hippophae L.

1. argentea *Nutt.* & & North America
fig. 1208. 1327

The silvery-leaved Shepherdia.
Hippophae argentea *Pursh.*
Missouri Silver Leaf, Buffalo Berry Tree,
Amer.
Rabbit Berry, Beef Suet Tree, American
Indians.
Graisse de Buffle, Buffalo Fat, French
Traders.

2. canadensis *Nutt.* & North America
fig. 1209. 1327

The Canadian Shepherdia.
Hippophae canadensis L.

Aristolochiæcæ. & & & & 1328

- I. ARISTOLOCHIA *L.* & & & & 1328

THE BIRTHWORT.
Aristolochie, Fr.
Osterluzcy, Ger.

1. siphon *L'Hérit.* & N. Am. f. 1210. 1329
The Siphon-like, or Tube-flowered, Birthwort.

A. macrophylla *Lam.*
Aristolochie Syphon, Fr.
Grossblättrige Osterluzcy, Ger.
Pipe Vine, or Birthwort, Amer.

2. tomentosa *Sims.* & North America
fig. 1211. 1329

The tomentose Birthwort.

- App. i. Half-hardy Species of Aristolochia.
1329

1. sempervirens *L.* & Candia.
4. glauca *Desf.* & Barbary.
1. altissima *Desf.*
1. caudata *Desf.*
1. trilobata *Willd.* South America.

	Page
Euphorbiacæ.	1330
♣ ♣ ♣ ♣ □ □ □ □	
I. EUPHORBIA L. ♣ □ □ □ □	1331
THE EUPHORBIA, or <i>Spurge</i> . <i>Tithymalus</i> Tourn. <i>Euphorbe</i> , Fr. <i>Wolfsmilch</i> , Ger.	
<i>E. Characias L.</i> □ Britain f. 1212.	1331
<i>E. aléppica</i> Hort.	
<i>E. spinosa L.</i> □ □ South of Europe	fig. 1213. 1331
App. i. <i>Half-hardy Species of Euphorbia.</i> 1332	
<i>E. dendroïdes L.</i> □ Italy.	
<i>E. mellifera</i> Ait. ♣ □ Madeira. fig. 1214.	
Other species.	
II. STILLINGIA Garden ♣ 1330. 1332	
THE STILLINGIA.	
<i>ligustrina Willd.</i> ♣ Carolina -	1332
The Privet-leaved Stillingia.	
III. BUXUS Tourn. ♣ ♣ □ 1330. 1332	
THE BOX TREE.	
<i>Buis</i> , Fr. <i>Buxbaum</i> , <i>Buchsbaum</i> , Ger.	
<i>1. sempervirens L.</i> ♣ ♣ Europe	1333
The evergreen, or common, Box Tree. <i>Buxus</i> Ray, and other authors. <i>Buis commun</i> , <i>Bois béni</i> , Fr. <i>Buchsbaum</i> , Ger. <i>Busso</i> , <i>Bossolo</i> , <i>Mortel</i> , Ital.	
<i>Varieties and Subvarieties</i> ♣	
♣ □ - - - 1333	
<i>1 arboréscens Mill. Dict.</i> ♣	
<i>1 argentea Hort.</i> ♣	
<i>2 aurea Hort.</i> ♣	
<i>3 marginata Hort.</i> ♣	
<i>2 angustifolia Mill. Dict.</i> ♣	
<i>1 variegata Hort.</i> ♣	
<i>3 suffruticosa Mill. Dict.</i> □ f. 1215	
<i>B. humilis</i> Dod.	
<i>B. s. nana</i> N. Du Ham.	
<i>Buis nain</i> , <i>Buis à Bordurcs</i> , <i>Buis d'Artois</i> , <i>Buis de Hollande</i> , <i>Petit Buis</i> , Fr.	
<i>Zwerch Buchsbaum</i> , Ger.	
<i>4 myrtifolia Lam.</i> ♣	
<i>2. balearica Willd.</i> ♣ Minorca	fig. 1220, 1221. 1341
The Balearic Box. <i>B. s. var. gigantea</i> N. Du Ham. <i>Minorca Box</i> . <i>Buis de Minorque</i> , <i>Buis de Mahon</i> , Fr. <i>Balcarischer Buchsbaum</i> , Ger.	
App. i. <i>Half-hardy Species of Buxus.</i> 1341	
<i>B. chinensis Lk.</i> ♣ □ China.	
App. I. <i>Half-hardy Species belonging to the Order Euphorbiacæ.</i> - 1341	

	Page
Urticacæ. ♣ ♣ ♣ □ ♣ ♣	1342
I. MORUS Tourn. ♣ ♣ ♣ □ ♣	1342, 1343
THE MULBERRY TREE. <i>Mûrier</i> , Fr. <i>Maulbeere</i> , Ger.	
<i>1. nigra Poir.</i> ♣ Persia	pl. 221, 222. fig. 1222. 1343
The black-fruited, or common, Mulberry. <i>Morus</i> Dod. <i>M. fructu nigro</i> Bauh.	
<i>Variety</i> - - - 1344	
♀ <i>laciniata Mill. Dict.</i> ♣	
<i>2. álba L.</i> ♣ China pl. 223, 224. 1348	
The white-fruited Mulberry Tree. <i>M. cándida</i> Dod. <i>M. fructu álbo</i> Bauh. <i>M. álba fructu minóri álbo insúlso</i> Du Ham.	
<i>Varieties</i> ♣ ♣ - - 1348	
<i>2 multicáulis Perrottet.</i> ♣ ♣	
<i>M. tatárica</i> Desf., not of L. or Pall. <i>M. bullata</i> Balbis. <i>M. cucullata</i> Hort. <i>Chinese black Mulberry</i> , Amer. <i>Perrottet Mulberry</i> , <i>Many-stalked Mulberry</i> . <i>Mûrier Perrottet</i> , Fr. <i>Mûrier à Tiges nombreuses</i> , <i>Mûrier des Philippines</i> , Ann. des Sci. <i>Moro delle Filippine</i> , Ital.	
<i>3 Morrettiána Hort.</i> ♣	
<i>Dandolo's Mulberry</i> .	
<i>4 macrophýlla Lodd. Cat.</i> ♣	
<i>M. a. latifolia</i> Hort. <i>M. hispánica</i> Hort. <i>Mûrier d'Espagne</i> , <i>Feuille d'Espagne</i> , Fr.	
<i>5 romana Lodd. Cat.</i> ♣	
<i>M. a. ovalifolia</i> . <i>Mûrier romain</i> , Fr.	
<i>6 nervosa Lodd. Cat.</i> ♣	
<i>M. nervosa</i> Bon. Jard. <i>M. subálba nervosa</i> Hort.	
<i>Subvariety.</i> ♣	
<i>2 longifolia Bon. Jard.</i> ♣	
<i>7 itálica Hort.</i> ♣	
<i>M. itálica</i> Lodd. Catt.	
<i>Subvariety.</i> ♣	
<i>2 rubra</i> ♣	
<i>M. rubra</i> Lodd. Cat.	
<i>8 ròsea Hort.</i> ♣	
<i>Mûrier rose</i> , <i>Feuille rose</i> , Fr.	
<i>9 columbássa Lodd. Cat.</i> ♣	
<i>Columba</i> , Fr.	
<i>10 membranacea Lodd. Cat.</i> ♣	
<i>Mûrier à Feuilles de Parchemin</i> , Fr.	
<i>11 sínensis Hort.</i> ♣	
<i>M. sínensis</i> Hort. <i>M. chinensis</i> Lodd. Cat. <i>The Chinese white Mulberry</i> , Amer.	
<i>12 pumila Nois.</i> ♣	
<i>M. a. nana</i> Hort. Brit.	
<i>Other Varieties</i> - - 1350	
<i>Wild Mulberries.</i>	
<i>La Feuille rose.</i> <i>M. a. s. ròsea</i> .	
<i>1a Feuille dorée.</i> <i>M. a. lucida</i> Hort. <i>M. incana</i> Hort.	

Plagiánthus divaricatus Forst. ♣ □ New Zealand.
P. siccoides Hook.

Clusia alaternoides Bot. Mag. ♣ □ Cape of Good Hope

Page

La Reine hâtarde.
? *Fuglia zarda*, Ital.
La Femelle.
Grafted Mulberries.
La Reine.
La grosse Reine.
M. a. macrophylla subar.
La Feuille d'Espagne.
M. a. macrophylla.
La Feuille de floes.
? *Fuglia doppia*, Ital.

3. (a.) constantinopolitana Poir. ♀ Turkey 1358

The Constantinople Mulberry Tree.
M. byzantina Sieb.

4. (a.) tatarica Pal. ♀ Banks of the Volga - - fig. 1225. 1358
The Tartarian Mulberry Tree.

5. rubra L. ♀ N. America pl. 225. 1359
The red-fruited Mulberry Tree.
M. virginica Pluk.
M. pennsylvanica Nois.

? *Variety* ♀ - - - 1360
canadensis Lam. ♀

6. (r.) scabra Willd. ♀ N. America 1360
The rough-leaved Mulberry Tree.
M. canadensis Poir.

App. i. Half-hardy Species of *Morus*. 1360

M. indica L. ♀ □ East Indi s.
M. mauritiana Jacq. ♀ □ Madagascar.
M. latifolia Willd. Isle of Bourbon.
M. australis Willd. Isle of Bourbon.
M. reticulata Thunb. Quito.
M. corbifolia Thunb. Quito.
M. ede irgalli Cam. New South Wales.
M. atropurpurea. Nepal.
M. parvifolia. Nepal.
M. serrata. Nepal.
M. heterophylla.
M. lasiocata viridis. Nepal.
M. scandens. Nepal.

II. BROUSSONETIA Vent. 1342. 1361
THE BROUSSONETIA.

Morus Sieb *Kempf.*
Papirus Encyc. Bot.

1. papyrifera Vent. ♀ China pl. 226. 1361
The paper-bearing Broussonetia, or Paper Mulberry.

Morus papyrifera L.
Variety ♀ - - - 1361
2 *cucullata* ♀
B. cucullata Bon Jard.
B. spatulata Hort. Brit.
B. navicularis Lodd. Cat.

III. MACLURA Nutt. ♀ 1342. 1362
THE MACLURA.
Törylon Rafinesque.

1. aurantiaca Nutt. ♀ North America fig. 1226, 1227, 1228. 1362
The orange-like-fruited Maclura, or Osage Orange.
Bow-wood, Yellow-wood, N. Amer.

IV. FIGUS Tourn. ♀ 1343. 1365
THE FIG TREE.
Figuier, Fr.
Feigenbaum, Ger.

1. Carica L. ♀ S. Europe pl. 227. 1365
The common Fig Tree.
F. communis Bauth.
F. humilis and *F. sylvestris* Tourn.
Figuier commun, Fr.
Feigenbaum, Ger.
Varieties - - - 1366
Garden Varieties

V. BORYA W. ♀ - 1343. 1370
THE BORYA.
Adelia Michx.
Bigelövia Sm.

1. ligustrina Willd. ♀ N. America 1370
The Privet-like Borya.
Adelia ligustrina Michx.
Bigelövia ligustrina Sm.

2. (? l.) acuminata Willd. ♀ Carolina fig. 1229. 1371
The acuminate-leaved Borya.
Adelia acuminata Michx.
Bigelövia acuminata Sm.

3. (l.) porulosa Willd. ♀ Georgia. 1371
The pore-like-dotted-leaved Borya.
Adelia porulosa Michx.
Bigelövia porulosa Sm.
? *Bigelövia ovata* Lodd. Cat.

4. distichophylla Nutt. ♀ E. Tennessee 1371
The two-rowed-leaved Borya.

Ulmiceæ. ♀ □ □ - 1371

I. ULMUS L. ♀ - - 1371. 1373
THE ELM.

Orme, Fr.
Ulm, or *Rüster*, Ger.
Olmo, Ital.

1. campestris L. ♀ Britain pl. 228, 229 fig. 1232. 1238, 1239. 1374
The English, field, or common small-leaved, Elm.
U. Atinia Pliny.
U. minor, folio angusto scabro, Ger. Emac.

Varieties ♀ - - - 1375
A. Timber Trees.

- 1 vulgaris ♀
U. campestris Hort. Dur.
- 2 latifolia Hort. ♀
- 3 alba Masters. ♀
- 4 acutifolia Masters. ♀
- 5 stricta Hort. Dur. ♀ pl. 230.
Red English Elm.
- 6 virens Hort. ♀
Kidbrook Elm.
- 7 cornubiensis Hort. ♀
The Cornish Elm.
U. stricta Lindl.
Subvarieties. ♀
- 2 parvifolia Lindl. ♀
U. s. microphylla Lodd. Cat.
aspera Lodd. Cat.
crispa Lodd. Cat.
- 8 sarniensis ♀
The Jersey Elm.
U. sarniensis Lodd. Cat.
- 9 tortuosa ♀
The twisted Elm.
U. tortuosa Lodd. Cat.
? *Orme tortillard*, Fr.

B. Ornamental, or curious, Trees.

- 10 foliis variegatis Lodd. Cat. ♀
- 11 betulæfolia ♀
U. betulæfolia Lodd. Cat
- 12 viminalis ♀ pl. 231.
U. viminalis Lodd. Cat.

Page

Page

- 13 parvifolia \checkmark fig. 1250.
U. parvifolia Jacq.
U. microphylla Pers.
U. pumila var. β (*transbaicalensis*)
 Pall.
U. pumila Willd.
U. p. foliis parvis, &c., Pluk.
U. hionitis Enum.

- 14 planifolia \checkmark pl. 232.
U. planifolia Hort.

- 15 chinensis \checkmark fig. 1231.
U. chinensis Pers.
Thé de l'Abbé Gallois, Orme nain,
 Fr.

- 16 cucullata Hort. \checkmark

- 17 concavifolia Hort. \checkmark

- 18 foliis aureis Hort. \checkmark

Other Varieties.

- U. c. nana* Lodd. Cat.
U. c. foliis maculatis Lodd. Cat.
U. dubia Lodd. Cat.
U. viscosa Lodd. Cat.

French Varieties - 1378

- L'Orme à Feuilles larges et rudes.
 L'Orme Tall, l'Orme Têteul, l'Orme de Hollande.
The British, or Lime Tree, Elm.

- L'Orme à Feuilles lisses et glabres.
The dwarf Elm.

- L'Orme à Feuilles lisses et glabres.
The shining smooth-leaved Elm.

- Le petit Orme à Feuilles panachées de blanc.
The shining silvery-leaved Elm.

- Le petit Orme à Feuilles panachées de jaune.
The dwarf golden-leaved Elm.

- L'Orme à petites Feuilles, l'Orme mâle, l'Orme
 pyramidal.

- The small-leaved Elm.*

- L'Orme à très-grandes Feuilles, l'Orme femelle,
 l'Orme de Trison.

- The large-leaved Elm.*

- L'Orme de Hollande à grandes Feuilles panachées.
The variegated Dutch Elm.

- L'Orme tortillard.
U. torosa Lodd. Cat.

- The twisted Elm.*

2. (c.) suberosa *Mench.* \checkmark England
 pl. 233. fig. 1240. 1395

The cork-barked Elm.

- U. campestris* Woods.

- U. campestris* and *Theophrasti Du Ham.*

- U. vulgatissima folio l'eto scabra* Ger.
 Emac.

- U. montana* Cam.

- Common Elm Tree, Hunt Evel.*

- L'Orme Liège, l'Orme-fongeur, Fr.*

Varieties \checkmark - - 1395

- 1 vulgaris \checkmark

The Dutch cork-barked Elm.

- U. suberosa* Hort. Dur.

- 2 foliis variegatis Lodd. Cat. \checkmark

U. suberosa variegata Hort. Dur.

- 3 alba \checkmark

U. suberosa alba Masters.

- 4 erecta Lodd. Cat. \checkmark

- 5 var. \checkmark

The broad-leaved Hertfordshire Elm,
Wood.

- 6 var. \checkmark

The narrow-leaved Hertfordshire Elm,
Wood.

3. (c.) major *Sm.* \checkmark England
 pl. 234, 235. fig. 1241. 1395

The greater, or Dutch cork-barked, Elm.

- U. hollandica* Mill.

- U. major hollandica*, &c., Pluk.

- U. major, ampliore folio*, &c., Du Ham.

- Filia mas* Matth.

- U. latifolia* Michx.

4. carpinifolia *Lin.* \checkmark England - 1396
 The Hornbeam-leaved Elm.

5. effusa *Willd.* \checkmark England
 pl. 236, 237. fig. 1242. 1397

The spreading-branched Elm.

- U. ciliata* Ehrh.

- U. pedunculata* Lam.

- U. octandra* Schk.

- U. folio latissimo*, &c., Buxb.

- U. laevis* Pall.

- L'Orme pédonculé, Fr.*

6. montana *Bauh.* \checkmark England
 fig. 1243, 1244. 1398

The mountain, Scotch, or Wych, Elm.

- U. glabra* Huds.

- U. effusa* Sibth.

- U. scabra* Mill.

- U. nuda* Ehrh.

- U. campestris* Willd.

- Wych Hazel* of old authors.

Varieties \checkmark - - 1398

A. Timber Trees.

- 1 vulgaris \checkmark

- 2 rugosa *Masters.* \checkmark

U. rugosa Lodd. Cat.

- 3 major *Masters.* \checkmark pl. 238.

- 4 minor *Masters.* \checkmark

- 5 cebennensis *Hort.* \checkmark

The Cevennes Elm.

- 6 nigra \checkmark

The black *Irish* Elm.

U. nigra Lodd. Cat.

- 7 australis *Hort.* \checkmark

B. Ornamental or curious Varieties.

- 8 pendula \checkmark pl. 239.

U. pendula Lodd. Cat.

U. glabra decumbens Hort. Dur.

U. horizontalis Hort.

U. rubra Hort. Soc. Gard.

- 9 fastigiata *Hort.* \checkmark pl. 240.

U. glabra replicata Hort. Dur.

U. Fördii Hort.

U. exoniensis Hort.

- 10 crispa \checkmark

The curled-leaved Elm.

? *U. crispa* Willd.

Other Varieties.

7. (m.) glabra *Mill.* \checkmark Eng. f. 1245. 1403
 The smooth-leaved, or Wych, Elm.

- U. montana* β Fl. Br.

- U. folio glabro* Ger. Emac.

- U. campestris* var. 3. With.

- The feathered Elm.*

Varieties \checkmark - - 1404

A. Timber Trees.

- 1 vulgaris \checkmark

The common smooth-leaved Elm.

- 2 vegeta \checkmark

U. americana vegeta Hort Soc. Gard.

U. americana Masters.

The Haulingdon Elm, the Chi-

chester Elm, the American Elm,

? the Scampston Elm.

- 3 var. \checkmark

The Scampston Elm.

- 4 major \checkmark

U. glabra major Hort. Dur.

The Canterbury Seedling.

- 5 glandulosa *Lindl.* \checkmark

- 6 latifolia *Lindl.* \checkmark

- 7 microphylla *H. S.* \checkmark

? *U. parvifolia* Hort. Soc. Gard

- Page
- B. Ornamental or curious Trees.
- 8 pëndula \checkmark
- U. campestris pëndula* Hort. Dur.
The Downton Elm.
- 9 variegata H. S. \checkmark
- 10 ramulosa Booth. \checkmark
8. álba *Kül.* \checkmark Hungary - - 1405
The whitish-leaved Elm.
9. americana L. \checkmark North America
pl. 241, fig. 1246. 1406
The American Elm.
The white Elm, Amer.
Canadian Elm, American white Elm.
- Varieties* \checkmark - - 1406
- 1 rùbra *Ait.* \checkmark
- 2 álba *Ait.* \checkmark
- U. mollifolia* Roem. et Schult.
- 3 pëndula *Pursh.* \checkmark
- 4 incisa H. S. \checkmark pl. 242.
10. (a.) fúlva *Michx.* \checkmark North America
fig. 1247. 1407
The tawny-budded, or stippery, Elm.
U. rubra Michx.
Orme gras, Fr.
Red Elm, Red-wooded Elm, Moose Elm.
11. alata *Michx.* \checkmark Virginia f. 1248. 1408
The Wahoo, or cork-winged, Elm.
U. pùmila Walt.
Wahoo, Indians of N. America.
- App. i. *Doubtful Sorts of Ulmus.* - 1409
- U. pubescens* Walt.
U. fruticosus Willd.
U. integrifolia. Himalayas.
U. virgata. Himalayas.
U. lanceifolia Royle. Himalayas.
U. crósa Royle. Himalayas.
U. levigata Royle. Himalayas.
U. virgata Royle. Himalayas.
- II. PLANERA Gmel. \checkmark 1372. 1409
THE PLANERA.
Rhamnus Pall., Güldenst.
Ulmus, various authors, as to *Planera* Richardi.
1. Richardi *Michx.* \checkmark West of Asia
pl. 243, 244, fig. 1249, 1250. 1409
Richard's Planera, or Zelkova Tree.
P. crenata Michx., Desf.
P. carpinifolia Wats.
Rhamnus carpinifolius Pall.
Rhamnus ulmóides Güldenst.
Ulmus crenata Hort. Par.
Ulmus parvifolia Willd.
Ulmus campestris Walt.
Ulmus polygama Richard.
Ulmus nemorátis Ait.
Ulmus foliis crenatis basi aequalibus fructu oróide, non compresso, Poir.
Le Zelkova, Orme de Sibérie, Fr.
Richard's Planera, Ger.
2. Gmelini *Michx.* \checkmark N. Am. f. 1251. 1413
Gmelin's Planera.
P. ulmifolia Michx.
P. aquatica Willd.
Anonymus aquaticus arbor, &c., Walt.
- ² *P. Abelicea* Schultes *Crete.* - - 1415
The Abelicea of Ctesius.
- III. CELTIS *Tourn.* \checkmark \square \square \square
THE CELTIS, or Nettle Tree. 1372. 1413
Lóbus of Lobel and other authors.
Micoaculicr. Fr.
Züngelbaum. Ger.
- Page
1. austrális L. \checkmark S. Europe f. 1252. 1414
The southern Celtis, or European Nettle Tree.
Lóbus árbor Lob.
Lóbus síve Celtis Cam.
Lote Tree.
Micoaculicr austral. *Micoaculicr de Provence, Fabreaculicr.* *Faubreguier des Provençaux, S. Du Ham.*
Lote, Ital.
- Variety* \checkmark - - 1414
With variegated leaves, *Brotero.* \checkmark
2. (a.) caucásica *Willd.* \checkmark Caucasus 1415
The Caucasian Celtis.
3. Tournefortii *Lam.* \checkmark \square Armenia
pl. 245. 1416
Tournefort's Celtis.
C. orientális minor, foliis minoribus et crassioribus fructu flavo. Tourn.
C. orientális Mill., not of L.
Micoaculicr du Levant, Micoaculicr d'Orient, Fr.
Morgelandischer Züngelbaum, Ger.
4. (T.) sínensis *Pers.* \checkmark \square China 1416
The China Celtis.
5. Willdenoviana *Schultes.* \checkmark China 1416
Willdenow's Celtis.
C. sinensis Willd.
6. occidentális L. \checkmark North America
pl. 246, 247. 1417
The Western Celtis, or North American Nettle Tree.
C. fructu obscuro purpurascente, Tourn.
C. obliqua Mönch.
Nettle Tree, Sugar Berry, Amer.
Bois inconnu, Illinois.
Micoaculicr de Virginie, Fr.
- Varieties* \checkmark - - 1417
- 2 cordata *Willd.* \checkmark
- 3 scabriuscula *Willd.* \checkmark
- C. austrális* Willd.
C. ? o. β tenuifolia Pers.
C. áspera Lodd. Cat.
C. orientális Hort.
7. crassifolia *Lam.* \checkmark N. Am. f. 1254. 1418
The thick-leaved Celtis, or Hackberry.
C. cordifolia L'Herit.
C. cordata Desfont.
Hagberry, Hoop Ash, Amer.
Micoaculicr à Feuilles en Cœur, Fr.
8. levigata *Willd.* \square Louisiana - 1420
The glabrous-leaved Celtis.
9. pùmila *Ph.* \square Maryland - - 1420
The dwarf Celtis.
- App. i. *Species of Celtis half-hardy, or not yet introduced.* - - 1420
- C. orientális* L. ? \square Himalayas fig. 1255
C. tetrandra *Roarb.* Himalayas.
C. alpina *Royle.* Himalayas.
C. Inglesii *Royle.* Himalayas.
-
- Juglandæcæ. \checkmark - 1420
- I. JUGLANS L. \checkmark - 1420, 1421
THE WALNUT TREE.
Noyer, Fr.
Walnuts, Ger.

Page

1. *règia* L. ☿ Persia
 pl. 248, 249, 250. fig. 257. 1423
 The royal, or common, Walnut Tree.
Nux Jùglans Dod.
Nux Jùglans, seu règia vulgàris, Bauh.
Noyer commun, Fr.
Nosguier, Provence.
Gemeine Walnuss, Ger.
Varieties ☿ - - 1423
 2 *màxima* ☿
Nux Jùglans fructu máximo,
 Bauh.
Noix de Jauge, Bon Jard.
Bannut, Warwickshire.
 3 *ténèra* ☿
 The skin-shelled, or titmouse, Walnut.
Nux Jùglans fructo ténèro et frágile
putámine Bauh.
Noyer à Coque tendre, Noyer Mé-
sange, Bon Jard.
Noyer de Mars, in Dauphin.
 4 *scrótina* Desf. ☿
 The late-vegetating Walnut.
Nux Jùglans fructo scrótino Bauh.
Noyer tardif, Noyer de la Saint
Jean, Bon Jard.
Noyer de Mai, in Dauphiné.
 5 *laciniata* ☿
 The Fern-leaved Walnut.
Nux Jùglans foliis laciniatis Re-
 neaum.
Jùglans heterophylla Hort.
J. trilobifolia Lodd. Cat.
Other Varieties.
 The Highflie.
 The Yorkshire Walnut.

2. *nigra* L. ☿ North America
 pl. 251, 252. fig. 1260. 1435
 The black wooded Walnut Tree.
The black-Walnut, the black Hickory Nut,
 N. America.
Noyer noir, Fr.
 3. *cinèrea* L. ☿ N. Am. p. 253. f. 1262.
 1439
 The grey-branched Walnut Tree, or *Butter-nut*.
J. cathartica N. Amer. Sylv.
J. obtùnga Mill.
Oil-nut, White Walnut, Amer.
Noyer cendré, Fr.
Graue Walnuss, Ger.

II. CARYA Nutt. ☿ - 1421. 1441
 THE CARYA, or Hickory Tree.
Jùglans sp. L.
Hicòrius Rafinesque.
Hickory, Amer.

1. *olivæfórmis* Nutt. ☿ North America
 fig. 1263. 1441
 The olive-shaped Caryá, or *Pecanè-nut Hickory*.
Jùglans rubra Gærtn.
Jùglans cylindrica Lam.
Jùglans Pècan Mùhlenb.
Jùglans angustifolia Ait.
Jùglans olivæfórmis Michx.
Pècan-nut, Illinois Nut, Amer.
Pècanier, Pacanus, Noyer Pècanier, Fr.
 2. *amàra* Nutt. ☿ N. Amer. f. 1264. 1443
 The bitter-nut Caryá.
Jùglans amàra Michx.
Bitter-nut, White Hickory, Swamp Hickory,
 Amer.
 3. *aquática* Nutt. ☿ North America
 fig. 1265, 1266. 1414
 The aquatic Caryá, or *Water Bitter-nut Hickory*.
Jùglans aquática Michx.

Page

4. *tomentòsa* Nutt. ☿ N. Am. f. 1267. 1444
 The tomentose Caryá, or *Mocker-nut Hickory*.
Jùglans álba L.
C. álba Mill.
Jùglans tomentòsa Michx.
White-heart Hickory, Common Hickory,
 Amer.
Noyer dur, Illinois.
Variety ☿ - - 1445
 2 *màxima* Suet. Hort. Brit. ☿
 5. *álba* Nutt. ☿ North America
 pl. 254. fig. 1269. 1446
 The white-nutted Caryá, or *Shell-bark Hickory*.
Jùglans álba Michx.
Jùglans álba ovàta Marsh.
Jùglans squamòsa Michx.
Jùglans compressa Gærtn.
Shag-bark Hickory, Scaly-bark Hickory,
Kisky Thomas Nut, Amer.
Noyer tendre, Illinois.
 6. *sulcàta* Nutt. ☿ N. Am. f. 1271. 1448
 The furrowed-fruited Caryá.
Jùglans laciniòsa Michx. Arb.
Jùglans mucronàta Michx. Fl. Bor. Amer.
Jùglans sulcàta Willd.
Thick Shell-bark Hickory, Springfield Nut,
Gloucester Nut, Amer.
 7. *porcìna* Nutt. ☿ North America
 fig. 1272, 1273, 1274. 1419
 The Pig-nut Caryá.
Jùglans porcìna *æ* *obcordàta* Michx. Arbr.
Jùglans porcìna var. with fruit round and
somewhat rough, Michx. N. Amer. Syl.
Jùglans obcordàta Mùhlenb.
Big-nut, Hog-nut, Broom Hickory.
Variety ☿ - - 1449
 2 *glàbra* ☿ f. 1272. b, and 1274. b.
Jùglans porcìna ßficifórmis Michx.
 Arb.
Jùglans glàbra Mùhl.
 8. *myristicæfórmis* Nutt. ☿ South Carolina
 fig. 1275. 1451
 The Nutmeg-like-fruited Caryá, or *Nutmeg Hickory*.
Jùglans myristicæfórmis Michx.
 9. *microcárpa* Nutt. ☿ N. America 1451
 The small-fruited Caryá.
 10. *integrifólia* Spreng. ☿ - - 1451
 The entire leaf (let) ed Caryá, or *Hickory*.
Hicòrius integrifolius Rafinesque.
 App. i. *Other kinds of Caryá.* - 1451
C. ambigua. North America.
Jùglans ambigua Michx.
C. pubescens Lk.
C. rigida.
Jùglans vigida Lodd. Cat.

III. PTEROCARYA Kunth. ☿
 THE PTEROCARYA. 1421. 1451
Jùglans sp. L.

1. *caucásica* Kunth. ☿ Caucasus
 pl. 255. fig. 1276. 1452
 The Caucasian Pterocarya.
Jùglans pterocárpa Michx.
Rhus obscurum Bieb.
Jùglans Fraxinifolia Lamond MS.
Fraxinus levigata Hort. Par.

Salicàcææ. ☿ ☿ ☿ ☿ ☿ 1453

- I. *SALIX* L. ☿ ☿ ☿ ☿ ☿ - - 1453
 THE WILLOW.

	Page
<i>Harab</i> , Hebrew.	
<i>Hea</i> , Gr.	
<i>Säliz</i> , Lat.	
<i>Sauze</i> , Fr.	
<i>Weide</i> , Felber, Ger.	
<i>Salcio</i> , Ital.	
<i>Sauze</i> , Span.	
<i>Willd.</i> , Swe.L.	
<i>Willg.</i> , Flem.	
<i>Willg.</i> , Anglo-Sax.	
<i>Willow</i> , <i>Withy</i> , <i>Sallow</i> , <i>Osier</i> , Eng.	
<i>Saugb.</i> , Scotch.	
Group i. <i>Purpurea</i> Koch, Borrer. ♀ ♂ 1490	
<i>Osier Willows, with one Stamen in a Flower.</i>	
1. <i>purpurea</i> L. ♂ Britain fig. 1294., and fig. 1. in p. 1603. - - 1490	
The purple Willow.	
♂. <i>purpurea</i> ♂ Koch Comm.	
<i>Varieties</i> ♀ - - 1490	
1 ♂	
♂. <i>purpurea</i> Smith, Willd.	
2 ♂	
♂. <i>Lambertiana</i> Smith, Willd.	
3 ♂	
♂. <i>Helix</i> Willd. Em.	
1 monalephica Koch. ♂	
♂. <i>sericea</i> K. ch. ♂	
♂. <i>monandra vertice</i> Ser. Sal. Helv.	
♂. <i>bractea rubra</i> Koch. ♂	
2. <i>Helix</i> L. ♂ ♂ Britain fig. 2. in p. 1603. - - 1491	
The <i>Helix</i> , or <i>Rose</i> , Willow.	
♂. <i>purpurea</i> var. Koch Comm.	
? ♂. <i>oppositifolia</i> Host Sal. Austr.	
3. <i>Lambertiana</i> Sm. ♂ ♂ England fig. 3. in p. 1603. - - 1492	
Lambert's, or the <i>Poyt'n</i> , Willow.	
♂. <i>purpurea</i> ♂ Koch Comm.	
4. <i>Woollgariana</i> Borr. ♂ ♂ England fig. 4. in p. 1603. - - 1492	
Woollgar's Willow.	
♂. <i>monandra</i> Sal. Wob.	
♂. <i>monandra</i> var. Hoffm. Hist. Sal.	
5. <i>Forbyana</i> Sm. ♂ ♂ England fig. 5. in p. 1603. - - 1492	
Forby's Willow, or the <i>fine Basket Osier</i> .	
♂. <i>fissa</i> Lin. Soc. Trans., not of Hoffm.	
♂. <i>rubra</i> ♂ Koch Comm.	
6. <i>rubra</i> Hud. ♂ ♂ Britain fig. 6. in p. 1604. - - 1493	
The <i>red</i> , or <i>green-leaved</i> , Willow, or <i>Osier</i> .	
♂. <i>rubra</i> , in part. Koch Comm.	
♂. <i>fissa</i> Hoffm. Sal.	
♂. <i>concolor</i> Host Sal.	
♂. <i>viridescens</i> Vill. Dauph.	
♂. <i>lividaris</i> Walker's Essays.	
App. i. <i>Purpurea</i> of which Plants have been introduced, but not described. 1493	
♂. <i>elliptica</i> Todd. Cat.	
App. ii. <i>Purpurea</i> described by Authors, but not yet introduced, or of doubtful Identity with Species already in the Country. 1493	
♂. <i>concolor</i> , mas. et fem., H. et Sal.	
♂. <i>mutabilis</i> <i>fragilis</i> <i>longissima</i> , &c., Ray.	
♂. <i>rubra</i> Engl. Fl.	
♂. <i>Helix</i> L.	
♂. <i>Helix</i> , mas. et fem., H. et Sal.	
♂. <i>oppositifolia</i> , mas. et fem., H. et Sal.	
♂. <i>Helix</i> L.	
♂. <i>purpurea</i> , mas. et fem., H. et Sal.	
♂. <i>mutabilis</i> , mas. et fem., H. et Sal.	
♂. <i>concolor</i> , mas. et fem., H. et Sal.	
♂. <i>mutabilis</i> , mas. et fem., H. et Sal.	

	Page
Group ii. <i>Acutifolia</i> Borrer. ♂ ♂ 1494	
<i>Prunöse</i> Koch.	
<i>Willows with dark Bark, covered with a fine Bloom.</i>	
7. <i>acutifolia</i> Willd. ♂ ♂ <i>Podolia</i> fig. 25. in p. 1607. - - 1494	
The pointed-leaved Willow.	
♂. <i>violacea</i> Andr. Bot. Rep., not of Willd. nor S. <i>ca-pica</i> Hort.	
8. <i>daphnoides</i> Villars. ♂ ♂ Switzerland fig. 1295., and fig. 26. in p. 1608. 1494	
The <i>Daphne-like</i> Willow.	
♂. <i>præcox</i> Hoppe.	
♂. <i>bigemmis</i> Hoffm.	
♂. <i>cinerea</i> Host Sal.	
9. <i>pomeranica</i> Willd. ♂ ♂ Pomerania 1496	
The <i>Pomeranian</i> Willow.	
♂. <i>daphnoides</i> var. Villars.	
Group iii. <i>Triandra</i> Borrer. ♂ ♂ 1496	
<i>Amgdalina</i> Koch.	
<i>Osier Willows, with three Stamens in a Flower.</i>	
10. <i>undulata</i> Koch, Hook. ♂ ♂ England fig. 1296., and fig. 13. 14. in p. 1605. 1497	
The <i>wavy-leaved</i> Willow.	
♂. <i>undulata</i> Ehrh. Beytr.	
? ♂. <i>No. 38.</i> , Trev. Obs. Bot.	
♂. <i>lancolata</i> Sm.	
<i>Varieties</i> ♂ ♂ - - 1497	
2 <i>undulata</i> Forbes. ♂ ♂	
3 <i>lancolata</i> Smith. ♂ ♂	
4 Having the catkins androgynous ♂ ♂	
11. <i>hippophæfolia</i> Thuillier. ♂ ♂ Silesia, &c. 1498	
The <i>Sea-Buckthorn-leaved</i> Willow, or <i>Osier</i> .	
♂. <i>undulata</i> Trev. Obs. Bot.	
? ♂. <i>undulata</i> var. Borr. in a letter.	
12. <i>triandra</i> L. ♂ ♂ Britain fig. 1297., and fig. 15. in p. 1605. - - 1498	
The <i>three-stamened-flowered</i> Willow, or <i>Osier</i> .	
♂. <i>amygdalina</i> , part of, Koch Comm.	
<i>Varieties</i> , ♂ ♂	
? 2 The <i>French willow</i> ♂ ♂ 1499	
♂. <i>triandra</i> Curt. Fl. Lond.	
? ♂. <i>Hoppeana</i> Willd.	
? 3 <i>Hoppeana</i> ♂ ♂ - - 1500	
♂. <i>androgyna</i> Hoppe.	
♂. <i>Hoppeana</i> Willd. Sp. Pl.	
♂. <i>triandra androgyna</i> Seringe.	
♂. <i>amygdalina</i> , part of, Koch Comm.	
? 4 ♂ ♂	
♂. <i>triandra undulata</i> Mertens.	
13. <i>Hoffmanniana</i> Sm. ♂ ♂ Britain fig. 16. in p. 1606. - - 1500	
Hoffmann's Willow, or <i>Osier</i> .	
♂. <i>triandra</i> Hoffm.	
14. <i>amygdalina</i> L. ♂ ♂ Britain fig. 1298., and fig. 18. in p. 1606. - - 1500	
The <i>Almond-leaved</i> Willow, or <i>Osier</i> .	
♂. <i>amygdalina</i> , part of, Koch Comm.	
15. <i>Villarsiana</i> Flugge et Willd. ♂ ♂ Dauphiné fig. 17. in p. 1606. - - 1502	
Villars's Willow, or <i>Osier</i> .	
♂. <i>triandra</i> Villars Delph.	
♂. <i>amygdalina</i> var. Koch Comm.	

- Page
- App. i. *Triándra* of which there are Plants
in the Country not described. 1502
- S. tenuifolia* Lodd. Cat. and G. not of Sm.
- App. ii. *Triándra* described, but not yet
introduced, or of doubtful Identity with Species
in the Country. - - - 1502
- S. spectábilis*, mas et fem., *Host Sal.*
S. semperlórens, mas et fem., *Host Sal.*
S. tenuifóra, mas et fem., *Host Sal.*
S. venusta, mas et fem., *Host Sal.*
S. vária, mas et fem., *Host Sal.*
S. amygdáliná, mas et fem., *Host Sal.*
S. lignístiná, mas et fem., *Host Sal.*
S. speciosa, mas et fem., *Host Sal.*
- Group iv. *Pentándra* Borrer. \checkmark 1503
- Trees, having Flowers with 3—5 Stamens.
16. *pentándra* L. \checkmark Britain fig. 1299. a,
and fig. 34. in p. 1610. - 1503
- The five-stamened-flowered Willow.
S. pentándra, part of. Koch Comm.
The sweet Willow, Bay-leaved Willow.
- Variety \checkmark - - - 1503
- 2 hermaphroditica \checkmark
- S. hermaphroditica* L.
17. *Meyeriana* Willd. \checkmark Pomerania
f. 1300., and f. 33. in p. 1610. 1504
- Meyer's Willow.
- S. cuspidata* Schultz.
S. tinctoria Sm. in Rees's Cycl.
S. pentándra β L.
S. hexándra Ehrh.
S. Ehrhartiana Sm. in Rees's Cycl.
S. tetrandra Willd.
18. *lúcida* Mühlenb. \checkmark North America
f. 1301., and f. 32. in p. 1610. 1504
- The shining-leaved Willow.
S. Forbésii Swt. Hort. Brit.
- Group v. *Frágiles* Borrer. \checkmark 1507
- Trees, with their Twigs mostly brittle at the Joints.
19. *babylónica* L. \checkmark Asia pl. 256. fig. 22.
in p. 1607. - - - 1507
- The Babylonian, or Weeping, Willow.
S. propéncus Sering. Sal. Helv.
S. orientális, &c., Tourn.
S. arabica, &c., C. Bauh.
Sauz pleurcur, Parasol du grand Seigneur,
Fr.
Trauer Weide, *Thränen Weide*, Ger.
- Varieties \checkmark - - - 1513
- 1 *vulgáris* fem. Hort. \checkmark
- 2 *Napoleóna* Hort. \checkmark
- 3 *crispa* Hort. \checkmark pl. 257. fig. 21.
in p. 1606.
20. *decípiens* Hoffm. \checkmark Britain pl. 258.
f. 1309., and f. 29. in p. 1609. 1515
- The deceptive, White Welch, or varnished, Wil-
low.
S. americana Walker's Essays.
S. frágilis, part of. Koch Comm.
21. *montána* Forbes \checkmark Switzerland
fig. 19. in p. 1606. - - - 1515
- The Mountain Willow.
22. *frágilis* L. \checkmark Britain fig. 1310., and
fig. 27. in p. 1606. - - - 1516
- The brittle-twigged, or Crack, Willow.
S. frágilis, in part. Koch Comm.
23. *monspeliensis* Forbes. \checkmark $\&$ Montpe-
lier fig. 30. in p. 1609. - - - 1517
- The Montpellier Willow.
 β *S. frágilis* var. Borr. in a letter.
24. *Russelliana* Sm. \checkmark Britain fig. 1311.,
and fig. 28. in p. 1608. - - - 1517
- The Russell, or Duke of Bedford's, Willow.
 β *S. frágilis* Woody.
The Dishley or Leicestershire, Willow; the
Huntingdon Willow.
S. péndula Ser. Sal. Helv.
S. viridis Fries Nov.
S. rubens Schrank Baier. Fl.
- Varieties - - - - 1521
25. *Purshiana* Borrer. \checkmark N. Amer. 1522
- Pursh's Willow.
S. ambigua Pursh, Sm., Forbes, Hook.
- App. i. *Frágiles* introduced, but not yet de-
scribed, or of doubtful Identity. 1522
- S. adscéndens* Donnell's Nursery.
S. bigemmis Lodd. Cat.
S. decípiens, fem., Lodd. Cat.
S. frágilis Lodd. Cat.
S. murina Lodd. Cat.
S. rubra G. Lodd.
- App. ii. *Frágiles* described, but not yet in-
troduced, or of doubtful Identity with in-
troduced Species. - - - 1522
- S. frágilis*, mas et fem., *Host Sal.*
S. fragilior, mas et fem., *Host Sal.*
S. fragilissima, mas et fem., *Host Sal.*
 β *Frágilis* Host Syn.
S. palustris, mas et fem., *Host Sal.*
S. capensis Thunb. Fl. Cap.
S. subserrata Willd. Sp. Pl.
S. Salsaf *batledi* Forsk. Cat. Pl. Egypt.
- Group vi. *Alba* Borrer. \checkmark 1522
- Trees of the largest Size, with the Aspect of the
Foliage whitish.
26. *álba* L. \checkmark Europe pl. 259, 260. f. 1314,
1315., and f. 136. in p. 1629. 1522
- The whitish-leaved, or common white, Willow.
S. álba, part of. Koch Comm.
The Huntingdon, or Swallow-tailed, Wil-
low.
- Varieties \checkmark - - - 1523
- 2 *carúlea* \checkmark fig. 137. in p. 1629.
- Blue Willow.
S. álba var. Sm. Fl. Br.
S. carúlea Sm. Eng. Bot.
S. álba β Sm. Eng. Fl.
The upland, or red-twigged, Wil-
low, Pontey.
The Leicester Willow, Davy's Agr.
Chem.
- 2 3 *crispa* \checkmark - - - 1525
- 4 *rosea* Lodd. Cat. \checkmark
27. *vitellina* L. \checkmark Britain pl. 261. fig. 20.
in p. 1606. - - - 1528
- The yolk-of-egg-coloured, or yellow, Willow, or
Golden Osier.
S. álba Koch Comm.
- Variety \checkmark - - - 1528
- With reddish branchlets. *Sm* \checkmark

	Page		Page
App. i. <i>Albe</i> described, but which, probably, have not been introduced into Britain. 1528		39. <i>Lyònu?</i> Schl. & Switzerland fig. 10. in p. 1604. - - - 1532	
<i>S. excelsior</i> Host Sud.		Lyon's Willow.	
Group vii. <i>Nigræ</i> . & ¶ - 1529		40. <i>Houstoniàna</i> Pursh. & Virginia and Carolina fig. 11. in p. 1604. 1532	
<i>Extra-European Kinds allied to the Kinds of one or all of the three preceding Groups.</i>		Houston's Willow.	
28. <i>nigra</i> Mùhlenb. ¶ N. America fig. 152. in p. 1630. - - - 1529		<i>S. tristis</i> Lodd. Cat.	
The black, or dark-branched American, Willow.		41. <i>falcata</i> Pursh. & N. America fig. 148. in p. 1630. - - - 1533	
<i>S. caroliniana</i> Michx.		The Sick-leaved Willow.	
<i>S. pentandra</i> Walt.		42. <i>grisea</i> Willd. & Pennsylvania - 1533	
<i>S. vulgaris</i> Clayt. Fl. Virg.		The grey Willow.	
29. <i>Humboldtiana</i> Willd. & Peru fig. 8. in p. 1604. - - - 1529		<i>S. sericea</i> Mùhlenb.	
Humboldt's Willow.		? <i>S. pennsylvànica</i> Forbes.	
30. <i>Bonplandiàna</i> Humb. et Bonpl. & ? ¶ Mexico fig. 9. in p. 1604. - 1529		<i>Variety</i> & - - - 1534	
Bonpland's Willow.		2 <i>glabra</i> &	
App. i. <i>Nigræ</i> described, but not yet introduced. - - - 1530		43. <i>petiolàris</i> Sm. & Scotland fig. 1319., and fig. 23. in p. 1607. - 1533	
<i>S. ligustrina</i> Michx. North America.		The long-petioled Willow.	
<i>S. occidentalis</i> Bosc. Iste of Cuba.		<i>S. grisea</i> Willd.	
<i>S. octandra</i> Sieb. Egypt.		<i>S. grisea</i> var. β <i>subglabrata</i> Koch Comm.	
Group viii. <i>Prinoides</i> Borrer. & ¶ 1530		44. <i>pennsylvànica</i> Forbes. & ? N. America fig. 95. in p. 1620. - - - 1534	
<i>Shrubs, mostly Natives of North America, and used in Basket-making.</i>		The Pennsylvanian Willow.	
31. <i>rigida</i> Mùhlenb. & North America fig. 141. in p. 1630. - - - 1530		? <i>S. petiolàris</i> Sm.	
The stiff-leaved Willow.		? <i>S. grisea</i> Willd.	
<i>S. cordata</i> Michx.		? <i>S. pedicellàris</i> Spreng. Syst., Pursh.	
<i>S. cordifolia</i> Herbs. Banks. MSS.		45. <i>Mùhlenbergiana</i> Willd. & Pennsylvania fig. 145. in p. 1630. - - - 1534	
32. <i>prinoides</i> Pursh. & ¶ North America f. 1317., and f. 40. in p. 1612. 1530		Mùhlenberg's, or the brown American, Willow.	
The Prinos-like Willow.		<i>S. alpina</i> Walt.	
33. <i>discolor</i> Mùhlenb. & North America fig. 147. in p. 1630. - - - 1530		<i>S. incana</i> Michx.	
The two-coloured Willow.		<i>S. flava</i> Schœpfl.	
34. <i>angustata</i> Pursh. & N. America 1531		<i>S. tristis</i> Mùhlenb.	
The narrowed, or tapered-leaved, Willow.		46. <i>tristis</i> Ait. & North America fig. 150. in p. 1630. - - - 1534	
35. <i>confòrmis</i> Forbes. & North America fig. 24. in p. 1607. - - - 1531		The sad, or narrow-leaved American, Willow.	
The uniform-leaved Willow.		47. <i>cordata</i> Mùhlenb. & North America fig. 142. in p. 1630. - - - 1534	
Group ix. <i>Griseæ</i> Borrer. ¶ & & & & 1531		The heart-leaved Willow.	
<i>Chiefly Shrubs, Natives of North America.</i>		Group x. <i>Rosmarinifolia</i> Borrer. 1535	
36. <i>virèscens</i> Forbes. & Switzerland f. 1318., and f. 7. in p. 1604. 1531		& & &	
The greenish-leaved Willow, or verdant Osier.		<i>Low Shrubs, with narrow Leaves.</i>	
<i>S. hippophaefolia</i> Lodd.		48. <i>rosmarinifolia</i> L. & North America f. 1320., and f. 87. in p. 1618. 1535	
37. <i>reflexa</i> Forbes. & North America fig. 94. in p. 1619. - - - 1532		The Rosemary-leaved Willow.	
The reflexed-catkin'd Willow.		<i>S. rosmarinifolia</i> , part of, Koch Comm.	
38. <i>virgata</i> Forbes. & North America fig. 12. in p. 1605. - - - 1532		49. <i>angustifolia</i> Borrer, Hook., ? Wulf. & & Scotland fig. 1321., and fig. 86. in p. 1618. - - - 1535	
The twiggy Willow.		The narrow-leaved Willow.	
		<i>S. arbuscula</i> Sm.	
		<i>S. rosmarinifolia</i> & Koch Comm.	
		<i>S. incubæca</i> L.	
		50. <i>decumbens</i> Forbes. & ? Switzerland fig. 88. in p. 1618. - - - 1536	
		The decumbent Willow.	
		51. <i>fuscata</i> Pursh. & North America 1536	
		The dark-brown-branched Willow.	

	Page		Page
Group xi. <i>Fúscæ</i> Borrer. ☼ ☼ ☼ ☼	1536	57. <i>alaternoides</i> Forbes. ☼	Switzerland
<i>Mostly procumbent Shrubs.</i>		fig. 76. in p. 1618.	- - 1542
52. <i>fúscæ</i> L. ☼ fig. 83. in p. 1618.	1536	The Alaternus-like Willow.	
The brown Willow.		58. <i>proteafolia</i> Schl. ☼ ? ☼	Switzerland
S. <i>rèpens</i> Hook.		fig. 75. in p. 1617.	- - 1542
S. <i>rèpens</i> Koch, <i>part of</i> , Koch Comm.		The Protea-leaved Willow.	
<i>Varieties</i> ☼ ☼ - -	1537	Group xiii. <i>Reticulàte</i> Borrer. ☼	1542
1 <i>vulgàris</i> ☼ Britain fig. 83. in		59. <i>reticulàta</i> L. ☼ England fig. 1323.,	
p. 1618.		and fig. 67. in p. 1616.	- 1542
<i>var. α</i> Hook.		The netted, or wrinkled, leaved Willow.	
S. <i>fúscæ</i> Sm.		App. i. <i>Reticulàte</i> described, but not yet in-	
S. <i>rèpens</i> Koch β Koch Comm.		troduced. - -	1543
2 <i>rèpens</i> ☼ Britain fig. 84. in		S. <i>vestita</i> Pursh. Labrador.	
p. 1618.		Group xiv. <i>Glaúcæ</i> Borrer. ☼ ☼ ☼ ☼	1543
<i>var. β</i> Hook.		<i>Small, upright, with soft silky Leaves.</i>	
S. <i>rèpens</i> L.		60. <i>clæagnoides</i> Schleicher. ☼ Europe	
S. <i>rèpens</i> Koch α Koch Comm.		fig. 69. in p. 1616. - -	1544
3 <i>prostràta</i> ☼ Britain fig. 82. in		The Eleagnus-like Willow.	
p. 1618.		S. <i>clæagnifolia</i> Forbes.	
<i>var. γ</i> Hook.		S. <i>glauca</i> <i>var.</i> Koch.	
S. <i>prostràta</i> Sm.		61. <i>glauca</i> L. ☼ Scotland fig. 1324., and	
4 <i>fu'cida</i> ☼		fig. 68. in p. 1616. - -	1544
<i>var. δ</i> Hook.		The glaucous Mountain Willow.	
S. <i>fu'cida</i> Sm.		S. <i>appendiculàta</i> Fl. Dan.	
? <i>Subvarieties</i> ☼		62. <i>sericea</i> Villars. ☼ Switzerland fig. 74.	
S. <i>ascendens</i> Sm. Eng. Bot. ☼ Britain		in p. 1617. - -	1544
fig. 80. in p. 1618.		The silky Willow.	
S. <i>fu'cida</i> , <i>exclusive of β</i> , Sm. Eng. Fl.		S. <i>glauca</i> Koch Comm.	
S. <i>repens</i> Koch <i>var.</i> Koch Comm.		S. <i>Lappònum</i> Sm.	
S. <i>parvifolia</i> Sm. Eng. Bot. ☼ Britain		63. <i>Lappònum</i> L. ☼ Lapland fig. 1325.,	
fig. 81. in p. 1618.		and fig. 73. in p. 1617. - -	1545
S. <i>fu'cida</i> β Sm. Eng. Fl.		The Laplanders' Willow.	
S. <i>repens</i> Koch <i>var.</i> Koch Comm.		S. <i>arenaria</i> Fl. Dan.	
5 <i>incubæca</i> ☼ England fig. 79.		64. <i>obtusifolia</i> Willd. ☼ ☼ Lapland 1545	
in p. 1618.		The blunt-leaved Willow.	
S. <i>incubæca</i> L.		65. <i>arenaria</i> L. ☼ Scotland fig. 70. in	
6 <i>argentea</i> ☼ England fig. 78. in		p. 1617. - -	1545
p. 1618.		The sand Willow.	
S. <i>argentea</i> Sm.		S. <i>limosa</i> Wahlenb.	
S. <i>rèpens</i> Koch γ Koch Comm.		<i>Variety</i> ☼ - -	1546
53. <i>Doniàna</i> Sm. ☼ Britain fig. 1322.,		? <i>leucophýlla</i> ☼	
and fig. 85. in p. 1618. - -	1540	S. <i>leucophýlla</i> Schl.	
Don's, or the rusty-branched, Willow.		66. <i>obovàta</i> Pursh. ☼ Labrador fig. 144.	
Group xii. <i>Ambígua</i> Borrer. ☼ ☼ ☼ ☼	1540	in p. 1630. - -	1546
<i>Shrubs.</i>		The obovate-leaved, or Labrador, Willow.	
54. <i>ambígua</i> Ehrh. Borrer. ☼ ☼ Engl. 1540		67. <i>canescens</i> Willd. ? ☼ ? ☼ ? Germany 1546	
The ambiguous Willow.		The greyish Willow.	
S. <i>ambigua</i> Koch, <i>part of</i> , Koch Comm.		S. <i>limosa</i> Wahlenb. <i>var.</i> Koch Comm.	
<i>Varieties</i> ☼ ☼ ☼ - -	1541	68. <i>Stuartiàna</i> Sm. ☼ Scotland fig. 72.	
1 <i>vulgàris</i> ☼ ☼		in p. 1617. - -	1546
<i>var. α</i> Borrer in Eng. Bot. Suppl.		Stuart's, or the small-leaved shaggy, Willow.	
2 <i>màjor</i> ☼		S. <i>arenaria masculina</i> Sm.	
β <i>màjor</i> Borrer.		S. <i>Lappònum</i> Walker.	
S. <i>ambigua</i> β Hook.		S. <i>limosa</i> Wahlenb. <i>var. foliis angustor-</i>	
S. <i>versifolia</i> Neesing-		<i>ribus lanceolàtis</i> Koch Comm.	
<i>Sauze de la Suisse.</i>		<i>Variety</i> - -	1547
3 <i>spathulàt</i> ☼			
<i>var. γ</i> <i>spathulàta</i> Borrer.			
S. <i>ambigua</i> γ Hook.			
S. <i>spathulàta</i> Willd.			
4 <i>undulàta</i> ☼			
<i>var. δ</i> <i>undulàta</i> De R.			
S. <i>spathulàta</i> Willd., <i>var. undulàta</i> on Pro-			
fessor Mertens.			
55. <i>finmàrcheica</i> Willd. ? ☼ ? ☼ Finmark	1544		
The Finmark Willow.			
56. <i>versicolor</i> Forbes. ? ☼ ? ☼ Switzer-			
land fig. 77. in p. 1618. - -	1544		
The various-coloured Willow.			

	Page		Page
69. <i>pyrenæica</i> Gouan. ✚ Pyrenees 1547	1547	80. <i>Micheliàna</i> Forbes. ✚ ? Ý fig. 135. in p. 1629. - 1552	1552
The Pyrenean Willow.		Michel's Willow.	
<i>Variety</i> ✚ - - 1547	1547	<i>S. holosericea</i> Willd.	
2 <i>elliptica</i> Dec. ✚		? <i>S. holosericea</i> var. Borrer.	
<i>S. pyrenæica</i> & <i>elliptica</i> Dec.			
70. <i>Waldsteiniàna</i> Willd. ✚ Alps - 1547	1547	81. <i>ferruginea</i> Anderson. Ý Scotland fig. 128. in p. 1627. - 1552	1552
Waldstein's Willow.		The ferruginous-leaved Sallow, or Willow.	
Group xv. <i>Viminàles</i> Borrer. ✚ Ý 1547	1547		
<i>Willows and Osiers.—Mostly Trees, or large Shrubs, with long pliant Branches, used for Basket-making.</i>		82. <i>acuminata</i> Sm. Ý England fig. 1330., and fig. 131. in p. 1628. - 1553	1553
71. <i>subalpina</i> Forbes. ✚ Switzerland fig. 93. in p. 1619 - 1547	1547	The acuminate-leaved, or large-leaved, Sallow, or Willow.	
The subalpine Willow.		<i>S. lanceolata</i> Seringe.	
72. <i>càndida</i> Willd. ✚ North America f. 1326., and f. 91. in p. 1619. 1548	1548	App. i. <i>Viminàles</i> in the Country, but not described. - 1553	1553
The whitish Willow.		<i>S. trichocarpa.</i>	
<i>Varieties</i> - - - 1548	1548	Group xvi. <i>Cinèrea</i> Borrer. ✚ Ý 1553	1553
73. <i>incana</i> Schrank. ✚ ? Ý Pyrenees, &c. f. 1327., and f. 90. in p. 1619. 1548	1548	<i>Sallows.—Trees and Shrubs, with roundish shaggy Leaves, and thick Catkins.</i>	
The hoary-leaved Willow, ? or Osier.		83. <i>pàllida</i> Forbes. ✚ Switzerland fig. 96. in p. 1620. - 1555	1555
<i>S. riparia</i> Willd., &c.		The pale Willow.	
<i>S. lananulæfolia</i> Lapeyr., &c.		84. <i>Willdenoviàna</i> Forbes. ✚ fig. 41. in p. 1613. - 1555	1555
<i>S. angustifolia</i> Poir., &c.		Willdenow's Willow.	
<i>S. rosmarinifolia</i> Gouan. &c.		85. <i>Pontederàna</i> Willd. ✚ Switzerland f. 1331., and f. 43. in p. 1613. 1555	1555
<i>S. viminalis</i> Will.		Pontedera's Willow.	
74. <i>linèaris</i> Forbes. ✚ Switzerland f. 1328., and f. 89. in p. 1619. 1549	1549	<i>S. pinnata alpina nigricans, folio oleagineo serrata</i> Ponted. Comp.	
The linear-leaved Willow.		<i>S. Pontedèræ</i> Bellardi.	
? <i>S. incana</i> var. <i>linèaris</i> Borrer.		86. <i>macrostipulacea</i> Forbes. Ý Switzerland fig. 130. in p. 1627 - 1557	1557
75. <i>viminalis</i> L. ✚ Ý England fig. 1329., and fig. 133. in p. 1629. - 1549	1549	The large-stipuled Sallow.	
The twiggy Willow, or common Osier.		87. <i>incanescens</i> ? Schl. ✚ Ý Switzerland fig. 120. in p. 1625. - 1557	1557
<i>S. longifolia</i> Lam.		The whitish-leaved Sallow.	
<i>Varieties</i> - - - 1550	1550	88. <i>pannosa</i> Forbes. ✚ Ý Switzerland fig. 123. in p. 1626. - 1557	1557
Bark of the branchlets brownish yellow.		The cloth-leaved Sallow.	
Bark of the branchlets dark brown.		? <i>Variety</i> - - - 1558	1558
Velvet Osier.		Leaf, catkin, ovary, and bractea larger.	
76. <i>stipularis</i> Sm. ✚ Ý England fig. 132. in p. 1628. - 1550	1550	89. <i>mutabilis</i> Forbes. ✚ Switzerland 1558	1558
The stipuled, or auricled-leaved, Osier, or Willow.		The changeable Willow, or Sallow.	
77. <i>Smithiàna</i> Willd. Ý England fig. 134. in p. 1629. - 1550	1550	90. <i>cinèrea</i> L. Ý England fig. 1332., and fig. 125. in p. 1626. - 1558	1558
Smith's Willow, or the silky-leaved Osier.		The grey Sallow, or Ash-coloured Willow.	
<i>S. mollissima</i> Sm.		<i>S. cinèrea</i> var. Koch Comm.	
<i>S. acuminata</i> , with narrower leaves. Koch Comm.		<i>Varieties</i> Ý - - - 1559	1559
<i>S. acuminata</i> ? Lindl. Synops.		1 With variegated leaves Ý	
78. <i>mollissima</i> Ehrh. Ý Germany 1551	1551	2 Ý	
The softest, or faced Willow, or Osier.		<i>S. cinèrea</i> Sm., according to Koch.	
<i>S. pubera</i> Koch.		3 Ý	
79. <i>holosericea</i> Hook., ? Willd. ✚ Ý Germany - - - 1551	1551	<i>S. aquatica</i> Sm. according to Koch.	
The velvety, or "soft-shaggy-flowered," Willow, or Osier.		4 Ý	
<i>S. Smithiàna rugosa</i> Forbes.		<i>S. olcypsha</i> Sm., according to Koch.	
? <i>S. acuminata</i> , the var. mentioned by Sm. in Eng. Fl.			
<i>S. acuminata</i> var. <i>rugosa</i> Sm. MSS.			
? <i>S. rubra</i> Walker's Essays.			

- | | Page |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 91. aquática <i>Sm.</i> † England fig. 127. in p. 1627. - - - 1559
The Water Sallow, or Willow.
<i>S. cinerea</i> var. Koch Comm. | |
| 92. olcifolia <i>Sm.</i> † England fig. 126. in p. 1626. - - - 1559
The Olive-leaved Willow, or Sallow.
<i>S. cinerea</i> var. Koch Comm. | |
| 93. geminata <i>Forbes.</i> † ? Britain fig. 129. in p. 1627. - - - 1560
The twin-catkin Sallow, or Willow. | |
| 94. crispa <i>Forbes.</i> † fig. 42. in. p. 1613. 1560
The crisp-leaved Willow. | |
| 95. aurita <i>L.</i> † England fig. 124. in p. 1626. - - - 1560
The round-eared, or trailing Sallow, or Willow.
<i>S. uliginosa</i> Willd.
The trailing Sallow, Norfolk.
Varieties † - - - 1560
<i>S. cladostemma</i> Hayne, according to Koch. †
microphylla † Lodd. †
<i>S. caprea pumila</i> , folio subtundo, subtus incano, <i>Dill.</i> , according to Smith. † | |
| 96. latifolia <i>Forbes.</i> † fig. 118. in p. 1625. 1561
The broad-leaved Willow, or Sallow. | |
| 97. caprea <i>L.</i> † Britain fig. 1333, 1334, 1335., and fig. 122. in p. 1626. 1561
The Goat-Willow, or the great round-leaved Sallow. | |
| 98. sphacelata <i>Sm.</i> † Britain fig. 121. in p. 1625. - - - 1563
The withered pointed-leaved Willow, or Sallow. | |
| Group xvii. <i>Nigricantes</i> Borrer. † † † 1563
<i>Shrubs, with long Branches, or small Trees. Mostly Sallows.</i> | |
| 99. australis <i>Forbes.</i> † Switzerland fig. 103. in p. 1621. - - - 1565
The southern Sallow, or Willow. | |
| 100. vaudensis <i>Forbes.</i> † Switzerland fig. 117. in p. 1624. - - - 1565
The Vaudois Sallow, or Willow. | |
| 101. grisophylla <i>Forbes.</i> † Switzerland fig. 119. in p. 1625. - - - 1565
The grey-leaved Willow, or Sallow. | |
| 102. lacustris <i>Forbes.</i> † Switzerland fig. 116. in p. 1624. - - - 1566
The Lake Willow, or Sallow. | |
| 103. crassifolia <i>Forbes.</i> † fig. 115. in p. 1624. - - - 1566
The thick-leaved Willow, or Sallow. | |
| 104. cotinifolia <i>Sm.</i> † Britain fig. 1336., and fig. 114. in p. 1624. - - - 1566
The Cotinus, or Quince, leaved Sallow, or Willow.
<i>S. spadicca</i> Vill.
<i>S. phyticifolia</i> var. Koch Comm. | |
| 105. hirta <i>Sm.</i> † † Britain fig. 113. in p. 1623. - - - 1567
The hairy-branched Sallow, or Willow.
<i>S. plicata</i> Schl. is the fem. of <i>S. hirta</i> <i>Forbes.</i> | |
| 106. rivularis <i>Forbes.</i> † Switzerland fig. 102. in p. 1621. - - - 1567
The River Willow, or Sallow. | |
| 107. atropurpurea <i>Forbes.</i> † Switzerland 1567
The dark-purple-branched Willow, or Sallow. | |
| 108. coriacea <i>Forbes.</i> † Switzerland fig. 112. in p. 1623. - - - 1568
The coriaceous-leaved, or leathery, Willow, or Sallow. | |
| 109. nigricans <i>Sm.</i> † Britain. fig. 37. in p. 1611. - - - 1568
The dark broad-leaved Willow.
<i>S. phyticifolia</i> β L. | |
| 110. Andersoniana <i>Sm.</i> † Scotland fig. 109. in p. 1623. - - - 1568
Anderson's Willow, or the green Mountain Sallow.
<i>S. phyticifolia</i> var. Koch.
Varieties - - - 1569 | |
| 111. damascena <i>Forbes.</i> † - - - 1569
The Damson-leaved Willow, or Sallow.
<i>S. damascenifolia</i> Anderson MSS.
<i>S. phyticifolia</i> L., a state of, Lindl. Syn. | |
| 112. Ansoniana <i>Forbes.</i> † Switzerland fig. 107. in p. 1622. - - - 1569
Anson's Sallow, or Willow. | |
| 113. helvetica <i>Forbes.</i> † Switzerland 1570
The Swiss Willow, or Sallow. | |
| 114. firma <i>Forbes.</i> † fig. 106. in p. 1622. 1570
The firm-leaved Sallow, or Willow. | |
| 115. carpinifolia <i>Schl.</i> † Germany 1570
The Hornbeam-leaved Sallow, or Willow. | |
| 116. rotundata <i>Forbes.</i> † † Switzerland fig. 1337. in p. 1572., and fig. 104. in p. 1621. - - - 1571
The round-leaved Willow, or Sallow.
? <i>S. rotundifolia</i> Host. | |
| 117. dura <i>Forbes.</i> † fig. 105. in p. 1622 1571
The hardy Sallow, or Willow. | |
| 118. Forsteriana <i>Sm.</i> † † Britain fig. 110. in p. 1623. - - - 1571
The glaucous Mountain Sallow, or Forster's Willow.
<i>S. phyticifolia</i> var. Koch. | |

- | | Page |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------|
| 119. <i>rupēstris</i> Donn. ✕ Scotland fig. 111.
in p. 1623. - - 1573
The silky Rock Willow, or Sallow. | 111.
1573 |
| 120. <i>tennifolia</i> L. ☼ England fig. 50, in
p. 1611. - - 1573
The thin-leaved Willow.
<i>S. arbuscula</i> Wahlenb. var. Koch.
<i>S. l.</i> of Eng. Bot. is <i>S. bicolor</i> of Hook.
? <i>Variety</i> - - 1573
With silky hairs on the upper
half of the ovary, and towards
the base of its stalk. | 50.
1573
1573 |
| 121. <i>propinqua</i> Borr. ☼ ✕ Britain 1574
The nearly related, or flat-leaved, upright,
Mountain Willow. | 1574 |
| 122. <i>petraea</i> Anders. ☼ Britain fig. 97.
in p. 1620. - - 1574
The Rock-Sallow, or Willow.
<i>S. arbuscula</i> Wahlenb. | 97.
1574 |
| 123. <i>Ammanniana</i> Willd. ✕ Alps 1575
Ammann's Willow. | 1575 |
| 124. <i>atrovirens</i> Forbes. ☼ Switzerland
fig. 108, in p. 1622. - 1575
The dark-green Sallow, or Willow. | 1575 |
| 125. <i>strépida</i> Forbes. ☼ Switzerland
fig. 100, in p. 1621. - 1575
The creaking Willow, or Sallow. | 1575 |
| 126. <i>sórdida</i> Forbes. ☼ Switzerland
fig. 101, in p. 1621. - - 1576
The sordid Sallow, or Willow. | 1576 |
| 127. <i>Schleicheriana</i> Forbes. ☼ Switzer-
land fig. 98, in p. 1620. - 1576
Schleicher's Willow, or Sallow. | 1576 |
| 128. <i>grisonensis</i> Forbes. ☼ Grisons fig. 99,
in p. 1620. - - 1576
The Grisons Sallow, or Willow. | 1576 |
| Group xviii. <i>Bicolores</i> Borrer. ✕ ☼ ✕ ✕ 1577
<i>Bushy Shrubs, with Leaves dark green above, and
glaucous beneath.</i> | 1577 |
| 129. <i>tennior</i> Borrer. ☼ Scotland 1577
The narrower-leaved intermediate Willow.
<i>S. laurina</i> Sm., according to Sm.
<i>S. bicolor</i> Sm. Eng. Bot., according to Sm. | 1577 |
| 130. <i>laxiflora</i> Borrer. ☼ Britain 1578
The loose-catkined Willow. | 1578 |
| 131. <i>laurina</i> Sm. ✕ Britain fig. 1338.,
and fig. 38, in p. 1612. - 1578
The Laurel-leaved, or shining dark green,
Willow.
<i>S. bicolor</i> Sm.
<i>S. arbuscula</i> Wahlenb. var. Koch
Comm. | 1338.,
1578
1578 |
| 132. <i>pátens</i> Forbes. ☼ fig. 39, in p. 1612.
1578
The spreading-branched Willow. | 1578 |
| 133. <i>radicans</i> Sm. ✕ Britain fig. 46, in
p. 1614. - - 1579
The rooting-branched Willow.
<i>S. phyticifolia</i> Ltn. Fl. Lap.
<i>S. arbuscula</i> Wahlenb. var. Koch
Comm. | 46, in
1579 |
| 134. <i>Borreriana</i> Sm. ☼ Scotland fig. 1339.
1579
Borrer's, or the dark upright, Willow. | 1339.
1579 |
| 135. <i>Davalliana</i> Sm. ☼ Scotland fig. 47,
in p. 1614. - - 1580
Davall's Willow.
<i>S. tetrapla</i> Walker.
<i>S. phyticifolia</i> Willd.
<i>S. thymelæoides</i> Schl.
<i>Variety</i> ☼ - - 1580
<i>S. Davalliana</i> Sm., the Swiss
kind. ☼ | 47,
1580
1580
1580
1580 |
| 136. <i>tétrapla</i> Sm. ☼ Scotland fig. 49, in
p. 1614. - - 1580
The four-ranked Willow. | 49, in
1580 |
| 137. <i>ramifúca</i> Forbes. ☼ Britain fig. 53.
in p. 1615. - - 1581
The brown-branched Willow.
? <i>S. tetrapla</i> mas Borrer. | 53.
1581 |
| 138. <i>Forbesiana</i> ☼ f. 51, in p. 1615. 1581
Forbes's Willow.
<i>S. Weigeliana</i> Forbes. | 1581 |
| 139. <i>Weigeliana</i> Borr. ☼ Scotland f. 1340.,
and f. 48, in p. 1614. - 1582
Weigel's Willow.
<i>S. Wulfcuiana</i> Sm.
<i>Variety</i> - - 1582
Leaves more conspicuously toothed,
rather silky when young; the
shoots more downy, and the
ovary pube scent towards the
point only. (<i>Borr.</i>) | 1340.,
1582
1582
1582 |
| 140. <i>nitens</i> Anders. ☼ Britain fig. 1341.,
and fig. 44, in p. 1613. - 1582
The glittering-leaved Willow. | 1341.,
1582 |
| 141. <i>Croweana</i> Sm. ☼ Scotland fig. 52.
in p. 1615. - - 1583
Crowe's Willow.
<i>S. arbuscula</i> Wahlenb. var. Koch
Comm.
<i>S. humilis</i> Schl., <i>sem.</i> of <i>S. Croweana</i>
Sm.
? <i>S. heterophylla</i> Host. | 52.
1583
1583 |
| 142. <i>bicolor</i> Ehrh. ☼ Britain fig. 54, in
p. 1615. - - 1583
The two-coloured Willow.
<i>S. tenuifolia</i> Sm.
<i>S. floribunda</i> Forbes.
<i>S. livida</i> Wahlenb. | 54, in
1583 |
| 143. <i>phillyreifolia</i> Borr. ☼ Scotland 1584
The Phytrea-leaved Willow. | 1584 |
| 144. <i>Dicksoniana</i> Sm. ✕ Scotland fig. 55,
in p. 1615. - - 1584
Dickson's Willow.
<i>S. nigriflora</i> Sm., not of L. | 55,
1584 |

- | | Page | | Page |
|------------------------------------------------------------|-----------------------------------------|--------------------------------------------------------------------------------------|-------------------------------------------------------|
| Group xix. <i>Vacciniifolia</i> Borr. ☼ ☼ ☼ | 1585 | 156. <i>retusa</i> L. ✕ Alps | fig. 1346., and fig. 139. in p. 1630. - 1589 |
| <i>Small, and generally procumbent, Shrubs.</i> | | The <i>retuse-leaved</i> Willow. | |
| 145. <i>vacciniifolia</i> Walker. ✕ Scotland | f. 1342., and f. 57. in p. 1615. 1585 | S. <i>retusa</i> Koch, part of, Koch Comm. | |
| The Vaccinium-leaved Willow. | | S. <i>serpyllifolia</i> Jacq. | |
| S. <i>prunifolia</i> , part of, Koch Comm. | | ? <i>Varieties</i> - - - 1589 | |
| 146. <i>carinata</i> Sm. ☼ Scotland | fig. 59. in p. 1615. - - - 1585 | ? S. <i>Kitaibeliana</i> Scop. | |
| The Keeled, or folded-leaved, Willow. | | ? S. <i>Uva-ursi</i> Scop. | |
| S. <i>prunifolia</i> , part of, Koch Comm. | | ? S. <i>serpyllifolia</i> Scop. | |
| 147. <i>prunifolia</i> Sm. ☼ Scotland | fig. 1615. 1585 | 157. <i>Kitaibeliana</i> Willd. ✕ Carpathian | Mountains f. 64. in p. 1616. 1589 |
| The Plum-leaved Willow. | | Kitaibel's Willow. | |
| S. <i>Myrsinites</i> Lightf., not of L. | | S. <i>retusa</i> Koch, β <i>major</i> Koch Comm. | |
| S. <i>prunifolia</i> , part of, Koch Comm. | | ? S. <i>Uva-ursi</i> Pursh. | |
| <i>Variety</i> ✕ - - - 1586 | | ? S. <i>retusa</i> var. L. | |
| <i>stylò longiøre</i> Koch. ☼ | | 158. <i>Uva-ursi</i> Pursh. ☼ Labrador | fig. 151. in p. 1630. - - - 1590 |
| S. <i>prunifolia</i> Ser. | | The Bearberry-leaved Willow. | |
| S. <i>formosa</i> Willd. | | ? S. <i>Kitaibeliana</i> Willd. | |
| S. <i>foetida</i> Schl. | | ? S. <i>retusa</i> var. L. | |
| S. <i>alpina</i> Sut. | | 159. <i>serpyllifolia</i> Scop. ✕ France, &c. | fig. 1347, 1348., and fig. 65. in p. 1616. - - - 1590 |
| 148. <i>venulosa</i> Sm. ☼ Scotland | fig. 56. in p. 1615. - - - 1586 | The Wild-Thyme-leaved Willow. | |
| The veiny-leaved Willow. | | S. <i>retusa</i> Koch, γ Koch Comm. | |
| S. <i>prunifolia</i> , part of, Koch Comm. | | S. <i>retusa</i> var. L. | |
| 149. <i>cæ'sia</i> Villars. ☼ Dauphiné | fig. 66. in p. 1616. - - - 1586 | 160. <i>cordifolia</i> Parsh. ✕ Labrador | fig. 143. in p. 1630. - - - 1590 |
| The grey-leaved Willow. | | The heart-leaved Labrador Willow. | |
| S. <i>myrtillodes</i> Willd. | | Group xxii. <i>Herbæcæ</i> Borrer. ☼ 1590 | |
| S. <i>prostrata</i> Ehrh. | | <i>Very low Shrubs, scarcely rising an Inch above the Ground.</i> | |
| Group xx. <i>Myrtillodes</i> Borrer. ☼ | 1587 | 161. <i>herbæcæ</i> L. ☼ Britain | fig. 1349., and fig. 62. in p. 1615. - 1590 |
| <i>Small Bilberry-like Shrubs, not Natives of Britain.</i> | | The herbaceous-looking Willow. | |
| 150. <i>myrtillodes</i> L. ☼ Poland, &c. | f. 1343. 1587 | ? <i>Varieties</i> - - - 1591 | |
| The Myrtillus-like, or Bilberry-leaved Willow. | | 162. <i>polâris</i> Wahlb. ☼ Lapland | f. 1350, 1351., and f. 63. in p. 1615. 1591 |
| S. <i>elegans</i> Besser. | | The Polar Willow. | |
| 151. <i>pedicellâris</i> Pursh. ☼ Catskill Moun- | tains - - - 1587 | Group xxiii. <i>Hastatæ</i> Borrer. ☼ ☼ ☼ | 1592 |
| The long-stalked-capsuled Willow. | | <i>Low Shrubs, with very broad Leaves, and exceedingly shaggy and silky Cutkins.</i> | |
| 152. <i>planifolia</i> Pursh. ☼ Labrador | 1587 | 163. <i>hastata</i> L. ☼ Lapland | fig. 1352., and fig. 35. in p. 1611. - 1592 |
| The flat-leaved Labrador Willow. | | The halberd-leaved Willow. | |
| Group xxi. <i>Myrsinites</i> Borrer. ☼ ✕ | 1587 | S. <i>hastata</i> Koch, part of. | |
| <i>Small bushy Shrubs.</i> | | <i>Varieties</i> ☼ - - - 1592 | |
| 153. <i>Myrsinites</i> L. ☼ Scotland | fig. 1344. 1588 | 2 <i>serrulata</i> ☼ | |
| The Whortleberry-leaved Willow. | | S. <i>hastata</i> Willd. | |
| S. <i>Myrsinites</i> β Sm. | | 3 <i>malifolia</i> ☼ | |
| S. <i>arbutifolia</i> Willd. | | S. <i>malifolia</i> Sm. | |
| ? S. <i>Macnabiana</i> Macgillivray. | | S. <i>hastata</i> , part of, Koch. | |
| S. <i>Myrsinites</i> Koch, part of, Koch Comm. | | S. <i>hastata</i> Hook., Borr. | |
| S. <i>dûbia</i> Suter. | | 4 <i>arbûscula</i> ☼ ? ✕ fig. 1353., and | fig. 138. in p. 1630. - 1592 |
| 154. <i>betulifolia</i> Forster. ☼ Britain | f. 1345., and f. 60. in p. 1615. - 1588 | S. <i>arbûscula</i> Wahl. | |
| The dwarf Birch-leaved Willow. | | S. <i>arbûscula</i> β Lin. Fl. Suec. | |
| S. <i>Myrsin. es</i> Sm. Koch, part of, Koch Comm. | | S. <i>arbûscula</i> γ Lin. Sp. Pl. | |
| 155. <i>procumbens</i> Forbes. ✕ Scotland | fig. 61. in p. 1615. - 1588 | | |
| The procumbent Willow. | | | |
| S. <i>le'pis</i> Hook. | | | |
| S. <i>retusa</i> Wither. | | | |

- | | Page | | Page |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|--------------------------------------------------------------|------|
| 164. lanata L. ♀ Lapland fig. 1354, and fig. 71. No. 2. in p. 1617 - 1593 | 1593 | 179. cerasifolia Schk. ♀ - - - 1597 | 1597 |
| The woolly-leaved Willow. | | The Cherry-leaved Willow. | |
| S. lanata, the kind No. 2., Forbes. | | 180. chrysanthos Ed. ♀ Finmark 1597 | 1597 |
| S. lanata, at least part of, Koch Comm. | | The golden-flowered Norway Willow. | |
| ? S. caprea Fl. Dan., t. 215 | | ? S. lanata var. | |
| ? S. chrysanthos Fl. Dan., t. 1057. | | 181. cinnamomea Schk. ♀ Switzerland 1597 | 1597 |
| Varieties, according to Koch | | The Cinnamon Willow. | |
| ♂ - - - 1594 | 1594 | 182. clethrafolia Schk. ♀ Switzerland 1597 | 1597 |
| 2 glabræscens ♀ | | The Clethra-leaved Willow. | |
| S. chrysanthos Vahl Fl. Dan. | | 183. conifera Wangeh. ♀ N. Amer. 1597 | 1597 |
| 3 glandulosa Wahlb. Fl. Lapp. ♀ | | The cone-bearing Willow. | |
| 4 depolita Koch. ? ♀ | | S. longirostris Micht. | |
| S. depressa Lin. Fl. Suec. | | 184. coruscans Willd. ♀ Styria - 1597 | 1597 |
| Group xxiv. <i>Miscellaneæ</i> A. ♀ ♂ - - - 1594 | 1594 | The glittering Willow. | |
| Kinds of <i>Salix</i> described in Sal. Wob., and not included in any of the preceding Groups. | | ? S. arbuscula Jacq., ? Wahlb. | |
| 165. ægyptiaca L. ♀ Egypt fig. 146. in p. 1630. - - - 1594 | 1594 | 185. cydoniaefolia Schk. ♂ Switzerland 1597 | 1597 |
| The Egyptian Willow. | | The Quince-leaved Willow. | |
| Café and Ban, Alpin. Egypt. | | 186. dubia Hort. ♂ - - - 1597 | 1597 |
| 166. alpina ? Forbes. ? ♀ ? ♂ ? - - - fig. 149. in p. 1630. - - - 1595 | 1595 | The doubtful Willow. | |
| The alpine Willow. | | 187. eriantha Schk. ♂ Switzerland 1597 | 1597 |
| 167. herberifolia Pall. ♀ Dauria fig. 1355., and fig. 140. in p. 1630. - 1595 | 1595 | The woolly-flowered Willow. | |
| The Herberry-leaved Willow. | | 188. fagifolia W. et K. ♂ Croatian Alps 1597 | 1597 |
| 168. tetrasperma Roxb. ♀ India fig. 31. in p. 1609. - - - 1595 | 1595 | The Beech-leaved Willow. | |
| The four-seeded Willow. | | 189. finmarchica Lodd. Cat. ♀ Sweden 1598 | 1598 |
| 169. ulmifolia Forbes. ♀ Switzerland 1595 | 1595 | The Finmark Willow. | |
| The Elm-leaved Willow, or Sallow. | | 190. foliolosa Afzel. ♂ Lapland - 1598 | 1598 |
| 170. villosa Forbes. ♂ Switzerland fig. 92. in p. 1619. - - - 1595 | 1595 | The many-leaved Willow. | |
| The villous-leaved Willow. | | ? S. foliosa Loud. Hort. Brit. | |
| Group xxv. <i>Miscellaneæ</i> B. ♀ ♂ - - - 1596 | 1596 | S. alpina myrifolia Rudl. | |
| Kinds of <i>Salix</i> introduced, and of many of which there are Plants at Messrs. Loddiges's, but which we have not been able to refer to any of the preceding Groups. | | S. arbuscula β L. | |
| 171. albescens Schk. ♀ Switzerland 1596 | 1596 | 191. formosa Willd. ♀ Swiss Alps 1598 | 1598 |
| The whitish-leaved Willow. | | The elegant Willow. | |
| S. stylosa & Dec. | | S. glauca Willd. | |
| 172. alnifolia Host. ♀ - - - 1596 | 1596 | ? S. alpina Scop. | |
| The Alder-leaved Willow. | | 192. fuscata Pursh. ♂ New York - 1598 | 1598 |
| 173. Ammanniana Willd. ♀ Salzburg 1596 | 1596 | The brown-stemmed Willow. | |
| Ammann's Willow. | | 193. glabrata Schk. ♀ Switzerland - 1598 | 1598 |
| S. Marsinate Hoffm. | | The glabrous Willow. | |
| S. hirtata Hopp., not of L. | | 194. heterophylla Deb. ♀ Europe 1598 | 1598 |
| 174. angustata Pursh. ♀ New York 1596 | 1596 | The various-leaved Willow. | |
| The tapered leaved American Willow. | | 195. humilis Dec. ♂ - - - 1598 | 1598 |
| 175. angustifolia Willd. ♀ Near the Caspian Sea - - - 1596 | 1596 | The humble Willow. | |
| The narrow leaved Caspian Willow | | 196. Jacquini Host. ♀ Alps - 1598 | 1598 |
| 176. betulina Host. ? ♀ - - - 1596 | 1596 | Jacquin's Willow. | |
| The Birch-like Willow. | | S. fusca Jacq. | |
| 177. candidula Host. ? ♀ - - - 1596 | 1596 | S. alpina Scop. | |
| The whitish Willow | | S. Jacquiniina Willd. | |
| 178. canescens Lodd. ? ♀ - - - 1596 | 1596 | 197. livida Wahlb. ♀ - - - 1598 | 1598 |
| The hoary Willow | | The livid-leaved Willow | |
| | | S. arbuscula γ Lin. Fl. Suec. | |
| | | ? S. arbuscula β Lin. Sp. Pl. | |
| | | S. straboana Willd. | |
| | | ? S. foliolosa Afzel. | |
| | | S. multiflora Bess. | |
| | | S. bicolor Ehrh. | |
| | | S. livida Hook. | |
| | | 198. longifolia Mühlb. ♂ Banks of the Susquehanna - - - 1599 | 1599 |
| | | The long-leaved Willow. | |
| | | 199. mespilifolia Schk. ♀ Switzerland 1599 | 1599 |
| | | The Mespilus-leaved Willow. | |
| | | 200. murina Schk. ♀ Switzerland - 1599 | 1599 |
| | | The Mouse Willow. | |
| | | 201. myricoides Mühlb. ♀ N. Am. 1599 | 1599 |
| | | The Myrica-like Willow. | |

	Page
202. <i>nervosa</i> Schl. ☼ Switzerland	1599
The <i>nerved-leaved</i> Willow.	
203. <i>obtusa</i> Link. ☼ Switzerland	1599
The <i>blunt-leaved</i> Willow.	
204. <i>obtusifolia</i> Willd. ☼ Lapland	1599
The <i>obtusely-veined</i> Lapland Willow.	
S. <i>foliis oblongis</i> , &c., Lin. Fl. Lapp.	
S. <i>caprea</i> β Lin. Sp. Pl.	
S. <i>Olea sylvestris</i> , &c., Rudd.	
205. <i>obtusiserratis</i> Schl. ☼ Switzerland	1599
The <i>obtusely-serrated-leaved</i> Willow	
206. <i>pallascens</i> Schl. ☼ Switzerland	1599
The <i>pale</i> Willow.	
207. <i>paludosa</i> Link. ☼ -	1600
The <i>Marsh</i> Willow.	
208. <i>persicaefolia</i> Hort. ☼ -	1600
The <i>Peach-tree-leaved</i> Willow.	
209. <i>pyrenaica</i> Gouan. ✕ Pyrenees	1600
The <i>Pyrenean</i> Willow.	
210. <i>pyrifolia</i> Schl. ☼ Switzerland	1600
The <i>Pear-tree-leaved</i> Willow.	
211. <i>recurvata</i> Pursh. ☼ N. America.	1600
The <i>recurved-cathinet</i> Willow.	
212. <i>salviaefolia</i> Lk. ☼ Portugal	1600
The <i>Sage-leaved</i> Willow.	
S. <i>pyralis</i> Seringe Sal. Helv.	
S. <i>oleifolia</i> Ser. Sal. exsicc.	
S. <i>oleifolia</i> Vill.	
S. <i>Fluggiana</i> Willd.	
213. <i>Schraderiana</i> Willd. ✕ -	1600
Schrader's Willow.	
S. <i>dicolor</i> Schrad.	
214. <i>septentrionalis</i> Host. ☼ -	1600
The <i>northern</i> Willow.	
215. <i>silesiaca</i> Willd. ☼ Sweden, &c.	1601
The <i>Silesian</i> Willow.	
216. <i>Starkeana</i> Willd. ☼ Silesia	1601
Starke's <i>Sallow</i> , or the <i>Marsh Silesian</i> Willow.	
217. <i>tetrandra</i> Host. ☼ -	1601
The <i>four-stamened</i> Willow.	
218. <i>thymelæoides</i> Host. ✕ -	1601
The <i>Wild-Olive-like</i> Willow.	
219. <i>Trevirana</i> Lk. ☼ -	1601
Treviranus's Willow.	
220. <i>velutina</i> Willd. ☼ -	1601
The <i>velvet</i> Willow.	
221. <i>versifolia</i> Spreng. ✕ -	1601
The <i>twining-leaved</i> Willow.	
222. <i>vaccinioides</i> Host. ✕ -	1601
The <i>Vaccinium-like</i> Willow.	
223. <i>Waldsteiniانا</i> Willd. ☼ Croatian Alps	1601
Waldstein's Willow.	
S. <i>alpestris</i> Host.	
224. <i>Wulfeniana</i> Willd. ☼ Carinthia	1601
Wulfen's Willow.	
S. <i>hastata</i> var. Koch Comm.	
? S. <i>phylicifolia</i> Wulf.	

App. i. *Kinds of Salix described or recorded in Botanical Works, but not introduced into Britain, or not known by these Names in British Gardens.* - - 1602

☼ *arctica* R. Br. ☼ North America.
 ☼ *desertorum* Rich. ☼

	Page
S. <i>rostrata</i> Rich. ☼	1599
S. <i>cinerascens</i> Link M.S.S. ☼ Portugal.	
The <i>ash-coloured Portuguese</i> Willow.	
? S. <i>grandifolia</i> Ser.	
S. <i>grandifolia</i> Ser. Sal. Helv. ☼	1599
S. <i>stipularis</i> Ser. Sal. exsicc.	
? S. <i>cinerascens</i> Link.	
S. <i>divaricata</i> Pall. ✕ Dauria.	1599
S. <i>hirsuta</i> Thunb. ☼ Cape of Good Hope.	1599
S. <i>pedicellata</i> Desf. ? ♀ Barbary.	1599
The <i>stalked Barbary</i> Willow.	
S. <i>integra</i> Thunb. ☼ Japan.	1599
? The <i>entire-leaved</i> <i>Japan</i> Willow.	
S. <i>japonica</i> Thunb. ♀ Japan.	1599
S. <i>Rin. vulgo Ajuki</i> , Kämpfer.	
S. <i>mucronata</i> Thunb. ☼ Cape of Good Hope.	1599
S. <i>rhannifolia</i> Pall. ✕ Siberia.	1599
S. <i>pumila</i> , <i>foliis ovalibus</i> , &c., Gmel.	
S. <i>Seringiana</i> Gaulin in Ser. Sal. Helv. ☼ Switzerland.	1599
S. <i>laevigata</i> Ser. Sal. exsicc., No. 70.	
S. <i>Konderiana</i> Ser. Sal. exsicc., No. 12.	
S. <i>longifolia</i> Schl. Cal.	
S. <i>serotina</i> Pall. ☼ ♀ Wolga.	1599
S. <i>caprea</i> y Lin. Sp. Pl.	
S. <i>oxytraca</i> Willd.	
S. <i>Gmeliniana</i> Willd. Sp. Pl.	

App. ii. *Kinds of Salix figured or described in the Salicetum Woburnense.* - 1631

App. iii. *Koch's Arrangement of the Species of Salix indigenous to Europe; including, also, some extra-European Species, with references to the pages in this work, where each species is described.* - - 1633

App. iv. *Kinds of Salix described in Host's Flora Austriaca, and figured in Host's Salix.* - - - 1635

II. *POPULUS* Tourn. ♀ 1454. 1636

THE *POPLAR.*

- Peuplier*, Fr.
- Pappel*, Ger.
- Pioppo*, Ital.
- Poplicr*, Dutch.
- Alamo*, Span.

1. *alba* L. ♀ Europe pl. 262. f. 1507. 1638

The *white Poplar*, or *Ablec Tree*.

- P. *alba latifolia* Lob.
- Populus* No. 1634. Hall. Hist.
- P. *major* Mill.
- P. *nivea* Willd.
- P. *alba nivea* Mart.
- Leuké* Dioscorides.
- The *great white Poplar*, *great Aspen*, *Dutch Beech*.
- Peuplier blanc*, *Ypreiau blanc de Hollande*, *Frauc Picard*, Fr.; *Aubo*, or *Aoub. ro.* in some Provinces.
- Weisse Pappel*, *Silber Pappel*, *Weisse Aspe*, *Weissalber Baum*, Ger.
- Abelboom*, Dutch.

Varieties - - 1639

2. (a.) *canescens* Sm. ♀ Britain pl. 268. fig. 1508. 1639

The *grey*, or *common white*, *Poplar*.

- P. *alba* Mill.
- P. *alba foliis minutis* Ray.
- P. *alba foliis minore* Bauth.
- P. No. 1634 β Hall. Hist.
- Peuplier grisaille*, Fr.

Varieties ♀ - - 1640

2 *hybrida* Bieb. Fl. Tour.

- P. *alba* Bieb. l. c.
- ? P. *intermedia* Mertens.
- ? P. *a. crassifolia* Mertens.
- P. *grisea* Lodd. Cat.

- 3 *acerifolia* Y
P. acerifolia Lodd. Cat.
P. quercifolia Hort.
P. palmata Hort.
- 4 *aremburgica* Y
P. aremburgica Lodd. Cat.
- 5 *belgica* Y
P. belgica Lodd. Cat.
- 6 *candicans* Y
P. candicans Lodd. Cat.
- 7 *nivea* Y
P. nivea Lodd. Cat.
- 8 *egyptiaca* Hort. Y
P. a. pallida Hort.
Other Varieties.
- 9 *pandula* Y
P. a. var. gracilis ramis pendulibus Mertens.
3. *trémula* L. Y Europe
 pl. 26 t. fig. 1509. 16 15
 The trembling-leaved Poplar, or *Aspen*.
P. No. 1633. Hall. Hist.
P. libyca Ray.
P. hybrida Dod.
P. nigra Traut.
P. pendula Du Roi.
Le Tremble, Fr.
La Tremula, Alberatta, Alberetto, Ital.
Zitter Pappel, Espe, Ger.
- Varieties* Y - - 1509
- 1 *monticola* Y
P. monticola Mertens.
 ? *P. trémula* L.
- 2 *parvifolia* Mertens. Y
- 3 *grandifolia* Mertens. Y
- 4 *rotundifolia* major Mertens. Y
- 5 *minor* Mertens. Y
- 6 *oxyodonta* Y
P. oxyodonta Mertens.
- 7 *stricta* Y
P. stricta Mertens.
- 8 *pendula* Y pl. 265.
P. pendula Lodd. Cat.
- 9 *supina* Y
S. supina Lodd. Cat.
- 10 *lavigata* Y
P. lavigata Ait.
4. (t.) *trépida* Willd. Y North America
 fig. 1510. 16 19
 The North American trembling-leaved Poplar,
 or American *Aspen*.
P. tremuloides Michx.
5. (l.) *grandidentata* Michx. Y Canada
 1651
 The large-toothed-leaved Poplar, or North American large *Aspen*.
- Variety* Y - - 1651
- 2 *pendula* Michx. Y
6. *græca* Ait. Y Greece
 pl. 266. fig. 1512. 1651
 The Grecian, or Athenian, Poplar.
7. *nigra* L. Y Europe pl. 267, 268, 269.
 fig. 1513, 1511. 1652
 The black-barked, or common black, Poplar.
P. No. 1632. Hall. Hist.
P. alba Traut.
P. romanica Du Ham.
Agnetos, Greek.
Kabaki, Modern Greek.

- The old English Poplar*, Suffolk.
The Willow Poplar, Cambridgeshire.
Water Poplar.
Cotton Tree, fem. of *P. nigra* so called at Bury St. Edmunds.
Peuplier noir, Peuplier hard, Osier Blanc, Fr.
Schwarze Pappel, Ger.
- Varieties* Y - - 1652
- 2 *viridis* Lindl. Y
P. viridis Lodd. Cat.
- 3 *salicifolia* Y
P. salicifolia Lodd. Cat.
8. (n.) *canadensis* Michx. Y N. America
 fig. 1515. 1655
 The Canadian Poplar.
P. lavigata Willd., not of Hort. Kew.
P. monilifera Hort. Par.
Cotton Wood, Michx.
Peuplier de Canada, Fr.
9. (n.) *betulifolia* Pursh. Y Banks of the Hudson - fig. 1516. 1656
 The Birch-leaved Poplar.
P. nigra Michx. Fl.
P. hudsonica Michx. Arb.
P. hudsoniana Bosc.
American Black Poplar, Amer.
Peuplier de la Baie d'Hudson, Fr.
10. *monilifera* Ait. Y ? Canada
 pl. 270, 271, 272. fig. 1517. 1657
 The Necklace-bearing, or black Italian, Poplar.
P. virginiana L.
P. glandulosa Mench Meth.
P. carolinensis Mench Weissenst.
P. nigra italica Lodd. Cat.
P. nigra americana Lodd. Cat.
P. acledusa Lindl.
 ? *P. marylandica* Bosc.
Virginian Poplar, Swiss Poplar, Canadian, or Berry-bearing, Poplar, Mill.
Peuplier Suisse, Peuplier triphylon, Peuplier de Virginie, Dumont.
- Varieties* Y - - 1657
- 2 *Lindleyana* Booth. Y
 The new wave-leaved Poplar,
 Hort.
- 3 *foliis variegatis* Hort. Y
11. *fastigiata* Y Italy
 pl. 273. fig. 1519, 1520. 1660
 The fastigate, or Lombardy, Poplar.
P. dilatata Ait.
P. nigra italica Du Roi.
P. italica Mench Weissenst.
P. italica dilatata Willd.
P. pyramidata Hort.
P. pannonica Jacq.
P. italica var. carolinensis Burgsdorf.
Cypress Poplar, Turin Poplar, Po Poplar, Peuplier d'Italie, Peuplier pyramidale, Fr.
Lombardische Pappel, Italianische Pappel, Ger.
Pioppo Cypressino, Ital.
12. *angulata* Ait. Y North America
 pl. 274, 275. fig. 1533. 1670
 The angled-branched, or Carolina, Poplar.
P. angulosa Michx.
P. heterophylla Du Roi.
P. macrophylla Lodd. Cat.
P. balsamifera Mill.
Mississippi Cotton Tree, Amer
- Varieties* Y - - 1671
- 2 *nova* Audibert Y
- 3 *Medusa* Booth. Y

13. heterophýlla L. \checkmark N. Am. f. 1534. 1672
 The various-shaped-leaved Poplar Tree.
P. magna, foliis amplis, albis cordiformibus, albis subrotundis, primoribus tomentosis, Gron.
P. cordifolia Burgsdorf.
P. argentea Michx.
Cotton Tree, Michx. N. A. S.

14. balsamifera L. \checkmark North America
 pl. 276. fig. 1535, 1536. 1673
 The balsam-bearing, Poplar, or Tacamahac Tree.
P. Tacamahaca Mill.
The Tacamahac, Amer.
Le Baumier, Fr.
Peuplier liard, Tacamahac, in Canada.
Balsam Pappel, Ger.

Varieties \checkmark - - 1673
 2 viminalis \checkmark Altai.
P. viminalis Lodd. Cat.
P. salicifolia Hort.
P. longifolia Fischer.
 3 latifolia Hort. \checkmark
 4 intermedia Hort. \checkmark Dahuria
 5 suaveolens \checkmark
P. suaveolens Fischer, and Lodd. Cat.
 6 foliis variegatis Mill. \checkmark

15. canadensis Mill. \checkmark North America
 pl. 277. fig. 1557. 1676
 The whitish-leaved balsam-bearing, or Ontario, Poplar.
P. macrophylla Lindl.
P. latifolia Mönch Meth.
P. ontariensis Desf.
P. cordata Lodd. Cat.
P. canadensis Mönch Weissenst., not of Michx.
Balm of Gilead Tree, Boston.
Peuplier liard, Canada.
Peuplier à Feuilles vernissées, Fr.

Betulaceæ. \checkmark ☉ - 1677

I. ALNUS Tourn. \checkmark ☉ - - 1677
 THE ALDER.
Bétule Species L.
Aunc, Fr.
Erle, Ger.
Ontano, Ital.
Aliso, Span.

1. glutinosa Gerbn. \checkmark Europe
 pl. 278. fig. 1538. 1678
 The glutinous, or common, Alder.
Bétulus Alnus L.
Bétula emarginata Ehrh.
Alnus Ray.
Aunc, Fr.
Geveine Elsc, Elscr, Schwartz Erle, Ger.
Eiscboom, Dutch.
Alno, Ontano, Ital.
Aliso, Alamo nigro, Span.

Varieties \checkmark - - 1678
 2 emarginata Willd. \checkmark
 3 laciniata Willd. \checkmark
A. g. incisa Hort. fig. 1538.
 4 quercifolia Willd. \checkmark
 5 oxycanthæfolia \checkmark fig. 1539.
A. oxycanthæfolia Lodd. Cat.
 6 macrocarpa \checkmark
A. macrocarpa Lodd. Cat.
 7 foliis variegatis Hort. \checkmark
 Other Varieties.

2. (g.) oblongata Willd. ☉ \checkmark Hungary
 1687

The oblong-leaved Alder.
Alnus fol. oblong., ☉ ., Bauh.
A. fol. ovato-lanceol., ☉ ., Mill.
Langliche Elsc, Ger.
Variety ☉ \checkmark - - 1687
 2 foliis ellipticis Ait. ☉ \checkmark
A. pinnata Lodd. Cat.

3. incana Willd. \checkmark N. Am. f. 1543. 1687

The hoary-leaved Alder.
Bétula Alnus var. incana L. Sp. Pl.
Bétula incana L. Supp.
A. folio incano., ☉ ., Bauh.
Bétula viridis Vill.
Weisse Erle, Graue Elsc, Weisse Eller, Ger.

Varieties \checkmark - - 1688
 2 laciniata Lodd. Cat. \checkmark
 3 glauca \checkmark
A. glauca Michx.
Bétula incana var. glauca Ait.
Black Alder, Amer.
 4 angulata Ait. \checkmark
Other Varieties.
 ? 1.
A. americana Lodd. Cat.
 ? 2.
A. canadensis Lodd. Cat.
 ? 3.
A. rubra Lodd. Cat.

4. serrulata Willd. ☉ North America
 fig. 1544. 1688

The saw-leaved Alder.
Bétula serrulata Ait.
Bétula rugosa Ehrh.
 ? *A. americana* Lodd. Cat.
 ? *A. canadensis* Lodd. Cat.
Common Alder, Amer.
Hazel-leaved Alder.

5. undulata Willd. ☉ Canada - 1689

The waved-leaved Alder.
Bétula crispa Ait.
Bétula Alnus var. crispa Michx.
A. crispa Pursh.

6. cordifolia Lodd. \checkmark Calabria
 pl. 281. fig. 1545. 1689

The heart-leaved Alder.
A. cordata Tenore.

7. viridis Dec. ☉ Hungary f. 1546. 168

The green-leaved Alder.
A. ovata Lodd. Bot. Cab.
A. fruticosa Schmidt.
Bétula ovata Schrank.
Bétula Alno-Bétule Ehrh.
Bétula viridis Hort.

App. i. Other Species of Alnus. 1690

A. obtusifolia Royle. Banks of the Jumna.
A. elongata Royle. Cashmere.
A. nepalensis Royle. Nepal.

II. BETULA Tourn. ☉ \checkmark - 1690

THE BIRCH.
Bouleau, Fr.
Betula, Ital.
Abedul, Span.
Betulla, Port.
Birke, Ger.
Berk, Dutch.
Birk, Danish and Scotch.
Björk, Bork, Swedish.
Bereza, Russian.
Brzoza, Polish.

	Page
<i>Leaves small. Natives chiefly of Europe.</i>	
1. <i>alba</i> L. † Europe † 1517. 1550. 1691	
The white, or common, Birch.	
B. <i>pubescens</i> Ehrh.	
B. <i>betula</i> Kay.	
B. <i>artensis</i> Rafin.	
Bouleau commun, Fr.	
Gewöhnliche Birke, Gr.	
Varieties † - - - 1691	
2 <i>péndula</i> Sm. † pl. 282.	
The weeping Birch.	
B. <i>péndula</i> Roth.	
B. <i>verrucosa</i> Ehrh.	
B. <i>péndula virgulus</i> Loes.	
3 <i>pubescens</i> † fig. 1548.	
B. <i>pubescens</i> Ehrh.	
4 <i>póntica</i> † fig. 1549.	
B. <i>póntica</i> Lodd. Cat.	
5 <i>articifolia</i> †	
B. <i>articifolia</i> Lodd. Cat.	
6 <i>dalecarlica</i> L. Supp. †	
7 <i>macrocarpa</i> Willd. †	
8 <i>foliis variegatis Dumont.</i> †	
Other Varieties.	
2. <i>dáurica</i> Pall. † † Dauria † 1556. 1704	
The Daurian Birch.	
B. <i>excelsa canadensis</i> Wang.	
Bouleau de Sibérie, Fr.	
Variety † † - - - 1705	
2 <i>parvifolia</i> † †	
3. <i>fruticosa</i> Pall. † † Eastern Siberia, &c.	
fig. 1557. 1705	
The shrubby Birch.	
B. <i>humilis</i> Schrank.	
B. <i>quebecensis</i> Schrift. der Ges. Naturf.	
4. <i>pùmila</i> L. † † Canada fig. 1558. 1705	
The hairy dwarf Birch.	
B. <i>nana</i> Kalm.	
5 <i>nàna</i> L. † † Lapland, &c. f. 1559. 1705	
The dwarf Birch.	
B. <i>nana Succorum</i> Bromel.	
B. No. 259. Amm. Ruth.	
B. <i>palustris pùmila</i> , &c., Cels.	
Variety † - - - 1706	
2 <i>stricta</i> Lodd. Cat. †	
6. <i>glandulosa</i> Michx. † † Canada - 1707	
The glandular-branched Birch.	
<i>Leaves large. Natives of North America.</i>	
7. (a.) <i>populifolia</i> Mill. † † North America	
pl. 283. fig. 1560. 1707	
The Poplar-leaved Birch.	
B. <i>acuminata</i> Ehrh.	
B. <i>lenta</i> Du Roi.	
White Birch and Oldfield Birch. Amer.	
Varieties † - - - 1707	
2 <i>laciniata</i> †	
B. <i>laciniata</i> Lodd. Cat.	
3 <i>péndula</i> †	
B. <i>péndula</i> Lodd. Cat.	
8. <i>papyracea</i> Mill. † † North America	
pl. 284. fig. 1561. 1708	
The Paper Birch.	
B. <i>papyrifera</i> Michx.	
B. <i>lancolata</i> Hort.	
B. <i>cubra</i> Lodd. Cat.	
B. <i>canadensis</i> Lodd. Cat.	
B. <i>nigra</i> of the Paris Nursery	
Canoe Birch, White Birch Amer.	

	Page
<i>Varieties † - - -</i>	
2 <i>fusca</i> † Carolina.	1708
B. <i>fusca</i> Bosc.	
3 <i>trichoclada</i> Hort. †	
4 <i>platyphylla</i> Hort. †	
9. <i>nigra</i> L. † † North America	
pl. 285, 286. fig. 1562, 1563. 1710	
The black Birch.	
B. <i>lanulosa</i> Michx. Fl.	
? B. <i>cubra</i> Michx. Arb.	
B. <i>angulata</i> Lodd. Cat.	
Red Birch, Amer.	
10. <i>excelsa</i> H. Kew. † † North America	
fig. 1564, 1565. 1711	
The tall Birch.	
B. <i>lutea</i> Michx.	
? B. <i>nigra</i> Du Roi.	
Yellow Birch, Amer.	
11. <i>lenta</i> L. † † N. Amer. † 1566. 1713	
The pliant Birch.	
B. <i>carpinifolia</i> Ehrh.	
B. <i>nigra</i> Du Roi.	
Black Birch, Cherry Birch, Canada Birch.	
Sweet Birch, Mountain Mahogany, Amer.	
Bouleau Méricier, Fr.	
App. i. <i>Species of Birch not yet introduced.</i>	1714
B. <i>rhizophylla</i> Wall. Kamaon.	
B. <i>acuminata</i> Wall. Nepal.	
B. <i>nitida</i> Kamaon.	
B. <i>cyathostachya</i> Kamaon.	
B. <i>resinifera</i> Kunawar.	

Corylaceæ, or Cupuliferae. 1715

† † † † †

I. *QUERCUS* L. † † † † † 1715. 1717

THE OAK.

Mer Tourn.
Süder Tourn.
Dere, Celtic.
Aauck, Ac. Saxon.
Al, Alou, Allan, Hebrew.
Drus, Greek.
Chêne, Fr.
Eiche, Ger.
Eik, Dutch.
Quercia, Ital.
Encina, Span.

A. *Leaves decutuous.*

§ i. *Röbur.* British Oaks. † 1730, 1731

1. *pedunculata* Willd. † † Britain

pl. 287, 288, 289. fig. 1567. 1581, 1582. - - - 1731. 1710

The common, or peduncled, British Oak.

Q. *Röbur* L.
Q. R. *pedunculatum* Mart.
Q. *faucina* Roth.
Q. *racemosa* S. Du Ham.
Q. *emula longa pedunculo* Bauh.
Q. *hincris* Dalech.
Quercus Fuchs.
Q. *navalis* Burnet.
Chêne blanc, Secondat.
Chêne à Grappes, Chêne femelle, Gravelin Fr.
Stiel Eiche, Irub Eiche, That Eiche, Loh Eiche, Wald Eiche, Ger.

	Page
<i>Varieties</i> † -	1731
2 pubescens Lodd. Cat. †	
3 fastigiata † pl. 290.	
<i>Q. fastigiata</i> Lam. Dict.	
<i>Q. pyramidatis</i> Hort.	
<i>Chêne Cyprés, Chêne des Pyrénées,</i>	
<i>Fr.</i>	
4 péndula † fig. 1568.	
The weeping Oak.	
<i>Q. péndula</i> Lodd. Cat.	
5 heterophylla † fig. 1560. 1570.	
<i>Q. salicifolia</i> Hort.	
<i>Q. laciniata</i> Lodd. Cat.	
<i>Q. filicifolia</i> Hort.	
<i>Q. Fennesi</i> Hort.	
6 foliis variegatis Lodd. Cat. †	
7 purpurea †	
<i>Q. purpurea</i> Lodd. Cat.	
8 Hodginsii Lodd. Cat. †	
9 dulcis. †	
<i>Chêne à Feuilles caduques, presque sessiles,</i>	
<i>Dralet.</i>	
Other Varieties.	
2. sessiliflora Sal. † Britain pl. 291, 292,	
293. fig. 1572, 1585, 1586 - 1736	
The sessile-flowered Oak.	
<i>Q. Robur</i> Willd.	
<i>Q. R. var. sessile</i> Mart.	
<i>Q. sessilis</i> Ehrh.	
<i>Q. platyphyllos, mas et firm.,</i> Dalech.	
<i>Q. latifolia mas, &c.,</i> Bauh.	
<i>Q. regalis</i> Burnet.	
<i>Chêne mâle, Secondat.</i>	
<i>Chestnut Oak, Bay Oak.</i>	
<i>Chêne rouge or rouvre, Durclin, Fr.</i>	
<i>Steine Eiche, Gemeine Eiche, Spät Eiche,</i>	
<i>Winter Eiche, Durr Eiche, Roth Eiche,</i>	
<i>Berg Eiche, Ger.</i>	
<i>Quercia vera, Ital.</i>	
<i>Roble, Span.</i>	
<i>Varieties</i> † -	1736
2 pubescens † fig. 1573.	
<i>Q. s. var. β</i> Sm.	
<i>Q. pubescens</i> Willd.	
<i>Q. R. lanuginosum</i> Lam. Dict.	
<i>The Darmast, Mart. Fl. Rust.</i>	
Other Varieties -	1737
1 Le Chêne à Trochets, or Chêne	
à petits Glands, <i>Bosc.</i>	
2 Le Chêne à Feuilles découpées,	
<i>Bosc.</i>	
3 Le Chêne laineux, or Chêne des	
Collines, <i>Bosc.</i>	
4 Le Chêne noirâtre, <i>Bosc.</i>	
Mr. Bree's Varieties. -	1738
3. pyrenaica Willd. † Pyrenees	
fig. 1696. 1842	
The Pyrenean Oak.	
<i>Q. Tauzin Pers.</i>	
<i>Q. nigra</i> Thore.	
<i>Q. Tosa Bosc.</i>	
<i>Q. stolonifera</i> Lapeyr.	
<i>Chêne noir, Secondat.</i>	
<i>Varieties</i> -	1844
1 With large acorns, on pedun-	
cles, axillary and terminal,	
<i>N. Du Ham.</i>	
2 With axillary acorns of a middle	
size, <i>N. Du Ham.</i>	

	Page
3 With small acorns, on long ra-	
cemes, <i>N. Du Ham.</i>	
<i>Q. Tauzin laciniata Desvaur.</i>	
<i>Q. Tauzin digitata Desvaur.</i>	
4. apennina Lam. † Lyons fig. 1698. 1844	
The Apennine Oak.	
<i>Q. congloubrata Pers.</i>	
<i>Chêne hirsut, Fr.</i>	
5. E'sculus L. † South of Europe	
pl. 294. fig. 1699, 1670. 1844	
The Esculus, or Italian, Oak.	
<i>Phagus E'sculus mas et firm.</i> Dalech.	
<i>Chêne grec, Fr.</i>	
<i>Varieties</i> -	1845
§ ii. Cérris. Mossy-cupped, or Turkey, Oaks.	
† †	1730. 1846
6. Cérris L. † France	
pl. 295, 296, 297. fig. 1702. 1846	
The bitter, or mossy-cupped, Oak.	
<i>Q. crinita α and β</i> Lam. Dict.	
<i>Q. Haliphlaeos</i> Juss.	
<i>Q. burgundica, &c.,</i> Bauh.	
<i>Q. Cérris Phnii, &c.,</i> Lob.	
<i>Cérrus</i> Dalech.	
<i>The Turkey Oak, the Iron or Wainscot</i>	
<i>Oak.</i>	
<i>Chêne Cérris, Chêne de Bourgogne, Fr.</i>	
<i>Burgundische Eiche, Cerr-eiche, Ger.</i>	
<i>Varieties</i> † -	1847
* <i>Foliage deciduous.</i>	
a. <i>Leaves pinnatifid or sinuately. Caps</i>	
<i>of the Acorns mossy.</i>	
1 vulgaris † fig. 1702.	
<i>Q. C. frondosa</i> Mill.	
<i>Subvarieties</i> †	
See fig. 1703, 1704, 1705, and	
1706. in p. 1846, 1847.	
2 péndula Neill. † fig. 1707.	
The pendulous, or weeping, Turkey	
Oak.	
3 variegata Lodd. Cat. †	
b. <i>Leaves dentate. Caps of the Acorns</i>	
<i>bristly.</i>	
4 austriaca † fig. 1708. pl. 298.	
<i>Q. austriaca</i> Willd.	
<i>Q. Cérris</i> Host, α and β No. 28.	
<i>Q. crinita γ Cérris</i> L.	
<i>Q. calyce hispido, &c.,</i> Bauh.	
<i>Cérrus</i> Chus.	
<i>Cérris minoris ramulus cum flore,</i>	
<i>Ger.</i>	
<i>Cérris Phnii minore glande</i> Lob.	
<i>E'gilops minore glande</i> Dod.	
<i>Haliphlaeos Cérris femina</i> Da-	
<i>lech.</i>	
5 cæna major † fig. 1709.	
<i>Q. cæna major</i> Lodd. Cat.	
6 cæna minor †	
<i>Q. cæna minor</i> Lodd. Cat.	
7 Râgnal †	
The Râgnal Oak.	
<i>Q. Râgnal</i> Lodd. Cat.	
** <i>Foliage subevergreen. Leaves dentate.</i>	
<i>Acorns with bristly Caps.</i>	
8 fulhamensis † pl. 299, 300,	
301. fig. 1710, 1711.	
The Fulham Oak.	
<i>Q. C. dentata</i> Wats.	
<i>Q. C. hybrida var. dentata</i> Swt	

- | | Page |
|---------------------------------------------------------------------------------------------------------------|----------------|
| 9 <i>Lucombecina</i> Y pl. 300. 303, 304, 305. fig. 1712, 1713, 1714. | 303, |
| The Lucombe Oak. | |
| Q. <i>Lucombecina</i> Swt. | |
| Q. <i>ezoniensis</i> Lodd. Cat. | |
| The Evergreen Turkey Oak, the Devonshire Oak, the Exeter Oak. | |
| *** Foliage evergreen, or very nearly so. Leaves varying from dentate to sinuate. Cups of the Acorns bristly. | |
| 10 L. <i>crispa</i> I pl. 306. fig. 1715. 1717. c. 1718. | 1715. |
| The new Lucombe Oak. | |
| Q. <i>Lucombecina crispa</i> Hort. | |
| 11 L. <i>suberosa</i> I fig. 1717a. | 1717a. |
| Q. L. <i>suberosa</i> Hort. | |
| 12 L. <i>incisa</i> I fig. 1717. b. | 1717. b. |
| Q. L. <i>incisa</i> Hort. | |
| 13 L. <i>dentata</i> I fig. 1716. | 1716. |
| Q. L. <i>dentata</i> Hort. | |
| 14 heterophylla I fig. 1719. | 1719. |
| Q. L. <i>heterophylla</i> Hort. | |
| Other Varieties. | |
| Q. C. <i>bullata</i> . | |
| The blistered, or rough-leaved, Turkey Oak. | |
| Q. C. <i>dentata pendula</i> . | |
| 7. <i>Ægilops</i> L. Y Greece, &c. pl. 307, 308. fig. 1721. 1861 | 1861 |
| The Ægilops, or Falonia, Oak. | |
| Q. <i>orientalis</i> , &c., Tourm. | |
| <i>Ægilops</i> sive <i>Cerrus</i> más C. Bauh. Velani Tourm. | |
| <i>Gilans Cérri</i> Dalech. | |
| The Great Prickly-cupped Oak. | |
| <i>Chêne Velani</i> , Fr. | |
| <i>Chêne Velanide</i> , Bose. | |
| <i>Knopper Eiche</i> , Ger. | |
| Varieties Y - - 1862 | 1862 |
| 2 <i>pendula</i> Y | |
| 3 <i>latifolia</i> Hort. Y | |
| § iii. <i>Alba</i> . White American Oaks. Y 1730. 1863 | 1863 |
| 8. <i>alba</i> L. Y North America pl. 309. fig. 1722. a, 1723. 1726, 1727. 1864 | 1864 |
| The American white Oak. | |
| Q. <i>alba virginiana</i> Park. | |
| Q. <i>a. pinnatifida</i> Walt. | |
| Q. <i>pubestris</i> Marsh. | |
| <i>Chêne blanc de l'Amérique</i> , Fr. | |
| <i>Weisse Eiche</i> , Ger. | |
| Varieties Y - - 1864 | 1864 |
| 1 <i>pinnatifida</i> Michx. Y f. 1723. a. | 1723. a. |
| Q. <i>alba</i> Ban. | |
| Q. <i>virginiana</i> Catesh. | |
| Q. <i>a. pubestris</i> Marsh. | |
| 2 <i>repanda</i> Michx. Y fig. 1723. b, 1724. | 1723. b, 1724. |
| 9. <i>olivæformis</i> Michx. Y Banks of the Hudson - fig. 1730. 1869 | 1869 |
| The olive-shaped-fruited American Oak. | |
| The mossy-cupped Oak, Amer. | |
| 10. <i>macrocarpa</i> Willd. Y United States pl. 310. fig. 1722. b, 1731. 1869 | 1869 |
| The large-fruited American Oak. | |
| The over-cup white Oak, Bur Oak Amer | |
| <i>Chêne à gros Glands</i> , <i>Chêne blanc</i> , Fr. | |
| <i>Gross-fruchtige Eiche</i> , Ger. | |
| 11. <i>obtusiloba</i> Michx. Y North America pl. 311. fig. 1752. 1722. c. 1870 | 1870 |
| The blunt-lobed-leaved, or Post, Oak. | |
| Q. <i>stellata</i> Willd. | |
| <i>Iron Oak</i> , <i>Box white Oak</i> , <i>American Turkey Oak</i> , <i>Upland white Oak</i> , Amer. | |
| 12. <i>lyrata</i> Walt. Y North America fig. 1733, 1734. 1871 | 1871 |
| The lyrate, or over-cup, Oak. | |
| Swamp Post Oak, Water white Oak, Amer. | |
| § iv. <i>Prinus</i> . Chestnut Oaks. Y 1730. 1872 | 1872 |
| 13. <i>Prinus</i> L. Y North America 1872 | 1872 |
| The <i>Prinus</i> , or Chestnut-leaved, Oak. | |
| Varieties. Y | |
| 1 <i>palustris</i> Michx. Fl. Y pl. 312. fig. 1735. - 1872 | 1872 |
| Q. P. <i>palustris</i> Michx. Syl. | |
| Q. <i>Prinus</i> L. | |
| Q. <i>castaneaefolia</i> , &c., Pluk. | |
| The Swamp Chestnut Oak, the Chestnut white Oak. | |
| 2 <i>monticola</i> Michx. Fl. Y pl. 313. fig. 1736. - 1873 | 1873 |
| Q. P. <i>monticola</i> Michx. fil. | |
| Q. <i>montana</i> Willd. | |
| Q. <i>Prinus</i> Sm. | |
| The Rock Chestnut Oak. | |
| 3 <i>acuminata</i> Michx. Fl. Y f. 1737. 1875 | 1875 |
| Q. P. <i>acuminata</i> Michx. fil. | |
| Q. <i>Castanea</i> Willd. | |
| The yellow Oak. | |
| 4 <i>pumila</i> Michx. Fl. Y fig. 1738. 1722. e. - 1875 | 1875 |
| Q. P. <i>Chinquapin</i> Michx. fil. | |
| Q. <i>Chinquapin</i> Pursh. | |
| Q. <i>prinoides</i> Willd. | |
| The <i>Chinquapin</i> , or Dwarf Chestnut, Oak. | |
| 5 <i>tomentosa</i> Michx. Fl. Y pl. 314. fig. 1722. d, 1739. 1876 | 1876 |
| Q. P. <i>discolor</i> Michx. fil. | |
| Q. <i>bicolor</i> Willd. | |
| Q. <i>Michauxii</i> Nutt. | |
| The Swamp white Oak. | |
| § v. <i>Rubra</i> . Red American Oaks. Y 1730. 1877. | 1877. |
| 14. <i>rubra</i> L. Y North America pl. 315, 316, 317. fig. 1740. to 1744. 1877 | 1877 |
| The red, or Champion, Oak. | |
| Q. <i>Escaulti divisura</i> , &c., Pluk. | |
| Varieties Y - - 1877 | 1877 |
| Q. <i>rubra latifolia</i> Y | |
| Q. <i>rubra</i> L. | |
| The Champion Oak. | |
| Q. <i>rubra montana</i> Y | |
| The mountain red Oak. | |
| 15. <i>coccinea</i> Willd. Y South America pl. 318, 319. fig. 1746, 1747, 1748. 1879 | 1879 |
| The scarlet Oak. | |
| Q. <i>rubra</i> ß Ait. | |
| 16. <i>ambigua</i> Willd. Y North America pl. 320. fig. 1719. 1881 | 1881 |
| The ambiguous, or grey, Oak. | |
| Q. <i>borcatis</i> Michx. | |

17. falcata Michx. ♀ North America Page
1894
fig. 1750, 1751. 1882
The sickle-shaped, or Spanish, Oak.
Q. discolor Ait.
Q. elongata Willd.
Q. lyrata Lodd. Cat.
Q. cuneata Wang.
Q. triloba Willd.
The downy-leaved Oak.
18. tinctoria Willd. ♀ North America Page
1894
pl. 321. fig. 1753, 1754. 1884
The Quercitron, or Dyer's Oak.
Q. virginiana, &c., Pluk.
Q. discolor Willd.
The black Oak, Amer.
Chêne des Teinturiers, Fr.
Varieties ♀ - - 1885
1 angulosa Michx. ♀ f. 1753, 1754
Q. nigra Pursh.
Q. americana Pluk.
Q. velutina Lam. Diet.
Q. tinctoria Bart.
2 sinuosa Michx. ♀ fig. 1755,
1756, 1757.
Q. nigra Wang.
19. palustris Willd. ♀ North America Page
1894
pl. 322. fig. 1758, 1759. 1887
The Marsh, or Pin, Oak.
Q. montana Lodd. Cat.
Q. Banisteri Lodd. Cat.
20. Catesbeii Willd. ♀ North America Page
1894
fig. 1762, 1763. 1889
The barren Scrub Oak.
Q. rubra β Abb. and Sm.
Q. Esculi divisiara, &c., Cat.
- § vi. Nigra. Black American Oaks. ♀ 1890
1730. 1890
21. nigra L. ♀ North America Page
1890
fig. 176½, 1765. 1890
The black Jack Oak.
Q. marylandica, &c., Ray.
Q. ferruginea Michx.
Q. aquatica Lodd. Cat.
Barrens Oak, Amer.
22. aquatica Soland. ♀ North America Page
1892
fig. 1767. 1892
The Water Oak.
Q. foliis cuneiformibus, &c., Gron.
Q. folio non serrato, &c., Cat.
Q. nigra Willd.
Q. uliginosa Wangh.
Varieties ♀ - - 1892
2 nana ♀ fig. 1767.
The dwarf jagged Oak.
Q. aquatica Sm.
Q. a. elongata Ait.
Q. dentata Bart.
Q. nana Willd.
3 maritima Michx. ♀
Q. hemisphaerica Willd.
Other Varieties.
See fig. 1767, 1768.
23. ilicifolia Wangh. ♂ North America Page
1893
fig. 1770. 1893
The Holly-leaved, or Bear, Oak.
Q. Banisteri Michx.
? Q. aquatica Abb. et Sm.
Black Scrub Oak. Dwarf red Oak, Amer.
24. heterophylla Michx. ♀ N. Amer. Page
1894
The various-leaved, or Bartram's, Oak.
25. agrifolia Willd. ♂ North America Page
1894
The prickly-leaved American Oak.
- § vii. Phellos. Willow Oaks. 1730. 1894
♀ 1894
26. Phellos L. ♀ North America Page
1894
pl. 323. fig. 1771. 1894
The Willow Oak.
Q. virginiana, &c., Pluk.
Q. Flex marylandica Ray.
Varieties ♀ 1895
1 sylvaticus Michx. ♀ fig. 1774.
2 latifolius Lodd. Cat. ♀ pl. 324.
3 humilis Pursh. ♂
4 sericeus ♂
Q. Phellos Sm.
Q. P. pumilus Michx.
Q. humilior sulcis foliis brevior.
The Highland Willow Oak.
Q. sericea Willd.
Q. pumila Michx.
The running Oak.
5 cinereus. ♀ ♂ fig. 1773.
Q. P. γ L.
Q. P. β cinereus Ait.
Q. humilis Walt.
Q. cinerea Willd.
The Upland Willow Oak.
6 maritimus Michx.
Q. maritima Willd.
27. (P.) laurifolia Willd. ♀ N. America Page
1897
fig. 1776. 1897
The Laurel-leaved Oak.
The Laurel Oak, Swamp Willow Oak.
Variety ♀ - - 1897
2 hybrida Michx. ♀ fig. 1775.
Q. l. 2. obtusa Ait.
28. imbricata Willd. ♀ North America Page
1898
fig. 1777. 1898
The Shingle Oak.
Q. latifolia Hort.
Laurel Oak, Field-Cup Oak, Jack Oak,
Black Jack Oak, Amer.
Chêne à Lattes, Fr.
B. Leaves evergreen.
- § viii. Flex. Holm, or Holly, Oaks. ♂ 1899
- a. Natives of Europe.
29. Flex L. ♀ South of Europe Page
1899
pl. 325. fig. 1781. 1899
The common evergreen, or Holm, Oak.
Flex arborea Bauh.
L'Yeuse, or Chêne vert, Fr.
Steine Eiche, Ger.
Elicc, Ital.
Encina, Spain.
Varieties ♂ - - 1899
1 integrifolia ♂
Santal, Dutch.
Saber secundus Matth.
2 serratifolia Lodd. Cat. ♂ f. 1778.
Flex Matth.
3 fagifolia Lodd. Cat. ♂ f. 1779.
Phellodrys Matth.
Flex, No. 3. Du Ham.
4 crispa Lodd. Cat. ♂
5 latifolia Lodd. Cat. ♂ pl. 396.
fig. 1780.
Q. l. oblonga Hort.

	Page
6 longifolia Lodd. Cat. ♀	1900
pl. 327.	
Q. l. salicifolia Hort.	
7 variegata Hort. ♀	
Other Varieties -	1900
30. Bullata Desf. ♂ Barbary	
fig. 1783, 1784. 1905	
The sweet Acorn Oak.	
? Flex major Clus.	
Chêne à Glands doux, Chêne Ballote, Fr.	
31. gramuntia L. ♀ ♂ Spain, &c.	
pl. 328. fig. 1787, 1788. 1906	
The Holly-leaved Grammont Oak.	
? Flex foliis rotundioribus, &c., Magn.	
Chêne de Grammont, Fr.	
Halleblättrige Eiche, Ger.	
Encina dulce, Gouette, Span.	
32. cocifera L. ♂ South of Europe	
fig. 1789, 1790, 1791, 1792. 1908	
The Kermes, or Berry-bearing, Oak.	
Flex cocifera Cam.	
Flex aculeata coccifera Garid.	
Flex cocigera, Ger.	
Chêne aux Kermes, Fr.	
Kermes Eiche, Ger.	
33. pseudo-cocifera Desf. ♀ ♂ Algiers	
fig. 1794. 1911	
The false berry-bearing, or Kermes, Oak.	
Chêne à faux Kermes, Fr.	
Stechernde Eiche, Ger.	
34. Süber L. ♀ South of Europe	
pl. 329, 330. fig. 1797, 1798. 1800.	
1911	
The Cork Tree.	
Süber Cam.	
Süber Prinus Matth.	
Süber latifolium, &c., Du Ham.	
Chêne Liège, Fr.	
Kork Eiche, Ger.	
Alcornoque, Span.	
Varieties ♀ - - 1911	
2 latifolium ♀	
Süber latifolium, &c., Bauh.	
3 angustifolium ♀ fig. 1798.	
Süber angustifolium Bauh.	
4 dentatum ♀ pl. 331. fig. 1797.	
Q. Pseudo-Süber of Muswell Hill.	
35. Pseudo-Süber Desf. ♀ Tuscany, &c.	
fig. 1801. 1917	
The False-Cork Oak.	
Chêne faux Liège, Chêne de Gibraltar, Fr.	
Fäuchte Kork Eiche, Ger.	
b. Natives of North America.	
§ ix. Virétes. Live Oaks. ♀ 1730. 1918	
36. virens Ait. ♀ North America	
pl. 332. fig. 1802, 1803. 1918	
The green, or Live, Oak.	
Q. Phellos L.	
Q. sempervirens Banister.	
37. myrtifolia Willd. ♀ Carolina - 1920	
The Myrtle-leaved Oak.	

	Page
c. Natives of Nepal.	
§ x. Lanata. Woolly, or downy, leaved Oaks. ♀	1730. 1920
38. lanata Sm. ♀ Upper Nepal	
fig. 1804. 1920	
The woolly-leaved Nepal Oak.	
Q. lanuginosa D. Don.	
Q. Banja Ham. MSS.	
? Q. oblongata D. Don.	
? Q. incana Royle.	
39. annulata Sm. ♀ Upper Nepal	
fig. 1805. 1921	
The ring-cupped Oak.	
Q. Phallata Ham. MSS.	
? Q. Kamroopi D. Don.	
Q. glauca Lodd. Cat.	
? Q. glauca Thunb.	
? Q. acuminata Hort.	
App. i. Oaks in British Gardens, not referable, with certainty, to any of the above Sections. ♀ ♀ - - - 1922	
40. Turneri Willd. ♀ Thibet f. 1806. 1922	
Turner's Oak.	
Q. hybrida Hort.	
Chêne de Turner, Fr.	
Turnersche Eiche, Ger.	
41. hybrida nana ♀ Hybrid	
fig. 1810, 1811. 1924	
The dwarf hybrid Oak.	
Q. hybrida Lodd. Cat.	
A hybrid between Q. pedunculata and Q. Flex, Hort. Soc. Gard.	
Q. hinnitis Hort.	
Q. nana Hort.	
42. Fontanesii Guss. ♀ Calabria	
fig. 1813. 1925	
Desfontaine's Oak.	
Q. pseudo-cocifera of Catros, &c.	
43. ? australis Link. ♀ Gibraltar	
fig. 1814. 1925	
The southern Oak.	
44. Cookii ♀ Gibraltar fig. 1815. 1926	
Captain Cook's Oak.	
45. falkenbergensis Benth. ♀ Falkenberg	
1926	
The Falkenberg Oak.	
App. ii. European kinds of Oaks, not yet introduced. - - - 1926	
Q. faginea Lam. Spain. fig. 1816	
Q. aculeatissima Lam. Diet.	
Phellobos, alba angustifolia, &c., Dalech.	
Q. argilifolia Pers. Spain	
Q. hispanica S. Lam. Diet.	
Chêne à Feuilles d'Églisop, Bose.	
Q. Breton Bose. Between Ferrogneux and Bordeaux.	
Chêne Breton, at Nantes.	
Chêne marin, Banani.	
Q. umbratica Bose, Java	
Chêne Soule, Chêne Oair, Chêne de Baie, Fr.	
Q. aspera Bose.	
Le Chêne épre, Fr.	
Chêne Lerermien, Bose.	
Chêne Castellan, Bose. Spain.	
Q. lusitanica Lamb. Portugal. fig. 1817.	
Q. rotundus Fr.	
Robur 1., and Robur 2., C. P. S.	
Galho, sive Robur majus, Ger., and Galho minor Ger.	
The Portuguese Gall Oak.	
Chêne du Portugal, Fr.	

	Page
<i>Q. pérsiana</i> Pers. Portugal. fig. 1818.	
<i>Q. glauca</i> Bosc.	
<i>Q. calycina</i> Poir. France	
<i>Q. expansa</i> Poir.	
<i>Q. rotundifolia</i> Lam. Spain.	
<i>The round-leaved Spanish Oak.</i>	
<i>Chêne à Feuilles rondes.</i>	
<i>Q. humilis</i> Lam. Portugal.	
<i>The dwarf Portuguese Oak.</i>	
<i>Q. pædem viræ superans</i> Bauh.	
<i>Robur 7., sive Q. pumila</i> Clus.	
<i>Chêne pygmée, Fr.</i>	

App. iii. African Oaks which have not yet been introduced. - 1928

Q. obtecta Poir.

App. iv. Oaks of Asia Minor and Persia not yet introduced. - 1928

<i>Q. infectoria</i> Oliv. fig. 1819, 1820.	
<i>Q. caricusis</i> Willd.	
<i>Chêne à Galles, Fr.</i>	
<i>Elber Eiche, Ger.</i>	
<i>Q. Libani</i> Oliv. Mount Lebanon. fig. 1826.	
<i>Q. rigida</i> Willd. Caramania.	
<i>Q. Mex aculeata, &c., Tourn.</i>	
<i>Q. ibérica</i> Stev. Georgia.	
<i>Q. castaneaefolia</i> C. A. Meyer. Mazanderan.	
<i>Q. mongolica</i> Fisch. Tartary.	

App. v. Himalayan Oaks not yet introduced. 1933

<i>Q. spicata</i> Sm. Nepal. fig. 1828.	
<i>Q. spatulata</i> Bosc.	
<i>Q. A'rcula</i> Ham. MSS.	
<i>Q. obtusifolia</i> D. Don. Nepal	
<i>Q. grandifolia</i> D. Don. Nepal. fig. 1829.	
<i>The Magnolia-leaved Oak.</i>	
<i>Q. velutina</i> Lindl. Tavoy. fig. 1830.	
<i>Q. lamellosa</i> Sm. Nepal. fig. 1831.	
<i>Q. induricata</i> Ham. MSS.	
<i>Q. semicarpifolia</i> Sm. Nepal. fig. 1832.	
<i>Cassia Ham. MSS.</i>	

App. vi. Oaks of Japan, Cochín-China, and China, which have not yet been introduced. 1935

<i>Q. glabra</i> Thunb. Japan.	
<i>Q. concentrica</i> Lour. Cochín-China	
<i>Q. acuta</i> Thunb. Japan.	
<i>Q. serrata</i> Thunb. Japan.	
<i>Q. glauca</i> Thunb. Japan.	
<i>Kaz no Ki</i> Kæmpf.	
<i>Q. cuspidata</i> Thunb. Japan.	
<i>Sui, vulgo Sui no Ki, Kæmpf.</i>	
<i>Fagus folio Fraxini</i> Kæmpf.	
<i>Q. dentata</i> Thunb. Japan.	
<i>Koku Kæmpf.</i>	
<i>Q. chinensis</i> Baugé. China.	
<i>Q. obovata</i> Bunge. Near Pekin.	

App. vii. Oaks of Java, Sumatra, and the Molucca Isles, not yet introduced. - 1936

<i>Q. sundaiæ</i> Blume. Java. fig. 1833, 1831.	
<i>The Sunda Oak.</i>	
<i>Q. prunosa</i> Blume. fig. 1836.	
<i>The frosty Oak.</i>	
<i>Q. angustata</i> Blume. Gedé. fig. 1835.	
<i>The narrow-leaved Oak.</i>	
<i>Q. pallida</i> Blume. Gedé. fig. 1837, 1838.	
<i>The pale Oak.</i>	
<i>Q. elegans</i> Blume. Bantam. fig. 1832.	
<i>The elegant Oak.</i>	
<i>Q. placenteria</i> Blume. Gedé. fig. 1810.	
<i>The placenta-cupped Oak.</i>	
<i>Q. glaberrima</i> Blume. Gedé. fig. 1812.	
<i>The smoothest-leaved Oak.</i>	
<i>Q. costata</i> Blume. fig. 1811, 1843.	
<i>The ribbed-cupped Oak.</i>	
<i>Q. rotundata</i> Blume. fig. 1844.	
<i>The round-fruited Oak.</i>	
<i>Q. platycarpa</i> Blume. Bantam. fig. 1846.	
<i>The broad-fruited Oak.</i>	
<i>Q. daphnoides</i> Blume. Bantam. fig. 1845.	
<i>The Daphne-like Oak.</i>	
<i>Q. racemosa</i> Hook. Sumatra.	
<i>Q. gemelliflora</i> Blume. Salak, &c. fig. 1847.	
<i>The twin-flowered Oak.</i>	
<i>Q. induta</i> Blume. Gedé. fig. 1845.	
<i>The cloth-cupped Oak.</i>	
<i>Q. urceolaris</i> Hook. Sumatra.	

	Page
<i>Q. pseudo-molucca</i> Blume. Java. fig. 1849.	
<i>The false Molucca Oak.</i>	
<i>Q. molucca</i> L. Molucca Isles.	
<i>The Molucca Oak.</i>	
<i>Q. turbinata</i> Blume. Salak. fig. 1850.	
<i>The top-shaped-cupped Oak.</i>	
<i>Q. lineata</i> Blume. Java. fig. 1851.	
<i>The parallel-veined Oak.</i>	

App. viii. Mexican Oaks not yet introduced. 1941

<i>Q. xalapensis</i> Humb. et Boep. New Spain. fig. 1852.	
<i>Roble de Dueto, Span.</i>	
<i>Q. glaucescens</i> Humb. et Boep. Mexico. fig. 1853.	
<i>Q. obtusa</i> Humb. et Boep. New Spain. fig. 1851.	
<i>Q. pandurata</i> Humb. et Boep. fig. 1853, 1856.	
<i>Q. repanda</i> Humb. et Boep. New Spain. fig. 1857.	
<i>Q. laurina</i> Humb. et Boep. New Spain. fig. 1858.	
<i>Q. sideroxyla</i> Humb. et Boep. New Spain. fig. 1860.	
<i>Q. mexicana</i> Humb. et Boep. Mexico. fig. 1861.	
<i>Q. crassipes</i> angustifolia Humb. et Boep. Ariz. fig. 1859.	
<i>Q. crassipes</i> Humb. et Boep. Ariz. fig. 1862.	
<i>Q. lanceolata</i> Humb. et Boep. New Spain. fig. 1863.	
<i>Q. reticulata</i> Humb. et Boep. New Spain. fig. 1865.	
<i>Q. chrysophylla</i> Humb. et Boep. fig. 1864.	
<i>Q. pulchella</i> Humb. et Boep. New Spain. fig. 1866.	
<i>Q. spicata</i> Humb. et Boep. Nabajas. fig. 1867.	
<i>Q. stipularis</i> Humb. et Boep. New Spain. fig. 1868.	
<i>Q. crassifolia</i> Humb. et Boep. Mexico. fig. 1869.	
<i>Q. depressa</i> Humb. et Boep. Mexico. fig. 1871.	
<i>Q. ambigua</i> Humb. et Boep. Mexico. fig. 1870.	
<i>Q. confertifolia</i> Humb. et Boep. New Spain. fig. 1872.	
<i>Q. tridens</i> Humb. et Boep. New Spain. fig. 1873.	
<i>Q. acutifolia</i> Willd. New Spain. fig. 1874.	
<i>Q. elliptica</i> Willd. Mexico.	
<i>Q. mucronata</i> Willd. New Spain.	
<i>Q. Castanea</i> Nee.	
<i>Q. tomentosa</i> Willd. New Spain.	
<i>Q. pedunculata</i> Nee.	
<i>Q. circinata</i> Willd. New Spain.	
<i>Q. splendens</i> Willd. New Spain.	
<i>Q. rugosa</i> Willd. Huifaulica and Ociula.	
<i>Q. macrophylla</i> Willd. New Spain.	
<i>Q. diversifolia</i> Willd. New Spain.	
<i>Q. cándicans</i> Willd. New Spain.	
<i>Q. microphylla</i> Willd. New Spain.	
<i>Q. lobata</i> Willd. New Spain.	
<i>Q. magnoliaefolia</i> Willd. Mexico.	
<i>Q. lutea</i> Willd. Mexico.	
<i>The yellow-leaved Mexican Oak.</i>	
<i>Q. salicifolia</i> Willd. Mexico.	
<i>The Willow-leaved Mexican Oak.</i>	

II. FAGUS L. † † - 1715. 1949
THE BEECH.

<i>Fagus</i> of the Romans.	
<i>Oxva</i> of the Greeks.	
<i>Castanea</i> Tourn.	
<i>Hêtre, Fr.</i>	
<i>Buche, Ger.</i>	
<i>Beuke, Dutch.</i>	
<i>Bog, Dan.</i>	
<i>Bok, Swed.</i>	
<i>Buk, Russ. and Pol</i>	
<i>Faggio, Ital.</i>	
<i>Haya, Span.</i>	
<i>Faya, Port.</i>	

A. Capsule muricate, capsuliform. Ovaries included. Young leaves plicate.

Natives of Europe, and of North and South America.

a. Species in Cultivation in British Gardens.

1. sylvatica L. † Europe - 1950

<i>The Wood, or common, Beech.</i>	
<i>Castanea Fagus, Scop.</i>	
<i>Fagus</i> Bauh.	
<i>F. sylvestris</i> Michx.	
<i>O'zga, Greek.</i>	
<i>Fagus, Latin.</i>	
<i>Hêtre commun, Fr.</i>	
<i>Gemeine Buche, Ger.</i>	
<i>Rood-beuke, Dutch.</i>	

Varieties † - - 1950

2 purpurea Ait. †	
<i>The purple Beech.</i>	
<i>F. s. 2. atro-rubens</i> Du Roi.	
<i>Hêtre noir, Fr.</i>	
3 cuprea Lodd. Cat. †	
<i>The copper-coloured Beech.</i>	

	Page
4 fölliis variegätis Lodd. Cat. Y	1875, 1876.
5 heterophýlla Y fig. 1875, 1876.	1877.
The various, or cut, leaved Beech.	
F. s. <i>laciniata</i> Lodd. Cat.	
F. s. <i>asplenifolia</i> Lodd. Cat.	
F. s. <i>incisa</i> Hort.	
F. s. <i>salicifolia</i> Hort.	
<i>Hêtre à Feuilles de Saule</i> , Fr.	
6 cristata Lodd. Cat. pl. 334.	1877.
The crested, or curled-leaved, Beech.	
F. s. <i>crispa</i> Hort.	
<i>Hêtre Crête de Coq</i> , Fr.	
7 péndula Lodd. Cat. Y pl. 335,	336.
The weeping Beech.	
<i>Hêtre Parasol</i> , Fr.	
Other British Varieties or Variations.	

8 americana Y	1917, 1980
F. <i>syloëstris</i> Michx.	
White Beech, Amer.	
2. ferruginea Ait. Y N. Am. f. 1917, 1980	1915.
The American ferruginous-wooded Beech.	
F. <i>americana latifolia</i> Du Roi.	
Red Beech, Amer.	
Varieties. Y	
2 caroliniana Y fig. 1915.	1916.
F. <i>caroliniana</i> Lodd. Cat.	
3 latifolia Y fig. 1916.	Lee.
F. <i>latifolia</i> Lee.	

b. Species not yet introduced.

3. obliqua Mirb. Y Chili fig. 1919. 1982	1919.
The oblique-leaved Beech.	
B. Cupul: involucriform; Segments narrow, laciniate. Ovaries laterally inserted. Young leaves not plicate.	

a. Species introduced into Britain.

4. betuloides Mirb. ♀ Terra del Fuego	1920, 1982
The Birch-like, or evergreen, Beech.	
<i>Betula antarctica</i> Forst.	
5. antarctica Forst. Y Terra del Fuego 1982	1982
The antarctic Beech.	
b. Species not yet introduced into British Gardens.	
6. Dombèyi Mirb. Y Chili fig. 1921. 1982	1921, 1982
Dombéy's, or the Myrtle-headed, Beech	
7. dubia Mirb. ♀ Straits of Magellan	1932, 1933
The dubious Beech.	

III. CASTANEA Tourn. Y 1716 1983

THE CHESTNUT.

Fagus L. and others.
Chataignier, Fr.
Kastanie, Ger.
Castagno, Ital.
Castano, Span.
Castanhoiro, Port.
Castaniecræ, Swed. and Dan.
Keschton, Russ.

1. véscæ Gartn. Y Asia Minor	1937, 338. 1983
The eatable, sweet, or Spanish, Chestnut.	
<i>Fagus Castanea</i> L.	
<i>Castanea sativa</i> Mill.	
<i>Castanea vulgaris</i> Lam.	

Varieties Y	-	Page
		1984

A. Botanical Varieties.

2 asplenifolia Lodd. Cat. Y	1927, 1928. 2002
C. <i>heterophylla</i> Hort.	
C. <i>laciniata</i> Hort.	
C. <i>salicifolia</i> Hort.	
3 cochleata Lodd. Cat. Y	1927, 1928. 2002
C. v. <i>foliis lucidis</i> Hort.	
4 glabra Lodd. Cat. Y	1927, 1928. 2002
C. v. <i>foliis aureis</i> Lodd. Cat.	
5 glauca Y	1927, 1928. 2002
C. <i>glauca</i> Hort.	
6 variegata Y	1927, 1928. 2002
C. v. <i>foliis aureis</i> Lodd. Cat.	
7 americana Y	1927, 1928. 2002
C. <i>véscæ</i> Michx.	

B. Fruit-bearing Varieties. (See p. 1984.)

2. pumila Willd. Y ♂ North America	1927, 1928. 2002
The Dwarf Chestnut, or Chincapin.	
<i>Fagus pumila</i> L.	
<i>Castanea pumila virginiana</i> , &c., Pluk.	
<i>Chataignier Chincapin</i> , Fr.	
<i>Zwerch Kastanie, or Castanje</i> , Ger.	

App. i. Species of Castanea not yet introduced into European Gardens. 2003

C. <i>indica</i> Roxb. Nepal.	1929.
C. <i>Roxburghii</i> Lindl. Chittagong.	1929.
<i>Quercus castaneicarpa</i> Roxb.	1929.
C. <i>spheroarpa</i> Lindl. Sikkim.	1929.
C. <i>tribuloides</i> Lindl. Upper Nepal.	1929.
<i>Quercus tribuloides</i> Sm.	1929.
<i>Quercus castanea</i> Hana. MSS.	1929.
<i>Quercus ferox</i> Roxb.	1929.
C. <i>martabanica</i> Wall. Martaban. fig. 1929.	1929.
C. <i>argentea</i> Blume. Java. fig. 1831.	1929.
C. <i>Tinggorot</i> Blume. Bantam. fig. 1930.	1929.
C. <i>javanica</i> Blume. Gede. fig. 1932.	1929.
var. C. j. <i>montana</i> Blume.	1929.
C. <i>montana</i> Blume. Bajador.	1929.
C. j. <i>fusciceps</i> Blume.	1929.
C. <i>inermis</i> Lindl. Singapore.	1929.
C. <i>chinensis</i> Spreng. China.	1929.

IV. CARPINUS L. Y ♂ 1716. 2004
 THE HORNBAM.
Charme, Fr.
Hagnbuche, or Hainbuche, Ger.

1. Bétulus L. Y Britain	1937, 339. 2004
The Birch, or common, Hornbeam.	
<i>Carpinus</i> Matth.	
<i>Ostrya</i> Bauh. Pin.	
<i>Oryus</i> Trag.	
<i>Fagus</i> Bauh. Hist.	
<i>Bétulus</i> Lob.	

Varieties Y - - 2005

2 incisa Lodd. Cat. Y	1936. 2013
C. v. <i>quercifolia</i> Desf.	
C. v. <i>heterophylla</i> Hort.	
3 variegata Lodd. Cat. Y	1936. 2013
2. (B.) americana Michx. Y N. America	1936. 2013
The American Hornbeam.	
C. <i>virginiana</i> Michx.	
3. (B.) orientalis Lam. Y ♂ Asia Minor	1937. 2014
The Oriental Hornbeam.	
C. <i>dumensis</i> Scop.	

- Page
 App. i. *Species or Varieties of Cárpinus not yet introduced into European Gardens.* 2014
C. (B.) Carpinizza Hort. Transylvania.
C. viminea Lindl. Nepal. fig. 1938.
C. faginea Lindl.
- V. *O'STRYA* Willd. ♀ - 1716. 2015
 THE HOP HORNBEAM.
Cárpinus L. and others.
Hopfenbuche, Ger.
1. *vulgàris* Willd. ♀ Italy
 pl. 340, 341. fig. 1939. 2015
 The Hop Hornbeam.
Cárpinus O'strya Hort. Cliff.
O'strya carpinifolia Scop.
O'strya Bauh.
O'strya italica, &c., Michx.
2. (v.) *virgínica* Willd. ♀ North America
 pl. 342. 1940. 2015
 The Virginian Hop Hornbeam.
Cárpinus virginiana Abb.
Cárpinus O'strya virginiana Michx. Fl.
Cárpinus O'strya Michx. Syl.
Iron Wood, Lever Wood, Amer.
Bois dur, Illinois.
- VI. *CO'RYLUS* L. ♀ 1716. 2016
 THE HAZEL.
Coudrier, Fr.
Haselnuss, Ger.
1. *Avellàna* L. ♀ Europe fig. 1941. 2017
 The common Hazel Nut.
Coudrier Noisetier, Fr.
Haselstrauch, Nussbaum, Ger.
Avellano, Nocciole, Ital.
Avellano, Span.
Varieties ♀ - - 2017
 A. *Botanical Varieties.*
- 1 *sylvéstris* Ait. ♀ fig. 1941
C. Avellàna Svensk.
C. sylvéstris Bauh.
- 2 *pùmilus* ♀
C. pùmilus Lodd. Cat.
- 3 *heterophýlla* ♀
 The various, or Nettle, leaved Hazel.
C. heterophýlla Lodd. Cat.
C. laciniàta Hort.
C. urticifolia Hort.
- 4 *purpurea* ♀
C. purpurea Lodd. Cat.
C. atro-purpurea Hort.
- B. *Varieties cultivated for their Fruit.*
- 5 *tubulòsa* fig. 1942. ♀
C. tubulosa Willd.
C. maxima Mill.
C. sativa Bauh.
C. s. rubra Ait.
Red Filbert.
Langbartnuss, or Lambertnuss,
 Ger.
Noisetier franc à Fruit rouge, Fr.
- 6 *tubulòsa álba* ♀
C. sativa álba Ait.
C. A. álba Lodd. Cat.
White Filbert.
Weisse Langbartnuss, Ger.
- 7 *crispa E. of Pl.* ♀ fig. 1943.
 The frizzled Filbert.
- 8 *ténuis* Lodd. Cat. ♀
 The thin-shelled, or *Cosford*, Nut.
- 9 *glomeràta* Bauh. ♀
C. glomerata Lodd. Cat.
Cluster Nut, Hort. Soc. Cat.
Noisetier à Grappes, Fr.
- 10 *barcelonénsis* Lodd. Cat. ♀
C. sativa grândis Bauh.
C. A. grândis Lodd. Cat.
The Cob Nut.
The Barcelona Nut.
Downton large Nut, &c., Hort. Soc. Cat.
- 11 *Lambérti*. ♀
C. Lamberti Lodd. Cat.
The Spanish Nut.
Large Bond Nut, Lambert's Nut,
Lambert's large Nut, Token Nut,
 &c., Hort. Soc. Cat.
Other Varieties.
 The great Cob Nut, Hort. Soc. Cat.
 The Downton large square Nut,
 Hort. Soc. Cat.
 The Northampton Nut, Hort.
 Soc. Cat.
 The Northamptonshire Prolific,
 Hort. Soc. Cat.
2. *Colúrna* L. ♀ Turkey
 pl. 343, 344. fig. 1948. 2029
 The Constantinople Hazel.
C. byzantina Herm.
Avellana peregrina humilis Bauh.
A. pùmila byzantina Clus.
C. arborea Hort.
Le Noisetier de Bizance, Fr.
Byzantinische Haselnuss, Ger.
Varieties ♀ - - 2029
 2 *intermèdia* ♀
C. intermèdia Lodd. Cat.
 3 *arboréscens* Fisch. ♀
3. *rostràta* Ait. ♀ North America 2030
 The beaked, *American, or Cuckold*, Hazel.
C. sylvéstris, &c., Gron.
C. cornuta Hort.
4. *americàna* Michx. ♀ N. America 2030
 The American Hazel.
C. americàna humilis Wang.
Dwarf Cuckold Nut, wild Filbert, Amer.
- App. i. *Species of Córnylus not yet introduced.* 2031
C. ferox Walt. Nepal. fig. 2250.
-
- Garryàceæ.* - 2031
- I. *GA'RRYA* Doug. ♀ - - 2031
 THE GARRYA.
1. *ellíptica* Doug. ♀ North Carolina
 fig. 1951. 2032
 The elliptic-leaved Garrya.
-
- Platanàceæ.* ♀ ♀ - 2032
- I. *PLA'TANUS* L. ♀ ♀ - 2033
 THE PLANE TREE.
Platane, Fr.
Platanus, Ger.

1. *orientalis* L. ♀ Levant
pl. 345, 346. fig. 1954, 1955. 2033

The Oriental Plane.

P. orientalis *verus* Park.
Platane de l'Orient, Fr.
Morgenlandischer Platannus, Ger.
Doolb, Arabic.
Chinar, Persian.

Varieties ♀ - - 2034

2 *acerrifolia* Ait. ♀ pl. 347, 348.

The Maple-leaved Plane Tree.

P. o. Aceris folio Tourn.
P. acerrifolia Willd.
P. intermedia Hort.

3 *hispanica* ♀

The Spanish Maple.

P. hispanica Lodd. Cat.
P. macrophylla Cree.

4 *cuneata* ♀ pl. 349.

P. o. undulata Ait.
P. cuneata Willd.

Other Varieties.

2. *occidentalis* L. ♀ ♂ North America
pl. 350. fig. 1959. 2043

The Western Plane.

P. occidentalis seu virginensis Park.
Button-wood, *Water Beech*, *Sycamore*,
Cotton Tree, Amer.
Platane de Virginie, Fr.

Variety ♀ - - 2043

2 *tortuosa* ♀

Platane tortillard, Fr.

Balsamiferae. ♀ - 2048

I. LIQUIDAMBAR L. ♀ - 2049

THE LIQUIDAMBAR.

Allingia Noronha.
Liquidambar, Fr.
Ambarbaum, Ger.

1. *Styraciflua* L. ♀ North America

pl. 351, 352. fig. 1961. 2049

The Sweet Gum Liquidambar.

Liquidambar arbor Pluk.
Styrax Aceris folio Ray.
Liquidambar resinosa, *Copalme de l'A-*
merique, *Liquidambar Copal*, Fr.
Pliesender Ambarbaum, Ger.

2. *imbérbe* Willd. ♀ Levant f. 1963. 2053

The beardless, or Oriental, Liquidambar.

L. orientalis Mill.
? *Platanus orientalis* Pocko.
L. imberbis Sm.

App. i. *Species of Liquidambar not yet introduced.* - - 2054

I. *Allingia* Blume. Java. fig. 1965.
Allingia Liquidambar.
Allingia excelsa Noronha.
Lignum papuanum Rumph.

Myricaceae. ♂ ♀ 2055

I. MYRTICA L. ♂ ♀ - 2055

THE CANDLEBERRY MYRTLE.

Gale, Fr.
Wachs Strauss, Ger.

1. *Gale* L. ♂ Europe f. 1996, 1967. 2056

The Sweet Gale, Sweet Willow, Candleberry Myrtle, or Dutch Myrtle.

Gale Ray.
Eleagnus Card.
Myrtus brabantica, Ger.
Rhus myrtifolia belgica Bauh.
Rhus sylvestris altera Dalech.
Rhus sylvestris Park.
Myrica palustris Lam.
Gale, *Pimento Royal*, Fr.
Gemeine Wachs Strauch, Ger.

2. *cerifera* L. ♂ North America 2057

The common Wax-bearing, or American, Candleberry Myrtle.

M. cerifera angustifolia Ait.
Myrtus brabantica, sc. Pluk.
Cérrier de la Louisiane, Fr.

Varieties ♂ ♀ - - 2057

2 *latifolia* Ait. ♂ fig. 1668.

The broad-leaved American Candleberry Myrtle.

M. c. media Michx.
M. carolinensis Willd.
M. pennsylvanica Lam.
M. c. sempervirens Hort.
Myrtus brabantica Cat.
Cérrier de Pennsylvanie, Fr.
Carolinischer Wachstrauch, Ger.

3 *pumila* Michx. ♂

App. i. *Half-hardy Species of Myrica cultivated in British Gardens.* - 2058

M. Faya Ait. Madeira. fig. 1969.
The Azores Candleberry Myrtle.

M. serrata Lam. C. G. H.

M. athiopica L.

M. confusa Burm.

M. quercifolia L. C. G. H.

Laurus africana, Sc., Com.

M. q. hirsuta Mill.

M. cordifolia L. C. G. H.

Alaternoides Ylicia folio, Sc., Walt.

M. foliis subcordatis, Sc., Burm.

Gale capensis, Sc., Petiv.

Coriutrigemulaleudra Ylicia neulota folio Pluk.

M. capensis Lodd. Cat.

App. ii. *Half-hardy Species of Myrica not yet introduced.* - 2059

M. spatulata. Madagascar. fig. 1970.

II. COMPTONIA Banks. ♂ - 2059

THE COMPTONIA.

Liquidambar L. Sp.

Myrica L. Hort. Cliff.

Gale Petiv.

Comptonia, Fr.

Comptonia, Ger.

1. *asplenifolia* Banks. ♂ North America
fig. 1971. 2059

The Asplenium-leaved Comptonia.

Liquidambar asplenifolium L. Sp.

Liquidambar pergrinum L. Syst.

Myrica L. Hort. Cliff.

Gale mariana Pet.

Myrtus brabantica affinis Pluk.

The sweet Fern Bush, Amer.

Casuarieae. 2060

Casuarina equisetifolia East Indies
fig. 1972. 2060

C. littorea Rumph.
Swamp Oak, Austral.
Filao d' Feuilles de Prê, Fr.

C. nodiflora Forst. New Caledonia.
C. distyla Vent. New Holland.
C. stricta Ait. New Holland.
C. torulosa Ait. Holland.
The Cork-barked Casuarina

Page
Gnetaceæ. ♁ ♀ - 2062

I. *E'PHEDRA* L. ♁ ♀ - 2062
 THE EPHEDRA.

1. *distàchya* L. ♁ Spain
 fig. 1973, 1974. 2063
 The two-spiked Ephedra.
Great shrubby Horsetail, or Sea Grape.
E'phedra vulgaris Rich.
Polygonum marinum Tabern.
Polygonum quartum Plinii Clus.
Polygonum lamifolium, &c., Bauh.
E. maritima major Tourn.
 Tragos Cam.
Raisin de Mer, Ephedre multiflore, Fr.
Zueghrigger Ross Schwanz, Ger.

2. *monostàchya* L. ♀ Siberia
 fig. 1975, 1976. 2063
 The one-spiked Ephedra, or Small shrubby
Horsetail.
E. petiolis sæpe plùribus, &c., Gmel.
E. minima, &c., Amm.
E. polygonoides Pall.
Ephedre mineure, Ephedre de Sibérie, Fr.

3. *altissima* Desf. ♁ Barbary
 fig. 1977, 1978, 1979. 2064
 The loftiest Ephedra.
E. sine Arabiæ Bellonii, &c., Tourn.
Polygonum maritimum scandens Bauh.

4. *frágilis* Desf. ♁ Spain - 2065
 The fragile Ephedra.
E. cretica Tourn.
Episcetum montanum creticum Alp.]

5. *americana* Willd. ♁ Quito f. 1980. 2065
 The American Ephedra.

Taxaceæ. † † ♁ ♀ 2065

I. *TA'XUS* L. † ♁ ♀ 2065, 2066
 THE YEW.

1. *baccàta* L. † Eur. pl. 353, 354. 2066
 The berried, or common, Yew.
Taxus No. 1663, Hall. Hist.
Efenbaum, Ithenbaum, or Eichenbaum, Ger.
Taxo, Ital.
Tero, Span.

Varieties † ♁ ♀ - 2066

2 *fastigiàta* † pl. 355. fig. 1981,
 1982.

T. fastigiàta Lindl.
T. hibernica, Hook.
The upright, or Florence Court,
Yew; the Irish Yew.

3 *procùmbens* ♁ ♀
T. procùmbens Lodd. Cat.

4 *erècta* ♁
 The upright Yew.
T. b. fastigiàta.

5 *fòliis variegàtis* Lodd. Cat. ♁

6 *frùctu lùteo* ♁
 Other Varieties.

2. (b.) *canadènsis* Willd. ♁ N. Amer. 2093
 The Canada, or North American, Yew.
T. b. minor Michx.

Page
II. SALISBURIA Sm. † 2065. 2094
 THE SALISBURIA.
Ginkgo Kæmpf., L., and others.

I. *adiantifolia* Sm. † Japan
 pl. 356, 357. fig. 1992, 1993. 2094
 The Maiden-hair-leaved Salisburia, or *Ginkgo*
Trec.
Ginkgo, Gin-an, Itsjo, Kæmpf.
Ginkgo biloba L.
Noyer du Japon, Arbre aux quarante
E'cus.

App. i. *Half-hardy Genera belonging to the*
Order Taxaceæ. - 2100

Podocarpus L'Hérit.
P. macrophyllus Sieb., Japan.
 The long-leaved Japan Yew.
Taxus macrophylla Thunb.
P. latifolius Wall. Pundna. fig. 1995.
P. spinulosus Spreng. Port Jackson
P. excelsus Lodd. Cat.
Taxus spinulosus Smith.
P. nùcifer Pers., Japan.
Taxus nucifera Kæmpf.
P. elongatus L'Hérit. C. G. H. fig. 1997.
Taxus elongatus Ait.
P. chilinus Rich., Chili. fig. 1996.
P. coriàceus Rich. Island of Montserrat. fig. 1998.
P. taxifolius Kunth. Peru. fig. 1999, 2000.
P. montanus Lodd. Cat.
Taxus montana Willd.
P. excelsus Lodd.
P. nerifolius Lodd.
Taxus japonica Lodd.
Dacrydium Solander.
D. cupressinum Sol. New Zealand. fig. 2001.
Thalassia cupressina Spreng.
D. taxifolium Sol.
D. excelsum Don. New Zealand.
The Kahikatea, or Swamp Pine.
D. ? plumosum D. Don. New Zealand.
The Karaka of the New Zealanders.
D. elatum Wall. Pulo-Penang.
Juniperus elata Roxb.
Phyllocladus Rich.
Podocarpus sp. Labill.
P. rhomboidalis Rich. Cape Van Diemen. fig. 2002, 2005.
Podocarpus asplenifolius Labill.
P. trichomanoides R. Br. New Zealand.
The Tanakau of the New Zealanders.

Coniferae, or Pinaceæ. 2103

Sect. I. *ABIE'TINE* Richard. 2104. 2106

I. *PINUS* L. † † † † † † † † † †
 2104. 2152

THE PINE.
Le Pin, Fr.
Fichte, Pynbaum, or Kiefer, Ger.
Pybaom, Dutch.
Pino, Ital. and Span.
Pinn, Anglo-Saxon.
Pinnua, Welsh.
Peigne, Erse.

Sect. i. *Binae*. — *Leaves generally 2 in a*
Sheath.

‡ i. *Sylvestres*. † ♁

A. *Cones having the Scales without Prickles.*

1. *sylvé'tris* L. † Europe
 fig. 2043, 2044, 2045. 2153
 The wood, or Scotch, Pine, or Scotch Fir.
P. fòliis binis, &c., Hall.
P. rubra Mill. Dict.
P. sylvé'tris communis Ait.

	Page
P. No. 29. Umel.	
<i>Pin Sauvage, Pin d'Ecosse, Fr.</i>	
<i>Gemeine Föhre, gemeine Fichte, Kiefer, Tanne, and 55 other names given in Hayne's Abbildung, Ger.</i>	
<i>Pyndam, Dutch.</i>	
<i>Pino sylvestico, Ital.</i>	
<i>Pino sylvestre, Span.</i>	
<i>Fyrre, Dan. and Swed.</i>	
<i>Sosna, Pol., Boh., and Russ.</i>	
Varieties † - - 2153	
a. <i>Timber Trees.</i> 2154	
1 <i>vulgäris</i> † fig. 20-16.	
The common wild Pine.	
2 <i>horizontalis</i> †	
<i>P. horizontalis</i> Don of Forfar.	
<i>P. s. var. montana</i> Sang.	
? <i>P. rubra</i> Mill. Diet. and N. Du Ham.	
<i>The Spycide Pine, Hort. Soc.</i>	
<i>The Highland Pine, Grigor.</i>	
<i>The horizontal-branched wild Pine, Laws.</i>	
<i>The red-wooded Scotch Pine, Sang.</i>	
3 <i>uncinata</i> Don of Forfar †	
fig. 20-17.	
The hooked-coned wild Pine.	
<i>Mar Forest wild Pine, Hort. Soc. Gard.</i>	
4 <i>haguenensis</i> †	
<i>Pin de Haguenau, Fr.</i>	
5 <i>rigensis</i> †	
<i>Pin de Riga Desf.</i>	
<i>Pin de Russie, Pin de Mûture, Fr.</i>	
Other Timber Tree Varieties.	
b. <i>Varieties curious or ornamental.</i> 2158	
6 <i>genevensis</i> †	
The Geneva wild Pine.	
7 <i>monophylla</i> Hodgins. †	
8 <i>scariosa</i> †	
<i>P. scariosa</i> Lodd. Cat.	
9 <i>intermedia</i> †	
10 <i>altaica</i> Ledebour †	
11 <i>tortuosa</i> Don of Forfar †	
Other Varieties, of curious or botanical interest.	
2. (s.) <i>pumilio</i> Hanke. † † Europe	
fig. 2057, 2058, 2063. 2186	
The dwarf, or Mountain, Pine.	
<i>P. sylvestris montana</i> γ Ait. Hort. Kew.	
<i>P. s. humilis</i> γ Neal.	
<i>P. cönis crectis</i> Tourn., &c.	
<i>P. humilis, sγ</i> Tourn.	
<i>P. succäticus seu carpäticus</i> Ungarish Mag.	
<i>Pinaster cönis crectis</i> Bauh.	
<i>P. tatärica</i> Mill. in Herb. Banks.	
<i>P. p. montanus</i> Park.	
<i>P. quirtus austriacus</i> Clus.	
<i>Pin nain, Fr.</i>	
<i>Krumholz, Ger.</i>	
Varieties † † - - 2186	
2 <i>rubreflora</i> † †	
3 <i>Fischeri</i> Booth. † †	
4 <i>Mughos</i> = fig. 2059, 2060, 2061. - - 2187	
The Mughos wild Pine.	
<i>P. s. Mughos Matt. Camer.</i>	
<i>P. montana</i> Baum.	
<i>P. Mughos Jacq. Poir., and N. Du Ham.</i>	
<i>P. echinata</i> Hort.	
<i>P. uncinata</i> Dec., Lodd. Cat.	
<i>Pin Mughos, Torchepin, Pin suffis, Pin crin, Pin du Briançonnais, Pin de Montagne, Fr.</i>	
<i>Bergfichte, Ger.</i>	
5 <i>M. nana</i> † † fig. 2062.	
The Kuce Pine of the Styrian Alps.	
Other Varieties.	
3. <i>Banksiana</i> Lamb. † North America	
fig. 2064, 2065, 2066, 2067. 2191	
Banks's, or the Labrador, Pine.	
<i>P. sylvestris divaricata</i> Ait.	
<i>P. rupëstris</i> Michx.	
<i>P. hudsonica</i> Lam.	
<i>Scrub Pine, Grey Pine, Hudson's Bay Pine, Ypres, Canada.</i>	
B. <i>Cones large, having the Scales furnished with Prickles.</i>	
4. <i>inops</i> Ait. † North America	
fig. 2068, 2069, 2070, 2071. 2192	
The Jersey, or poor, Pine.	
<i>P. virginiana</i> Du Roi., Mill. Diet., Wagh. Belt.	
<i>Pin chëtif, Fr.</i>	
5. <i>mitis</i> Michx. † North America	
f. 2072, 2073, 2074, 2075, 2076. 2195	
The soft-leaved, or yellow, Pine.	
<i>P. variabilis</i> Pursh; H. B., 23592.	
? <i>P. echinata</i> Mill. Diet.	
<i>New York Pine, Spruce Pine, Short-leaved Pine, Amer.</i>	
6. <i>pungens</i> Michx. † North Carolina	
fig. 2077, 2078, 2079, 2080. 2197	
The prickly-coned, or Table Mountain, Pine.	
§ ii. <i>Lariciönes.</i> †	
7. <i>Laricio</i> Poir. † Corsica	
fig. 2081, 2082, 2083, 2084. 2200	
The Corsican, or Larch, Pine.	
<i>P. sylvestris v. maritima</i> Ait. Hort. Kew., ed. 1.	
<i>P. maritima</i> Ait. Hort. Kew., ed. 2.	
Varieties † - - 2201	
1 <i>corsicana</i> †	
<i>Laricio de l'Île de Corse, Delamarre.</i>	
2 <i>subviridis</i> N. Du Ham. †	
3 <i>caramanica</i> †	
<i>P. caramanica</i> Bose.	
<i>P. caramanensis</i> Bon Jard.	
? <i>P. romäna, Lon. Hort. Soc. Gard.</i>	
<i>Laricio de Caramanic, ou de l'Asie Mineure, Delamarre.</i>	
4 <i>caläbrica</i> †	
<i>Laricio de Mont Sila en Calabre, Delamarre.</i>	
5 <i>austriaca</i> †	
<i>Laricio d'Autriche, ou de la Hongrie, Delamarre.</i>	
6 <i>pyrenaica</i> †	
<i>P. hispänica</i> Cook.	
? <i>P. pyrenäica</i> Lap.	
7 <i>täurica</i> Lodd.	
Other Varieties.	
8. (L.) <i>austriaca</i> Höss. † Austria	
fig. 2005. 2205	
The Austrian, or black, Pine.	
<i>P. nigricans</i> Hort.	
<i>P. nigrescens</i> Hort.	
<i>Schwartz Föhre, Ger.</i>	

9. (L.) *Pallasiana* Lamb. ♀ Siberia Page
 fig. 2086, 2087, 2088, 2089. 2206
 Pallas's, or the Tartarian, Pine.
P. taurica Hort.
P. tatrica, in the Hammersmith Nursery
 in 1797.
P. maritima Pall.
P. Pinca Habl. Taur.
P. halepensis Bieb., (exclusive of the Syno-
 nymes, except those of Pall. and Habl.).
P. Laricio Bieb. (Ditto.)
Tzaam, in the Tartar language.
10. (L.) *pyrenaica* Lap. ♀ S. of Spain Page
 fig. 2090, 2091, 2092, 2093. 2209
 The Pyrenean Pine.
P. hispanica Cook's Sketches in Spain.
Pinaster hispanica Roxas di San Clemente.
P. penicellus Lap. Hist. des Plantes des
 Pyrénées.
P. halepensis major Annales d'Hort. de
 Paris.
Pin Nazaron, *Pin pincau*, Fr.
11. *resinosa* Ait. ♀ Upper Canada Page
 fig. 2091, 2095, 2096, 2097. 2210
 The resinous, or red, Pine.
P. canadensis bifolia conis mediis ovatis
 Du Ham.
P. rubra Michx.
Norway Pine, in Canada.
Yellow Pine, in Nova Scotia.
Le Pin rouge de Canada, Fr.
- App. i. *Doubtful Species, apparently belong-*
ing to § ii. Lariciones. - 2213
- P. canadensis bifolia*, fol. brevioribus et tenuioribus, Du Ham. ♀
 fig. 2098, 2099. 2213
Le Petit Pin rouge de Canada, Fr.
- § iii. *Pinastri*. ♀
12. *Pinaster* Ait. ♀ South of Europe Page
 fig. 2100, 2101, 2105. 2213
 The Pinaster, or Cluster, Pine.
P. sylvestris γ Lin. Syst.
P. maritima altera Du Ham., Du Roi.
P. maritima N. Du Ham.
P. syrtica Thore Prom. sur les Côtes de
 Gascogne.
Pin de Bordeaux, *Pin des Landes*, Fr.
- Varieties* - - - 2214
 2 *escarenus* ♀ Nice.
P. escarena Risso.
 3 *Lemonianus* ♀ fig. 2102, 2103.
P. Lemoniana Ecnth.
 4 *minor* ♀ France f. 2104.
P. maritima minor N. Du Ham.
Pin Pinsot, *Pin de Mans*, *Pin à*
Trochet, Fr.
 5 *foliis variegatis* ♀
 6 *maritimus* ♀
 7 *chinensis* ♀
 8 *nepalensis* ♀
 9 *novus hollandicus* ♀
P. Nova Hollandica Lodd. Cat.,
 1836.
P. nova zealandica, in the Kew
 Arboretum.
 10 st. *helicus* ♀
 11 *Massoniâna* ♀
P. Massoniâna Lamb.

13. *Pinca* L. ♀ South of Europe Page
 fig. 2106, 2107, 2108, 2109. 2224
 The Stone Pine.
P. salva Bauh., Blackw., Du Ham.
P. domestica Matth.
Pin Pignon, *Pin bon*, *Pin cultivé*, *Pin*
Pinier, Fr.
Genéissere Fichte, Ger.
- Varieties* ♀ - - 2225
 ? 2 *fragilis* N. Du Ham. ♀
 3 *crœtica* Hort. ♀
 4 *americana* Hort. ♀
- § iv. *Halepenses*. ♀
14. *halepensis* Ait. ♀ Syria Page
 fig. 2110, 2111, 2112, 2113. 2231
 The Aleppo Pine.
P. hierosolymitana Du Ham.
P. maritima prima Mathiolius.
Pin de Jérusalem, Fr.
- Varieties* ♀ - - 2231
 2 *minor* ♀
 3 *maritima* ♀
P. maritima Lamb.
 4 *genuensis* ♀
P. genuensis Cook.
15. *brütia* Ten. ♀ Page
 fig. 2114, 2115, 2116. 2234
 The Calabrian Pine.
P. conglomerata Graefer Pl. Exsicc.
- App. i. *Species of Pine having 2 Leaves*
in a Sheath, which we cannot with certainty
refer to any of the preceding Sections.
- P. Massoniâna* Lamb., N. Du Ham., Willd., Laws., China 2236
 ? *P. nepalensis* Cels.
P. sp. from Nepal. fig. 2117.
- Sect. ii. *Ternutæ.* — *Leaves 3 in a*
Sheath. - - - 2236
- A. *Cones hardly so long as the Leaves; the Scales*
with Prickles.
- § v. *Tædæ*. ♀
16. *Tæda* L. ♀ North America Page
 f. 2118, 2119, 2120, 2121, 2122. 2237
 The Frankincense, or Lobolly, Pine.
P. foliis ternis Gron. Virg.
P. virginiana tenuifolia tripilis Pluk.
White Pine, at Petersburg and Richmond,
 in Virginia; *Oldfield Pine*, Amer.
Pin de l'Encens, Fr.
- Variety* ♀ - - 2237
 2 *alopeuroïdea* Ait. ♀
 The Fox-tail Frankincense Pine.
17. *rigida* Mill. ♀ North America Page
 fig. 2123, 2124, 2125, 2126. 2239
 The rigid, or Pitch, Pine.
P. Tæda rigida β Ait. Hort. Kew., &c.
P. canadensis trifolia Du Ham.
 ? *P. Tæda* α Poir.
 ? *Three-leaved Virginian Pine*, *Sap Pine*,
Black Pine.
Pin hérissé, *Pin rude*, Fr.

Variety	Page	Page
<i>P. Teda</i> var. \varnothing <i>alopecuröidea</i>	- 2239	§ x. <i>Canariensis</i> . $\text{♂} \text{♀}$
<i>Mil.</i> is by Mr. Lambert considered as a var. of <i>P. rigida</i> .		25. <i>canariensis</i> <i>C. Smith</i> . $\text{♂} \text{♀}$ Canaries
18. (r.) <i>serotina</i> <i>Michx.</i> $\text{♂} \text{♀}$ N. America	fig. 2127, 2128, 2129, 2130. 2212	f. 2162, 2163, 2164, 2165, 2166. 2261
The late, or <i>Pond.</i> Pine.		The Canary Pine.
? <i>Teda alopecuröidea</i> Ait.		? <i>P. adunca</i> Bosc.
<i>P. variabilis</i> <i>Lamb.</i> fig. 2131. - - - 2213		26. <i>sinensis</i> <i>Lamb.</i> $\text{♂} \text{♀}$ China
§ vi. <i>Ponderosa</i> . $\text{♂} \text{♀} \text{♂}$		fig. 2167, 2168, 2169. 2264
19. <i>ponderosa</i> <i>Doug.</i> $\text{♂} \text{♀}$ North America	fig. 2132, 2133, 2134, 2135, 2136, 2137. showing <i>Arceuthobium Oxycedri</i> <i>Bieb.</i> , <i>Fiscum Oxycedri</i> <i>Des.</i> , attached - - - 2243	The Chinese Pine.
The heavy-wooded Pine.		27. <i>insignis</i> <i>Doug.</i> $\text{♂} \text{♀}$ California
B. <i>Cones having the Scales hooked.</i>		fig. 2170, 2171, 2172. 2265
§ vii. <i>Sabiniæna</i> .		The remarkable Pine.
20. <i>Sabiniæna</i> <i>Doug.</i> $\text{♂} \text{♀}$ North America	f. 2138, 2139, 2140, 2142, 2143. 2246	28. <i>Teocote</i> <i>Schiede et Deppe</i> $\text{♂} \text{♀}$ Mount
Sabine's, or the great prickly-coned, Pine.		Orizaba - fig. 2173, 2174. 2266
21. <i>Coulteri</i> <i>D. Don.</i> ♀ Santa Lucia	fig. 2141, 2145, 2146, 2147. 2250	The Teocote, or twisted-leaved, Pine.
Coulter's, or the great hooked, Pine.		29. <i>pátula</i> <i>Schiede et Deppe</i> <i>MSS.</i> $\text{♂} \text{♀}$
<i>P. Sabiniæna</i> var. <i>Hort.</i>		Mexico - fig. 2175, 2176. 2267
<i>P. macrocarpa</i> <i>Lindl.</i> <i>MS.</i>		The spreading-leaved Pine.
22. <i>longifolia</i> <i>Rarb.</i> $\text{♂} \text{♀}$ Nepal	fig. 2149, 2150, 2151, 2152. 2252	§ xi. <i>Llaveana</i> . $\text{♂} \text{♀}$
The long-leaved Indian Pine.		30. <i>Llaveana</i> <i>Otto.</i> $\text{♂} \text{♀}$ Mexico
§ viii. <i>Gerardiæna</i> . $\text{♂} \text{♀}$		fig. 2177, 2178, 2179, 2180. 2267
23. <i>Gerardiæna</i> <i>Wall.</i> $\text{♂} \text{♀}$ East Indies	f. 2153, 2154, 2155. 2254	La Llave's Pine.
Gerard's, or the short-leaved Nepal, Pine.		App. i. <i>Species of 3-leaved Pines which cannot with certainty be referred to any of the preceding Sections, but of which there are living Plants in England.</i> - 2268
<i>P. Nèsa</i> <i>Govan.</i>		31. <i>californiæna</i> <i>Lois.</i> $\text{♂} \text{♀}$ California
<i>Eatable-seeded Pine of the East Indies.</i>		2268
? <i>Chilghöza</i> <i>Elphinstone.</i>		The Californian Pine.
C. <i>Cones long, slightly tubercled.</i>		<i>P. montereyensis</i> <i>Godefroy.</i>
§ ix. <i>Australis</i> . $\text{♂} \text{♀}$		<i>P. adunca</i> <i>Bosc.</i>
24. <i>australis</i> <i>Michx.</i> ♀ United States	f. 2156, 2157, 2158, 2159, 2160. 2255	<i>P. mouteragensis</i> <i>Hort. Soc. Gard.</i>
The southern Pine.		<i>Pm de Monterey</i> <i>Bon Jard.</i>
<i>P. palustris</i> <i>Willd., Mill., Ait., Ph., Lamb.</i>		<i>Fraseri</i> <i>Lodd. Cat.</i> - - - 2269
<i>P. americana palustris</i> , <i>Sc.</i> <i>Hort.</i>		<i>timoriensis</i> <i>Hort.</i> <i>Timor</i> - - - 2269
<i>P. serotina</i> <i>Hort.</i>		App. ii. <i>Pines supposed to have 3 Leaves, but of which the Cones only have been seen in Britain. The Cones are hooked or tubercled.</i> - - - 2269
<i>Long-leaved Pine, Yellow Pine, Pitch Pine, Amer.</i>		32. <i>muricæta</i> <i>D. Don.</i> $\text{♂} \text{♀}$ California
<i>Broom Pine, Southern States.</i>		fig. 2180. 2269
<i>Southern Pine and Red Pine, Northern States.</i>		The smaller prickly-coned Pine.
<i>Yellow Pine, Pitch Pine, Middle States.</i>		<i>Obispo, Span.</i>
<i>Georgia Pitch Pine of the Timber Merchants in England and the West Indies.</i>		33. <i>tuberculata</i> <i>D. Don.</i> $\text{♂} \text{♀}$ California
<i>Variety</i> - - - 2255		fig. 2181. 2270
2 <i>excelsa</i> ♀ North America.		The tuberculated Pine.
<i>P. palustris excelsa</i> <i>Booth.</i>		34. <i>radiata</i> <i>D. Don.</i> $\text{♂} \text{♀}$ California
		fig. 2182. 2270
		The radiated-scaled Pine.
		Sect. iii. <i>Quinæ</i> . — <i>Leaves 5 in a Sheath.</i>
		§ xii. <i>Occidentales</i> . $\text{♂} \text{♀} \text{♂} \text{♀}$
		35. <i>occidentalis</i> <i>Swartz.</i> $\text{♂} \text{♀}$ West Indies
		fig. 2183. 2271
		The West-Indian Pine.
		<i>P. folia quinæ, &c.</i> , <i>Plum. Cat., &c.</i>
		<i>Lirix americana</i> <i>Tourn.</i>

36. *Montezuma* Lamb. † † Mexico Page
 fig. 2184, 2185. 2272
Montezuma's, or the rough-branched Mexican, Pine.
P. occidentalis Kunth, Deppé in Schl. Lin.

§ xiii. *Leiophylla*. † †

37. *leiophylla* Schiede et Deppé MSS.
 † † Mexico f. 2186, 2187, 2189. 2273
 The smooth-leaved Pine.

§ xiv. *Cembra*. †

38. *Cembra* L. † Switzerland
 f. 2188, 2189, 2190, 2191, 2192. 2274
 The Cembran Pine.

- P. foliis quinis*, &c., Gmel., &c.
- P. sativa* Amm. Ruth.
- P. sylvestris*, &c., Bauh.
- P. sylvestris* Cembra Cam. Epit.
- Larix sempervirens*, &c., Breyh.
- Pinaster Aleo*, &c., Bell. Conif.
- Tæda arbor*, Cembra *Italorum*, Dale.
- Aphernousli Pine*, *Five-leaved Pine*, the *Siberian Stone Pine*, the *Swiss Stone Pine*.
- Arolcs*, in Savoy.
- Alrics*, in Switzerland.
- Cembra*, in Dauphiné.
- Cembra*, *Evau*, *Tinier*, Fr.
- Zürbelkiefer*, Ger.
- Kedr*, Russ. (See *Pall. Fl. Ross.*)

Varieties † - - 2275

- 1 *sibirica* †
 The Siberian Stone Pine, or Siberian Cedar.
P. Cembra Lodd. Cat.
Kedr, Pall.
- 2 *pygmæa* †
P. C. pumila Pall. Ross.
Stanz, Russ.
- 3 *helvetica* Lodd. Cat. ✱
 The Swiss Cembran, or Stone, Pine.

§ xv. *Ströbi*. †

39. *Ströbus* L. † North America
 fig. 2193, 2194, 2195, 2196. 2280
 The Strobus, or Weymouth, Pine.

- P. foliis quinis*, &c., Gron.
- P. canadensis quinquefolia* Du Ham.
- P. virginiana* Pluk.
- Larix canadensis* Tourn.
- New England Pine*, *White Pine*, *Pumpkin Pine*, *Apple Pine*, *Sawling Pine*, Amer.
- Pin du Lord*, *Pin du Lord Weymouth*, Fr.

Varieties † - - 2280

- 2 *alba* Hort. †
- 3 *brevifolia* Hort. †
- 4 *compræssa* Booth.
Floetbeck Weymouth Pine.
P. S. nova Lodd. Cat., ed. 1836.

40. *P. (S.) excelsa* Wallich † Nepal
 fig. 2197, 2198, 2199. 2285
 The lofty, or Bhotan, Pine.

- P. Dicksonii* Hort.
- Chilla*, or *Chylla*, Himalayas.
- Kuel*, Sirmoné and Gurhwal.
- Lemshing*, Bhotea.
- Ræsula*, or *King of the Firs*, Hindostan.

41. (*S.*) *Lambertiàna* Dougl. † N. Amer. Page
 fig. 2203, 2207. 2288
 The gigantic, or Lambert's Pine.

42. (*S.*) *monticola* Dougl. † Columbia
 fig. 2208, 2209. 2291
 The Mountain, or short-leaved Weymouth, Pine.

App. i. *Species of Pine which are not yet introduced, and of which little is known.* 2292

- P. contorta* Dougl. N. W. Amer. figs. 2210, 2211.
 The twisted-branched Pine.
- P. squamösa* Bosc. Lower Alps.
 ? *P. sylvestris* var.
- P. turbinata* Bosc. N. Amer.

II. *ABIES* D. Don. † ✱ ✱ 2105. 2293
 THE SPRUCE FIR.

- Pinus* of L. and others, in part.
- Picea* Lk.
- Picea* of the ancients.
- Sapin epicea*, Fr.
- Fichtenbaum*, Ger.
- Abiete*, Ital.
- Abieto*, Span.

§ i. *Leaves tetragonal, awl-shaped, scattered in insertion.* D. Don.

1. *excelsa* Dec. † Norway f. 2212. 2293
 The lofty, or Norway, Spruce Fir.

- A. communis* Hort.
- A. Picea* Mill., Michx.
- A. foliis solitariis*, &c., Hort. Cliff., &c., Hall.
- P. Abies* L. Sp. Pl., &c.
- P. Picea* Du Roi.
- P. excelsa* Lam.
- Common Spruce*, *Prussian Fir*.
- Faux Sapin*, *Epicea*, *Sapin-Pessc*, *Scrcnte*, *Sapin gentil*, *Pinessc*, Fr.
- Lafic*, in the Vosges.
- Gemeine rothe Tanne*, Ger.

Varieties † ✱ ✱ - - 2294

- 1 *communis* †
 The common Spruce, or White Fir of Norway.
- 2 *nigra* † fig. 2213.
 The black-leaved Spruce, or Red Fir of Norway.
- 3 *carpatica* †
 The Carpathian Spruce Fir.
A. carpatica Hort., and Hort. Brit.
- 4 *péndula* †
 The pendulous-branched, or weeping, Norway Spruce Fir.
A. communis péndula Booth.
Pinus Abies péndula Lodd. Cat.
- 5 *foliis variegatis* †
 Blotched-leaved Spruce Fir.
Pinus Abies foliis variegatis Lodd. Cat.
- 6 *Clanbrasiliana* ✱
Lord Clanbrasil's Spruce Fir.
P. Clanbrasiliana Lodd. Cat.
- 7 *Clanbrasiliana stricta* ✱
 Upright-growing Lord Clanbrasil's Spruce Fir.
- 8 *pygmæa* ✱
 The pygmy Spruce Fir.
A. nova Lond. Hort. Soc. Gard.
A. elegans Sm. of Ayr.
- 9 *tenuifolia* ✱
 The slender-leaved Spruce Fir.
A. tenuifolia Sm. of Ayr.

	Page	Page
10 gigantèa † The gigantic Spruce Fir. A. gigantèa Sm. of Ayr.		
11 monstrosa ■ The monstrous Spruce Fir. A. monstrosa Hort.		
Other Varieties. -	- 2295	
2. álba Michx. † N. Amer. f. 2224. 2310		
The white Spruce Fir. Pinus álba Ait., W., Lamb. Pinus láxa Ehrh. Pinus canadénsis Du Roi. Abies Picca fóliis, &c., Hort. Angl., Duh. A. currifolia Hort. Single Spruce, Amer. E'pinette blanche, Fr. Sapinette blanche, Fr.		
Variety † -	- 2311	
2 nána Dickson † Other Varieties.		
3. nìgra Ait. † North America fig. 2225, 2226, 2227. 2311		
The black Spruce Fir. Pinus nìgra Ait., W., Lamb., Du Roi. Pinus mariàna Ehr. Abies mariàna Wagh. Double Spruce.		
4. (n.) rùbra Poir. † Nova Scotia fig. 2228. 2316		
The red Spruce Fir, or Newfoundland red Pine. Pinus americàna rùbra Wagh. Pinus rùbra Lamb.		
Variety † -	- 2316	
2 cærùlea † A. cærùlea Booth.		
5. Smithiana Wall. † Himalayas fig. 2229. 2317		
Smith's, or the Himalayan, Spruce Fir. Pinus Smithiana Wall., Lamb. Pinus Khùtrow Royle. A. Morinda Hort. Kaga, or Kaggoc, in the Parbutee language.		
Varieties -	- 2317	
6. (? c.) orientális † Trebisonde - 2318		
The Oriental Spruce Fir. Pinus orientális Lamb., L., Vitman.		
§ ii. Leaves flat, generally glaucous beneath, imperfectly 2-roued.		
7. Douglàsii Lindl. † North America fig. 2230. 2319		
Douglas's, or the trident-bracted, Spruce Fir. Pinus taxifolia Lamb., Pursh. A. californica Hort. Pinus Douglàsii Sabine MSS., Lamb. Pin., iii. 90. The Nootka Fir, Sm. in Rees's Cyc.		
Varieties † -	- 2319	
2 taxifolia † The Yew-leaved trident-bracted Spruce Fir.		
8. Menzièsii Dougl. † California fig. 2232. 2321		
Menziès's, or the warped-branched, Spruce Fir. Pinus Menzièsii Lamb.		
9. canadénsis L. † N. America - 2322		
The Canada Pine, or Hemlock Spruce Fir. Pinus canadénsis L. Pinus americàna Du Roi, Ait., Lamb., Sm. Pinus Abies americàna Marsh. Perusse, by the French in Canada. Sapin du Canada, Fr. Schiertings Fichte, Ger.		
10. dumòsa Lamb. † Nepal fig. 2233, 2234. 2325		
The bushy alpine Spruce Fir. Pinus dumòsa Lamb. A. Bruoniàna Lindl. in Penny Cyc. Pinus decidua Wall, MSS. Pinus Bruoniàna Wall. Pl. As. Rar.		
11. cephalónica † Cephalonia fig. 2235, 2236. 2325		
The Cephalonian Silver Spruce Fir. Koukounaria, and also Elatos, in Cepha- lonia. A. taxifolia Hort. A. Inscombèna Hort. The Mount Enos Fir.		
App. i. Species of Abies of which little more is known than their Names. - 2329		
A. obovata D. Don. MS. Altai Mountains. Picea obovata Led. ? A. Smithiana var. D. Don. A. Mertensiana Bong. N. Amer. A. sitchensis Bong. N. Amer. A. trigona Raf. Oregon country. A. heterophylla Raf. Oregon country. A. aromatica Raf. Oregon country. A. microphylla Raf. Oregon country. A. obliquata Raf. Oregon country. A. filicèta Raf. Oregon country. A. hirtella Thunb. A. Kämpferii Thunb. A. Thunbergii Thunb. A. Mitrni Sieb. A. Torino Sieb. A. Araragi Sieb.		
III. PICEA D. Don. † - 2105. 2329		
THE SILVER FIR. Pinus L., in part. Abies Link. Nees Von Esenbeck, and Led. Abies Du Roi, in part. Sapin, Fr. Tannen, Ger.		
1. pectinàta † Europe f. 2237, 2238. 2329		
The comb-like-leaved Silver Fir. Abies of Pliny. Pinus Picca L., &c. Pinus Abies Du Roi. Abies álba Mill. Diet., &c. Abies Vári fólio Tourn., &c. Abies vulgàris Poir. Abies pectinàta Dec., Duh. A. bies taxifolia Hort. Par. Abies Picca Lindl. in Penny Cyc. A. bies cæcisa Lk. Spanish Fir. Sapin commun, Sapin à Feuilles d'If, Sapin blanc, Sapin argenté, Sapin cu Peigne, Sapin de Normandie, Fr. Weiss Tanne, Edeltanne, Ger.		
Varieties † -	- 2330	
2 tortuòsa † The twisted-branched Silver Fir.		
3 fóliis variegàtis † The variegated-leaved Silver Fir.		
4 cinèrea † The cinereous Silver Pine. Pinus Picea cinèrea Baum.		

2. (p.) *Pichta* ♀ Altai Mountains 2338 Page
 The Pitch Silver Fir. 2338
Pinus Pichta Lodd. Cat.
Pinus sibirica Hort.
Abies sibirica Led., Lindl. in Penny Cyc.
Abies Pichta Fischer.
Pichta, Russ.
3. balsamea L. ♀ North America
 fig. 2240, 2241. 2339
 The Balm of Gilead, or American, Silver Fir.
Pinus balsamea L., &c.
Abies balsamea Marsh.
Abies Taxis folio, &c., Hort. Angl., &c.
Abies balsamifera Michx.
Balsam Fir.
Le Beaucme de Giléad, le Sapin Baumier de
Giléad, Fr.
Balsam Fichte, Balsam Tanne, Ger.
- Variety ♀ - - 2339
- 2 longifolia Booth. ♀
 The long-leaved Balm of Gilead Silver
 Fir.
4. (b.) Fraseri Ph. ♀ Carolina
 fig. 2243, 2244. 2340
Fraser's, or the double Balsam, Silver Fir.
Pinus Fraseri Ph., Lamb.
Abies Fraseri Lindl. in Penny Cyc.
5. grandis Dougl. ♀ N. W. America
 fig. 2245, 2246. 2341
 The great Silver Fir.
Pinus grandis Dougl. MS., Lamb.
Abies grandis Lindl. in Penny Cyc.
The great Californian Fir.
6. amabilis Dougl. ♀ ? California
 fig. 2247, 2248. 2342
 The lovely Silver Fir.
Pinus amabilis Dougl. MS.
7. nobilis Dougl. ♀ North America
 fig. 2249, 2250. 2342
 The noble, or large-bracted, Silver Fir.
Pinus nobilis Dougl. MS., Lamb.
Abies nobilis Lindl. in Penny Cyc.
8. Webbiana Wall. ♀ Nepal
 fig. 2251, 2252, 2253. 2344
 Webb's purple-coned Silver Fir.
Pinus Webbiana Wall., Lamb. Pin.
Pinus spectabilis Lamb. Monog.
Abies Webbiana Lindl. in Penny Cyc.
Chilrow, and the Oonum, or purple-conced
Fir, in the Himalayas.
9. Pindrow Royle ♀ Kamaon
 fig. 2254, 2255. 2346
 The Pindrow, or tooth-leaved, Silver Fir.
Pinus Pindrow Royle, Lamb.
Taxus Lambertiana Wall. Cat.
Pindrow, and sometimes Morinda, in the
 Himalayas.
10. bracteata D. Don. ♀ California
 fig. 2256. 2348
 The leafy-bracted Silver Fir.
Pinus bracteata Lin. Trans., Lamb.
Pinus venusta Dougl. in Comp. to Bot. Mag.

11. religiosa H. et K. ♀ Mexico
 fig. 2257. 2349
 The sacred Mexican Silver Fir.
Pinus religiosa Humb. et Kunth, &c.
Abies religiosa Lindl. in Penny Cyc.
- ? P. hirtella - - - 2349
Abies hirtella Lindl. in Penny Cyc.
Pinus hirtella Thunb. et Kunth.
- IV. LARIX Tourn. ♀ 2105. 2350
 THE LARCH.
Pinus L. and others.
Abies Rich.
Melèze, Fr.
Lerchenbaum, Ger.
Laricio, Ital.
1. europæa Dec. ♀ South of Europe
 fig. 2258. 2350
 The European, or common, Larch.
Pinus Larix L., &c.
Abies Lin. Hort. Cliff.
Larix decidua Mill. Dict.
Larix folio decidua, &c., Bauh.
Larix Bauh. Pin., Dod., Cam.
Abies Larix Lam.
Melèze commune, Fr.
Lörche, Lorcher-Fichte, Gemeiner Lerchen-
baum, Terbentinbaum, Europäische Ce-
der, Weisser Lerchenbaum, Ger.
- Varieties ♀ - - 2350
- 1 communis Laws. ♀
 The common European Larch.
- 2 laxa Laws. ♀
 The loose-headed European Larch.
- 3 compacta Laws. ♀
 The compact, or crowded-branched,
 Larch.
- 4 pendula Laws. ♀
 The weeping-branched European
 Larch.
The weeping Larch from the Tyrol,
 Hort. Trans.
- 5 repens Laws. ♀
 The creeping-branched European
 Larch.
- 6 flore rubro ♀
 The red, or pink, flowered common
 Larch.
- 7 flore albo ♀
 The white-flowered Larch from the
 Tyrol.
- 8 sibirica ♀
 The Russian Larch.
L. sibirica Fisch.
 ? *L. archangelica* Laws.
L. rossica Sab.
Pinus L. sibirica Lodd. Cat.
- 9 dahurica ♀
 The Dahurian Larch.
L. dahurica Laws.
- 10 intermedia ♀
 The intermediate, or Altaian, Larch.
L. intermedia Laws.
Pinus intermedia Lodd. Cat.
- Other Varieties - - 2353
- L. Fraseri Comp. to Bot. Mag.
2. americana Mx. ♀ N. America 2399
 The American Larch.
Pinus laricina Du Roi.
Pinus microcarpa W.
Abies microcarpa Poir.
Hackmatack, Amer.
Tamarack, by the Dutch in New Jersey.
E'pinette rouge, in Canada.

	Page
<i>Varieties</i> † - -	2400
1 <i>rùbra</i> †	
The small red-coned American Larch.	
L. <i>microcárpa</i> Laws.	
Pinus <i>microcárpa</i> Pursh, Lodd. Cat.	
Epinette rouge, Canada.	
2 <i>péndula</i> †	
The black pendulous-branched American Larch.	
L. <i>péndula</i> Laws.	
Pinus <i>péndula</i> Ait., &c.	
Pinus <i>intermèdia</i> Du Roi.	
Pinus <i>lârix nigra</i> Marsh.	
A <i>bies péndula</i> Poir.	
Tamarack, Amer.	
3 <i>prolifera</i> †	
The proliferous-branched Larch.	
L. <i>prolifera</i> Maleolin.	

V. CÉDRUS Barrel. † ■ 2105. 2402

THE CEDAR.
Pinus L., in part.
A *bies* Poir., in part.
Lârix Tourn., in part.
Cèdre, Fr.
Ceder, Ger.

1. Libani Barr. † Mount Lebanon fig. 2267. 2402

The Cedar of Lebanon.
Pinus Cèdrus L. and others.
Pinus *foliis fasciculatis*, &c., Du Roi.
Lârix Cèdrus Mill. Dict.
Lârix *orientâlis* Tourn., Duh.
Cèdrus *mûgna* Dod.
C. *confera* Bauh., Ray.
C. *phœnicea* Reneauh.
Cèdrus Bell.
A *bies Cèdrus* Poir., N. Du Ham., Lindl.

Varieties † ■ - - 2402

2 *foliis argenteis* †

The silvery-leaved

3 *nana* † ■

The dwarf Cedar.

Other Varieties.

2. Deodâra Roxb. † Nepal fig. 2283, 2284, 2285, 2286. 2428

The Deodâra, or Indian, Cedar.
Pinus Deodâra Lamb.
A *bies* Deodâra Lindl.
Devadâra, or Deodâra, Hindostance.
The sacred Indian Fir.

Varieties, or nearly allied Species 2429

Shinlik, Moorcroft. Forests of Ladakh.

Christa rooro, Moorcroft. Forests of Ladakh.

VI. ARAUCARIA R. et P. † 2105. 2432

THE ARAUCARIA.
Eutassa Sal.
Colymbèa Sal.
Dombèya Lamb.
Cupressus Forst.
The southern Pine.

1. imbricatâ Pav. † Chili fig. 2286, 2293. 2432

The imbricate-leaved Araucaria, or Chili Pine.
A. Dombèyi Rich., Lindl. in Penny Cic.
Pinus Araucaria Mol.
Dombèya chilensis Lam.
Pino de Chili, Span.
Peghuén, in the Andes.
Sir Joseph Banks's Pine.

2. brasiliâna Rich. † Brazils fig. 2294, 2295, 2296. 2439

The Brazil Araucaria, or Brazil Pine.

3. excelsa Ait. † Norfolk Island fig. 2297. 2301. 2440

The lofty Araucaria, or Norfolk Island Pine.
Eutassa heterophylla Sal.
Cupressus columnaris, &c., Forst.
Dombèya excelsa Lamb.
Attingia excelsa Loud. Hort. Brit.
Pin de Norfolk, Fr.

4. Cunninghami Ait. † New Holland fig. 2303. 2305. 2443

Cunningham's Araucaria, or the Morcton Bay Pine.
Attingia Cunninghami G. Don, in Loud. Hort. Brit.

VII. CUNNINGHAMIA R. Br. † ▭ 2105. 2445

THE CUNNINGHAMIA.
Pinus Lamb.
Bêtis Salisb.

1. sinensis Rich. † ▭ China fig. 2306, 2307. 2445

The Chinese Cunninghamia, or broad-leaved Chinese Fir.
Bêtis jacutifolia Salisb.
Pinus lanceolata Lamb.
A *bies major sinensis*, &c., Pluk.
Cunninghamia lanceolata R. Br.
Araucaria lanceolata Hort.

VIII. DAMMARA Rumph. † ▭ 2105. 2447

THE DAMMARA, or Amboyna, Pine.
Pinus Lamb.
Agathis Sal.

1. orientâlis Lamb. † ▭ Amboyna fig. 2308, 2309. 2447

The Oriental Dammara Pine, or Amboyna Pitch Tree.
Pinus Dammara W., Lamb., Ait.
Dammara albi Rumph.
Agathis toranrifolia Sal. in Lin. Trans.
A. Dammara Rich.
Arbor javanensis, &c., Ray.

2. australis Lamb. † ▭ New Zealand fig. 2310, 2311. 2448

The Southern Dammara, or Kauri, Pine.
Agathis australis Lindl.
Kauri Tree, New Zealand Pitch Tree, Kauri Pine.

App. i. A tabular View of the principal Pinctums, or Collections of Abietina, in Europe. - - - 2449

Sect. II. CUPRESSINÆ. - 2453

† † ▭ ▭ ■

IX. THUJA L. † † ▭ ▭ ■ 2105. 2454

THE ARBOR VITÆ.
Thuja, or Arbre de Vie, Fr.
Lebensbaum. Ger.

§ i. *Thuja vèra* D. Don.

Page

1. occidentàlis L. ♀ Canada
 fig. 2312. 2314. 2454
 The western, or American, Arbor Vitæ.
Thuja Theophrasti Bauh.
Arbor Vitæ Clus.
 White Cedar, Amer.
Cèdre américain, *Cèdre blanc*, *Arbre de*
Fig., Fr.
Gemeiner Lebensbaum, Ger.
Albero de Vita, Ital.

Varieties ♀ - - 2454

- 2 variegata ♀
 The variegated-leaved Arbor Vitæ.
T. o. foliis variegatis Lodd. Cat.

- 3 odorata Marsh. ♀
 The sweet-scented Arbor Vitæ.

2. (o.) plicata Donn. ♀ N. Amer. 2458
 The plicate, or Nee's, Arbor Vitæ.

3. chilensis Lamb. ♀ Chili - 2458
 The Chili Arbor Vitæ.
Cupressus thyoides Pavon MSS.

§ ii. *Biota* D. Don.

4. orientalis L. ♀ China fig. 2215. 2459
 The Oriental, or Chinese, Arbor Vitæ.

Varieties ♀ - - 2459

- 2 stricta Hort.
 The fastigate Arbor Vitæ.
T. pyramidalis Bauh. Cat.

- 3 tatarica ♀
 The Tartarian Arbor Vitæ.
T. tatarica Lodd. Cat.

§ iii. *Cyprissa* D. Don.

5. cupressoides L. ♀ C. G. H.
 fig. 2316. 2460
 The Cypress-like, or African, Arbor Vitæ.
T. aphylla Burm.

6. pensilis Lamb. ♀ China - 2460
 The pensile Arbor Vitæ.

7. pendula Lamb. ♀ Tartary
 fig. 2317, 2318. 2461
 The pendulous-branched, or weeping, Arbor
 Vitæ.

App. i. Species not sufficiently known to be
 referred to any of the preceding Sections.

T. dolobrata Lin. Suppl., Thunb., W., Lamb. Japan - 2462
Quai, vulgo *Fi no ki*, and *Ibuki*, Kamf.
 Other Species

- X. CALLITRIS Vent. ♀ 2105. 2462
 THE CALLITRIS.
Thuja L., in part.
Fresnelia Mirbel.

Page

1. quadrivalvis Vent. ♀ Barbary
 fig. 2319. 2462

The four-valved Callitris.
Thuja articulata Desf.
Cupressus fructu quadrivalvi Shaw.
Thuja ströbilis tetragonis, &c., Vahl.

- C. Fothergillii ♀ - - - 2464
 ? *Cupressus Fothergillii*.
 C. triquetra ♀ C. G. H. - - - 2464
Cupressus triquetra Lodd. Cat.
 C. cupressiformis Vent. ♀ N. Holl. - - - 2464
 C. macrostachya Hort. ♀ - - - 2464

App. i. Species of *Callitris* Natives of Aus-
 tralia, and not yet introduced into Britain.

- C. rhomboides R. Br. ♀ N. Holl. - - - 2464
 C. oblonga Rich. ♀ Pt. Jackson - - - 2464
C. fruticosa R. Br.

XI. CUPRESSUS L. ♀ 2105. 2464

THE CYPRESS.

Cyprès, Fr.
Cypresse, Ger.
Cipresso, Ital.
Ciproste, Port.
Cypros, Hungarian.

1. sempervirens L. ♀ S. Europe
 fig. 2320. 2464

The evergreen, or common, Cypress.
C. pyramidalis Hort.
Cyprès pyramidal, *Cyprès ordinaire*, Fr.
Gemeine Cypressenbaum, Ger.
The Italian Cypress.

Varieties ♀ - - - 2465

- 1 stricta Mill. Dict.
Cyprès mâle, Fr.
 2 horizontalis Mill. Dict.
C. horizontalis N. Du Ham.
C. expansa Hort. Par.

2. thyoides L. ♀ N. Amer. f. 2327. 2475
 The Thuja-like Cypress, or White Cedar.

C. nana marihana, &c., Pluk.
Thuja sphaeroidalis, Rich.
Cyprès faux Thuja, Fr.

Variety ♀ - - - 2475

- 2 foliis variegatis ♀

3. lusitânica Tourn. ♀ Goa f. 2328. 2477
 The Portuges Cypress, or Cedar of Goa.

C. glauca Brot.
C. pendula L'Herit., ? not of Thunb.
Cedar of Bussaco.

4. torulosa Lamb. ♀ Nepal
 fig. 2329. 2331. 2478
 The twisted, or Bhotan, Cypress.

5. pendula Thunb. China
 fig. 2332. 2334. 2479
 The weeping Cypress.

App. i. Kinds of *Cupressus* of which there
 are Plants in British Gardens, but of which
 very little is known. - - - 2480

C. horizontalis Audibert.
 C. expansa Audibert.
 ? *C. expansa* Hort. Par.
 C. Fothergillii Lee.
 C. thurifera Hort. Soc. Gard.
 C. Tournefortii Audibert.
 C. hacciformis Willd.
 C. australis Pers.

App. ii. *Kinds of Cupressus not yet introduced.* - Page 2480

C. nootkatensis Lamb.
C. japonica Thunb.

XII. TAXODIUM Rich. ☿ ♀ ♀ ♂ 2105. 2480

THE TAXODIUM, or *Deciduous Cypress.*
Cupressus L.
Schubertia Mirb.
Condylocarpus Salisb.

1. *distichum* Rich. ☿ N. Amer. f. 2335. 2481

The two-ranked-leaved Taxodium, or *Deciduous Cypress.*

Cupressus disticha L., &c.
Cupressus americana Cat. Carol.
Cupressus virginiana Comm. Hort.
Schubertia disticha Mirb.
Bald Cypress, Cypress, Amer.
Cyprès de l'Amérique, Cyprès chauve, Fr.
Zweyzeitige Cypress, Ger.

Varieties ☿ - - 2481

- 1 *pätens* Ait. ☿
- 2 *nütans* ☿ fig. 2336—2338.
The long-leaved *Deciduous Cypress.*
T. d. pëndulum Lond. Hort. Brit.
- 3 *excëlsum* Booth. ☿
- 4 *sinënsë* ☿
T. sinënsë Noisette.
- 5 *sinënsë pëndulum* ☿
T. sinënsë pëndulum Hort.

T. sempervirens Lamb. ♀ N. Am. figs. 2339, 2340. 2487
The evergreen Taxodium.

T. capënsë ♀ ♂ C. G. H. - - 2487
The Cape, or *African, Taxodium.*
Cupressus juniperoides L.

XIII. JUNIPERUS L. ♀ ♀ ♂ ♂ ♂ ♂ 2106. 2487

THE JUNIPER.
Sabina Bauh.
Cedrus Tourn.
Genévrier, Fr.
Wachholder, Ger.

§ i. *Oxycedri.*—*Leaves spreading in the adult Plants.* D. Don.

1. *communis* L. ♂ Europe, America, and Asia - fig. 2343, 2344. 2489
The common Juniper.

J. No. 1661 a., Hall.
J. vulgaris, &c., Ray, &c.
J. minor Fuchs, Dalech.
J. communis saxatilis Pall.
J. alpina Clus.
J. minor montana C. Bauh.
Genévrier commun, Fr.
Gemeiner Wachholder, Ger.

Varieties ♂ - - 2489

- 1 *vulgäris* Park. ♂
J. v. fruticosa Bauh.
J. c. cretensis Pursh.
- 2 *succëica* Murl. ♂ fig. 2343.
The Swedish, or *trunc.* Juniper.
J. succëica Mill. Dict.
J. vulgaris arbor Bauh.

- 3 *nana* Willd. ♂ fig. 2344.
J. communis β Fl. Br., &c.
J. c. saxatilis Pall.
J. No. 1661. Hall.
J. alpina Ray, &c.
J. alpina minor Ger. Emac.
J. minor montana, &c., Bauh., N. Duh.
J. nana Sm. Eng. Fl.
J. sibirica Hort.
J. daurica Hort.
J. c. montana Ait.
 - 4 *oblönga* ♂ fig. 2346.
J. oblönga Hort.
 - 5 *oblönga pëndula* ♂ fig. 2345.
 - 6 *canadënsis* ♂ fig. 2347.
J. canadënsis Lodd. Cat.
 - 7 *depressa* Pursh. ♂
J. canadënsis Lodd. Cat.
- Other Varieties.

2. *Oxycedrus* L. ♂ Spain, Portugal, and France - fig. 2351, 2352. 2494

The Sharp Cedar, or *brown-berried, Juniper.*
J. major Cam. Epit.
J. m. monspeliënsium Lob. Ic.
J. phœnicea, &c., J. Bauh.
J. major, &c., C. Bauh., &c.
Cedrus phœnicea Matth.
Oxycedrus Clus.
Oxycedrus phœnicea Dod.
The prickly Cedar.
Le Cade, Fr.
Spanische, Ger.

3. *macrocarpa* Sm. ♂ Greece f. 2353. 2494
The large-fruited Juniper.
J. major, bicca cerulea, Tourn.

4. *drupacea* Lab. ♂ ♂ Syria fig. 2354, 2356. 2495.
The drupaceous, or *large-fruited, Juniper.*

5. *virginiana* L. ♀ North America fig. 2357. 2495

The Virginian Juniper, or *Red Cedar.*
J. major americana Ray.
J. maxima, &c., Sloane.

Varieties ♀ - - 2495

- 2 *hümilis* Lodd. Cat. ♂
 - 3 *caroliniana* ♀
J. caroliniana Du Roi.
- Other Varieties

6. *bermudiäna* L. ♀ Bermudas fig. 2357. 2498

The Bermudas Cedar.
Cedrus Bermuda Ray's Letters.

J. nepalënsis Hort. - - 2499
Cupressus nepalënsis Hort.

§ ii. *Sabina.*—*Leaves of the adult Plant imbricated.* D. Don.

7. *Sabina* L. ♂ Spain, &c. fig. 2359—2363. 2499

The common Savin.

Varieties ♂ ♂ - - 2499

- 1 *cupressifolia* Ait. ♂ fig. 2359.
The Cypress-leaved Savin.
J. lusitanica Mill. Dict.
Sabina Dod.
Sabina folio Cupressi Bauh., Duh., Ray.
La Sabine mâle, Fr.

	Page
2 tamariscifolia Ait. * fig. 2360.	2500
The Tamarisk-leaved, or berry-bearing, Savin.	
Sabina folio Tamarisci Dioscoridis Bauh.	
J. Sabina Mill. Diet.	
La Sabine femelle, Fr.	
3 foliis variegatis Mart.	
4 prostrata * fig. 2361, 2362.	
J. prostrata Michx.	
J. repens Nutt.	
J. hudsonica Lodd. Cat.	
5 alpina * fig. 2363.	
J. alpina Lodd. Cat.	
8. daurica Pall. * Dauria	2500
fig. 2364, 2365.	
The Daurian Juniper.	
9. phœnicea L. † S. Europe, &c.	2501
fig. 2366.	
The Phœnician Juniper.	
J. major Dioscoridis Clus.	
Cedrus phœnicea media Lob.	
Cedrus lycia reclusa J. Bauh.	
Cedrus folio Cupressi major, &c., C. Bauh.	
Oxycedrus lycia Dod.	
Genévrier de Phénicie, Fr.	
Dichtnadeliger Wachholder, Ger.	
10. (p.) lycia L. * South of Europe	2502
fig. 2367.	
The Lycian Juniper.	
J. p. β lycia N. Du Ham.	
Cedrus phœnicea altera Plinii et Theophrasti Lob.	
C. folio Cupressi, &c., C. Bauh.	
11. thurifera L. † Spain	2503
fig. 2368.	
The incense-bearing, or Spanish, Juniper.	
J. hispánica Mill. Diet.	
Cedrus hispánica, &c., Tourn.	
12. excelsa Willd. † Himalayas	2503
fig. 2369.	
The tall Juniper.	
J. Sabina var. Pall.	
Himalaya Cedar-wood.	
13. squamata D. Don. * Nepal	2504
- The scaled Juniper, or creeping Cedar.	
14. recurva Ham. * Nepal	2504
f. 2370.	
The recurved Nepal Juniper.	
15. uvifera D. Don. * Cape Horn	2504
- The grape-bearing, or large-fruited, Juniper.	
16. barbadensis L. † West Indies	2504
- The Barbadoes Juniper.	
J. bermudiana Hort.	
Jamaica Berry-bearing Cedar.	
17. chinensis L. * China	2505
fig. 2371, 2372.	
The Chinese Juniper.	
J. c. 2 Smithii ? †	2505
- fig. 2373.	
App. ii. Kinds of Juniperus mentioned in Books, but of which very little is known. 2505	

J. frigidissima Willd.
J. capensis Lam.

Empetræcæ. * * 2506

I. EMPETRUM L. * - 2506
THE CROWBERRY.

1. nigrum L. * Britain
fig. 2374, 2375. 2507
The black Crowberry, or Crakeberry.
E. montanum fructu nigro Ray.
Erica Matth.
Erica baccifera Cam. Epit.
E. coccifera procumbens Ger. Emac.
E. Còris folio undecima Clus.
- Variety * - 2507
- 2 scoticum Hook. * -

2. rubrum L. * South America
fig. 2376, 2377. 2507
The red-fruited Crowberry.
Cranberry of Staten Island.

II. COREMA D. Don. * - 2506. 2508
THE COREMA.
Empetrum L., in part.

1. alba D. Don. * Portugal 2508
The white-berried Corema.
Empetrum album L., &c.
Empetrum lusitanicum, &c., Tourn.
Erica cretæis, &c., Bauh. Pin.
The white-berried Heath.
Portugal Crakeberry.

III. CERATIOLA Mr. * 2506. 2508
THE CERATIOLA.

1. ericoides * South Carolina
fig. 2378, 2379. 2509
The Erica-like Ceratiola.

Smilacæcæ. * † † 2509

I. SMILAX L. * † † - 2510
THE SMILAX.

§ i. Stems prickly and angular.

1. aspera L. † S. France f. 2380. 2510
The rough Smilax.
Rough Bindweed.
Smilar, Fr. and Ger.

Varieties † - 2510

2 auriculata Ait. †
Leaves ear-shaped at the base.

3 mauritanica
S. mauritanica Poir.

2. excelsa L. † Syria fig. 2381. 2511
The tall Smilax.
S. orientalis, &c., Tourn.
S. aspera Alp. Ægypt.

3. rubens Wals. † N. Amer. f. 2382. 2511
The red-berried Smilax.

	Page
4. <i>Sarsaparilla</i> L. \mathbb{A} N. and S. America fig. 2383. 2511	16. <i>virginiana</i> Mill. \mathbb{A} Virginia - 2515 The Virginian Smilax.
The medicinal Smilax, or Sarsaparilla. <i>S. aspera peruviana</i> Sarsaparilla Bauh. <i>S. peruviana</i> Park. <i>S. glauca</i> Michx. The glaucous-leaved Smilax.	17. <i>pubera</i> Willd. \mathbb{A} North America 2515 The downy Smilax. <i>S. pumila</i> Walt.
5. <i>hastata</i> Willd. \mathbb{A} Carolina - 2512	App. i. <i>Kinds of Smilax which are either not introduced, or of which we have not seen the Plants.</i> - - - 2515
The spear-shaped-leaved Smilax. <i>S. Bona nox</i> Michx. <i>S. aspera</i> var. Lam.	<p><i>S. ovata</i> Ph. Georgia <i>S. alba</i> Ph. Carolina. <i>S. pandurata</i> Ph. N. Amer. <i>S. nigra</i> W. Spain. ? <i>The black-berried var. of S. aspera.</i> <i>S. catalonica</i> Poir. Spain. <i>S. horrida</i> Desf. N. Amer. <i>S. glauca</i> Walt. N. Amer. ? <i>S. Sarsaparilla</i> var. <i>S. alpina</i> W. Greece.</p>
Variety \mathbb{A} - - 2512	
2 lanceolata Ph. \mathbb{A} fig. 2383. <i>S. lanceolata</i> Walt. ? <i>S. longifolia</i> Wats.	
6. <i>Walteri</i> Ph. \mathbb{A} Virginia - 2512	
Walter's Smilax. <i>S. China</i> Walt.	
§ ii. <i>Stem prickly, round.</i>	
7. <i>China</i> L. \mathbb{A} China - - 2513	
The Chinese Smilax. <i>China radix</i> Bauh., &c. <i>C. vulgaris officinarum</i> Ger. Emac. <i>Smilax aspera</i> , minor Plum. Sankira, vulgo Quaquara, <i>Kempf.</i>	
8. <i>rotundifolia</i> L. \mathbb{A} N. America 2513	
The round-leaved Smilax.	
9. <i>aurifolia</i> L. \mathbb{A} N. America - 2513	
The Laurel-leaved Smilax. <i>S. altera</i> , &c., Plum. Ic. <i>S. lucida</i> , &c., Catesb.	
10. <i>lanuoides</i> L. \mathbb{A} N. America - 2513	
The black Bryony-like Smilax. <i>S. Bryonia nigra</i> , &c., Catesb.	
11. <i>caduca</i> L. \mathbb{A} S. America - 2514	
The deciduous Smilax.	
§ iii. <i>Stalks unarmed, 4-angled.</i>	
12. <i>Bona nox</i> L. \mathbb{A} N. America - 2514	
The Bona-nox, or ciliated, Smilax. <i>S. aspera India occidentalis</i> Bauh. <i>S. foliis latis</i> , &c., Pluk. <i>S. variegata</i> Walt.	
13. <i>latifolia</i> R. Br. \mathbb{A} N. Holl. - 2514	
The broad-leaved Smilax.	
14. <i>quadrangularis</i> Michx. \mathbb{A} N. America fig. 2385. 2514	
The four-angled Smilax.	
§ iv. <i>Stems unarmed, round.</i>	
15. <i>lanceolata</i> L. \mathbb{A} Virginia, &c. 2515	
The lanceolate-leaved Smilax. <i>S. non-spinosa</i> , &c., Cat.	
Liliaceæ. \mathbb{A} \mathbb{B} \mathbb{C} \mathbb{D} \mathbb{E} 2515	
I. ASPARGAGUS L. \mathbb{B} \mathbb{C} \mathbb{D} - 2516	
THE ASPARGAGUS. <i>Sarmentaceæ</i> , part of, Juss., Nees Von Esenbeck. <i>Asperges</i> , Fr. <i>Spargel</i> , Ger. <i>Coraleruyt</i> , Dutch. <i>Sperage</i> , Old English.	
1. <i>scandens</i> Willd. \mathbb{E} C. G. H. - 2516	
The climbing Asparagus.	
2. <i>albus</i> L. \mathbb{C} Candia - - 2516	
The white Asparagus. <i>A. spinosus</i> , &c., Park. <i>A. sylvestris</i> , &c., Ger. <i>Corruda tertia</i> Clus. <i>Wild Thong Sperage.</i>	
3. <i>aphyllus</i> \mathbb{B} S. Europe - 2517	
The prickly Asparagus. <i>Corruda altera</i> Clus. <i>A. petraea</i> , <i>Prickly Roche Asparagus</i> , Park. Variety - - - 2517 2 <i>creticus fruticosus</i> , &c., <i>Tourn.</i>	
4. (a.) <i>acutifolius</i> L. \mathbb{B} Spain - 2517	
The acute-leaved Asparagus. <i>A. Corruda Scop.</i> , &c. <i>A. foliis acutis</i> Bauh. <i>A. sylvestris</i> Cam. <i>A. petraeus</i> Ger. <i>Corruda l. Clus.</i>	
5. <i>horridus</i> L. \mathbb{E} ? Spain - - 2517	
The horrid-spined Asparagus. <i>A. hispanica</i> Tourn.	
II. RUSCUS L. \mathbb{B} \mathbb{C} \mathbb{D} \mathbb{E} 2516. 2517	
THE BUTCHER'S BROOM. <i>Fragon</i> , Fr. <i>Mäusdorn</i> , Ger.	
1. <i>aculeatus</i> L. \mathbb{C} Britain - - 2518	
The prickly, or common, Butcher's Broom. <i>Ruscus</i> No. 1238. <i>Hall.</i> , &c. <i>Myricaantha</i> , Lob.	

	Page
<i>R. f. Brúscus</i> Ger.	
<i>R. myrtifólius aculéatus</i> Tourn., &c.	
<i>Rúscus</i> Dod.	
<i>Box Holly, Knce Holly, Wild Myrtle, Prickly Pettigree.</i>	
<i>Houz Frelon, Petit Houz, Buis piquant, Fragon épineux, Fragon piquant, Fr. Stechender Mäusedorn, Ger.</i>	
<i>Varieties</i> - - -	2518
2 rotundifólius Barr. -	
<i>R. vulgáris fólio-ampliore</i> Dill.	
3 láxus Sm.	
<i>R. láxus</i> Lodd, Cat.	
? <i>R. flexuosus</i> Mill. No. 6.	
2. hypophýllum <i>L.</i> - Italy and Africa	fig. 2387. 2519
The under-leaf Ruscus, or broad-leaved Butcher's Broom.	
<i>R. latifólius, &c.,</i> Tourn.	
<i>Laurus alexandrina</i> Lob., Bauh.	
<i>L. a. Chamedóphne</i> Col.	
<i>L. a. genuína</i> Tourn.	
<i>L. a. vera</i> Clus., Ray.	
<i>L. Chamedóphne vera</i> Dioscóridis Park.	
<i>L. a. altera</i> Matth.	
<i>Fragon sans Folióle, Fr. Breitblättriger Mäusedorn, Ger.</i>	
<i>Variety</i> - - -	2519
2 trifoliátum - Zante.	
<i>R. trifoliátus</i> Mill.	
3. (h.) Hypoglóssum <i>L.</i> - Italy, &c.	2519
The Under-tongue Ruscus, or double-leaved Butcher's Broom.	
<i>R. angustifólius, &c.,</i> Tourn.	
<i>Hypoglóssum</i> Lob.	
<i>Laurus alexandrina, fructu pediculo insidente,</i> Bauh., Ray.	
<i>Uendária</i> Brunf.	
<i>Fragon à Folióle, Fr. Zungen Mäusedorn, Ger.</i>	
4. racemósus <i>L.</i> - Portugal	fig. 2388, 2389. 2520
The branchy Ruscus, or Alexandrian Laurel.	
<i>R. angustifólius, fructu sámmis ramulis, &c.,</i> Tourn.	
<i>Fragon à Grappes, Fr. Trauben Mäusedorn, Ger.</i>	
<i>R. andrógnynus L.</i> - Canary Isles	- 2520
III. YU'CCA <i>L.</i> - 2516. 2521	
THE YUCCA, or Adam's Needle.	
1. gloriósa <i>L.</i> - Virginia, &c. f. 2390. 2521	
The glorious Yucca, or Adam's Needle.	
<i>Y. fólius A'loes</i> Bauh.	
<i>Y. canadéna</i> Ald. Hort.	
<i>Y. índica, &c.,</i> Barr. Rar.	
<i>Yucca peruána</i> Ger., &c.	
<i>Y. nóva gloriósa, &c.,</i> Lob.	
<i>Cordylinc, &c.,</i> Ray.	
The superb Yucca.	
<i>Yucca nain</i> Bon. Jard.	
<i>Yucca à Feuilles entières, Fr. Prüchtiger Yukka, Ger.</i>	
<i>Variety</i> - - -	2521
2 fólius variegátis Lodd. Cat.	
2. (g.) supérba <i>Haw.</i> - fig. 2391. 2523	
The superb Yucca, or Adam's Needle.	
<i>Y. gloriósa</i> And. Bot. Rep.	
3. alóifólia <i>L.</i> - South America	fig. 2392, 2393. 2523.
The Aloe-leaved Yucca, or Adam's Needle.	
<i>Y. arboréscens, &c.,</i> Dill.	
<i>A'loe Yucca fólius, cauléscens,</i> Pluk.	
<i>A. americána Yucca fólius, arboréscens,</i> Com.	
<i>Y. cauléscens, fólius lineári-lanceolátis, &c.,</i> Michx.	
<i>Variety</i> - - -	2523
2 péndula <i>Cat. Hort. Par.</i> -	
The pendent-leaved Aloe-leaved <i>Y.</i>	
4. dracónis <i>L.</i> - S. Carolina f. 2394. 2525	
The Dragon Yucca, or drooping-leaved Adam's Needle.	
<i>Y. dracónis fólio serrátò refléto</i> Dill.	
<i>Dracóni árbori, &c.,</i> Bauh.	
<i>Tacóri fólio Dracóni árboris stímili</i> Bauh.	
5. stricta <i>Sims</i> - Carolina f. 2395. 2525	
The upright Yucca, or Lyon's narrow-leaved Adam's Needle.	
6. recurvifólia <i>Salisb.</i> - Georgia - 2526	
The recurved-leaved Yucca.	
<i>Y. recurva</i> Hort.	
7. filamentósa <i>L.</i> - Virginia f. 2396. 2526	
The filamentose Yucca, or thracady Adam's N.	
<i>Y. fólius filamentósis</i> Moris. Hist.	
<i>Y. virginiana, &c.,</i> Pluk.	
8. (f.) angustifólia <i>Ph.</i> - North America	fig. 2397. 2526
The narrow-leaved Yucca.	
9. fláccida <i>Haw.</i> - Georgia f. 2398. 2527	
The flaccid-leaved Yucca.	
10. glaucéscens <i>Haw.</i> - North America	fig. 2399. 2527
The glaucescent Yucca.	
—	
<i>Half-hardy Monocotyledonous Plants, deserving a place in the British Arboretum.</i> - - - 2527	
Foucroya longæva <i>Karw.</i> - Mexico	fig. 2400, 2401. 2527
F. gigantæa <i>Vent.</i> - S. America	2527
Littæa gemmiflora <i>Brig.</i> - Peru	fig. 2402. 2522
<i>Agave gemmiflora</i> Ker.	
<i>Bonapartea júncea</i> Haw.	
Agave americána <i>L.</i> - S. America	2529
The American Aloe.	
Phormium tenax <i>L.</i> - N. Zealand	2529
The New Zealand Flax.	
Chamærops humilis <i>L.</i> - S. Europe	fig. 2403. 2530
The dwarf Fan Palm, or Palmetto.	
<i>Phoenix humilis</i> Cav.	
<i>Pálma humilis</i> Bauh.	
<i>Palmist: E'cetail, Fr. Zwergpalme, Ger.</i>	
C. serrulata <i>Willd.</i> - Georgia	2532
C. hystrix <i>Ph.</i> - Georgia	- 2532
C. Palmétto <i>Willd.</i> - Carolina	- 2532
<i>Corypha Palmétto</i> Walt.	
The Cabbage Tree, Amer.	
Bambusa nigra <i>Lodd. Cat.</i> - India	2532
B. ? arundinæcea - India	- 2532
Arundo Donax <i>Willd.</i> - S. Europe and Mount Aëtna - - - 2532	

SUPPLEMENT,















CONSISTING OF ADDITIONS AND CORRECTIONS.

The Species or Varieties not included in the body of the work, but here added, have the sign of addition prefixed, thus +. Those in which the name has been altered have the mark || prefixed.

Part I. HISTORY AND GEOGRAPHY, &c.	-	-	-	-	-	Page
Part II. OF THE SCIENCE OF THE STUDY OF TREES	-	-	-	-	-	2534
Part III. THE ARBORETUM ET FRUTICETUM BRITANNICUM.	-	-	-	-	-	2534
						Page
<i>Clematidææ.</i>	-	2534		<i>Additional Species of</i>	<i>Berberis.</i>	2537
<i>Clematis triternata</i> Dec.	-	2534		<i>B. Coriaria</i> Wall.		
+ <i>C. nepalensis</i> Dec. Mouuts				<i>B. buxifolia</i>		
Choor, Urukta, &c.	-	2534		<i>Mahonia</i> Nutt.	-	2537
The Nepal Clematis.				<i>M. fascicularis</i> Dec.	-	2537
<i>C. montana</i> D. Don.				<i>M. repens</i> G. Don	-	2537
<i>C. montana</i> var. Ham. MS.					§ Nandina.	2537
+ <i>C. Hendersoni</i> Hort. Hybrid		2535		+ <i>Nandina domestica</i> Humb.		
<i>C. florida</i>	-	2535		China	fig. 2410.	2537
+ <i>C. f.</i> 3 Sieboldti D. Don.						
Japan		2535				
<i>C. Sieboldii</i> Parl.				<i>Cruciæææ.</i>	-	2538
<i>C. bicolor</i> Hort.				<i>Cheiranthus Cheiri</i> fruticulosus	-	2538
+ *13. <i>C. cærulea</i> Lindl. Japan		2535		<i>Iberis sempervirens</i>	-	2538
fig. 2407.						
The blue, or violet, flowered Clematis.				<i>Cistææææ Lindl.</i>	-	2538
<i>C. asiæca grandiflora</i> Sieb.				<i>Cistus latifolius</i>	-	2538
<i>C. grandiflora</i> Hort.				<i>Helianthemum umbellatum</i>	-	2538
<i>C. Viticella</i> L.	-	2535				
<i>C. V.</i> 5 baccata	-	2535		<i>Polygalææææ.</i>	-	2538
<i>C. campaniflora</i> Brot.	-	2535		<i>Polygala Chamæbuxus</i> L.	-	2538
<i>C. balearica</i> Rich.	fig. 2408.	2535				
<i>C. montana</i>	-	2535		<i>Malvæææææ.</i>	-	2538
				<i>Hibiscus syriacus</i>	-	2538
<i>Pæoniæææææ.</i>	-	2535		+ <i>Malva Munroana</i> D. Don		
				Columbia	fig. 2411.	2538
<i>Magnoliæææææ.</i>	-	2536		Munro's Mallow.		
<i>Magnolia grandiflora</i>	-	2536				
<i>M. glauca</i>	-	2536		<i>Tiliæææææ.</i>	-	2538
<i>M. tripetala</i>	-	2536		<i>Tilia</i>	-	2538
<i>M. purpurea</i>	-	2536		<i>T. europæa</i>	-	2538
<i>Liriodendron Tulipifera</i>	-	2536		<i>Græwia occidentalis</i>	fig. 2412.	2540
<i>Anonæææææ.</i>	-	2536		<i>Ternströmiæææææ.</i>	2540	
<i>Asimina</i> Adans.	-	2536		<i>Gordonia</i>	-	2540
				<i>Camellia reticulata</i>	fig. 2413.	2540
<i>Berbericæææææ.</i>	-	2536				
<i>Berberis vulgaris</i>	fig. 2409.	2536		<i>Aurantidæææææ.</i>	-	2540
<i>B. cuneatrifolia</i> Lam.	-	2537				
<i>B. floribunda</i>	-	2537				
<i>B. Lycium angustifolium</i> Royle.						
<i>B. asiatica</i> Roxb.	-	2537				
<i>The Lycium of Dioscorides.</i> Royle.						
<i>B. dealbata</i> Lindl.	-	2537				

	Page
Hypericàcæ.	2541
<i>Hypéricum canariense</i> L. fig. 2414.	2541
<i>H. chinense</i> L.	2541
? <i>H. nepalense</i> Hort.	
<i>H. Kalmianum</i> Lam.	2541
<i>H. foliosum</i> Ait.	
<i>H. prolificum</i> L.	fig. 2415. 2541
<i>H. ægyptiacum</i> L.	fig. 2416. 2541
+ <i>H. nepalense</i> Hort. ✱	2541
Aceràcæ.	2541
<i>Acer oblongum</i> Wall. fig. 2417, 2418.	2541
<i>A. Pseudo-Plátanus</i>	2542
<i>A. campêtre</i>	2542
+ var. 5 lævigatum. †	
The smooth-leaved Field Maple.	
+ <i>A. c. 6 nanum</i> Hort. ✱	2542
The dwarf Field Maple.	
<i>Doubtful Species of Acer</i>	2543
<i>Anticipated Species of Acer.</i>	
<i>A. ibéricum</i>	2543
<i>A. lævigatum</i>	2543
<i>A. villòsum</i>	2543
+ <i>Negundo</i> f. 3 violaceum Booth †	2543
Æsculàcæ.	2543
<i>Æsculus</i>	2543
<i>Æ. Hippocàstanum</i>	2543
+ <i>Æ. H. 4 fòliis argenteis</i> †	2543
<i>Æ. (H.) ohioënsis</i>	2543
<i>Æ. (H.) rubicunda</i>	2543
<i>Æ. (H.) Lyòni</i>	2543
<i>Pàvia hýbrida</i>	2543
<i>P. macrostàcha</i>	2543
<i>Other Varieties of Pàvia</i>	2543
Vitàcæ.	2544
<i>Ampelòpsis hirsùta</i>	2544
Rutàcæ.	2544
<i>Rùta gravèolens</i>	2544
<i>Half-hardy Species of Rùta</i>	2554
App. 1. <i>Half-hardy Species of Rutàcæ.</i>	
+ <i>Corra'a álba</i> Sm. ✱ □ fig. 2419.	2544
<i>C. cotinifolia</i> Sal.	
+ <i>C. a. 2 rotundifolia</i> Dec. ✱ □	2554
<i>C. rufa</i> Vent.	
+ <i>Cròwea saligna</i> Sm. ✱ □ fig. 2420.	2544
+ <i>Borònia serrulàta</i> Sm. ✱ □	fig. 2421, 2422. 2544
Xanthoxylàcæ.	2544
<i>Ptèlea trifoliàta</i>	2544
Coriàcæ.	2545
<i>Coriària myrtifolia</i>	2545
<i>C. microphýlla</i>	2545

	Page
Celastràcæ.	2545
<i>Euònymus europæus</i>	2545
<i>E. sarmentòsus</i>	2545
<i>E. garciniaefolius</i>	2545
<i>E. grandiflorus</i>	2545
+ <i>E. caucàsicus</i> Lodd. Cat. ✱	fig. 2423. 2545
? <i>E. nana</i> Bieb.	
<i>E. japònicus</i>	fig. 2425. 2545
<i>Celastrus scandens</i>	2545
+ <i>C. nepalensis</i> ✱	2545
+ <i>C. pyracanthifolia</i> ✱	2545
<i>Nemopánthes canadensis</i>	fig. 2424. 2545
Aquifoliàcæ.	2545
<i>Ilex Aquifolium</i>	2545, 2546
+ <i>I. A. 24 fructu nigro</i> Hort. †	2545
The black-fruited common Holly.	
<i>I. baleàrica</i>	fig. 2426. 2546
+ <i>I. magellanica</i> Lodd. Cat. †	2546
<i>Prinos deciduus</i>	2546
<i>P. verticillatus</i>	2546
<i>P. glàber</i>	fig. 2428. 2546
<i>P. coriàceus</i>	fig. 2427. 2546
Rhamnàcæ.	2546
<i>Zizyphus sinënsis</i>	2546
<i>Z. Jujuba</i>	fig. 2429. 3546
+ <i>Paliurus virgatus</i> D. Don. ✱	fig. 2430. 2547
The twiggy Christ's Thorn.	
<i>Rhàmnus</i>	2547
<i>R. hýbridus</i>	2547
<i>R. Erythròxylon</i>	2547
<i>R. E. angustissimum</i>	2647
<i>R. persicùolius</i> Bert.	2547
+ <i>R. glandulosus</i> Hort. ✱	2547
<i>Ceanòthus azureus</i>	2547
<i>C. intermèdius</i>	2547
+ <i>C. collinus</i> Dougl. ✱	fig. 2431. 2547
The Hill-side Ceanothus.	
<i>Collètia hòrrida</i> Willd.	fig. 2432c. 2548
+ <i>C. alicina</i> Gil. ✱	fig. 2432a. 2548
<i>Retanilla E'phedra</i> Bron.	fig. 2432b. 2548
<i>Pomadèrris</i>	2548
Anacardiàcæ.	2548
<i>Pistàcia</i>	2548
<i>Rhús</i>	2548
<i>R. snavèolens</i>	2548
<i>Rhús aromàtica</i>	2548
<i>R. snavèolens</i> Ait.	
<i>Myrica trifoliàta</i> Hort.	
<i>Toricodendron crenatum</i> Mill.	
+ <i>R. caustica</i> Hook †	2548
<i>Laúrus caustica</i> Mol.	
<i>Duvaña ovàta</i>	fig. 2433. 2549

	Page		Page
<i>Leguminosæ.</i>	2549	+ <i>K. ovata Sims</i>  fig. 2444.	2554
Sect. I. <i>SORNO`REÆ.</i>	2549	<i>K. cordata Lindl.</i>	
<i>Sophora</i>	2549	+ <i>K. inophylla, K. Comptoniana, K. prostrata,</i>	
+ <i>Baptisia tinctoria R. Br.</i> 	fig. 2434. 2549	<i>K. bracteata, and K. sericea</i>	2554
+ <i>Anagyris foetida L.</i> 	fig. 2435. 2549	Sect. V. <i>CASSIÆ.</i>	2554
+ <i>A. f. 2 glauca Dec.</i> 	2549	<i>Half-hardy Species.</i>	
+ <i>A. latifolia Willd.</i> 	2549	<i>Poinciana pulcherrima</i>	2554
<i>Brachysemum latifolium</i>	2549	<i>Other Half-hardy Leguminosæ.</i>	
Sect. II. <i>LO`TEÆ.</i>	2549	<i>Acacia Cavènia</i>	2554
<i>Ulex europæa</i>	2549	<i>A. Julibrissin</i>	2554
<i>Spartium junceum, + 3 odoratissimum D. Dou.</i> 	fig. 2436. 2550	<i>Rosaceæ.</i>	2554
+ <i>S. acutifolium Lindl.</i> 	2550	Sect. I. <i>AMYGDALÆÆ.</i>	2554
<i>Genista parviflora</i>	2550	<i>Amygdalus nana</i>	2554
<i>G. umbellata</i>	2550	<i>A. n. 3 campestris</i>	2554
<i>G. monosperma</i>	2550	<i>A. n. 4 incana</i>	2554
<i>G. æthiœnsis</i>	2550	<i>A. n. 5 sibirica</i>	2554
<i>G. anxantica</i>	2550	<i>Persica</i>	2554
<i>G. scaridsa Viv.</i>		<i>P. vulgaris 6 compressa</i>	2554
<i>G. januensis Viv.</i>		<i>Armeniaca</i>	2554
<i>G. genuensis Pers.</i>		<i>A. pedunculata Hort.</i>	2554
<i>G. sagittalis</i>	2550	<i>Prunus</i>	2554
<i>G. procumbens</i>	2550	<i>P. domestica myrobâlana</i>	2555
<i>G. canariensis</i> fig. 2437.	2551	<i>P. cœndicans</i>	2555
+ <i>G. splendens Webb et Berth.</i> 	2551	<i>Cerasus</i>	2555
+ <i>G. stenopétala Webb et Berth.</i> 	2551	<i>C. sylvestris</i>	2555
<i>G. microphylla</i>	2551	<i>C. chîcasa</i>	2555
<i>Cytisus Laburnum</i>	2551	<i>C. pubescens</i>	2555
<i>C. scoparius</i>	2551	<i>C. Pâdus 4 bracteosa</i>	2555
<i>C. Weldenii</i>	2551	<i>C. mollis</i>	2555
<i>C. proliferus</i>	2551	<i>C. lusitânica</i>	2555
+ <i>C. æoliœus Guss.</i> 	fig. 2438. 2551	<i>C. l. 2 Hiau</i>	fig. 2445. 2555
+ <i>C. elegans Ait.</i> 	2551	<i>Pâdus racemosa</i>	2555
+ <i>C. racemosus Marnock.</i> 	2552	Sect. II. <i>SPIRÆÆÆ.</i>	2555
<i>Adenocarpus</i>	2552	<i>Kerria</i> fig. 2446.	2555
<i>Amorpha glabra</i>	2552	<i>Spiræa chamædrifolia media f.</i>	2449. 2556
<i>Robinia</i>	2552	<i>S. adnifolia phyllantha</i> fig. 2448.	2556
<i>R. dubia</i>	2552	<i>S. crataegifolia</i>	fig. 2450. 2556
<i>R. hispida</i>	2552	<i>S. betulæfolia</i>	fig. 2447. 2556
<i>Caragana microphylla</i>	2552	<i>S. alpina</i>	2556
<i>Colutea arborescens</i> fig. 2440.	2552	<i>S. a. latifolia</i>	2556
<i>Half-hardy Lôtææ.</i>		<i>S. thalictroides</i>	2556
<i>Bossia rufo</i>	fig. 2441. 2552	<i>S. hypericifolia var. flava</i>	2556
<i>Anthyllis erinæcea</i>	2552	<i>S. ceanothifolia</i> fig. 2452.	2556
Sect. III. <i>HEDYSÆ`REÆ.</i>	2552	<i>S. hypericifolia Besseriâna</i> fig. 2451.	2556
<i>Half-hardy Hedysæreæ.</i>		<i>S. salicifolia alpestris</i> fig. 2453.	2556
<i>Desmodium polycarpum Dec.</i>	fig. 2442. 2552	+ <i>S. nutans</i>	2556
<i>Hedysarum polycarpum Poir.</i>		+ <i>S. taûrica</i>	2557
+ <i>D. nutans Wall.</i> 	fig. 2443. 2553	Sect. III. <i>POTENTILLÆÆ.</i>	2557
Sect. IV. <i>PHASOLÆÆ.</i>	2553	<i>Rubus</i>	2557
<i>Wistaria chinensis</i>	2553	<i>R. idæns</i>	fig. 2454. 2557
<i>Half-hardy Phasolææ.</i>		<i>R. fruticosus</i>	2557
+ <i>Kennedya rubicunda Vent.</i> 	2553	<i>Potentilla fruticosa</i>	2557
<i>Glycine rubicunda Curt.</i>		<i>P. f. 2 dahurica</i>	2557
<i>Cærtua rubicunda Moench.</i>			

	Page		Page
P. f. 3 tenuiloba - - -	2557	XIII.* STRANVÆSLA Lindl.	2563
P. floribunda - - -	2557	The Stranvæsia.	
+ X.* COWANIA D. Don.	2557	1. S. glaucescens Lindl. †	2563
The Cowania.		fig. 562, 563. in p. 845.	
+ 1. C. plicata D. Don $\frac{1}{2}$ fig. 2455.	2557	Cratægus glauca Wall.	
Sect. IV. ROSEÆ.	2558	Cotonæster (v.) laxiflora -	2563
Ròsa - - -	2558	+ C. (v.) 1. 2 uniflora Fisch. $\frac{1}{2}$	2564
Ròsa microphýlla - - -	2558	C. nummulària - - -	2564
+ R. m. álba $\frac{1}{2}$ - - -	2558	C. elliptica Hort.	
R. alpina - - -	2558	Eriobótريا elliptica Lindl.	
+ R. a. 15 speciòsa $\frac{1}{2}$ - - -	2558	Mespilus Cule Hort.	
R. Sabini - - -	2559	+ C. lævis Lodd. $\frac{1}{2}$ - - -	2564
+ R. S. grácilis $\frac{1}{2}$ - - -	2559	Pýrus commúnis - - -	2564
R. Doniana - - -	2559	P. variolòsa - - -	2564
+ R. D. hórrida $\frac{1}{2}$ - - -	2559	P. Michaúxii - - -	2564
R. Wilsoni Borr. - - -	2559	P. (Málus) prunifolia - - -	2564
R. damascèna - - -	2559	P. (M.) dioica - - -	2564
+ R. d. subálba $\frac{1}{2}$ fig. 2456.	2559	P. (c.) angustifolia - - -	2565
R. centifolia - - -	2559	+ P. Schótti Led. $\frac{1}{2}$ - - -	2565
+ R. c. muscòsa cristàta Hook. $\frac{1}{2}$	2559	+ P. stipulàcea Hort. $\frac{1}{2}$ - - -	2565
R. pomponia - - -	2559	P. Aria angustifolia - - -	2566
R. gállica - - -	fig. 2457. 2559	P. A. longifolia Hort. - - -	2566
R. álba - - -	fig. 2458. 2559	P. A. edulis Hort. - - -	2566
R. lítea - - -	2559	Cratægus græca Hort.	
+ R. l. 4 flòre plèno $\frac{1}{2}$ - - -	2559	P. rivularis - - -	2566
+ R. l. 5 Hóggü $\frac{1}{2}$ - - -	2560	P. americana - - -	2566
R. rubiginòsa - - -	2560	P. canadensis Hort.	
+ R. Lyónii $\frac{1}{2}$ - - -	2560	P. Sórbus - - -	2566
R. canina - - -	fig. 2459. 2560	P. spúria - - -	2566
R. indica - - -	2560	P. (arbutifolia) melanocarpa - - -	2566
+ R. i. 12 flavescens Hort. $\frac{1}{2}$ - - -	2560	Mespilus capitata Lodd.	
+ R. i. 13 Blaíri D. Don. $\frac{1}{2}$ - - -	2560	M. floribunda Lodd.	
R. símica - - -	fig. 2460. 2560	M. pübens Lodd.	
R. hýstrix Lindl.		P. (a.) m. 2 subpubescens Lindl. - - -	2566
R. levigata Michx.		P. m. xanthocarpa Hort.	
R. macrophýlla - - -	2560	P. grandifolia - - -	fig. 2465. 2566
Cratægus coccinea - - -	2562	P. Chamæmèspilus - - -	2566
+ C. e. 5 neapolitana Hort. $\frac{1}{2}$ - - -	2562	Eriobótريا - - -	2566
Mespilus constantinopolitana Godefroy.		E. elliptica - - -	2566
Cratægus punctata - - -	2562	Kagenéckia cratægoides fig. 2466.	2566
C. p. 4 brevispina Doug. f. 2462.	2562	Calycánthus levigatus - - -	2566
C. ovalifolia - - -	2562	Chimonánthus frágans - - -	2566
C. Douglásii - - -	2562	+ C. f. 4 parviflorus Hort. $\frac{1}{2}$ - - -	2566
C. trilobata - - -	2562		
C. Arònia - - -	2562	Granatæcæ.	2566
C. heterophýlla - - -	2562	Punica Granatum - - -	2566
C. Oxyacantha - - -	2562		
C. O. 9 purpúrea - - -	2562	Onagræcæ.	2566
C. prunifolia - - -	2562	Fúchsia microphýlla - - -	2566
C. O. 21 strícta - - -	2562	F. excorticata - - -	2567
C. mexicæna - - -	2562	+ F. fulgens Dec. $\frac{1}{2}$ - - -	2567
C. Lambertiana Hort.			
§ xvi. Glúca.	2625	Philadelphæcæ.	2567
Synopsis of the Species of Cratægus -	2562	+ Philadelphus Gordonianus Lindl. $\frac{1}{2}$	2567
Additional Species of Cratægus.		+ 6.* P. speciòsus Schrad. $\frac{1}{2}$ - - -	2567
+ C. florentina Zucch. $\frac{1}{2}$ - - -	2563	The showy-flowered Philadelphus, or Mock Orange.	
Mespilus florentina Bert.		P. grandiflorus - - -	2567
+ C. opaca Hook. $\frac{1}{2}$ - - -	2563	Deutzia scabra - - -	2567
		D. Bruònia - - -	2567
		+ D. grandiflora $\frac{1}{2}$ - - -	2567

	Page		Page
<i>Myrtàcææ.</i>	- 2567	<i>Cornàcææ.</i>	- 2571
Eucalýptus robústa fig. 2467.	2567	Córnus - - -	2571
E. amygdáлина - - -	2567		
+ E. alpína Hort. \blacksquare -	2567	<i>Loranthàcææ.</i>	- 2571
Leptospermum lanigerum	2567	Viscum - - -	2571
L. baccátum - - -	2567	Aícuba - - -	fig. 2479. 2571
L. flexuósum - - -	2567	Loranthus europæus	- 2571
L. grandiflórum - - -	2567		
L. stellátum - - -	2568	<i>Caprifoliàcææ.</i>	- 2572
+ L. scopàrium Forst. \mathfrak{z} fig. 2468.	2568	Sambucus - - -	2572
<i>The Broom Tree, or Dog-wood Tree,</i>		Viburnum - - -	2572
<i>of Van Diemen's Land.</i>		V. levigátum - - -	2572
<i>New Zealand Tea Plant.</i>		V. carolinianum Hort.	
Myrtus communis fig. 2469, 2470.	2568	V. Lantàna - - -	2572
<i>Crassulàcææ.</i>	- 2568	V. L. 2 grandifolia - - -	2572
Sedum populifolium fig. 2471.	2568	V. cotinifolium - - -	2572
		V. Mullaha Ham.	
<i>Reaumuriàcææ.</i>	2569	V. O'pulus - - -	2572
+ Reaumuria hypericoides Willd.		V. O. 4 nana Hort. \mathfrak{z} - - -	2572
\mathfrak{z} \square fig. 2472, 2473.	2569	Lonicera - - -	2572
<i>Hypericum alternifolium Lab.</i>		L. parviflora fig. 2480, 2481.	2572
+ R. vermiculata Linn. \mathfrak{z} \square -	2569	L. sempervirens - - -	2572
<i>Cactiàcææ.</i>	- 2569	L. s. 2 major - - -	2572
Opuntia vulgaris fig. 2474.	2569	+ L. hispídula \mathfrak{z} fig. 2483.	2572
		<i>Caprifolium hispidulum Dougl.</i>	
<i>Grossulàcææ.</i>	- 2569	L. longiflora fig. 2482.	2572
Ribes - - -	2569	+ L. montana Hort. \mathfrak{z} - - -	2572
R. lacustre - - -	2569	Symphoricarpus racemósus	- 2572
R. multiflórum - - -	2569		
R. vivifolium.		<i>Rubiàcææ.</i>	- 2573
R. prostrátum - - -	2569	Cephalánthus - - -	2573
R. resinósum - - -	2569	+ Luculia gratíssima D. Don \mathfrak{z} \square	
R. rigens - - -	2570	fig. 2484.	2573
R. punctátum - - -	2570	<i>Cinchona gratissima Wall.</i>	
R. flóridum - - -	2570	<i>Mussaenda Luculia Ham.</i>	
R. (n.) f. 2 grandiflórum Hort.	2570	<i>Luculi Swa. Nepalese.</i>	
R. rigens Michx.		<i>Compósiteæ.</i>	- 2573
R. inebrians - - -	2570	Artemisia - - -	2573
R. cæcum - - -	2570	Onocitium - - -	2573
R. viscosissimum fig. 738.	2570	Pyræthrum - - -	2573
R. flavum fig. 2476.	2570	+ Anthemis pórrigens Hort. \mathfrak{z}	
Mr. Gordon's List of the different		fig. 2486.	2573
species of Ribes.	2570	Eriocéphalus africanus	fig. 2487. 2573
<i>Escallonniàcææ.</i>	- 2570		
+ Escallonia glandulosa \mathfrak{z} \square -	2570	<i>Epacridiàcææ.</i>	- 2573
+ E. illinita \mathfrak{z} \square fig. 2477.	2570	Stenanthèra pinifolia	fig. 2488. 2573
<i>Saurifràgææ.</i>	- 2570		
Hydrangea - - -	2570	<i>Ericàcææ.</i>	- 2574
<i>Umbellàcææ.</i>	- 2570	Erica - - -	2574
Bupleurium fruticosum fig. 2478.	2570	E. Tétralix - - -	2574
<i>Hederàcææ.</i>	- 2570	E. T. 4 Mackaiana - - -	2574
Hedera Helix - - -	2570	E. Mackaui - - -	2574
<i>Hammulidàcææ.</i>	- 2570	E. arborea - - -	2574
Hamanulis virginica - - -	2570	E. ciliaris - - -	2574
		Calluna vulgaris - - -	2574
		Directions for the Culture of Cape	
		Heaths - - -	2574
		Andrómèda - - -	2574

	Page
Cassiope - - -	2574
Lyonia - - -	2574
L. marginata - - - fig. 2489.	2574
L. mariana - - - fig. 2490.	2574
L. racemosa - - - fig. 2491.	2574
Arbutus Unedo - - -	2575
A. hybrida - - -	2575
A. Andrichne - - - fig. 2492.	2575
+ A. speciosa ♀ ♂ - - -	2575
+ A. nepalensis ♀ ♂ - - -	2575
Arctostaphylos Uva-ursi - - -	2575
+ A. U. 2 austriaca Lodd. ♀ - - -	2575
Pernettia mucronata - - -	2575
Gaultheria Shallon - - -	2575
Epigaea repens - - -	2575
+ E. r. rubicunda D. Don ♀ - - -	2575
Rhododendron - - -	2576
R. ponticum azaleoides - - -	2576
+ R. p. fragrans ♀ - - -	2576
R. maximum hybridum - - - fig. 2494.	2576
R. Purshii - - -	2576
R. punctatum - - - fig. 2495.	2576
R. p. 2 majus - - -	2576
R. nudiflorum rutilans - - - fig. 2496.	2576
R. viscosum Cartonia - - - fig. 2497.	2576
R. Rhodora - - - fig. 2498.	2576
Kalmia - - -	2576
Menziesia globularis - - - fig. 2499.	2576
Azalea procumbens - - - fig. 2530.	2576
Ledum - - -	2576
Vaccinium - - -	2576
V. grandiflorum - - - fig. 2501.	2577
V. padiflorum - - - fig. 2502.	2577
Oxycoccus - - -	2577

Half-hardy Species of Ericaceæ.

+ Cyrilla racemosa N. Du Ham. ♀ ♂ - - -	fig. 2503. 2577
-----------------------------------------	-----------------

- C. racemiflora L.
- C. caroliniana Michx.
- Andromeda plumata Marsh.
- Vicia caroliniana L'Herit.
- I. Cyrilla Swt.
- I. racemiflora Lam.

+ Myrsinaceæ. - 2578

+ Myrsine africana L. ♀ ♂ fig. 2504.	2578
M. glabra Gært.	
Vitis idæa æthiopica Com. Hort.	
Bauxis africana Pluk.	
+ Manglilla Milleriana Pers. ♀ ♂ - - -	fig. 2505. 2578
Sideroxylon nitè L.	
Myrsine nitis Spr.	

Sapotaceæ. - 2578

Bumelia lycioides - - - fig. 2506.	2578
B. salicifolia - - - fig. 2507.	2578

Ebenaceæ. - 2578

Diospyros - - -	2578
D. intermedia, D. dignia, and D. stricta - - -	2578

Oleaceæ. - 2578

Ligustrum - - -	2578
L. spicatum - - -	2578
+ L. s. 2 glabrum Hook. ♀ - - -	2578
L. lucidum - - -	2579
L. l. 2 floribundum - - - fig. 2508.	2579
L. japonicum - - -	2579
Phillyrea - - -	2579
Olea - - -	2579
O. capensis - - - fig. 2509.	2579
+ Notelæva ligustrina Vent. ♀ - - -	fig. 2510. 2579
Syringa - - -	2579
S. Josikæa - - -	2579
S. Emodi - - -	2579
Fraxinus - - -	2579
F. excelsior - - -	2580
F. e. aurea - - -	2580
F. purpurascens - - -	2580
F. purpurea - - -	2580
F. (e.) angustifolia - - - fig. 2511.	2580
F. lentiscifolia - - - fig. 2512.	2580
F. epiptera - - -	2580

Jasminaceæ. - 2581

Jasminum - - -	2581
----------------	------

Apocynaceæ. - 2581

Vinea - - -	2581
+ V. acutifolia Bert. ♀ - - -	2581

Asclepiadaceæ. - 2581

Half-hardy Species of Periploca.

+ Physianthus albicans Hort. ♀ ♂ - - -	2581
P. undulatus Hort.	

Bignoniaceæ. - 2581

Bignonia - - -	2581
Tecoma radicans - - -	2581
T. r. major - - -	2581

Solanaceæ. - 2581

Grabowska - - -	2581
Solanum - - -	2581
S. bonariense - - -	2582
S. Balbisii - - - fig. 2514.	2582

- || S. decurrens Balb.
- || S. bruceifolium Jacq.
- || S. mauritanicum Willd.
- || S. viscosum Dec.

S. littorale Hort. - - -	2582
Lycium lanecolatum f. 2513. 2516.	2582
L. turbinatum - - - fig. 2517.	2582

Scrophulariaceæ. - 2582

Half-hardy Species.

+ Pentstemon Scouleri Doug. ♀ - - -	fig. 2518. 2582
+ P. atropurpurea G. Don. ♀ - - -	2582

	Page		Page
<i>Iubiùceæ</i> -	2582	<i>Ulmæcæ.</i> -	2586
<i>Phymus grandiflorus</i> fig. 2515.	2582	<i>Ulmus</i> -	2586
<i>Prostanthera lasianthos</i> fig. 2519.	2583	<i>U. c. nana</i> * -	2587
<i>Verbenæcæ.</i> -	2583	<i>U. americana</i> -	2587
<i>Fitex incisa</i> -	2583	+ <i>U. a. foliis variegatis</i> †	2587
<i>Clerodendron speciosissimum</i> -	2583	<i>Juglandæcæ.</i> -	2587
[†] <i>C. squamatum</i> Vahl.		<i>Juglans</i> -	2587
<i>Aloysia citriodora</i> -	2583	<i>Carya álba</i> -	2587
<i>Plumbaginæcæ.</i> -	2583	<i>Pterocarya caucásica</i> -	2587
<i>Plumbago capensis</i> fig. 2520.	2583	<i>Salicæcæ.</i> -	2587
<i>Chenopodiæcæ.</i> -	2583	<i>Sáliz</i> -	2587
<i>Atriplex Halimms</i> -	2583	<i>S. babilónica</i> -	2588
<i>Kochia prostrata</i> -	2583	<i>S. b. Napoleóna</i> -	2588
<i>Polygonæcæ.</i> -	2583	<i>S. nigra</i> fig. 2527 a.	2588
+ <i>Tragopyrum maritimum</i> Dougl. †	2583	<i>S. ligustrina</i> fig. 2527 b.	2588
<i>Lauræcæ.</i> -	2583	<i>S. versicolor</i> fig. 2528.	2588
<i>Laurus nobilis</i> -	2583	+ <i>S. coluteoides</i> Mirb. * fig. 2529.	2588
<i>L. Sássafras</i> -	2583	+ <i>S. sitchénsis</i> Hort. * -	2588
<i>Proteæcæ.</i> -	2584	<i>Pópulus</i> -	2588
+ <i>Banksia latifolia</i> R. Br. * †	2584	<i>P. nigra</i> and <i>P. fastigiata</i> -	2588
fig. 2522.	2584	<i>P. álba</i> -	2589
<i>Thymelæcæ</i> -	2584	<i>P. tremula</i> -	2589
<i>Daphne Mezereum</i> -	2584	<i>P. fastigiata</i> -	2589
<i>Eleagnæcæ.</i> -	2584	<i>P. balsamifera</i> fig. 2530.	2589
<i>Eleagnus horténsis orientális</i> -	2584	<i>Betulæcæ.</i> -	2589
<i>E. conferta</i> -	2584	<i>Alnus</i> -	2589
<i>Hippophæ Rhamnoides</i> -	2584	+ <i>A. acuminata</i> H. et B. † † f. 2531.	2589
<i>Aristolochiæcæ.</i> -	2584	+ <i>A. castaneafolia</i> Mirb. † † f. 2532.	2590
<i>Euphorbiæcæ.</i> -	2584	<i>Bétula álba</i> -	2590
<i>Euphorbia spinosa</i> -	2585	<i>B. fruticosa</i> -	2590
+ <i>E. Myrsinites</i> † -	2585	<i>B. papyracea</i> -	2590
+ <i>E. rigida</i> † -	2585	<i>Corylæcæ.</i> -	2590
<i>Búxus sempervirens</i> -	2585	<i>Quercus</i> -	2590
<i>Half-hardy Species.</i>		<i>Q. sessiliflora</i> -	2590
<i>Plagiánthus divaricatus</i> fig. 2524.	2585	<i>Q. Cérris</i> -	2591
+ <i>Cróton rosmarinifolia</i> Cunn. * †	2585	+ <i>Q. C. laciniata</i> † fig. 2534.	2591
fig. 2523.	2585	<i>Q. heterophýlla</i> -	2591
+ <i>Adèlia Acidòton</i> * † fig. 2525.	2585	<i>Q. Phéllós</i> -	2591
<i>Urticæcæ.</i> -	2586	<i>Q. Ballota</i> -	2591
<i>Morus nigra</i> fig. 2526.	2586	<i>Q. granuntia</i> -	2591
<i>M. álba</i> -	2586	<i>Q. hispánica</i> -	2591
<i>M. a. Morettiána</i> -	2586	<i>Q. virens</i> -	2591
<i>M. a. macrophýlla</i> -	2586	[†] <i>Q. hemispha'rica</i> Bart. Bot. Gard.	
<i>M. hispánica</i> -	2586	<i>Q. serrata</i> -	2591
<i>Broussonètia papyrifera</i> -	2586	<i>Q. glabérrima</i> -	2592
+ <i>B. p. 3 fructu álbo</i> * -	2586	Mexican Oaks -	2592
<i>Bòrya acuminata</i> -	2586	<i>Q. xalapénsis</i> -	2592
<i>B. ligustrina</i> -	2586	<i>Fágu</i> -	2593
<i>Ficus</i> -	2586	<i>Fágu cochinchinénsis</i> Lour. -	2593
		<i>F. antártica</i> Forst. -	2593
		<i>F. betuloides</i> -	2593
		<i>F. sylvática</i> fig. 2536.	2593
		<i>Castánea</i> -	2595
		<i>Cárpinus</i> -	2595
		<i>O'strya virginica</i> fig. 2537.	2595

	Page		Page
<i>Corylus Avellana</i>	2595	<i>P. Pinaster</i>	2598
<i>C. A. purpurea</i>	2595	<i>P. P. Lemoniana</i>	2599
<i>C. rostrata</i>	2526	<i>P. halepensis</i>	2599
<i>Platanaceæ.</i>	2597	<i>P. h. maritima</i>	2599
<i>Platanus</i>	2597	<i>P. brütia</i>	2599
<i>Balsamaceæ.</i>	2597	<i>P. variabilis</i>	fig. 2540. 2599
<i>Liquidambar</i>	2597	<i>P. Llaveana</i>	2599
<i>Myricaceæ.</i>	2597	<i>Abies</i>	2599
<i>Myrica Gale</i>	2597	<i>A. excelsa</i>	2599
<i>Gnetaceæ.</i>	2597	<i>Picea vulgaris</i> Link.	
<i>Ephedra americana</i>	fig. 2539. 2597	<i>A. e. viminalis</i>	2599
<i>Taxaceæ.</i>	2597	<i>P. viminalis</i> Alström.	
<i>Taxus baccata</i>	2597	<i>Hangetanne</i> , Swedish.	
<i>Phyllocladus trichomanoides</i>	2597	<i>A. e. stricta</i>	2600
<i>Coniferae, § Abietinae.</i>	2591	<i>A. Smithiana</i>	2601
<i>Pinus</i>	2597	<i>A. cephalonica</i>	2601
<i>P. sylvestris</i>	2597	<i>Picea</i>	2601
<i>P. pumilio</i>	2598	<i>P. pectinata</i>	fig. 2542. 2601
<i>P. humilis</i> Link.		<i>P. religiosa</i>	fig. 2543. 2602
<i>P. p. Mughus</i>	2598	<i>Larix europæa</i>	2602
<i>P. pumilio</i> Link.		<i>L. americana</i>	2603
<i>P. uncinata</i>	2859	<i>Cedrus Libani</i>	2603
<i>P. rotundata</i> Link.		<i>C. Deodara</i>	2603
<i>Pinus Laricio</i>	2598	<i>Araucaria excelsa</i>	2603
<i>P. austriaca</i>	2598	<i>A. Cunninghamii</i>	fig. 2545. 2603
<i>P. nigra</i> Link.		<i>Dammara australis</i>	2603
<i>P. nigra</i>	2598	<i>Cupressus sempervirens</i>	2605
		<i>Juniperus</i>	2605
		<i>J. hemisphærica</i> Presl	2605
		<i>Agave americana</i>	fig. 2546. 2605

APPENDIXES.

APP. I. Form of Return Paper	2609
APP. II. List of Trees and Shrubs growing in Italy, with their systematic and popular Italian Names	2510
APP. III. Priced Catalogues of Trees and Shrubs, contributed by British and Continental Nurserymen	2617
I. Catalogue of American and other Tree and Shrub Seeds, imported for Sale by George Charwood	2618
II. Catalogue of Forest and Ornamental Trees, American Plants, and Flowering Shrubs, sold by Richard Forrest	2620
III. A List of Trees, Plants, &c., sold by Peter Lawson and Son, Edinburgh	2626
IV. Catalogue of Hardy Trees and Shrubs cultivated for Sale in the Nursery of the Brothers Baumann, at Bollwyller	2635
V. List of Trees and Shrubs taken from the Retail Catalogue of James Booth and Sons, Hamburg	2646

INDEXES.

Index to Genera, including the English Names and scientific Synonymes	2655
Index to Miscellaneous Subjects	2667
Index to Persons and Places	2672

LIST OF THE PORTRAITS OF TREES

FORMING VOLS. V. VI. VII. AND VIII.

IN THE ORDER IN WHICH THEY ARE PLACED IN THESE VOLUMES, WITH THE NAME OF THE PLACE WHERE EACH TREE GREW, AND OF THE ARTIST WHO TOOK THE PORTRAIT.

The number on the left hand merely shows the series; that on the right extremity of the column, in parenthesis, is the number printed on the plate; and the number preceding it refers to that page of the body of the work where the tree is treated of.

VOL. V.

MAGNOLIAÆ CEE.

- | | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>1. <i>Magnolia grandiflora</i> - p. 261. (1.)
Brentford Nursery. <i>Miss Eliza Ronaldis.</i></p> <p>2. <i>M. g. exoniensis</i> - p. 261. (3.)
The Chiswick Villa. <i>Miss M. L.</i></p> <p>3. <i>M. glauca</i> - - p. 267. (4.)
Mile End Nursery. <i>Mr. R. Varden.</i></p> <p>4. <i>M. gl. Thompsoniana</i> - p. 267. (5.)
Mile End Nursery. <i>R. Varden.</i></p> <p>5. <i>M. tripétala</i> - - p. 269. (6.)
Mile End Nursery. <i>R. Varden.</i></p> <p>6. <i>M. macrophýlla</i> - - p. 271. (7.)
The Chiswick Villa. <i>Miss M. L.</i></p> <p>7. <i>M. acuminata</i> - - p. 273. (10.)
Messrs. Loddiges, Hackney. <i>R. Varden.</i></p> <p>8. <i>M. acuminata</i>, full-grown tree - (9a.)
Duke of Northumberland, Syon. <i>G. R. Lewis.</i></p> <p>9. <i>M. cordata</i> - - p. 275. (11.)
Messrs. Loddiges, Hackney. <i>R. Varden.</i></p> <p>10. <i>M. auriculata</i> - - p. 276. (8.)
Messrs. Loddiges, Hackney. <i>R. Varden.</i></p> <p>11. <i>M. (a.) pyramidata</i> - p. 277. (9.)
Messrs. Loddiges, Hackney. <i>R. Varden.</i></p> <p>12. <i>M. conspicua</i> - - p. 278. (12.)
Mile End Nursery. <i>R. Varden.</i></p> <p>13. <i>Liriodéndron Tulipífera</i> p. 284. (13.)
Hort. Soc. Gard. <i>F. Rauch.</i></p> <p>14. <i>L. Tulipífera</i>, full-grown tree (12a.)
Syon House. <i>G. R. Lewis.</i></p> | <p>22. <i>T. americana</i> } p. 373. (24. or 25.)
(syn. <i>álba</i>) }
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>23. <i>T. (a.) laxiflora</i> - - p. 374. (23.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>24. <i>T. (a.) pubescens</i> p. 374. (22. or 27.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

ACERAÆ CEE.

- | | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>15. <i>Tilia europæa</i> - - p. 364. (15.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>16. <i>T. e.</i>, full-grown tree - (8a.)
Studley Park. <i>H. W. Jukes.</i></p> <p>17. <i>T. (e.) platyphýlla</i> - p. 365. (19.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>18. <i>T. (e.) p. minor</i> - - p. 365. (20.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>19. <i>T. (e.) laciniata</i> - p. 366. (21. or 22.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>20. <i>T. (e.) parvifolia aurea</i> p. 366. (25.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>21. <i>T. (e.) álba</i> (syn. <i>ar-</i>
<i>géntea</i>) - - } p. 372. (18.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> | <p>25. <i>Acer tataricum</i> - - p. 406. (26.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>26. <i>A. spicatum</i> - - p. 406. (30.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>27. <i>A. striatum</i> - - p. 407. (28.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>28. <i>A. macrophýllum</i> - - p. 408. (34.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>29. <i>A. platanoides</i> - - p. 408. (35.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>30. <i>A. p. laciniatum</i> - - p. 409. (36.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>31. <i>A. saccharinum</i> - - p. 411. (37.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>32. <i>A. Pseudo-Platanus</i> - p. 414. (33.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>33. <i>A. Pseudo-Platanus</i>, full-
grown tree - } (33a.)
Studley Park. <i>H. W. Jukes.</i></p> <p>34. <i>A. obtusatum</i> (syn. <i>hý-</i>
<i>bridum</i>) - - } p. 420. (38.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>35. <i>A. opulifolium</i> - - p. 420. (31.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>36. <i>A. O'palus</i> - - p. 420. (31a.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>37. <i>A. eriocarpum</i> - - p. 423. (40.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>38. <i>A. e.</i>, full-grown tree - (39a.)
Bot. Gard., Kew. <i>L. Martin.</i></p> <p>39. <i>A. rubrum</i> - - p. 424. (39.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>40. <i>A. r.</i>, full-grown tree - (38a.)
Bot. Gard., Kew. <i>L. Martin.</i></p> <p>41. <i>A. monspessulanum</i> - p. 427. (27.)
Hort. Soc. Gard. <i>Miss M. L.</i></p> <p>42. <i>A. m.</i>, full-grown tree - (27a.)
Ham House, Essex. <i>H. W. Jukes.</i></p> |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

43. *A. campéstre* - p. 428. (32.)
Hort. Soc. Gard. *Miss M. L.*
44. *A. c.*, full-grown tree, 4to - (32a.)
Studley Park. *H. W. Jukes.*
45. *A. créticum*, full-grown tree } p. 430. (40a.)
Synon House, Middlesex. *G. R. Lewis.*
46. *Negúndo fraxinifólium* p. 460. (41.)
Hort. Soc. Gard. *Miss M. L.*
47. *N. f. crispum* - p. 460. (42.)
Hort. Soc. Gard. *Miss M. L.*

ÆSCULA'CEÆ.

48. *Æ'sculus Hippocástanum* p. 463. (43.)
Hort. Soc. Gard. *Miss M. L.*
49. *Æ. H.*, full-grown tree - (44.)
Forty Hill, Enfield. *W. J. Nesfield.*
50. *Æ. (H.) rubicúnda* } p. 467. (46.)
(syn. *cárnea*) -
Hort. Soc. Gard. *Miss M. L.*
51. *Pàvia rubra* - p. 469. (48.)
Hort. Soc. Gard. *Miss M. L.*
52. *P. r.*, full-grown tree - (48.)
Synon House. *G. R. Lewis.*
53. *P. r. humilis péndula* - p. 470. (52.)
Hort. Soc. Gard. *Miss M. L.*
54. *P. flàva* - p. 471. (50.)
Hort. Soc. Gard. *Miss M. L.*
55. *P. f.*, full-grown tree - (51.)
Synon House. *G. R. Lewis.*
56. *P. macrocárpa* - p. 473. (52.)
Hort. Soc. Gard. *Miss M. L.*

SAPINDA'CEÆ.

57. *Kölreutèria paniculàta* - p. 475. (53.)
Hort. Soc. Gard. *Miss M. L.*

XANTHOXYLA'CEÆ.

58. *Xanthóxyllum fraxíneum* p. 488. (54.)
Hort. Soc. Gard. *H. Le Jeune.*
59. *Ptèlea trifoliàta* - p. 489. (55.)
Hort. Soc. Gard. *Miss M. L.*
60. *Ailántus glandulòsa* - p. 490. (56.)
Hort. Soc. Gard. *Miss M. L.*
61. *A. g.*, full-grown tree - (55a.)
Synon House. *G. R. Lewis.*

CELASTRA'CEÆ.

62. *Euónymus europæus* - p. 496. (57.)
Hort. Soc. Gard. *Miss M. L.*
63. *E. latifólius* - p. 498. (58.)
Hort. Soc. Gard. *Miss M. L.*

AQUIFOLIA'CEÆ.

64. *Ilex Aquifólium* - p. 505. (59.)
Bayswater. *H. Le Jeune.*
65. *I. A.*, full-grown tree - (59a.)
Studley Park. *H. W. Jukes.*
66. *I. opàca* - p. 516. (60.)
Messrs. Loddiges. *Miss M. L.*
67. *I. opàca*, full-grown tree - (59b.)
Synon House. *G. R. Lewis.*

RHAMNA'CEÆ.

68. *Palínurus aculeàtus* - p. 527. (61.)
Hort. Soc. Gard. *Miss M. L.*
69. *P. a.*, full-grown tree - (60b.)
Synon House. *G. R. Lewis.*
70. *Rhámnus cathárticus* - p. 531. (62.)
Hort. Soc. Gard. *Miss M. L.*
71. *R. alpinus* - p. 536. (62c.)
Hort. Soc. Gard. *Miss M. L.*
72. *R. Frángula* - p. 537. (62a.)
Hort. Soc. Gard. *Miss M. L.*
73. *R. latifólius* - p. 538. (62b.)
Hort. Soc. Gard. *Miss M. L.*

HOMALINA'CEÆ.

74. *Aristotèlia Mácqui* - p. 543. (63.)
Fulham Nursery. *H. W. Jukes.*

LEGUMINO'SÆ.

75. *Sophòra japónica* - p. 563. (64.)
Hort. Soc. Gard. *Miss M. L.*
76. *S. j.*, full-grown tree, 4to - (63a.)
Synon House. *G. R. Lewis.*
77. *S. j. péndula* - p. 564. (65.)
Hort. Soc. Gard. *Miss M. L.*
78. *Virgília lútea* - p. 565. (66.)
Hort. Soc. Gard. *Miss M. L.*
79. *Cýtistus Labúrnum* - p. 590. (67.)
Hort. Soc. Gard. *Miss M. L.*
80. *C. L. quercifólium* (syn. } p. 590. (68.)
incisum) -
Hort. Soc. Gard. *Miss M. L.*
81. *C. alpinus* - p. 591. (70.)
Hort. Soc. Gard. *Miss M. L.*
82. *C. a. péndulus* - p. 591. (70a.)
Messrs. Loddiges. *H. Le Jeune.*
83. *Robínia Pseud-Acàcia* - p. 609. (71.)
Hort. Soc. Gard. *Miss M. L.*
84. *R. P.-A.*, full grown tree - 71.*
Kenwood. *W. J. Nesfield.*
85. *R. P.-A. umbraculifera* p. 610. (71a.)
Bayswater. *H. Le Jeune.*
86. *R. P.-A. tortuòsa* - p. 610. (72.)
Hort. Soc. Gard. *H. Le Jeune.*
87. *R. viscòsa* - p. 626. (73.)
Hort. Soc. Gard. *Miss M. L.*
88. *R. hispida macrophýlla* p. 628. (73a.)
Hort. Soc. Gard. *H. Le Jeune.*
89. *Caragàna arboréscens* - p. 629. (75.)
Hort. Soc. Gard. *Miss M. L.*
90. *Gledítischia triacánthos* p. 650. (76.)
Hort. Soc. Gard. *Miss M. L.*
91. *G. t.*, full-grown tree - p. 650. (75a.)
Synon House. *G. R. Lewis.*
92. *G. inermis* - p. 650. (77b.)
Hort. Soc. Gard. *Miss M. L.*
93. *G. i.*, full-grown tree - (76a.)
Synon House. *G. R. Lewis.*
94. *G. sinénsis* (syn. *hórrida*) p. 654. (78.)
Hort. Soc. Gard. *H. Le Jeune.*
95. *G. (s.) japónica* - p. 654. (82.)
Messrs. Loddiges. *H. Le Jeune.*

96. *G. (s.) nana* - p. 654. (80.)
Messrs. Loddiges. *H. Le Jeune.*
97. *G. (s.) purpurea* - p. 654. (78b.)
Messrs. Loddiges. *H. Le Jeune.*
98. *G. (s.) macracantha* - p. 654. (78.)
Messrs. Loddiges. *H. Le Jeune.*
99. *Gymnocladus cana-*
densis - } p. 656. (82a.)
Hort. Soc. Gard. *Miss M. L.*
100. *G. c.*, full-grown tree - (82.)
Syon House. *G. R. Lewis.*
101. *Céreis Siliquástrum* - p. 657. (83.)
Hort. Soc. Gard. *Miss M. L.*
102. *C. S.*, full-grown tree - (82b.)
Syon House. *G. R. Lewis.*
103. *C. canadensis* - p. 659. (83.)
Hort. Soc. Gard. *Miss M. L.*
104. *Acácia dealbata* - p. 666. (74.)
Hort. Soc. Gard. *Miss M. L.*

VOL. VI.

ROSACEÆ.

105. *Amýgdalus commúnis* p. 674. (85.)
Hort. Soc. Gard. *Miss M. L.*
106. *Pérsica vulgáris* - p. 680. (86.)
Hort. Soc. Gard. *Miss M. L.*
107. *Armeniaca vulgáris* - p. 682. (87.)
Hort. Soc. Gard. *Miss M. L.*
108. *A. sibírica* - p. 683. (88.)
Hort. Soc. Gard. *Miss M. L.*
109. *Prúnus spinósa* - p. 684. (90.)
Hort. Soc. Gard. *H. W. Jukes.*
110. *P. insítia* - p. 687. (91.)
Hort. Soc. Gard. *H. W. Jukes.*
111. *P. doméstica myrobá-*
lana - } p. 688. (93.)
Hort. Soc. Gard. *Miss M. L.*
112. *Cérasus sylvéstris* }
(syn. *durácina*) - } p. 693. (95.)
Hort. Soc. Gard. *H. Le Jeune.*
113. *C. s.*, full-grown tree - (95a.)
Studley Park. *H. W. Jukes.*
114. *C. vulgáris* (syn. *ávium*) p. 693. (94.)
Hort. Soc. Gard. *H. Le Jeune.*
115. *C. v. semperflórens* - p. 701. (96.)
Hort. Soc. Gard. *H. Le Jeune.*
116. *C. Maháleb* - - p. 707. (98.)
Hort. Soc. Gard. *H. Le Jeune.*
117. *C. Párus* - p. 709. (99.)
Hort. Soc. Gard. *H. Le Jeune.*
118. *C. serótina* - p. 712. (101.)
Hort. Soc. Gard. *H. Le Jeune.*
119. *C. lusitánica* - p. 714. (97.)
Syon House. *H. Le Jeune.*
120. *C. l.*, full-grown tree - (97a.)
Syon House. *H. Le Jeune.*
121. *Cratrégus cocéinea* - p. 816. (102.)
Hort. Soc. Gard. *H. Le Jeune.*
122. *C. glandulósa* - p. 817. (102a.)
Hort. Soc. Gard. *H. Le Jeune.*
123. *C. punctáta* - p. 818. (104.)
Hort. Soc. Gard. *H. Le Jeune.*
124. *C. pyrífólia* - p. 819. (105.)
Hort. Soc. Gard. *H. Le Jeune.*
125. *C. macracantha* - p. 819. (105.)
Hort. Soc. Gard. *H. Le Jeune.*
126. *C. Crús-gállí* - p. 820. (103.)
Hort. Soc. Gard. *Miss E. Ronalds.*
27. *C. C.*, full-grown tree - (106a.)
Fulham Palace. *H. Le Jeune.*
128. *C. C. pyracanthæfólia* p. 820. (108.)
Hort. Soc. Gard. *H. Le Jeune.*
129. *C. C. salicifólia* - p. 820. (108a.)
Hort. Soc. Gard. *H. Le Jeune.*
130. *C. C. ovalifólia* - p. 821. (108b.)
Hort. Soc. Gard. *H. Le Jeune.*
131. *C. C. prunifólia* - p. 821. (109.)
Hort. Soc. Gard. *H. Le Jeune.*
132. *C. nígra* - p. 822. (110.)
Hort. Soc. Gard. *H. Le Jeune.*
133. *C. purpurea* - p. 822. (111.)
Hort. Soc. Gard. *H. Le Jeune.*
134. *C. Douglási* - p. 823. (112.)
Hort. Soc. Gard. *H. Le Jeune.*
135. *C. fláva* - p. 823. (113.)
Hort. Soc. Gard. *H. Le Jeune.*
136. *C. apiifólia* - p. 824. (115.)
Messrs. Loddiges. *H. Le Jeune.*
137. *C. cordáta* - p. 825. (109a.)
Hort. Soc. Gard. *H. Le Jeune.*
138. *C. spatuláta* - p. 825. (115b.)
Hort. Soc. Gard. *H. Le Jeune.*
139. *C. Azarólus* - p. 826. (116.)
Messrs. Loddiges. *H. Le Jeune.*
140. *C. maroccána* - p. 827. (116a.)
Hort. Soc. Gard. *H. Le Jeune.*
141. *C. Arónia* - p. 827. (117.)
Hort. Soc. Gard. *H. Le Jeune.*
142. *C. orientális* (syn. }
odoratíssima) - } p. 827. (117a.)
Hort. Soc. Gard. *H. Le Jeune.*
143. *C. tanacetifólia* - p. 828. (117b.)
Hort. Soc. Gard. *H. Le Jeune.*
144. *C. t. glábra* - p. 828. (117c.)
Messrs. Loddiges. *H. Le Jeune.*
145. *C. t. Lecána* - p. 828. (117d.)
Leyton Nursery. *H. Le Jeune.*
146. *C. heterophýlla* - p. 829. (117d.)
Hort. Soc. Gard. *H. Le Jeune.*
147. *C. Oxyacantha* - p. 829. (118.)
Hort. Soc. Gard. *H. Le Jeune.*
148. *C. O.*, full-grown tree - (118.)*
Studley Park. *H. W. Jukes.*
149. *C. O. obtusáta* - p. 830. (117e.)
Hort. Soc. Gard. *H. Le Jeune.*
150. *C. O. laciniáta* - p. 830. (118a.)
Hort. Soc. Gard. *H. Le Jeune.*
151. *C. O. eriocárpa* - p. 831. (118b.)
Hort. Soc. Gard. *H. Le Jeune.*

152. *C. O. Oliveriàna* - p. 831. (118c.)
Hort. Soc. Gard. *H. Le Jeunc.*
153. *C. O. melanocárpa* p. 831. (118d.)
Hort. Soc. Gard. *H. Le Jeunc.*
154. *C. O. strícta* - p. 832. (118c.)
Hort. Soc. Gard. *Miss E. Ronatds.*
155. *C. O. reginæ* - p. 832. (118f.)
Messrs. Loddiges. *H. Le Jeunc.*
156. *C. mexicàna* - p. 843. (119.)
Messrs. Loddiges. *H. Le Jeunc.*
157. *Photínia serrulàta* - p. 868. (120.)
Syon House. *H. Le Jeunc.*
158. *Cotoneáster frígida* p. 871. (122a.)
Hort. Soc. Gard. *H. Le Jeunc.*
159. *C. affinis* - p. 871. (122a.)
Hort. Soc. Gard. *H. Le Jeunc.*
160. *C. acuminàta* - p. 872. (122c.)
Fulham Nursery. *Miss M. L.*
161. *C. nummulària* - p. 872. (122b.)
Hort. Soc. Gard. *H. Le Jeunc.*
162. *Amelánchier Botry-* }
àpium - } p. 874. (121.)
Hort. Soc. Gard. *H. Le Jeunc.*
163. *A. Botryap.*, full-grown tree (121a.)
Syon House. *G. R. Lewis.*
164. *Méspilus germànica* - p. 877. (123.)
Hort. Soc. Gard. *H. Le Jeunc.*
165. *M. Smíthii* (syn. }
grandiflora) - } p. 878. (124.)
Hort. Soc. Gard. *H. Le Jeunc.*
166. *Pýrus communis* - p. 880. (125.)
Hort. Soc. Gard. *H. Le Jeunc.*
167. *P. c.*, full-grown tree, 4to - (127.)
Hamley Cottage. *G. R. Lewis.*
168. *P. c. var. glout morceau* p. 881. (126.)
Bayswater. *H. Le Jeunc.*
169. *P. sinàica* - p. 889. (130.)
Hort. Soc. Gard. *H. Le Jeunc.*
170. *P. sinénsis* - p. 889. (130.)
Hort. Soc. Gard. *Miss M. L.*
171. *P. bollwylleriàna* - p. 890. (129.)
Hort. Soc. Gard. *H. W. Jukes.*
172. *P. variolòsa* - p. 891. (131.)
Hort. Soc. Gard. *H. W. Jukes.*
173. *P. Málus* - p. 891. (132.)
Hort. Soc. Gard. *H. Le Jeunc.*
174. *P. M.*, full-grown tree - (133.)
Tibberton. *G. R. Lewis.*
175. *P. baccàta* - p. 892. (134.)
Hort. Soc. Gard. *Miss M. L.*
176. *P. (M.) coronària* - p. 908. (137.)
Hort. Soc. Gard. *H. W. Jukes.*
177. *P. (M.) angustifolia* p. 909. (138.)
Hort. Soc. Gard. *H. Le Jeunc.*
178. *P. (M.) spectàbilis* p. 909. (132a.)
Hort. Soc. Gard. *H. Le Jeunc.*
179. *P. Aria undulàta* - p. 910. (139a.)
Hort. Soc. Gard. *H. W. Jukes.*
180. *P. vestità* - p. 912. (139.)
Messrs. Loddiges. *Miss M. L.*
181. *P. torminàlis* - p. 913. (139.)
Hort. Soc. Gard. *H. W. Jukes.*
182. *P. pinuacífida* - p. 915. (137d.)
Hort. Soc. Gard. *H. Le Jeunc.*
183. *P. aucupària* - p. 916. (143.)
Hort. Soc. Gard. *H. Le Jeunc.*
184. *P. americàna* - p. 920. (142.)
Hort. Soc. Gard. *H. W. Jukes.*
185. *P. Sòrbus* - p. 921. (145.)
Messrs. Loddiges. *Miss M. L.*
186. *P. S. lanuginòsa* - p. 924. (146.)
Hort. Soc. Gard. *H. W. Jukes.*
187. *P. spùria* - p. 924. (143.)
Hort. Soc. Gard. *H. W. Jukes.*
188. *Cydònia vulgàris* - p. 929. (146.)
Hort. Soc. Gard. *H. Le Jeunc.*
189. *C. sinénsis* - p. 931. (948.)
Hort. Soc. Gard. *H. Le Jeunc.*
- MYRTACEÆ.
190. *Eucalyptus robùsta* - p. 959. (148.)
Stamford Hill. *L. Martin.*
- CORNA'CEÆ.
191. *Córnus más* - p. 1014. (149a.)
Hort. Soc. Gard. *H. Le Jeunc.*
- CAPRIFOLIA'CEÆ.
192. *Sambucus nigra* - p. 1027. (150.)
Hort. Soc. Gard. *H. Le Jeunc.*
193. *Vibúrnum prunifolium* p. 1034. (151.)
Hort. Soc. Gard. *H. Le Jeunc.*
- ERICA'CEÆ.
194. *Arbutus hýbrida* - p. 1119. (152.)
Hort. Soc. Gard. *H. Le Jeunc.*
195. *A. Andràchne* - p. 1120. (152a.)
Messrs. Loddiges. *H. Le Jeunc.*
- HALESIA'CEÆ.
196. *Halèsia tetràptera* - p. 1190. (154.)
Hort. Soc. Gard. *H. Le Jeunc.*
197. *H. t.*, full-grown tree - (153.)
Syon House. *G. R. Lewis.*
- EBENA'CEÆ.
198. *Diospýros Lótus* - p. 1194. (155a.)
Bayswater. *H. Le Jeunc.*
199. *D. L.*, full-grown tree - (155.)
Syon House. *G. R. Lewis.*
200. *D. virginiana* - p. 1195. (156a.)
Hort. Soc. Gard. *H. Le Jeunc.*
201. *D. v.*, full-grown tree - (155b.)
Syon House. *G. R. Lewis.*
- OLEA'CEÆ.
202. *Fráxinus excélsior* - p. 1214. (157.)
Hort. Soc. Gard. *H. Le Jeunc.*
203. *F. e.*, full-grown tree - (156b.)
Kensington Gardens. *J. Martin.*
204. *F. e.*, full-grown winter }
tree, 4to - } (156c.)
Studley Park. *H. W. Jukes.*
205. *F. e. péndula* - p. 1214. (158.)
Hort. Soc. Gard. *H. Le Jeunc.*

206. *F. simplicifolia* - p. 1228. (159.)
Hort. Soc. Gard. *H. Le Jeune.*
207. *F. parvifolia* - p. 1229. (63.)
Hort. Soc. Gard. *H. Le Jeune.*
208. *F. leutiscifolia* - p. 1231. (166.)
Hort. Soc. Gard. *H. Le Jeune.*
209. *F. americana* - p. 1232. (161.)
Hort. Soc. Gard. *H. Le Jeune.*
210. *F. juglandifolia*, full-
grown tree, 4to } p. 1236. (163.)
Pope's Villa, Twickenham. *H. Le Jeune.*
211. *F. épiptera* (syn. }
láncea) - } p. 2137. (162.)
Hort. Soc. Gard. *H. Le Jeune.*
212. *F. pannosa*, full-
grown tree - } p. 1240. (162a.)
Ham House, Essex. *L. Martin.*
213. *O. rnus europæa* p. 1241. (167a.)
Hort. Soc. Gard. *H. Le Jeune.*
214. *O. e.*, full-grown tree - (167.)
Syon House. *G. R. Lewis.*

VOL. VII.

BIGNONIACEÆ.

215. *Catálpa syringæfolia* p. 1261. (169.)
Bayswater. *H. Le Jeune.*
216. *C. s.*, full-grown tree - (170.)
Syon House. *G. R. Lewis.*

LAURACEÆ.

217. *Laúrus nobilis* - p. 1297. (171.)
Hort. Soc. Gard. *H. Le Jeune.*
218. *L. Sássafras* - p. 1301. (172a.)
Kensington Nursery. *Miss M. L.*
219. *L. S.*, full-grown tree - (172.)
Syon House. *G. R. Lewis.*

SANTALACEÆ.

220. *Nýssa triflora*, full-
grown tree - } p. 1317. (173.)
Thompson's Villa, Richmond. *H. Le Jeune.*

ELEGNACEÆ.

221. *Elaëagnus orientalis* p. 1321. (175.)
Hort. Soc. Gard. *H. Le Jeune.*
222. *Hippóphæ Rham-*
noides angustifolia } p. 1325. (174a.)
Messrs. Loddiges. *C. Rauch.*

URTICACEÆ.

223. *Morus nigra* - p. 1343. (182.)
Messrs. Loddiges. *H. Le Jeune.*
224. *M. n.*, full-grown tree - (182a.)
Battersea. *H. W. Jukes.*
225. *M. álba* - p. 1348. (181.)
Hort. Soc. Gard. *H. Le Jeune.*
226. *M. a.*, full-grown tree, 4to (181a.)
Syon House. *H. Le Jeune.*
227. *M. rubra* (syn. }
pennsylváica) } p. 1359. (183.)
Hort. Soc. Gard. *Miss M. L.*
228. *Broussonétia papy-*
rifera - } p. 1361. (180.)
Kew Botanic Gardens. *H. L. Jeune.*
229. *Ficus Carica* - p. 1365. (178.)
Mile End Nursery. *H. W. Jukes.*

ULMACEÆ.

230. *Ulmus campestris* p. 1374. (185.)
Hort. Soc. Gard. *Miss M. L.*
231. *U. c.*, full-grown tree - (184.)
Kensington Gardens. *J. Martin.*

232. *U. c. strícta*, full-
grown tree - } p. 1375. (184a.)
Muswell Hill. *Miss M. L.*

233. *U. c. viminalis* - p. 1376. (185a.)
Hort. Soc. Gard. *H. W. Jukes.*

234. *U. planifolia* - p. 1377. (191b.)
Hort. Soc. Gard. *H. W. Jukes.*

235. *U. (c.) suberosa* - p. 1395. (186.)
Hort. Soc. Gard. *H. W. Jukes.*

236. *U. maior* - p. 1396. (188.)
Hort. Soc. Gard. *H. Le Jeune.*

237. *U. m.*, full-grown tree - (187.)
Studley Park. *H. W. Jukes.*

238. *U. effusa* - p. 1397. (191.)
Hort. Soc. Gard. *H. W. Jukes.*

239. *U. e.*, winter tree - (191a.)
Hort. Soc. Gard. *H. W. Jukes.*

240. *U. montana maior*, p. 1398. (188 a.)
Hort. Soc. Gard. *H. W. Jukes.*

241. *U. m. pendula* (syn. }
rubra) - } p. 1398. (190.)
Hort. Soc. Gard. *C. Rauch.*

242. *U. m. fastigiata* p. 1399. (187a.)
Hort. Soc. Gard. *H. Le Jeune.*

243. *U. americana* - p. 1406. (189.)
Hort. Soc. Gard. *H. Le Jeune.*

244. *U. a.*, incisa - p. 1406 (189a.)
Hort. Soc. Gard. *H. Le Jeune.*

245. *Plánera Richárdi* p. 1409 (184a.)
Hort. Soc. Gard. *H. Le Jeune.*

246. *P. R.*, full-grown tree - (184.)
Syon House. *G. R. Lewis.*

247. *Céltis Tournefórtii* p. 1416. (194.)
Hort. Soc. Gard. *Miss M. L.*

248. *C. occidentalis* p. 1417. (193.)
Hort. Soc. Gard. *Miss M. L.*

249. *C. o.*, full-grown tree - (192a.)
Syon House. *G. R. Lewis.*

JUGLANDACEÆ.

250. *Juglans régia* - p. 1423. (196.)
Hort. Soc. Gard. *Miss M. L.*

251. *J. r.*, full-grown tree - (195a.)
Chiswick Villa. *W. A. Nesfield.*

252. *J. r.*, full-grown winter tree (196a.)
Studley Park. *H. W. Jukes.*

253. *J. nigra* - p. 1435. (197.)
Hort. Soc. Gard. *C. Rauch.*

254. *J. n.*, full-grown tree, 4to - (197a.)
Syon House. *G. R. Lewis.*
255. *J. cinèrea* - p. 1439. (198.)
Hort. Soc. Gard. *C. Rauch.*
256. *Càrya álba*, full-
grown tree - } p. 1446. (200.)
Purser's Cross. *L. Martin.*
257. *Pterocàrya caucásica* p. 1451 (199.)
Hort. Soc. Gard. *C. Rauch.*
- SALICA'CEE.
258. *Sàlix babylónica* - p. 1507. (207.)
Hort. Soc. Gard. *H. W. Jukes.*
259. *S. b. críspa* - p. 1514. (208.)
Hort. Soc. Gard. *H. W. Jukes.*
260. *S. frágilis* - p. 1516. (205.)
Hort. Soc. Gard. *H. Le Jeune.*
261. *S. álba* - - p. 1522. (209.)
Hort. Soc. Gard. *H. Le Jeune.*
262. *S. a.*, full-grown tree - (209a.)
Turnham Green. *H. W. Jukes.*
263. *S. vitellina* - p. 1528. (206.)
Hort. Soc. Gard. *H. W. Jukes.*
264. *Pópulus álba* - p. 1638. (214.)
Hort. Soc. Gard. *Miss J. L.*
265. *P. canéscens* - p. 1639. (215.)
Hort. Soc. Gard. *Miss J. L.*
266. *P. trémula* - p. 1645. (217.)
Hort. Soc. Gard. *Miss J. L.*
267. *P. t. péndula* - p. 1646. (218.)
Messrs. Loddiges. *H. Le Jeune.*
268. *P. græ'ca* - p. 1651. (219.)
Hort. Soc. Gard. *Miss J. L.*
269. *P. nìgra* - p. 1652. (229.)
Hort. Soc. Gard. *H. Le Jeune.*
270. *P. n.*, winter tree - - (220.)
Hort. Soc. Gard. *H. Le Jeune.*
271. *P. n.*, full-grown tree, 4to - (221.)
Lambeth Palace. *Miss M. L.*
272. *P. monilífera* - p. 1657. (222.)
Hort. Soc. Gard. *Miss J. L.*
273. *P. m.*, winter tree - - (222.)
Bayswater. *H. Le Jeune.*
274. *P. m.*, full-grown tree, 4to. (222a.)
Syon House. *G. R. Lewis.*
275. *P. fastigiàta* } p. 1660. (221. or 216.)
(syn. *dilatàta*) }
Hort. Soc. Gard. *Miss J. L.*
276. *P. angulàta* - p. 1670. (224.)
Hort. Soc. Gard. *Miss J. L.*
277. *P. a.*, full-grown tree - (224.)
Syon House. *G. R. Lewis.*
278. *P. balsamífera* - p. 1673. (225.)
Leyton Nursery. *H. Le Jeune.*
279. *P. b.*, full-grown tree - (224.)
Syon House. *H. Le Jeune.*
280. *P. cándicans* - p. 1676. (227.)
Hort. Soc. Gard. *Miss J. L.*
- BETULA'CEE.
281. *A'lnus glutinòsa* - p. 1678. (229.)
Hort. Soc. Gard. *Miss M. L.*
282. *A. g. laciniàta* - p. 1678. (239.)
Hort. Soc. Gard. *H. W. Jukes.*
283. *A. g. l.*, full-grown tree, 4to (231.)
Syon House. *W. A. Nesfield.*
284. *A. cordifòlia* - p. 1689. (232.)
Hort. Soc. Gard. *Miss M. L.*
285. *Bétula álba péndula* p. 1691. (233.)
Hort. Soc. Gard. *Miss M. L.*
286. *B. populifòlia* - p. 1707. (237.)
Hort. Soc. Gard. *Miss M. L.*
287. *B. papyràca* - p. 1708. (241.)
Hort. Soc. Gard. *Miss M. L.*
288. *B. nìgra* - p. 1710. (234.)
Hackney Arboretum. *H. Le Jeune.*
289. *B.*, winter tree - - (235.)
Fulham Nursery. *L. Martin.*
- CORYLA'CEE.
290. *Quércus pedunculàta* p. 1731. (282.)
Hort. Soc. Gard. *Miss M. L.*
291. *Q. p.*, full-grown tree - (281a.)
Kensington Gardens. *J. Martin.*
292. *Q. p.*, winter tree - - (282a.)
Studley Park. *H. W. Jukes.*
293. *Q. p. fastigiàta* - p. 1731. (282b.)
Hort. Soc. Gard. *Miss M. L.*
294. *Q. sessiliflòra* - p. 1736. (281.)
Hort. Soc. Gard. *Miss M. L.*
295. *Q. s.*, full-grown tree, 4to (280**)
Woburn Abbey. *W. A. Nesfield.*
296. *Q. s.*, winter tree - (281*)
Studley Park. *H. W. Jukes.*
297. *Q. E'sculus* - p. 1844. (261.)
Hort. Soc. Gard. *Miss M. L.*
298. *Q. Cérris* - p. 1846. (275.)
Hort. Soc. Gard. *Miss M. L.*
299. *Q. C.*, full-grown tree, 4to (277.)
Syon House. *G. R. Lewis.*
300. *Q. C.*, winter tree, 4to - (276.)
Muswell Hill. *Miss M. L.*
301. *Q. C. austriàca* - p. 1848. (279.)
Hort. Soc. Gard. *Miss M. L.*
302. *Q. C. fulhaménsis* } p. 1850. (278.)
(syn. *dentàta*) }
Hort. Soc. Gard. *Miss M. L.*
303. *Q. C. f.*, full-grown tree - (278a.)
Fulham Nursery. *L. Martin.*
304. *Q. C. f.*, winter tree, 4to - (278b.)
Fulham Nursery. *H. Le Jeune.*
305. *Q. C. Lucombeàna* p. 1857. (280.)
Hort. Soc. Gard. *Miss M. L.*
306. *Q. C. L.*, winter tree - (280a.)
Fulham Nursery. *H. Le Jeune.*
307. *Q. Æ'gilops* - p. 1861. (266.)
Hackney Arboretum. *H. Le Jeune.*
308. *Q. Æ.*, full-grown tree - (267.)
Syon House. *G. R. Lewis.*
309. *Q. álba* - p. 1864. (252.)
Hackney Arboretum. *H. Le Jeune.*

Vol. VIII.

310. *Q. macrocarpa* - p. 1869. (254.)
Hackney Arboretum. *H. Le Jeune.*
311. *Q. obtusiloba* - p. 1870. (253.)
Hort. Soc. Gard. *Miss M. L.*
312. *Q. Prinus palustris* p. 1872. (262.)
Leyton Nursery. *Miss M. L.*
313. *Q. P. monticola* - p. 1873. (263.)
Leyton Nursery. *Miss M. L.*
314. *Q. P. tomentosa* - p. 1876. (262.)
Leyton Nursery. *Miss M. L.*
315. *Q. rubra* - - p. 1877. (269.)
Hort. Soc. Gard. *Miss M. L.*
316. *Q. r.*, full-grown tree, 4to - (271.)
Syon House. *G. R. Lewis.*
317. *Q. r.*, winter tree - (270.)
Fulham Nursery. *L. Martin.*
318. *Q. coccinea* - p. 1879. (273.)
Purser's Cross. *Miss M. L.*
319. *Q. c.*, full-grown tree - (273*a*.)
Muswell Hill. *Miss M. L.*
320. *Q. ambigua* - p. 1881. (272.)
Hort. Soc. Gard. *Miss M. L.*
321. *Q. tinctoria* - p. 1884. (268.)
Hackney Arboretum. *H. Le Jeune.*
322. *Q. palustris* - p. 1887. (274.)
Leyton Nursery. *H. Le Jeune.*
323. *Q. palustris*, full-grown tree (274*a*.)
Syon House. *H. Le Jeune.*
324. *Q. Phellos*, 4to - p. 1894 (258.)
Syon House. *G. R. Lewis.*
325. *Q. P. latifolia* - p. 1895. (249.)
Hackney Arboretum. *L. Martin.*
326. *Q. Ilex*, full-grown } p. 1899. (255.)
tree - }
Syon House. *G. R. Lewis.*
327. *Q. I. latifolia* - p. 1899. (255.)
Hort. Soc. Gard. *Miss M. L.*
328. *Q. I. longifolia* } p. 1900. (256.)
(syn. *angustifolia*) }
Hort. Soc. Gard. *Miss M. L.*
329. *Q. gramuntia*, full- } p. 1906. (260.)
grown tree - }
Purser's Cross. *Miss M. L.*
330. *Q. Süber*, full-grown } p. 1911. (257.)
tree - }
Fulham Nursery. *L. Martin.*
331. *Q. S.*, full-grown tree p. 1911. (258.)
Muswell Hill. *H. Le Jeune.*
332. *Q. S. deutata*, full- } p. 1913. (259.)
grown tree - }
Muswell Hill. *H. Le Jeune.*
333. *Q. virens* - p. 1918. (260.)
Chiswick Villa. *L. Martin.*
334. *Fagus sylvatica* - p. 1950. (283.)
Hort. Soc. Gard. *H. Le Jeune.*
335. *F. s. cristata* (syn. } p. 1952. (283*a*.)
crispa) - }
Hort. Soc. Gard. *H. Le Jeune.*
336. *F. s. pendula* - p. 1952. (284.)
Kensington Nursery. *Miss M. L.*
337. *F. s. p.*, full-grown tree - (285.)
Oundle. *M. J. B.*
338. *Castanea vesca* - p. 1983. (286.)
Hort. Soc. Gard. *Miss M. L.*
339. *C. v.*, full-grown tree - (287.)
Muswell Hill. *Miss M. L.*
340. *Carpinus Bétulus* - p. 2004. (244.)
Hort. Soc. Gard. *Miss M. L.*
341. *C. B.*, full-grown tree (243.)
Chiswick Villa. *L. Martin.*
342. *Ostrya vulgaris* - p. 2015. (244*a*.)
Hort. Soc. Gard. *Miss M. L.*
343. *O. v.*, full-grown tree - (245.)
Bot. Gard. Kew. *Miss M. L.*
344. *O. virginica* - p. 2015. (246.)
Hort. Soc. Gard. *C. Rauch.*
345. *Corylus Colurna* - p. 2029. (247.)
Mile End Nursery. *H. W. Jukes.*
346. *C. C.*, full-grown tree - (246*a*.)
Syon House. *G. R. Lewis.*
- PLATANACEE.
347. *Platanus orientalis* p. 2033. (288.)
Hort. Soc. Gard. *Miss M. L.*
348. *P. o.*, winter tree - - (287.)
Hort. Soc. Gard. *Miss M. L.*
349. *P. o. acerifolia* - p. 2034. (289.)
Hort. Soc. Gard. *H. W. Jukes.*
350. *P. o. a.*, full-grown tree - (289*a*.)
Elmhurst, Finchley. *W. A. Nesfield.*
351. *P. o. cuneata* - p. 2034. (288.)
Hort. Soc. Gard. *Miss M. L.*
352. *P. occidentalis* - p. 2043. (289*a*.)
Hort. Soc. Gard. *H. W. Jukes.*
- BALSAMACEE.
353. *Liquidambar Styra-* } p. 2049. (291.)
ciflua - }
354. *L. S.*, full-grown tree - - (290.)
Syon House. *G. R. Lewis.*
- TAXACEE.
355. *Taxus baccata* - p. 2066. (293.)
Hort. Soc. Gard. *Miss M. L.*
356. *T. b.*, tree at Harlington - (393*a*.)
Harlington. *Miss M. L.*
357. *T. b.*, tree at Darley - (393*b*.)
Darley. *J. E. Bowman.*
358. *T. b. fastigiata* - p. 2066. (294.)
Hort. Soc. Gard. *Miss M. L.*
359. *Salisbùria adiantifolia*. p. 2094. (292*a*.)
Messrs. Loddiges. *Miss M. L.*
360. *S. a.*, full-grown tree - (292.)
Mile End Nursery. *H. W. Jukes.*
- CONIFERE, § ABIÉTINE.
361. *Pinus sylvestris* - p. 2153. (312*a*.)
Hort. Soc. Gard. *H. Le Jeune.*
362. *P. s.*, full-grown tree - (312.)
Pain's Hill. *H. Le Jeune.*

363. *P. s.*, full-grown tree - (313.)
Muswell Hill. *W. A. Nesfield.*
364. *P. s. genevènsis* - p. 2158 (313a.)
Hort. Soc. Gard. *H. Le Jeune.*
365. *P. Larício* - p. 2200. (314.)
Hort. Soc. Gard. *H. Le Jeune.*
366. *P. L.*, full-grown tree - (315.)
Kew. *H. Le Jeune.*
367. *Pallasiana* - p. 2206. (318.)
Hort. Soc. Gard. *G. R. Lewis.*
368. *P. Pináster* - p. 2213. (316.)
Hort. Soc. Gard. *H. Le Jeune.*
369. *P. P.*, full-grown tree - (316a.)
Hendon. *L. Martin.*
370. *P. Pinea* - p. 2224. (319.)
Hort. Soc. Gard. *H. Le Jeune.*
371. *P. halepénsis* - p. 2231. (321.)
Hort. Soc. Gard. *G. R. Lewis.*
372. *P. Tæ'da*, full-grown tree, - } p. 2237. (325.)
Syon House. *G. R. Lewis.*
373. *P. rígida* - p. 2239. (326a.)
Dropmore. *H. Le Jeune.*
374. *P. rígida*, full-grown tree - (226.)
Pain's Hill. *H. Le Jeune.*
375. *P. serótina* - p. 2242. (327.)
Dropmore. *H. Le Jeune.*
376. *P. Cémbra* - p. 2274. (331.)
Kew. *L. Martin.*
377. *P. Stròbus* - p. 2280. (329.)
Hort. Soc. Gard. *H. Le Jeune.*
378. *Abies excélsa* - p. 2293. (338.)
Hort. Soc. Gard. *H. Le Jeune.*
379. *A. e.*, full-grown tree - (337.)
Syon House. *H. Le Jeune.*
380. *A. e.*, full-grown tree - (336.)
Studley Park. *H. W. Jukes.*
381. *A. e. nigra* - p. 2294. (338a.)
Studley Park. *H. W. Jukes.*
382. *A. álba* - p. 2310. (339.)
Hort. Soc. Gard. *H. Le Jeune.*
383. *A. Smithiana* - p. 2317. (340.)
Hort. Soc. Gard. *H. Le Jeune.*
384. *A. Douglasi* - p. 2319. (333.)
Hort. Soc. Gard. *H. Le Jeune.*
385. *A. canadénsis* - p. 2322. (335.)
Syon House. *H. Le Jeune.*
386. *A. c.*, full-grown tree - (335a.)
Whitton Park. *H. Le Jeune.*
387. *A. c.*, full-grown tree - (335b.)
Studley Park. *H. W. Jukes.*
388. *Picea pectinata* - p. 2329. (332.)
Leyton Nursery. *H. Le Jeune.*
389. *P. p.*, full-grown tree - (332a.)
Syon House. *H. Le Jeune.*
390. *P. p.*, full-grown tree - (332b.)
Rosenath. *Copied from Strutt.*
391. *P. balsàmea* - p. 2339. (334.)
Hort. Soc. Gard. *H. Le Jeune.*
392. *Làrix europæ'a* - p. 2350. (344.)
Hort. Soc. Gard. *H. Le Jeune.*
393. *L. e.*, large tree - - (344a.)
Kenwood. *G. R. Lewis.*
394. *L. microcarpa ame-* }
ricana - - } p. 2399. (347.)
Hort. Soc. Gard. *H. Le Jeune.*
395. *L. a.*, full-grown tree - (346.)
Syon House. *H. Le Jeune.*
396. *Cèdrus Libàni* - p. 2402. (348.)
Hort. Soc. Gard. *H. Le Jeune.*
397. *C. L.*, full-grown tree, 4to - (349.)
Syon House. *G. R. Lewis.*
398. *C. L.*, full-grown tree - (350.)
Kenwood. *G. R. Lewis.*
399. *C. L.*, full-grown tree - (351.)
Foxley Hall. *G. R. Lewis.*
400. *C. L.*, glàuca, full-grown tree (352.)
Thompson's Villa. *H. Le Jeune.*

CUPRE'SSINÆ.

401. *Thùja occidentàlis* p. 2454. (302.)
Hort. Soc. Gard. *Miss M. L.*
402. *T. orientàlis stricta* p. 2459. (303.)
Hort. Soc. Gard. *Miss M. L.*
403. *Cupressus semper-* }
virens - - } p. 2464. (306.)
Hort. Soc. Gard. *Miss M. L.*
404. *C. s. horizontàlis* - p. 2464. (307.)
Hort. Soc. Gard. *Miss M. L.*
405. *C. lusitànica* - p. 2477. (308.)
Hort. Soc. Gard. *Miss M. L.*
406. *Taxodium distichum* p. 2481. (295a.)
Hort. Soc. Gard. *Miss M. L.*
407. *T. d.*, full-grown tree - (295.)
Syon House. *G. R. Lewis.*
408. *Juniperus phænicea* p. 2501. (301.)
Hort. Soc. Gard. *Miss M. L.*
409. *J. p.*, full-grown tree - (300.)
Syon House. *G. R. Lewis.*
410. *J. virginiana* - p. 2495. (298.)
Hort. Soc. Gard. *Miss M. L.*
411. *J. v.*, full-grown tree - (207a.)
Syon House. *G. R. Lewis.*
412. *J. chinénsis (excélsa)* p. 2505. (297.)
Hort. Soc. Gard. *Miss M. L.*

LIST OF THE PORTRAITS OF ENTIRE TREES AND SHRUBS
GIVEN ALONG WITH THE TEXT IN VOLS. I. II. III. AND IV.

		Figure	Page			Figure	Page
<i>Magnoliàcææ.</i>				<i>Betulàcææ.</i>			
<i>Magnòlia conspícua</i>	-	35	281	<i>Álmus glutinòsa laciniàta</i>	-	1542	1686
<i>Rosàcææ.</i>				<i>Bétula álba</i>	-	1550	1693
<i>Ròsa índica</i>	-	546	809	<i>Corylàcææ.</i>			
<i>Crataégus Crús-gállí</i>	} 551. to 553		820	<i>Quércus pedunculàta</i>	-	1580	1741
<i>salicifòlia</i>				-	556	833	-
<i>Oxyacántha regína</i>	-	641	914	-	1582	1742	
<i>Pýrus torminàlis</i>	-	644	922	-	1587	1746	
<i>Sórbus</i>	-			<i>pedunculàta péndula</i>	-	1568	1732
<i>Myrtàcææ.</i>				<i>sessiliflòra</i>	-	1585, 1586	1744 1745
<i>Melaleúca linearifòlia</i>	-	685	957	<i>pedunculàta or sessiliflòra</i>			
<i>Eucalýptus resinífera</i>	-	691	959	<i>Oak shattered by lightning</i>	1643	1813	
<i>robústa</i>	-	2467	2567	<i>Herne's Oak</i>	- 1588,	1589	1755
<i>piperita</i>	-	690	959	<i>Wootton Oak</i>	-	1590	1756
<i>Angéphora lanceolàta</i>	-	702	961	<i>Meavy Oak</i>	-	1591	1757
<i>Leptospérmum scopàrium</i>	-	2468	2568	<i>Flitton Oak</i>	-	1592	1758
<i>Hamamelidàcææ.</i>				<i>Hatfield Broad Oak</i>	-	1593	1759
<i>Hamamélis virgínica</i>	-	757	1008	<i>Shordley Oak</i>	-	1594	1760
<i>Ericàcææ.</i>				<i>Moceas Park Oak</i>	-	1595	1762
<i>Árbutus Andràehne</i>	-	2493	2575	<i>Panshanger Oak</i>	-	1596	1762
<i>Oleàcææ.</i>				<i>Majesty Oak</i>	-	1597	1762
<i>Ligústrum læidum</i>	-	1023	1202	<i>Stately Oak</i>	-	1598	1762
<i>Chionánthus virgínica</i>	-	1030	1206	<i>Sir Philip Sydney's Oak</i>	-	1599	1763
<i>Fráxinus excélsior</i>	-	1019	1226	<i>Nannau Oak</i>	-	1600	1763
<i>péndula</i>	-	1045	1216	<i>Chandos Oak</i>	-	1601	1763
<i>Órnus europæa</i>	-	1067	1243	<i>Merton Oak</i>	-	1602	1764
<i>Solanàcææ.</i>				<i>Winfarthing Oak</i>	- 1603.	1623	1764 1775
<i>Brugmànsia suavèolens</i>	-	1120	1275	<i>Gog Oak</i>	-	1604	1765
<i>Elæagnàcææ.</i>				<i>Magog Oak</i>	-	1605	1765
<i>Elæágnus argénteá</i>	-	1204	1323	<i>Salecy Forest Oak</i>	-	1606	1766
<i>Urticàcææ.</i>				<i>The Duke's Walkingstick</i>	1607	1766	
<i>Mórus nígra</i>	-	1222	1345	<i>Greendale Oak</i>	- 1608,	1609	1767
<i>at Canterbury</i>	-	2526	2586	<i>Parliament Oak</i>	-	1610	1767
<i>Ulmàcææ.</i>				<i>Shelton Oak</i>	-	1611	1768
<i>Ulmus campéstris</i>	-	1238, 1239	1392 1394	<i>Swilear Lawn Oak</i>	-	1612	1769
<i>montàna</i>	-	1243, 1244	1402 1403	<i>Squith Oak</i>	-	1613	1769
<i>Plánera Richárdí</i>	-	1250	1412	<i>Beggar's Oak</i>	-	1614	1769
<i>Salicàcææ.</i>				<i>Queen Elizabeth's Oak</i>	-	1615	1770
<i>salix babylònica</i>	-	1308	1514	<i>Bull Oak</i>	-	1616, 1625	1770 1780
<i>Russelliàna</i>	-	1312, 1313	1520 1521	<i>Gospel Oak</i>	-	1617, 1628	1771 1781
<i>álba</i>	-	1316	1527	<i>Creeping Oak</i>	-	1618, 1632	1771 1792
<i>Pópulus nígra</i>	-	1514	1655	<i>King Oak</i>	-	1619, 1633	1771 1792
<i>fastigiàta</i>	-	1519	1660	<i>Cowthorpe Oak</i>	-	1620, 1624	1771 1776
				<i>Wallace Oak</i>	-	1621	1772
				<i>Chapel Oak of Allonville</i>	1622	1773	
				<i>Specimen of an old Oak</i>	-	1635	1794
				<i>Cobham Oak</i>	-	1636	1794
				<i>Ashton Oak</i>	-	1693	1838
				<i>Weishaden Oak</i>	-	1695	1842

	Figure	Page		Figure	Page	
<i>Q. Cériss Lucombeana</i>	- { 1712	1852	<i>Pinus resinosa</i>	- - -	2097 2212	
<i>crispa</i>	- - -	1713 1853	<i>Pinea</i>	- - -	2108 2227	
<i>álba</i>	- - -	1718 1858	<i>Tæ'da</i>	- - -	2122 2239	
<i>palústris</i>	- - -	1727 1866	<i>ponderosa</i>	- - -	2136 2245	
<i>Süber</i>	- - -	1761 1888	<i>Sabiniàna</i>	- - -	2141 2249	
<i>Fâgus sylvática</i>	- - -	1080 1916	<i>Coúlteri</i>	- - -	2147 2252	
<i>Inosculated Beech</i>	1881. 1884	1878 1955	<i>longifolia</i>	- - -	2152 2253	
<i>Great Beech at Windsor</i>	{ 1907	1976	<i>austrális</i>	- - -	2160 2258	
<i>Queen Beech</i>	- - -	1908 1977	<i>canariënsis</i>	- - -	2166 2263	
<i>Ashridge Beeches</i>	1910. 1911	1909 1977	<i>leiophýlla</i>	- - -	2187 2274	
<i>Eccles Beech</i>	- - -	1909 1977	<i>Cembra</i>	- - -	2192 2277	
<i>Earl's Mill Beech</i>	- - -	1912 1978	<i>Stròbus</i>	- - -	2196 2282	
<i>Grange Hall Beech</i>	- - -	1913 1979	<i>excélsa</i>	- 2201. 2202	{ 2287	
<i>Marriage Beech</i>	- - -	1914 1979	<i>Lambertiàna</i>	- - -	{ 2288	
<i>Castânea véscã</i>	- - -	2536 2594	<i>Abies excélsa</i>	- - 2214. 2218	{ 2290	
<i>Tortworth Chestnut</i>	- - -	1923 1986	<i>nigra</i>	- - 2226. 2227	{ 2313	
<i>Cobham Chestnut</i>	- - -	1924 1988	<i>Picea pectinàta</i>	at Roseneath - 2239	2332	
<i>Waldershaw Chestnut</i>	- - -	1925 1989	<i>at Studley</i>	- 2542	2602	
<i>Cárpinus Bétulus</i>	- 1933. 1935	{ 2005	<i>Webbiàna</i>	- - -	2253 2345	
		{ 2007	<i>Làrix europæ'a</i>	at Syon - 2259	2355	
			<i>Dunkeld</i>	- 2260	2355	
<i>Platanàccea.</i>			<i>Dalwick</i>	- { 2261	{ 2356	
<i>Tabernæmontàna alternifolia</i>	- 1956	2036	<i>Cèdrus Libàni</i>	at Syon - 2268	2404	
<i>Plátanus orientális</i>	- 1957	2037	<i>Enfield</i>	- 2269	2404	
<i>occidentális</i>	- - -	1960 2044	<i>Chelsea</i>	- 2270	2405	
<i>Casuaràccea.</i>			<i>Croome</i>	- 2271	2405	
<i>Casuarina equisetifolia</i>	- - -	1972 2061	<i>Hammersmith</i>	- 2272	2406	
<i>Taxàccea.</i>			<i>Gray</i>	- 2273	2406	
<i>Taxus baccèata fastigiàta</i>	at	} 1982 2067	<i>Quenby Hall</i>	- 2544	2603	
<i>Comber</i>	- - -		<i>Deodàra</i>	- - -	2285 2429	
<i>T. baccèata</i>	at Fountains Abbey	} 1983 2069	<i>Araucària imbricàta</i>	- - -	2293 2435	
<i>Buckland</i>	- 1984		2074	<i>excélsa</i>	- - -	2302 2443
<i>Harlington</i>	- 1985		2075	<i>Cunninghàmii</i>	- - -	2305 2445
<i>Mambhilad</i>	- - -		1987 2077			
<i>Gresford</i>	- - -		1988 2078			
<i>Fortingal</i>	- 1989		2079			
<i>Westfelton</i>	- - -		1990 2083			
<i>Studley</i>	- - -		1991 2093			
<i>Salisbùria adiantifolia</i>	in the		} 1994 2099			
<i>Leyden Botanic Garden</i>	- - -					
<i>Conífera, § Abiétina.</i>						
<i>Pinus sylvéstris</i>	at Studley	- 2050	2163			
	at Dunmore	{ 2051	2163			
		{ 2056	2185			
	Group of, in	{ 2052	2164			
	Strathspey	{ 2055	2185			
(s.) <i>pumílio Mâghus</i>	- - -	2061	2188			
<i>pumílio</i>	- - -	2063	2190			
<i>Banksiàna</i>	- - -	2067	2191			
<i>inops</i>	- - -	2071	2194			
<i>púngens</i>	- - -	2080	2199			

LIST OF LANDSCAPES

INTRODUCED IN THE TEXT IN ORDER TO SHOW THE EFFECT OF PARTICULAR SPECIES OF TREES IN SCENERY.

Ericàcææ.

	Figure	Page
An <i>Ericacætum</i> to show the effect of a Garden of American <i>Ericàcææ</i> in a bottom, surrounded by hilly and woody Scenery } <i>Euphorbiàcææ.</i> <i>Báucus sempervirens.</i> Part of a Parterre of embroidery in Dwarf Box - - - - - }	1006	1185
Château de Richelieu, to show the effect of embroidered Parterres in Box - - - - - }	1217	1338
	1218	1338

Salicàcææ.

<i>Sàlix babylónica.</i> Villa of Consequa at Canton - - - - -	1302	1508
Churchyard in Baden - - - - -	1303	1509
Vale of Tombs in China - - - - -	1304	1510
Views in the Park at Monza, in Lombardy - - - - -	1305. to 1307	1512, 1513
<i>Pópulus fastigiàta.</i> Bridge to show the contrast between its horizontal lines and the perpendicular lines of the Lombardy Poplars - - - - - }	1521	1663
Stable Offices, &c., with Poplars, to show the contrast between the horizontal lines of the Buildings and the vertical ones of the Trees - - - - - }	1522	1664
Groups of round-headed Trees broken by Poplars - - - - -	1523, 1524, 1524a	1664, 1665
Waterfall in the Gardens of Schwезingen, in Baden, with Lombardy Poplars - - - - - }	1525	1665
Ettlinger Thor Carlsruhe - - - - -	1526	1666
Tivoli Gardens, Vienna - - - - -	1527	1666
Château de Neuville, near Nancy - - - - -	1528	1667
Landscape by Domenichino, showing Lombardy Poplars supporting the effect of a Tower - - - - - }	1529	1667
Poplars in the Cemetery at Père la Chaise - - - - -	1530, 1531	1668, 1669
Entrance to the Botanic Garden at Munich - - - - -	1532	1670
Poplars with a modern Villa - - - - -	2277	2421

Betulàcææ.

<i>Bétulus.</i> Landscape near the Lake Petrovskoyi, near Moscow, composed principally of Birch Woods - - - - - }	1551	1694
-------------------------------------------------------------------------------------------------------------------	------	------

Corylàcææ.

<i>Quercus pedunculàta</i> and <i>sessiliflora.</i> Groups of Oak Trees by Strutt - - - - - }	1629. to 1631	1791
Scenes in Savernake Forest, by Strutt - - - - -	1632, 1633	1792
Old Oak and Rock, by Strutt - - - - -	1634	1793
Oaks torn up by a Whirlwind - - - - -	1644	1814
Wistman's Wood - - - - -	1694	1839
Q. <i>Plex.</i> St. Michael's Mount, Cornwall - - - - -	1782	1903

Coníferææ.

<i>Abiétinæ.</i> Views in Pine Forests in America - - - - -	2004, 2005	2117, 2118
Pine Forest on fire - - - - -	2011	2138
<i>Abies excelsa</i> on the Pass of Kroglevin, in Norway - - - - -	2219	2301
Other Views in Norway, showing the effect of the Spruce Fir alone, and combined with other Trees - - - - - }	2220. to 2222	2302, 2303
<i>Làrix europæa.</i> Views in the Tyrol, showing the effect of Larches in Mountain Scenery - - - - - }	2263, 2264	2357, 2358
<i>Cèdrus Libàni.</i> Cedars on Mount Lebanon - - - - -	2274	2411
Landscapes showing the effect of Cedars in composition with Architectural Scenery - - - - - }	2275, 2276	2420, 2421
Grove of Cedars - - - - -	2278	2422
Cedars in Landscapes, contrasted with other Trees - - - - -	2279. to 2282	2422. to 2425

LIST OF ENGRAVINGS

OF BOTANICAL SPECIMENS OF TREES AND SHRUBS GIVEN
ALONG WITH THE TEXT.

<i>Ranunculàcææ.</i>		Figure	Page		Figure	Page
<i>Clématis Flammula</i>	- -	9	234	<i>Bér. dúlcis</i>	- -	51 305
<i>orientális</i>	- -	10	234	<i>heterophýlla</i>	- -	52 306
<i>gláucia</i>	- -	11	235	<i>dealbàta</i>	- -	53 307
<i>Vitálba</i>	- -	12	236	<i>aristàta</i>	- -	54 307
<i>virginiàna</i>	- -	13	237	<i>buxifòlia</i>	- -	55 308
<i>Vjórna</i>	- -	14	238	<i>Mahònia fasciculàris</i>	- -	56 309
<i>cylíndrica</i>	- -	15	239	<i>Aquifòlium</i>	- -	57 310
<i>Símsü</i>	- -	16	240	<i>nervosa</i>	- -	58 310
<i>reticulàta</i>	- -	17	240	<i>rèpens</i>	- -	59 311
<i>flórída</i>	- -	18	241	<i>Nandína doméstica</i>	Suppl. 2410	2537
<i>cerùlea</i>	- - Suppl.	2407	2525	<i>Cruciàcææ.</i>		
<i>Viticélla</i>	- -	19	242	<i>Vélla Pseudò-Cýtisis</i>	- -	60 312
<i>baleárica</i>	- - Suppl.	2408	2535	<i>Alýssum saxátile</i>	- -	61 313
<i>campaniflòra</i>	- -	20	242	<i>Ibèris sempervirens</i>	- -	62 313
<i>críspa</i>	- -	21	243	<i>Capparidàcææ.</i>		
<i>cirrhòsa</i>	- -	22	244	<i>Cápparis spinòsa</i>	- -	63 314
<i>montàna</i>	- -	23, 24	245	<i>Cistàcææ.</i>		
<i>Atrágene alpína</i>	- -	25	247	<i>Cístus purpùreus</i>	- -	64 318
<i>sibíríca</i>	- -	26	248	<i>heterophýllus</i>	- -	65 318
<i>americàna</i>	- -	27	248	<i>cretícus</i>	- -	66 320
<i>Pæoniá Moítan papaveràcæa</i>	- -	28	250	<i>incànus</i>	- -	67 320
<i>Báncsü</i>	- -	29	250	<i>Heliánthemum scabròsum</i>	- -	68 331
<i>Xanthorhiza apiifòlia</i>	- -	31	255	<i>glomeràtum</i>	- -	69 333
<i>Winteràcææ.</i>				<i>brasilièncè</i>	- -	70 334
<i>Illicium floridànum</i>	- -	32	257	<i>lignòsum</i>	- -	71 335
<i>Magnoliàcææ.</i>				<i>Fumàna</i>	- -	72 336
<i>Magnòlia conspícua</i>	- -	34	279	<i>grandiflòrum</i>	- -	73 344
<i>purpùrea</i>	- -	36	282	<i>maeránthum múltiplex</i>	- -	74 348
<i>Dilleniàcææ.</i>				<i>hyssopifòlium múltiplex</i>	- -	75 352
<i>Hibbértia dentàta</i>	- -	37	292	<i>Hudsonia ericòides</i>	- -	76 354
<i>volùbilis</i>	- -	38	292	<i>Polygalàcææ.</i>		
<i>Anonàcææ</i>				<i>Polýgala Chamæbáxus</i>	- -	77 356
<i>Asímína tríloba</i>	- -	39	294	<i>oppositifòlia màjor</i>	- -	78 356
<i>pygmæà</i>	- -	40	294	<i>Pittosporàcææ.</i>		
<i>Schizandràcææ.</i>				<i>Billardièra longiflòra</i>	- -	79 357
<i>Schizàndra cocéinea</i>	- -	41	295	<i>mutábilis</i>	- -	80 357
<i>Menispermàcææ.</i>				<i>Sóllya heterophýlla</i>	- -	81 357
<i>Menispèrmum canadèncè</i>	- -	42	296	<i>Pittosporum Tobìra</i>	- -	82 858
<i>däuricum</i>	- -	43	297	<i>undulàtum</i>	- -	83 358
<i>smilácinum</i>	- -	44	297	<i>Caryophyllàcææ.</i>		
<i>Cócculus carolinus</i>	- -	45	298	<i>Diánthus arbúscula</i>	- -	84 359
<i>Berberàcææ.</i>				<i>Drýpis spinòsa</i>	- -	85 359
<i>Bérberis vulgàris</i>	- - Suppl.	2409	2536	<i>Linàcææ.</i>		
<i>sibíríca</i>	- -	46	301	<i>Linum arbòreum</i>	- -	86 360
<i>canadèncis</i>	- -	48	303	<i>Malvàcææ.</i>		
<i>emarginàta</i>	- -	49	304	<i>Lavàtera marítima</i>	- -	87 361
<i>ibérica</i>	- -	50	304	<i>Hibíscus syriacus</i>	- -	88 362
				<i>Sida pulchèlla</i>	- -	89 363
				<i>Málva Munroàna</i>	- - Suppl.	2411 2538

	Figure	Page		Figure	Page
<i>Sterculiàceæ.</i>			<i>Acer</i>		
<i>Sterculia</i>	-	90	<i>circinatum</i>	-	127
<i>platanifolia</i>	-	363	<i>palmatum</i>	-	128
<i>Tiliàceæ.</i>			<i>eriocarpum</i>	-	129
<i>Grèwia</i>	Suppl. 2412	2541	<i>rubrum</i>	-	130
<i>Ternstroemiàceæ.</i>			<i>monsperulanum</i>	-	131
<i>Malachodendron</i>	-	91	<i>campèstre</i>	-	132
<i>ovatum</i>	-	377	<i>ereticum</i>	-	132
<i>Stuàrtia</i>	-	92	<i>Æsculàceæ.</i>		
<i>virginica</i>	-	378	<i>Æsculus</i>		
<i>Gordonia</i>	-	93	<i>glabra</i>	-	133
<i>lasianthos</i>	-	380	<i>pallida</i>	-	134
<i>pubescens</i>	-	94	<i>Pavia</i>		
<i>Camèllia</i>	-	95	<i>rubra arguta</i>	-	135
<i>japonica</i>	-	382	<i>neglecta</i>	-	136
<i>anemoniflora</i>	-	96	<i>macrostachya</i>	-	137
<i>reticulata</i>	Suppl. 2413	2541	<i>Meliàceæ.</i>		
<i>maliflora</i>	-	97	<i>Melia</i>		
<i>Sasànqua</i>	-	98	<i>Azederach</i>	-	138
<i>Kissi</i>	-	99	<i>Vitàceæ.</i>		
<i>oleifera</i>	-	100	<i>Vitis</i>		
<i>euryoides</i>	-	101	<i>vinifera</i>	-	139
<i>Thèa</i>	-	102	<i>apiifolia</i>	-	140
<i>viridis</i>	-	393	<i>Labrusca</i>	-	141
<i>Bohèa</i>	-	103	<i>æstivalis</i>	-	142
<i>Aurantiàceæ.</i>			<i>cordifolia</i>	-	143
<i>Seville Orange</i>	-	104	<i>riparia</i>	-	144
<i>Hypericàceæ.</i>			<i>indica</i>	-	145
<i>Hypericum</i>			<i>Ampelopsis</i>		
<i>hircinum</i>	-	105	<i>hederacea</i>	-	146
<i>olympicum</i>	-	106	<i>Cissus</i>		
<i>canariense</i>	Suppl. 2414	2541	<i>antarctica</i>	-	147
<i>Uratum</i>	-	107	<i>ritiginea</i>	-	148
<i>prolificum</i>	Suppl. 2415	2541	<i>quinquefolia</i>	-	149
<i>ægyptiacum</i>	Suppl. 2416	2541	<i>Geraniàceæ.</i>		
<i>empetrifolium</i>	-	108	<i>Pelargonium</i>		
<i>Androsæmum</i>	-	109	<i>cucullatum</i>	-	150
<i>Aceràceæ.</i>			<i>zonale</i>	-	151
<i>Acer</i>			<i>inquinans</i>	-	152
<i>oblongum</i>	-	-	<i>Zygophyllàceæ.</i>		
<i>circinatum</i>	-	112	<i>Meliánthus</i>		
<i>Leaves of Accers.</i>			<i>major</i>	-	153
<i>Acer</i>			<i>Zygophyllum</i>		
<i>oblongum</i>	{	113	<i>sessilifolium</i>	-	154
	Suppl. 2417, 2418	2542	<i>Rutàceæ.</i>		
<i>tataricum</i>	-	114	<i>Ruta</i>		
<i>spicatum</i>	-	115	<i>graveolens</i>	-	155
<i>striatum</i>	-	116	<i>angustifolia</i>	-	156
			<i>Aplophyllum</i>		
			<i>limifolium</i>	-	157
			<i>Corræa</i>		
			<i>alba</i>	Suppl. 2419	2544
			<i>Cròwea</i>		
			<i>saligna</i>	Suppl. 2420	2544
			<i>Boronia</i>		
			<i>serrulata</i>	Suppl. {2421	2544
				2422}	
<i>macrophyllum</i>	-	117, 118	<i>Xanthoxylàceæ.</i>		
			<i>Xanthoxylon</i>		
			<i>fraxineum</i>	-	158
<i>platanoides</i>	-	119	<i>Ailantus</i>		
			<i>glandulosa</i>	-	159
<i>Lobèlii</i>	-	120	<i>Coriàceæ.</i>		
<i>platanoides</i>	-	121	<i>Coriaria</i>		
			<i>myrtifolia</i>	-	160
<i>saccharinum</i>	-	122	<i>Staphylèacèæ.</i>		
<i>Pseudo-Plátanus</i>	-	123	<i>Staphylèa</i>		
			<i>trifolia</i>	-	161, 162
			<i>pinnata</i>	-	163
<i>obtusatum</i>	-	124	<i>Celastràceæ.</i>		
			<i>Euonymus</i>		
<i>barbatum</i>	(opulifolium)	125	<i>europæus</i>	-	164
<i>Opalus</i>	-	126	<i>verrucosus</i>	-	165

	Figure	Page		Figure	Page
<i>Eu. latifolius</i>	-	- 166	498	<i>Condàlia microphýlla</i>	- - 216 541
<i>atropurpureus</i>	-	- 167	499	<i>Collètia spinòsa</i>	- - 217 541
<i>americànus</i>	-	- 168, 169	500	<i>hòrrida</i>	Suppl. 2432c 2548
<i>caucàsius</i>	-	Suppl. 2423	2545	<i>alicina</i>	Suppl. 2432a 2548
<i>japònicus</i>	-	Suppl. 2425	2545	<i>Pomadèrris elliptica</i>	- - 218 542
<i>echinàtus</i>	-	- 170	501	<i>Phýlica cricòides</i>	- - 219 542
<i>Celàstrus scàndens</i>	-	- 171	502	<i>Retanilla obeordàta</i>	Suppl. 2432b 2548
<i>Nemopánthes canadènsis</i>	{	172	503		
	Suppl. 2424	2545		<i>Homalinàceæ.</i>	
<i>Màytenuus chilènsis</i>	-	- 173	503	<i>Azàra dentàta</i>	- - 220 544
<i>Cassìne capènsis</i>	-	- 174	504		
				<i>Anacardiàceæ.</i>	
<i>Aquifoliàceæ.</i>				<i>Pistàcia vèra</i>	- - 221 546
<i>Mygínda myrtifolia</i>	-	- 175	505	<i>Lentiseus</i>	- - 222 548
<i>Ilex Aquifolium marginàtum</i>	-	- 176	507	<i>Rhús Còtinus</i>	- - 223 549
<i>laurifolium</i>	-	- 177	507	<i>typhina</i>	- - 224 550
<i>erassifolium</i>	-	- 178	507	<i>glàbra</i>	- - 225 551
<i>ciliàtum</i>	-	- 179	507	<i>venenàta</i>	- - 226 553
<i>fèrox</i>	-	- 180	507	<i>Coriària</i>	- - 227, 228 554
<i>recúrvm</i>	-	- 181	507	<i>copállina</i>	- - 229 554
<i>serratifolium</i>	-	- 182	507	<i>radicans</i>	- - 230 555
<i>balàrica</i>	{	183	516	<i>Toxicodèndron</i>	- - 231 556
	Suppl. 2426	2546		<i>Duvaína depèndens</i>	- - 232 559
<i>Cassìne</i>	-	- 184	517	<i>ovàta</i>	Suppl. 2433 2549
<i>angustifolia</i>	-	- 185	518	<i>latifolia</i>	- - 233 559
<i>vomitòria</i>	-	- 186	518	<i>Schìnus Millii</i>	- - 234 560
<i>maderènsis</i>	-	- 187	520		
<i>chinènsis</i>	-	- 188	520	<i>Burseràceæ.</i>	
<i>paraguariènsis</i>	-	- 189	520	<i>Balsamodèndron gileadense</i>	235, 236 561
<i>Prinos ambíguus</i>	-	- 190	521		
<i>verticillàtus</i>	-	- 191	521	<i>Leguminòsæ.</i>	
<i>lævigàtus</i>	-	- 192	522	<i>Piptánthus nepalènsis</i>	- 237, 238 566
<i>glàber</i>	-	Suppl. 2428	2546	<i>Edwàrdsia chilènsis</i>	- - 239 567
<i>coriàceus</i>	-	Suppl. 2427	2546	<i>grandiflora</i>	- - 240 567
				<i>microphýlla</i>	- - 241 567
<i>Rhamnàceæ.</i>				<i>Podalýria sericea</i>	- - 242 567
<i>Zizyphus vulgàris</i>	-	- 193	525	<i>Choròzema Henchmànnii</i>	- - 243 568
<i>Lòtus</i>	-	- 194	526	<i>Podolòbium trilobàtum</i>	- 244, 245 568
<i>Jijuba</i>	-	Suppl. 2429	2546	<i>Oxylòbium arborèscens</i>	- 246 568
<i>Paliùrus aculeàtus</i>	-	- 195	527	<i>Brachysèma latifolium</i>	- - 247 568
<i>virgàtus</i>	-	Suppl. 2430	2547	<i>Callistachys ovàta</i>	- - 248 568
<i>Berehènta volùbilis</i>	-	- 196	528	<i>Viminària demudàta</i>	- - 249, 250 568
<i>Rhàmnus Alatèrnus</i>	-	- 197	530	<i>Sphærolòbium vimineum</i>	- 251, 252 569
<i>eathàrticus</i>	-	- 198	532	<i>Aòtus ericòides</i>	- - 253 569
<i>tinetòrius</i>	-	- 199	532	<i>Eutàxia myrtifolia</i>	- - 254, 255 569
<i>infectorius</i>	-	- 200	533	<i>Dillwýnia glabèrrima</i>	- - 256 569
<i>saxàtilis</i>	-	- 201	533	<i>Euchilus obeordàtus</i>	- 257, 258 569
<i>oleòides</i>	-	- 202	534	<i>Pultenæa strìcta</i>	- - 259 570
<i>buxifolius</i>	-	- 203	534	<i>Mirbènta reticulàta</i>	- - 260 570
<i>Erythròxylon</i>	-	- 204	534	<i>Davièsia latifolia</i>	- - 261 570
<i>angustissimum</i>	-	- 205	535	<i>Anagýris fœ'tida</i>	- Suppl. 2435 2549
<i>alnifolius</i>	-	- 206	536	<i>Baptisia tinetòria</i>	- Suppl. 2434 2549
<i>frangulòides</i>	-	- 207	536	<i>Ulex europæa</i>	- - 262, 263 571
<i>alpìnus</i>	-	- 208	536	<i>nàna</i>	- - 264, 265 575
<i>Fràngula</i>	-	- 209	537	<i>Spàrtium júnceum</i>	- - 266 576
<i>latifolius</i>	-	- 210	538	<i>odoratissimum</i>	Suppl. 2436 2550
<i>Purshianus</i>	-	- 211	538	<i>Genista càndicans</i>	- - 267 578
<i>carpinifolius</i>	-	- 212	538	<i>triquetra</i>	- - 268 578
<i>Ceanòthus azureus</i>	-	- 213	539	<i>radiàta</i>	- - 269 579
<i>americànus</i>	-	- 214	539	<i>ànglica</i>	- - 270 581
<i>velutinus</i>	-	- 215	540	<i>germànica</i>	- - 271 581
<i>collinus</i>	-	Suppl. 2431	2548	<i>monospërma</i>	- - 272 582

	Figure	Page		Figure	Page
<i>G. aethnensis</i>	-	- 273	<i>Anthyllis Bárba Jovis</i>	-	- 337 641
<i>auxántica</i>	-	- 274, 275	<i>Medicàgo arborea</i>	-	- 338 642
<i>tinetòria</i>	-	- 276	<i>Lòtus jacobæus</i>	-	- 339 642
<i>sagittàlis</i>	-	- 277	<i>Psoràlea glandulòsa</i>	-	- 340, 341 642
<i>prostràta</i>	-	- 278	<i>Indigòfera denudata</i>	-	- 342 643
<i>pilòsa</i>	-	- 279	<i>austràlis</i>	-	- 343 643
<i>linitòlia</i>	-	- 280	<i>Swainsonia galegifolia</i>	-	- 344 643
<i>canariensis</i>	Suppl.	2437 2551	<i>Lessertia frutescens</i>	-	- 345 643
<i>virgata</i>	-	- 281	<i>Sutherlandia frutescens</i>	-	- 346 643
<i>Cytisus álbus</i>	-	- 282	<i>Coronilla E'merus</i>	-	- 347 644
<i>nigricans</i>	-	- 283	<i>júncea</i>	-	- 348 644
<i>sessilifolius</i>	-	- 284, 285	<i>stipularis</i>	-	- 349 645
<i>triflorus</i>	-	- 286	<i>gláuca</i>	-	- 350 645
<i>scoparius</i>	-	- 287	<i>Hippocrèpis baleàrica</i>	-	- 351, 352 645
<i>leucanthus</i>	-	- 288	<i>Adesmia microphylla</i>	-	- 353, 354 645
<i>purpureus</i>	-	- 289, 290	<i>Loudonia</i>	-	- 355, 356 646
<i>austriacus</i>	-	- 291	<i>viscosa</i>	-	- 357 646
<i>supinus</i>	-	- 292	<i>Desmodium polycarpum</i>	Suppl.	2442 2553
<i>oleucus</i>	-	- 293	<i>nütans</i>	Suppl.	2443 2553
<i>nanus</i>	-	- 294	<i>Clíanthus puniceus</i>	-	- 358 646
<i>proliferus</i>	-	- 295	<i>Wistaria frutescens</i>	-	- 359 647
<i>Adenocarpus intermedius</i>	-	- 296	<i>chinensis</i>	-	- 360 648
<i>parvifolius</i>	-	- 297	<i>Lupinus arboreus</i>	-	- 361 649
<i>telonensis</i>	-	- 298	<i>Marshallianus</i>	-	- 362 649
<i>Ononis frutescens</i>	-	- 299	<i>Kennedyia ovata</i>	Suppl.	2444 2554
<i>rotundifolia</i>	-	- 300	<i>Mucuna macrocarpa</i>	-	- 363 649
<i>Natrix</i>	-	- 301	<i>Gleditschia monosperma</i>	-	- 364 653
<i>peduncularis</i>	-	- 302	<i>Ceratonia Siliqua</i>	-	- 365, 366 660
<i>Amorpha frutescens</i>	-	- 303	<i>Cassia Barclayana</i>	-	- 367 660
<i>fragrans</i>	-	- 304	<i>australis</i>	-	- 368 661
<i>erocœo-lanata</i>	-	- 305	<i>Schötia latifolia</i>	-	- 369 661
<i>Robinia Pseud-Acacia</i>	-	- 306	<i>Chamaefistula corymbosa</i>	-	- 370 661
<i>viscosa</i>	-	- 307	<i>Acacia alata</i>	-	- 371 662
<i>hispida</i>	-	- 308	<i>armata</i>	-	- 372 662
<i>Caragana Altajana</i>	-	- 309	<i>juniperina</i>	-	- 373 662
<i>arenaria</i>	-	- 310	<i>diffusa</i>	-	- 374, 375 663
<i>frutescens</i>	-	- 311	<i>stricta</i>	-	- 376, 377 663
<i>pygmaea</i>	-	- 312, 313	<i>melanoxylon</i>	-	- 378, 379 663
<i>spinosa</i>	-	- 314	<i>myrtifolia</i>	-	- 380 663
<i>Chandagu</i>	-	- 315	<i>suaveolens</i>	-	- 381 663
<i>Halimodendron argenteum</i>	-	- 316	<i>Oxycedrus</i>	-	- 382, 383 664
<i>Calophaea wolgarica</i>	-	- 317	<i>verticillata</i>	-	- 384 664
<i>Colutea arborescens</i>	{	Suppl. 2440 2552	<i>pulchella</i>	-	- 385, 386 664
<i>eruenta</i>	-	- 318	<i>lophantha</i>	-	- 387 664
<i>nepalensis</i>	-	- 319	<i>farnesiana</i>	-	- 388 665
<i>Astragalus Tragacantha</i>	-	- 320	<i>nigricans</i>	-	- 389, 390 665
<i>ereticus</i>	-	- 321	<i>Julibrissin</i>	-	- 391 665
<i>aristatus</i>	-	- 322	<i>mollissima</i>	-	- 392, 393 666
<i>brevifolius</i>	-	- 323			
<i>Hovea Celsi</i>	-	- 324	<i>Rosaceae.</i>		
<i>latifolia</i>	-	- 325	<i>Amygdalus nana</i>	-	- 394, 395 674
<i>Platylobium formosum</i>	-	- 326	<i>Persica vulgaris</i>	-	- 396 679
<i>Bossia rufa</i>	Suppl.	2441 2552	<i>compressa</i>	-	- 397 680
<i>Goodia latifolia</i>	-	- 327	<i>Armeniaea vulgaris ovalifolia</i>	-	- 398 682
<i>Scotia dentata</i>	-	- 328	<i>cordifolia</i>	-	- 399 682
<i>Templetonia glauca</i>	-	- 329, 330	<i>dasycarpa</i>	-	- 400, 401 683
<i>Rafnia triflora</i>	-	- 321, 332	<i>persicifolia</i>	-	- 402 683
<i>Viborgia obovata</i>	-	- 333	<i>brigantiaea</i>	-	- 403 684
<i>Loddigesia ovalifolia</i>	-	- 334	<i>Prunus candicans</i>	-	- 404, 405 690
<i>Aspalathus callosa</i>	-	- 335	<i>Cerasus serrulata</i>	-	- 406 701
<i>Sarcophyllum carnosum</i>	-	- 336	<i>Pseudo-Cerasus</i>	-	- 407 701
			<i>Chamaecerasus</i>	-	- 408 702

	Figure	Page		Figure	Page
Cér. prostrata	-	409	Potentilla fruticosa	-	468
borealis	-	410	frut. tenuiloba	-	469
nigra	-	411, 412	glabra	-	470
japónica	-	413, 414	Cowanìa plicata	Suppl.	2455
múltiplex	-	415, 416	Rosa ferox	-	471
sinénsis	-	417	kamtschática	-	472
virginiàna	-	418	bracteata	-	473
serótina	-	419	microphýlla	-	474
Capóllim	-	420	involverata	-	475
lusitànica	-	421	lúcida	-	476
Hixa	Suppl.	2455	nítida	-	477
Laurocérasus	-	422	Ràpa	-	478
caroliniàna	-	423	parviflora flore pleno	-	479
Púrshia tridentata	-	424, 425	fraxinifolia	-	480
Kérria japónica	{	426	cinnamomea	-	481
Spiræa opulifolia	Suppl.	2446	alpina	-	482
chamadriifolia	-	427, 428	laevis	-	483
mèdia	Suppl.	2449	suavis	-	484
ulmifolia	-	430	lutescens	-	485
phyllantha	Suppl.	2448	sulphurea	-	486
cratægifolia	Suppl.	2450	spinosissima	-	487
betulaefolia	Suppl.	2447	grandiflora	-	488
trilobata	-	431	myriacantha	-	489
alpina	-	432	damascena	{	490
hypericifolia	-	433	centifolia	Suppl.	2456
acuta	-	434	muscosa	-	491
crenata	-	435	gállica	{	493
savrànica	-	436	parvifolia	Suppl.	2457
Besseriàna	Suppl.	2451	turbinata	-	494
thalictroides	-	437	álba	{	495
ceanothifolia	Suppl.	2452	lutea	Suppl.	2458
corymbosa	-	438	punicea	-	497
racciniifolia	-	439	rubiginosa	-	498
bèlla	-	440	caucæsa	-	499
salicifolia	-	443	canina	Suppl.	2459
alpèstris	Suppl.	2453	aciphýlla	-	501, 502
latifolia	-	441	rubrifolia	-	503
grandiflora	-	442	índica	-	504
tomentosa	-	444	Noissettàna	-	505
laevigata	-	445	odoratissima	-	506
ariæfolia	-	446, 447	longifolia	-	507
sorbifolia	-	448	semperflorens	-	508
Rùbus micranthus	-	449, 450, 450a	Lawrenceàna	-	509
occidentalis	-	451	arvensis	-	510
idæus	{	452	perpervirens	-	511
laciniatus	Suppl.	2554	multiflora	-	512
caesius	-	453	Grevillei	-	513
parvifolius	-	454	moschata	-	514
corylifolius glandulosus	-	455	nivea	-	515
corylifolius	-	456	rubifolia	-	516
spectabilis	-	457	sínica	{	517
fruticòsus	-	458	Banksia	Suppl.	2460
fruticòsus	-	459	lutea	-	518
pompònius	-	460	microcarpa	-	519
tomentosus	-	461	macrophýlla	-	520
hispidus	-	462	sericea	-	521
odoratus	-	463	moschata nepalensis	-	522
nutkanus	-	464	hýstrix	-	523
macropetalus	-	465			524
reflexus	-	466			
coronarius	-	467			

	Figure	Page		Figure	Page
<i>Lòwea berberifòlia</i>	- - -	549 813	<i>Cr. parvifòlia</i>	- - -	614 867
<i>Cratægus glandulòsa subvillòsa</i>	- - -	550 818	flòrida	- - -	613 867
lobàta	- - -	554 824	grossulariaefòlia	- - -	615 867
<i>Oxyacantha sibírica</i>	- - -	555 830	virginica	- - -	616 867
parvifòlia	- - -	557 842	mexicana	- - -	617 867
flòrida	- - -	558 842	<i>Photinia arbutifòlia</i>	- - -	619 869
grossulariaefòlia	- - -	559 842	<i>Cotoneaster vulgaris</i>	- - -	620 870
virginica	- - -	560 843	laxiflòra	- - -	621, 622 871
<i>Pyracantha</i>	- - -	561 844	rotundifòlia	- - -	623, 624 872
glàucia (<i>Stranvèsia</i>)	562, 563	845	microphýlla	- - -	625 873
<i>Leaves of Cratægus.</i>			<i>Amelanchier vulgaris</i>	- - -	626 874
coccinea	- - -	564 851	<i>Botryopium</i>	- 627, 628, 629	{ 874 875
corállina	- - -	565 852	sanguinea	- - -	630, 631 875
indentàta	- - -	566 852	ovàlis	- - -	632 876
glandulòsa	- - -	567 853	flòrida	- - -	633, 634 876
subvillòsa	- - -	568 853	<i>Pyrus erenàta</i>	- - -	638, 639 890
punctàta rùbra	- - -	569 854	intermèdia angustifòlia	- - -	640 912
aúrea	- - -	570 854	rivularis	- - -	642, 643 915
brevispina	- Suppl.	2461 2562	spùria	- - -	645 925
myrtifòlia	- - -	571 854	arbutifòlia	- - -	646 926
macracantha	- - -	572 855	pùmila	- - -	647, 648 926
minor	- - -	573 855	melanocarpa	- - -	649 927
<i>Crús galli</i>	- - -	574 855	grandifolia	- {	650 928
splendens	- - -	575 856	=	{ Suppl. 2465	2566
prunifòlia	- - -	576 856	<i>Chamaemèspilus</i>	- - -	651 928
lineàris	- - -	577 856	<i>Cydònia japònica</i>	- - -	652 931
salicifòlia	- - -	578 856	<i>Raphiòlepis indica</i>	- - -	653 932
ovalifòlia	- - -	579 856	salicifòlia	- - -	654 933
pyracanthifòlia	- - -	580 856	<i>Eriobòtrya japònica</i>	- - -	655, 656 933
nigra	- - -	581 857	<i>Kagenèkia crategoides</i>	{ Suppl.	657 934 2466 2566
purpùrea	- - -	582 857	<i>Potèrium caudatum</i>	- - -	658 934
altàica	- - -	583 858	<i>Calycanthàceæ.</i>		
Douglasii	- - -	584 858	<i>Calycanthus flòridus</i>	- - -	659 936
flàva	- - -	585 859	glàucus	- - -	660 937
lobàta	- - -	586 859	laevigatus	- - -	661 937
trilobàta	- - -	587 860	<i>Chimonánthus fràgrans</i>	- - -	662 938
apiifòlia minor	- - -	588 860	fràgrans grandiflòrus	- - -	663 938
apiifòlia	- - -	589 860	<i>Granatàceæ.</i>		
cordàta	- - -	590 861	<i>Pùnica Granatum rùbrum</i>	- - -	664 940
spathulàta	- - -	591 861	nàna	- - -	665 942
<i>Azaròtus</i>	- - -	592 862	<i>Onagràceæ.</i>		
<i>Arònia</i>	- - -	593 862	<i>Fúchsia microphýlla</i>	- - -	666 944
maroceàna	- - -	594 862	coccinea	- - -	667 944
orientàlis	- - -	595 863	excorticàta	- - -	668 945
sanguinea	- - -	596 863	<i>Lythràceæ.</i>		
tanaetifòlia	- - -	597 863	<i>Heimìa salicifòlia</i>	- - -	669 945
glàbra	- - -	598 863	<i>Lagerstrømìa indica</i>	- - -	670 945
Leeana	- - -	599 864	<i>Tamaricàceæ.</i>		
heterophýlla	- - -	600 864	<i>Tamarix gállica</i>	- - -	671 947
<i>Oxyacantha</i>	- - -	602 864	<i>Myricària germanica</i>	- - -	672 949
obtusàta	- - -	601 865	<i>Philadelphàceæ.</i>		
laciniatà	- - -	603 865	<i>Philadelphus coronàrius</i>	- - -	673 951
pteridifòlia	- - -	604 865	inodòrus	- - -	674 952
melanocarpa	- - -	605 865			
<i>Oliveriana</i>	- - -	606 865			
erioearpa	- - -	607 865			
quercifòlia	- - -	608 866			
múltiplex	- - -	609 866			
aúrea	- - -	610 866			
purpùrea	- - -	611 866			
ròsea	- - -	612 866			

	Figure	Page		Figure	Page
<i>Ph. verrucòsus</i>	-	675	953	<i>Ribes alp. pùmilum</i>	- 726 979
<i>latifòlius</i>	-	676	953	<i>petræum</i>	- 727 979
<i>lâxus</i>	-	677	953	<i>spicatum</i>	- 728 980
<i>hirsùtus</i>	-	678, 678a	954	<i>multiflorum</i>	- 729 980
<i>Decumària bárbara</i>	-	679, 680	955	<i>procumbens</i>	- 730 981
<i>Deutzia scàbra</i>	-	681	956	<i>prostratum</i>	- 731 981
				<i>resinosum</i>	- 732 981
Myrtàcæ.				<i>punctatum</i>	- { 733 982
<i>Tristània nereifòlia</i>	-	682	957		Suppl. 2475 2569
<i>Beaufòrtia decussàta</i>	-	683	957	<i>nigrum</i>	- 734 983
<i>Calothámnus villòsa</i>	-	684	957	<i>flòridum</i>	- 735 985
<i>Endémia tetragòna</i>	-	686	958	<i>inèbrians</i>	- 736 986
<i>Melaleuca hypericifòlia</i>	-	687	958	<i>cèream</i>	- 737 986
<i>Eucalýptus resinifera</i>	-	688, 689	959	<i>viscosíssimum</i>	- 738 987
<i>viminàlis</i>	-	692, 693	960	<i>sanguineum</i>	- 739 988
<i>amygdàlina</i>	-	694, 695	960	<i>glutinòsum</i>	- 740 988
<i>piperità</i>	-	696	960	<i>malvaceum</i>	- 741 988
<i>pulverulénta</i>	-	697	960	<i>aúream</i>	- 742 989
<i>cordàta</i>	-	698	960	<i>seròtinum</i>	- 743 989
<i>Angòphora cordifòlia</i>	-	699	960	<i>tenuiflorum</i>	- 744 990
<i>Callistèmon lanceolàtus</i>	-	700	960	<i>flàvum</i>	- Suppl. 2476 2569
<i>Leptospernum grandifòlium</i>	-	701	961		
<i>scopàrium</i>	-	Suppl. 2468	2568	Escalloniàcæ.	
<i>Fabrcia myrtifòlia</i>	-	703	961	<i>I' tea virginica</i>	- 745 992
<i>Bæ'kia virgàta</i>	-	704	961	<i>Escallònia rubra</i>	- 746 993
<i>Metrosidèros corifòlius</i>	-	705	961	<i>montevidènsis</i>	- 747 993
<i>Psidium Cattleyanum</i>	-	706	962	<i>illinita</i>	- Suppl. 2477 2570
<i>Mýrtus communis</i>	Suppl. 2469, 2470	2568	2568		
<i>tomèntosa</i>	-	707	964	Saxifragæ.	
<i>Cálythrix glàbra</i>	-	708	964	<i>Hydránga arborèscens</i>	- 748 995
				<i>cordàta</i>	- 749 995
Passifloràcæ.				<i>nivea</i>	- 750 995
<i>Passiflora carúlea</i>	-	709	965	<i>quercifolia</i>	- 751 996
<i>incarnàta</i>	-	710	965	<i>Horténsia</i>	- 752 996
<i>Disémma adiantifòlia</i>	-	711	965		
				Umbellàcæ.	
Crassulàcæ.				<i>Bupleurum fruticòsum</i>	{ 753 998
<i>Sèdum populifòlium</i>	Suppl. 2471	2568	2568		Suppl. 2478 2570
<i>Sempervivum arbòreum</i>	-	712	965		
				Araliàcæ.	
Nitrariàcæ.				<i>Aràlia spinòsa</i>	- 754 999
<i>Nitrària Schòberi sibirica</i>	-	713	966	<i>Hèdera Hèlix vulgàris</i>	- 755 1000
<i>Schòb. cáspica</i>	-	714	967		
				Hamamelàcæ.	
Cactàcæ.				<i>Hamamèlis virginica</i>	- 756 1007
<i>Opúntia vulgàris</i>	-	Suppl. 2474	2569	<i>Fothergilla alnifòlia màjor</i>	- 758 1009
				<i>alnif. obtùsa</i>	- 759 1009
Reamuriàcæ.				Cornàcæ.	
<i>Reamurià hypericòides</i>	Sup. { 2472 } 2569	2569	2569	<i>Còrnus alternifòlia</i>	- 760 1010
	{ 2473 }			<i>sanguinea</i>	- 761 1011
Grossulàcæ.				<i>álba</i>	- 762 1011
<i>Ribes oxyacanthiòides</i>	-	715	969	<i>stricta</i>	- 763, 764 1012
<i>setosum</i>	-	716	969	<i>paniculàta</i>	- 765 1013
<i>triflorum</i>	-	717	969	<i>sericea</i>	- 766 1013
<i>niveum</i>	-	718	970	<i>circinàta</i>	- 767 1014
<i>Cynòsbati</i>	-	719	970	<i>más</i>	- 768 1015
<i>divaricatum</i>	-	720	971	<i>flòrida</i>	- 769 1017
<i>irriguum</i>	-	721	971	<i>Benthàmia fragifera</i>	- 770 1020
<i>speciosum</i>	-	722	975		
<i>Diacántha</i>	-	723	976	Loranthàcæ.	
<i>lacústre</i>	-	724	976	<i>Viscum álbum</i>	- 771 1021
<i>alpinum</i>	-	725	979	<i>Arceuthòbium Oxýcedri</i>	- 2137 2246
				<i>Aicuba japònica</i>	- Suppl. 2479 2571

	Figure	Page		Figure	Page
<i>Caprifoliàcea.</i>			<i>Art. argentea</i> - - - 839 1069		
<i>Sambucus nigra</i> - - -	773	1028	<i>Helicrhysum fruticans</i> - - -	840	1070
<i>nig. laciniata</i> - - -	774	1028	<i>congestum</i> - - -	841	1070
<i>follis argenteis</i> - - -	775	1028	<i>Cineraria eruenta</i> - - -	842	1071
<i>canadensis</i> - - -	776	1030	<i>lanata</i> - - -	843	1071
<i>racemosa</i> - - -	777	1031	<i>geifolia</i> - - -	844	1071
<i>Viburnum Tinus</i> - - -	778	1032	<i>anelloides</i> - - -	845, 846	1071
<i>rugosum</i> - - -	779	1033	<i>Othonna virginea</i> - - -	847	1072
<i>Lentago</i> - - -	780	1034	<i>Osteospermum pisiferum</i> 848. to 850		1072
<i>pyrifolium</i> - - -	781, 782	1034	<i>Calendula chrysanthemifolia</i> - - -	851	1072
<i>nudum</i> - - -	783	1035	<i>Mutisia latifolia</i> - - -	853	1072
<i>squamatum</i> - - -	784	1035	<i>Sonchus fruticosus</i> - - -	853	1073
<i>Lantana</i> - - -	785	1036	<i>Aster argophyllus</i> - - -	854	1073
<i>lantanoïdes</i> - - -	786	1037	<i>aculeatus</i> - - -	855	1073
<i>cotinifolium</i> - - -	787, 788	1037	<i>Chrysocoma Comaurea</i> - - -	856	1073
<i>dentatum</i> - - -	789, 790	1038	<i>Podanthus Mitiqui</i> - - -	857	1074
<i>odoratissimum</i> - - -	791	1039	<i>Culeitium salicinum</i> - - -	858	1074
<i>O'pulus</i> - - -	792	1039	<i>Pyræthrum feniculaceum</i> {	859	1074
<i>acerifolium</i> - - -	793	1040	{ <i>Sup.</i> 2485	2573	
<i>Diervilla canadensis</i> - - -	794, 795	1042	<i>Eriocephalus africanus</i> Suppl.	2487	2574
<i>Lonicera Periclymenum serotinum</i> - - -	797	1044	<i>A'nthemis porrigenis</i> - Suppl.	2486	2573
<i>Caprifolium</i> - - -	798	1046	<i>Epacridàcea.</i>		
<i>etrusca</i> - - -	799	1046	<i>Stenanthera pinifolia</i> Suppl.	2458	2574
<i>impléxa</i> - - -	800	1046	<i>Leucopogon albiliflorus</i> - - -	860	1075
<i>flava</i> - - -	801	1047	<i>Trochocarpa laurina</i> - - -	861	1075
<i>pubescens</i> - - -	802	1047	<i>Epacris grandiflora</i> - - -	862	1075
<i>parviflora</i> {	803, 804	1048	<i>Andersonia sprengeloides</i> - - -	863	1075
{ <i>Suppl.</i> 2480, 2481	2572		<i>Ericàcea.</i>		
<i>grata</i> - - -	805	1048	<i>Erica Tétralix</i> - - -	864	1079
<i>sempervirens</i> - - -	806	1049	<i>cinerea</i> - - -	865	1080
<i>majior</i> - - -	807	1049	<i>codonodes</i> - - -	866	1081
<i>confusa</i> - - -	808	1050	<i>australis</i> - - -	867	1081
<i>longiflora</i> - - -	Suppl. 2482	2572	<i>stricta</i> - - -	868	1081
<i>japonica</i> - - -	809, 810	1051	<i>ciliaris</i> - - -	869	1082
<i>tatarica</i> - - -	811, 812	1053	<i>Gypsocallis vagans</i> - - -	870	1082
<i>nigra campaniflora</i> - - -	813, 814	1053	<i>multiflora</i> - - -	871	1083
<i>punicea</i> - - -	815	1054	<i>carnea</i> - - -	872	1083
<i>Xylosteum</i> - - -	816	1055	<i>Callista acuminata</i> - - -	873	1089
<i>involutrata</i> - - -	817, to 819	1056	<i>Cerámia serpyllifolia</i> - - -	874	1090
<i>alpigena</i> - - -	820, 821	1056	<i>urecolaris</i> - - -	881	1092
<i>oblongifolia</i> - - -	822	1057	<i>Désmia conferta</i> - - -	875	1090
<i>curvula</i> - - -	823, 824	1057	<i>Erica aggregata</i> - - -	876	1090
<i>hispidula</i> - - -	Suppl. 2483	2572	<i>Gypsocallis intertexta</i> - - -	877	1090
<i>Symphoricarpos vulgaris</i> - - -	825	1059	<i>Pachysa physodes</i> - - -	878, 879	1091
<i>racemosus</i> - - -	826	1059	<i>Blaëria ericoides</i> - - -	880	1091
<i>Leycesteria formosa</i> - - -	827	1060	<i>Eurylepis albens</i> - - -	882	1093
<i>Rubiàcea.</i>			<i>Eurystegia triiceps</i> - - -	883	1093
<i>Cephalanthus occidentalis</i> 828, 829	1061		<i>Lophandra cubica</i> - - -	884	1093
<i>Lucilia gratissima</i> - Suppl.	2484	2573	<i>Lamprotis calycina</i> - - -	885	1093
<i>Pinckneya pubens</i> - - -	830	1062	<i>Andróméda polifolia</i> - - -	889	1106
<i>Sérissa fe'tida</i> - - -	831	1062	<i>pol. angustifolia</i>	890	1106
<i>Compositæ.</i>			<i>grandiflora</i>	891	1106
<i>Stachelia dubia</i> - - -	832	1064	<i>latifolia</i> - - -	892	1106
<i>Baccharis halimifolia</i> - - -	833	1065	<i>minima</i> - - -	893	1106
<i>Fva frutescens</i> - - -	834	1066	<i>Cassiope hypnoides</i> - - -	894	1107
<i>Santolina Chamæcyparissus</i> - - -	835	1067	<i>tetragona</i> - - -	895	1107
<i>rosmarinifolia</i> - - -	836	1067	<i>Cassindra calyculata</i> - - -	896	1108
<i>Artemisia Abrótanum</i> - - -	837	1068	<i>angustifolia</i> - - -	897	1108
<i>Santónica</i> - - -	838	1069	<i>Zenobia speciosa</i> - - -	898	1109
			<i>speciosa nitida</i> - - -	899	1109

	Figure	Page		Figure	Page
<i>Z. speciosa pulverulenta</i>	- -	900 1109	<i>Rh. indicum</i>	- -	955 1148
<i>Lyonia marginata</i>	{	902 1110	<i>ledifolium</i>	- -	956 1149
<i>marg. rubra</i>		Suppl. 2489 2575	<i>sinense</i>	- -	957 1149
<i>mariana</i>	{	901 1110	<i>Farrera</i>	- -	958 1150
<i>paniculata</i>		903 1110	<i>Kalmia latifolia</i>	- -	959 1151
<i>salicifolia</i>	Suppl. 2490 2575	<i>angustifolia</i>	- -	960 1152	
<i>racemosa</i>	- -	904 1111	<i>glauca</i>	- -	961 1152
<i>multiflora</i>	- -	905 1112	<i>hirsuta</i>	- -	962 1153
<i>capreaefolia</i>	Suppl. 2491 2575	- -	<i>Menziesia ferruginea</i>	- -	963 1153
<i>Leucothoe axillaris</i>	- -	906 1112	<i>globularis</i>	Suppl. 2494 2577	
<i>spinulosa</i>	- -	907 1112	<i>Azalea procumbens</i>	{	964 1154
<i>acuminata</i>	- -	908 1113	<i>Suppl. 2500 2577</i>		
<i>floribunda</i>	- -	910 1114	<i>Leiophyllum thymifolium</i>	- -	965 1154
<i>spicata</i>	- -	911 1114	<i>Ledum palustre</i>	- -	966 1155
<i>Pieris ovalifolia</i>	- -	912 1114	<i>latifolium</i>	- -	967 1155
<i>japonica</i>	- -	913 1115	<i>canadense</i>	- -	968 1156
<i>Phyllodoce taxifolia</i>	- -	914 1115	<i>Vaccinium Myrtillus</i>	- -	969 1157
<i>empetriformis</i>	- -	915 1115	<i>uliginosum</i>	- -	970 1158
<i>Daboecia polifolia</i>	- -	916 1116	<i>tenellum</i>	- -	971 1159
<i>Arbutus Unedo</i>	- -	917, 918 1116	<i>stamineum</i>	- -	972 1160
<i>hybrida</i>	- -	919 1117	<i>dumosum</i>	- -	973 1160
<i>Andrachne</i>	{	920 1117	<i>corymbosum</i>	- -	974, 975 1161
<i>canariensis</i>		Suppl. 2492, 2493 2575	<i>virgatum</i>	- -	976 1161
<i>Arctostaphylos Uva-ursi</i>	- -	921 1120	<i>fuscatum</i>	- -	977 1161
<i>Pernettia mucronata</i>	- -	922 1122	<i>marianum</i>	- -	978 1162
<i>Gaultheria procumbens</i>	- -	923 1123	<i>grandiflorum</i>	{	979 1162
<i>Shallon</i>	- -	924 1124	<i>Suppl. 2501 2577</i>		
<i>Clèthra alnifolia</i>	- -	925 1125	<i>elongatum</i>	- -	980 1162
<i>tomentosa</i>	- -	926 1126	<i>minutiflorum</i>	- -	981 1162
<i>arborea</i>	- -	927 1128	<i>glabrum</i>	- -	982 1162
<i>Rhododendron ponticum</i>	- -	928, 929 1128	<i>resinosum</i>	- -	983 1163
<i>maximum</i>	- -	930 1129	<i>Arctostaphylos</i>	- -	984 1164
<i>hybridum</i>	- -	931 1131	<i>padifolium</i>	{	985, 986 1164
<i>catawbiense</i>	Suppl. 2494 2576	<i>Suppl. 2502 2577</i>	<i>Suppl. 2502 2577</i>		
<i>caucasicum</i>	- -	932 1154	<i>Tritis idæa</i>	- -	987 1165
<i>punctatum</i>	- -	933 1135	<i>buxifolium</i>	- -	988, 989 1165
<i>ferrugineum</i>	- -	934 1136	<i>nitidum</i>	- -	990 1166
<i>hirsutum</i>	- -	935 1137	<i>ovatum</i>	- -	991 1166
<i>lapponicum</i>	- -	936 1137	<i>Oxycoccus palustris</i>	- -	992 1168
<i>dauricum</i>	- -	937 1137	<i>macrocarpus</i>	- -	993 1170
<i>camtschaticum</i>	- -	938 1138	<i>erectus</i>	- -	994 1171
<i>Chamaecistus</i>	- -	939 1138	<i>Bryanthus Gmelini</i>	- -	995 1171
<i>flavum</i>	- -	940 1139	<i>Stelleri</i>	- -	996 1172
<i>nudiflorum</i>	- -	941 1139	<i>Enkianthus quinqueflorus</i>	997, 998	1173
<i>Govenidium</i>	- -	942 1140	<i>Cyrilla racemosa</i>	Suppl. 2503	2577
<i>rossa</i>	- -	943 1140			
<i>calendulaceum</i>	- -	944 1141			
<i>viscosum</i>	- -	945 1142			
<i>hispidum</i>	- -	946 1142			
<i>Cartonia</i>	- -	947 1143			
<i>nitidum</i>	- -	948 1144			
<i>speciosum</i>	- -	949 1144			
<i>Rhodora</i>	- -	950 1145			
<i>arboresum</i>	- -	951 1146			
<i>altaclerense</i>	- -	952 1147			
<i>campanulatum</i>	- -	953 1148			
<i>anthopogon</i>	- -	954 1148			

*Symplocaceæ.**Symplocos sinica* - - 1007 1186*Styracææ.*

Styrax officinale - - 1008 1187
grandifolium - - 1009 1188
laevigatum - - 1010 1189
pulverulentum - - 1011 1189

Halesiææ.

Halesia tetraptera - - 1012 1190
parviflora - - 1013 1191
diptera - - 1014 1191

Myrsinææ.

Myrsine africana - Suppl. 2504 2578
Manglilla Milleriana - Suppl. 2505 2578

	Figure	Page		Figure	Page
Sapotàceæ.			<i>Vinea</i> minor -	- 1084	1256
<i>Argània</i> Sideróxylo -	- 1015	1192	<i>Gelsèmium</i> nítidum -	- 1085	1256
<i>Bumèlia</i> lyeiöides	{ Suppl. 2506	2578	<i>Nèrium</i> Oleánder -	- 1086	1256
tènax -	- 1017	1193	Asclepiadàceæ.		
salicifölia -	Suppl. 2507	2578	<i>Períploca</i> græ'a -	- 1087, 1088	1257
Oleàceæ.			angustifölia -	- 1089	1258
<i>Ligústrum</i> vulgàre -	1019, 1020	1199	Bignoniàceæ.		
vul. sempervirens -	- 1018	1199	<i>Bignònia</i> capreolàta -	- 1090	1259
spicàtum -	- 1020	1201	<i>Tècoma</i> radicans -	- 1091	1260
licidum -	Suppl. 2508	2579	grandiflöra -	- 1092	1260
floribúndum -	- 1024	1202	austrális -	- 1093	1261
<i>Phillyrea</i> angustifölia -	- 1025	1204	capénsis -	- 1094	1261
ang. rosmarinifölia -	- 1026	1204	<i>Eccremocárpus</i> longiflörus -	- 1095	1263
mèdia -	- 1027	1204	<i>Calámpelis</i> scàbra -	- 1096, 1097	1263
latifölia -	- 1028	1205	Cobæàceæ.		
<i>Chionánthus</i> virgínea -	- 1029	1206	<i>Cobæ'a</i> scándens -	- 1098	1264
<i>O'lea</i> europæ'a -	- 1031	1207	Convolvulàceæ.		
eur. satíva longifölia -	- 1032	1207	<i>Convólulus</i> Dorýenium -	- 1100	1264
ferrugínea -	- 1033	1208	<i>Cneörum</i> -	- 1099	1264
americàna -	- 1034	1208	Boraginàceæ.		
frágrans -	- 1035	1208	<i>Lithospérnum</i> fruticòsum -	- 1101	1265
capénsis -	Suppl. 2509	2579	<i>E'chium</i> cándicans -	- 1102	1265
<i>Notelá'a</i> ligústrina -	Suppl. 2510	2579	Cordiàceæ.		
<i>Syrínga</i> vulgàris -	- 1036	1210	<i>Ehrètia</i> serràta -	- 1103	1265
Josikæ'a -	- 1037	1210	Solanàceæ.		
pérsica -	- 1039	1211	<i>Solànum</i> Dulcamàra -	- 1104	1266
laciniàta -	- 1040	1211	erispium -	- 1105	1268
rothomagénsis -	- 1041	1212	bonariénse -	- 1106	1268
<i>Emòdi</i> -	- 1042	1212	Balbísii -	Suppl. 2514	2582
<i>Fontanèsia</i> phillyreöides -	- 1043	1213	ligústrinum -	- 1107	1269
<i>Fráxinus</i> excélsior -	- 1044	1215	<i>Lýcium</i> europæ'um -	- 1108	1270
heterophýlla -	- 1050	1228	bárbarum -	- 1109	1270
variegàta -	- 1051	1229	chinénse -	- 1110, 1111	1271
parviflöra -	- 1052	1230	ruthénicum -	- 1112	1272
oxyeárpa -	- 1053	1230	lanceolàtum -	Suppl. 2513, 2516	2582
lentiscifölia -	- 1054	1231	turbinàtum -	{ Suppl. 2517	2582
americàna -	- 1055	1232	àfrum -	- 1114, 1115	1273
pubéscens -	- 1056	1234	<i>Grabòwskia</i> boerhaaviæfölia -	- 1116	1274
sambucifölia -	- 1057, 1058	1235	<i>Nicotiàna</i> gláuca -	- 1118	1274
quadrangulàta -	- 1059, 1060	1236	<i>Brugnànsia</i> sanguínea -	- 1117	1275
juglandifölia -	1061, 1062	{ 1236	snavèolens -	- 1120	1275
platyeárpa -	- 1063, 1064	1238	<i>Céstrum</i> nocturnum -	- 1119	1275
angustifölia -	Suppl. 2511	2580	<i>Párqui</i> -	- 1122	1275
lentiscifölia -	Suppl. 2512	2581	<i>Véstia</i> lyeiöides -	- 1121	1275
<i>O'rnus</i> europæ'a -	- 1065	1242	Scrophulariàceæ.		
rotundifölia -	- 1069	1244	<i>Búddlea</i> globösa -	- 1123, 1124	1276
americàna -	- 1070	1244	<i>Hallèria</i> lúcida -	- 1125	1277
floribúnda -	- 1072	1245	<i>Cóisia</i> lanàta -	- 1126	1277
Jasminàceæ.			<i>Calceolària</i> integrifölia -	- 1127, 1128	1277
<i>Jasminum</i> fruticans -	- 1073	1248			
húmile -	- 1074	1249			
heterophýllum -	- 1075	1249			
revolutum -	- 1076	1250			
pubígerum -	- 1077	1250			
officinále -	- 1078	1251			
Apocynàceæ.					
<i>Vinea</i> májor -	- 1082, 1083	1255			

	Figure	Page		Figure	Page
<i>Verónica</i> decussata	- 1129, 1130	1278	<i>Cinnamònum</i> <i>Cámphora</i>	- 1174	1305
<i>Penstèmon</i> <i>Scouleri</i>	Suppl. 2518	2582	<i>verum</i>	- 1175	1305
<i>Labiàceæ.</i>			<i>Proteàceæ.</i>		
<i>Saturèja</i> montana	- 1131	1278	<i>Grevillea</i> <i>rosmarinifolia</i>	- 1176	1306
<i>Thymus</i> <i>vulgaris</i>	- 1132	1278	<i>acuminata</i>	- 1177, 1178	1306
<i>grandiflorus</i>	- Suppl. 2515	2582	<i>Hàkea</i> <i>pugionifòrmis</i>	- 1179	1306
<i>Hyssòpus</i> <i>officinàlis</i>	- 1133	1278	<i>Bánksia</i> <i>latifolia</i>	- Suppl. 2522	2584
<i>Teuèrium</i> <i>fruticans</i>	- 1135, 1136	1279	<i>Thymelàceæ.</i>		
<i>Màrum</i>	- 1134	1279	<i>Dáphne</i> <i>Mezereum</i>	- 1180	1308
<i>Phlòmis</i> <i>fruticosa</i>	- 1137	1279	<i>altàica</i>	- 1181	1309
<i>purpùrea</i>	- 1138	1279	<i>alpina</i>	- 1182	1309
<i>Rosmarinus</i> <i>officinàlis</i>	- 1139	1280	<i>Laurèola</i>	- 1183	1309
<i>Lavándula</i> <i>Spica</i>	- 1140	1281	<i>póntica</i>	- 1184	1310
<i>Sálvia</i> <i>officinàlis</i>	- 1141	1282	<i>Thymelà'ea</i>	- 1185	1311
<i>var.</i>	- 1142	1282	<i>Tárton-raira</i>	- 1186	1311
<i>Hablitziana</i>	- 1143	1283	<i>collina</i>	- 1187	1312
<i>pomifera</i>	- 1144	1283	<i>neapolitana</i>	- 1188	1312
<i>Audibértia</i> <i>incana</i>	- 1145	1283	<i>oleoides</i>	- 1189	1312
<i>Lavándula</i> <i>dentata</i>	- 1146	1283	<i>Gnídium</i>	- 1190	1313
<i>pinnata</i>	- 1147	1283	<i>Cneòrum</i>	- 1191	1313
<i>Stæ'chas</i>	- 1149	1284	<i>odora rubra</i>	- 1192	1314
<i>Plectrànthus</i> <i>fruticòsus</i>	- 1148	1284	<i>hýbrida</i>	- 1193	1314
<i>Sphàcele</i> <i>campanulata</i>	- 1150	1284	<i>Dírca</i> <i>palustris</i>	- 1194	1315
<i>Sálvia</i> <i>Gràhami</i>	- 1151	1284	<i>Santalàceæ.</i>		
<i>Prostanthèra</i> <i>lasianthos</i>	Suppl. 2519	2583	<i>Nýssa</i> <i>biflora</i>	- 1195, 1196	1317
<i>Verbenàceæ.</i>			<i>villòsa</i>	- 1197, 1198	1318
<i>Fítex</i> <i>A'gnus</i> <i>cástus</i>	- 1152	1285	<i>cándicans</i>	- 1199	1318
<i>Clerodéndron</i> <i>inèrme</i>	- 1153	1286	<i>grandidentata</i>	- 1200, 1201	1319
<i>Aloýsia</i> <i>citriodòra</i>	- 1154	1286	<i>Osýris</i> <i>álba</i>	- 1202	1320
<i>Globulariàceæ.</i>			<i>Elæagnàceæ.</i>		
<i>Globulària</i> <i>longifolia</i>	- 1155	1287	<i>Elæágnus</i> <i>horténsis</i>	- 1203	1322
<i>Plumbaginàceæ.</i>			<i>argénteá</i>	- 1204	1323
<i>Plumbàgo</i> <i>capénsis</i>	- Suppl. 2520	2583	<i>salicifolia</i>	- 1205	1324
<i>Chenopodiàceæ.</i>			<i>Hippóphàe</i> <i>rhamnoides</i>	- 1206	1325
<i>Chenopòdium</i> <i>fruticòsum</i>	1156, 1157	1289	<i>salicifolia</i>	- 1207	1326
<i>A'triplex</i> <i>Hálinus</i>	- 1158	1290	<i>Shephérdia</i> <i>argénteá</i>	- 1208	1327
<i>portulacòides</i>	- 1159	1290	<i>canadénsis</i>	- 1209	1328
<i>Diòtis</i> <i>Ceratòides</i>	- 1160	1291	<i>Aristolochiàceæ.</i>		
<i>Polygonàceæ.</i>			<i>Aristolòchia</i> <i>sìpho</i>	- 1210	1329
<i>Tragopýrum</i> <i>lanceolatum</i>	- 1161	1293	<i>tomentosa</i>	- 1211	1329
<i>buxifolium</i>	- 1162	1293	<i>Euphorbiàceæ.</i>		
<i>polygamum</i>	- 1163	1294	<i>Euphórbia</i> <i>Charàcias</i>	- 1212	1331
<i>Atraphàxis</i> <i>spinosa</i>	- 1164	1294	<i>spinosa</i>	- 1213	1331
<i>Calligonum</i> <i>Pallásii</i>	- 1165, 1166	1295	<i>mellifera</i>	- 1214	1332
<i>Rùmex</i> <i>sp.</i>	- 1167	1295	<i>Búxus</i> <i>sempervirens nana</i>	- 1215	1333
<i>Lauràceæ.</i>			<i>baleàrica</i>	- 1220, 1221	1341
<i>Laúrus</i> <i>Borbònia</i>	- 1168	1299	<i>Plagiánthus</i> <i>divaricatus</i>	Suppl. 2524	2585
<i>carolinénsis</i>	- 1169	1300	<i>Cròton</i> <i>rosmarinifolia</i>	Suppl. 2523	2585
<i>aggregata</i>	- 1170	1301	<i>Adèlia</i> <i>acidòton</i>	- Suppl. 2525	2585
<i>Benzòin</i>	- 1171	1303	<i>Urticàceæ.</i>		
<i>Diospýrus</i>	- 1172	1304	<i>Mòrus</i> <i>álba multicaúlis</i>	- 1223	1348
<i>geniculata</i>	- 1173	1304	<i>tatàrica</i>	- 1225	1358
			<i>Maclùra</i> <i>aurantiaca</i>	- 1226, to 1228	{ 1363 1364
			<i>Bòrya</i> <i>acuminata</i>	- 1229	1371

	Figure	Page		Figure	Page
<i>Ulmàcæ.</i>			<i>Sal. acuminàta</i>	- - - 1330	1554
<i>Ulmus campestris parvifolia</i>	- 1230	1377	<i>Pontederàna</i>	- - - 1331	1556
<i>cam. clinicensis</i>	- - - 1231	1377	<i>cinerea</i>	- - - 1332	1558
<i>suberosa</i>	- - - 1240	1395	<i>càprea</i>	- 1333. to 1335	1562
<i>màjor</i>	- - - 1241	1396	<i>cotinifolia</i>	- - - 1336	1566
<i>carpinifolia</i>	- - - 1242	1397	<i>rotundàta</i>	- - - 1337	1572
<i>glabra</i>	- - - 1245	1403	<i>làurina</i>	- - - 1338	1578
<i>americàna</i>	- - - 1246	1406	<i>Borreriàna</i>	- - - 1339	1579
<i>fùlva</i>	- - - 1247	1408	<i>Weigeliàna</i>	- - - 1340	1582
<i>alàta</i>	- - - 1248	1409	<i>nitens</i>	- - - 1341	1582
<i>Plànera Richàrdi</i>	- - - 1440	1410	<i>racciniifolia</i>	- - - 1342	1585
<i>Gmelini</i>	- - - 1251	1413	<i>myrtilloides</i>	- - - 1343	1587
<i>Celtis australis</i>	- - - 1252	1414	<i>Myrsinites</i>	- - - 1344	1588
<i>crassifolia</i>	- - - 1254	1419	<i>betulifolia</i>	- - - 1345	1588
<i>orientalis</i>	- - - 1255	1420	<i>retusa</i>	- - - 1346	1589
<i>Juglandàcæ.</i>			<i>serpyllifolia</i>	- 1347, 1348	1590
<i>Juglans règia</i>	- - - 1257	1425	<i>herbacea</i>	- - - 1249	1591
<i>nigra</i>	- - - 1260	1436	<i>polaris</i>	- - - 1350, 1351	1592
<i>cinerea</i>	- - - 1262	1439	<i>hastata</i>	- - - 1352	1592
<i>Carya olivæformis</i>	- - - 1263	1442	<i>arbuscula</i>	- - - 1353	1593
<i>amara</i>	- - - 1264	1443	<i>lanata</i>	- - - 1354	1594
<i>aquatica</i>	- - 1265, 1266	1444	<i>coluteoides</i>	- Suppl. 2529	2588
<i>tomentosa</i>	- - - 1267	1445	<i>berberifolia</i>	- - - 1355	1595
<i>alba</i>	- - - 1269	1446	<i>Leaves of Salix.</i>		
<i>sulcata</i>	- - - 1271	1449	<i>purpurea</i>	- - - 1	} 1603
<i>porcina</i>	- - 1272. to 1274	1449	<i>Helix</i>	- - - 2	
<i>myristicæformis</i>	- - - 1275	1451	<i>Lambertiàna</i>	- - - 3	
<i>Pterocarya caucásica</i>	- - - 1276	1452	<i>monàndra</i>	- - - 4	
<i>Salicàcæ.</i>			<i>Forbyàna</i>	- - - 5	} 1604
<i>Salix purpurea</i>	- - - 1294	1490	<i>rubra</i>	- - - 6	
<i>daphnoides</i>	- - - 1295	1495	<i>virescens</i>	- - - 7	
<i>undulata</i>	- - - 1296	1497	<i>Humboldtiana</i>	- - - 8	
<i>triàndra</i>	- - - 1297	1499	<i>Bonplandiàna</i>	- - - 9	} 1605
<i>amygdalina</i>	- - - 1298	1501	<i>Lyoni</i>	- - - 10	
<i>pentàndra</i>	- - - 1299	1503	<i>Houstoniana</i>	- - - 11	
<i>Meyeriàna</i>	- - - 1300	1505	<i>virgata</i>	- - - 12	
<i>lucida</i>	- - - 1301	1506	<i>undulata</i>	- - - 13	} 1606
<i>decipiens</i>	- - - 1309	1515	<i>lanecolata</i>	- - - 14	
<i>fragilis</i>	- - - 1310	1516	<i>triàndra</i>	- - - 15	
<i>Russelliàna</i>	- - - 1311	1518	<i>Hoffmanniana</i>	- - - 16	
<i>alba</i>	- - 1314, 1315	1523	<i>Villarsiana</i>	- - - 17	} 1606
<i>nigra</i>	} - Suppl. 2527	1524	<i>amygdalina</i>	- - - 18	
<i>ligustrina</i>		2588	<i>montana</i>	- - - 19	
<i>prinoides</i>	- - - 1317	1530	<i>vitellina</i>	- - - 20	
<i>virescens</i>	- - - 1318	1531	<i>annularis</i>	- - - 21	} 1607
<i>petiolaris</i>	- - - 1319	1533	<i>babylonica</i>	- - - 22	
<i>rosmarinifolia</i>	- - - 1320	1535	<i>petiolaris</i>	- - - 23	
<i>angustifolia</i>	- - - 1321	1536	<i>conformis</i>	- - - 24	
<i>Doniana</i>	- - - 1322	1540	<i>violacea</i>	- - - 25	} 1608
<i>versicolor</i>	- Suppl. 2528	2588	<i>præcox</i>	- - - 26	
<i>reticulata</i>	- - - 1323	1543	<i>fragilis</i>	- - - 27	
<i>glauca</i>	- - - 1324	1544	<i>Russelliana</i>	- - - 28	
<i>Lapponium</i>	- - - 1325	1545	<i>decipiens</i>	- - - 29	} 1609
<i>candida</i>	- - - 1326	1548	<i>monspeiciensis</i>	- - - 30	
<i>incana</i>	- - - 1327	1548	<i>tetrasperma</i>	- - - 31	
<i>linearis</i>	- - - 1328	1549	<i>lucida</i>	- - - 32	
<i>viminialis</i>	- - - 1329	1549	<i>Meyeriàna</i>	- - - 33	} 1610
			<i>pentandra</i>	- - - 34	
			<i>hastata</i>	- - - 35	
			<i>malifolia</i>	- - - 36	
			<i>nigricans</i>	- - - 37	} 1611

	Figure	Page		Figure	Page
Sål. bicolor	38	1612	Sål. strépida	100	1621
påtens	39		sórdida	101	
prinöides	40		rivulàris	102	
Willdenoviàna	41	1613	austràlis	103	1622
crísipa	42		rotundàta	104	
Pontederàna	43		dùra	105	
nítens	44		fírma	106	
Borrieriàna	45	1614	Ansoniàna	107	1623
phylcifòlia	46		atrovirens	108	
Davalliàna	47		Andersoniàna	109	
Wulfeniàna	48		Forsteriàna	110	
tétrapla	49	1615	rupéstris	111	1624
tenuifòlia	50		coriàcea	112	
Weigeliàna	51		hírta	113	
Crowcàna	52		cotinfòlia	114	
ramifúsa	53	1616	crassifòlia	115	1625
floribúnda	54		lacústris	116	
Dicksoniàna	55		vandénsis	117	
prunifòlia	56		latifòlia	118	
vacciniifòlia	57	1617	grisophýlla	119	1626
venulòsa	58		incanésceus	120	
carinàta	59		sphacelàta	121	
Myrsinítes	60		càprea	122	
procúmbens	61	1618	pannòsa	123	1627
herbàcea	62		aurita	124	
polàris	63		cinèrea	125	
Kitaibelàna	64		oleifòlia	126	
serpyllifòlia	65	1619	aquática	127	1628
myrtillöides	66		ferrugínea	128	
reticulàta	67		germinàta	129	
gláuca	68		macrostipulàcea	130	
elæagnifòlia	69	1620	acuminàta	131	1629
arenària	70		stipulàris	132	
lanàta	71		viminàlis	133	
Stuartiàna	72		Smithiàna	134	
Lappònum	73	1621	Micheliàna	135	1630
seríceea	74		álba	136	
proteafolia	75		cærùlea	137	
alaternöides	76		arbúscula	138	
versicolor	77	1622	retùsa	139	1631
argénteaa	78		berberifòlia	140	
incubàcea	79		rígida	141	
adscéndens	80		cordàta	142	
parvifòlia	81	1623	cordifòlia	143	1632
prostràta	82		obovàta	144	
fúsa	88		Muhlenbergiàna	145	
rèpens	84		ægyptiaca	146	
Doniàna	85	1624	díscolor	147	1633
arbúscula	86		falcàta	148	
rosmarinifòlia	87		alpína	149	
decúmbens	88		trístis	150	
lineàris	89	1625	U'va-úrsi	151	1634
incàna	90		nigra	152	
cándida	91		Pópulus álba	1507	
villòsa	92		canésceus	1508	
subalpína	93	1626	trémula	1509	1635
refléxa	94		trépida	1510	
penusylvànica	95		grandidentàta	1511	
pállida	96		græ'ca	1512	
petræ'a	97	1627	nigra	1513	1636
Schleicheriàna	98		canadénsis	1515	
grisonénsis	99		betulæfòlia	1516	

	Figure	Page
<i>Póp. monilifera</i>	- - 1517	1657
<i>fastigiata</i>	- - 1520	1661
<i>angulata</i>	- - 1533	1671
<i>heterophýlla</i>	- - 1534	1672
<i>balsamifera</i>	{ 1535, 1536	1674
	{ Suppl. 2530	2589
<i>candicans</i>	- - 1537	1676

Betulàceæ.

<i>Alnus glutinosa</i>	- - 1540	1677
<i>laciniata</i>	- - 1538	1678
<i>oxyacanthæfolia</i>	- - 1539	1679
<i>incana</i>	- - 1543	1687
<i>serrulata</i>	- - 1544	1688
<i>cordifolia</i>	- - 1545	1689
<i>viridis</i>	- - 1546	1690
<i>acuminata</i>	- Suppl. 2531	2589
<i>castaneæfolia</i>	- Suppl. 2532	2589
<i>Bétula álba</i>	- - 1547	1691
<i>pubescens</i>	- - 1548	1691
<i>póntica</i>	- - 1549	1692
<i>dãurica</i>	- - 1556	1704
<i>fruticosa</i>	- - 1557	1705
<i>pàmila</i>	- - 1558	1706
<i>nana</i>	- - 1559	1706
<i>populifolia</i>	- - 1560	1707
<i>papyracea</i>	- - 1561	1709
<i>nigra</i>	- - 1562, 1563	{ 1710 1711
<i>excelsa</i>	- - 1564, 1565	1712
<i>lenta</i>	- - 1566	1713

Corylàceæ.

<i>Quercus pedunculata</i>	- - 1567	1731
<i>ped. heterophýlla</i>	1569. to 1571	{ 1733 1734
<i>sessiliflora</i>	- - 1572	1736
<i>pubescens</i>	- - 1573	1737
<i>sessiliflora var.</i>	- 1574. to 1579	{ 1738 1739
<i>pedunculata</i>	- - 1583	1743
<i>sessiliflora</i>	- - 1584	1743
<i>pyrenaica</i>	- - 1696	1843
<i>apennina</i>	- - 1697	1844
<i>E'sculus</i>	- - 1699. to 1701	1845
<i>Cérris and vars.</i>	1702. to 1706	{ 1846 1847
<i>péndula</i>	- - 1707	1848
<i>austriaca</i>	- - 1708	1848
<i>cana majior</i>	- - 1709	1849
<i>fulhamensis</i>	- 1710, 1711	{ 1850 1851
<i>laciniata</i>	- Suppl. 2534	2591
<i>Lucombeana</i>	- 1714	1854
<i>crispa</i>	- 1715, 1717c	1856
<i>dentata</i>	- 1716	1856
<i>suberosa</i>	- 1717a	1857
<i>incisa</i>	- 1717b	1857
<i>heterophýlla</i>	- 1719	1859

	Figure	Page
<i>Qu. A'gilops</i>	- - 1721	1862
<i>Acorns of American Oaks</i>	1722	1863
<i>álba</i>	- - 1723, 1726	{ 1864 1866
<i>repanda</i>	- - 1724	1865
<i>squamosa</i>	- - 1725	1865
<i>oliveforinis</i>	- - 1730	1869
<i>macrocarpa</i>	- - 1731	1870
<i>obtusiloba</i>	- - 1732	1870
<i>lyrata</i>	- - 1733, 1734	1871
<i>Prius palustris</i>	- - 1735	1873
<i>monticola</i>	- - 1736	1874
<i>acuminata</i>	- - 1737	1875
<i>pàmila</i>	- - 1738	1875
<i>tomentosa</i>	- - 1739	1876
<i>rùbra</i>	- - 1740. to 1744	{ 1877 1878
<i>coccinea</i>	- - 1746. to 1748	1880
<i>ambigua</i>	- - 1749	1882
<i>falcata</i>	- - 1750, 1751	1883
<i>tinctoria</i>	- 1753. to 1757	{ 1885 1886 1887
<i>palustris</i>	- 1758. to 1760	{ 1888 1888
<i>Catesbæi</i>	- - 1762, 1763	1889
<i>nigra</i>	- - 1764, 1765	{ 1890 1891
<i>aquatica</i>	- - 1767, 1768	1892
<i>ilicifolia</i>	- - 1770, 1771	{ 1894 1895
<i>Phellos</i>	- - 1774	1896
<i>aurifolia</i>	- - 1776	1897
<i>hýbrida</i>	- - 1775	1897
<i>imbriçaria</i>	- - 1777, 1778	{ 1898 1899
<i>Flex</i>	- - 1781	1900
<i>Ballota</i>	- - 1783. to 1786	{ 1905 1906
<i>gramuntia</i>	- - 1787, 1788	1907
<i>coccifera</i>	- - 1789. to 1792	{ 1908 1909
<i>pseudo-coccifera</i>	- - 1794	1911
<i>Süber</i>	- - 1795. to 1798	{ 1912 1913
<i>Pseudo-Süber</i>	- - 1801	1917
<i>virens</i>	- - 1802, 1803	{ 1918 1919
<i>lanata</i>	- - 1804	1921
<i>annulata</i>	- - 1805	1922
<i>Túrneri</i>	- 1806. to 1809	{ 1923 1924
<i>hýbrida nana</i>	- 1810. to 1812	1924
<i>Fontanèsii</i>	- - 1813	1925
<i>australis</i>	- - 1814	1925
<i>Coókii</i>	- - 1815	1926
<i>faginea</i>	- - 1816	1926
<i>lusitânica</i>	- - 1817	1927
<i>prásina</i>	- - 1818	1928
<i>infectoria</i>	- - 1819, 1820	1928
<i>Libani</i>	- - 1826	1932
<i>castaneæfolia</i>	- - 1827	1933

Oaks not yet introduced.

		Figure	Page	Figure	Page
<i>Himalayan.</i>					
Qu.	<i>spicàta</i>	-	- 1828	1930	2003
	<i>grandifolia</i>	-	- 1829	1931	2004
	<i>velutina</i>	-	- 1830	1932	2004
	<i>lamellosa</i>	-	- 1831	1938	2014
	<i>semicarpifolia</i>	-	- 1832	1939	2015
<i>Javanese.</i>					
	<i>sundàica</i>	-	- 1833, 1834	1940	2016
	<i>pruinosa</i>	-	- 1835	1941	2016
	<i>angustata</i>	-	- 1836	1942	2017
	<i>pallida</i>	-	- 1837, 1838	1943	2018
	<i>élegans</i>	-	- 1839	1948	2029
	<i>placentària</i>	-	- 1840	1949	2029
	<i>glaberrima</i>	-	- 1842	1950	2031
	<i>costata</i>	-	- 1841, 1843	1938	2032
				1939	
	<i>rotundata</i>	-	- 1844	1939	
	<i>daphnoides</i>	-	- 1845	1939	
	<i>platycarpa</i>	-	- 1846	1939	
	<i>gemelliflora</i>	-	- 1847	1940	
	<i>induta</i>	-	- 1848	1940	
	<i>pseudomolucca</i>	-	- 1849	1940	
	<i>turbinata</i>	-	- 1850	1941	
	<i>lineata</i>	-	- 1851	1941	
<i>Mexican.</i>					
	<i>xalapensis</i>	-	- 1852	1941	
	<i>glaucescens</i>	-	- 1853	1941	
	<i>obtusa</i>	-	- 1854	1942	
	<i>pandurata</i>	-	- 1855, 1856	1942	
	<i>repanda</i>	-	- 1857	1943	
	<i>laurina</i>	-	- 1858	1943	
	<i>sideroxyla</i>	-	- 1860	1943	
	<i>mexicana</i>	-	- 1861	1944	
	<i>crassipes</i>	-	- 8862	1944	
	<i>angustifolia</i>	-	- 1859	1943	
	<i>lanceolata</i>	-	- 1863	1944	
	<i>reticulata</i>	-	- 1865	1945	
	<i>chrysophylla</i>	-	- 1864	1945	
	<i>pulchella</i>	-	- 1866	1945	
	<i>spicata</i>	-	- 1867	1945	
	<i>stipularis</i>	-	- 1868	1946	
	<i>crassifolia</i>	-	- 1869	1946	
	<i>ambigua</i>	-	- 1870	1946	
	<i>depressa</i>	-	- 1871	1946	
	<i>confertifolia</i>	-	- 1872	1947	
	<i>tridens</i>	-	- 1873	1947	
	<i>acutifolia</i>	-	- 1874	1947	
<i>Fagus</i>	<i>sylvatica heterophylla</i>		{ 1875 } { 1876 }	1952	
	<i>sylv. cristata</i>	-	- 1877	1952	
	<i>ferruginea</i>	-	- 1917	1981	
	<i>caroliniana</i>	-	- 1915	1980	
	<i>latifolia</i>	-	- 1916	1980	
	<i>obliqua</i>	-	- 1919	1982	
	<i>betuloides</i>	-	- 1920	1982	
	<i>Dombeyi</i>	-	- 1921	1983	
	<i>dubia</i>	-	- 1922	1983	
<i>Castanea</i>	<i>pumila</i>	-	- 1927, 1928	2002	
	<i>martabanica</i>	-	- 1929	2003	
<i>Cas.</i>	<i>Tungurrut</i>	-	-	1930	2003
	<i>argentea</i>	-	-	1931	2004
	<i>javanica</i>	-	-	1932	2004
<i>Carpinus</i>	<i>americana</i>	-	-	1936	2013
	<i>orientalis</i>	-	-	1937	2014
	<i>viminea</i>	-	-	1938	2014
<i>Ostrya</i>	<i>vulgaris</i>	-	-	1939	2015
	<i>virginica</i>	-	-	1940	2016
				{ Suppl. 2537	2595
<i>Corylus</i>	<i>Avellana</i>	-	-	1941	2017
	<i>Av. tubulosa</i>	-	-	1942	2018
	<i>crispa</i>	-	-	1943	2018
	<i>Colurna</i>	-	-	1948	2029
	<i>arborescens</i>	-	-	1949	2029
	<i>ferox</i>	-	-	1950	2031
<i>Garryaceae.</i>					
<i>Garrya</i>	<i>elliptica</i>	-	-	1951	2032
<i>Platanaceae.</i>					
<i>Platanus</i>	<i>orientalis</i>	-	- 1952. to 1954	{ 2033	
	<i>occidentalis</i>	-	- 1959	{ 2035	2043
<i>Balsamaceae.</i>					
<i>Liquidambar</i>	<i>Styraciflua</i>	-	-	1961	2049
	<i>imbérbe</i>	-	-	1963, 1964	2054
	<i>Altingia</i>	-	-	1965	2054
<i>Myricaceae.</i>					
<i>Myrica</i>	<i>Gale</i>	-	-	1966, 1967	{ 2056
					{ 2057
	<i>cerifera latifolia</i>	-	-	1968	2057
	<i>Faya</i>	-	-	1969	2059
	<i>spathulata</i>	-	-	1970	2059
<i>Comptonia</i>	<i>asplenifolia</i>	-	-	1971	2060
<i>Gnetaceae.</i>					
<i>Ephedra</i>	<i>distachya</i>	-	-	1973, 1974	2063
	<i>monostachya</i>	-	-	1975, 1976	2064
	<i>altissima</i>	-	-	1977, 1978, 1979	2064
	<i>americana</i>	-	-	1980	2065
				{ Suppl. 2539	2597
<i>Taxaceae.</i>					
<i>Taxus</i>	<i>baccata fastigiata</i>	-	-	1981	2066
	<i>Salisburia adiantifolia</i>	-	-	1982, 1983	2095
<i>Podocarpus</i>	<i>latifolius</i>	-	-	1995	2100
	<i>chilinus</i>	-	-	1996	2101
	<i>elongatus</i>	-	-	1997	2101
	<i>coriaceus</i>	-	-	1998	2101
	<i>taxifolius</i>	-	-	1999, 2000	2101
	<i>Daerydium cupressinum</i>	-	-	2001	2102
<i>Phyllocladis</i>	<i>rhomboidalis</i>	2002,	2003	2102	
<i>Coniferae.</i>					
<i>Pinus</i>	<i>sylvestris</i>	-	-	{ 2043	2153
				{ to	to
				{ 2046	2155
				{ 2048	2160
	<i>syl. uncinata</i>	-	-	2047	2157
	<i>pumilio</i>	-	-	{ 2057,	2058
				{ 2062	2186
					2189

	Figure	Page		Figure	Page
<i>Pin. pum. Mûghus</i>	- 2059, 2060	2187	<i>Pin. Llavedàna</i>	- 2177. to 2179	{ 2267 2268
<i>Banksiàna</i>	- 2064. to 2066	{ 2190 2191	<i>muricàta</i>	- - 2180	2269
<i>inops</i>	- 2068. to 2070	{ 2192 2193	<i>tuberculàta</i>	- - 2181	2270
<i>mitis</i>	- 2072. to 2076	{ 2195 2196	<i>radiàta</i>	- - 2182	2271
<i>púngens</i>	- 2077. to 2079	{ 2198 2200	<i>occidentàlis</i>	- - 2183	2272
<i>Larício</i>	- 2081. to 2084	{ 2200 to 2202	<i>Montezumæ</i>	- 2184, 2185	{ 2272 2273
<i>austriaca</i>	- - - 2085	{ 2205 2206	<i>leiophýlla</i>	- - 2186	{ 2274 2274
<i>Pallasiàna</i>	- 2086. to 2089	{ to 2208	<i>Cembra</i>	- 2188. to 2191	{ to 2277
<i>pyrenàica</i>	- 2090. to 2093	{ 2209 2210	<i>Stròbus</i>	- - 2193, 2195	{ 2280 2281
<i>resinòsa</i>	- 2094. to 2096	{ 2210 2211	<i>excèlsa</i>	- - 2197, 2200	{ 2286 2287
<i>canadènsis bifòlia</i>	2098, 2099	{ 2213 2214	<i>Lambertiàna</i>	- 2203, 2206	{ 2288 2290
<i>Pináster</i>	- 2100, 2101. 2105	{ 2215 2218	<i>montícola</i>	- - 2208, 2209	{ 2291 2292
<i>Lemoniànus</i>	- 2102, 2103	2216	<i>contórta</i>	- - 2210, 2211	2292
<i>minor</i>	- 2104	2217	<i>Abies excèlsa</i>	- - - 2212	2293
<i>Pínea</i>	- 2106, 2107. 2109	{ 2225 to 2227	<i>excèlsa nigra</i>	- - - 2213	2294
<i>halépènsis</i>	- 2110. to 2113	{ 2231 to 2233	<i>álba</i>	- - - 2224	2311
<i>brútia</i>	- 2114. to 2116	2235	<i>nigra</i>	- - - 2225	2312
<i>sp. from Nepal</i>	- - 2117	2236	<i>rùbra</i>	- - - 2228	2316
<i>Tæ'da</i>	- 2118. to 2121	{ 2237 2238	<i>Smithiàna</i>	- - - 2229	2317
<i>rígida</i>	- 2123. to 2126	{ 2239 2241	<i>Douglàsü</i>	- 2230, 2231	{ 2319 2320
<i>serótina</i>	- 2127. to 2130	{ 2242 2243	<i>Menzièsü</i>	- - - 2232	2321
<i>variàbilis</i>	- { 2131 Suppl. 2540	{ 2242 2599	<i>dumòsa</i>	- - 2233, 2234	2325
<i>ponderòsa-</i>	2132. to 2135	{ 2244 2245	<i>cephalónica</i>	- - 2235, 2236	2326
<i>Arcuthòbium Oxýcedri</i>	- 2137	2246	<i>Picea pectinàta</i>	- 2337, 2338	{ 2330 2331
<i>Pinus Sabiniàna</i>	- 2138. to 2142	{ 2247 to 2249	<i>balsànea</i>	- 2240. to 2242	{ 2339 2340
<i>Coultèri</i>	- 2144. to 2146	{ 2250 2251	<i>Fràseri</i>	- - 2243, 2244	2340
<i>longifolia</i>	- 2148. to 2151	{ 2252 2253	<i>grándis</i>	- - 2245, 2246	2341
<i>Gerardiàna</i>	- 2153 to 2155	{ 2254 2255	<i>amàbilis</i>	- - 2247, 2248	2342
<i>austràlis</i>	- 2156. to 2159	{ 2256 to 2258	<i>nóbilis</i>	- - 2249, 2250	2343
<i>canariènsis</i>	- 2162. to 2165	{ 2261 2263	<i>Webbiàna</i>	- 2251, 2252	{ 2344 2345
<i>sinènsis</i>	- 2167. to 2169	2264	<i>Pindrow</i>	- - 2254, 2255	{ 2346 2347
<i>insignis</i>	- - 2170, 2172	2265	<i>bracteàta</i>	- - 2256	2348
<i>Trocòte</i>	- - 2073, 2074	2266	<i>religiòsa</i>	- { 2257 Suppl. 2543	{ 2349 2602
<i>pátula</i>	- 2175, 2176	2267	<i>Làrix europæa</i>	- - - 2258	2353
			<i>Cèdrus Libàni</i>	- - - 2267	2403
			<i>Deodàra</i>	- 2283, 2284. 2286	{ 2428 to 2430
			<i>Araucària imbricàta</i>	- 2286. to 2292	{ to 2433 2435
			<i>brasiliana</i>	- 2294. to 2296	{ 2439 2440
			<i>excèlsa</i>	- - 2297. to 2301	{ 2441 2442
			<i>Cunninghãmi</i>	{ 2303, 2304 Suppl. 2545	{ 2444 2604
			<i>Cunninghãmia sinènsis</i>	2306, 2307	2446

	Figure	Page		Figure	Page
<i>Dámmara orientális</i>	- 2308, 2309	{ 2447 2448	<i>Jun. lýcia</i>	- - 2367, 2368	{ 2502 2503
<i>austrális</i>	- - 2310, 2311	{ 2448 2449	<i>thurífera</i>	- - - 2369	2503
<i>Thùja occidentális</i>	- - 2312	2455	<i>excélsa</i>	- - - 2370	2504
<i>orientális</i>	- - 2315	2459	<i>reécúrva</i>	- - - 2371	2504
<i>cupressóides</i>	- - 2316	2460	<i>chinénsis</i>	- - 2372, 2373	2505
<i>péndula</i>	- - 2317, 2318	2461	<i>Smíthii</i>	- - - 2374	2505
<i>Cállítris quadriválvis</i>	- - 2319	2463	<i>E'mpetrum nígrum</i>	- 2375, 2376	2507
<i>Cupréssus sempervirens</i>	- 2320	2465	<i>rùbrum</i>	- - 2377, 2378	2508
<i>thyóides</i>	- - 2327	2475	<i>Ceratiola ericóides</i>	- 2379, 2380	2509
<i>lusitánica</i>	- - 2328	2478	<i>Smilax áspera</i>	- - - 2381	2510
<i>torulósa</i>	- - 2329. to 2331	2479	<i>excélsa</i>	- - - 2382	2511
			<i>rùbens</i>	- - - 2383	2511
			<i>Sarsaparílla</i>	- - - 2384	2512
<i>péndula</i>	- - 2332. to 2334	{ 2479 2480	<i>hastàta lanceolàta</i>	- - - 2385	2512
<i>Taxóidium distichum</i>	- 2335	2481	<i>quadranguláris</i>	- - - 2386	2514
<i>dist. nútans</i>	- - 2338	2482	<i>Rúscus aculeátus</i>	- - - 2387	2518
<i>sempervirens</i>	- 2340, 2341	2487	<i>hypophýllum</i>	- - - 2388	2519
<i>Juníperus commúnis</i>	} 2343. to 2349	{ 2489 to 2491	<i>racemósus</i>	- 2389, 2390	2520
and Varieties				<i>Fúcca gloriósa</i>	- - - 2391
<i>Oxýcedrus</i>	- 2351, 2352	2494	<i>supérba</i>	- - - 2392	2523
<i>macrocarpa</i>	- - 2353	2494	<i>aloiífolia</i>	- - - 2393	2523
<i>drupácea</i>	- - 2354. to 2356	2495	<i>dracónis</i>	- - - 2395	2525
<i>virginiána</i>	- - 2357	2496	<i>strícta</i>	- - - 2396	2526
<i>bermudiána</i>	- - 2358	2498	<i>filamentósa</i>	- - - 2397	2526
			<i>angustifólia</i>	- - - 2398	2526
<i>Sabína and Vars.</i>	2359. to 2363	{ 2499 2500	<i>fláccida</i>	- - - 2399	2527
<i>dáurica</i>	- - 2364, 2365	2501	<i>glaucescens</i>	- - - 2400	2527
<i>phaénicea</i>	- - 2366	2502	<i>Fourcroya longa</i>	- - - 2401	2527
			<i>Littæa gemmiflora</i>	- - - 2403	2529
			<i>Agave americana</i>	- Suppl.	2546 2606
			<i>Chamaerops humilis</i>	- - 2404	2530

LIST OF FUNGI, LICHENS, MOSSES, &c.

<i>Æcidium Pini</i>	- - 2031	2148	<i>Bulgária sarcóides</i>	- - 1681	1835
<i>Berberidis</i>	- - - 47	303	<i>Cantharellus cibarius</i>	- - 2037	2149
<i>Agáricus androsæceus</i>	- 1690	1837	<i>Craterium leucocéphalum</i>	- 1898	1974
<i>ostreátus</i>	- - 1667	1833	<i>Cenangium quercinum</i>	- 1682	1835
<i>campanélla</i>	- - - 2019	2146	<i>Dacrymyces stellatus</i>	- 2024	2148
<i>deliciósus</i>	- - - 2034	2149	<i>Daedalea betulina</i>	- - 1553	1703
<i>dryinus</i>	- - - 1665	1833	<i>gibbosa</i>	- - - 1886	1973
<i>fláccidus</i>	- - - 2021	2147	<i>quercina</i>	- - - 1668	1833
<i>hypothëjus</i>	- - - 2032	2149	<i>Ditiola radicata</i>	- - 2023	2147
<i>lepideus</i>	- - - 2020	2147	<i>Exidia auriculæ Judæ</i>	- 1686	1835
<i>maculátus</i>	- - - 2036	2149	<i>glandulosa</i>	- - 1687	1835
<i>mastrucátus</i>	- - - 1882	1971	<i>Fistulina hepática</i>	- - 1674	1834
<i>mùcidus</i>	- - 1883	1971	<i>Geoglossum cucullatum</i>	- 2039	2149
<i>multifórmis</i>	- - - 2033	2149	<i>viride</i>	- - - 1899	1974
<i>muscarius</i>	- - - 1555	1704	<i>Helvella esculenta</i>	- - 1904	1975
<i>palmátus</i>	- - - 1666	1833	<i>píleus</i>	- - - 1887	1973
<i>phlebóphorus</i>	- - - 1885	1973	<i>Hýdnum auriscálpium</i>	- 2030	2148
<i>rútilans</i>	- - - 2018	2146	<i>corallóides</i>	- - 1889	1973
<i>spínipes</i>	- - - 2028	2148	<i>Erinæceus</i>	- - 1675	1834
<i>strobilinus</i>	- - - 2029	2148	<i>imbricatum</i>	- - 2035	2149
<i>Bolëtus edulis</i>	- - 1689	1836	<i>Hýpnum cupressifórmis</i>	- - 1658	1831
<i>imbricatús</i>	- - 1888	1973	<i>denticulátum</i>	- - 1656	1831
<i>velutinus</i>	- - 1670	1833	<i>Hysterium rugosum</i>	- 1897	1974

	Figure	Page		Figure	Page
<i>Lecidea córnea</i>	- - 1664	1833	<i>Rhizopogon álbus</i>	- - 1905	1976
<i>Léskca complanáta</i>	- - 1657	1831	<i>Sclerodérma cútrinum</i>	- - 1688	1836
<i>Lóphium elátum</i>	- - 2027	2148	<i>Sclerótium quercinum</i>	- - 1691	1837
<i>mytilinum</i>	- - 2026	2148	<i>Spathulária flávida</i>	- - 2040	2149
<i>Lichen pulmonárius</i>	- - 1659	1832	<i>Sphæria agaricifórmis</i>	- - 2042	2149
<i>articulátus</i>	- - 2533	2590	<i>alutácea</i>	- - 2041	2149
<i>Morchélla esculénta</i>	- - 1902	1974	<i>deústa</i>	- - 1893	1973
<i>pátula</i>	- - 1903	1975	<i>discifórmis</i>	- - 1894	1973
<i>Nidulária erucibulum</i>	- - 2025	2148	<i>fragifórmis</i>	- - 1892	1973
<i>Opégrapha venósa</i>	- - 1906	1976	<i>Peziza</i>	- - 1895	1974
<i>Peziza aciculáris</i>	- - 1679	1834	<i>Spilonía fuliginósum</i>	- - 1662	1833
<i>eruginósa</i>	- - 1891	1973	<i>Sticta scrobiculáta</i>	- - 1660	1832
<i>aurántia</i>	- - 1678	1834	<i>Stictis radiáta</i>	- - 1683	1835
<i>fructígena</i>	- - 1900	1974	<i>Stilbum turbinátum</i>	- - 1896	1974
<i>melástoma</i>	- - 1890	1973	<i>Theléphora hírsúta</i>	- - 1677	1834
<i>polymórpha</i>	- - 1680	1834	<i>laciniáta</i>	- - 2038	2149
<i>Phacidium coronátum</i>	- - 1692	1837	<i>quercina</i>	- - 1676	1834
<i>Polýporus fomentárius</i>	- - 1672	1834	<i>sanguinolénta</i>	- - 2022	2147
<i>igniárius</i>	- - 1671	1834	<i>Tremella intumescens</i>	- - 1685	1835
<i>squamósus</i>	- - 1669	1833	<i>mesentérica</i>	- - 1684	1835
<i>versicolórus</i>	- - 1554	1703	<i>Tüber cibárium</i>	- - 1901	1974
<i>deformed sp.</i>	- - 1673	1834	<i>U'snea plicáta</i>	- - 1661	1832
<i>Puccinia Búxi</i>	- - 1219	1340	<i>Verrucaria analépta</i>	- - 1663	1833

LIST OF INSECTS.

<i>Acherontía A'tropos</i>	- - 1081	1253	<i>Ph. Auróra</i>	- - 1729	1868
<i>Balaninus núcum</i>	- - 1947	2028	<i>lúna</i>	- - 1962	2053
<i>Bómbyx móri</i>	- - 1224	1356	<i>lúcida</i>	- - 1766	1891
<i>Callidum flexuósum</i>	- - 1268	1446	<i>neógama</i>	- - 1261	1438
<i>Cántharis vesicatória</i>	- - 1047	1224	<i>neústria</i>	- - 1769	1893
<i>Catocála cloeáta</i>	- - 1293	1484	<i>Polyphémus</i>	- - 2535	2591
<i>Cerúra fúrcula</i>	- - 1518	1659	<i>stigma</i>	- - 1752	1884
<i>Clýtus Arictis</i>	- - 1541	1680	<i>tesseláris</i>	- - 1918	1981
<i>Cóssus Lignipérda</i>	- - 1233	1386	<i>Fir Cone attacked by Phalæna</i>		
<i>Jaws of the Larva of Cóssus</i>	1234	1387	<i>Tinea</i>	- - 2016	2145
<i>Lignipérda</i>	- -		<i>Fir Bud attacked by Eudórea</i>		
<i>Cýnips gállæ tinctoriæ</i>	- - 1822	1929	<i>resínea</i>	- - 2017	2146
<i>Galls of Cýnips gállæ tinctoriæ</i>	1821	1929	<i>Trees attacked by Pissódes pini</i>		
<i>Dead Sea Apples</i>	- 1823, 1824	1931	and <i>P. notátus</i>	- - 2012	2141
<i>Cýnips producing Dead Sea Apples</i>	1825	1932	<i>Scarabæus Melolóntha</i>	- - 1648	1822
<i>Galls of Diplólepis umbráculus</i>	1697	1843	<i>Scólýtus destrúctor</i>	- - 1235	1387
<i>Dóreus parallelopípedus</i>	- 635	886	<i>Silpha grísea</i>	- - 1288	1479
<i>Little Grey Ermine Moth</i>	{ 2463	2564	<i>Sinodéndron cylíndricum</i>	- - 1048	1229
<i>Scotch Pine shoots perforated</i>	{ 2464	2565	<i>Sphínx coniferárum</i>	- - 2161	2260
by <i>Hylúrgus pinipérda</i>	- 2013	2142	<i>drupiferárum</i>	- - 1253	1418
<i>Larvæ of Ladybird and Sýrphus</i>	547	811	<i>jasmíneárum</i>	- 1079, 1080	{ 1252
<i>Liméxylon nasále</i>	- - 1646	1818	<i>juglándis</i>	- - 1270	1447
<i>Lucánus Cérvus</i>	- - 1645	1816	<i>ligústri</i>	- - 1021	1201
<i>Megachile centunculáris</i>	- 548	812	<i>Ravages of Tómicus chalcógráphus</i>	- - 2015	2144
<i>Nitídula grísea</i>	- - 1289	1480	<i>Ravages of Tómicus typógráphus</i>	- - 2014	2143
<i>Noctua hastulífera</i>	- - 1544	1688	<i>Tórtrix viridána</i>	- - 1647	1818
<i>Oak Galls and Apples</i>	1649. to 1651	{ 1824	<i>Trochílium crabronifórmis</i>	- - 1290	1481
<i>Oak Spangles</i>	- - 1652	1827	<i>Larvæ of Trochílium crabronifórmis</i>	- - 1296	1482
<i>Papílio Favónius</i>	- - 1745	1879	<i>Zeuzèra æ'sculi</i>	- - 656	887
<i>Papílio Hiónæus</i>	- - 2521	2584	<i>Section of a Pear Tree perforated by Zeuzèra æ'sculi</i>	- - 637	887
<i>Phalæna álbigfrons</i>	- - 1728	1868			

LIST OF DIAGRAMS, &c.

	Figure	Page		Figure	Page
Effect of the Sea-Breeze on the Oak - - -	1	195	Basket-making - - 1277. to 1285	1473	1473
Parallelogram for drawing Trees - - -	2	205	Fences of Willow - 1286, 1287	1476	1476
Touch of the Oak - - -	3	208	Willow pierced by Trochilium crabroniforme } 1291, 1292	1482	1482
Touch of the Leaves of the wild Pear - - -	4	208	Huts of Birch Bark in Lapland	1552	1698
Outline of an entire Tree, a young Oak - - -	5	209	Oak Trees with inosculated Trunks - }	1626, 1627	1780
Outline of a young Tree of the Bird-Cherry - - -	6	210	Spray of the Oak - 1637. to 1641	1794	1794
Details of leafing, the Oak - - -	7	210	Touch of the Oak, by Strutt - 1642	1796	1796
Details of leafing, the Bird-Cherry - - -	8	210	Barnacles on the Oak 1653. to 1655	1829	1830
Chinese Characters signifying "Moutan" - - -	30	252	Barnacles attached to a Piece of Spruce Fir - - }	2541	2600
Mode of protecting Magnolia grandiflora - - -	33	266	Branching of Turkey Oak - 1720	1859	1859
Cotyledons of Acer Pseudo-Platanus - - -	110	415	Acorns of different Species of American Oaks - - }	1722	1863
Mode of planting Robinia hispida - - -	111	416	Axe for detaching Cork from the Trunk of the Cork Tree	1799	1915
Designs for a Rosarium { 525. to 527	{	{ 795	Spray of the Beech - 1879, 1880	1967	1967
Arch for training Roses - - -	528	797	Inosculated Beeches - - -	1881	1971
Trellis for ditto - - -	529. to 531	798	Rustic Garden Seat of Hazel Rods - - -	1884	1972
Standards for training Roses - - -	532. to 536	799	Rustic Flower-Basket of Hazel Rods - - -	1944	2023
Mode of training Roses on Walls	537	800	Rustic Arbour of Hazel Rods -	1946	2025
Budding Roses - - -	538. to 542	{ 802	Plane Tree Leaf, and Map of the Morea - - -	1958	2038
Pruning Roses - - -	543. to 545	{ 805	Log House of Pine Trunks -	2006	2123
Training standard Roses - - -	546	809	American Worm Fence -	2007	2123
Germination of the Mistletoe -	772	1024	Swedish Wooden Fence -	2008	2124
Honeysuckle, architectural ornament - - -	796	1043	Grafting the Abietinae - 2009, 2010	2130	2131
Plans for Ericetums, or Heatheries - - -	{ 886	{ 1096	Skeleton of a Scotch Pine at Gordon Castle - - -	2049	2162
Brick and Tile Drains - 999. to	1001	1175	Mode of cutting out Deals from the Trunks of Pine and Fir Trees	2053, 2054	2170
Designs for Ericetums, or American Grounds - - -	{ 1002	{ 1178	Section of Larch Board, having a Branch with the Bark on embedded in it - - -	2265	2380
Spray of the Ash - - -	1046	1222	Sections of Larch affected by Rot - - -	2266	2386
Grafting Ornus europæa - - -	1066. to 1068	{ 1242	Frame and Cover for protecting Half-hardy Plants - - -	2405, 2406	2531
Grafting Ornus americana -	1071	1245	Ground Plan of an Espalier for training the Locust -	2439	2552
French Parterres - 1216, 1217	{	{ 1337	Black Mulberry at Canterbury	2526	2586
Spray of the Elm - - -	1232	1382	Design for a Labyrinth - - -	2538	2599
Elms injured by Scolytus destructor - - -	1236, 1237	{ 1388			
Birds (Titmouse) on the Walnut	1256	1424			
Budding the Walnut - 1258, 1259	{	{ 1431			
		1432			

LIST OF AUTHORITIES

FOR GENERIC AND SPECIFIC NAMES, &c.

- A.
- A. et S.* - *Albertini et Schwinitz.* Moravian missionaries resident in America.
- Abb. et Smith.* *Abbott and Smith.* The first an amateur, the second the founder of the Linnaean Society.
- Abbott.* - *Abbott.* See *Abbott et Smith.*
- Abel.* - *Abel.* Author of "Personal Observations, &c. in China."
- Ach.* - *Acharius.* A Swedish professor, and writer upon Lichens.
- Ach. Rich.* - *Acholle Richard.* A French botanist.
- Adans.* - *Adanson.* A French systematic botanist.
- A. De C.* - *Alphonse De Candolle.* Author of "Monographie des Campanulées," &c.; and son of the distinguished De Candolle.
- Afzel.* - *Afzelius.* A Swedish professor.
- Ait.* - *Aiton.* Director of the Royal Garden at Kew.
- Ab. et Schw.* *Albertini and Schwinitz.* Writers upon Fungi.
- All. } - Allioni.* An Italian botanist.
Allioni. }
- Alpin. } - Prosper Alpini.* An Italian physician, and author of "De Plantis Ægypti et de Balsamo," &c.
- Anders. } - Anderson.* A London merchant; published a paper on Pæonies.
Anderson. }
- Andr. } - Henry Andrews.* A botanical draughtsman, and editor of the "Botanical Repository," &c.
Andrews. }
- Audib. } - Audibert.* A nurseryman at Tarascon, in the south of France.
Audibert. }
- Austin.* - *Austin.* A nurseryman at Glasgow.
- B.
- Bab.* - *Babington.* An English botanist.
- Backhouse.* - *Backhouse.* A nurseryman at York.
- Balb. } - Balbis.* A French professor of botany.
Balbis. }
- Banister.* - *Banister.* A botanical author, and traveller.
- Barrcl. } - Barrelier.* A French botanist.
Barrelier. }
- Bartr. } - Bartram.* Formerly a nurseryman at Philadelphia.
Bartram. }
- Bat. } - Bastard or Batard.* A writer upon the Flora of France.
Bast. }
- Batsch.* - *Batsch.* A writer upon Fungi.
- Bauh. } - Bauhin.* Brothers, professors of medicine, published in 1620—1650.
Bauhin. }
- Benth. } - Bentham.* An English botanist, secretary to the Horticultural Society of London.
Bentham. }
- Bergius.* - *Bergius.* A Swedish writer upon Cape Plants.
- Berk.* - *Rev. M. J. Berkeley.* An English cryptogamic botanist.
- Bert. } - Bertoloni.* A writer upon the Flora of France.
Bertol. }
- Bess. } - Besser.* A Russian professor, resident in the Crimea.
Besser. }
- Bieb.* - *Bieberstein.* A Russian botanist of great note.
- Bigcl. } - Jacob Bigelow, M.D.* Professor of botany at Boston, U. S., and author of "American Medical Botany," and "Florula Bostoniensis."
Bigelow }
- Big.* }
- Blume.* - *Blume, M.D.* A Dutch botanist.
- Boerh.* - *Boerhaave.* An old Dutch botanist.
- Bolt.* - *Bolton.* An English writer on Fungi.
- Booth.* - *Booth.* Brothers, nurserymen at Hamburg.
- Banpl.* - *Bonpland.* A French traveller in South America, and botanist.
- Booth.* - *W. Beattie Booth.* Describer of the Camelias figured in Chandler's "Illustrations of the Camellæe."
- Bork.* - *Borkhausen.* A German botanical author.
- Borrer.* - *William Borrer.* A writer on British Plants, and one of the authors of "Lichenographia Britannica."
- Bosc.* - *Bosc.* A French botanist, and traveller in North America.
- Breyn.* - *Breyn.* Author of "Exoticarum Plantarum Centuria," &c.
- Brong.* - *A. Brongniart.* A French botanist.
- Brot.* - *Brotero.* A Portuguese botanist.
- Broussonet.* *Broussonet.* A French botanist, and traveller.
- Buch.* - *Von Buch.* A German botanist, an author of a Flora of the Canaries.
- Buchanan.* *Buchanan,* formerly Hamilton. An English physician, and traveller in Nepal.
- Bull.* - *Bulliard.* A French writer upon Fungi.
- Bunge.* - *Bunge.* One of the authors of "Flora Altaica."
- Burch.* - *Burchell.* An English botanist, and traveller at the Cape of Good Hope.
- Burgsdorf.* *Burgsdorf.* A German botanist.
- Busch.* - *Busch.* A German gardener, once a nurseryman at Brentford, Middlesex; and afterwards gardener to the Empress Catherine, at Zarskojesele.
- Burb.* - *Burbaum, M.D.* A Russian

- botanist; traveller through Armenia.
- C.
- Cæsalp.* - *Cæsalpinus*. A famous old Italian botanist.
- Camb.* - *Cambessedes*. One of the authors of "Flora Brasiliæ meridionalis."
- Camer.* - *Camerarius*. A German botanist, author of "Hortus Medicus et Philosophicus," &c.
- Cass.* - *H. Cassini*. A French botanist.
- Castagne.* - *L. Castagne*. A French botanist?
- Catesb.* } - *M. Catesby*. A botanist, and
Catesby. } traveller in North America.
Catros. } - *Catros*. A nurseryman at Bordeaux.
- Cav.* - *Cavanilles*. A Spanish professor and botanist.
- C. Bauh.* - *Caspar Bauhin*. A celebrated botanist of the 16th century.
- Cels.* - *Celsius, D.D.*, Greek professor at Upsal, and friend of Linnæus.
- Cels.* - *Cels*. A nurseryman in Paris.
- C. G. Nees Von Esenbeck.* } - *C. G. Nees Von Esenbeck*. A German botanist.
- Chaill.* - *M. Chaillot*. A Swiss botanist.
- Cham.* - *Chamisso*. A German traveller round the world.
- Chois.* - *Choisy*. A Swiss botanist.
- Clus.* - *Clusius*. An old French botanist and traveller.
- Colebr.* - *Colebrooke*. A celebrated English writer upon Indian Plants.
- Colla.* - *Colladon*. A Genevese botanist.
- Com.* - *Commelin*. A Dutch botanist.
- Cook.* - *Capt. S. E. Cook*. A naturalist and traveller.
- Correa.* - *Correa de Serra*. A Portuguese botanist and diplomatist.
- Crantz.* - *Crantz*. An Austrian botanist.
- Cullum.* - *T. G. Cullum*. A botanical amateur.
- Cunn.* } - *A. Cunningham*. Colonial botanist at Sydney.
Cunningham. }
Curt. - *Curtis*. An English writer upon Plants.
- D.
- Dalech.* - *Daléchamps*. Author of "Historia generalis Plantarum." 1586, 1587.
- Darlington.* - *Darlington*. A writer in "Amer. Lyc. N. H." of New York.
- Davies.* - *H. Davies, D.D.* A Welsh botanist.
- D. Don.* - *David Don*. Professor of botany in King's College, London, and librarian to the Linnæan Society.
- Deb.* } - *Debray*. A botanist of Frankfurt.
De Bray. }
Dec. } - *A. P. De Candolle*. The celebrated French systematic botanist.
De Candolle. }
- De Geer.* - *C. De Geer*. Author of "Mémoires pour servir à l'Histoire des Insectes," &c.
- Dejean.* - *Le Comte Dejean*. A French naturalist.
- Del.* - See *Delile*.
- Delamarre.* - *Delamarre*. A French writer on plants.
- Delarb.* - *Ant. Delarbec*. A French botanist, author of "Flore d'Auvergne."
- Delile.* - *Delile*. A French professor, and traveller in Egypt.
- Descemet.* - *Descemet*. Director of the Botanic Garden at Nikitka, in the Crimea.
- Desf.* - *Desfontaines*. A French botanical author, and traveller in Barbary.
- Desrous.* - *Desrousseaux*. A French botanist.
- Desm.* - *Desmazières*. A French cryptogamic botanist.
- Desv.* - *Desvaux*. A French professor of botany.
- Dickson.* - *Dickson*. An English cryptogamic botanist.
- Dios.* - *Dioscorides*. An ancient classic author and botanist.
- Dod.* } - *Dodonæus, or Dodoens*. A
Dodon. } botanist of the 16th century.
- Domb.* - *Dombey*. A French traveller in South America.
- Don of Forfar.* - *Don of Forfar*. A Scotch botanist.
- Donn.* - *Donn*. Formerly curator of the Cambridge Botanic Garden.
- Dougl.* } - *David Douglas*. The celebrated botanical collector and martyr.
Douglas. }
- Dralct.* - *Dralct*. A French writer on the oak.
- Duby.* - *Duby*. A French botanist.
- Dufour.* - *Dufour*. A French botanist.
- Dumont.* - *Dumont de Courset*. A writer upon French garden plants.
- Dun.* } - *Dunal*. A French botanist.
Dunal. }
- Dupont.* - *Dupont*. A French botanist, author of "Double Flore Parisienne." 1805.
- Dur.* } - *Durand*. A French botanist.
Durand. }
- Du Roi.* - *Du Roi*. A German writer upon plants.
- E.
- Ehr.* - See *Ehrenberg*.
- Ehrenberg.* - *Ehrenberg*. A German traveller in Arabia, &c.
- Ehrh.* - *Ehrhart*. A German botanist.
- Elliot.* - *Elliot*. An American botanist.
- Ellis.* - *Ellis*. A London merchant and botanist.
- Eschsch.* - *Dr. Eschscholtz*. A German botanist.
- Esper.* - *Esper*. A German writer on Fungi.
- F.
- Fab.* } - *P. C. Fabricius*. A German
Fabr. } botanist.
Falla. - *Falla*. A nurseryman at Gateshead, near Newcastle.
- Fing.* - *Fingerhuth*. A German botanist.
- Fisch.* } - *Dr. Fischer*. A Russian botanist.
Fischer. }
- Floy.* - *Floy*. A nurseryman in New York.
- Flügge.* - *Flügge*. A German writer upon grasses.
- Forbes.* - *Forbes*. Gardener to His Grace the Duke of Bedford, at Woburn.
- Forsk.* - *Forsk. Skth.*. A Danish naturalist, and traveller in Arabia.
- Forst.* - *Forsters (Father and Son)*. Travellers in the South Seas with Captain Cook.
- Fourc.* - *A. F. Fourcroy*. A French botanist.
- Fries.* - *Fries*. A Swedish botanist, and writer upon Fungi.

Fuch. - *Fuchs*. A celebrated German botanist.
G.
Gart. - *Gartner*. A celebrated German carpologist.
Garden. - *Garden*. A Scotch physician resident at Charleston.
Gaudichaud. - *Gaudichaud*. A French botanist.
G. Don. - *Geo. Don*. A botanist, and editor of "Don's Miller."
Geoffr. - *Geoffroy*. A French botanist of the 18th century.
Ger. et Lob. - See *Gérard*, and also *Lobel*.
Ger. }
Gérard. } - *Gérard*. A French botanist.
Germar. } - *Germar*. A German entomologist.
Gesn. - *Conrad Gesner* of Zurich, a famous botanist.
Gill. - *Dr. Gillies*. A botanist and traveller.
Gill. et Hook. - See *Gill*, and also *Hook*.
Gmel. - *Gmelin*. A Russian botanist, and traveller in Siberia.
Godefroy. - *Godefroy*. A nurseryman at Ville d'Avray, near Paris.
Goldie. - *Goldie*. A nurseryman at Ayr, in Scotland.
Gordon. - *George Gordon*. Superintendent of the Arboretum in the Horticultural Society's Garden.
Gouan. }
Grah. } - *Gouan*. A French botanist.
Graham. } - *Graham, M.D.* Regius professor of Botany at Edinburgh
Grev. - *Dr. Greville*. An English botanist, and writer upon cryptogamous plants.
Grigor. - *Grigor*. A nurseryman at Elgin.
Güldenst. - *Güldenstedt*. A Russian botanist.
Guss. - *Joannes Gussone, M.D.* Director of the Royal Botanic Garden at Palermo, and a botanical author.
Gyll. - *L. Gyllenhall*. Author of "Insecta Succica descripta." 1808, 1827.

II.

Henke. - *Henke*. A German botanical writer.
Hakluyt. - *Hakluyt*. Author of "Two Remembrances of Things to be undiscovered in Turkey, &c." 1592.
Hall. - *Haller*. A Swiss botanist.
Hall, fil. - *Haller the younger*.
Ham. }
Hamill. } - *Hamilton*. A botanist, and traveller in the East Indies.
Hanbury. - *Hanbury*. A writer upon gardening.
Harris. - *Harris*. An entomologist.
Hart. - *Hartweg*. Author of "Florus Carlsruhensis."
Hartig. - *Hartig*. A writer on the culture of forest trees.
Hassclq. - *Hassclquist*. A botanist, and traveller in the Levant.
Hayne. - *Hayne*. A German botanist.
H. B. - *Humboldt and Bonpland*. Famous travellers and botanists.
H. B. et Kth. - *Humboldt, Bonpland, and Kunth*. German botanists.
H. H. - *N. C. Heltenus*. A Swedish botanist.
H. et Kth. - *Humboldt and Kunth*. German botanists.
H. L. W. - *Henry Lee Warner, Esq.*, Tibberton Court, Herts.

Hodgins. - *Hodgins*. Nurseryman, Dunganstown, near Wicklow.
Hoffmans - *Hoffmannsegg*. A botanist of Dresden.
Holmsk. - *Holmskiöld*. A Danish botanist.
Hook. - *Sir W. J. Hooker*. Regius professor of botany in the University of Glasgow.
Hook. et Arn. - *Sir W. J. Hooker and Arnott*, authors of "The Botany of Captain Beechey's Voyage to the Pacific," &c.
Hoppe. }
Hope. } - *Hoppe*. A German botanist and collector of plants.
Horn. - *Hornemann*. A Danish botanist and professor.
Hort. - *Hortulanorum*. Of the Gardeners.
Hort. Par. - *Of the Paris Garden*.
Host. - *Host*. An Austrian botanist, and botanical author.
Howison. - *Howison*. A writer in the "Edin. Phil. Journ."
Huds. - *Hudson*. An English writer upon British plants.
Hull. - *Hull*. A English botanist.
Humb. et Bonp. }
Humb. et Bonpl. } - *Humboldt and Bonpland*. — See *H. B.*
I.
Illig. }
Illiger. } - *C. Illiger*. A writer on natural history.
J.
Jacq. - *Jacquin*. An Austrian traveller in South America, and botanist.
Jaume St. Hilaire. - *Jaume Saint Hilaire*. A French botanist, and author of "Plantes de la France," &c.
J. B. }
J. Bauh. } - *Jo. Bauhin*, brother of Caspar Bauhin, and author of "Historia Plantarum universalis." 1650, 1651.
John Grigor. - *John Grigor*, nurseryman of Forres.
J. O. W. - *J. O. Westwood*. An English entomologist.
Juss. - *Jussieu*. A celebrated French systematic botanist.
Juv. - *Juvenal*. A classic author.
K.
Kæmpf. - *Kæmpfer*. A traveller in Japan.
Karw. et Zuccar. - *Baron Karwinski and Zuccarini*. Zealous promoters of botany in Germany.
Kalm. - *Kalm*. Professor of botany at Abo, in Sweden.
Ker. - *Ker*. A describer of plants in "Bot. Reg."
Kerner - *J. S. Kerner*. Author of "Figures (et Descriptions) des Plantes économiques." 1786—1794.
Kit. }
Kitaib. } - *Kitaibel*. A Hungarian botanist.
Klotzsch. - *Klotzsch*. A German writer on Fungi.
Knowl. et West- }
cott. } - *Knowles (G. B.) and Westcott (F.)*. Conductors of the "Floral Cabinet."
Koch. - *Koch*. A professor at Erlangen.
Kochl. - *Koehler*. A writer on German and French grasses.
Krause. }
Krauss. } - *Krause*. A Dutch botanist, and author of "Boomen en Heestar."
Kunth. - *Kunth*. A Prussian botanist.

L.

L. - *Linnaeus*. The celebrated reformer of natural history.

Lab. } - *Labillardière*. A French botanist.

Labill. }
Lag. }
La Gasca. }
Lam. }
Lamarck. }
Lamb. }
Lambert. }
Lap. }
Lapey. }
Lat. } - *Latrille*. A French entomologist.

Latr. }
Laur. } - *Lawrence (Miss)*. An English flower-painter.

Laxm. - *Laxmann*. A German writer on Siberian plants.

Leach. - *W. E. Leach*. An English entomologist and author.

Lecch. - *Lechenault*. A French botanist.

Led. }
Ledebour. }
Lec. } - *Lec*. A nurseryman at Hammersmith.

Lehm. - *Lehmann*. A German botanist and professor at Hamburg.

Lejeunc. - *Lejeune*. A French botanist.

Lessing. - *Lessing*. A writer on Composite, and resident at Berlin.

Lewis. - *Lewis*. An American traveller.

L. fil. - *Linnaeus the younger*.

L'Hér. }
L'Hérit. }
Lighfoot. } - *C. L. L'Héritier*. A French botanist and author.

Lin. - *Linnaeus*. The celebrated reformer of natural history.

Lindl. - *Dr. Lindley, F.R.S., &c.* Professor of botany in University College, London.

Link. }
Lk. }
Link et Otto. }
Lk. et Otto. }
Lob. }
Lobel. }
L'Obel. }
Lodt. }
Loddiges. } - *Messrs. Loddiges*. Celebrated English nurserymen and botanists, Hackney.

Lois. }
Lois. Destlong- }
champs. }
Loudon. } - *J. C. Loudon, F.L.S., &c.* Author of this work, and of various works on gardening, agriculture, &c.

Lour. - *Lourcuro*. A Portuguese traveller in Cochiu-China.

L. pat. - *Linnaeus the elder*.

Lysons - *Lysons*. Author of "Envi-
 rons of London," &c.

M.

Macgillivray. - *Macgillivray*. A botanist, and editor of a small edition of "Withering's Botany," &c.

Mackie. - *Mackie*. A nurseryman at Norwich.

Main. - *J. Main, A.L.S.* A botanical and gardening author.

Mal. }
Malp. } - *Malpighi*. A physiological botanist.

Malcolm. - *Malcolm*. Late nurseryman at Kensington.

Marsh. }
Marshall. }
Marsham. } - *Marshall*. A writer on American trees.

Masters. - *W. Masters* of the Canterbury Nursery.

Math. - *Mathiolus*. An Italian physician.

Mayes. - *Mayes*. A writer in the "West of England Journal of Science," &c.

Medik. - *Medicus*. A German botanist of the last century.

Menzies. - *Menzies*. A Scotch botanist, and traveller round the world with Vancouver.

Mertens. - *Mertens*. A French botanist.

Meycr. - *Meyer*. A German botanist.

Michx. - *Michaux*. A French botanist, and traveller in N. Amer., and author of "Flora Borealis Americana."

Michx. fil. }
Michx. jun. }
Mill. } - *Michaux the younger*. Also a botanist and traveller in N. America, author of "Histoire des Arbres de l'Amérique."

Mill. - *Miller*. An English gardener and botanist.

Mirb. - *Mirbel*. A French physiological botanist.

M. J. B. - *The Rev. M. J. Berkeley, F.L.S., &c.* An eminent English cryptogamist.

M'Nab. - *M'Nab*. Superintendent of the Edinburgh Botanic Garden.

Moc. - *Mocino*. A Mexican botanist.

Moc. et Scssc. - *Mocino and Sesse*. Two Mexican botanists.

Monch. - *Monch*. A German botanist.

Mol. - *Molina*. An Italian writer upon the natural history of Chili.

Mor. - *Morison*. An old writer on plants.

Moug. et Nest. - *Mougeot*, a German cryptogamist; and *Nestler*, a botanist of Strasburg.

Mont. - *J. Monti*. An Italian botanist.

Mr. G. Lindley. - *Mr. George Lindley*, late nurseryman, Norwich.

Mühl. }
Mühlentb. }
Mühlberg. }
Munch. } - *Mühlenberg*. A North American botanist.

Muntl. }
Munting. }
Murr. } - *Baron Otto von Munchausen*. A German botanist.

Mutis. - *Mutis*. A Spanish botanist, resident in New Grenada.

Mx. - *Michaux*. See *Michx.*

N.

Neck. }
Necker. }
Nec. } - *Necker*. A German writer upon botanical affairs.

Nees. }
Nees von Escn- }
beck. }
Ncill. } - *Louis Nec*. A Spanish botanist.

Nees. }
Nees von Escn- }
beck. }
Ncill. } - *Nees von Esenbeck*. A German botanist.

Ncill. }
Nestl. } - *Dr. Neill* of Canon Mills. A zealous botanist, and promoter of horticulture.

Nestl. - *Nestler*. A botanist of Strasburg.

- Niss.* } - *Nissolle.* A French botanist.
Nissole. }
Nois. } - *Noisette.* A French nurseryman.
Noisette. }
Noronha. } - *Noronha.* A Spanish botanist who visited Madagascar.

Nutt. } - *Nuttall.* A North American botanist.
Nuttall. }

O.

Old. - *Older.* A Danish botanist.
Oldaker. - *Oldaker.* Formerly gardener to Sir Joseph Banks.
Oliv. } - *Olivier.* A French botanist, and traveller in Persia.
Olivier. }
Opiz. } - *Opiz.* A German botanist.
Or. }
Ort. } - *Ortega.* A Spanish botanist.
Otto. } - *Otto.* Director of the Royal Botanic Garden, Berlin.
Otto et Hayne. - *Otto and Hayne.* Two German botanists.

P.

Pall. - *Pallas.* A Russian traveller and naturalist.
Parm. - *Parmentier.* A French nurseryman.
Parkinson. - *Parkinson.* An old botanical author.
Penny. - *George Penny.* A botanist and nurseryman.
Perrotlet. - *Perrotlet.* A French botanist.
Pers. - *Persoon.* A French botanist and botanical author.
Pesch. - *Peschier.* Author of "Disp. inaug. de Irritabilitate Animalium et Vegetabilium." 1697.
Petit Thouars. - *Aubert du Petit Thouars.* A French botanist and botanical author.
Ph. - *Pursh.* A Prussian botanist, and traveller in North America.
Phillips. - *Phillips.* Author of "Sylva Florifera," &c.
Pliny. - *Pliny.* An ancient naturalist and classic author.
Plumier. - *Plumier.* A French botanist, and traveller in the West Indies.
Pohl. - *Pohl.* A German botanist.
Poir. - *Poir.* A French botanical writer.
Poit. et Turp. - *Poitau and Turpin.* French botanical draughtsmen.
Poll. - *Pollich.* A German writer on the plants of the Palatinate.
Potts. - *J. Potts.* A collector of plants in China.
Pourr. - *Pourret.* A French botanist.
Presl. - *Presl.* A Bohemian botanist.
Puer. - *Puceri.* A professor at Copenhagen.
Pursh. - *Pursh.* A Prussian botanist, and traveller in North America.

R.

R. et P. - *Ruiz and Pavon.* Spanish botanists, and travellers in Peru and Chile.

Raf. }
Rafin. } - *Rafinesque Schmalz.* A botanical author.
Rafinesque. }
Rand. } - *Rand.* Author of "Index Plantarum officinalium Horti Chelseani," &c. 1720.

Ratzeburg. - *Ratzeburg.* An entomologist.
Ray. - *John Ray.* A celebrated botanist and naturalist.
Reb. - *J. F. Reichenbach.* A German botanist.
R. Br. }
R. Brown. } - *Dr. Robert Brown, F.R.S., &c.* A distinguished English botanist and traveller in New Holland.

Red. }
Redouté. } - *Redouté.* A French botanical draughtsman.
Reich. } - *Reichenbach.* A German botanist.

Renault. - *Renault.* A French botanical writer.
Recaulm. - *P. Recaulme.* Author of "Specimen Historiæ Plantarum." 1611.
Retz. - *Retzius.* A German botanist.
Reyn. - *Reynier.* A botanist of Lausanne.
Rich. - *Richard.* A French botanist.
Richards. }
Richardson. } - *Dr. Richardson.* A traveller in the northern parts of British America, and author of the Appendix on Natural History to Franklin's "Travels."

Risso. - *Risso.* An Italian writer upon oranges.
Robert T. Pincc. - *Robert T. Pinck.* of the Exeter Nursery.
Robs. - *Robson.* An English botanist.
Rochel. - *Rochel.* Superintendent of the garden at Pesth.
Röm. et Schult. }
Ram. et Schult. } - *Harmer,* a German botanist; and *Schultz,* a Bavarian botanist.

Roll. - *Rollinson.* A nurseryman near London.
Ronalds. - *Ronalds.* A nurseryman at Brentford.
Rossmüssler. - *Rossmüssler.* A German entomologist.
Rott. - *Rottler.* A German missionary.

Roxas di San }
Clemente. } - A Spanish botanist.

Roxb. - *Rozeburgh.* An Indian botanist.
Royle. - *Dr. Royle, F.R.S., &c.,* Prof. of Mat. Med. in King's College. Author of "Illustrations of the Natural History and Botany of the Himalayas," &c.

Rudge. - *Rudge.* An English writer upon botanical subjects.
Rudolphi. - *Rudolphi.* A German cryptogamic botanist.

Ruiz et Pav. }
Ruiz et Pavon. } - *Ruiz and Pavon.* Spanish botanists, and travellers in Peru and Chile.

Rumph. - *Rumphius.* Author of "Herbarium Amboinensis."

Russ. - *Russell.* A botanist of Aleppo.

S.

Sab. }
Sabinc. } - *J. Sabine, F.R.S., &c.* A great encourager of natural history, botany, and gardening.

Sal. }
Salisb. } - *Salisbury.* An eminent English botanist.

Salzm. } - *Salzmann.* Author of "Enumeratio Plantarum rariorum, quas collegit." 1818.

Samouelle. - *Samouelle.* A British entomologist and author.

Santi. - *Santi.* An Italian botanist.

Savi. - *Savi.* An Italian botanist.

Schæff. - *Schæffer.* A German writer upon Fungi.
Schiede et. Deppe. *Schiede and Deppe.* Writers on the botany of Mexico.
Schl. - *Schleicher.* A Swiss plant collector.
Schlecht. - *Schlechtendahl.* A celebrated German botanist.
Schlecht. et Cham. *Schlechtendahl and Chamisso.* Two German botanists.
Schleicher. - *Schleicher.* A Swiss plant-collector.
Schmidt. - *Schmidt.* A German botanical author.
Schneevoogl. *Schneevoogl.* Author of "Icones Plantarum rariorum." Haarlem, 1793.
Schober. - *Schober.* A Russian physician.
Schonberg. *Schonberg.* A German botanist.
Schott. - *Scholl.* A traveller in Brazil.
Schousb. *Schousboe.* A writer upon the Flora of Morocco.
Schouw. - *Schouw.* A celebrated Danish botanist, and professor at Copenhagen.
Schrad. } - *Schrader.* A German botanist.
Schrader. }
Schrank. } - *Schrank.* A Bavarian botanist.
Schreb. - *Schreber.* A German botanist.
Schubert. - *Schubert.* A Polish botanist.
Schult. } - *Schultes.* A Bavarian botanist.
Schultes. }
Schum. } - *Schumacher.* A Danish botanist.
Schwein. - *Schweinitz.* An American botanist.
Scop. - *Scopoli.* An Italian botanist.
Scouler. - *D. Scouler.* Professor at Dublin.
Secondat. - *Secondat.* A French naturalist.
Ser. } - *Seringe.* A Swiss botanist.
Seringe. }
Sèvres. } - *Sèvres.* A French writer on agriculture.
Sibth. - *Sibthorp.* An English botanist, and professor of botany at Oxford, traveller in Greece, and author of "Flora Græca."
Sieb. - *Sieber.* A botanical collector.
Sieb. - *Siebold (Dr. von).* A Bavarian botanist, who has imported many valuable plants from Japan.
Sievers. - *Sievers.* A German botanist.
Sims. - *Sims.* An English garden botanist.
Smith. - *Sir J. E. Smith.* Founder of the Lin. Soc., and author of several botanical works.
Smith of Ayr. *Smith.* Nurseryman of Ayr.
Sol. } - *Solander.* A Swedish botanist, and companion of Sir Joseph Banks in Cook's voyage round the world.
Soland. }
Somm. - *Sommerschelt.* A German writer on Fungi.
Soulange-Bodin. *Soulange-Bodin.* A French botanist and horticulturist.
Sow. - *Sowerby.* An English botanical draughtsman.
Spenser. - *Spenser.* The celebrated poet.
Steele. - *Steele.* A writer in the "Gardener's Magazine."
Steph. } - *Stephens.* A British entomologist.
Stephens. }
Stev. } - *Steven.* A Russian botanist.
Steven. }
Stev. et Fisch. } - *Steven and Fischer.* Russian botanists.
Stokes. - *Stokes.* An English physician.
Strab. - *Strabo.* The geographer.
Straus. - *Strauss.* A German botanist.

Swt. - *Sweet.* An English botanist.
Swz. } - *Swartz.* A Swedish botanist, and traveller in the West Indies.
Swartz. }

T.

Tabern. - *Tabernaemontanus.* A German botanist.
Talc. - *Tate.* Nurseryman, Sloane Street, Chelsea.
Tausch. - *Tausch.* A German botanist.
Tav. - *Tavernier.* A traveller in the East.
Ten. } - *Tenore.* A Neapolitan botanist.
Tenore. }
Theoph. - *Theophrastus.*
Thib. - *Thibaud De Chanvalon.* A French botanist.
Thomp. } - *Robert Thompson.* Superintendent of the fruit department in the Chiswick Garden.
Thompson. }
Thorc. - *Thorc.* A French botanist.
Thory. - *Thory.* A French botanist.
Thouin. - *Thouin.* A French botanist.
Thuil. } - *Thuilleier.* A French botanist.
Thuillier. }
Thun. } - *Thunberg.* A Swedish botanical traveller.
Thunb. }
Tighe. - *Tighe.* An English poet.
Tineo. - *Tineo.* A professor at Palermo.
T. Nées ab E. *T. Nées ab Esenbeck.* A German botanist.
Tode. - *Tode.* A German writer on Fungi.
Torrey. - *Torrey.* An American botanist.
Tourn. - *Tournefort.* An old French botanist, and traveller in Greece and Asia Minor.
Trag. - *Tragus or Bock.* A German botanist.
Trat. - *Trattinik.* A botanist of Vienna.
Turner. - *Dr. Turner.* A British botanist and author.
Turp. - *Turp.* A French botanist and draughtsman.

V.

Vahl. - *Vahl.* A botanical author.
Vail. - *Vaillant.* A French botanist and traveller.
Vcat. - *Ventnat.* A French botanist.
Vest. - *Vest.* A Styrian botanist.
Fill. } - *Villars.* A French botanist.
Villars. }
Vilmorin. - *Vilmorin.* A nurseryman and botanist at Paris.
Virg. - *Virgil.*
Vitn. - *Vitmann.* A German botanist.
Viv. - *Viviani.* An Italian botanist.

W.

W. - *Willdenow.* A German botanist, and editor of an edition of Linnaeus's "Species Plantarum," &c.
Wahlenb. - *Wahlenberg.* A Swedish botanist.
Waitz. - *Waitz.* A German collector of roses.
Wal. } - *Waldstein.* A noble German patron of botany.
Waldst. }
Waldst. et Kil. *Waldstein and Kitabel.* Authors of the "Flora of Hungary."

- | | | | | | |
|------------------------|---|-----------------------------------------------------------------------------------------------------------------|----------------------|---|-----------------------------------------------------------------------------------------------------|
| <i>Walker.</i> | - | <i>Dr. Walker.</i> A Scotch writer on plants. | <i>Westw.</i> | - | <i>J. O. Westwood, F.L.S.</i> An eminent English entomologist, and secretary to the Lond. Ent. Soc. |
| <i>Wall.</i> | - | <i>Wallich.</i> Superintendent of the botanical garden at Calcutta. | <i>W. Henderson.</i> | - | <i>W. Henderson.</i> Crum Castle. |
| <i>Wallr.</i> | - | <i>Wallroth.</i> A German botanist. | <i>Whitley.</i> | - | <i>Whitley.</i> A nurseryman at Fulham. |
| <i>Walt.</i> | - | <i>Walter.</i> A writer on the "Flora of Carolina." | <i>Wicks.</i> | - | <i>Wickstrom.</i> A German botanist. |
| <i>W. et N.</i> | - | <i>Weiche and Nees.</i> German writers on Rûbi. | <i>Willd.</i> | - | See <i>W.</i> |
| <i>Wang.</i> | } | <i>Wangenheim.</i> A German botanist. | <i>Winch.</i> | - | <i>Winch.</i> A botanist, and writer on the flora of Northumberland and Durham. |
| <i>Wangenh.</i> | | | <i>With.</i> | - | <i>Withering.</i> An English botanist. |
| <i>Wats.</i> | } | <i>Watson.</i> An English writer upon trees and shrubs. | <i>Wood.</i> | } | <i>Woods.</i> An English writer on roses. |
| <i>Watson.</i> | | | <i>Wormsk.</i> | | |
| <i>W. B.</i> | - | <i>W. Borrer, F.L.S.</i> Accelerated English botanist, and writer upon English plants, more especially willows. | <i>Wulf.</i> | - | <i>Wulfen.</i> A German botanist. |
| <i>Weiche.</i> | - | <i>Weiche.</i> A German writer on Rûbi. | Z. | | |
| <i>Weiche et Nees.</i> | - | <i>Weiche and Nees.</i> Two German writers on Rûbi. | <i>Zeyh.</i> | - | <i>Zeyher.</i> A German gardener. |
| <i>Wend.</i> | - | <i>J. W. F. Wenderoth.</i> A German botanist. | <i>Zucc.</i> | - | <i>Zuccagni.</i> Superintendent of the Botanic Garden at Florence. |
| <i>Wendl.</i> | - | <i>Wendland.</i> A German garden botanist. | | | |

LIST OF BOOKS REFERRED TO,

THE TITLES OF WHICH ARE ABRIDGED IN THE TEXT.

The number following each contracted title refers to the page in which the book is first cited.

A.

- Abbildung der deutschen Holzarten, &c.* 190. F. Guimpel, C. L. Willdenow, and F. G. Hayne, Abbildung der deutschen Holzarten, &c. Berlin, 1850, 4to, pl. 216.
- Abbildung der fremden in Deutschland ausdauernden Holzarten, &c.* 190. F. Guimpel, Abbildungen der fremden in Deutschland ausdauernden Holzarten, mit Angabe der Cultur von F. Otto, und Beschreibung von F. G. Hayne. Berlin, 1819—1825, 4to.
- Abb. Inst.* 1877. See *Abbott and Smith's Natural History of Georgia.*
- Abb. and Smith, Ins. of Georgia.* 1659. See *Abbott and Smith's Natural History of Georgia.*
- Abbott and Smith's Insects of Georgia.* 1418. See *Abbott and Smith's Natural History of Georgia.*
- Abbott and Smith's Natural History of Georgia.* 1658. *Abbott and Smith's Natural History of the Lepidopterous Insects of Georgia.* London, 1797. 2 vols. fol.
- Abel's Chin.* 391. Personal Observations made during the Progress of the British Embassy to China, and on its Voyage to and from that Country in the years 1816 and 1817. By Clerk Abel. Lond. 1818. 4to.
- Abhandl. Königl. Akad. Wissens. Berlin.* 2293. Abhandlungen der Physikalischen Klasse der Königlich-Preussischen Akademie der Wissenschaften aus den Jahren 1820 und 1821. Berlin, 1822. 4to.
- Account of the Larch Plantations on the Estates of Athol and Dunkeld.* 2358. A Paper published in the Transactions of the Highland Society of Scotland, vol. xi. p. 169.
- Achar. Acad. Handl.* 797. Acharius in Königl. Vetenskaps-Academiens Handlingar. 1741, and continued. 8vo.
- Ach. Rich. Monogr.* 1831. Richard (A.), Monographie de la Famille des Eléagnées. Paris, 1832. 4to.
- Ach. Syn.* 1976. Erik Acharius, Synopsis Methodica Lichenum, sistens omnes hujus Ordinis naturalis detectas Plantas, quas, secundum Genera, Species, et Varietates, disposuit, &c. Lundæ, 1814. 8vo.
- A Collection of Roses from Nature.* 749. A Collection, &c., by Miss Lawrence. Lond. 1799. fol. pl. 90.
- Act. Haarlem.* 1276. Verhandelingen uitgegeeven door de Hollandse Maatschappij der Weetenschappen te Haarlem. Haarlem, 1754, and continued. 8vo.
- Act. Nat. Sernt. Berl.* 471. Beschäftigungen der Berlinischer Gesellschaft Naturforschender Freunde. Berlin, 1775, and continued. 8vo.
- Act. Petrop.* 1291. Commentarii Academiæ Scientiarum Imperialis Petropolitanae. Petropoli, 1728—1751. 14 vols. 4to.
- Act. Soc. Batar.* 2055. Verhandelingen van het Bataafsche Genootschap der Proefondervindelyke wysbegeerte te Rotterdam. Rotterdam, 1774, and continued. 4to.
- Act. Succ.* 1342. Acta Literaria Succia. Upsal, 1730, &c. 4to.
- Adans. Fam.* 292. Familles des Plantes, &c. By Michael Adanson. Paris, 1763. 2 vols. 8vo.
- A Description of the Genus Pinus.* 2104. A Description, &c. By Aylmer Bourke Lambert, F.R.S., Pres. of the Lin. Soc., &c. Lond. 1803. folio. A second volume in 1829, folio; and a new edition in 1832, in 2 vols. 8vo; a third vol. in 1836, in folio.

- Æncid.* 962. Virgil's *Æneid*. Dryden's Translation. Carey's ed. Lond. 1806. 3 vols. 8vo.
- Afbeeldingen der Fraaiste, Meestwitheemsche Boomen en Heesters*, &c. 189. By J. C. Krause. Amsterdam, 1802. Royal 4to, pl. 120.
- Afz. Ros. Succ. Tent.* 769. Afzelius (Ad.), *De Rosis Suecanis Tentamina*. Upsal, 1804—1807. 4to.
- Agric. of Hamp.* 1798. A General View of the Agriculture of Hampshire, including the Isle of Wight. By Charles Vancouver. London, 1813. 8vo.
- Agricult. Manual.* 2157. The Agriculturist's Manual; being a familiar Description of the Agricultural Plants cultivated in Europe, including Practical Observations, &c. By P. Lawson and Son. Edinb. 1836. 8vo.
- Agricul. of Clydesdale.* 93. J. Naismith's Agricultural Survey of the County of Clydesdale. Lond. 1806. 8vo.
- Agricultural Report.* 2327. See *Napier's Colonies*.
- Agricultural Report for the County of Durham.* 1404. Bailey's General View of the Agriculture of Durham, &c. Lond. 1811. 8vo.
- Ait. Hort. Kew.* 1892. W. Aiton's *Hortus Kewensis*. A Catalogue of Plants Cultivated in the Royal Gardens at Kew. Lond. 1810. 5 vols. 8vo.
- Alb. Acad. Ann.* 237. Albinus (B.), *Academicarum Annotationum Libri 8*. Leyden, 1754—1783. 4to.
- A Letter on the Acacia.* 615. By Dr. Pulteney, published in the Gentleman's Magazine for 1801, p. 1098.
- Alex. Trav.* 2467. Travels from India to England. By J. E. Alexander. Lond. 1827. 4to.
- Allgem. Gartenzeit.* 1211. Allgemeine Gartenzeitung. By F. Otto and Dr. Dietrich. Berlin. Begun in 1832, and continued.
- All. Ped.* 233. Allioni (C.), *Flora Pedemontana, sive Enumeratio methodica Stirpium indigenarum Pedemontii*. Aug. Taur. 1785. 3 vols. fol. pl. 92.
- All's Well that Ends Well.* 486. One of Shakespeare's plays.
- Alp. Eg.* 526. Prosper Alpinus, *De Plantis Egypti Liber*. Venetiis, 1592. 4to.
- Alp. Exot.* 430. Alpinus (P.), *De Plantis Exoticis Libri 2*. Edidit (A.) Alpinus. Venetiis, 1629. 4to.
- Alpin. Exot.* 1312. Id. *De Plantis Exoticis, libri duo*. Venetiis, 1629. 4to.
- Am. Acad.* 1706. *Amœnitates Academicæ seu Dissertationes variae Physicæ, &c.* By Linnæus and his Pupils. Erlangen, 1790. 10 vols.
- Amœnitates Exoticæ.* 2105. See *Kiempfer's*, &c.
- Aman. Quer.* 1741. See *Amœnitates Quercæ*.
- Amœnitates Quercæ.* 1722. By the late Professor Burnet, published in Nos. 5. and 6. of Burgess's *Eidodendron*. 1833. folio.
- American Gardener's Magazine.* 1357. The American Gardener's Magazine and Register of all useful Discoveries and Improvements in Horticulture and Rural Affairs. Conducted by C. M. Hovey and P. B. Hovey, jun. Boston. Begun 1835, and continued in monthly Nos. 8vo.
- American Museum.* 283. The American Museum, a Repository of Fugitive Pieces in Verse and Prose. Philadelphia, 1790—1792. 8vo.
- American Ornithology.* 86. By Alexander Wilson. Lond. 1832. 3 vols. 8vo.
- Amer. Lyc. N. H. of New York.* 704. Annals of the Lyceum of Natural History of New York. New York, 1824, and continued. 8vo.
- American Sylva.* 189. An English translation of Michaux's *Histoire des Arbres Forestiers de l'Amérique Septentrionale*. Paris, 1817.
- Ann. Ruth.* 631. See *Ammann's Stirp. Ruth.*
- Ammann Stirp. Ruth.* 683. Ammann (Johan.), *Stirpium rariorum in Imperio Rutheno sponte provenientium Icones et Descriptiones*. Petrop. 1739. 4to.
- Annulet.* 1830. An annual publication, edited by S. C. Hall. Lond. 1830. 12mo.
- Anales de las Cienc. Nat.* 1942. See *Anales de las Ciencias Naturales*.
- Anales de las Ciencias Naturales.* 1945. Madrid, 1798—1804. 7 vols. En la Imprenta Real.
- An Autumn near the Rhine.* 2115. Lond. 8vo.
- Anderson MS.* 1552. Geo. Anderson, Esq., Tooley Street and North Mimms, Herts.
- Andr. Bot. Rep.* 240. The Botanist's Repository for New and Rare Plants. By H. C. Andrews. Lond. 1797, et seq. 10 vols. 4to.
- Andr. Heaths.* 1081. Coloured Engravings of Heaths, with botanical descriptions. Id. Lond. 1802—1809. 3 vols. fol.
- An. Hort. Soc. Par.* 278. See *Annales de la Société d'Hort.*, &c.
- Anleit. die Bäume und Sträucher Oesterreichs*, &c. 2201. Gemeinfaßliche Anleitung die Bäume und Sträucher Oesterreichs aus den Blättern zu erkennen. Von Franz Höss. Vienna, 1830. 12mo.
- Annales de Chimie.* 1459. Annales de Chimie et de Physique. Par M.M. Gay-Lussac et Arago. Commenced in 1816. 8vo.
- Ann. d'Hort.* 2403. See *Annales de la Société d'Horticulture de Paris*.
- Annales d'Hort. de Paris.* 138. See *Annales de la Société d'Horticulture de Paris*.
- Annales de la Société d'Horticulture de Paris.* 143. In monthly Nos. 8vo. 1827, and continued; amounting, in 1837, to 21 vols. 8vo.
- Annales de Fromont.* 1359. Annales Horticoles de Fromont. Paris. In monthly Nos. 8vo. Commenced in 1829.
- Annales de l'Institut de Fromont.* 140. See *Annales de Fromont*.
- Annales de la Soc. Entomol. de France.* 1818. Annales de la Société Entomologique de France. Commenced in 1832. 8vo.
- Annales du Musé.* 2485. Annales du Muséum d'Histoire Naturelle. Paris. 8vo. Begun in 1802, and continued.
- Annales du Muséon.* 135. See *Annales du Musé.*
- Annales du Muséum de Paris.* 2104. See *Annales du Musé.*
- Annales Forestières.* 1397. Annales Forestières; faisant suite au Mémorial Forestier. Paris. Begun in 1808, and continued. 8vo.
- Annals of the Muséum.* 2104. See *Annales du Musé.*
- Ann. du. Mus. d'Hist. Nat. de Paris.* 424. See *Annales du Musé.*
- Ann. Lyc.* 661. See *Annals of the Lyceum of Natural History of New York*.
- Ann. of Bot.* 1534. Annals of Botany. By C. König and J. Sims. Lond. 1805, 1806. 2 vols. 8vo.
- Ann. Sc. Nat.* 714. See *Annales des Sciences*, &c.
- An Ode Thrice newly reviv'd*, &c. 836. A black letter tract, published in 1612.
- Anthonne's (M. d') Cypologie du Chêne rouge*, in the *Nouv. Journ. de Physique*. 1823. See *Nouv. Journ.*, &c.
- Antiquary*, 2021. One of Sir Walter Scott's Novels.

- Antiquities of the Common People.* 511. Bourne's (H.) *Antiquitates Vulgares, or Antiquities of the Common People.* Newcastle, 1725. 8vo.
- Ant. of the Jews.* 528. Hydriotaphia, &c., together with the Garden of Cyrus, or Quineuncial, Lozenge, or Network Plantation, of the Ancients, artificially, naturally, and mystically considered. By Sir Thomas Browne, M.D. Lond. 1684. 8vo.
- A Peasant's Voice to Landowners, on the best Means of benefiting Agricultural Labourers, &c.* 1888. By Mr. John Denson, sen. Cambridge and Lond. 1830. 8vo.
- Apollonius Rhodius.* 1795. Apollonii Rhodii Argonautica. 1st ed. Florent. 1496—1498. 4to.
- Appendix to Captain Franklin's Tour to the North Pole.* 2311. By Dr. Richardson and J. Sabine, Esq. Lond. 1823. 4to.
- Archæologia.* 23. Archæologia; or, Miscellaneous Tracts relating to Antiquity. Lond. 1779—1824. 20 vols. 4to.
- Archæol. Attic.* 1724. Rous (F.), *Archæologia Atticæ Libri vii.* Oxon. 1637. 4to.
- Archery Revived.* 2084. See *Shotterel and Durfey's Archery, &c.*
- Archives annuelles de la Normandie.* 1774. Caen, 1824.
- Arch. Mag.* 23. The Architectural Magazine, &c. Conducted by J. C. Loudon, F.L.S., &c. Lond. 1834. In monthly Nos. 8vo, continued.
- Art. Mem.* 533. Pietro Arduini, Memoria di Osservazioni ed Sperienze sopra la Coltura e gli Usi di varie Piante. Padova, 1776. 4to.
- Arnol's Hist. of Edin.* 95. Arnot's History of Edinburgh. Edin. 1779. 4to.
- Asmolean Appendix to Ray's Historia Insectorum.* 1480. See *Ray.*
- Asiat. Research.* 1930. Asiatic Researches; or, Transactions of the Society instituted in Bengal, for enquiring into the History, the Antiquities, the Arts and Sciences, and Literature of Asia. Calcutta, 1788, continued.
- As. Jour.* 2111. The Asiatic Journal and Monthly Register for British and Foreign India, China, and Australia. In monthly Nos. 8vo
- Asso Ar.* 587. See *Asso Syn. Arr.*
- Asso Syn. Arr.* 534. Ignatius De Asso, Synopsis Stirpium indigenarum Arragoniæ. Marsilia, 1779. 4to.
- Athenæum.* 1774. A weekly publication, on Literature and Art. Lond.
- A Topographical and Historical Description of Suffolk.* 1770. 1829.
- A Treatise on the Common Acacia.* 614. Bordeaux, 1763. Pamp.
- Actuarium ad Floram Pedemontanam.* 973. Carolus Allioni, Taurini, 1789. 8vo. 1 fasc.
- Audubon's Birds of America.* 1729. The Birds of America, &c., from Drawings made during a Residence of Twenty-five Years in the United States and its Territories. By J. J. Audubon. 1828, &c. Atlas folio.

B.

- Bailey's Survey of Durham.* 2320. See *Agricultural Report for the County of Durham.*
- Balb. Cat. Taur.* 690. Balbis (J. B.), *Catalogi Horti Botanici Taurinensis, 1805—1813.* 8vo.
- Banist. Cat. Stirp. Virg.* 1864. Herbarium Virginianum; or, an Account of such Plants as J. Banister sent the Designs of to the Bishop of London published in James Petiver's Memoirs for the Curious. 1767.
- Banks. Herb.* 1149. The Banksian Herbarium, now in the British Museum.
- Banks Icon. Kæmpf.* 1200. Josephus Banks, Icones selectæ Plantarum quas in Japonia collegit et delineavit E. Kæmpfer. Lond. 1791. Fol.
- Barrel. Icon.* 322. See *Barrelier Plantæ, &c.*
- Barrelier Observatæ, &c.* 2402. See *Barrelier Plantæ, &c.*
- Barrelier Plantæ per Galliam, &c.* 2402. Plantæ per Galliam Hispaniam et Italiam observatæ. By Jacob Barrelier. Paris, 1714. Fol.
- Barl. Fl. Vir.* 503. Barton (W. P. C.), A Flora of North America. Philadelphia, 1831. 8vo.
- Barl. Ord. Nat.* 2509. Barthing (F. T.), Ordines Naturales Plantarum, eorumque Characteres et Affinitates; adjectâ Generumque Enumeratione. Götting. 1830. 8vo.
- Bartram's Travels.* 277. Travels through South and North Carolina. By William Bartram. Philadelphia, 1791. 1 vol. 8vo.
- Bartram Trav., ed Germ.* 995. William Bartram's Riesen durch Nord- und Sud-Carolina, aus dem Englischen von E. A. W. Zimmermann. Berol. 1793. 8vo.
- Bast. Ess.* 768. Essai sur la Flore du Département de Maine et Loire. By T. Bastard, or Batard. Angers, 1809. 12mo.
- Bast. Suppl.* 768. See *Batard Suppl.*
- Batard Suppl.* 763. Supplément à l'Essai sur la Flore du Département de Maine et Loire. Id. Angers, 1812. 1 fasc., 12mo.
- Bat. Fl. Main.* 767. See *Bast. Ess.*
- Bat. Fl. Main. et Loire Suppl.* 772. See *Batard Suppl.*
- Bath Soc. Pap.* 840. See *Bath Society's Transactions.*
- Bath Soc. Papers.* 1641. See *Bath Society's Transactions.*
- Bath Agricultural Society's Papers.* 1550. See *Bath Society's Transactions.*
- Bath Society's Transactions.* 2381. Letters and Papers of the Bath Agricultural Society, &c. Bath, 1792, continued. 8vo. 15 vols. to 1838.
- Batsch. Conf.* 2148. Flenchus Fungorum. By A. J. G. C. Batsch. Halle. 3 fasc. i, 1783; ii, 1786; iii, 1789. 4to.
- Baudrill. Traités, &c.* 421. Baudrillart, one of the authors of *Traité Général des Eaux et Forêts, Chasses, et Pêches.* Paris, 1825. 5 vols. 4to.
- Bauh. (J.) Hist.* 546. Historia Plantarum universalis. By J. Bauhin. Ebroduni. 1650, 1651. 3 vols. folio.
- Bauh. Pin.* 297. Pinax Theatri Botanici. By C. Bauhin. Basileæ. Ed. 1., 1623; ed. 2., 1674. 4to.
- Baum. Cat.* 675. *Catalogi Horti Bollwylleriani.* By the brothers Baumann. 1810—1814. Baumann's Catalogue for 1838 forms an article in an Appendix to this Arboretum.
- Baxt. Brit. Fl. Pl.* 829. British Pænogamous Botany; or, Figures and Descriptions of the Genera of British Flowering Plants. By W. Baxter, A.L.S., Curator of the Oxford Botanic Garden. Oxford. In monthly Nos. 8vo. 5 vols. published in 1837.
- Beauties of England, Dorsetshire.* 36. See *Beauties of England and Wales.*
- Beauties of England, Kent.* 24. See *Beauties of England and Wales.*
- Beauties of England and Wales, Shropshire.* 1768. The Beauties of England and Wales; or, Deline-

- ations, Topographical, Historical, and Descriptive, of each County. By John Britton, F.S.A. &c. and E. M. Brayley. Lond. Begun in 1801, and a volume published yearly. 8vo.
- Bechst. Forst.* 767. Bechstein (J. M.), Forstbotanik, &c. Gotha, 1821. 8vo.
- Bechstein et Scharffenburg Forst. Ins.* 1818. Bechstein and Scharffenburg, Vollständige Naturgeschichte der Schädlichen Forst Insecten. Leipsig, 1805. 2 vols. 4to.
- Beckman's Hist.* 464. See *Beckman's History of Inventions*.
- Beckman's Hist. of Invent.* 1914. History of Inventions of Discoveries. By John Beckman. Translated by Johnson. Lond. 1797. 3 vols. 8vo.
- Bedf. Hort. Eric. Woburn.* 1079. See *Hort. Eric. Woburn*.
- Beech. Voy. Pt. Bot.* 1037. Voyage to the Pacific and Behring's Straits, to cooperate with the Polar Expeditions in 1825—1828. Lond. 1831. In 2 parts. 4to.
- Belknap's History of New Hampshire.* 2282. Belknap (Jeremy), Description of the White Mountains in New Hampshire, in *Transact. of the Amer. Phil. Soc.*, vol. ii. Philadelphia, 1786. 4to.
- Bell. App. Fl. Pedem.* 739. See *Bellardi App. ad Fl. Ped.*
- Bellardi App. ad Fl. Ped.* 1555. Appendix ad Floram Pedemontanam. 1790 and 1791. 8vo.
- Bell. Conifer.* 2274. De Arboribus Coniferis resiniferis aliisque sempiterna fronde virentibus. By Pierre Belon, or Bellon. Paris, 1553. 4to.
- Bell. Gall.* 1725. Cæsar (Julius), Commentaria de Bello Gallico.
- Bellon. Obs.* 2495. Bellonii Observaciones in Clusii exoticarum Libri x. 1605. Fol. pl. 237.
- Berg. Pl. Cap.* 1080. Descriptiones Plantarum ex Capite Bonæ Spei. By P. J. Bergius. Stockholm, 1767. 8vo.
- Berk. Eng. Fl.* 1976. Volume V. Part II. of Smith's English Flora. By the Rev. M. J. Berkeley, M.A. Lond. 1836. 8vo.
- Berl. MSS.* 972. Berlandier MSS.
- Berlinische Baumzucht.* 100. Berlinische Baumzucht, oder Beschreibung der in Königlichen Botanischen Garten bei Berlin im Freien ausdauernden Bäume und Sträucher, &c. Von Carl Ludwig Willdenow. Berlin, 1811. 8vo.
- Bert. Pl.* 583. See *Bertol. Pl. Gen.*
- Bertol. Pl. Gen.* 599. Plantæ Genuenses quas annis 1802, 1803 observavit et recensuit. By A. Bertoloni. Genue, 1804. 8vo.
- Bertoloni's Flora Italica.* 164. Bertolonii (A.) Flora Italica. Bononia, 1833—1835. 8vo.
- Besl. Eyst.* 1209. See *Besl. Hort. Eyst.*
- Besl. Hort. Eyst. Fern. Ord.* 754. Hortus Eystettensis. By Basilius Besler. Nuremberg, 1612. 2 vols. fol.
- Bess. Cat. Hort. Crem.* 766. Catalogus Horti Botanici Wolhyniensis Cremenici. 1811—1816. 8vo.
- Besser En. Pl. Volhyn.* 1587. See *Besser Enum. Pl. Volh.*, &c.
- Besser Enum. Cont.* 584. See *Besser Enum. Pl. Pot.*, &c.
- Besser Enum. Pl. Pod.*, &c. 728. See *Besser Enum. Pl. Volh.*
- Bess. Enum. Pl. Volh.* 631. Enumeratio Plantarum Volhyniæ, Podoliæ, &c. By W. J. J. Besser. Vilna, 1822. 8vo.
- Bess. Galic.* 1598. See *Bess. Prim. Fl. Gal.*
- Bess. Prim. Fl. Gal.* 768. Primitiæ Floræ Galiciæ. Id. Vienna, 1809. 2 vols. 12mo.
- Biblia Pauperum.* 1366. A very curious work, published between 1400 and 1450, with 40 woodcuts, illustrated by texts of Scripture.
- Bibliotheca Geoponica.* 187. A posthumous work of the late Mr. Forsyth, not yet published.
- Bibliothèque Physico-économique.* 2408. Bibliothèque Physico-Economique, Instructive, et Amusante. 1794.
- Bibl. Univ.* 2094. See *Bibliothèque Universelle de Genève*
- Bibl. Univ. de Genève.* 2350. See *Bibliothèque Universelle de Genève*.
- Bibliothèque Universelle de Genève.* 162. Geneva, 8vo.
- Bieb. Casp.* 1299. Beschreibung der Länder zwischen den Flüssen Terek und Kur am Caspischen Meere. By L. B. F. Marschal de Bieberstein. Frankfurt am Mayn, 1800. 8vo.
- Bieb. Cent. Pl. Rar.* 1058. Centuria Plantarum rariorum Rossæ meridionalis, præsertim Tauriæ et Caucasi. Id. Charkov. 1810. fol.
- Bieb. Fl. Taur.-Cauc.* 339. Flora Taurico-Caucasica, exhibens stirpes phanogamas in Chersoneso-Taurica et Regionibus Caucasica sponte crescentes. Id. Charkov. 1808—1819. 3 vols. 8vo.
- Bieb. Fl. Taur. Suppl.* 499. See *Bieb. Suppl.*
- Bieb. Suppl.* 661. Flora Taurico-Caucasica. Vol. III, seu Supplementum. Id. Charkov. 1819. 8vo.
- Bigelow's Account of the Plants of Boston and its Vicinity.* 1650. 1854.
- Bigelow's Amer. Bot.* 1019. American Medical Botany. By J. Bigelow. Boston, 1817—1821. 8vo.
- Bigel. Fl. Bost.* 974. Flora Bostoniensis. Id. Boston, 1814. 8vo.
- Bigel. Med. Bot.* 1017. See *Bigelow's Amer. Bot.*
- Bigelow's Med. Bot.* 268. See *Bigelow's Amer. Bot.*
- Big. Med. Bot.* 552. See *Bigelow's Amer. Bot.*
- Birkbeck's Notes, &c.* 1867. Morris Birkbeck's Notes on a Journey in America, from the Coast of Virginia to the Territory of Illinois. London, 1819. 8vo.
- Biv. St. Sic. Mant.* 582. Stirpium rariorum in Sicilia proventium Manip. Panorm. 1813—1816. 4 vols. 8vo.
- Black. Herb.* 314. A curious Herbal; containing 600 Cuts of the Useful Plants. By Elizabeth Blackwell. London, 1757. 2 vols. fol.
- Black. Icon.* 506. See *Black. Herb.*
- Blomfield's Norfolk.* 33. Essay towards a Topographical History of the County of Norfolk, by the Rev. F. Blomfield. 1799—1810. 11 vols. 8vo.
- Bluff et Fing. Fl. Germ.* 1540. Bluff et Fingerhuth's Floræ Germaniæ. Norimb. 1825. 12mo.
- Blum. Bjdr.* 569. See *Blume Bjdr. Fl. Ind.*
- Blum. Bjdr. Fl. Ind.* 295. Bjdragen tot de Flora van nederlandse Indië. By C. L. Blume. Batavia, 1825, 1826.
- Blume Fl. Jav.* 1936. Floræ Javæ et Insularum adjacentium. By C. L. Blume and J. B. Fischer. Bruxellis, 1828. Fol.
- Bobart's Hist. Pl. Oxon.* 53. Catalogus Horti Botanici Oxoniensis, &c. By D. Bobart and W. Broune, A.M. Oxonii, 1658. 16mo.
- Bocc. Muss.* 545. Museo di Piantæ rare della Sicilia, Malta, Corsica, Italia, Piemonte, e Germania. By Paulo Boccone. Venetia, 1697. 4to.
- Bocc. Sic.* 316. Icones et Descriptiones rariorum Plantarum Sicilia, Melitæ, Galliæ, et Italiæ. Id. Lond. 1674. 4to.

- Boerh. Ind. Alt.* 67. Index alter Plantarum quæ in Horto Acad. Lugduno-Batavo aluntur. By H. Boerhaave. Lugd. Bat. 1720. 2 vols. 4to.
- Boerh. Lugd.* 469. Index Plantarum quæ in Horto Acad. Lugduno-Batavo reperuntur. Id. Lugd. Ba. 1710. 4to.
- Boltwyler Catalogue.* 742. See *Baum. Cat.*
- Bolt. Fun.* 1823. An History of Funguses growing about Halifax. By James Bolton. Huddersfield, 1788, 1789. 3 pts. Append. 1791. 4to.
- Bonafous's Mémoire sur une Education de l'ers à Soie,* &c. 1358. Paris.
- Bonod on Mulberries.* 1355. Bonoed (John), A Treatise on the Art of making Silk: together with Instructions how to plant Vines. London, 1699. 4to.
- Bon Jard.* 443. Le Bon Jardinier, contenant les Principes généraux de Culture, &c. Paris, 12mo. A volume yearly.
- Bonninghausen Fl. Monaster.* 151. Bonninghausen (C. M. F.), Prodrômus Floræ Monasteriensis. Westphalorum. Monast. 1824. 8vo.
- Bonpl. Malmaison* 262. See *Bonpl. Pl. Rar.*
- Bonpl. Pl. Rar.* 251. Description des Plantes rares cultivées à Malmaison et à Navarre. By A. Bonpland. Paris, 1813—1816. Fol.
- Bon Sens.* 1996. A French periodical.
- Book of Trades.* 1336. Frankfurt, 1334.
- Bork. Holz.* 754. Beschreibung der in den Hessen-Darmstädtischen Lauden im Freien wachsenden Holzarten. By M. B. Borkhausen. Franef. Main, 1790. 8vo.
- Bosc Act. Soc. Hist. Nat. Par.* 955. Actes de la Société d'Histoire Naturelle de Paris. Paris, 1792. Fol.
- Bosc Dict. d'Agric.* 732. Nouveau Cours complet d'Agriculture théorique et pratique. New edition. Paris, 1821—1823. 16 vols. 8vo.
- Bosc Mém. sur les Chênes.* 1847. Mémoires sur les différentes espèces de Chênes qui croissent en France. By L. Bosc. Paris, 1808. 4to.
- Botanicum Menseliense.* Botanicum Menseliense. By P. Magnol. Menselii, 1686. 12mo.
- Botanist.* 1172. The Botanist. Conducted by B. Maund, Esq., F.L.S., assisted by Professor Henslow. In monthly Nos. 8vo.
- Botaniste Cultivateur; or Description, Culture, and Use of the greatest Part of the Plants, Foreign and Indigenous, which are cultivated in France and England, according to the Method of Jussieu,* 143. By Dumont De Courset. Paris, 1802, 5 vols. 8vo.; ed. 2. Paris, 1811, 6 vols. 8vo. Supplement on vol. vii. 1814.
- Botanologia Universalis Hibernica.* 107. By J. Kéoch, A. B. Corke, 1735. 4to.
- Bot. Arrangem.* 700. A Botanical Arrangement of all the Vegetables naturally growing in Great Britain, &c. By W. Withering. Birmingham, 1776, 2 vols. 8vo.; 3d ed. greatly enlarged, Lond. 1790, 4 vols. 8vo.
- Bot. Cab.* 339. See *Lodd. Bot. Cab.*
- Bot. Cult.* 1497. See *Botanist Cultivateur.*
- Bot. Gard.* 254. See *Maund's Bot. Gard.*
- Bot. Mag.* 98. See *Curt Bot Mag.*
- Bot. Reg.* 250. Edwards's Botanical Register. Lond. 8vo. Begun in 1815, and continued monthly. Conducted now by Dr. Ländley.
- Bot. Rep.* 229. See *Andr. Bot. Rep.*
- Bouché.* Naturgeschichte der Schädlichen und Nützlichen Garten Insecten. Berlin, 1833. 12mo.
- Boucher's Treatise on raising Forest Trees.* 105. Edinb. 1775. 4to.
- Boyce's Account of the Parish of Kensington.* 47. 1705.
- Boyceau's Jardinage,* &c. 1338. J. Boyceau, Traité au Jardinage, &c. Paris, 1714. Fol.
- Bradley's New Improvements of Planting and Gardening.* 98. Lond. 1718. 8vo.
- Bradley's Treat. on Husbandry and Gardening.* 98. A Treatise on Husbandry and Gardening, &c. By Richard Bradley, F.R.S. Lond. 1721. 8vo.
- Bramham Park.* 2009. A poem by Fawkes.
- Brande's Journ.* 124. A Journal of Travels in Peru, &c. By C. Brande. Lond. 8vo.
- Brand's Popular Antiquities.* 900. Brand's (Rev. J.), Observations on Popular Antiquities. Newcastle, 1777. 8vo.
- Bray's (Mss.) Borders of the Tamar and the Tavy.* 900. Description of the Part of Devonshire bordering on the Tamar and Tavy; its Nat. Hist., &c. London, 1836. 3 vols. 8vo.
- Brewster's Edin. Journ.* 124. See *Ed. Phil. Journ.*
- Breyne Cent.* 601. Icontearium Plantarum Centuria, By Jacobus Breynius, Gedani, 1678. Fol.
- Breyne Icon.* 1259. Icones rariorum Plantarum, conjunctum cum priori a J. Ph. Breynio editæ. 1789. 4to.
- Breyne Obs.* 2274. Prodrômus Fasciculi rariorum Plantarum in Hortis Hollandiæ observatarum. By Jacobus Breynius. Gedani, vol. i. 1680; vol. ii. 1689. 2 vols. 8vo.
- Bright's Travels in Hungary.* 369. Dr. Bright's Travels from Vienna, through Lower Hungary, in 1815. Edinb. 1818. 4to.
- Brit Ent.* 1876. See *Curtis Brit. Ent.*
- Brit. Fungi.* 2147. Consisting of dried Specimens of the Species described in Vol. V. Pt. II. of the English Flora, together with such as may hereafter be discovered indigenous to Britain. By the Rev. M. J. Berkeley, A. M. Lond. 1876. Fasc. 1. and 2.
- Britton's Wiltshire.* 128. Wilt-hire. See *Beauties of England.*
- Brit. Topog.* 31. Gough (R.), Anecdotes of British Topography; or, an Historical Account of what has been done for illustrating the Topographical Antiquities of Great Britain and Ireland. London, 1768. 4to.
- Bromel. Chl. Goth.* 1706. Chloris Gothica. By Olaus Bromelius. Gothoburgi, 1691. 8vo.
- Brong. Mém. Rham.* 524. Mémoire sur la Famille des Rhamnées; ou Histoire Naturelle et Médicale des Genres qui composent ce Group de Plantes. Par Adolphe Brongniart. Paris, 1826. 4to.
- Brota Flor. Lus.* 242. See *Brotero's Flora Lusitanica.*
- Brotero's Pl. Lus.* 1926. See *Brotero's Flora Lusitanica.*
- Brotero's Flora Lusitanica.* 194. Olisiponæ, 1804. 2 vols. 8vo.
- Brota Phyt.* 580. Phytographia Lusitania selector. By Felix Avellar Brotero. Olisiponæ, 1801. Fol.
- Brown's Edit of White's Selborne.* 1578. White's Natural History of Selborne, and Observations on various Parts of Nature; with the Naturalist's Calendar, and notes, by Capt. Thomas Brown, F.L.S. London, 1833. 8vo.

- Brown Prod.* 1307. Prodrum Floræ Novæ Hollandiæ et Insulæ Van Diemen. By Robert Brown. Lond. 1810. 8vo.
- Browne's Journ.* 1129. The Civil and Natural History of Jamaica. By Patrick Browne. Lond. 1756. Fol. Ed. 2. 1789.
- Brug. Voy.* 2410. Bruguère (J. G.), Voyages dans l'Empire Ottoman. Paris. 3 vols. 4to.
- Butbul Nameh.* 792. A curious fragment, by the Persian poet Attar.
- Bulletin des Sciences.* 1999.
- Bulletin des Sciences de la Société Philomatique.* 2104. Bulletin des Sciences, par la Société Philomatique de Paris. Paris, from 1791.
- Bull. de la Soc. d'Ag. du Département de l'Hérault.* 2095. Bulletins de la Société d'Agriculture du Département de l'Hérault, s'étant à Montpellier. Montpellier, 1803—1813. 8vo.
- Bulletin Universel.* 1133. Bulletin Universel des Sciences et de l'Industrie. Paris, 1825. 24 vols. 8vo.
- Bull. Fl. Par.* 1082. Flora Parisiensis. By Bulliard. Paris, 1776—1780. 5 vols. 8vo.
- Bull. Fr.* 365. Herbar de la France. Id. 1780, et seq. Fol. pl. 600.
- Bull. Herb.* 581. See *Bull. Fr.*
- Bulliard's Hist. des Champ. de France.* 1975. Histoire des Champignons de la France. Id. Paris, 1798. Fol.
- Bull. Sci.* 587. See *Bulletin des Sciences, &c.*
- Bunge Mém. Acad. Scien. Petersb.* 1936. Dr. Bunge in Mem., &c. See *Mém. Acad. Scien., &c.*
- Bunge's List of Plants of the North of China.* 432. Enumeratio Plantarum quas in China boreali collegit Dr. Al. Bunge. 1831. Published in the Mem. Acad. Scien. Petersb., &c., 1833.
- Burgess's Eidodendron.* 1722. Burgess's (H. W.), Eidodendron. Views of the General Character and Appearance of Trees, Foreign and Indigenous. Lond. 1827. Folio.
- Burgsd. Adelit.* 1657. Burgsdorf (F. A. L.), Anleitung zur sichern Erziehung und zweckmässigen Anpflanzung der einheimischen und fremden Holzarten welche in Deutschland und unter ähnlichen Klima im freien Fortkommen. 1795. 2 vols. 8vo.
- Burmeister Handb. der Ent.* 1862. Burmeister (Dr. H.), Manual of Entomology. Translated from the German by W. E. Shuckard. London, 1835. 8vo.
- Burm. Fl. Ind.* 602. N. L. Burmann's Flora Indica. Lugd. Bat. 1768. 4to.
- Burm. Prod.* 2059. See *Burm. Fl. Ind.*
- Burm. Rar. Plant. Afr.* 504. Rariorum Africanarum Plantarum Decades 10. By J. Burmann. Amsterdami, 1758, 1739. 4to.
- Burnett's Amœn. Quer.* 1837. See *Amœnitates Quercæ.*
- Burnett's Inaugural Address to the Medico-Botanical Society.* 1459. Lond. 1831.
- Burton's Monast.* 2073. Burton, J., Monasticon Eboracense: and the Ecclesiastical History of Yorkshire. York, 1758. Folio.
- Buzb. Cent.* 319. J. C. Buxbaum, Plantarum minus cognitarum Centuriæ quinquæ. Petropoli, 1728—1740. 5 vols. 4to.
- Buzb. Hal.* 1397. Id., Enumeratio Plantarum in Agro Hallense crescentium. Halæ, 1721. 8vo.

C.

- Cæsar's Commentaries.* 21.
- Calendarium Hortense.* 44.; or Gardener's Almanack. By John Evelyn. Lond. 1664, 1669, 1676, 1706. 8vo.
- Camb. Ann. Sci. Nat.* 722. See *Cambessedes Mon. Spir.*, &c.
- Camb. Bot.* 1046. Enumeratio Plantarum quas in Insula Balcaribus collegit J. Cambessedes. Published in the Mémoires du Muséum, tom. xiv. 1857.
- Cambessedes Mon. Spir. in Ann. Sci. Nat.* 72. Cambessedes's Monograph of the Spiræas, in the Annales des Sciences Naturelles, tom. 1.
- Camden's Account of the New Forest.* 1761. See *Camden's Britannia.*
- Camden's Britannia.* 1391. Britannia, sive florentissimorum Regnorum Angliæ, &c., Chorographica Descriptio. Lond. 1697. Fol. Translated by Edward Gibson. 1722. 2 vols. fol.
- Camden's Life of Queen Elizabeth.* 948. Camden's (W.), Life of Queen Elizabeth. Oxford, 1717. 5 vols. 8vo.
- Can. Epit.* 1010. Joachin Camerarius, De Plantis Epitome utilissima. Francof. Men. 1586. 4to.
- Camellieæ Britannicæ.* 386. See *Chandl. Cam. Brit.*
- Camer. Hort.* 2063. Hortus Medicus et Philosophicus. Id. Francof. Men. 1588. 4to.
- Carrington's Dartmoor.* 1757. A poem. 8vo.
- Carstele's Traité sur le Mûrier blanc.* 1538. Paris.
- Catal. of Brit. Plants.* 729. Catalogue of British Plants. By Professor Henslow. 8vo. pamph.
- Catalogue of the Leyden Garden.* Boerhaave (H.), Index Plantarum quæ in Horto Acad. Lugd.-Bat. reperiuntur. Lugd. Bat. 1710. 8vo.
- Catal. of Tradescant's Museum.* Museum Tradescantianum; or a Collection of Rarities preserved at South Lambeth. By John Tradescant. 1656. 12mo
- Catal. of Trees and Shrubs, both Exotic and Domestic, which are propagated for Sale in the Gardens near London.* 60. By the Society of Gardeners. London, 1730. Fol.
- Catalogue.* 44. Catalogue of Plants observed in North America. By John Banister. 1680.
- Catalogus Plantarum rariorum Angliæ et Insularum adjacentium.* 1001. By John Ray. Lond. 1670. Ed. 2. 1677. 8vo.
- Catesb. Carol.* 297. Catesby's Natural History of Carolina, Florida, and the Bahama Islands. Lond. 1741—1743. 2 vols. folio.
- Cat. Hort. Vindob.* 674. See *Jacq. Hort. Vin.*
- Can. Anal.* 579. See *Anal. de las Venetas Naturales.*
- Can. Diss.* 561. A. J. Cavanilles, Monadelphicæ Classis Dissertationes Decem. Matrit. 1790. 4to. pl. 297.
- Can. Ic.* 312. Id., Icones et Descriptiones Plantarum quæ aut sponte in Hispania crescent, aut in Hortis hospitantur. Matrit. 1791—1801. 6 vols. fol. pl. 601.
- Celtic Researches and Inquiry into the Mythology of the Druids.* 1753.
- Chab. Scieg.* 586. Dominicus Chabrey, Omnium Stirpium Scægraphia. Coliniæ Allobrogum, 1666. Ed. 2. 1677. Fol.
- Chambers's (Sir W.) Dissertations on Oriental Gardening.* 80. Lond. 1774. 4to.
- Chandl. Cam. Brit.* 387. See *Chandl. Ill.*
- Chandl. Ill.* 582. Illustrations by Alfred Chandler, and Descriptions by W. Beattie Booth, of the Camellieæ. Lond. 1831. Imp. 4to.

- Charlwood's Catalogue*. 1316. A Catalogue, by George Charlwood, Nursery and Seedsman, Covent-Garden. Published yearly.
- Childe Harold* 1917. A poem by Lord Byron.
- Chois Prod. Hyp.* 598. J. D. Choisy, Prodromus d'une Monographie des Hypéricinées. Geneva, 1821. 4to.
- Choul. De var. Querc. Hist.* 1721. Joh. Du Choul, De variâ Quercus Historiâ, accessit *Plati Montis Descriptio*. Lugduni, 1555. 8vo.
- Chron Sax* 1747. See *Saxon Chronicles*.
- Church History of England*. 1753. Dodd (C.), Church History of England, chiefly regarding the Roman Catholics in England, from 1737 to 1742. Brussels, 3 vols. folio.
- Churchill's Medical Botany*. 553. Medical Botany; or, Illustrations and Descriptions of the Medicinal Plants of the London, Edinburgh, and Dublin Pharmacopœias. By J. Stephenson, M.D., &c., and James Morss Churchill, F.L.S. Lond. 1831. 4 vols. 8vo.
- Cider*. 901. A poem by Phillips.
- Cistinea*. 316. See *Sot. Cist*.
- Civil and Natural History of Chili*. 2456. See *Molina's History, &c.*
- Clayt. Flor. Virgin* 516. Gronovius (J. F.), Flora Virginica, exhibens Plantas quas J. Clayton in Virginia collegit. Lugduni Batavorum, 1743. 8vo.
- Clus. Hist.* 323. See *Clusius's Rariorum Plantarum*.
- Clus. Pan.* 1155. Carolus Clusius, Rariorum aliquot Stirpium per Pannoniam observ. Historia. 1583. 8vo.
- Clusius's Rariorum Plantarum*. 147. Id., Rariorum Plantarum Historia. Antverpiæ, 1605. Fol. 2.
- Cobb's Manual of the Mulberry Tree, &c.* Massachusetts.
- Cold. Novc.* 2039. Colden (C.), Plantæ Coldenhamiæ in Provincia Novboracensi Americæ sponte crescentes; in Act. Societ. Upsal. 1743.
- Colla Clout's come Home again*. 1684. A ballad by Spenser.
- Colla Hort. Rip* 473. See *Colla Hort. Ripul.*
- Colla Hort. Ripul.* 863. Colla, Hortus Ripulensis. Turin, 1820—1837. 4to.
- Collection of Poems*. 2022. Dublin, 1801.
- Colonies, &c.* 2327. See *The Colonies*.
- Colum. Ic.* 2519. Columna: F., Phytobasanos, sive Plantarum aliquot Historia, cum Annotationibus Jani Planci. Mediolani, 1774. 4to.
- Com. Dict.* 316. See *McCull. Dict.*
- Com. Hort.* 1074. See *Comm. Hort. Anst.*
- Comm. Goett.* 1982. Commentaria Societatis regię Goettingensis. 1751, 1754, 1769—1816. 4 vols. 4to.
- Comm. Hort. Anst.* 398. Joh. Commelyn, Horti medici Amstelodamensis rariorum Plantarum Descriptio et Icones Pars prima. Amstelod. 1697. Fol. Pars altera. By Casp. Commelyno. 1703. Fol.
- Comm. Præcl.* 2523. Casp. Commelyn, Præudia botanica ad Publicas Plantarum Demonstrationes. Lugd. Bat. 1706. 4to.
- Commentaries*. 189. See *Cæsar's Commentaries*.
- Commons's Complaint, wherein is contained Two special Grievances*. 1751. By Arthur Standish. 1611.
- Communications to the Board of Agriculture*. 2354. London, 1797—1819. 7 vols. 4to. New Series, 8vo.
- Comp. Bot. Mag.* 1269. See *Companion to the Botanical Magazine*.
- Companion to the Botanical Magazine*, 2249; being a Journal containing such interesting botanical information as does not come within the prescribed Limits of the Magazine; with occasional figures. By Sr W. J. Hooker. Lond. Begun in 1835.
- Compend. Fl. Ger.* 1635. Bluff (M. J.) et Fingerhuth (C. A.), Compendium Floræ Germaniæ. Norimb. 1825. 8vo.
- Complete Husbandman*. 1639. By S. Hartlib, London, 1579. 4to.
- Confessio Amantis*. 2017. An old poem by Gower.
- Cook's (Captain J.) First Voyage*. London 1777. 2 vols. 8to.
- Cook's (Captain J.) Second Voyage*. 1442. Cook's Second Voyage. London, 1784. 4to.
- Cord. Hist.* 1168. Cordus (C.), Historia Plantarum, ed. à C. Gesnero. Argentorati, 1651. Folio.
- Coriolanus*. 1724. By Shakspeare.
- Corn. Can.* 1259. Jacob Cornuti, Canadensium Plantarum aliarumque nondum editarum Historia. Paris, 1636. 4to.
- Corsica*. 1118. A poem by Mrs. Barbauld. 1769.
- Cosmography*. 1828. Munster (F.), La Cosmographie Universelle, Paris, 1552. Folio.
- Coun. Rep.* 87. Maitland's (W.) History of Edinburgh, from its Foundation to the present Time. Edinburgh, 1753. Folio.
- Court of Love*. 829. A poem by Chaucer.
- Cowley on Plants*. 1346. A poem.
- Cowper's Yardley Chase* 1785. A poem.
- Crantz Austr.* 339. Henr. Joh. Nepom. Crantz, Stirpes Austriacæ. 1762. 8vo.
- Croker's Edition of Boswell's Johnson*. 1519.
- Cullen's Trans. Clar.* 2051. C. Cullen's History of Mexico, translated from the Italian of Clavigero. London, 1787. 2 vols. 4to.
- Cullum's Hawsted*. 33. History and Antiquities of Hawsted and Hardwick. By the Rev. Sir John Cullum, Bart. 2d ed., with notes. Lond. 1813. 4to.
- Cultivateur Provençal*. 2412. A periodical quoted in Lenseleur Deslongchamps's Histoire du Cèdre du Liban.
- Culture of Forests, &c.* 1992. By Lieutenant-Colonel Emmerich. Lond. 1789. 8vo.
- Cupan. Pamph.* 706. See *Cup. Pamph. Sic.*
- Cup. Pamph. Sic.* 587. Franciscus Cupani, Pamphytum Siculum, seu Historia Plantarum Siciliae. fol.
- Curt. Bot. Mag.* 241. The Botanical Magazine, &c. By W. Curtis; continued by Dr. Sims; and, since the death of Dr. Sims, by Sir W. J. Hooker. Lond. Begun in 1787, and continued in monthly Nos. 8vo.
- Curtis Brit. Ent.* 2142. See *Curtis's British Entomology*.
- Curtis's British Entomology*. 1484. British Entomology; being Illustrations and Descriptions of the Genera of Insects found in Great Britain and Ireland, &c. By J. Curtis, F.L.S., &c. Lond. 1824—1836. 13 vols. 8vo.
- Curt. Fl. Lond.* 313. See *Curt. Lond.*
- Curt. Lond.* 769. Flora Londinensis; or, Plates and Descriptions of such Plants as grow wild in the Environs of London, with their places of Growth, &c. By W. Curtis. Lond. 1777. 3 vols. fol. pl. 432.

D.

- Dalech. Hist.* 1678. See *Dalechamp's Hist. Plant.*
Dalechamp's Hist. Plant. 1845. Jacques Dalechamp's *Historia generalis Plantarum.* Lugd. 1586, 1587. 2 vols. folio.
Dalib. Paris. 1983. T. F. Dalibard, *Floræ Parisiensis Prodomus.* Paris, 1749. 12mo.
Dandolo's Dell'Arte di governare i Buechi da Seta. 1358. Milan.
Davies's Welsh Bards. 899. Edw. Davies's *Celtic Researches*; London, 1801, 8vo: and Edw. Davies's *Rites of the British Druids*; London, 1809, 8vo.
Davies's Welsh Bot. 764. *Welsh Botany.* By H. Davies. Lond. 1813. 8vo.
Davis's (Dennis's) Landscape-Gardener. 129. *The Landscape-Gardener.* By the Rev. Prebendary Dennis. London, 1836. 8vo.
Davy's Agricultural Chemistry. 152. *Elements of Agricultural Chemistry.* By Sir Humphry Davy. In a Course of Lectures for the Board of Agriculture. London, 1813. 4to.
Dean's Croome Guide. 73. 8vo.
De Arboribus Coniferis 2407. See *Bell. Confif.*
De Arboribus Coniferis Resiniferis, &c. 187. See *Bell. Confif.*
De Arborum, Fruticum, et Herbarum, &c. 187. Joann. Meursius, *Arboretum Sacrum, sive de Arborum, Fruticum, et Herbarum Consecratione, Proprietate, &c.* Lugd. Bat. 1642. 8vo, pl. 10.
Death of Acis. 2122. A poem by Barry Cornwall.
De Bell. Gall. 2069. Cæsar (Julius), *Commentaria de Bello Gallico.*
Dec. Astr. 635. Augustin Pyramus De Candolle, *Astragalologia.* Paris, 1802. 4to et fol.
Dec. Fl. Fr. 328. Id. et Lamarck, *Floræ Française.* Paris, 1805—1815. 5 vols. 8vo.
Dec. Fl. Fr. Suppl. 330. See *Dec. Fl. Fr.*
Dec. Hort. Monsp. 472. See *Dec. Cat. Hort. Monsp.*
Dec. Cat. Hort. Monsp. 372. Id., *Catalogus Plantarum Horti Botanici Mospeliensis, addito Observationum circa Species novas aut non satis cognitæ Fasciculo.* Monsp. 1813. 8vo.
Dec. Lég. M. M. 561. Id., *Mémoire sur la Famille des Légumineuses.* Paris, 1825. 4to. pl. 70.
Dec. Mém. Soc. Gen. 503. De Candolle in *Mémoires de la Société de Physique et Histoire Naturelle de Genève.* 1821, and continued. 4to.
Dec. Mém. inéd. in Soc. Phys. Gen. 634. De Candolle's MSS. in the *Mémoires de la Soc. de Phys. et d'Hist. Nat. de Genève.*
Dec. Phys. Vég. 675. See *De Candolle's Physiologie Végétale.*
De Candolle's Physiologie Végétale, 162; ou *Exposition des Forces et des Fonctions vitales des Végétaux.* Id. Paris, 1832. 2 vols. 8vo.
Dec. Pl. rar. Jard. Gen. 2261. Id., *Plantes rares du Jardin de Genève.* Genev. 1825—1827. 4to, pl. 24.
Dec. Prod. 233. *Prodomus Systematis naturalis Regni Vegetabilis; seu Enumeratio methodica Ordinum, Generum, Specierumque Plantarum hucusque cognitarum.* Paris, 1824—1830. 4 vols. 8vo.
Dec. Syst. 234. Id., *Regni vegetabilis Systema naturale; sive Ordines, Genera, et Species Plantarum, secundum Methodi naturales Norinus digestarum et descriptarum.* Paris, 1818—1821. 2 thick vols. 8vo.
De Chabral's Statistiques de Savone, &c. 1854.
De Fruct. et Sem. Plant. 2103 J. Gärtner, *De Fructibus et Seminibus Plantarum; continuat. à C. F. Gärtner sub titulo "Carpologia."* Leipzig, 3 vols. 4to. i., 1788; ii, 1791; iii, or Supplementum *Carpologiae.* 1805.
De l'Agric. du Gatinis, &c. 2158. See *De l'Agriculture du Gatinis.*
De l'Agriculture du Gatinis, &c. 2121. By M. A. Pavis. Paris, 1833. 8vo.
Delamarre's Traité Pratique de la Culture des Pins. 3149. Paris, 1834. 3d. ed.
Delarb. Auvergn. 979. See *Delarb. Fl. Auver.*
Delarb. Fl. Auver. 1053. *Antoine de l'Arbre. Floræ d'Auvergne.* Ed. 1., 1 vol. 8vo, Clermont-Ferrand, 1795. Ed. 2., Riom et Clermont, 1800, 2 vols. 8vo.
Delauy. Bon Jard. 755. See *Bon Jard.*
Delauy Herb. Anat. 338. *Delauy in Herbar de l'Amateur,* a French periodical.
Delless. Icon. 297. See *Delless. Icon. Sel.*
Delless. Icon. sel. 306. Benj. Delessert, *Icones selectæ Plantarum in System. univers. descriptarum.* Paris, 1820. 4to.
Deleuze's Hist., &c. 136. *History and Description of the Royal Museum of Natural History, &c.* Translated by M. Deleuze. Paris, 1833. 8vo.
Del. Fl. Egypt. 534. Alir. Rafen Delle, *Floræ Egyptiæ Illustratio.* Paris, 1813. Fol.
Delicæ Sylvarum, 1792; or, *Grand and Romantic Forest Scenery in England and Scotland.* Drawn from Nature, and etched by J. G. Strutt. Lond. Fol.
Delile Descri. de l'Égypte. 1289. A. R. Delile, *Mémoires botaniques, extraits de la Description de l'Égypte.* Paris, 1813. Fol.
Dello Influsso dei Boscchi, &c. 220. *Dello Influsso dei Boscchi sullo Stato fisico dei Paesi, e sulla Prosperità della Nazione.* By G. Gautieri. Milan, 1817. 8vo.
Den Denkschrift d. Regensb. Bot. Ges. 1635. *Denkschriften der Königlich-Bairischen Botanischen Gesellschaft in Regens-burg.* Regensburg, 1815, and continued. 4to.
Dendrographia. 187. By John Johnston, Frankfurt, 1662. Fol.
Den. Brit. 235. See *Dendrologia Britannica.*
Dendrologia Britannica, 188; or, *Trees and Shrubs that will live in the open Air of Britain.* By P. W. Watson. Lond. 2 vols. 8vo. pl. 172.
Dendrologia Naturalis. 187. Ulysses Aldrovandus, *Dendrologia Naturalis, Libri duo.* Bononiæ, 1607. Fol.
Dendrology, 834; in which are Facts, Experiments, and Observations, demonstrating that Trees and Vegetables derive their Nutrimt independently of the Earth, &c. By John Wallis, Timber Surveyor. Lond. 1833. 8vo.
Derbyshire Report 2023. *General View of the Agriculture of Derbyshire.* By John Farey, sen. London, 1817. 2 vols. 8vo.
Der Hausvater. 145. By Baron Otto Von Münchhausen of Schwöbbache, 1765.
Der Trüffelbau. 1975. *Der Trüffelbau, oder Anweisung die schwarzen und weissen Trüffeln in Wäldern, Lustgebüsch, und Gärten, durch Kunst zu ziehen, und grosse Anlagen dazu zu machen.* Von Alexander Von Bornholz. Quedlinburg und Leipsig, 1825.
De Re Rustica. 835. L. J. M. Colmella, *De Re Rustica Libri duodecim.* Paris, 1543. 8vo.
De Saticibus Europæis Commentatio. 1456. By G. D. J. Koch. Erlangen, 1828. 8vo.
Description of the Scenery of the Lakes. 2374. By W. Wordsworth.
Deserted Village. 839. A poem by Goldsmith.

- Desf. Act. Par.* 526. Actes de la Société d'Histoire Naturelle de Paris. Paris, 1792. Folio.
Desf. Ann. Mus. 423. See *Annales du Muséum*.
Desf. Allan. 1817. See *Desf. Fl. Atl.*
Desf. Fl. Atl. 316. R. L. Desfontaines, Flora Atlantica. Paris, 1798, 1799. 2 vols. 4to.
Desf. Cor. 744. Id., Choix des Plantes du Corollaire des Instituts de Tournefort. Paris, 1808. 4to.
Desf. Hist. des Arbres et Arbustes. 1809. Id., Histoire des Arbres et Arbustes qui peuvent être cultivés en pleine Terre sur le Sol de la France. Paris, 1809. 2 vols. 8vo.
Desf. Cat. Hort. Paris. 989. See *Desf. Catal. Hort. P.*
Desf. Catal. Hort. P. 304. Id., Catalogus Plantarum Horti regii Parisiensis. Paris, 1829. 8vo. 3d ed.
Desf. Hort. Par. 258. See *Desf. Catal.*, &c.
Desf. Hort. Par. et Arb. 1240. See *Desf. Catal.*, &c., and *Desf. Hist. des Arbres*, &c.
Desv. Journ. Bot. 755. Desvieux, Journal de Botanique. Paris, 1808—1814. 5 vols. 8vo.
Dicks. Crypt. 1956. See *Dicks. H. Sicc. Fasc.*
Dicks. H. Sicc. 1706. See *Dicks. H. Sicc. Fasc.*
Dicks. H. Sicc. Fasc. 2066. Plantarum cryptogamicarum Eritanniæ Fasciculi quatuor. Lond. 1785—1801. 4to.
Dict. Clas. d'Hist. Nat. 1212. See *Dict. Classique d'Hist. Nat.*
Dict. Classique d'Hist. Nat. 1385. Dictionnaire Classique d'Histoire Naturelle, par MM. Audouin, Bourdon, Brongniart, Edwards, De Férussac, Desprez, Florens, Jussieu, Lucas, Richard, Bory De St. Vincent, &c. Paris, 1824—1830. 17 vols. 8vo.
Dict. des Eaux et des Forêts. 654. See *Baudrillart Traité*.
Dict. Gén. des Eaux et Forêts. 236. See *Baudrillart Traité*.
Dictionnaire. 612. See *Baudrillart Traité*.
Dictionnaire Général des Eaux et Forêts. 564. See *Baudrillart Traité*.
Dictionnaire des Sciences Naturelles. 2111. Publié par les Professeurs du Jardin du Roi. Paris, 1816—1829. 60 vols. 8vo.
Dictionnaire Encyclopédique. 2504. Lamarek (J. B.), Dictionnaire Méthodique, Partie Botanique. Paris, 1789. 8 vols. 4to.
Dict. Fl. Taur. 768. See *Bieb. Fl. Taur.-Cauc.*
Dict. Hist. et Crit. 1720. Dictionnaire Historique et Critique. Par P. Bayle. Rotterdam, 1697—4 vols. fol.
Dict. of Comm. 2113. See *M'ulloch's Dict.*
Dictionary of Commerce. 1335. See *M'ulloch's Dict.*
Dictionnaire Universel. 1989. Vahmont de Bonare, Dictionnaire Raisonné Universel d'Histoire Naturelle. Lyons, 1791. 8vo.
Dietr. Lec. Suppl. 1050. Dietrichs (J. G.), Vollständiges Lexicon der Gärtnerer und Botanik. Weimar, 1801. 2 vols. 8vo.
Dill. Eth. 234. Joh. Jac. Dillenius, Hortus Flthamensis. Lond. 1732. 2 vols. fol.
Dill. Mus. 1832. Id., Historia Muscorum. Ed. 1., Oxonii, 1741; ed. 2., Lond., 1763. 4to.
Dissertation. 2104. In G. Forster's De Plantis Escentibus. Berlin, 1786. 8vo.
Dissertations on Oriental Gardening. 80. See *Chambers's (Sir J.) Dissertations, &c.*
Dobell's Travels. 1500. Travels in Kamtschatka and Siberia. Lond. 2 vols. small 8vo.
Dod. Pempt. 878. See *Dodon. Pempt.*
Dodon. Pempt. 765. Rambertus Dodonæus seu Dodoens, Stirpium Historiæ Pemptades Sex, sive Libri xxx. Antverpæ, 1616. Fol.
Domestic Gardener's Manual. 624. The Domestic Gardener's Manual. London, 8vo.
Don's Mill. 252. See *Don's Miller's Dictionary*.
Don's Miller's Dictionary. 222. A general System of Gardening and Botany, &c., founded on Miller's Dictionary, and arranged according to the Natural System. By George Don, F.L.S. In 4 vols. 4to. 3 vols. published in 1837.
Don's Prod. 291. See *D. Don Prod. Nep.*
D. Don Prod. Nep. 1100. David Don, Prodromus Floræ Nepalensis. Lond. 1825. Small 8vo.
D. Don MSS. 1849. Professor Don MSS.
Donald's Cat. 1262. A Catalogue of Trees and Shrubs contained in Robert Donald's Arboretum, at Goldworth Nurseries, Woking, near Ripley, Surrey. Folio sheet.
Donn Hort. Can. 250. See *Donn Hort. Cantab.*
Donn Hort. Cantab. 2158. James Donn, Hortus Cantabrigiensis; or, a Catalogue of Plants, indigenous and foreign, cultivated in the Walkerian Botanic Garden at Cambridge. Cambridge, 1796. 8vo.
Doomsday-Book. 1573. Doomsday-Book seu Liber Censualis Willelmi Primi, Regis Angliæ, inter Archivos Regni in Domo Capitulari Westmonasterii asservatus; jubente Rege augustissimo Georgio Tertio prælo mandatus typis. 2 tom. folio. Lond. 1783.
Dougl. MS. 988. See *Douglas's MSS.*
Douglas MS. 248.
Drake's Shakespeare. 1345. Drake's edition of Shakespeare.
Drury Introd. 1817. Drury (D.), Exotic Insects. London, 1770—1782. 3 vols. 4to.
Dryden's Virgil. 1223. Dryden's Translation of Virgil.
Dublin Soc. Trans. 902. Transactions of the Dublin Society. Dublin, 1800—1810. 6 vols. 8vo.
Duby et Decand. Bot. Gallie. 1572.
Duby and De Candolle's Botanicum Gallicum. 132. J. E. Duby et A. P. De Candolle, Botanicum Gallicum, seu Synopsis Plantarum in Flora Gallica descriptorum. Paris, 1828—1830. 2 vols. 8vo.
Duh. Arb. Fr. 477. See *Du Hamel's Arbres*.
Du Hamel's Arbres Fruitiers. 984. Traité des Arbres Fruitiers. By Henri Louis Du Hamel du Monceau. Paris, 1768. 3 vols. 8vo.
Du Ham. Arb. Nouv. 261. See *Duh. Ed. Nouv.*
Duh. Ed. Nouv. Du Hamel's Traité des Arbres et Arbustes qui se cultivent en France en pleine Terre. A new edition. By Michel. Paris, 1811—1816. 5 vols. fol.
Dum. Bot. Cult. 2153. See *Botaniste Cultivateur*.
Dum. Cours. 503. See *Botaniste Cultivateur*.
Dum. Cours. Bot. Cult. 531. See *Botaniste Cultivateur*.
Dum. Cours. Suppl. 821. See *Botaniste Cultivateur*.
Duncead. 1701. A poem by Pope.
Dun. incd. 522. Dunal's MSS.
Dun. Man. Anon. 293. See *Dunal Manog.*
Dunal. Monog. 293. Mich. Felix Dunal, Monographie de la Famille des Anonacées. Paris, 1817. 4to.

- Dur. Bourg.* 585. Durande, Flore de Bourgogne. Dijon, 1782. 2 vols. 8vo.
Du Roi Harbk. 401. Joh. Phil. Du Roi, Die Harbkesche wilde Baumzucht. Braunschweig, 1771, 1772. 2 vols. 8vo.
D'Uro. Enum. 586. Enumeratio Plantarum quas in Insulis Archipelagi, Anno 1819, 1820, collegit Dumont D'Urville. Paris, 1822. 8vo.
Dwight's Travels in New England. 185. Travels in New England and New York. By Dr. Dwight. Lond. 1823. 2 vols. 8vo.

E.

- Eaton Man. Bot. Ed.* 1047. Id., A Manual of Botany for the Northern and Middle States of America. Albany, 1824. 12mo.
Econ. Hist. 101. Economical History of the Hebrides and Highlands of Scotland. 88. By Dr. Walker. Edinb. 1812.
Ed. Phil. Journ. 2110. Edinburgh Philosophical Journal. Conducted by Dr. Brewster and Professor Jameson. Edinb. 1819—1824. 10 vols. 8vo. Continued by Prof. Jameson alone, under the same name, from 1824; and from 1826, called "The Edinburgh New Philosophical Journal." Edinburgh, 1819, and continued. 8vo.
Edw. Ornith. 2402. Natural History of uncommon Birds, and of some other rare and undescribed Animals, Quadrupeds, Reptiles, Fishes, Insects, &c. By Geo. Edwards. Lond. 1743—1751. 4 vols. 4to, pl. 210.
Ehret Act. Angl. 1120. Ehret (G. D.), in the Philosophical Transactions.
Ehret P.-cl. 936. G. D. Ehret, Plantæ et Papiliones rariores. Lond. 1748—1759. Fol.
Ehrh. A. bor. 772. See *Ehrh. Beitr.*
Ehrh. Beitr. 683. Friedrich Ehrhart, Beiträge zur Naturkunde. Hanover et Osnabruk, 1787—1792. 7 vols. 8vo.
Eichwal Plantæ Caspico-Caucasice. 1932. Wilna and Leipsig, 1831. folio.
Elegy. 176. Gray's Elegy in a Country Churchyard.
Elementary Art. 203. See *Harding's Elementary Art.*
Elements of Naval Architecture. 140. Eléments de l'Architecture Navale. By Du Hamel du Monceau. Paris, 1752. 4to.
Elench. Sard. 707. Moris (J. H.), Stirpium Sardoarum Elenchus. Carali, 1827—1829.
Elles's Account of the Parish of St. Leonard, Shoreditch. 77. The History and Antiquities of the Parish of St. Leonard, Shoreditch, and Liberty of Norton Folgate, in the Suburbs of London. London, 1798. 4to.
Elliott Fl. S. Car. 825. A Sketch of the Botany of South Carolina and Georgia. By Steph. Elliott. Charleston, 1821—1824. 2 vols. 8vo, pl. 12.
Elliott Journ. Acad. Sci. Philad. 647. See *Journ. Acad. Scien. Phil.*
Ellis's Nat. Hist. of Corallines. 70. Essay towards a Natural History of Corallines, and other Marine Productions of the like Kind, commonly found on the Coast of Great Britain and Ireland. By John Ellis. Lond. 1755. 4to, pl. 39.
Ellis's Nat. Hist. of Zoophytes, &c. 70. Natural History of many curious and uncommon Zoophytes, collected from various Parts of the Globe, systematically arranged and described. By Dr. D. Solander. 1d. Lond. 1786. 4to, pl. 62.
Ell. Sketch. 1047. See *Elliott Fl. S. Car.*
En. Pl. 237. See *Encyc. of Plants.*
Encyc. Amer. 85. Encyclopædia Americana. A popular Dictionary of Arts, Sciences, Literature, History, Politics, and Biography, brought down to the present time; including a copious Collection of original Articles in American Biography; on the basis of the Seventh Edition of the German Conversations-Lexicon. Edited by F. Lieber and E. Wrigglesworth. Philadelphia, 1829. 13 vols. 8vo.
Encyc. Bot. 1361. See *Lam. Encyc.*
Encyc. of Agr. 572. An Encyclopædia of Agriculture, &c. By J. C. Loudon, F.L.S., &c. Lond. 1831. 8vo.
Encyc. of Cot. Arch. 897. An Encyclopædia of Cottage, Farm, and Villa Architecture, Furniture, &c. By J. C. Loudon, F.L.S., &c. Lond. 1833. 8vo.
Encyc. of Gard. 76. An Encyclopædia of Gardening; comprising the Theory and Practice of Horticulture, Floriculture, Arboriculture, and Landscape-Gardening, &c. By J. C. Loudon, F.L.S., &c. Lond. 1835. Ed. 2. 8vo.
Encyc. of Plants. 4. An Encyclopædia of Plants; comprising the Description, Specific Character, Culture, History, Application in the Arts, &c. By J. C. Loudon, F.L.S., &c. Lond. 1831. New ed. 1833. 8vo.
Encyclopædia of Arboriculture. 1805. See *Arboretum Britannicum.*
Encyclopédie Méthodique. 1452. Encyclopédie; ou Dictionnaire raisonné des Sciences, des Arts, et des Métiers, &c. Lausanne, 1781. 39 vols. 8vo.
Eng. Bot. 236. English Botany. By Sir J. E. Smith and Messrs. Sowerby. Lond. 1790-1814. 36 vols. 8vo. An abridged edition, with partially coloured plates, is now (1838) publishing.
Eng. Bot. Suppl. 766. Supplement to the English Botany of Sir J. E. Smith and Messrs. Sowerby; the Descriptions, &c., by Sir W. J. Hooker, LL.D., and other eminent Botanists. Lond., 1831. 8vo. Continued.
Eng. Fl. 344. See *English Flora.*
Engl. Gard. Cat. 1043. A Catalogue of the Plants cultivated in the Gardens near London. By a Society of Gardeners. London, 1730. Folio.
English Bowman; or Tracts on Archery. 2087. See *Robert's English Bowman, &c.*
English Flora. 27. English Flora. By Sir J. E. Smith. London, 1824—1828. 4 vols. 8vo.
Entomological Magazine. 903. Lond. 1831. In quarterly Nos. 8vo.
Entomological Society's Transactions. 1450. Transactions of the Entomological Society of London. Lond. Begun in 1834, and continued in 8vo.
Enum. Pl. Volhyn. 1598. See *Besser En. Pl. Volhyn.*
Enum. Stip. Ruth. 1377. See *Ammann Stip. Ruth.*
Enumeration of the Plants of the Berlin Royal Garden. 1416. See *Willdenow's Enumeration, &c.*
E'pitaphe de Colardeau à Du Hamel de Denainvillers, le Frère de Du Hamel. 2474.
Epithalamium. 1430. See *Herrick's Epithalamium.*
Esch. Mém. Acad. Scienc. Peters. 510. Eschscholtz in Mémoires of the Academy of Sciences at St. Petersburg.

- Essai sur les Harmonies Végétales et Animales du Chêne.* 1722. By M. Marquis. (? Paris, 1828.)
Essay on Arceuth. 2087. See *Moseley's Essay on Arceuth.*
Essay on Enclosure and Planting, &c., in Scotland. 87. See *Sutherland's Essay, &c.*
Essay on Plantations. 1988. Francis Bacon, Viscount St. Alban, Silva Silvarum. London, 1627.
Essay on Planting. 188. Essay on Planting. By the Rev. W. Hanbury. London, 1758. 8vo.
Essay on Trees in Landscape. 14. See *Kennion's Essay, &c.*
Essays. 836. See *Essays on Natural History.*
Essays on Natural History. 710. By Dr. Walker. London, 8vo.
Essays on Husbandry. 2278. By the Rev. J. Harte. London, 1764. 8vo.
Études de la Nature. 1307. See *St. Pierre's Études de la Nature.*
Evelyn's Diary. 59. Memoirs of John Evelyn, Esq., F.R.S.; comprising his Diary from 1641 to 1705-6, and a Selection of his familiar Letters. To which is subjoined the Private Correspondence between King Charles I. and Sir Edward Nicholas, &c. Edited by W. Bray, Esq., F.A.S. London, 1827. 5 vols. 8vo.
Exot. Fl. 306. See *Hook. Ex. Fl.*
Exploitation des Bois. 570. H. L. Du Hamel du Monceau, De l'Exploitation des Bois. Paris, 1764. 2 vols. 4to.
Ex. Cent. 683. Jacobus Breynius, Exoticarum Plantarum Centuriam. Gedani, 1678. Fol.

F.

- Fab. Ent.* 1884. Jo. Christ. Fabricii Entomologia systematica emendata et aucta sec. Classes, Ordines, Genera, et Species. Hafnia, 1792-1799. 9 vols. 8vo.
Fabric. Helms. 2464. P. C. Fabricii Enumeratio methodica Plantarum Horti Medici Helmstadtensis. Helmstadt, 1759. 8vo.
Faerie Queene. 677. A poem by Spenser.
Fairchild's City Gardener. 77. The City Gardener; containing the most experienced Method of cultivating and ordering such Evergreens, Fruit Trees, Flowering Shrubs, &c., as will be ornamental and thrive best in the London Gardens. By Thomas Fairchild 1722. Pamph. 8vo.
Faithful Shepherdess. 1958. A poem by Beaumont and Fletcher.
Familles des Plantes. 2163. See *Adan. Fam.*
Farm. Mag. 101 The Farmer's Magazine. Edinb. 8vo.
Faun. Ins. Germ. 1824. G. W. F. Panzer, Fauna Insectorum Germanicæ Initia. Nürnberg. 1793-1833. 8vo.
Fawkes's Bramham Park. 2703. A poem.
Feuillée It. 1276. Feuillée (Louis). Journal des Observations Physiques, Mathématiques, et Botaniques faites dans l'Amérique Méridionale, &c. Paris, 1714-1745. 4to
First Additional Supplement to the Encyclopædia of Agriculture. 186. By J. C. Loudon, F.L.S., &c. Lond. 1834. Pamph. 8vo.
First Additional Supplement to Hort. Brit. 1073. By J. C. Loudon, F.L.S., &c. Lond. 1832. Pamph. 8vo.
First Annual Report of the Edinburgh Botanical Society. 2590. Edinb 1837. 8vo.
Fisch. Hort. Gorenk. 631. F. Fischer, Catalogue du Jardin des Plantes de Gorenki, près de Moseou. 1808. 12mo.
Fisch. Misc. Itasp. 1-94. Fischer (F.), Spanische Miscellen &c. Dresden, 1804. 8vo.
Fisch. MSS. 345. Fischer's MSS.
Fischer's Anleit. zur Trüffeljagd, &c. 1975. V. F. Fischer, Anleitung zur Trüffeljagd, ein Beitrag zur Forst- und Jagd-Wissenschaft. Karlsruhe, 1812. Pamph.
Five Hundred Points of Good Husbandry, 836; together with a Book of Huswifery, &c. By Thomas Tusser. New ed. Lond. 1812. 8vo.
Fl. Br. 237. Sir J. E. Smith's Flora Britannica. Lond. 1800-1804. 3 vols. 8vo.
Fl. Cab. 2547. The Floral Cabinet. By G. B. Knowles and F. Wescott. London and Birmingham. 1837, 1838. 4to.
Fl. Carolinaana. 120. T. Walter. Flora Carolinaana. Lond. 1788. 8vo.
Fl. Dan. 343. See *Flora Danica.*
Fl. de Virg. 1956. See *Flore de Virgûl.*
Fl. Fr. 337. See *Flore Française.*
Fl. Gr. 322. See *Flor. Græca.*
Fl. Hib. 1087. See *Mackay Fl. Hibern.*
Fl. Ind. Occid. 5271. Swartz (Olof), Flora Indiæ Occidentalis. Erlangen, 1797, 1800, 1806. 3 vols. 8vo.
Fl. Mex. Icon. ined. 713. Sesse et Mocino, Flora Mexicana, Icon. ined. Iconibus ab ipso Sesse et Cervantesio curatis nondum editis usus est Candolleus.
Fl. Nap. 583. See *Tenore's Flora Neapolitana.*
Fl. Pyr. 778. Philippe Picot De la Peyrouse, Figures de la Flore des Pyrénées. Paris, 1795-1801. Fol.
Fl. Wett. 581. Flora der Wetterau von G. Gärtner. By Meyer und J. Scherbius. 1799-1801. 4 vols. 8vo.
Flor. Græc. 211. John Sibthorp. Flora Græca. Edited by Sir J. E. Smith, and continued by Sir W. J. Hooker, and Dr. Lindley.
Flora. 178. See *Pursh's Fl. Amer. Sept.*
Flora Americana Septentrionalis. 2119. See *Pursh Fl. Amer. Sept.*
Flora Anglica. 27. See *Huds. Fl. Angl.*
Flora Carniolica. 1456. Scopoli J. A., Flora Carniolica. Vienna, 1760. 8vo.
Flora Danica. 153. Flora Danica, sive Icones Plantarum sponte nascentium in Regnis Daniae et Norvegiae, &c. By Vahl, Horneman, and Müller. Hafniae. 1762-1839. 9 vols. fol.
Flora Domestica. 1282. Flora Domestica; or, the Portable Flower-Garden. London, 1823. 8vo.
Flora Edunensis. 2540. See *Grey. Fl. Ed.*
Flora Germanica. 146. See *Roth's Flora Germanica.*
Flora Helvetica. 160. See *Suter's Flora Helvetica*, and also *Gaudin's Flora Helvetica.*
Flora Hungarica. 146. Waldstein (F.) et Kitaibel (P.), Descriptiones et Icones Plantarum rariorum Hungariæ Vienna, 1802-1812. 4 vols. fol.
Flora Italica. 164. See *Bertoloni's Flora Italica.*
Flora Lapponica. 153. Carolus Linnæus s. von Linné, Flora Lapponica. Amstelodami, 1737. 8vo, 2d ed. By Sir J. E. Smith. Lond. 1792.

- Florula Louisiana.* 518. *Florula of Louisiana.* By Professor Rafinesque. New York, 1817.
- Flora Lusitania.* 1414. See *Brot. Flor. Lus.*
- Flora Metropolitana.* 882. *Flora Metropolitana; or, Botanical Rambles within Thirty Miles of London.* By Daniel Cooper. Lond. 1836. 12mo.
- Flora Mex. ic. and MSS.* 713. See *Fl. Mex. icon. ined.*
- Flora Neapolitana.* 164. See *Tenore's Flora Neapolitana.*
- Flora of Belgium.* 143. Lejeune and Courtois, Horti Botanici Academicæ Leodiensis Discretioni adjunctus; conjunctis studiis ediderunt: Compendium Floræ Belgicæ. Liege, 1828—1831. 2 vols. 8vo.
- Flora of Berwick upon Tweed.* 512. See *Johnston's Flora of, &c.*
- Flora of Nantes.* 263. Bonamy, *Flora Nannetensis Prodrromus.* Nann. 1782. 12mo.
- Flora of the Faroe Islands.* 153. On the Vegetation and Temperature of the Faroe Islands. By W. C. Trevelyan, Esq. Published in the Edinb. Phil. Journ. for January, 1835.
- Flora Peruviana.* 1502. Ruiz et Pavon, *Flora Peruviana et Chilensis.* Matriti, 1798, 1799. 5 vols. fol.
- Flora Scandinavia.* 153. A. J. Retzius, *Floræ Scandinaviæ Prodrromus.* Holmiæ, 1779. Ed. 2., Lipsiæ, 1795. 8vo.
- Flora Scotica.* 1437. See *Lightfoot's Flora Scotica.*
- Flora Scotica.* 123. See *Hook. Fl. Scotica.*
- Flora Silesiaca.* 1273. See *Krock. Fl. Sil.*
- Flora Suecica.* 153. C. Linnaeus, *Flora Suecica.* Holmiæ, 1745. Ed. 2., 1755. 8vo.
- Flora Transyl.* 1210. Baumgarten (J. C. G.), *Enumeratio Stirpium Magni Transylvaniæ Principatus.* Vienna, 1806. 3 vols. 8vo.
- Flora Taurico-Caucasica.* 255. L. B. F. Marschall de Bieberstein, *Flora Taurico-Caucasica.* Charokovia, 1818. 8vo.
- Floræ Siculæ Prodrromus.* 164. See *Gussone's Floræ Siculæ, &c.*
- Flor de Virgile.* 1956. A. L. A. Fée, *Flor de Virgile; ou, Nomenclature méthodique et critique des Plantes, Fruits, et Produits végétaux mentionnés dans ses Ouvrages.* Paris, 1802. 8vo.
- Flora Française.* 692. A. P. De Candolle et Lamarek, *Flora Française.* Paris, 1805—1815. 5 vols. 8vo.
- Floræ Javæ.* 2003. See *Blume Fl. Jav.*
- Florula Bostoniensis.* 1370. Jacob Bigelow. *Florula Bostoniensis.* Boston, 1814. 8vo.
- Flügge Ann. Mus.* 829. Flügge in *Ann. Mus.* See *Annales du Muséum.*
- Forbes's Hort. Tour.* 1926. *Journal of a Horticultural Tour through Germany, Belgium, and Part of France, in the Autumn of 1835.* By James Forbes, A.L.S., &c. London, 1837. 8vo.
- Forest Law.* 1751. See *Manwood on Forest Law.*
- Forest Trees, &c.* 695. *The Manner of raising, ordering, and improving Forest and Fruit Trees, &c.* By Moses Cook. London, 1676. 4to.
- Forster's Guide.* 1797. See *Monteath's Forster's Guide.*
- Forsk. Ægypt.* 235. See *Forsk. Ægypt. Arab.*
- Forsk. Ægypt. Descr.* 707. See *Forsk. Ægypt. Arab.*
- Forskaol Ægypt. Arab.* 1248. Petrus Forskaol, *Flora Ægyptiaco-Arabica.* Harniæ, 1775. 4to.
- Forster MS.* 1588. G. Forster's MS.
- For. Trees of Russ. in Jam. Journ.* 2175. A Paper on the Forest Trees of Russia. By Dr. Howison, in Jameson's Journal.
- Forsyth's Chronological and Biographical History.* 80. By W. Forsyth, Esq. Not yet published.
- Four Books of Husbandry.* 1774. See *Googe's Four Books of Husbandry.*
- Fr. El.* 2147. E. Fries, *Elenchus Fungorum, sistens Commentarium in Systema Mycologicum.* Gryphiswald, 1828. 2 vols. 8vo.
- Fragm. Fl. Ital.* 583. See *Viv. Fragn.*
- Franklin's First Journey.* 1675. See *Franklin's Narrative.*
- Franklin's (Captain John) Narrative of a Journey to the Shores of the Polar Sea in the Years 1819—1822, 2190.* Lond. 1823. 4to.
- Franklin's Journ. Append.* 638. Appendix to the above, on various Subjects relating to Natural History. By Dr. Richardson and J. Sabine, Esq. Lond. 1823. 4to.
- Fras. Cat.* 607. See *Fraser's Cat.*
- Fraser's Cat.* 989. A Catalogue of Plants cultivated in Fraser's Nursery, Chelsea.
- French Encyclopædia.* 612. See *Encyc. Méth.*
- Fries Novit.* 1598. E. M. Fries, *Novitiæ Floræ Suecicæ.* Lunda, 1814. 4to.
- Fries Syst. Mus.* 1703. See *Fries Syst. Myc.*
- Fries Syst. Myc.* 2147. *Systema Mycologicum sistens Fungorum Ordines, Genera, et Species huc usque cognitatas.* Gryphiswald et Lunda, 1821—1823. 3 vols. 8vo.
- Fuchs Hist.* 551. Leonhard Fuchs, *De Historiâ Stirpium Commentarii insignes.* Basileæ, 1542. Fol.
- Fumifugium.* 368. Evelyn's (John) *Fumifugium; or, the Inconvenience of the Air and Smoke of London dissipated; together with some Remedies humbly proposed.* London, 1661. 12mo.
- Furber's Nursery Catalogue.* 2015. Catalogue of English and Foreign Trees. By R. Furber. London, 1733. 8vo.

G.

- Gært. Fr.* 358. See *De Fruct. et Sem.*
- Gært. Fruct.* 257. See *De Fruct. et Sem.*
- Gært. Sem.* 1441. See *De Fruct. et Sem.*
- Gagnob. Act. Helvet.* 1706. Gagnebin (A.) in *Acta Helvetica Physico-Mathematico-Botanico-Medica.* 1751, and continued. 4to.
- Garcilasso.* 1502. Wiffen's Translation of Garcilasso.
- Gard. Mag.* 16. *The Gardener's Magazine.* Conducted by J. C. Loudon, F.L.S., &c. London, 13 vols. 8vo, to 1838. Continued monthly.
- Gardener's Dictionary.* 885. *The Planter's, Florist's, and Gardener's Dictionary; being a practical Collection from the most approved Authors in the English Language, relating to the above Three Parts of Gardening, founded on Experience, worthy of Notice, and adapted to the Climate of Scotland.* By James Gordon, Nurseryman and Florist, at Fountainbridge, near Edinburgh. Edinb. 1774. 8vo.
- Gardening.* 2008. See *James's.*
- Garid. Air.* 637. P. J. Garidel, *Histoire des Plantes qui naissent aux Environs d'Aix.* Aix, 1715. 2 vols. fol.

- Garten Ins.* 2144. See *Bouché Naturgesch.* &c.
Galer. Mont. 760. Gateraui, Description des Plantes qui croissent aux Environs du Montauban. Montauban, 1789. 8vo.
Gaudin's Flora Helvetica. 160. Gaudin (J.), Flora Helvetica. Tours, 1828—1830.
Genealogical History of the Ancient Family of Heron. 96. A Genealogical History, &c. By Sir R. Heron. London, 1863. 4to.
Gen. et Cat. N. A. P. 1301. Thomas Nuttall, The Genera of North American Plants, and a Catalogue of the Species. Philadelph. 1818. 2 vols. 12mo.
Gen. of N. Amer. Plants. 122. See *Gen. et Cat. N. A. P.*
Genera Plantarum. 2103. See *Juss. Gen.*
Genera Plantarum. 2103. See *Lia. Gen.*
General Report of Scotland. 89. By Sir John Sinclair. Edinb. 1791—1799. 21 vols. 8vo.
General Survey of the Agriculture of Scotland. 103. The present State of the Husbandry in Scotland. By Andrew Wight. Edinb. 1778—1790. 6 vols. 8vo.
Genl. Mag. 25. Gentleman's Magazine. London, Begun in 1731, and continued in monthly Nos. 8vo.
Geog. and Hist., &c. 2484. See *Geography and History of the Western States.*
Geography and History of the Western States. 2484. History and Geography of the Mississippi Valley; to which is appended a condensed Physical Geography of the Atlantic, United States, and the whole American Continent. By Timothy Flint. Cincinnati, 1832. 2 vols. 8vo.
Georgics. 1382. Dryden's Translation of Virgil.
Ger. Em. 693. See *Ger. Emac. App.*
Ger. Emac. 1010. See *Ger. Emac. App.*
Ger. Emac. Append. 526. John Gerard, The Herbal; or, General History of Plants gathered by John Gerard. Lond. 1597. Fol. Enlarged by Johnson, 1633.
Ger. Gallo-Prov. 336. Ludovic Gerarde, Flora Gallo-Provincialis Paris, 1761. 8vo.
Ger. Prov. 1064. See *Ger. Gallo-Pror.*
Gerard's Catalogue. 693. Jo. Gerard, Catalogus Arborum, Fruticum, ac Plantarum, tam indigenarum quam exoticarum, in Horto Gerardi nascentium. London, 1596. 8vo.
Gerard's Herb. 24. The Herbal; or, General History of Plants gathered by John Gerarde, Master in Chirurgerie. London, 1597. Folio.
Gerusalemme Liberata. 2472. A poem by Tasso.
Geschichte der Obstcultuur. 478. Allgemeine Geschichte der Obst-Cultur, von den Zeiten der Urwelten, bis auf die gegenwärtigen herab. By F. K. L. Siedler. Frankfurt, 1862. 8vo.
Gesn. Icon. Piet. 1943. Gesner (Conrad), Opera Botanica ex Bibliotheca C. J. Trew. Edidit et præfatus est C. C. Schmiedel. Norimb. 1751—1770. Folio.
Gilb. Bot. Prat. 580. Joh. Em. Gilbert, Histoire des Plantes d'Europe; ou Elémens de Botanique pratique. Lyons, 1798, 2 vols. 8vo. 2d ed., Lyons, 1806, 3 vols. 8vo.
Gilhes's MSS. 559. Dr. Gillies's MS.
Girardin's Essay on Landscape, &c. 1669. De la Composition des Paysages, ou des Moyens d'embellir la Nature autour des Habitations, en y joignant l'Utile à l'Agreable. Par M. L. R. Girardin, Vicomte d'Ermenouville. Paris, 8vo. Translated into English under the title of "An Essay on Landscape, with an Historical Introduction, &c. London, 1783. 12mo.
Gmel. Fl. Bad. 763. Gmel. Fl. Bad. Als. 768. Carol. Christ. Gmelin, Flora Badensis-Alsatia. Carlsruhe, 1805—1808. 5 vols. 8vo.
Gmel. Fl. Sibir. 729. Gmel. Sib. 247. Joh. Georg. Gmelin, Flora Sibirica. Petropoli, 1747—1769. 4 vols. 4to.
Gmel. Itin. 1211. Gmelin (J. G.), Travels through Siberia, between the Years 1733—1743. In German. Göttingen, 1751, 1752. 8vo.
Gmel. Syst. 240. Joh. Frid. Gmelin, Caroli Linnæi Systema Naturæ. Lugduni, 1796. 10 vols. 8vo.
Godefroy Cat. 756. The Catalogue of M. Godefroy, Nurseryman, Ville d'Avray, near Paris.
Goose's Four Books of Husbandrie. 1774. Four Books of Husbandrie, collected by Conradus Hereshbachius, &c.; containing the whole Art and Trade of Husbandrie, Gardening, &c. Newly Englished and increased. By Barnaby Googe. London, 1577. 4to.
Gorter Ingr. 734. David De Gorter, Flora Ingrica ex Schedulis Steph. Krascheninikow, &c. Petropoli, 1761. 8vo. Appendix, 1764.
Göthe's Versuch über die Metamorphose der Pflanzen. 225. J. W. Von Göthe. Versuch über die Metamorphose der Pflanzen. Stuttg. 1831. 8vo.
Gouan Descr. du Ginkgo. 2094. Ant. Gouan, Description du Ginkgo biloba. Montpellier, 1812. 4to. and 8vo.
Gouan Fl. Monsp. 357. Id., Flora Mompeliaca. Lugduni, 1762. 8vo.
Gouan Herb. 353. Id., Herborisations des Environs de Montpellier. Montpellier, 1796. 8vo.
Gouan Hort. 1046. Id., Hortus Regius Mompeliensis sistens Plantas tum indigenas tum exoticas, &c. Lugduni, 1762. 8vo, pl. 4.
Gouan Il. 369. Id., Illustrationes Botanicae. Tiguri, 1773. Fol.
Gouan Mons. 2464. See *Gouan Fl. Monsp.*
Græffer Pl. Essicc. 2234. Græffer (John), A Descriptive Catalogue of upwards of 1100 Species and Varieties of Herbaceous or Perennial Plants. London, 1789. 8vo.
Gray's Arr. 1123. Natural Arrangement of British Plants, according to their Relations to each other, as pointed out by Jussieu, De Candolle, Brown, &c., including those cultivated for use; with an Introduction to Botany, in which the Terms newly introduced are explained. By S. F. Gray. London, 1821. 2 vols. 8vo, pl. 21.
Green. Crypt. 1831. See *Green. Sc. Cr. Fl.*
Green. Sc. Cr. Fl. 2147. Scottish Cryptogamic Flora, or Descriptions of Plants belonging chiefly to the Order Fungi. By R. K. Greville. Edinb. 1825—1838. 6 vols. 8vo, pl. 360.
Green. Fl. Ed. 1834. Flora Ednensis. Id. Edinb. 1824. 8vo, pl. 3.
Grogner's Recherches Historiques et Statistiques sur le Mûrier, le Ver à Soie, et la Fabrication de la Soierie, &c. 1358. Lyons, 1826. Pamph. 8vo.
Gron. Orient. 1320. J. Fred. Gronovius, Flora Orientalis, sive Recensio Plantarum quas Rauvoisius collegit. Lugd. Bat. 1755. 8vo.
Gronov. et Wall. Fl. Car. 515. See *Fl. Caroliniana.*
Gronov. Virg. 1111. Id., Flora Virginica exhibens Plantas quas J. Clayton in Virginia collegit. Lugd. Bat. 1743, 4to. Ed. 2., Lugd. Bat. 1762, 4to.
Græcæ's Treatise on Ancient Arms and Armour. 2087. London, 1785—1786. 4to.
Guardian 1439. The Guardian. By Steele and others. London, 1714. 4to

- Guimp. Abb. Holz.* 926. F. Guimpel, Abbildung der deutschen Holzarten für Forstmänner und Liebhaber der Botanik. Berlin, 1815—1820. 2 vols. 4to, pl. 216.
- Güld. Itin.* 1058. Güldenstädt (T. A.), Reisen durch Russland, und in caucasiſchen Gebürge, herausgegeben von P. L. Pallas. St. Petersburg, 1787. 4to.
- Guss. Cat.* 587. Jo. Gussone, Catalogus Plantarum, quæ in Horto Regio asservantur. Neapol. 1821. 8vo.
- Gussone's Floræ Siculæ Proëromus.* 164. Neap. 1829.
- Gussone Pl. Rar.* 587. Id., Plantæ rariores per Regiones Samnii ac Apurtii collectæ. Neapoli, 1826. 4to, pl. 66.
- Gyll. Ins. Succ.* 1816. L. Gyllenhal, Insecta Suecica descripta. Scaris et Lips. 1808—1827. 4 vols. 8vo.

H.

- Habl. ined. Taur.* 948. Hablitz's Physikal. Beschr. d. Taurischen Statthalterſchaft nach ihrer Lage u. allen drey Naturreichen. Hann. and Osnab. 1797. 8vo.
- Hænke Beob.* 2186. Thadæus Hænke, Beobachtungen auf Reisen nach dem Riesen-gebürge, &c. Dresden, 1791. 4to.
- Hall. Helv.* 2499. Albert Von Haller, Historia Stirpium indigenarum Helvetiæ. Bernæ, 1768. 3 vols. fol.
- Hall. Hist.* 1375. See *Hall. Helv.*
- Halloween.* 2022. A poem by Burns.
- Hall's Sketches in Canada and the United States.* 2117. Forty Sketches of North America. By Capt. Basil Hall, R.N. London, 4to.
- Hall's Travels in Scotland.* 1681. Travels in Scotland by an unusual Route, with a Trip to the Orkneys and Hebrides; containing Hints for Improvement in Agriculture and Commerce, with Characters and Anecdotes, and a Map and Views. By the Rev. J. Hall. London, 1807. 2 vols. 8vo.
- Ham. MSS.* 245. See *Hamil. MSS.*
- Hamil. MSS.* 390. Hamilton MSS.
- Handbook for Travellers in Southern Germany.* 2115. London, 1827. 8vo.
- Harbkesche Baumzucht theils nord-amerikenscher und fremder, &c.* 190. By J. P. Du Roi. 1771—1772. 2 vols. 8vo.
- Harding's Elementary Art.* 203. Elementary Art, or the Use of the Lead-Pencil advocated and explained. By J. D. Harding. London, 1834. Folio.
- Harleian MS.* 512. Harleian Collection of MS. in the British Museum
- Hartig's Lehrbuch für Forster.* 2223. Hartig (G. L.), Instruction sur la Culture du Bois à l'Usage des Forestiers. Translated by J. Baudrillart. 1 vol. 8vo. Paris, 1805.
- Hasted's Kent.* 24. See *Beauties of England, &c.*
- Haworth Lep. Brit.* 1828. A. H. Haworth, Lepidoptera Britannica, sistens Digestionem novam Insectorum Lepidopterorum qua in Magni Britanniei reperitur, &c. Londini, 1803. 8vo.
- Haworth's Review of Entomology.* 1480. In the Entomological Society's Transactions, vol. i.
- Haworth Suppl.* 2527. A. H. Haworth, Supplementum Plantarum succulentarum. Lond. 1819. 8vo.
- Haworth Syn. Plant. Suc.* 965. Id., Synopsis Plantarum succulentarum. Lond. 1812. 8vo.
- Haw. Suppl.* 2525. See *Haworth Suppl.*
- Haw. Sup. Pl. Suc.* 2527. See *Haworth Suppl.*
- Hayley's Life and Posthumous Writings of William Cowper.* 1764. Life and Posthumous Writings of William Cowper, Esq.; with an Introductory Letter to the Right Hon. Earl Cowper. By W. Hayley, Esq. Chichester, 1803-4. 3 vols. 4to.
- Hayne Abbild.* 303. See *Hayne Abbild. der deut. Holz.*
- Hayne Abbild. der deut. Holz.* 586. Abbildung der deutschen Holzarten für Forstmänner und Liebhaber der Botanik, herausgegeben von F. Guimpel, entworfen und beschreibung von C. L. Willdenow, in letzter Rücksicht fortgesetzt von F. G. Hayne. Berlin, 1820. 4to.
- Hayne Den.* 233. See *Hayne's Dendrologische Flora.*
- Hayne Dendr.* 1455. See *Hayne's Dendrologische Flora.*
- Hayne's Dendrologische Flora.* 415. Dendrologische Flora, oder Beschreibung der in Deutschland im freien ausdauernden Holzgewächse. Von Dr. T. G. Hayne. Berlin, 1822. 8vo.
- Hayne Term. Bot.* 1014. F. G. Hayne et C. L. Willdenow, Terniini Botanici iconibus illustrati. Berolini, 1799. 4to.
- H. B. et Kunth Nov. Gen. Amer.* 503. Humboldt, Bonpland, et Kunth, Nova Genera et Species Plantarum. Paris, 1815—1825. 7 vols. 4to.
- Head's Continuation of a Home Tour.* 2549. Continuation of a Home Tour through the Manufacturing Districts of England, in the Summer of 1835. By Sir G. Head. London, 1837. 8vo.
- Headrick's Forfarshire.* 100. General View of the Agriculture of the County of Angus or Forfarshire, with Observations on the Means of Improvement, &c. By the Rev. James Headrick. London, 1813. 8vo.
- Heber's (Bishop) Journal.* 919. Heber's (Bishop) Journey through India. London, 1830. 2 vols. 4to, and 3 vols. 8vo.
- Hebrew Melodies.* 1463. By Lord Byron.
- Hedw. F. Gen.* 528. D. J. Hedwig, Filicum Genera et Species. Lips. 1799. Folio, pl. 6.
- Hebr.* Observations Entomologica, continens Metamorphoses Coleopterorum nonnullorum. Turici, 1836. 8vo.
- Heinsingla, eller Snorro Sturleson's Nordlänsche Konuga Sagor.* 2020.
- Hem. Diss.* 772. See *Herm. Diss.*
- Henry VI.* 839. One of Shakspeare's plays.
- Herb. Banks MSS.* 1530. MSS. in the Banksian Herbarium.
- Herb. Hænke.* 1059. Hænke's Herbarium.
- Herb. Reinw. dt.* 1168. Reinwardt's Herbarium.
- Herb. Thib.* 340. Thibaud's Herbarium.
- Herbal.* 1867. See *Parkinson's Herbal.*
- Herbal; or, Theatre of Plants, &c.* 52. See *Parkinson's Herbal, &c.*
- Herbarium.* 967. See *Willdenow's Herbarium.*
- Herbarium Mauritanium.* 542. Willemet (P. R.), Herbarium Mauritanium. Leipsig, 1796. 8vo.
- Hercfordshire Report.* 884. General View of the Agriculture of the County of Hercford, &c. By John Duncumb, A.M. London, 1805. 8vo.

- Herm. Diss.* 754. B. F. Hermann, Diss. de Rosa. Argentorati, 1762. 4to.
- Herm. Ludg.* 2498. Paulus Hermann, Floræ Lugduno-Batavæ Flores, ed. L. Zumbach. Lugd. Bat. 1690. 8vo.
- Herm. Par.* 424. Id., Paradisus Batavus, Opus posthumum, ed. G. Sherard. Lugd. Bat. 1698. 4to. Ed. 2., Lugd. Bat. 1705.
- Hern. Mex.* 713. See *Hern. Mex. Icon.*
- Hern. Mex. Icon.* 713. Hernandez (F. G.), Nova Plantarum, Animalium, et Mineralium Mexicorum Historia: a N. A. Reccho in Volumen digesta, a J. T. J. Fabro, et F. Columna, Lynceis, Notis et Additionibus illustrata. Romæ, 1651. Folio.
- Herrera's Agricultura.* 1298. Gabr. Alonso de Herrera, Agricultura general adicionada per la Real Sociedad Economica Matritense. Madrid, 1818. 4 vols. 8vo.
- Herrick's Epithalamium.* 1430. Herrick (R.), Hesperides; or, Works both Human and Divine. London, 1648. 8vo.
- Hierophyton*, &c. 1720. M. Hiller, Hierophyton, sive Commentarius in Loca Scripturæ Sacræ quæ Plantarum Mentionem faciunt. Trajecti ad Rhen. 1725. 4to.
- Highlands of Scotland*, &c. 1224. An Economical History of the Hebrides and Highlands of Scotland. By John Walker, D.D. Edinb. 1808. 2 vols. 8vo.
- Highland Soc. Trans.* 2161. See *High. Soc. Trans.*
- High. Soc. Trans.* 2165. Quarterly Journal of Agriculture; and the Prize Essays and Transactions of the Highland Society of Scotland. Edinb. 1828. In quarterly numbers. 8vo.
- Hints on the Planting and General Treatment of Hardy Evergreens.* 512. Hints on the Planting and General Treatment of Hardy Evergreens in the Climate of Scotland, &c. By William McNab. Edinb. 1831. Pamph. 8vo.
- Hist.* 717. See *History of the Arundel Family.*
- Hist. des Champs.* 1704. See *Roque's Hist. des Champs.*
- Hist. and Stat.* 258. Lang (J. D.), An Historical and Statistical Account of New South Wales. London, 1824. 2 vols. 12mo.
- Hist. des Plantes Pyr.* 2290. See *Lap. Hist. des Pl. des Pyrénées.*
- Hist. du Japon.* 1361. Kämpfer (Engelbert), History of Japan. Translated from High Dutch, by J. G. Scheuchzer, London, 1728, 2 vols. folio. Et traduite en Français sur la Version Anglaise de J. G. Scheuchzer, La Haye, 1729, 2 vols. folio.
- Hist. Nat. Hymenopt.* 2140. Lepelletier de Saint Fargeau, Histoire Naturelle des Insectes Hyménoptères, tom. 1. Paris, 1836. 8vo. (Part of the "Suites à Buffon.")
- Hist. Rei Herb.* 16. Kurt Sprengel, Historia Rei Herbariæ. Amstelodami, 1807. 2 vols. 8vo.
- Hist. of Incent.* &c. 147. History of Inventions and Discoveries. By John Beckman. Translated by Johnston. London, 1797. 3 vols. 8vo.
- Histoire des Arbres de l'Amérique.* 142. See *Michx. Arb.*, &c.
- Histoire des Arbres et Arbrisseaux.* 189. Histoire des Arbres et Arbrisseaux qui peuvent être cultivés en pleine Terre sur le Sol de la France. Paris, 1809. 2 vols. 8vo.
- Histoire des Arbres Forciers de l'Amérique Septentrionale.* 189. See *Michaux's Arbres Forciers*, &c.
- Histoire des Chênes de l'Amérique.* 189. See *Michaux's Histoire des Chênes*, &c.
- Histoire des Chênes de l'Amérique Septentrionale.* 142. See *Michaux's Histoire des Chênes*, &c.
- Histoire des Drogues.* 788. Pierre Pomet, Histoire Générale des Drogues. Paris, 1684. Fol.
- Histoire des Plantes.* 609. Jean Robin, Histoire des Plantes. Paris, 1620. 12mo. Printed with the 2d ed. of Lonicier's History of Plants.
- Histoire des Plantes du Dauphiné.* 963. D. Villars, Histoire des Plantes du Dauphiné. Grenoble, 1786—1788. 3 vols. 8vo.
- Histoire des Végétaux Fossiles.* 2111. By A. Brongniart. Paris, 1823. 8vo, pl. 117.
- Histoire du Cèdre.* 2411. See *Histoire du Cèdre du Liban.*
- Histoire du Cèdre du Liban.* A Paper by Loiseleur Deslongchamps, published in the Annales de l'Agrie. Franc. for 1837.
- Histoire Naturelle des Iles Canaries.* 1332. See *Webb and Bertholet's Histoire Naturelle*, &c.
- Historia Plantarum.* 16. See *Theophrastus's Historia Plantarum.*
- Historia Plantarum.* 48. See *Ray's Historia Plantarum.*
- Historia Salicum.* 1457. G. F. Hollmann, Historia Salicum iconibus illustr. Lipsiæ, 1755. Fol.
- Historical and Descriptive Account of Persia.* 2039. Fraser's.
- History of Craven.* 371. See *Whitaker's History of Craven.*
- History of Dublin.* 117. See *Walsh's History of Dublin.*
- History of Edinburgh.* 95. See *Arnol's History of Edinburgh.*
- History of Enfield.* 41. Dr. May's.
- History of Manchester.* 21. Whitaker's (Rev. J.) History of Manchester, in Four Books. London, 1771. 4to.
- History of Mexico.* 2051. The History of Mexico, collected from Spanish and Mexican Historians, &c. By the Abbé Clavigero. Translated by Charles Cullen. Lond. 1787. 2 vols. 4to.
- History of New England.* 2314. By J. Josselyn. London, 1672.
- History of Northamptonshire.* 1777. Whalley (Rev. P.), History and Antiquities of Northamptonshire. Oxford, 1791. 2 vols. folio. Compiled from the MS. of Mr. Bridges.
- History of Plants.* 1829. Ruellius (J.), De Natura Stirpium. Paris, 1536, fol. Basilii, 1543, fol.
- History of Selborne.* 1758. Natural History and Antiquities of Selborne. By the Rev. G. White. London, 1813. 4to.
- History of Somersetshire.* 1431. Collinson (Rev. J.), The History and Antiquity of the County of Somerset. Bath, 1791. 3 vols. 4to.
- History of the Arundel Family.* 717. The Antiquities of Arundel. The peculiar Privileges of its Castle and Lordship; with an Abstract of the Lives of the Earls of Arundel from the Conquest to the present Time. London, 1766. 8vo.
- History of the County of Down.* 116. The Ancient and Present State of the County of Down. By C. Smith. 1744. 8vo.
- History of the first Fourteen Years of King James.* 787. By Sir Thomas G. Cullum.
- History of the Mogul Empire.* 785. Histoire Generale de l'Empire du Mogul. Paris, 1705. 4to.
- History of the Rose.* 789. Rosenbergs (J. C.), De Rosa iatrica Animadversiones et Exercitationes Medica. Argent. 1624. 12mo.
- History of Willows.* 1456. See *Hoffm. Hist. Sal.*
- H. K.* 554. See *Art. Hort. Kew.*
- Hoffland's Description of White Knights.* 148. An Historical Description of White Knights. By Mr. and Mrs. Hoffland. Lond. 1820. Fol. ...

- Hoffm. Hist. Sal.* 1492. See *Hoffm. Sal.*
Hoffm. Sal. 1494. Georg. Franz. Hoffmann, Historia Salicum Iconibus illustr. Lipsiæ, 1785. Fol.
Hoffmansegg Fl. Port. 1207. Hoffmansegg et Link, Flore Portugaise. Rostoch et Berlin, 1806. Fol.
Hogg on the Classical Plants of Sicily. 164. See *Hogg's observations, &c.*
Hogg's Observations on the Classical Plants of Sicily. 164. An Article published in the Journal of Botany, 2d ser. p. 204.
Holland's Translation of Pliny's Natural History. 19. London, 1601. 2 vols. folio.
Holzbibliothek. 190. (Wood Library.) A term given by the Germans to a species of Publication, or rather portable Museum. A hundred or more sorts of wood, with dried specimens of the leaves, flowers, seed, and winter's wood, of each species, are put loose into little cases about the size of 8vo volumes, and these are finished exteriorly like books, and the back lettered with the name of the wood in different languages
Hom. Odys. 1723. Homer's Odysey, Pope's Translation, with Notes by Gilbert Wakefield. Lond. 1796. 8vo.
Hone's Every-Day Book. 900. Hone's (W.), Every-Day Book and Year Book. London, 1826. 3 vols. 8vo.
Hook. Bot. Miscel. 541. Botanical Miscellany; containing Figures and Descriptions of such Plants as recommend themselves by their Novelty, Rarity, or History. By Sir W. J. Hooker. London, 1830—1833. 3 vols. 8vo.
Hook. Bot. Comp. 1345. See *Hook. Comp. Bot. Mag.*
Hook. W. Jack. Comp. Bot. Mag. 1938. See *Comp to Bot. Mag.*
Hook. British Flora. 27. British Flora; comprising the Phanogamous or Flowering Plants, and the Ferns. 1d. Lond. 1830—1833. 2 vols. 8vo.
Hook. Ex. Fl. 1277. Exotic Flora. Id. Edinb. 1823—1827. 3 vols. 8vo, pl. 232.
Hook. Fl. Bor. Amer. 309. Flora Boreali Americana; compiled principally from the Plants collected on the Northern Land Expedition, under Command of Capt. Sir John Franklin. 1d. Lond. 1829—1834. 4to.
Hook. Fl. Scotica. 123. Flora Scotica; or, a Description of Scottish Plants, arranged both according to artificial and natural Methods. 1d. London, 1821. 8vo.
Hook. Lond. 769. Curtis's Flora Londinensis, continued by Sir W. J. Hooker. Lond. Fol. See *Curt. Lond.*
Hook. Scot. 235. See *Hook. Fl. Scot.*
Hoole's Tasso. 1958. London, 1767. 2 vols. 12mo.
Horn. Cat. Hort. Hafn. 345. Hornemann (J. W.), Enumeratio Plantarum Horti Botanici Hafniensis. Hafn. 1807. 8vo.
Horn. Hort. Reg. Haff. 2094. J. W. Hornemann, Hortus Regius Botanicus Hafniensis. Hafniæ, 1815. 8vo.
Horn. Hort. Hafn. Suppl. 584. J. W. Hornemann, Horti Regii Botanici Hafniensis Supplementum. Hafniæ, 1819. 12mo.
Horsfield's History, &c., of Sussex. 1838. Horsfield (T. W.), The History, Antiquities, and Topography of the County of Suffolk. Lewis, 1835. 4to.
Hort. Acad. Lugd. Bat. Cat. 45. Paulus Hermann, Horti Lugduno-Batavi Catalogus. Lugd. Bat. 1687. 8vo.
Hort. Angl. 647. Hortus Anglicus; or, the Modern English Flower-Garden, &c. By the Author of the British Botanist. London, 1822. 2 vols. 8vo.
Hort. Brit. 4. Hortus Britannicus. A Catalogue of all the Plants indigenous, cultivated in, or introduced to, Britain. Edited by J. C. Loudon, F.L.S., &c. Lond. 1830. 8vo.
Hort. Cliff. 293. Carolus Linnæus. Hortus Cliffortianus. Amsterd. 1737. Fol.
Hort. Eric. Woburn. 1079. Hortus Ericus Woburnensis. London, 1824.
Hort. Eyst. 1267. See *Besl. Eyst.*
Hort. Europæus Americanus. 76. By Mark Catesby. Lond. 1767. Fol.
Hort. Herrenh. 640. See *Hort. Herrenhaus.*
Hort. Herrenhaus. 750. Wendland (J. C.), Hortus Herrenhausanus. Hanover, 1778—1799. Folio.
Hort. Kew. 36. See *Ait. Hort. Kew.*
Hort. Kew. Icon. 1089. See *Ait. Hort. Kew.*
Hort. Par. 423. Hortus Parisiensis.
Hort. Lam. 319. See *Lam. Dict.*
Hort. Soc. Cat. of Fruits. 790. Catalogue of Fruits cultivated in the Garden of the London Horticultural Society. Lond. 1826. 8vo.
Hort. Trans. 251. Transactions of the London Horticultural Society. Lond. 1815—1831. 7 vols. 4to. New series commenced in 1831, and continued.
Hort. Worlitz. 1238. Description of the Gardens of Worlitz.
Horti Regii Hamptonensis. 40. MS. No. 3370. in the British Museum.
Hortis. 2008. Gesner (C.), Catalogus Plantarum Alphabeticus. Tiguri, 1542. 4to.
Hortus Addelestonensis. 987. By Mr. Cree. Lond. 1837. 12mo.
Hortus Americanus Europæus. 69. See *Hort. Europæus Americanus.*
Hortus Carlsruhanus. 152. Hortus Carlsruhanus, oder Verzeichniß sammtlicher Gewächse, &c. Herausgegeben von Garten-inspector Hartweg. Carlsruhe, 1825. 8vo.
Hortus Medicus Edinburgensis. 104. See *Sutherland's Hortus Medicus Edinburgensis.*
Hortus Elthamensis. 235. Joh. Jac. Dilleni, Hortus Elthamensis. Londini, 1732. 2 vols. fol.
Höss Anleit. 1045. See *Anleit. die Bäume und Sträucher Oesterreichs, &c.*
Höss's Gemeinssassliche Anleitung, &c. 2205. See *Anleit. die Bäume und Sträucher Oesterreichs, &c.*
Host Fl. Austr. 1393. See *Host's Fl. Austr.*
Host's Fl. Austr. 1454. N. T. Host, Flora Austriaca. Vienn. 1827—1831. 2 vols. 8vo.
Host's Sal. Austr. 1454. Host's Salix. Vienna. folio.
Host's Synops. 1498. Id., Synopsis Plantarum in Austrâ crescentium. Vindobonæ, 1797. 8vo.
Houel Voyage en Sicile. 1988. Houel (Jean), Voyage Pittoresque des Isles de Sicile, de Malte, et de Lipari, &c. Paris, 1782. 4 tom. folio.
Houst. Phil. Trans. et Reliq. Houst. 1276. Houston in Philosophical Transactions: and Josephus Banks, Reliquiæ Houstonianæ. Londini, 1781. 4to.
Houtt. Pflanz. 235. See *Houtt. Pfl. Syst.*
Houtt. Pfl. Syst. 648. Mark Houttuyn, Des Ritters von Linné Pflanzen-system nach der Anleitung des Houttuynschen Werks. Nürnberg, 1778—1788. 14 vols. 8vo.
Howell's Hist. of the World. 1333. An Institution of General History; or, the History of the World, from the Beginning, to the Death of Constantine Duca, Anno 1061. London, 1662. Folio.

- Howitt's Book of the Seasons.* 1480. W. Howitt's Book of the Seasons; or, the Calendar of Nature. Lond. 1831. 16mo.
- Huds. Fl. Angl.* 772. William Hudson, Flora Anglica. Lond. 1762, 1 vol. 8vo. Ed. 2., Lond. 1778, 2 vols. 8vo.
- Humb. et Bonpl. Pl. Equin.* 1173. Alexand. F. H. von Humboldt et Bonpland, Plantes Équinoxiales. Paris, 1805—1816. 2 vols. fol.
- Humb. et Bonpl. Nov. Gen. et Sp. Pl.* 1529. See *H. B. et Kunth Nov. Gen. &c.*
- Hunt's Foliage: Evergreens.* 2122. Leigh Hunt's Foliage, Poems. Lond. 12mo.
- Hunt. Ecol. Syst.* 1911. See *Hunter's Evelyn.*
- Hunter's Evelyn.* 45. Evelyn's Silva, with Notes. By A. Hunter, M.D. York, 1776. 2 vols. 4to.
- Hunter's (Admiral) Journal of Transactions at Port Jackson and Norfolk Island.* 2412. An Historical Account of the Transactions at Port Jackson and Norfolk Island. By Admiral John Hunter. Lond. 1793. 4to.
- Husbandry of Scotland.* 369. An Account of the System of Husbandry adopted in the more improved Districts of Scotland, &c. Edinb. 1812. 8vo.
- Hutchins's Account of Dorsetshire.* 1758. The History and Antiquities of the County of Dorset, &c. Lond. 1774. Folio.
- H. Wob.* 1087. See *Hort. Eric. Woburn.*
- Hymn to Delos.* 1725. Callimachus's Hymn and Epigrams.

I.

- Iconographia Rustica.* 80. Iconographia; or, the Nobleman, Gentleman, and Gardener's Recreation, &c. By Stephen Switzer. Lond. 1718. 3 vols. 8vo.
- Iliad.* 1382. Homer's Iliad, translated by Pope.
- Illustr. Brit. Ent.* 1483. See *Stephens's Illust.*
- Illustrations.* 386. See *Illustrations and Descriptions of the Camellieæ.*
- Illustrations and Descriptions of the Camellieæ.* 382. See *Chandl. Ill.*
- Illustrations of the Botany, &c., of the Himalayan Mountains.* 173. See *Royle Illust. Bot. Himalayas.*
- Incidents of Travels in Egypt, &c.* 2605. By an American. Lond. 1837. 2 vols. 8vo.
- Index Plantarum Agri Erjordiensis.* 1409. By J. Planer. Gothæ, 1788. 8vo.
- Insect Architecture.* 1826. Insect Architecture, forming Vols. VII. and VIII. of the Library of Entertaining Knowledge. Lond. 1830. 12mo.
- Insects of Georgia.* 1360. See *Abbott and Smith's Insects of Georgia.*
- Insect Transform.* 1819. Insect Transformations, being Vol. VI. of the Library of Entertaining Knowledge. Lond. 1330. 12mo.
- Ins. Succ.* 2141. L. Gyllenhal, Insecta Suecica descripta. Scaris et Lips. 1808—1827. 4 vols. 8vo.
- Institutiones.* 2103. Joseph Pitton de Tournefort, Institutiones Rei Herbariæ. Cur. A. De Jussieu. Paris, 1717—1719. 3 vols. 4to.
- Introd. to Bot.* 213. See *Lindl. Introd. to Bot.*
- Introduction to Botany.* 78. By James Lee. London, 1760. Ed. 2. 1765; ed. 3., 1776; ed. 4., 1791. 8vo.
- Irish Farmer's and Gard. Mag.* 108. Irish Farmer's and Gardener's Magazine. Conducted by Martin Doyle and Edmund Murphy. In monthly numbers. Dublin, 1834. 8vo.
- Irish Tour.* 1226. Tour in Ireland; with general Observations on the State of that Country, made in 1776—1779. By Arthur Young. Dublin, 1780. 2 vols. 8vo.
- Iring's Bracebridge Hall.* 1752. Bracebridge Hall; or, the Humourist. By Geoffry Crayon. Lond. 1816. 2 vols. 8vo.
- Isis.* 741. Isis, oder Encyclopädische Zeit., herausgegeben von Oken. Large 4to, monthly.
- Isle of Palms.* 1761. Wilson's (Professor) Isle of Palms, City of the Plague, &c. Lond. 2 vols. 8vo.
- Issues of the Erchequer.* 1353. Lond. 1837. 8vo.
- Itin. Curios.* 1768. Stukely (W.), Itinerarium Curiosum; or, an Account of the Antiquities and Remarkable Curiosities observed in Travels through Great Britain. 100 plates. Lond. 1724. Fol.

J.

- Jac. Sc.* 1277. See *Jacq. Hort. Schön.*
- J. cq. Austr.* 236. N. J. Von Jacquin, Floræ Austriacæ, sive Plantarum selectarum in Austria Archiducatu sponte crescentium Icones, ad vivam colorata, et Descriptionibus ac Synonymis illustratæ. Vienna, 1773—1778. 5 vols. fol. pl. 200.
- Jacq. Aust. Append.* 598. See *Jacq. Austr.*
- Jacq. Col.* 323. Id., Collectanea ad Botanicum, &c., spectantia. Vindobonæ, 1786—1790. 4 vols. 4to. Vol. V. sive Supplementum. 1795.
- Jacq. f. Ecl.* 238. See *Jacq. fl. Ecl.*
- Jacq. fl. Ecl.* 238. Jos. Franc. Von Jacquin. Eclogæ Botanicae. Vienna, 1811—1816. Fol.
- Jacq. Frag.* 400. N. J. Von Jacquin, Fragmenta Botanica. Vienna, 1800—1809. Fol.
- Jacq. Ger. Gall. Prov.* 336. See *Ger. Gallo-Prov.*
- Jacq. Hort. Schön.* 240. Id., Plantarum rariorum Horti Casarei Schönbrunensis. Vienna, 1797—1804. 4 vols. fol.
- Jacq. Hort. Vin.* 348. Id., Hortus Botanicus Vindobonensis. Vindobonæ, 1770—1776. 3 vols. fol.
- Jacq. Icon.* 297. See *Jacq. Icon. Rur.*
- Jacq. Icon. Rar.* 319. Id., Icones Plantarum rariorum. Vindobonæ, 1781—1793. 3 vols. fol.
- Jacq. Misc.* 1257. Id., Miscellanea Austriaca ad Botanicam, etc., spectantia. Vindobonæ, 1778—1781. 2 vols. 4to.
- Jacq. Obs.* 599. Id., Observationes Botanicae. Vindobonæ, 1764—1771. 4 fasc. fol.
- Jacquin Ueber den Ginkgo.* 149. Jos. Franc. Von Jacquin, Ueber den Ginkgo. Wien, 1819. 8vo.
- Jacq. Vind.* 345. See *Jacq. Hort. Vin.*
- Jam. Jour.* 2113. See *Jamieson's Journal.*
- Jamieson's Journal.* 2111. See *Edin. Phil. Journ.*
- Jamieson's Gardening.* 2608. The Theory and Practice of Gardening; wherein is fully handled all that relates to fine Gardens, commonly called Pleasure-Gardens. By John James of Greenwich. London, 1728. 4to.

- Janua Trilinguis*. 1339. *Janua Linguarum reserata*, Lat., Anglicè, et Gallicè. By John Amos Comenius. Lesnia, 1631. 8vo.
- Jesse's Gleanings*. 840. *Gleanings in Natural History*. By Edward Jesse, Esq. London, 1832. 8vo.
- Jewish Antiquities*. 1902. Moses and Aaron; or, the Civil and Ecclesiastical Rites used among the Ancient Hebrews, observed; &c. By Thomas Godwin, or Goodwyn. Oxford, 1622. 4to.
- Johnston's (Dr.) Flora of Berwick upon Tweed*. 512. *Flora of Berwick upon Tweed*. By G. Johnston, D.D. 1829 and 1831. 2 vols. 8vo. Vol. I. containing the Phanogamous, and Vol. II. the Cryptogamous, Plants.
- Johns. Ger.* 659. See *Ger. Em.*
- Johnson's Gerard*. 243. See *Ger. Em.*
- Jour. Phys.* 244. See *Journ. de Physique*.
- Journ. de Physique*. 1826. Paris, 1773, and continued. 4to.
- Jour. R. Inst.* 1491. *Journal of Science and the Arts*. Edited by W. T. Brande of the Royal Institution of Great Britain. Lond. 1816. 8vo. Continued annually.
- Journ. Acad. Sci. Philad.* 647. *Journal of the Academy of Natural Sciences of Philadelphia*. 1817, continued. 8vo.
- Journ. Hort. Tour.* 1965. See *Journal of a Horticultural Tour*.
- Journ. of Bot.* 1195. See *Journal of Botany*.
- Journ. of Geo. Soc.* 1982. *Journal of the Royal Geographical Society*. London, 1830—1837. 7 vols. 8vo.
- Journ. of Travels in Albania*. 2042. Some Account of a Journey into Albania, Romelia, and other Provinces of Turkey, in 1809 and 1810. By Sir John Cam Hobhouse. Lond. 1812. 4to.
- Journal*. 2430. See *Moorcroft's Journal*.
- Journal, &c.* 919. See *Bishop Heber's*.
- Journal d'Agriculture des Pays-Bas*. 1354. Brussels. In monthly numbers 8vo.
- Journal de Botanique*. 1844. Desvaux, *Journal de Botanique*. Paris, 1808—1814. 5 vols. 8vo.
- Journal de la Société d'Agriculture Pratique*. 1358. Paris, 1834. 8vo.
- Journal d'Hist. Nat.* 1843. *Journal d'Histoire Naturelle*. Paris, 1792. 8vo.
- Journal des Forêts*. 616. Paris. 8vo.
- Journal of a Horticultural Tour*. 1243. By Dr. Neill. Edinb. 1823. 8vo.
- Journal of a Residence in Norway*. 2173. See *Laing's Journal*.
- Journal of a Residence in Norway, during the Years 1834, 1835, and 1836*. 2304. See *Laing's Journal, &c.*
- Journal of Botany*. 1414. *The Journal of Botany*; being a Second Series of the Botanical Miscellany. By Sir W. J. Hooker, LL.D. Lond. 1834. 8vo.
- Journal of Science*. 1749. See *Jour. R. Inst.*
- Journal of the Asiatic Society of India*. 393. Lond. 1830. 8vo.
- Journal of the Transactions at Port Jackson and Norfolk Island*. 2442. See *Hunter's (Admiral) Journ.*
- Journey, &c.* 786. *Journey from India to England, in the Year 1797, &c.* By John Jackson. London, 1804. 8vo.
- Journey from Aleppo to Jerusalem*. 2410. *Journey from Aleppo to Jerusalem at Easter, 1697*. By Henry Maundrell, M.A. Oxford, 1703. 4to.
- Juss. Ann. du Mus.* 398. *Jussieu in Annales du Muséum*.
- Juss. Gen.* 377. Antoine-Laurent De Jussieu, *Genera Plantarum*. Paris, 1789. 8vo.
- Juss. Gen. ed. Usteri*. 1261. Id., Ditto ed. Usteri. Turici, 1791. 8vo.

K.

- Kæmp. Amœn. Exoticæ*. 253. Engelbert Kæmpfer, *Amœnitatum Exoticarum Politico-Physico-Medicarum Fasciculi quinque, &c.* Lengow, 1712. 4to.
- Kæmpf. Ic.* 283. Id., *Icones selectæ Plantarum, &c.* Londini, 1791. Fol.
- Kal.* 1478. See *Stang's Planter's Calendar*.
- Kalendar*. 2512. See *Calendarium Hortense*.
- Kalm Act. Succ.* 1359. Kalm in Ditto. See Ditto.
- Kalm Amœn.* 1125. See *Amœn. Acad.*
- Kalm It.* 1301. Kalm (P.), *Travels into North America*. Translated by J. Forster. Lond. 1770, 1771. 8vo.
- Kasthofer's Voyage dans les Petits Cantons, et dans les Alpes Rhétiques*. 2276. Geneva, 1827. 8vo.
- Keats's Poems*. 1320. London, 1816. 8vo.
- Fennion's Essay on Trees in Landscape*. 14. London, 1816.
- Kenrick's American Orchardist*. 989. *The New American Orchardist; or, Account of the most valuable Varieties of Fruits adapted for Cultivation in the United States*. By William Kenrick. Boston, 1837. 8vo.
- Kenrick's American Silk-Grower's Guide*. 1351. *The American Silk-Grower's Guide; or, the Art of raising the Mulberry and Silk on the System of Successive Crops in each Season*. Ed. Boston, U. S. 1835. Small 8vo.
- Ker Bot. Reg.* 358. See *Bot. Reg.*
- Key to Structural, Physiological, and Systematic Botany*. 222. See *Lindley's Key*.
- King's Survey of the Coast of Australia*. 2591. London, 1826. 8vo.
- Kirby and Spence's Introduction to Entomology*. 618. *An Introduction to Entomology; or, Elements of the Natural History of Insects, &c.* By William Kirby, M.A., &c., and William Spence, F.L.S., &c. London, 1815, 2 vols. 8vo. New ed., London, 1826, 4 vols. 8vo.
- Knipl. Cent.* 1614. Joh. Hier. Kniplhoff, *Botanica in Originali seu Herbarium vivum, Centuriæ 12*. Halle, 1757—1764. Fol.
- Knoop Pomol.* 891. Joh. Herm. Knoop, *Pomologia*. Ed. Holl., Leeuwarden, 1758. Ed. Gall., Amsterdam, 1771. Fol.
- Knorr Del.* 1049. Georg. Wolfgang Knorr, *Deiciæ Naturæ selectæ, oder Auserlesene, &c.* Nürnberg, 1766—1767. 2 vols. fol.
- Knorr His.* 238. See *Knorr Thes.*
- Knorr Thes.* 1249. Id., *Thesaurus Rei Herbariæ hortensisque universalis*. Nounberg, 1770—1772. 2 vols. fol.
- Kob.* 2145. *Walne Ursache der Baumtrökniss der Nadel-walder*. Nürnberg. 1786. 4to.

- Koch's Comm.* 1154. De Salicibus Europæis Commentatio. Auctore G. D. J. Koch. Erlangen, 1828. 12mo.
Krock. Fl. Scl. 755. Ant. Joh. Krocker, Flora Silesiaca renovata. Vratislaviæ, 1787—1790. 2 vols. 8vo.
Krock. Siles. 761. See *Krock. Fl. Sil.*
Kth. Nov. Sp. Amer. 509. See *H. B. et Kunth Nov. Gen., &c.*
Kunth Nov. Gen. Am. 481. See *H. B. et Kunth Nov. Gen. &c.*

L.

- Lab. Pl. Syr. Dec.* 702. See *Labillard. Ic. Plant., &c.*
Labill. Nov. Cal. 663. Labillardière (J. J.), Sertum Austro-Caledonicum. Paris, 1824. Fol.
Labill. Nov. Holl. 357. See *Labill. Specim.*
Labill. Pl. Syr. 1213. See *Labillard. Ic. Plant., &c.*
Labill. Specim. Nov. Holl. 2102. Jac. Jul. Labillardière, Novæ Hollandiæ Plantarum Specimen. Paris, 1804—1806. 2 vols. fol.
Labillard. Ic. Plant. Syr. 2491. Id., Icones Plantarum Syriæ rariorum. Decades v. Paris, 1791—1812. 5 vols. 4to.
Lady of the Lake. 2493. By Sir Walter Scott.
La Gérosalemme Liberata. 1382. See *Gérosalemme Liberata.*
Lag. Gen. et Spec. 333. Mar. Lagasera, Genera et Species Plantarum quæ aut Novæ aut nondum rectè cognoscuntur. Madriti, 1816. 4to.
Laing's Journal of a Residence in Norway. 2173. London, 1826. 8vo.
Laing's Norway. 1654. See *Laing's Journal.*
La Maison de Campagne. 138. Par Made. Aglaé Adanson, Paris, 1826. 8vo.
Lam. Diet. Encyc. 233. Jean Bapt. Monet de la Marek, Encyclopédie Méthodique Botanique. Paris, 1782—1796. 4 vols. 4to. Vol. V. et seq., see *Poir. Diet.*
Lam. Ill. 255. Id., Illustration des Genres. 2 vols. of text, and 900 pls.
Lamarck MS. 1452.
Lamb. Gen. Pin. 1934. See *Lambert's Monograph, &c.*
Lamb. Pin., ed. 2. 2208. See *Lambert's Monograph, &c.*
Lambert's Monograph of the Genus Pinus. 188. A Description of the Genus Pinus. By Aylmer Bourke Lambert, Esq., F.R.S., Pres. Linn. Soc. &c. London, 1822. 2 vols. 8vo.
Land Steward. 1223. The Nobleman, Gentleman, Land Steward, and Surveyor's Complete Guide, &c. London, 1787. 8vo.
Landscape-Gardener. 129. See *Davis's (Drnnis's) Landscape-Gardener.*
Language of Flowers. 1164. London, 1834. 12mo.
L'Annuaire du Cultivateur. 615. L'Annuaire, &c. Par G. Romme, Paris, 1795. 8vo.
Lanth. Ac. 420. Lanth (Th.), Dissertation de Acere. Strasbourg, 1781. 4to.
Lap. Hist. des Pl. des Pyrénées. 2209. See *La Peyr. Abr.*
Lapeyr. Abr. 553. See *La Peyr. Abr.*
La Peyr. Abr. 588. Philippe Picot de la Peyrouse, Histoire abrégée des Plantes des Pyrénées. Toulouse, 1813. 8vo.
La Peyrouse Suppl. Pl. Pyren. 2209. Id., Supplement to the above. Toulouse, 1818. 8vo.
La Rosière de Salency. 792. A Drama, in Madame de Genlis's Théâtre d'Éducation. London, 1788. 12mo.
Latr. Hist. Génér. 2140. I. atreille (P. A.), Considérations Générales sur les Animaux composant les Classes des Crustacés, des Arachnides, et des Insectes. Paris, 1810. 8vo.
Lauder's Gilpin. 90. Gilpin's Forest Scenery. Edited by Sir Thos. Dick Lauder. Edinburgh, 1834. 2 vols. 8vo.
Laur. Ros. 750. A Collection of Roses from Nature. By Miss Lawrence. London, 1799. Fol.
Laurie Todd. 2138. A novel by Galt.
Lawson's Manual. 2157. See *Agric. Manual.*
Laxm. Nov. Act. Pet. 632. Laxmann in Nova Acta Acad. Petr. See *Nova Acta, &c.*
Le Bon Jardinier. 189. See *Bon Jard.*
Le Botaniste Cultivateur. 189. See *Botaniste Cultivateur.*
Le Langage des Fleurs. 623. Paris, 12mo.
L'Echo du Monde Savant. 2529. A French periodical. Paris, 1831, and continued.
Led. Fl. Ross. Alt. Ill. 727. See *Led. Icon. &c.*
Led. Icon. Pl. Fl. Ross. 2338. C. F. Ledebour, Icones Plantarum novarum vel imperfectè cognitarum, Floram Rossicam, imprimis Altaicam, illustrantes. Riga, &c., 1829. Fol. pl. 100.
Ledeb. Ind. Hort. Dorp. Suppl. 691. Ledebour, Supplement to the Catalogue of Plants in Dorpat Garden for 1824.
Legacy, &c. 1353. By Samuel Hartlib. Ed. 2., 1651; ed. 3., 1655.
Lehm. Pot. Diss. 747. J. G. C. Lehman, Monographia Generis Potentillarum. Lipsiæ, 1820. 4to.
Leitch Ritchie's Journey to St. Petersburg and Moscow. 1836. 625. London. 8vo.
Lejeune Fl. Spa. 798. A. L. S. Lejeune, Flore des Environs de Spa. Liège, 1811—1813. 2 vols. 8vo.
Lempriere's Class. Diet. 1307. Lempriere's Classical Dictionary. 2084. London, 1792. 8vo.
Lepid. Javanica. 1820. A Descriptive Catalogue of the Lepidopterous Insects contained in the Museum of the East India Company. By Dr. T. Horsfield. London, 1828, 1829. 4to.
Les Agrémens de la Campagne, &c. 367. Leyden and Amsterdam, 1750. 4to.
Les Annales de Fromont. 189. See *Annales de Fromont.*
Les Jardins. 2172. A poem by l'Abbé Delille.
Les Obs., &c., en Grèce, Asie, et autres Pays étrangers. 1554. Pierre Belon, ou Bellon, Les Observations de plusieurs Singularités trouvées en Grèce, Asie, Judée, Égypte, Arabie. Paris, 1554. 4to. Ed. 2., Anvers, 1555. 8vo.
Les Plaisirs du Gentilhomme Campagnol. 911. A poem by Nicholas Rapin. 1883.
Les Roses. 749. Les Roses. Par J. P. Redouté; avec le Texte, par C. A. Thory. Paris, 1817. 3 vols. 4to.
Les Végétaux Résineux, &c. 2051. Les Végétaux Résineux, tant indigènes qu'exotiques. Par F. S. Duplessy. Paris, 1802. 4 vols. 8vo.
Lessing's Synopsis Generum Compositarum. 1063. Ferolmi, 1832. 8vo.
Letter on a new Way of planting Acorns. 1787. See *Method for decreasing the Consumption of Timber in the Navy, &c.* By T. Nichols.

- Letters*. 1770. Letters addressed to a young Gentleman, on the Subject of Literature, &c. By C. Davy. London, 1789. 8vo.
- Letters*. 57. Letters written by the late Jonathan Swift, and several of his Friends, from the Year 1703 to 1704. Published from the original, with Notes by J. Hawkesworth. London, 1767. 3 vols. 8vo.
- Lettre sur le Robinier*. 616. Lettre sur le Robinier, &c. By M. François de Neufchâteau. Paris, 1863. 12mo.
- L'Hér. Stirp. Nov.* 255. See *L'Hérit. Stirp. Nov.*
- L'Hérit. Corn.* 1010. C. L. L'Héritier, Cornus seu Specimen Botanicum, &c. Paris, 1788. Fol.
- L'Hérit. Diss.* 1062. Id., Dissertations, nempe Louchea, Buchozia, Michauxia, Hymenopappus, Virgilia, Michauxia. Paris. Fol.
- L'Hérit. Hort. Par.* 1419. L'Héritier in Hortus Parisiensis.
- L'Hérit. Serl.* 475. See *L'Hérit. Serl. Angl.*
- L'Hérit. Serl. Angl.* 1069. Id., Sertum Anglicum seu Plantæ rariores, &c. Paris, 1788. Fol.
- L'Hérit. Stirp.* Nov. 634. Id., Stirpes novæ aut minus cognitæ. Paris, 1784, 1785. 6 fasc. fol.
- L'Histoire Naturelle de la Rose*. 749. J. L. M. Guilleman, Histoire Naturelle de la Rose. Paris, 1800. 8vo.
- L'Homme des Champs*. 1507. L'Homme des Champs, ou les Georgiques Francaises, avec des Notes. Par J. Delisle. Strasbourg, 1799. 8vo.
- Life and Posthumous Writings of W. Cooper*. 1764. By Wm. Hayley. Chich. 1804. 4 vols. 8vo.
- Life of Constantine*. 1720. The History of the Church, and Eusebius's Life of Constantine. London, 1630. Fol.
- Life of Lord Kaims*. 94. See *Woodhouselee's Life of Lord Kaims*.
- Light*. Scot. 765. Flora Scotica; or, a Systematic Arrangement, in the Linnæan Method of the native Plants of Scotland and the Hebrides. By John Lightfoot. London, 1777, 2 vols. 8vo; 2d ed. 1789.
- Lin. Amæn.* 237. See *Lin. Amæn. Acad.*
- Lin. Amæn. Acad.* 1314. Carolus Linnæus s. Von Linné, Amœnitates academicæ, seu Diss. antehac seorsim editæ. Holmiæ et Lipsiæ, 1749 et seq., 10 vols. 8vo; ed. 2., Holmiæ, 1762 et seq.; ed. 3., cur. J. C. D. Schrebero, Erlangæ, 1787—1790.
- Lin. Eric.* 1087. Linnæus (Car.), de Erica, in Amæn. Acad. 1770.
- Lin. Fil. Supp.* 475. Carolus Linnæus filius, Supplementum Plantarum. Brunsvigæ, 1781. 8vo.
- Lin. Fl. Lapp.* 1106. Carolus Linnæus s. Von Linné, Flora Lapponica. Amstelodami, 1737, 8vo; ed. 2., cur. J. E. Smith, Eq. Londini, 1792.
- Lin. Gen.* 246. Id., Genera Plantarum. Leidæ, 1737; ed. 2., Leidæ, 1742; ed. 3., Leidæ, 1759; ed. 4., Holmiæ, 1754; ed. 5., Holmiæ, 1764; ed. 6., Viennæ, 1767; ed. 7., cur. J. J. Reichard, Francof. Man. 1775; each 1 vol. 8vo; ed. 8., cur. J. C. D. Schreber, Francof. Man. 1789—1791, 2 vols. 8vo.
- Lin. Gen. Pl. ed. Schreber*. 1390. See *Lin. Gen.*, ed. viii.
- Lin. Hort. Clif.* 1261. Id., Hortus Cliffortianus. Amstelodami, 1737. Fol.
- Lin. Hort. Ups.* 1292. Id., Hortus Upsaliensis. Stockholm, 1748. 8vo.
- Lin. Mant.* 256. Id., Mantissa Plantarum altera. Holm. 1771. 8vo.
- Lin. Mat. Med.* 532. Id., Materia Medica. Holmiæ, 1749. Ed. Schreber, 1772.
- Lin. Scb. Thes.* 542. See *Schb. Thesaur.*
- Lin. Sp.* 233. See *Lin. Sp. Pl.*
- Lin. Sp. Plan.* 522. Id., Species Plantarum. Ed. 1., Holmiæ, 1753, 2 vols. 8vo; ed. 2., Holmiæ, 1762, 1763, 2 vols. 8vo; ed. 3., Vindobonæ, 1764; ed. 4., by J. J. Reichard, Francof. Man. 1779, 1780, 4 vols. 8vo; ed. 5., by C. L. Willdenow, Berolini, 1797—1810, 5 vols. 8vo.
- Lin. Syst. Nat.* 1893. Linnæus (Carl.), Systema Naturæ. Lugd. Bat. 1755. Folio.
- Lin. Syst. Veg.* 595. J. A. Murray, C. Linnæi Systema Vegetabilium. Gottingæ et Gothæ, 1744; Gottingæ, 1784; Parisiis, 1798.
- Lin. Trans.* 55. Transactions of the Linnæan Society of London. London, 1791, and continued. 17 vols. 4to.
- Lind. Wicksb.* 1706. Linder seu Linderstolpe (J.), Flora Wicksbergensis. Stockholm, 1760, 8vo; ed. 2., 1728.
- Lindl. Coll.* 932. Collectanea Botanica; or, Figures, &c., of rare and curious Exotic Plants. By Dr. Lindley, F.R.S., &c. Lond. 1821. Fol.
- Lindl. Hort. Trans.* 701. Dr. Lindley in the Horticultural Society's Transactions. See *Hort. Trans.*
- Lindl. Intro. to Bot.* 694. Id., Introduction to Botany. London, 1832, 8vo; ed. 2., London, 1835.
- Lindl. Intro. to N. S.* 492. See *Lindley's Introduction to the Natural System*.
- Lindl. Lin. Trans.* 931. Dr. Lindley in the Linnæan Society's Transactions. See *Lin. Trans.*
- Lindl. MSS.* 1934. Dr. Lindley's MSS.
- Lindl. Nat. Syst. of Bot.* 935. See *Lindley's Introduction to the Natural System*.
- Lindl. Ros. Mon.* 748. Id., Rosarium Monographia. London, 1820. 8vo.
- Lindl. Synops.* 28. Id., A Synopsis of the British Flora, &c. London, 1829. Small 8vo. Ed. 2. 1835.
- Lindley's Introduction to the Natural System.* 222. Id., An Introduction to the Natural System of Botany. London, 1830, 8vo. Ed. 2. 1835.
- Lindley's Key.* 265. Id., A Key to Structural, Physiological, and Systematic Botany, for the use of Classes. By Dr. Lindley, F.R.S., &c. London, 1835. 8vo.
- Link Enum.* 327. H. F. Link, Enumeratio Plantarum Horti Regii Botanici Berolinensis altera. Berol. 1821, 1822. 2 vols. 8vo.
- Link et Otto Abbild.* 965. See *Abbild. der Deutschen Holzarten*.
- Link Bert. Abhand.* 2398. See *Abhand. Acad. Berl.*
- Link Jahrb.* 1204. H. F. Link, Jahrbücher der Gewächskunde. Berlin, 1820. 8vo.
- Linnaea.* 948. F. L. Von Schlechtendahl, Linnaea; ein Journal für die Botanik in ihrem ganzem Umfange. Berlin und Halle, 1836—1838. 10 vols. 8vo. Continued.
- Linnaean Correspondence.* 54. A Selection of the Correspondence of Linnæus and other Naturalists, from the Original Manuscripts. By Sir J. E. Smith. London, 1821. 2 vols. 8vo.
- List of the Plants collected in Ross's Voyage.* 1602. By R. Brown, in Ross's Voyage. London, 1836. 4to.
- Literary Panorama.* 1778. A weekly periodical. London, 1815.
- Lob. Adv.* 586. See *L'Obel's Adversaria*.
- Lob. Icon.* 549. Mathias De Lobel seu Lobelius, Stirpium Icones. Antverpiæ, 1591. 4to.
- L'Obel's Adversaria.* 55. Id., Stirpium Adversaria nova, &c. London, 1605. Fol.
- Lodd. Bot. Cab.* 230. The Botanical Cabinet. By Conrad Loddiges and Sons. London, 1817—1834. 20 vols. 12mo and 4to.

- Lodd. Cat.* 473. A Catalogue of Plants, &c., in the Hackney Establishment. By Messrs. Loddiges. Published annually. 12mo.
- Loeffl. Res.* 1079. Petr. Löfling, Rega til spanska länderna. Ed. Linn. Stockh. 1758, 8vo; ed. Germ. Berl. 1766, 8vo.
- Loes. Pruss.* 1678. Joh. Loeselius, Flora Prussica. Regiomonti, 1703. 4to.
- Lois. Fl. Gall.* 586. Loiseleur Deslongchamps, Flora Gallica. Parisii, 1806, 2 vols. 12mo.
- Lois. Herb. Annot.* 259. M. Loiseleur Deslongchamps, in *Herbier de l'Amateur*, a French periodical.
- Lois. Not.* 575. Id., Notice sur les Plantes à ajouter à la Flore de France. Paris, 1810. 8vo.
- Lois. Nouv. Dict.* 977. Loiseleur des Longchamps (J. L. A. M.), Nouveau Dictionnaire d'Histoire Naturelle. 1816, continued. 36 vols.
- Loiseleur Deslongchamps's Histoire du Cèdre.* 2411. See *Histoire du Cèdre*.
- London and Wise's Retired Gardener.* 361 The Retired Gardener: being a Translation of *Le Jardinier Solitaire*, &c. By George London and Henry Wise. London, 1766. 2 vols. 8vo.
- London Horticultural Society's Catalogue of Fruits.* 758. See *Hort. Soc. Cat. of Fruits*.
- London Horticultural Society's Transactions.* 103. See *Hort. Trans.*
- Lonicer's History of Plants.* 609. Lonicerus (A.), *Botanicon Plantarum Historia earundem ad vivum artificiosè expressis Iconibus.* Frankfurt 1665. Folio.
- London's H. B.* 389. See *Hort. Brit.*
- Lour. Coch.* 235. See *Lour. Fl. Coch.*
- Lour. Fl. Coch.* 679. Joannes De Loureiro, Flora Cochinchinensis. Ulyssipone, 1790, 2 vols. 4to; Ed. 2., cur. C. L. Willdenow, Berolini, 1793, 2 vols. 8vo.
- Louth's Trans.* 1717. New Translation of Isaiah, &c. By the Rev. R. Louth. London, 1791. 12mo.
- Ludw. Ectypa Veg.* 1343. C. G. Ludwig, Ectypa Vegetabilium Usibus medicis destinatum, &c. Hallæ, 1760. Fol.
- Lyon Herb.* 1114. Lyon's Herbarium.
- Lyonnet Mém. Posth.* 1821. Lyonnet (P.), Recherches sur l'Anatomie et les Métamorphoses des différentes Espèces d'Insectes. Ouvrage posthume, publié par M. W. De Haau. Paris, 1832. 4to.
- Lyson's Environs of London.* 43. Environs of London, &c. By the Rev. Daniel Lysons, A.M., &c. London, 1796. 4 vols. 4to.

M.

- Macbeth.* 919. One of Shakspeare's Plays.
- Mackay Fl. Hibern.* 2153. See *Mackay's Irish Flora*.
- Mackay's Flora Hibernica.* 1118. See *Mackay's Irish Flora*.
- Mackay's Irish Flora.* 107. Flora Hibernica; comprising the Flowering Plants, Ferns, Characeæ, Musci, Hepaticæ, Lichenes, and Algae of Ireland, &c. By J. T. Mackay, M.R.I.A., &c. Dublin, 1836. 8vo.
- Mackay's List of Irish Plants.* 575. A Catalogue of the Plants found in Ireland, with Descriptions of some of rarer sorts. Id. Dublin, 1825. 4to.
- Mag. of Bot. and Zool.* 2147. Magazine of Zoology and Botany, Conducted by Sir W. Jardine, Bart.; P. J. Selby, Esq.; and Dr. Johnston, in 18'6, 1837. Continued now (1838) under the title of *Annals of Natural History* (including Geology), superintended by the same gentlemen, in conjunction with Sir W. J. Hooker. London, 8vo. Published every two months.
- Magaz. Encyclop.* 1828. Millin's Magazin Encyclopédique, ou Journal des Sciences, &c. Paris, 1793, &c. 8vo.
- Magazine of Nat. Hist.* 21. Magazine of Natural History. Conducted by J. C. Loudon, F.L.S., &c. Lond. 1828. 8vo, publishing in monthly numbers. 10 vols. now published (1838).
- Magn. Bot.* 336. See *Magnol Bot.*
- Magn. Hort. Monsp.* 606. See *Magn. Monsp.*
- Magn. Monsp.* 1806. Id., Hortus Regius Monspeliensis. Monspeli, 1697. 8vo.
- Magnol Bot.* 889. Petrus Magnol, Botanicon Monspeliense. Monspeli, 1686. 12mo.
- Maitland's Hist. of Edin.* 87. Maitland's (W.) History of Edinburgh, from its Foundation to the present Time. Edinburgh, 1753. Folio.
- Malcolm's London.* 786. Londinum Redivivum; or, an Ancient History, and Modern Description, of London. Lond. 1802—1805. 4 vols. 4to.
- Man.* 2154. See *Agricult. Manual*.
- Manning and Bray's Surrey.* 25. The History and Antiquities of Surrey. By Owen Manning. Lond. 1804. Folio.
- Mant.* 692. See *Lin. Mant.*
- Mantissa.* 2093. See *Lin. Mant.*
- Manwood on Forest Law.* 1751. Treatise on the Laws of the Forest and of the Purlicu. By John Manwood. Lond. 1565. Sm. 1to.
- Marnock's Fl. Mag.* 2542. Marnock's (R.) Floricultural Magazine and Miscellany of Gardening. Lond. 1836. In monthly Nos. 8vo.
- Marol's Recueil des Plans, &c. de plusieurs des Châteaux, Grottes, &c.* 1338. Paris, 1661.
- Marquis, Essai sur les Harmonies Végétales et Animales du Chêne.* 1722. See *Essai sur les Harmonies, &c.*
- Marsh. Arb. Amer.* 499. Humphrey Marshall, Arbustum Americanum; the American Grove, &c. Philadelph. 1785. 8vo.
- Marsh. Plant.* 1309. See *Marshall, &c.*
- Marshall on Planting and Rural Ornament.* 1090. Planting and Rural Ornament, &c. Lond. 1785, 1 vol. 8vo; ed. 2., Lond. 1793, 2 vols. 8vo. Mr. Marshall's name is not put to either edition.
- Mart. Fl. Rust.* 1751. Flora Rustica. By Thomas Martyn. Lond. 1792—1794. 4 vols. 8vo.
- Mart. Mill.* 319. See *Martyn's Miller*.
- Martin et Rios, Disquisitionum Magicorum Libri Sex.* 2020. Lugd. 1608. Folio.
- Martyn's Mill.* 303. See *Martyn's Miller's Dictionary*.
- Martyn's Miller's Dictionary.* 280. Gardener's Dictionary; or, a complete System of Horticulture. By Philip Miller, F.R.S. Lond. 1759, 3 vols. fol.; improved edition, edited by Professor Martyn, Lond. 1807, 4 vols. fol.

- Masque of Queens.* 2365. By Ben Jonson.
- Materia Medica.* 1241. Fr. Geoffroy, Tractatus de Materiâ Medicâ. Paris, 1741, 3 vols. 8vo. Venetiis, 1742, 2 vols. 4to.
- Matthew on Naval Timber,* &c. 1380. On Naval Timber and Arboriculture; with critical Notes on Authors who have recently treated on the subject of Planting. By Patrick Matthew. Lond. 1831. Royal 8vo.
- Math. Hist.* 1313. Matthioli (P. A.), Commentaria in Dioscoridem de Materiâ Medicâ. Venice, 1558. Fol.
- Matth. Valgr.* 972. Matthioli Opera quæ extant omnia, edidit C. Bauhin. France, 1598. Fol.
- Maund's Botanic Garden.* 253. The Botanic Garden; or, Magazine of Hardy Flowering Plants cultivated in Great Britain. By B. Maund, Esq., F.L.S. Begun in 1824, and continued monthly.
- M'Cull. Dict.* 677. See *M'Culloch's Dictionary*, &c.
- M'Culloch's Dictionary of Commerce.* 2284. A Dictionary, Practical, Theoretical, and Historical, of Commerce, &c. By J. R. M'Culloch. Ed. 2. Lond. 1834. 8vo.
- Mech. Mag.* 465. Mechanic's Magazine. Lond. 1822—1837. 27 vols. 8vo. Continued.
- Med. Fl.* 479. Medical Flora; or, Manual of the Medical Botany of the United States of North America. By C. S. Ranneque. Philadelphia, 1828. 2 vols. 8vo.
- Med. Gesch.* 874. Medicus (F.), Geschichte der Botanik unterer Zeiten. Munich, 1793. 8vo.
- Medical Botany.* 465. See *Stephenson and Churchill's Medical Botany.*
- Medical and Philosophical Essays.* 2389. Essays, Medical and Experimental; to which are added Select Histories of Diseases. By J. Percival, M.D. Lond. 1767—1778. 3 vols. 8vo.
- Meerb. Leon.* 1012. Nicol. Meerburg, Plantarum selectarum leones pictæ. Lugd. Bat. 1798. Fol.
- Melville's (Lord Letter to Spencer Percival, Esq.* 1809. Pamph. 8vo. 1810.
- Mem. et Corr. of Pepys.* 2603. Memoirs of Samuel Pepys, Esq., &c. London, 1828. 5 vols. 8vo.
- Mém. de Desmarêts in Journ. de Physique.* 1996. See *Journal de Physique.*
- Mém. de l'Acad. des Scien.* 1722. See *Mémoires de l'Acad. des Sciences.*
- Mém. de la Russie Méridionale.* 687. By Descezet.
- Mém. de l'Institut.* 1385. Mémoires de l'Institut National des Sciences Mathématiques et Physiques. Paris, 1797, and continued. 4to.
- Mém. Mus.* 1982. Mémoires du Muséum d'Histoire Naturelle. Paris, 1815, 1816. 4to.
- Mém. Scien. Mathé. de l'Institut,* &c. 1757. See *Mém. de l'Inst.*
- Mém. Soc. Imp. Mosq.* 762. See *Mém. Soc. Imp. Nat. Mosc.*
- Mém. Soc. Nat. Cur. Mos.* 1432. See *Mém. Soc. Imp. Nat. Mosc.*
- Mém. Soc. Imp. Nat. Mosq.* 1932. Mémoires de la Société Impériale des Naturalistes de Moscou. Moscow, from 1785. 4to.
- Mém. sur l'Hist. Nat. du Chêne.* 1722. Mémoires, &c. Par M. De Secondât. Paris, 1782. Fol.
- Mém. Wern. Soc.* 1257. Memoirs of the Wernerian Natural History Society. Edinburgh, 1811, and continued. 8vo.
- Memoir.* 614. Memoir on the common Acacia. Paris, 1786.
- Memoir addressed to the Society of Arts, on the Planting and Rearing of Forest Trees,* &c. 617. By W. Withers, Esq.
- Memoir on the Date Palm.* 142. By A. Michaux.
- Memoir on the Roads of Cephalonia.* 2328. By Major-General Napier. Lond. 1835. 8vo.
- Mémoire sur les Chênes.* 1908. Louis Bosc, Mémoires sur les différentes Espèces de Chênes qui croissent en France. Paris, 1808. 4to.
- Mémoire sur le Figuier.* 1369. By Bernard, in the Journal de Physique. Tom. 29.
- Mémoire sur le Zéikoua.* 1409. See *Michx. Mém. sur le Zéikoua.*
- Mémoire sur les Marrons d'Inde.* 147. Par M. Bon de St. Hilaire. Paris, 12mo, and in 4to.
- Mémoire sur les Pins.* 140. By Fougeroux de Bondaroy; published in the Mémoires de l'Académie des sciences.
- Mémoires.* 2145. Mémoires pour servir à l'Histoire des Insectes. Par C. De Geer. Stockholm, 1752—1778. Tom. 7. 4to.
- Mémoires de l'Académie des Sciences.* 140. Mémoires de Mathématique et de Physique, présentés à l'Académie Royale des Sciences, par divers Savans. Paris, 1750, and continued. 4to.
- Mémoires d'Agriculture.* 135. Mémoires d'Agriculture, d'Economie Rurale et Domestique, &c. Paris, 1785—1791. 24 vols. 8vo.
- Mémoires présentés à l'Académie des Sciences de St. Petersburg.* 176. See *Nov. Act. Pet.*
- Mémoires sur les Conifères.* 2096. Mémoires sur les Conifères et les Cycadées; ouvrage posthume de L. C. Richard, terminé et publié par Achille Richard fils. Stuttgart et Paris, 1826. 8vo.
- Mémoires sur les différentes Espèces de Chêne qui croissent en France, et sur ces E'trangers à l'Empire qui se cultivent dans les Jardins et Pépinières des Environs de Paris,* &c. 1722. By Ludov. Bosc. Published in the Mém. de l'Institut. National de France, for 1807.
- Memoirs.* 2203. See *Walpole's Memoirs.*
- Memoirs of the American Academy of Arts.* 614. Memoirs of the American Academy of Arts and Sciences. Boston and Charlestown, 1785, and continued. 4to.
- Memoirs of the Literary Society of Manchester.* 1651. Memoirs of the Literary and Philosophical Society of Manchester. Warrington, 1785—1816. 8vo.
- Memoirs of the Missionaries in the Levant.* 2409. A Collection of the most interesting Missionary Voyages, &c. London, 1805. 12mo.
- Memoirs of the Royal Academy of Sciences at Madrid.* 2436. See *Anales de las Cienc. Nat.*
- Memoria Balfouriana.* 86. See *Sibbald's Memoria Balfouriana.*
- Memorials of Oxford.* 2076. By Dr. Ingram. Oxford, 1832—1837. 3 vols. 8vo.
- Mer. Fl. Par.* 765. F. V. Mérat. Nouvelle Flore des Environs de Paris. Paris, 1812. 8vo.
- Merry Wives.* 1755. Shakspeare's Merry Wives of Windsor.
- Methods for decreasing the Consumption of Timber in the Navy,* &c. 1787. By T. Nichols. London, 1793. 8vo.
- Methodus Plantarum.* 53. See *Ray's Methodus Plantarum.*
- Metzger's Castle of Heidelberg.* 147. Historical Description of the Castle of Heidelberg Gardens, &c. Heidelberg, 1830.

- Mexica*. 2185. Ward's (H. G.) Mexico in 1827. London, 2 vols. 8vo.
- Mey. Verz. Pflanz. Cauc*. 2587. Meyer C. A.'s, Verzeichniss der Pflanzen welche während der, &c. Petersburg, 1831. 8vo.
- McGregor's Sketches of the Maritime Colonies of British America*. 2116. London, 1828. 8vo.
- Mich. Arb.* 261. See *Michaux's Arbres forestiers, &c.*
- Mich. Fl. Bor. Amer.* 237. André Michaux, Flora Boreali-Americana. Parisiis, 1802. 2 vols. 8vo.
- Mich. Gen.* 1126. P. A. Micheli, Nova Plantarum Genera. Florentiæ, 1729. Fol.
- Michaux's Arbres de l'Amérique*. 135. See *Michaux's Arbres, &c.*
- Michaux's Arbres Forestières de l'Amérique*. 2119. André Franç. Michaux fils, Histoire des Arbres forestiers de l'Amérique Septentrionale. Paris, 1810—1813. 3 vols. 4to.
- Michaux's Histoire des Chênes Américains*. 1861. André Michaux, Histoire des Chênes de l'Amérique Septentrionale. Paris, 1801, fol.; ed. Germ. 3 Kerne, 1862.
- Michx. fil. N. Amer. Syl.* 1873. André François Michaux, North-American Sylva; or, a Description of the Forest Trees of the United States, Canada, and Nova Scotia. Translated from the French by A. L. Hillhouse. Paris, 1819. 3 vols. 8vo.
- Michx. Mém. sur le Zekoua*. 1109. A. Franç. Michaux, Mémoire sur le Zekoua. Paris, 1831.
- Michx. Quer.* 1869. See *Michaux's Histoire des Chênes*.
- Midsummer Night's Dream*. 901. One of Shakspeare's Plays.
- Mill. Ic.* 261. Figures of the Plants described in the Gard. Diet. By Philip Miller. London, 1760. 2 vols. fol.
- Mill. Illustr.* 339. Johan. Miller, Illustratio Systematis Sexualis Linnæi. Londini, 1777. Fol.
- Miller's Dictionary*. 54. The Gardener's Dictionary. By Philip Miller. London, 1731; ed. 2. 1733; ed. 3., 1737; ed. 4., 1741; ed. 5., 1747; ed. 6., 1752; ed. 7., 1759; ed. 8., 1768; ed. 9., see *Mortly's Miller*.
- Milne's Botanical Dictionary*. 77. A Botanical Dictionary. By Colin Milne. London, 1770. 8vo.
- Minutes of the Method adopted by Mr. Robert Turner of raising Oaks*. 1804. In Gard. Mag., vol. viii.
- Mirbel Mém. Mus.* 2462. Mirbel in Mémoires du Muséum. See *Mém. Mus.*
- Mirror*. 2021. The Mirror of Literature, Amusement, and Instruction. London, begun in 1822, and continued in weekly Nos. 8vo.
- Misc. Writings, &c.* 2114. Evelyn's Miscellaneous Writings, edited by Upeott. London, royal 4to. 1825.
- Miscellaneous Tracts relating to Natural History, &c.* 1332. By Benjamin Stillingfleet. London, 1759, 8vo; Ed. 2., 1762; Suppl. by F. J. Brand, 1781.
- Missionary Journal*. 2411. See *Wolf's Missionary Journal*.
- Mith. Gen.* 377. See *Mich. Gen.*
- Mith. Act. Acad. Nat. Cur.* 1007. See *Nov. Act. Nat. Cur.*
- McLeay's Report to the Treasury*. 1388. Report to the Treasury upon the State of the Elms in St. James's and Hyde Parks. Published in the Edin. Phil. Journ., No. xxxi. art. 12.
- McNab's Hints on the Planting and General Treatment of Hardy Evergreenes*. 1176. See *Hints on the Planting, &c.*
- McNab's Treatise on Cope Heaths*. 1091. A Treatise on the Propagation, Cultivation, and general Treatment of Cope Heaths in a Climate where they require Protection during the Winter Months. By William McNab, Superintendent of the Bot. Gard. Edin., &c. Edinb. and Lond. 1832. 8vo.
- Moc. et Sesse Fl. Mex. Icon. ined.* 843. See *Moc. et Sess. Fl. Mex.*
- Moc. et Sesse Fl. Mex. Ic. ined.* 713. See *Fl. Mex. Icon. ined.*
- Moc. Fl. Nuth.* Mocino (Josef), Drawings of Plants collected by him at Nootka Sound. Not published.
- Moench Lob. Adv.* 1243. Moench in ditto. See *Ditto*.
- Moench Meth.* 253. Conrad Moench, Methodus Plantas Horti et Agri Marburgensis describendi. Marburgi, 1794. 8vo.
- Moench Suppl.* 277. Id., Supplementum ad Methodum, &c. Marburgi, 1802. 8vo.
- Moench Weiss.* 772. See *Moench Weissenst.*
- Moench Weissenst.* 1657. Id., Verzeichniss ausländischer Bäume des Lust-chlosses Weissenstein. Francf. 1785. 8vo.
- Mol. Sag. sulla Stor. Nat. del Chili*. 243. Giov. Igu. Molina, Saggio sulla Storia naturale del Chili. Bologna, 1782. 8vo.
- Molina's History of Chili, &c.* 2436. London, 1809. 2 vols. 8vo.
- Monog. Hymen. Ichu. Affin.* 1827. See *Hymen. Monog.*
- Monog. Ros.* 748. Proneville A. De, Nomenclature raisonnée des Espèces, Variétés, et Sous-variétés du Genre Rosier. Paris, 1818. 1 vol. 8vo.
- Monograph.* 2432. See *Lamb. Gen. Pin.*
- Monograph of American Vines*. 481. Rafinesque's Monograph of the American Vines in his Medical Flora, vol. 2.
- Monteith's Forester's Guide*. 89. The Forester's Guide, &c. By — Monteith. Stirling, 1819. 12mo, plates.
- Monthly Review*. 1765. London, 1749, and continued till 1808.
- Moorcroft's Journal*. 2130. Moorcroft's Travels in Ladak, Kashwir, Bokhara, &c. London, 1837.
- Moore's American*. 791. Odes of Anacreon, translated into English Verse, with Notes. By Thomas Moore, Esq. London. 2 vols. 12mo.
- Mor. Hist.* 256. Joan Bapt. Morand, Historia Botanica practica. Mediolani, 1744. Fol.
- Mor. Præful.* 1241. Robert Morison, Præfulda Botanica, pars 1. London, 1669. 8vo.
- Morrel's Théorie des Jardins*. 188. Théorie des Jardins. By N. Morcl. Paris, 1776. 8vo. Improved edition by C. Morcl, 1802.
- Moris. Elench.* 586. See *Elench. Sard.*
- Moris. Oxon.* 931. Robert Morison, Plantarum Historia universalis Oxoniensis. Oxonii, 1680. 2 vols. folio.
- Moris Strip. Sard.* 578. See *Elench. Sard.*
- Morning Chronicle*. 141. A Common daily New-paper.
- Morris Fl. Consp.* 931. Richard Morris, Flora Conspiciua. London, 1820. 8vo.
- Musclay's Essay on Archery*. 2087. Worcester, 1799. 8vo.
- Much Ado about Nothing*. 1014. One of Shakspeare's plays.

- Mühl. Cat.* 479. See *Mühlenberg's Catalogue, &c.*
Mühl. Nov. Act. Scrut. Berol. 1234. Mühlberg in *Nov. Act., &c. See Act. Nat. Scrut., &c.*
Mühlenb. Nov. Act. Soc. Nat. Scrut. Berol. 1504. See *Mühl. Nov., &c.*
Mühlenb., Sims, et Kön. Ann. of Bot. 1504. Mühlberg, Sims, and König, in *Ann. Bot.* See *Ann. of Bot.*
Mühlenberg's Catalogue of North American Plants. 729. Henr. Mühlberg, *Catalogus Plantarum Americae Septentrionalis* Lancaster, 1813. 8vo.
Münch. Haus. 489. See *Münch. Haus.*
Münch. Hausv. 754. Barron Otto Von Münchäusen, *Monatliche Beschäftigungen für einen Baum- und Pflanzen-gärtner, als eine Zugabe zum 5ten Th. des Hausvater.* Hanover, 1771. 8vo.
Mundy's Needwood Forest. 1785. A poem.
Murray on the Silkworm. 1357. See *Murray's Observations, &c.*
Murray's Encyclopædia of Geography. 172. An Encyclopædia of Geography, comprising a complete Description of the Earth, physical, statistical, civil, and political, &c. By Hugh Murray. F.R.S.E., &c. London, 1833. 8vo.
Murray's Observations on the Silkworm. 1358. *Observations on the Silkworm, &c.* By John Murray. London.
Murr. Nov. Comm. Gött. 1048. J. A. Murray in *Nov. Comm., &c.* See *Nov. Comm. Gott.*
Museum Rusticum. 1380. *Museum Rusticum et Commerciale; or, Select Papers on Agriculture, Commerce, &c.* Revised and digested by several Members of the Society for the Encouragement of Arts, Manufactures, and Commerce. London, 1764. 6 vols. 8vo.
Museum Tradescantianum. 53. *Museum Tradescantianum; or, a Collection of Rarities preserved at South Lambeth, near London.* By John Tradescant. London, 1656. 12mo.
Mr. Fl. Bor. Amer. 303. See *Mich. Fl. Bor. Amer.*

N.

- Nat. Hist. Jorât.* 1955. *Histoire Naturelle du Jorât, &c.* Par M. le Comte G. De Razoumowsky. Lausanne, 1789. 2 vols. 8vo.
Nat. Hist. of the County of Dublin, 465. An Essay towards a Natural History of the County of Dublin, &c. By John Ruttly, M.D. Dublin, 1772. 2 vols. 8vo.
Nat. Syst. of Bot. 2558. See *Lindl. Nat. Syst. of Bot.*
Natural History. 696. *Caii Plinii Secundi Historiæ Naturalis.* Venice, 1496. Folio.
Natural History of Carolina. 69. See *Catesb. Carol.*
Natural History of Oxfordshire. 1221. By Dr. R. Plot. Oxford, 1705. Fol.
N. Cours d'Agr. 315. See *Nouveau Cours d'Agriculture.*
N. Duh. 278. See *Du Ham. Arb. Nour.*
N. D. Ham., 476. See *Du Ham. Arb. Nour.*
Neal Cat. Hort. Blackb. 2186. A Catalogue of Plants in the Garden of John Blackburn, Esq., &c. By Adam Neale, Gardener. London, 1779. 8vo.
Neck. Elem. 528. *Nat. Jos. De Necker, Elementa Botanica secundum Systema omericum, seu naturale.* Neowede ad Rhenum, 1790. 3 vols. 8vo.
Neck. Gall. 1078. See *Neck. Gallo-Bell.*
Neck. Gallo-Bell. 1266. Id., *Deliciae Gallo-Belgiæ sylvestris.* Argentorati, 1768. 2 vols. 12mo.
Needwood Forest. 1769. See *Mundy's Needwood Forest.*
Nees Hoya Phys. 538. C. G. Nees Von Esenbeck, *Hora Physicæ Berolinensis, collectæ ex Symbolis Virorum doctorum Link, Rudolphi, &c.* Bonna, 1820. Fol.
Nees Nov. Act. Nat. Cur. Bonn. 956. See *Nov. Act.*
Nees (T.) ab Esenb. Gen. Pl. Fl. Germ. 2032. See *Nees Von Esenbeck, &c.*
Nees Von Esenbeck Gen. Pl. 2549. *Genera Plantarum Floræ Germanicæ Iconibus et descriptionibus illustrata.* Auctore T. F. L. Nees ab Esenbeck, M.D. Bonna, 1833. 8vo.
Neill's Horticultural Tour. 145. *Neill's Horticultural Tour through France and the Netherlands.* Edinb. 1823. 8vo.
Nestl. Pot. 747. See *Nestl. Pot. Diss.*
Nestl. Pot. Diss. 747. C. G. Nestler, *Monographia de Potentilla.* Parisiis et Argentorati, 1816. 4to.
New Botanist's Guide, 882. *The New Botanist's Guide to the Localities of the rarer Plants of Britain.* London, By Hewett Cottrell Watson. London, 1835—1837. 2 vols. 8vo.
New Descriptive Catalogue of Roses cultivated in the Sawbridgeworth Nursery. 749. By T. Rivers, jun. Sawbridgeworth, 1831. Pamph. 8vo.
New Improvements. 2072. See *Bradley's New Improvements, &c.*
New Planter's Calendar. 2059. *The Planter's Calendar; or, the Nurseryman and Ferester's Guide in the Operations of the Nursery, the Forest, and the Grove.* By the late Walter Nicol, edited and completed by E. Sang. Edinb. 1830. 8vo.
New Principles of Gardening. 80. *New Principles of Gardening; or, the Laying out Pastures, Groves, Wildernesses, Labyrinths, Avenues, Parks, &c.* London, 1756. 4to.
New York Farm, 1356. *American Farmer.* In periodical Nos. New York. 4to.
Newman's Dictionary. 314. *A Marine Pocket Dictionary of the Italian, Spanish, Portuguese, and German Languages, &c.* By Henry Newman. London, 1799. 12mo.
Nich. Obs., &c. 1788. *Observations on the Propagation and Management of Oak Trees in general, but more particularly applying to His Majesty's New Forest in Hampshire.* Southampton, 1791. 8vo.
Nichol's Hist. Leicest. 2413. *Collections towards the History and Antiquities of the Town and County of Leicester.* London, 1798—1796. 2 vols. 4to.
Nichol's Illustrations. 786. *Nichols (J.), Illustrations of the Manners and Expenses of Ancient Times in England.* London, 1797. 4to.
Nicholai, Die Wäuder, oder Processions Raupe, &c. 1820. Berlin, 1833. 8vo.
Nicolson's Historical Scottish Library. 86. London, 1702. 8vo.
Newman's Dictionary. 303. See *Newman's Dictionary.*
Nois. Arb. Fruitiers. 1359. See *Nois. Jard. Fruit.*
Noisette Jard. Fruit. 675. *Louis Noisette, Le Jardin Fruitier.* Paris, 1813. 2 fasc. 4to.
Nomenclator, 1186. See *Steucl's Nomenclator.*
Nomenclature raisonnée. 749. *Aug. de Pronville, Nomenclature raisonnée des Espèces, Variétés, et Sous-variétés du Genre Rosier.* Paris, 1818. 8vo.
North-American Flora. 142. See *Mich. Fl. Bor. Amer.*
North-American Sylva. 1379. See *Mich. fl. N. Amer. Syl.*

- Note Historique sur le Ginkgo.* 2097. Recherches sur l'Histoire des Cyprès, et Note Historique sur le Ginkgo biloba. Par M. J. L. A. Loiseleur Deslongchamps. Paris, 1854. Pamph. 8vo.
- Nouv. Dict. d'Hist. Nat.* 2445. See *Nouveau Dictionnaire*, &c.
- Nouv. Duh.* 239. See *Du Ham Arb. Nouv.*
- Nouv. Journ. de Physique.* 1823. See *Journal de Physique.*
- Nouv. Mém. de la Soc. Imp. des Nat. de Moscou.* 365. See *Mém. Soc. Imp.*, &c.
- Nouveau Cours d'Agriculture.* 675. See *Nouveau Cours complet.*
- Nouveau Cours complet* 135. *Nouveau Cours complet, ou Dictionnaire raisonné d'Agriculture.* Par les Membres de la Section d'Agr. de l'Institut. Paris, 1809. 16 vols. 8vo.
- Nouveau Dictionnaire d'Histoire Naturelle.* 465. Paris, 1819. 36 vols. 8vo.
- Nouveau Du Hamel.* 138. See *Du Ham. Arb. Nouv.*
- Nov. Act. N. C. App.* 2350. See *Nov. Act. Nat. Cur.*, &c.
- Nov. Act. Nat. Cur. Bonn.* 936. See *Nova Acta Naturæ Curiosorum.*
- Nov. Act. Pet.* 632. *Nova Acta Academiarum imperialis Petropolitane.* Petropoli, 1783—1837. 4to.
- Nov. Act. Soc. Nat. Scrut. Brevol.* 1504. See *Act. Soc. Scrut. Brevol.*
- Nova Acta Naturæ Curiosorum.* 2141. *Nova Acta Physico-Medica Academiæ Cæsareæ Leopoldino-Carolinæ Naturæ Curiosorum.* Bonnæ et Nuremberg, 1757 to the present Time. 4to
- Nov. Comm. Göttl.* 1043. *Novi Commentarii Societatis Regiæ Scientiarum Göttingensis.* 1751 to the present Time. 4to.
- Nouv. Hollandiæ Plantarum Specimen.* 556. By J. J. La Billardière. Parisiis, 1791—1812. 5 vols. 4to.
- Nursery Catalogue.* 2015. See *Furber's Nursery Catalogue.*
- Nutt. Gen. Amer.* 303. Thomas Nuttall, *The Genera of North-American Plants, and a Catalogue of the Species.* Philadelphia, 1818. 2 vols. 12mo.

O.

- Observ. Entomol.* 2141. *Dr. Heer's Entomological Observations.* 1836.
- Observations des Singularités.* 1910. *Pierre Belon, ou Bellon, Les Observations de plusieurs Singularités trouvées en Grèce, &c.* Paris, 1554, 4to; ed. 2. Anvers, 1555, 8vo.
- Observations, &c., in Scotland.* 42. *Observations relative chiefly to Picturesque Beauty, made in the Year 1776, on several Parts of Great Britain, and particularly the Highlands of Scotland.* By W. Gilpin, A.M., &c. Lond. 1789. 2 vols. 8vo.
- Observations faites dans un Voyage, &c.* 2407. *Pallas (P. S.), Observations faites dans un Voyage entrepris dans les Gouvernemens Méridionaux de l'Empire de Russie pendant les Années 1793 et 1794.* Leipzig, 1801. 2 vols. 4to.
- Observations on Oak Trees, &c., in a Letter to the Earl of Chatham, 1787.* See *Nichols Obs.*
- Observations on the Coniferæ.* 2104. See *Observations on the Coniferæ of New Zealand.*
- Observations on the Coniferæ of New Zealand.* 2101. By Mr. G. Bennet, published in Lambert's *Pinus.*
- Observations on the Diseases of Trees.* 1213. *William Forsyth, Observations on the Diseases of Trees, &c.* London, 1791. 8vo.
- Odyssey.* 835. See *Hom. Odys.*
- Œd. Fl. Dan.* 364. *Icones Plantarum sponte nascentium in Regnis Daniæ et Norvegiæ, &c. Hafniæ, 1761—1770, vols. 1, 2, and 3, Auctore G. C. Eder, & vols. 4. and 5., Auct. O. F. Müller, 1771—1782; vols. 6. and 7., Auct. M. Vahl, 1787—1803; vol. 8., Auct. J. W. Horneman, 1806—1816.* 8 vols. fol.
- Ochlof. Abbild.* 2065. *C. C. Celhafen, Abbildung der wilden Bäume, &c.* Nürnberg, 1767—1773. 4to.
- Office of Woods, &c.* 1804. *A Parliamentary Report.*
- Oldfield's Anecdotes of Archery.* 2087. *Anecdotes of Archery, ancient and modern.* By H. G. Oldfield. London, 1791. 12mo.
- Oliv. Enc. Méth.* 1929. See *Encyc. Méth.*
- Oliv. Voy.* 1847. See *Olivier's Travels.*
- Oliv. Voy. dans l'Emp. Ottom.* 1928. See *Olivier's Travels.*
- Olivier's Travels.* 1860. *Travels in the Ottoman Empire, Egypt, and Persia.* By G. A. Olivier. Lond. 1801. 2 vols. and atlas. 4to.
- Olivier's Voyage.* 813. See *Olivier's Travels.*
- On Nav. Tim.* 1953. See *On Naval Timber, &c.*
- On Naval Timber.* 1400. See *Matthew on Naval Timber.*
- On Permanent Colours.* 1911. *Experimental Researches concerning the Philosophy of Permanent Colours, and the best Means of producing them, by dyeing, Calico-printing, &c.* By E. Bancroft, M.D. London, 1794. 8vo.
- Ort. Decad.* 559. *Cas. Gomez De Ortega, Novarum aut rariorum Plantarum Horti R. Matritensis Centurie.* Matriti, 1797—1798. 8 vols. 4to. Tit. gen. 1800.
- Osh. Itin.* 541. *Pehr Osbeck, Dagbock åter en Östendyck Resa.* Stöckholm, 1757. 8vo.
- Osterreiches Allgemeine Baumzucht.* See *Höss Aulcit.*
- Otto's Garten Zeitung.* 1365. See *Allgemeine Gartenzeitung.*
- Outl. Geog. Distr.* 7-5. See *Watson's Outlines.*
- Outlines, &c.* 257. See *Watson, &c.*
- Outlines of Botany.* 237. *Outlines of Botany; including a General History of the Vegetable Kingdom, &c.* By Professor Burnett. London, 1855. 2 vols. 8vo.
- Ovid. Met.* 141. *Ovid's Metamorphoses.* London, 1773. 2 vols. 12mo.

P.

- Pall. Astr.* 637. *Peter Simon Pallas, Species Astragalorum descriptæ et Iconibus illustratæ.* Lipsiæ, 1800. Fol.
- Pall. Flor. Ross.* 247. See *Pallas's Flora Rossica.*
- Pallas's Flora Rossica.* 156. 1d., *Flora Rossica.* Petropoli, 1784 et 1788. 2 vols. fol.
- Pall. Itin. ed. Gall.* 599. *Voyages dans l'Empire Russe.* Paris, 1793. 8 vols. 8vo, avec atlas. 4to.
- Pall. Itin. ed. Gall. Append.* 582. See *Pall. Itin. ed. Gall.*
- Pall. Nov. Act. Pet.* 632. *Pallas in Nov. Act. Pet.* See *Nov. Act. Pet.*

- Pall. Nord. Beytr.* 910. Id., Neue nordische Beiträge zur physikalischen, &c. Petersb. und Leipz. 1781—1796. 7 vols. fol.
- Panzer Fauna Ins. Germ.* 1483. See *Faun. Ins.*
- Paradise Lost.* 2122. A poem by Milton.
- Paradise of Plants.* 515. Adam in Eden; or, Nature's Paradise, &c. By William Cole, Herbalist. London, 1657. Fol.
- Paradise and the Peri.* 2040. A poem by Moore.
- Paradisus.* 717. See *Parkinson's Paradisi in sole.*
- Parad. Lond.* 813. *Paradisus Londinensis.* By R. A. Salisbury. London, 1805—1808. 2 vols. 4to.
- Parke's China.* 847. Historie of the Great and Mightie Kingdom of China, and the Situation thereof. Translated out of Spanish by R. Parke. London, 1588. 4to.
- Park, Hampstead.* 1391. The Topography and Natural History of Hampstead, in the County of Middlesex. With an Appendix of Original Records. By John James Park. London, 1814. 8vo.
- Parkinson's Herbal, or Theatre of Plants.* 52. The Theatre of Plants; or, an Herball of a large extent; with numerous wood engravings. By John Parkinson. Lond. 1640. Fol.
- Parkinson's Paradisi in sole.* 52. *Paradisi in sole Paradisus Terrestris;* or, a Garden of all Sorts of Pleasant Flowers, which our English Ayre will admitt to be noursed up; with a Kitchen Garden, Orcharde, &c. By John Parkinson, Apothecary of London. London, 1629. Fol.
- Parkinson's Theat. Bot.* 1908. See *Parkinson's Herbal, &c.*
- Par. Lon.* 283. See *Parad. Lond.*
- Pascal's (Dr.) Treatise on the Mulberry, &c.* 1558. New York, 1830. 8vo.
- Pausanias Lac.* 1441. Pausanias's Description of Greece
- Pav. Diss. in Mém. Acad. Reg. Med. Mot.* 2432. Jos. Pavon, Dissertacion botanica sobre los Generos Tovalia, Actinophyllum, &c. Madrid. 4to. As quoted in *Mém. Acad.*, &c.
- Paxton's Mag. of Bot.* 1149. Magazine of Botany, and Register of Flowering Plants. By J. Paxton, F.L.S., &c. London, begun in 1834, and continued in monthly Numbers, large 8vo.
- Pèlerinage à Jérusalem, &c.* 2412. par R. P. M. J. Geramb. Paris, 1854.
- Pen. Cyc.* 306. See *Penny Cyc.*
- Pennant's London.* 2070. Account of London. By Thomas Pennant. London, 1790. 4to.
- Pennywick's (Dr.) Works, in Prose and Verse.* 94 Geographical and Historical Description of the Shire of Tweeddale; with a Miscellany, and curious Collection of select Scottish Poems. 1715 and 1814. 4to.
- Penny Cyc.* 299. The Penny Cyclopædia of the Society for the Diffusion of Useful Knowledge. London. In monthly parts, folio. Begun in 1833, and 10 vols. published.
- Penny Magazine.* 1336. A periodical paper published in London, weekly. Begun in 1832.
- Perambulations of Kent.* 25. Perambulation of Kent; containing the Description, History, and Customs of that Shire.
- Perce Forest.* 786. A Romance.
- Perennial Calendar.* 1023. The Perennial Calendar and Companion to the Almanack, illustrating the Events of every Day in the Year, &c. By T. Forster, M.B., F.L.S., &c. London, 1824. 8vo.
- Pers. Ench.* 331. C. II Persoon, Synopsis Plantarum seu Enchiridium Botanicum. Paris, 1805 and 1807. 2 vols. 12mo.
- Pers. Ic. Pict.* 2148. Id., Icones pictæ Specierum rariorum Fungorum, &c. Paris et Strasb. 1803. 2 fasc. 4to, pl. 18.
- Pers. Myc. Eur.* 2147. Id., Mycologia Europæa. Erlang, 1822—1828. 3 vols. 8vo, pl. 30.
- Pers. Obs. Mycol.* 1836. Id., Observationes Mycologicæ. Lipsiæ, 1796. 8vo.
- Pers. Syn.* 243. See *Pers. Ench.*
- Persia in Miniature.* 786. By Sir Robert Ker Porter.
- Perthshire Report.* 2354. General View of the Agriculture in the County of Perth, &c. By Dr. James Robertson. Perth, 1799. 8vo.
- Pettiv. Gaz.* 1116. James Pettiver, *Gazophyllacium Naturæ et Artis.* Londini, 1702—1704. 5 decad. fol.
- Pettiv. Mus.* 2559. Id., *Musæi Pettiveriani Centuriæ Decem.* London, 1695. 8vo.
- Ph. Fl. Amer. Sept.* 259. Frederick Pursh, *Flora Americana Septentrionalis.* London, 1814. 2 vols. 8vo.
- Pharsalia.* 1724. Lucan's *Pharsalia.* Rowe's Translation.
- Phil. Mag.* 1388. Philosophical Magazine and Journal. By Alex. Tilloch and Richard Taylor. London, 1798—1826. 68 vols. 8vo, continued under the name of Philosophical Magazine and Annals of Chemistry. By R. Taylor and R. Phillips. London, 1827—1832. 11 vols. 8vo.
- Phil. Mag. 3 ser.* 738. A further Continuation of the Philosophical Magazine, under the name of the London and Edinburgh Philosophical Magazine and Journal of Science. By Sir David Brewster, R. Taylor, and R. Phillips. London, 1832, continued. 8vo.
- Phil. Trans.* 23. The Philosophical Transactions of the Royal Society of London. Commenced in 1665, and continued to the present Time. London. 4to.
- Phil. Trans. Abr.* 49. An Abridgment of the *Phil. Trans.* By C. Hutton, LL.D.; G. Shaw, M.D.; and R. Pearson, M.D. London, 1804—1809. 13 vols. 4to.
- Philosophical Conferences of the Virtuosi of France.* 1821. A Collection of Discourses of the Virtuosi of France upon Questions of Philosophy. Translated into English by G. Havers. London, 1664.
- Physical and Historical Researches into Rhabdomancy, &c.* 2020. Translated from the Italian of Carlo Amoretti into German, with supplementary Treatises, by Ritter. Berlin, 1809. 8vo.
- Phys. Vég.* 215. See *Physiologie Végétale.*
- Physiologie Végétale.* 215. Physiologie Végétale, ou Exposition des Forces et des Fonctions Vitales des Végétaux, &c. By M. A. P. De Candolle. Paris, 1832. 3 vols. 8vo.
- Physique Occult, ou Traité de la Baguette divinatoire.* 2020. La Physique Occulte, ou Traité de la Baguette Divinatoire. Par Pierre le Lorrain de Vallemont. Amst. 1693. 12mo.
- Phytographia.* 53. See *Pluk. Phyt.*
- Picture Bible.* 165. The Pictorial Bible; being the Old and New Testaments, according to the authorised Version, illustrated with many hundred woodcuts. London, 1836. In monthly parts, 8vo.
- Pièces relatives à la Culture et aux Usages de cet Arbre [Robinia].* 617. Lettre sur le Robinier, connus sous le Nom impropre de Faux Acacia; avec plusieurs Pièces relatives à la Culture et Usage de cet Arbre. Par M. François De Neufchâteau. Paris, 1803. 12mo.
- Pitt's Survey of Worcestershire.* 896. General Survey of the County of Worcester. London, 1810. 8vo.
- Pitt's Trans.* 2334. Pitt's Translation of Virgil.

- Pl. Off.* 1691. See *Ehrh. Pl. Off.*
Pl. Rar. Hort. Gen. 503. See *Dec. Pl. Rar. Jard. Gen.*
Plant. and Rur. Orn. 1774. See *Marshall on Planting and Rural Ornament.*
Plant. Amer. 274. See *Plum. Plant. Amer.*
Plant. Kal. 509. Planter's Kalender. By W. Nicol. Edited and completed by Edw. Sang. Edin. 1820. Fd. 2., 8vo.
Plante Equinoctiales. 1941. Humboldt et Bonpland, *Plante Equinoxiales.* Paris, 1808—1816. 2 vols. fol.
Plante per Galliam, Hispaniam, et Italiam observatae. 2415. By Jaco. Barrelier. A posthumous work, edited by Bern. Jussieu. Paris, 1714. Fol.
Planter and Forest Pruner. 188. The Forest Pruner, or Timber Owner's Assistant, &c. London, 1895. 8vo.
Planter's Calendar. 89. See *Plant. Kal.*
Planter's Guide. 188. The Planter's Guide; or, a Practical Essay on the best Method of giving immediate Effect to Wood, &c. By Sir Henry Steuart, Bart. &c. Edin. 1829. 8vo.
Plantes du Dauphiné. 2276. See *Villars's Plantes du Dauphiné.*
Planting and Rural Ornament. 188. See *Marshall on Planting and Rural Ornament.*
Plenk Icon. 463. J. J. Plenk, *Icones Plantarum medicinalium.* 1803 et seq. Fol.
Pluc. Epist. 1354. The Letters of Pliny the Consul, with occasional Remarks by William Melmoth, Esq. London, 1746. 2 vols. 8vo.
Plin. Hist. Nat. 1914. Plinii *Historia Naturalis.* Hardouin's edit. Paris, 1793. 3 vols. fol. Translated under the title of Pliny's Natural History of the World. By Philemon Holland. London, 1601. 2 vols. fol.
Plin's (Dr.) History of Oxfordshire. 1768. See *Natural History of Oxfordshire.*
Pluk. Alm. 1256. See *Pluk. Alm. Phyt.*
Pluk. Alm. Phyt. 68. Léonard Plukenett, *Almagestum Botanicum sive Phytologia Onomasticon.* Londini, 1796. 4to.
Pluk. Amath. 379. Id., *Amaltheum Botanicum.* Londini, 1705. 4to.
Pluk. Mant. 257. Id., *Almagesti Botanici Mantissa.* Londini, 1700. 4to.
Pluk. Phyt. 727. See *Plukenet's Phytographia.*
Plukenet's Phytographia. 53. Id., *Phytographia sive Stirpium illustriorum, &c.* Londini, 1691. 4 vols. 4to.
Plum. Cat. 2271. Plumier (C.), *Catalogus Plantarum Americanarum.* Paris, 1703. 4to.
Plum. Icon. 479. Plumier (C.), *Description des Plantes de l'Amérique, avec leurs Figures.* Paris, 1695. Folio.
Plum. Plant. Amer. 2271. Charles Plumier, *Description des Plantes de l'Amérique.* Paris, 1693 and 1713. Fol.
Pococke Itin. 2654. See *Pococke Orient.*
Pococke Orient. 480. Description of the East, and some other Countries. By R. Pococke. London, 1743—1745. 2 vols. fol.
Poeppig's Travels in the Peruvian Andes. 2434. Leipzig, 1830.
Pohl Böhem. 760. J. J. Pohl, *Versuch einer Flora Böhmens.* Prag. 1810 and 1814. 2 vols. 8vo.
Poir. Dict. 480. See *Poiret Encyc. Méth.*
Poir. Dict. Encyc. Suppl. 1928. See *Poiret Encyc. Suppl.*
Poir. Encyc. Méth. 1543. See *Poiret Encyc. Méth.*
Poir. Encyc. Suppl. 975. See *Poiret Encyc. Suppl.*
Poir. Suppl. 519. See *Poiret Encyc. Suppl.*
Poiret Encyc. Méth. 1-77. J. L. M. Poiret, *Encyclopédie Méthodique, ou Dictionnaire de Botanique.* (Voy. Lam. Dictionn. continué par Poiret.) Paris, 1804—1808; being the 5th, 6th, 7th, and 8th vols. of Lamareck's *Encyclopédie Méthodique, &c.*
Poiret Encyc. Suppl. 1415. Id., *Encyclopédie Méthodique Supplément au Dictionn. de Botanique.* Paris, 1810—1816. 4 vols. 4to.
Poiret's Hist. Phil. des Plantes. 1915. Id., *Histoire Philosophique, &c., des Plantes de l'Europe.* Paris, vols. 1—3. 8vo.
Poit. Arb. Fr. 475. See *Poit. et Turp. Arb. Fruit.*
Poit. et Turp. Arb. Fruit. 5018. Poiteau et Turpin, *Traité des Arbres fruitiers de Du Hamel.* Nouvelle édition. Paris, 1808 et seq. 15 fasc. Fol.
Political Register. 616. The Political Register; a periodical paper. Edited by Wm. Cobbett. London.
Poll. Fl. Fer. 584. Ciro Pollini, *Horti et Provinciae Veronensis Plantæ novæ vel minus cognitæ.* Pavie, 1816.
Pollich Pall. 1950. J. A. Pollich, *Historia Plantarum in Pallatinato sponte nascentium.* Mannheim, 1776. 3 vols. 4to.
Pom. Brit. 896. See *Pomarium Britannicum.*
Pom. Mag. 248. The Pomological Magazine. Lond. 1833. 3 vols. 8vo.
Pomarium Britannicum. 923. *Pomarium Britannicum; an Historical and Botanical Account of Fruits known in Great Britain.* By Henry Phillips. London, 1820. 8vo.
Pomona. 896. *Pomona Herefordiensis; or, a Descriptive Account of the old Cider and Perry Fruits of Herefordshire.* By T. A. Knight, F.R.S., &c. London, 1819. 4to.
Ponted. Comp. 1555. Jul. Pontedera, *Compendium Tabularum Botanicarum.* Patavii, 1719. 4to.
Pouty's Prof. Planter. 1522. The Profitable Planter, &c. By William Pouty. Huddersfield, 1800. 8vo.
Pope's Odyssey. 1654. Pope's Translation of Homer's *Odyssey.*
Porter's (Miss Jane) Scottish Cliffs. 1772. London. 5 vols. 8vo.
Porter's Sir Robert Ker Travels. 786. London, 2 vols. 4to.
Pott's Survey of Worcestershire. 897. See *Pitt's Survey, &c.*
Pouchet Bot. appliqué. 2453. Professor of Botany at Rouen.
Pourc. Act. Toul. 594. See *Pourc. Act. Toul.*
Pourc. Act. Toul. 326. Pourcet in Actis Academiæ Tolosanæ.
Pract. Hints on Planting. 1462. See *Pract. Treat. on Planting.*
Pract. Treat. on Planting. 849. Hayes.
Practic Husbandman. 98. Maxwell (R.), *The Practical Husbandman; being a collection of Miscellaneous Papers on Husbandry.* Edinb. 1757. 8vo.
Practical Planter. 188. *The Practical Planter; or, a Treatise on Forest Planting, &c.* By Walter Nicoll. Edinb. 1799. 8vo.
Price's Essay on the Picturesque. 744. *An Essay on the Picturesque, as compared with the Sublime and the Beautiful, &c.* By Uvedale Price, Esq. London, 1794. 8vo.

- Prince's Catalogue.* 310. A Catalogue of Fruit and Ornamental Trees and Plants (all of which are hardy) cultivated at the Linnean Botanic Garden, W. Prince, Proprietor, Flushing, Long Island, near New York. New York, 1839. 8vo.
- Proc. of Geol. Soc. of Lond.* 2110. Proceedings of the Geological Society of London.
- Prod.* 309. See *Dec. Prod.*
- Prodromé de la Monographie du Genre Rosier.* 749. Prodrome de la Monographie des Espèces et Variétés communes du Genre Rosier, &c. By C. A. Thory. Paris, 1820. 12mo.
- Prodromus.* 189. See *Dec. Prod.*
- Prodromus Floræ Nepalensis.* 1921. D. Don, Prodromus Floræ Nepalensis. London, 1825. 12mo.
- Pronv. Nom.* 1753. See *Pronville's Nomen. Rais.*
- Pronville's Nomen. Rais.* 749. Aug. De Fronville, Nomenclature raisonnée des Espèces, Variétés, et Sous-variétés du Genre Rosier. Paris, 1818. 8vo.
- Pursh Fl. Bor. Amer.* 238. Fred. Pursh, Flora Borealis Americana. London, 1814. 2 vols. 8vo.

Q.

- Quarterly Journal of Agriculture.* 182. The Quarterly Journal of Agriculture; and the Prize Essays and Transactions of the Highland Society of Scotland. Edin. 1828. In quarterly Nos. 8vo.
- Quarterly Review.* 22. The Quarterly Review. London. In quarterly Nos. 8vo.
- Quayle's Jersey and Guernsey.* 1286. General View of the Agriculture and present State of the Islands on the Coast of Normandy subject to the Crown of Great Britain. By Thomas Quayle, Esq. London, 1815. 8vo.
- Quip for an upstart Courtier.* 486. A Quip for an upstart Courtier; or, a quaint Dispute between Velvet Breeches and Cloth Breeches. London, 1592. 4to.
- Quintinge's Complete Gardener.* 46. The Complete Gardener, &c. Englished by John Evelyn. Lond. 1693. Fol.

R.

- R. et P. Fl. Per. et Chil. Prod.* 544. H. Ruiz et Jos. Pavon, Floræ Peruvianæ et Chilensis Prodromus. Matriti, 1794, fol.; ed. 2., Romæ, 1797, 4to.
- R. et P. Fl. Per. et Chil. Syst.* 544. Id., Systema Vegetabilium Floræ Peruvianæ et Chilensis. Matriti, 1798. 8vo.
- R. et P. Fl. Per. Syst.* 543. See *R. et P. Fl. Per. et Chil. Syst.*
- Raf. Med. Flor.* 518. Medical Flora; or, Manual of the Medical Botany of the United States of North America. By C. S. Rafinesque. Philadelphia, 1828. 2 vols. 8vo.
- Rafin. Journ. Phys.* 503. Rafinesque in Journal de Physique. Paris, 1773, and continued. 4to.
- Rafin. Méd. Bot.* 1017. See *Raf. Med. Flor.*
- Rafin. Préc. Déc. Som.* 588. Rafinesque-Schmaltz, Précis des Découvertes Somnologiques ou Zoologiques et Botaniques. Palerme, 1814. 18mo.
- Rafin. Ros. Amer. in Ann. Phys.* 766. Rafinesque on American Roses, in Annals of Physic. New York.
- Rafin. Speech.* 582. Rafinesque-Schmaltz, Specchio delle Scienze, o Giornale enciclopedico di Sicilia. 1814.
- Rape of Proserpine.* 2472. A poem by Claudian.
- Rau. Enum.* 765. Ambr. Rau, Enumeratio Rosarum circa Wirceburgum sponte crescentium. Norinbergæ, 1816. 8vo.
- Rauw. Itin.* 646. Rauwolf's Journey through Syria, Mesopotamia, Palestine, and Egypt. Translated from the English by Stapherst. 1693. 8vo.
- Ray's General History of Plants.* 53. See *Ray's Historia Plantarum.*
- Ray's Historia Plantarum.* 41. Joh. Ray, Historia Plantarum. Londini, 1686 et 1688. 2 vols. fol. Vol. 3., seu Supplementum, 1704.
- Ray's Letters.* 3c. 2414. Philosophical Letters between Mr. Ray and several Correspondents, &c. Lond. 1718. 8vo.
- Ray's Methodus Plantarum.* 53. Id., Methodus Plantarum nova. Londini, 1682. Amstel. 1682. 8vo.
- R. Br. Prod.* R. Brown, Prodromus Floræ Novæ Hollandiæ et Insulæ van Diemen. Londini, 1810. 8vo.
- Rehb. Icon. Exot.* 1062. Reichenbach (L.), Iconographia Botanica Exotica. Leips. 1827—1830. 4to.
- Rehb. Pl. Crit.* 1210. Reichenbach (W. G. L.), Iconographia Botanica, seu Plantæ Criticæ. Leips. 1825—1828. 4to.
- Réaumur. Mém.* 1971. See *Réaumur's Mémoires.*
- Réaumur Mémoires.* 1820. Mémoires pour servir à l'Histoire des Insectes. Par R. A. F. De Réaumur. Paris, 1734—1742. 6 vols. 4to.
- Rech. Hist.* 1717. See *Recherches Historiques, &c.*
- Recherch. sur l'Anat., &c., de différ. Espèces d'Insectes.* 1819. See *Lyonet Mém Post.*
- Recherches Historiques sur les Chênes.* 1722. By M. Marquis, Professor of Botany at Rouen.
- Recherches sur la Découverte de la Rose,* 3c. 787. Langles (L.), Recherches sur la Découverte de l'Essence de Rose. Paris, 1804. 12mo.
- Recreations of Agriculture.* 615. Recreations in Agriculture, Natural History, Arts, and Miscellaneous Literature. By Dr. Anderson. London, 1799. 5 vols. 8vo.
- Recueil des Plans, &c., de plusieurs des Châteaux, Grottes, &c.* 1338. See *Marot's Recueil des Plans, &c.*
- Red. Ros.* 750. See *Red. et Thor. Ros.*
- Red. et Thor. Ros.* 752. P. J. Redouté, Les Roses. Pl. 169. color.: avec le texte. par Cl. Ant. Thory. Paris, 1817—1824. 3 vols. fol.
- Rees's Cycl.* 582. See *Rees's Cyclopædia.*
- Rees's Cyclopædia.* 1917. London, 1819. 39 vols. 4to.
- Regensb. Bot. Zeit.* 1633. Regensburgs Botanische Zeitung. Regensburg, 1803. 8vo. 1 vol. per ann.

- Régn. Bot.* 2464. See *Régnauld Bot. Icon.*
Régnauld Bot. Icon. 1:566. De Regnauld, La Botanique mise à la Portée de tout le Monde. Paris, 1774. Fol.
Reich. Mag. 656. Heichenbach (G. C.), Magazin des Pflanzenreichs. 1793, 1794. 4to.
Reid's Hist. and Lit. Bot. 899. See *Reid's Historical and Literary Botany.*
Reid's Historical and Literary Botany. 486. Historical and Literary Botany; containing the Qualities, Anecdotes, and Superstitions of Plants, &c. To which is added, Flowers; or, the Artists: a Romantic Story, from the French of Madame De Genlis. By Eliza P. Reid Windsor, 1826. 3 vols. 12mo.
Rem. on For. Scen. 2047. See *Remarks on Forest Scenery.*
Remarks on Forest Scenery. 14. By William Gilpin. 2 vols. 8vo.
Remarkable Oaks, &c. 1781. See *Rooke's Remarkable Oaks.*
Renault Fl. de l'Orne. 1212. P. A. Renault, Flore du Département de l'Orne. Alençon, 1804. 8vo.
Report on the Natural Forests of Scotland. 2180. By Mr. Grigor, Nurseryman, of Elgin. Published in the Highland Society's Transactions, vol. xii.
Reports, &c. 2170. Parliamentary Reports on the Timber Trade for 1831—1835.
Residence in Constantinople. 2042. Walsh R, A Residence at Constantinople during the Commencement, Progress, and Termination of the Greek and Turkish Revolutions. London, 1856. 2 vols. 8vo.
Ret. Gard. 2008. See *Retired Gardener.*
Retired Gardener. 364. See *London and Wise's Retired Gardener.*
Retz. Obs. 255. See *Retz. Obs. Bot.*
Retz. Obs. Bot. 754. A. J. Retzous, Observationes Botanicæ. London, 1774. 6 fasc. 4to. Ed. 2. Lipsiæ, 1779—1791. 6 vols. 4to.
Rêverie du Promeneur solitaire. 13:6. By J. J. Rousseau.
Revue Entomologique. 2145. See *Silbermann's Revue Entomologique.*
Reyn. Act. Laus. 769. See *Reyn. Mém. Laus.*
Reyn. Mém. 585. See *Reyn. Mém. Laus.*
Reyn. Mém. Laus. 772. Reynier in Mém. de la Soc. de Lausanne.
Rhizotomi. 1724. By Sophocles.
Rich. Diss. 1061. G. G. Richter, Diss. de Muscorum Notis et Salubritate. Göttingæ, 1747. 4to.
Rich Mém. Conf. 2063. See *Richard Mém. sur les Conifères.*
Richard II. 436. One of Shakspeare's plays.
Richard Mém. sur les Conifères. 2103. See *Mémoires sur les Conifères et les Cycadées.*
Richard's Analyse du Fruit. 558. Louis-Claude Richard, Analyse du Fruit, considéré en général; publiée par Duval. Paris, 1808. 8vo.
Richard's Elements of Botany. 1024. New Elements of Botany, &c. By A. Richard, M.D.; translated, with Notes, by P. Clinton; Dublin, 1829. 8vo. Translation by W. Macgillivray; London, 1831, 8vo.
Riv. Mon. Irr. 1042. Aug. Quirinus Rivinus, Ordo Plantarum Flore irregulari Monopetalo. Lipsiæ, 1699. Fol.
R. Mal. 1420. Henricus Van Rheede, Hortus Indicus Malabaricus. 1678—1708. 12 vols. fol.
Robert's English Bowman; or, Tracts on Archery. 2087. London, 1801. 8vo.
Roderick, the Last of the Goths. 1917. A poem by Southey.
Rom. Arch. 1052. Joh. Jac. Roemer, Archiv für die Botanik. Leipzig, 1795—1805. 3 vols. 4to.
Rom. et Schult. Syst. Vég. 1414. Id. et Schultes, Ed. nov. Systematis Vegetabilium Linnæi. Turici, 1815. 8vo.
Rokeby. 2084. A poem by Sir Walter Scott.
Romance of Nature. 952. The Romance of Nature. By Louisa Anne Twamley. London, 1837. 8vo.
Romance of the Rose. 941. By Chaucer.
Rooke's Remarkable Oaks, &c. 1781. Description and Sketches of some remarkable Oaks in the Park at Welbeck, in the County of Nottingham, a seat of His Grace the Duke of Portland. By Hayman Rooke. London, 1790. 4to.
Roque's Hist. des Champ. 1704. Histoire des Champignons. Par M. Roque. Paris. 4to.
Rosarium Monographia. 749. See *Lindley's Rosarium Monographia.*
Rose Amateur's Guide. 2560. The Rose Amateur's Guide. By T. Rivers, jun. London, 1857. 8vo.
Rösel Ins. Belust. Suppl. 1826. Rösel, Der monatlich herausgegebenen Insecten Belustigung. Nürnberg, 1746—1761. 4 vols. 4to.
Rosig. Ros. 752. Rossignol (K. G.), Les Roses dessinées et enluminées d'après Nature, avec une Description botanique. Traduit de l'Allemand par M. De Laffite. Leips. 4to.
Rossmässler Forstins. 2140. Rossmässler, Naturgeschichte derjenigen Insecten, welche den bei uns angebauten Holzarten am meisten schädlich werden. Leipz. 1834. 8vo.
Roth Cat. 575. Alb. Wilh. Roth, Catalecta Botanica. Lipsiæ, 1797—1805. 3 vols. 8vo.
Roth Fl. Germ. 635. See *Roth's Flora Germanica.*
Roth Germ. 1731. See *Roth's Flora Germanica.*
Roth's Flora Germanica. 146. Id., Tentamen Floræ Germanicæ. Lipsiæ, 1788—1801. 3 vols. 8vo.
Roxb. Lucan. 1462. Rowe's Translation of Lucan.
Rox. Fl. Ind. 1201. See *Roxb. Fl. Ind. ined.*
Rox. Hort. Beng. 1934. W. Roxburgh, Hortus Bengalensis; or, a Catalogue, &c. Calcutta, 1814. 8vo.
Roxb. Cor. 645. See *Roxb. Corom.*
Roxb. Corom., 1595. Id., Plants of the Coast of Coromandel, published under the direction of Sir Joseph Banks. London, 1795—1798. 2 vols. fol.
Roxb. Fl. Ind. ined. 751. Flora Indica. By W. Roxburgh. Serampore, 1820—1824. 2 vols. 8vo.
Roy. Cours d'Agric. 258. Roy in Nouveau Cours d'Agriculture.
Roy. Lugdb. 1289. Adrianus Van Royen, Floræ Leydensis Prodromus, exhibens Plantas Horti Lugduno-Batavi. Lugd. Bat. 1740. 8vo. Auctor dicitur C. Linnæus.
Royal Society's Transactions. 77. See *Phil. Trans.*
Royl. Illustr. Bot. Himalaya. 245. Illustrations of the Botany and other Branches of the Natural History of the Himalayan Mountains, and of the Flora of Cashmere. London, 1833—1838. 4to.
Rubi Germanici. 733. See *Wache and Nees's Rubi Germanici.*

- Rudb. Lapp.* 1598. Rudbeck (Olaus), Mora-Samdad, sive Lapponia Illustrata. Succo-Germ. et Lat. Upsal, 1701. 4to.
- Rudge's Survey of Gloucestershire.* 1760. General View of the Agriculture of the County of Gloucester. By T. Rudge. London, 1807. Svo.
- Ruiz et Pav. Fl. Per.* 934. See *R. et P. Fl. Per.*
- Rumph. Amb.* 1940. See *Rumph. Amboyn.*
- Rumph. Amboyn.* 2447. Geo. Everh. Rumphius, Herbarium Amboinense. Amstelodami, 1750. 6 vols. fol. Auctarium, 1755.
- S.
- Sab. Hort.* 1259. Liberatus Sabbati, Hortus Romanus. Vol. I., Auct. G. Bonelli et L. Sabbati; Vols. II. III. IV. and V., Auct. N. Martelli et L. Sabbati; Vols. VI. and VII., Auct. Const. Sabbati et N. Martelli. Roma, 1772—1784. 7 vols. fol.
- Sal. Hort. Allert.* 813. See *Sal. Prod.*
- Sal. Par.* 282. See *Sal. Par. Lond.*
- Sal. Par. Lond.* 380. See *Parad. Lond.*
- Sal. Prod.* 490. R. A. Salisbury, Prodrum Stirpium in Horto ad Chapel Allerton. Londini, 1796. Svo.
- Sal. Wob.* 1454. Salictum Woburnense. By His Grace the Duke of Bedford. Royal Svo.
- Salices.* 188. Salices. By Dr. Wade. Dublin, 1811. Svo.
- Salicetum; or, the Botanical History and Cultivation of Willows.* 1458. By Dr. Walker. Contained in his Vol. of Essays.
- Salictum Woburnense.* 129. See *Sal. Wob.*
- Salish. Prod.* 233. See *Sal. Prod.*
- Sang's Nicol.* 1226. See *Plant. Kal.*
- Santi Viagg.* 1046. Geo. Santi, Viaggi al Mont Amiata. Pisa, 1795, 8vo; e per la Toscana, Vols. II. and III., 1798 and 1806.
- Sat. Mag.* 1851. The Saturday Magazine. A periodical paper, published in London, weekly. Begun in 1832.
- Sauv. Mousp.* 1520. Franç. Boissier De Sauvages, Methodus Foliorum S. Plantæ Mospelienses juxta Foliorum Ordinem digestæ. La Haye, 1751. 8vo.
- Sav. Alb. Tosc.* 929. Gaetano Savi, Trattato degli Alberi della Toscana. Firenze, 1801, 2 vols. 8vo; ed. 2. 1811.
- Sav. Trät.* 1241. See *Sav. Alb. Tosc.*
- Savi's Bibl. Ital.* Savi (C.) in Bibliotheca Italica.
- Savi Fl. Pis.* 765. Id., Flora Pisana. Pisis, 1798. 2 vols. 8vo.
- Savi's Botanicon Etruscum.* 164. Id., Botanicon Etruscum. Pisis, 1808 and 1815. 2 vols. Svo.
- Scandinavia.* 1591. Travels in various parts of Europe, Asia, and Africa. By E. D. Clarke. London, 1810—1823. 6 vols. 4to.
- Scheuchz. H.* 2186. Joh. Jac. Scheuchzer, Itineraria Alpina. Ed. 2. Lugd. Bat. 1723. 4to.
- Schëide et Deppe MSS.* 2373.
- Schkuhr H.* 296. Christ. Schkuhr, Botanisches Handbuch. Wittenberg, 1791—1803. 3 vols. 8vo.
- Schkuhr Handb.* 598. See *Schkuhr H.*
- Schlecht. Linnæa.* 586. A. Schlechtendal Filius, Linnæa. Berol. 1826, and continued. Svo.
- Schlechtend. Berl. Mag.* 748. A. Schlechtendal in Berl. Mag.
- Schleich. Cat.* 769. See *Schleicher Cat.*
- Schleicher Cat.* 1596. J. C. Schleicher, Catalogi Plantarum Helvetiæ exsiccatarum, 1794—1816.
- Schm. Arb.* 415. See *Schmidt Baum.*
- Schmidt Baum.* 493. F. Schmidt, Oesterreichs allgemeine Baumzucht. Wien, 1792 und 1794. 2 vols. fol.
- Schmidt Fl. Boh.* 1267. F. W. Schmidt, Flora Bohemica. Pragæ, 1793, 1794. 2 vols. fol.
- Schmidt Myc.* 2148. See *Schmidt S. M.*
- Schoepf. Mat. Med. Amer.* 1534. Joh. Dav. Schoepf, Materia Medica Americana. Erlang., 1787. Svo.
- Schoolmistress.* 1282. A poem by Shenstone.
- Schrad. Diss.* 951. See *Schrad. Diss. Philad.*
- Schrad. Diss. Philad.*, 952. Schrader in De Candolle's Prodrum.
- Schrad. Gen. Ill.* H. A. Schrader, Genera nonnulla Observationibus illustrata. Göttingæ, 1808. 4to.
- Schrader Index Sem. Hort. Acad. Gött.* 828. Schrader (C. F.), Index Plantarum Horti Botanici Pedagogii Regii Glauchensis. Hall., 1772. 12mo.
- Schrader Hort. Gött. MSS.* 1600. See above.
- Schranck Baicr. (Bavar.) Fl.* 1517. Franz Von Paula Schrank, Baiersche Flora. Munich, 1789. 2 band. 8vo.
- Schranck Salish.* 1548. Id., Primitiæ Floræ Salisburgensis. Francof. Man. 1792. 8vo.
- Schranck Fl. Mon.* 767. Id., Plantæ rariores Horti Monacensis. Munich, 1817—1819. Fol.
- Schreb. Dec.* 744. Joh. Christ. Dan. Von Schreber, Icones Plantarum minus cognitarum, Decas I. Halle, 1766. Fol.
- Schreb. Gen.* 1126. Id., Linnæi Genera Plantarum. Francof. 1786. 2 vols. 8vo.
- Schrift. der Gesells. Naturf. Freunde.* 1765. See *Act. Nat. Scrut. Berol.*
- Schubert's Catalogue.* 1199. Spis Roslin ogrodu Botanicznego Krolewskiego-Warszawskiego Oni-warsyjetu. Catalogue des Plantes du Jardin Botanique de l'Université Royale de Varsovie. Publié par M. Schubert. Warsaw, 1824. 8vo.
- Schutt. Estr. Fl.* 1405. F. A. Schultes, Estrichs Flora. Wien, 1794. 2 band. 8vo.
- Schultz Fl. Starg. Suppl.* 1504. Schultz (C. F.), Floræ Stargardiensis Supplementum Primum. Naubrand, 1819. 8vo.
- Scop. Carn.* 342. Joh. Ant. Scopoli, Flora Carniolica. Viennæ, 1760, 1 vol. 8vo; ed. 2. Viennæ, 1772, 2 vols. 8vo.
- Scop. Del. Flor. Insubr.* 1414. Id., Deliciæ Floræ Insubricæ. Ticini, 1786—1788. 3 vols. fol.
- Scotia Illustrata.* 86. See *Sibbald's Scotia Illustrata.*
- Scots Gardener.* 87. The Scots Gardener; whereunto is annexed the Gardener's Kalendar. By J. Reid. Edin. 1683. 4to.
- Scott's Marmion.* 1763. A poem by Sir Walter Scott.
- Scottish Cryptogamic Flora.* 302. By Dr. Greville. Edin. 1822. 8vo.
- Seasons.* 962. Thompson's Seasons.

- Seba Thesaur.* 1761. Alb. Seba, Locupletissimi Rerum Naturalium Thesauri Descriptio. Amst. 1734—1765. 4 vols. fol.
- Secondat Mém. du Chêne.* 1812. Mémoires sur l'Histoire Naturelle du Chêne. Par M. De Secondat. Paris, 1785. Fol.
- Segu. Fer.* 1231. Joh. Franc. Seguiet, Plante Veronenses. Verona, 1745. 3 vols. 8vo.
- Seligm. Ar. Ic.* 1301. Trew (C. J.), Hortus, per J. M. Seligmann. Nürnberg, 1768—1786. 3 tom. folio.
- Selkirkshire Rep.* 1226. General View of the Agriculture in the Counties of Roxburgh and Selkirk. By R. Douglas. Edinb. 1798. 8vo.
- Sentiment of Flowers.* 1252. London, 1837.
- Sentimental Journey.* 1643. Sentimental Journey through France and Italy. By the Rev. L. Sterne. London, 1768. 2 vols. 12mo.
- Sepp's Icones Lignorum.* 1952. J. C. Sepp. A Representation of Inland and Foreign Wood, as well Trees as Shrubs; set forth with the Dutch, German, English, French, and Latin Names together. Amsterd., 1773—1791. 2 vols. 4to.
- Ser. Mol.* 768. N. C. Seringe Mélanges Botaniques. Berne, 1818. 2 vols. 8vo.
- Ser. Mus. Helv.* 1491. Id., Musée Helvétique d'Histoire Naturelle. Partie Botanique. Berne, 1818 and 1820. 4 vols. 4to.
- Series of Facts, &c.* 1800. A Series of Facts, Hints, Observations, and Experiments on the different Modes of raising, pruning, and training young Trees in Plantations. By W. Hillington. Shrewsbury, 1830. 8vo.
- Sering. Sal. exsicc.* 1498. See *Sering. Saules de la Suisse*.
- Sering. Saules de la Suisse.* 1541. N. C. Seringe, Saules de la Suisse. Berne. 5 fasc. 4to.
- Seringe Monogr. des Saules de la Suisse.* 1574. Id., Essai d'unc Monographie des Saules de la Suisse. Berne, 1815. 8vo.
- Shakspeare's Hamlet.* 485.
- Shaw, I. c., Icon.*, 2462. See *Shaw's Afr.*
- Shaw's Afr.* 526. Shaw (T.), Travels or Observations relating to several Parts of Barbary and the Levant. Oxford, 1738, folio; and London, 1757, 4to.
- Shaw's Staffordshire.* 1492. The History and Antiquities of Staffordshire, &c. By the Rev. Stebbing Shaw. London, 1798—1801. Folio.
- Shepherd's Week.* 2022. A poem by Gray.
- Shottel and Durfey's Archery Reviv'd.* 2084. The Bowman's Excellence, or, Archery Reviv'd; a poem by R. Shottel and T. Durfey. London, 1676. 8vo.
- Sibbald's Memoria Balfouriana.* 86. By Sir Robert Sibbald, M.D. Edinburgh, 1699. 8vo.
- Sibbald's Scotia Illustrata.* 86. Scotia Illustrata, sive Prodomus Historiæ Naturalis in quo Regionis Naturæ, &c. By Dr. R. Sibbald. Edin. 1684. Fol.
- Sibthorp's Fl. Græc.* 16. See *Flor. Græc.*
- Sieb. Voy. de Crète.* 320. Sieber (F. G.), Reise nach der Insel Kreta. Leipzig, 1823. 8vo.
- Silbermann's Revue Entomologique.* 2145. Silbermann, Revue Entomologique. Strasbourg, 1833. 8vo, in parts.
- Silliman's Journal.* 552. The American Journal of Science and Art. Conducted by Benjamin Silliman, M.D. In quarterly Nos. 8vo.
- Sims et Kön. Ann. of Bot.* 1504. See *Ann. of Bot.*
- Simond's Travels through Italy in 1817.* 2471. Simond (L.), A Tour in Italy and Sicily. London, 1828. 8vo.
- Singularities, &c.* 2516. See *Observations des Singularités, &c.*
- Siris; or, a Chain of Philosophical Reflections and Enquiries concerning the Virtues of Tar Water.* 2126. By Dr. G. Berkeley, Lord Bishop of Cloyne. Dublin, 1744. 8vo.
- Sketches.* 2188. See *Sketches in Spain*.
- Sketches, &c.* 44. Historical and Biographical Sketches of the Progress of Botany in England, from its Origin to the Introduction of the Linnæan System. By R. Pulteney, M.D. London, 1796. 2 vols. 8vo.
- Sketches in Spain.* 170. See *Sketches in Spain, &c.*
- Sketches in Spain, &c.* 1353. Sketches in Spain. By Capt. S. E. Cook, R.N. London, 1834. 2 vols. 8vo.
- Sketches of the Mar. Col. of British America.* 2139. See *McGrigor's Sketches of the Maritime Colonies, &c.*
- Stoane Hist.* 480. See *Stoane Jan.*
- Stoane Jan.* 736. The Natural History of Jamaica. London, 1707—1725. 2 vols. fol.
- Smith's Agricultural Report of Argylshire.* 92. General View of the Agriculture of the County of Argyll. By J. Smith. Edinb. 1798. 8vo.
- Smith's Eng. Bot.* 342. See *Eng. Bot.*
- Smith's Eng. Fl.* 255. See *Eng. Fl.*
- Smith's Exot. Bot.* 377. Exotic Botany. By Sir J. E. Smith. London, 1804, 1805. 2 vols. 4to, pl. 120.
- Smith's Flor. Brit.* 366. See *Fl. Br.*
- Smith's Flor. Gr.* 243. See *Flor. Græc.*
- Smith's History of Nova Scotia.* 1125. Published in the Mag. Nat. Hist.
- Smith's Icon.* 1093. See *Smith Icon. Pict.*
- Smith Icon. Pict.* 995. Icones pictæ Plantarum rariorum. By Sir J. E. Smith. London, 1790—1793. Fol.
- Smith's Introd. to Bot.* 1818. An Introduction to Physiological and Systematical Botany. Id. London, 1814. Pl. 15.
- Smith's Prodomus of Sibthorp's Flora Græca.* 164. Floræ Græcæ Prodomus: sive Plantarum omnium Enumeratio quas in Provinciis aut Insulis Græciæ invenit Johannes Sibthorp, M.D.; cum Annotationibus. Id. London, 1806—1813. 2 vols. 8vo.
- Smith's (Sir J. E.) Translation of Linnæus's Lachesis Lapponica.* 1105. See *Tour in Lapland*.
- Smith's Spic.* 314. Spicilegium Botanicum. Id. Lat. et Engl. Fascic. duo. London, 1792. Fol. pl. 24.
- Smith's Tour on the Continent.* 145. A Sketch of a Tour on the Continent in the Years 1786, 1787. Id. London, 1791. 3 vols. 8vo.
- Solander MS. in Herb. Banks.* 1109. Solander's MS. in the Banksian Herbarium.
- Sopwith's Fountains Abbey.* 2073. Eight Views of Fountains Abbey, intended to illustrate the Architecture and Picturesque Scenery of that celebrated Ruin; with an Historical and Architectural Description. By Thomas Sopwith. Folio.
- Southey's Roderick.* 1223. Roderick, the Last of the Goths. By Southey.
- Sow. Eng. Bot.* 1832. See *Eng. Bot.*
- Spach Ann. des Scien. Nat.* 987. Spach in the Annales des Sciences Naturelles.

- Spectacle de la Nature*. 2051. *Spectacle de la Nature, ou Entretiens sur les Particularites de l'Histoire Naturelle*. Par Noel Pluche. Paris, 1764—1770. 8 vols. 12mo. A Translation, under the title of *Spectacle de la Nature*; or, *Nature Displayed*; being discourses on such Particulars of Natural History as were thought most proper to excite the Curiosity and form the Minds of Youth. Illustrated with Copperplates. By Humphreys. London, 1740. 7 vols. 8vo.
- Spenser's Shepherd's Calendar*. 1785. A poem by Spenser.
- Spreng. Antiq. Bot.* 1917. Kurt Sprengel, Antiquitatum Botanicarum Specimen. Lipsiæ, 1798. 4to.
- Spreng. Gen.* 1261. Sprengel's (C.) edition of Linnæus's *Genera Plantarum*. Gött. 1830. 2 vols. 8vo.
- Spreng. Mag.* 997. Sprengel (K.), *Neue Entdeckungen in ganzen Umfang der Pflanzenkunde*. Leip. 1820. 8vo.
- Spreng. Spec. Bot. Ant.* 1721. Sprengel (K.), *Antiquitatum Botanicarum Specimen*. Leip. 1798. 4to.
- Spreng. Syst. Append.* 325. See *Spreng. Syst.*
- Sprengel's Syst.* 249. *Systema Vegetabilium, Caroli Linnæi*. Cur. Curtio Sprengel. Göttingæ, 1827. 4 vols. 8vo.
- Stackh. Comm. de Theophrast.* 1643. See *Theophrastus's Historia Plantarum*.
- Standish's Commons' Complaint.* 836. *The Commons' Complaint, concerning the Waste of Woods, and the Dearth of Victuals*; with Four Remedies for the same. London, 1611. 4to.
- Stat. Account. Scot.* 1226. Sinclair (Sir G.), *Statistical Account of Scotland*. Drawn up from the Communication of the Ministers of the different Parishes. Edinb. 1791—1799. 21 vols. 8vo.
- Staunt. Embass.* 2460. An authentic Account of an Embassy from the King of Great Britain to the Emperor of China. By Sir G. Staunton, Bart. Lond. 1797. 2 vols. 4to, pl. 44.
- Steel's History of Peat Moss.* 1464. Steel (A.), *The Natural and Agricultural History of Peat Moss, or Turf Bog*. Edinb. 1826. 8vo.
- Stephenson and Churchill's Medical Botany.* 465. See *Churchill's Medical Botany*.
- Stephens Illust. Haust.* 1820. See *Stephens (J. F.), &c.*; the work being divided into two sections, *Mandibulata* and *Haustellata*.
- Stephens (J. F.), Illustrations of British Entomology.* London, 1828. 10 vols. and some parts, and continued. 8vo.
- Steucl. Nom. Bot.* 889. See *Steucl's Nomenclator*.
- Steucl's Nomenclator.* 1486. E. Steucl, *Nomenclator Botanicus, enumerans Ordine alphabetico Nomina atque Synonyma Plantis imposita*. Stuttgart. et Tubing. 1821—1824. 2 vols. 8vo.
- Stev. Obs. incd. in H. & Willd.* 745.
- St. Helena Gazette.* 1511. A periodical.
- St. Hilaire Livr.* 1444. Saint Hilaire (Auguste), *Histoire des Plantes les plus remarquables du Brésil et de Paraguay*. Paris, 1824. 3 vols. 4to.
- Stirpium Historiæ Pemptadcs.* 234. See *Dod. Pempt.*
- St. James's Chronicle.* 1507. A London newspaper.
- Stoke's Bot.* 1123. A Botanical Arrangement of British Plants, &c., by W. Withering, including a new set of references to Figures. By Jonathan Stokes, M.D. Birmingham, 1787. 2 vols. 8vo.
- Story of Rimini.* 2122. A poem by Leigh Hunt.
- St. Pierre's Etudes de la Nature.* 1307. J. Bernardin Henri De Saint Pierre, *E'tudes de la Nature*. Paris, 1800. 10 vols. 18mo.
- Strutt's Sylva.* 1221. See *Strutt's Sylva Britannica*.
- Strutt's Sylva Britannica.* 48. *Sylva Britannica*; or, *Portraits of Forest Trees distinguished for their Antiquity, &c.* By J. G. Strutt. Imperial 8vo, 50 plates.
- Strype's Life of Grindal.* 35. *Life and Acts of Bishop Grindal*; with an Appendix. By John Strype, M.A. London, 1710. Folio.
- Studies of Nature.* 1006. St. Pierre's *E'tudes de la Nature*, translated by Dr. H. Hunter. London, 1799. 3 vols. 8vo.
- Sturm D. Fl.* 1494. Jacob Sturm, *Deutschland Flora*. Pars. 1. (Phanog.), fasc. 1—28. Pars 2. (Cryptog.), fasc. 1—11. Nürnberg, 1798 et seq. 2 vols. 4to.
- Suburban Gardener.* 693. *The Suburban Gardener, and Villa Companion*: comprising the Choice of a Suburban or Villa Residence, or of a Situation on which to form one. By J. C. Loudon, F.L.S., H.S., &c. London, 1833. 8vo.
- Suppl. Encyc. Agri.* 738. Supplement to the *Encyclopædia of Agriculture*. By J. C. Loudon, F.L.S., &c. London, 1834. 8vo.
- Suppl. to Eng. Bot.* 735. See *Supplement to English Botany*.
- Supplement to English Botany.* 1433. See *Eng. Bot. Suppl.*
- Sur la Naturalisation des Arbres Forestiers de l'Amérique.* 142. *Mémoire sur la Naturalisation, &c.* Par F. A. Michaux. Paris, 1805. 8vo.
- Surrey.* 28. See *Munning and Bray's Surrey*.
- Survey of Dorsetshire.* 36. See *Coker's Survey of Dorsetshire*.
- Survey of Forfarshire.* 100. See *Headrick's Survey of Forfarshire*.
- Survey of London.* 512. *A Survey of London, containing the original Antiquities, Increase, moderne Estate, and Description of that City, written in the Year 1598*. By John Stowe, Citizen of London. London, 1598. Small 4to.
- Survey of Worcestershire.* 896. See *Pitt's Survey of Worcestershire*.
- Survey of the County of Nottingham.* 1520. *General View of the Agriculture of the County of Nottingham*. By R. Lowe. London, 1798. 8vo.
- Sut. Helv.* 1586. See *Suter's Flora Helvetica*.
- Suter's Flora Helvetica.* 160. J. R. Suter. *Flora Helvetica*. Turici, 1802. 2 vols. 12mo.
- Sutherland's Catalogue of Plants in the Edinburgh Botanic Garden.* 53.
- Sutherland's Hortus Medicus Edinburgensis.* 104. Jas. Sutherland, *Hortus Medicus Edinburgensis*. Edinb. 1683. 8vo.
- Svensk Bot.* 1043. *Svensk Botanik*. Stockholm, 1804—1819. 8 vols. 8vo.
- Swartz Ft. Ind. Occ.* 999. Olof von Olaus Swartz, *Flora Indiæ Occidentalis*. Erlanga, 1787—1806. 3 vols. 8vo.
- Swartz Prod.* 2271. Id., *Prodromus Descriptionum Vegetab. Indiæ Occidentalis*. Holmiæ, 1788. 8vo.
- Sw. Br. Fl.-Gard.* 242. *The British Flower-Garden*. Conducted by R. Sweet and afterwards by Professor Don, till 1837. London. 8vo.
- Sweet's H.B.* 388. See *Sweet's Hort. Brit.*
- Sweet's Hort. Brit.* 240. Sweet (R.), *Hortus Britannicus*. London, 1826. 8vo.
- Swt. Cist.* 518. *Cistaceæ*. *The Natural Order of Rock Rose*. By R. Sweet. London, 1830. 8vo.
- Swt. Fl. Austr.* 359. *Flora Australasica*. London, 1827, 1828. 8vo.

- Swz. Obs.* 450. Olof seu Olaus Swartz, *Observationes Botanicae*. Erlange, 1791. 8vo.
- Syl. Amer.* 263. See *Sylva Americana*.
- Syl. Sketches*. 1029. Sylvan Sketches; or, a Companion to the Park and the Shrubbery, &c. By Miss Kent, the Author of *Flora Domestica*. London, 1825. 8vo.
- Syllog. Nov. Pl. Soc. Bot. Nat.; Ratisb. edita*. 1487. Sylloge Plantarum Novarum a Societate Regia Ratisbonensi edita. Ratisbon, 1821, and continued.
- Sylva*. 44. Sylva; or, a Discourse on Forest Trees. By John Evelyn. London, 1664. 4to.
- Sylva Americana*. 190. By D. J. Brown. Boston, 1832. 8vo.
- Sylva Britannica*. 48. See *Strutt's Sylva Britannica*.
- Sylva Florifera*. 188. Sylva Florifera; the Shrubbery, containing an Historical and botanical account of the Flowering Shrubs and Trees which now ornament the Shrubbery, &c. By H. Philips, F.R.S. London, 1823. 2 vols. 8vo.
- Synopsis of the British Flora*. 734. See *Lindl. Synops.*
- Synopsis Methodica Stirpium Britannicarum*. 53. Joh. Ray, *Synopsis Methodica Stirpium Britannicarum*. Londini, ed. 2., 1696; ed. 3., 1724. 8vo.
- Syst. Eleuth.* 1150. J. C. Fabricius, *Systema Eleutheratorum secundum Ordines, &c.* Kiliae, 2 vols. 8vo.
- Syst. Veg. Querc.* 1925. See *Sprengel's Syst. Veg.*
- T.
- Tabern. Krauterb.* 1731. J. T. Tabernaemontanus, *Kräuterbuch*. Francof. Mein, 1588—1592. Ed. 2., Cur. C. Bauhino. Francof. Mein, 1613 und 1625. Basilia, 1664 and 1731.
- Tableau Historique*. 1233. Tableau Historique des Progrès de la Culture des Arbres à Oïessa. Par M. Descemet.
- Task*. 1308. Cowper's Task. A poem.
- Tenore*. 430. Tenore (M), Sylloge Plantarum. Naples. 8vo.
- Ten. Syll.* 1205. See *Tenore*.
- Tenore. Prod. Fl. Neap.* 337. Mich. Tenore, *Prodromus Floræ Neapolitanæ*. Neap. 1811—1813. 8vo.
- Tenore. Syn. Fl. Neap.* 352. Id., *Synopsis novarum Plantarum, quæ in Prodrómo describuntur*. Neap. 1815. 8vo.
- Tenore Syll. Fl. Neap.* 2494. See *Tenore*.
- Tenore's Flora Neapolitana*. 164. Id., *Flora Neapolitana*. Neapoli, 1811 et seq. 6 fasc. fol.
- Tentamen Floræ Nepalensis illustratæ*. 2100. Dr. Wallich, *Tentamen, &c.* Calcutt. et Scraip. 1824. Fol.
- Thalaba*. 2040. A poem by Southey.
- The Analyst*. 2082. *The Analyst*; a monthly (quarterly) Journal of Science, Literature, and the Fine Arts. London, 1834, and continued.
- The Botanist's and Gardener's New Dictionary*. 80. *The Botanist's and Gardener's New Dictionary*; containing the Names, Classes, Orders, Generic Characters, &c. By Jas. Wheeler, Nurseryman, Gloucester. London. 8vo.
- The Colonies*. 2327. *The Colonies: Treating of their Value generally; and of the Ionian Islands in particular*. By Col. C. J. Napier, C.B. London, 1833. 8vo.
- The Garden*. 1991. A poem by Ropin.
- The Language of Flowers*. 792. See *Language of Flowers*.
- The Laidley Worm of Spindleston Heughs*. 919. An ancient Song.
- The Manner of raising Forest Trees, &c.* 46. By Moses Cooke. London, 1676. 4to.
- The Pioneers*. 2138. A novel by Cooper.
- The School of Shooting*. 2086. See *Toxophilus*.
- The State of Eton School*. 839. A curious MS. 1560.
- The True Method of Improving an Estate by Plantations of Timber Trees*. 80. By Batty Langley. London, 1728. 4to.
- The Winter's Tale*. 486. One of Shakspeare's Plays.
- Theat. Bot.* 1902. See *Parkinson's Theat. Bot.*
- Théat. d'Agric.* 1339. *Olivier de Serres. Théâtre d'Agriculture*. Paris, 1600, fol.; 1611, 8vo. Genève, 1639; Lyon, 1675; Paris, 1801, 1805. 2 vols. 4to.
- Théâtre d'Éducation*. 792. By Madame De Genlis. Paris, 6 tom. 12mo.
- Théologie des Insectes*. 1816. F. C. Lesser, *Théologie des Insectes, ou Démonstration des Perfections de Dieu dans tout ce qui concerne les Insectes*. Trad. de l'Allemand: avec des Remarques de M. P. Lyonnet. A la Haye, 1742. 2 vols. 8vo, pl. 2.
- Theophrastus's Historia Plantarum*. 16. Theophrasti *Historia Plantarum*, Gr. et Lat. Amsterdam 1644. Fol. Translated by J. Staekhouse. Oxford, 1811. 4to.
- Théorie Élémentaire*. 215. A. P. De Candolle, *Théorie élémentaire de la Botanique*. Paris, 1813. 8vo.
- Thib. incd.* 37. Thibaud's MS.
- Thickness's For. Veg.* 1834. A Treatise on Foreign Vegetables, &c. By Ralph Thicknesse, M.D. London, 1749. 8vo.
- Thomp. Disp.* 677. Thompson's (A. T.) London Dispensatory.
- Thore (Chor. Land)*. 1482. J. Thore, *Essai d'une Chloris du Département des Landes*. Dax, 1803. 8vo.
- Thore Prom. sur les Côtes de Gascogne*. 2214. Id. Dax, 1804. 8vo.
- Thouin Ann. Mus.* 951. Thouin in *Annales du Musée*.
- Thouin Dict. Agric.* 258. Thouin in *Dict. Agric.*
- Three Years in North America*. 1919. By J. Stuart. Edinburgh, 1833. 12mo.
- Thoul. Fl. Par.* 415. J. L. Thuillier, *Flora des Environs de Paris*. Paris, 1790. 12mo.
- Thoul. Paris*. 1498. See *Thoul. Fl. Par.*
- Thun. Prod.* 504. C. P. Thunberg, *Prodromus Plantarum Capensium*. Upsal, 1794. 2 parts. 8vo.
- Thunb. Diss.* 1683. Id., *Diss. de Ericâ*. Upsal, 1785. 4to. Ed. 2. Cur. R. A. Salisbury. Featherstone, 1800.
- Thunb. Fl. Jap.* 235. See *Thunb. Jap.*
- Thunb. Jap.* 504. Id., *Flora Japonica*. Lipsie, 1784. 8vo.
- Thunb. Lan. Soc. Trans.* 235. Id., in the Transactions of the Linnæan Society.
- Tilli Cat. Hort. Pisani*. 1311. M. A. Tilli, *Catalogus Horti Pisani*. Florentia, 1723. Fol.
- Tillock's Phil. Mag.* 1388. See *Phil. Mag.*
- Tin. Pug.* 556. See *Tin. Pug. Sc.*

- Tin. Pug. Sic.* 557. Vinc. Tineo, Pugillus Plantarum Sicillæ rariorum. Panormi, 1817. 8vo.
- T. Nees ab Esenbeck Gen. Pl. Floræ Germanicæ.* 1816. See *Nees Von Esenbeck Gen. Pl. Toxic Fung. Meck.* 2148. H. J. Tode, Fungi Meklenburgensium selecti. Lincburgi, 1790. 2 fasc. 4to.
- Torrey Fl. U. S.* 555. Torrey's Flora of the Northern and Middle States. New York, 1826.
- Tour in Ireland.* 2080. See *Irish Tour.*
- Tour in Lapland.* 1105. Lachesis Laponica; or, a Tour in Lapland, now first published from the original MS. Journal of the celebrated Linnæus. By Sir J. E. Smith. London, 1811. 2 vols. 8vo.
- Tour in Scotland.* 901. By Pennant. Chester, 1771. 8vo.
- Tour on the Continent.* 145. See *Smith's Tour on the Continent.*
- Tourn. Cor.* 602. J. Pitton de Tournefort, Corollarium Institutionum Rei Herbariæ. Paris, 1703. 4to.
- Tourn. Inst.* 328. Id., Institutiones Rei Herbariæ. Paris. 1717 et 1719. 3 vols. 4to.
- Tourn. Itin.* 1310. Id., Relation d'un Voyage au Levant. Paris, 1717. 2 vols. 4to.
- Toxophilus.* 2086. Toxophilus; or Schole of Shotyng, conteyned in Two Books. By Roger Ascham. London, 1544. 4to.
- Tracts.* 57. Tracts on Practical Agriculture and Gardening, in which the Advantage of imitating the Garden Culture in the Field, &c. By Richard Weston. London, 1769. 4to.
- Tradescant's Catalogue.* 40. Published in Museum Tradescantianum. See *Museum Tradescantianum.*
- Trag.* 1198. Hieronimi Tragi, De Stirpium, maxime earum quæ in Germanica nostra nascuntur, &c. Argentorati, 1551. 4to.
- Traité de l'Aménagement des Bois et Forêts, &c.; suivi de Recherches sur les Chênes à Glands doux.* 1735. Par M. Dralet. Paris, 1812. Small 8vo.
- Traité de la Châtaigne.* 1996. Par A. A. Parmentier. Paris, 1770. 8vo.
- Traité de la Culture des Terres.* 137. H. L. Du Hamel du Monceau, Traité de la Culture des Terres. Paris, 1750—1761. 6 vols. 12mo. Vers. Germ., Dresden, 1752; Vers. Belg., Amsterdam, 1762.
- Traité Prat., &c.* 2155. See *Traité Pratique, &c.*
- Traité Pratique de la Culture des Pins.* 2121. See *Delamarre's Traité, &c.*
- Traité sur les Mûriers blancs.* 1350. Par C. Castellet. Aix, 1760. 12mo.
- Trans. Roy. Soc.* 2116. See *Phil. Trans.*
- Trans. Soc. Arts.* 1969. See *Transactions of the Society of Arts.*
- Transactions of the Entomological Society of London.* 906. London. Begun in 1834, in 8vo.
- Transactions of the Highland Society.* 100. The Quarterly Journal of Agriculture; and the Prize Essays and Transactions of the Highland Society of Scotland. Edin. 1828. In quarterly Nos. 8vo.
- Transactions of the Horticultural Society of London.* 172. London, 1st series, 1812—1830, 7 vols. 4to; 2d series, begun in 1831, and continued occasionally.
- Transactions of the Prussian Horticultural Society.* 149. Verhandlungen des Vereins zur Beförderung des Gartenbaues in den Königlich Preussischen Staaten. Berlin. 4to.
- Transactions of the Royal Irish Academy.* 111. Dublin. 4to. Published occasionally.
- Transactions of the Society for the Encouragement of Arts.* 1475. Transactions of the Society for the Encouragement of Arts, Manufactures, Commerce, &c. Lond. 1783. Published occasionally.
- Transactions of the Society of Arts.* 1550. See *Transactions of the Society for the Encouragement of Arts.*
- Transactions of the Society of Edinburgh.* 789. Edin. 1788. 4to. Published occasionally.
- Tratt. Alb. Tosc.* 2094. Trattinick in Savi (C.) Trattato degli Alberi della Toscana. Florence, 1801. 2 vols. 8vo.
- Tratt. Arch.* 406. Leop. Trattinick, Archiv. der Gewächskunde. Vienn. 1811—1812. 3 fasc. 4to.
- Tratt. Ros.* 742. Id., Rosacearum Monographia. Vindob. 1823, 1824. 4 vols. small 8vo.
- Travelling Memorandums.* 1662. Travelling Memorandums made in a Tour on the Continent of Europe in the Years 1786 and 1788. By Francis Garden, Lord Gardenstone. Edin. 1796. 3 vols. 12mo.
- Travels.* 556. Travels into North America; containing its Natural History, with the Civil, Ecclesiastical, and Commercial State of the Country. By P. Kalin. In Swedish. Stockh. 1753, 1756, and 1761. 3 vols. 8vo. Translated by J. R. Forster. Warrington, 1770. 3 vols. 8vo.
- Travels.* 786. See *Porter's (Sir Robert Ker) Travels.*
- Travels.* 1345. Dr. D. E. Clarke's Travels in various Countries of Europe, Asia, and Africa. Lond. 1810. 4 vols. 4to.
- Travels.* 2167. Travels in Poland, Russia, Sweden, &c. By W. Core, F.R.S. London, 1784. 2 vols. 4to.
- Travels.* 2408. Travels through the Southern Provinces of the Russian Empire, in the Years 1792—1794. By P. S. Pallas. 2 vols. 4to.
- Travels among the Arab Tribes.* 2411. Travels among the Arab Tribes of Syria. By J. S. Buckingham. Lond. 1825. 4to.
- Travels in America in 1753.* 1151. See *Travels, 556.*
- Travels in Asia Minor, &c.* 2411. B. J. Macdonald Kinnier. London, 1818. 8vo.
- Travels in Egypt.* 1930. See *Olivier's Travels.*
- Travels in Georgia and Florida.* 1018. See *Eartram's Travels.*
- Travels in Greece.* 511. By Dr. Chandler. London, 1766. 4to.
- Travels in New England.* 2260. See *Dwight's Travels in New England.*
- Travels in North America.* 2194. See *Travels, 556.*
- Travels in Persia.* 786. By Sir William Ouseley. London, 1819.
- Travels in Scotland.* 2175. See *Hall's Travels in Scotland.*
- Travels in the Ionian Isles, &c.* 2227. Travels in the Ionian Islands, Albania, Thessaly, &c. By Dr. Holland, in 1813—1814. Lond. 1815. 4to.
- Travels in the Levant.* 2201. See *Olivier's Travels.*
- Travels in the Mysore.* 2430. Burnes (A.), Travels into Bokhara, &c. London, 1824. 3 vols. 8vo.
- Travels in the Peruvian Andes.* 2434. See *Pæppig's Travels, &c.*
- Travels in the Tarentaisc.* 1430. Travels in the Tarentaisc, &c. By R. Bakewell, Esq. London, 1824. 8vo.
- Travels through Italy in 1817.* 2471. See *Simon's Travels, &c.*
- Treat., &c.* 1304. See *Boucher's Treatise on raising Forest Trees.*
- Treatise on Ancient Arms and Armour,* 2087. See *Grosz's, &c.*

- Treatise on Civil Arch.* 1441. *Treatise on Civil Architecture.* By Sir William Chambers. London, 2d ed., 1768. Folio.
Treatise on Country Residences. 2385. A Treatise on the forming, improving, and managing Country Residences, &c. By J. C. Loudon, F.L.S., &c. Lond. 1805. 2 vols. 4to.
Treatise on Forest Trees. 162. Earl Haddington. 1733.
Treatise on the Acacia. 1616. See *Withers's Treatise*.
Treatise on the Growth, Qualities, and Uses of the Acacia Tree. 617. See *Withers's Treatise on the Acacia*.
Treatise on the Mulberry, &c. 1858. See *Pascal's Treatise on the Mulberry*.
Tredgold's Carpentry. 1219. *Elementary Principles of Carpentry.* By Thomas Tredgold. London 4to.
Trev. Obs. Bot. 1497. See *Treviranus's Obs. Bot.*
Treviranus's Obs. Bot. 1454. *Beiträge zur Pflanzen Physiologie.* Göttingen, 1811. 8vo.
Trew Ehr. 841. See *Trew Sel.*
Trew Sel. 424. C. J. Trew, *Plantæ selectæ ab Ehret. pictæ.* 1750—1753. Fol.
Trüffelbau. 1975. See *Der Trüffelbau.*
Turner's Herbar. 25. See *Turner's Names of Herbes.*
Turner's Names of Herbes. 35. A New Herball; with the Names of Herbes in Greek, Latin, Dutch, French, &c.; with numerous wood-engravings. In 2 parts. Pt. 1., London, 1551, fol.; pt. 2., Collen, 1562, fol.
Turp. et Pôt. Fl. de Par. 1212. Poiteau et Turpin, *Flore Parisienne.* Paris, 1808, et suiv. Fol.

U.

- Ungarisch Mag.* 2186. *Ungrisches Magazin von H. C. Von Windisch.* Presburg, 1781—1783. 3 vols, 8vo.
Ups. 1229. C. Linnæus, *Hortus Upsaliensis.* Stockholm, 1748. 8vo.

V.

- Vahl Enum.* 1203. Martinus Vahl, *Enumeratio Plantarum.* Hauniae, 1805, 1806. 2 vols. 8vo.
Vahl Fl. Dan. 1594. See *Fl. Dan.*
Vahl Synb. Id., *Symbolæ Botanicae.* Hauniae, 1790—1794. 3 fasc. fol.
Vail. Herb. 601. Vaillant (S.), *Botanicon Parisiense.* Lug. Bat. 1723. 8vo.
Vail. Acad. Scien. 1203. Vaillant in *Acad. des Scien.*, &c.
Various Thoughts on Planting and Training Trees. 188. By Dr. Anderson, under the Name of Agricola. Edin. 1771. 8vo.
Vegetable Statistics. 1189. By S. Hales. London, 1727, 8vo; ed. 2., 1733; ed. 3., in *Statistical Essays*, 1738.
Veiled Prophet of Khorassan. 2040. A poem by Moore.
Vent. Cels. 259. E. P. Ventenat, *Description des Plantes nouvelles ou peu connues du Jardin de J. M. Cels.* Paris, 1800, fol.; ed. Germ., Remer, Zurich, 1802.
Vent. Choix. 330. Id., *Choix des Plantes.* Paris, 1803—1808. 10 fasc. fol.
Vent. Hort. Cels. 338. See *Vent. Cels.*
Vent. Ill. 662. See *Vent. Matm.*
Vent. Jard. Matm. 1266. See *Vent. Matm.*
Vent. Matm. 642. Id., *Jardin de la Malmaison*, 1803—1805. 2 vols. fol.
Vent. Tabl. du Règne Végét. 1562. Id., *Tableau du Règne Végétal.* Paris, 1797. 4 vols. 8vo.
Ventenat Dec. Nov. Gen. 2462. Id., *Decas Generum novorum, &c.* 1808. 4to.
Verhand. Batav. Genootsch. 2329. *Verhandelingen van het Bataviaasch Genootschap der Koustan an wetenschappen.* Batavia aut Rotterdam, 1770—1792. 6 vols. 8vo.
Versuch über die Metamorphose der Pflanzen. 225. See *Gütthe's Ditto*.
Vetusta Monumenta. 1748. *Vetusta Monumenta quæ ad Rerum Britannicarum Memoriam conservandam Societas Antiquariorum Londini sumptu suo edenda curavit.* London, 1747—1796. 3 tom. folio.
Vill. Cat. Strasb. 889. D. Villars, *Catalogue méthodique du Jardin de Strasbourg.* Strasbourg, 1807. 8vo.
Vill. Dauph. 421. See *Villars's Plantes du Dauphiné.*
Vill. Prosp. 587. Villars (M.), *Prospectus de l'Histoire des Plantes du Dauphiné.* Grenoble, 1779.
Villars's Plantes du Dauphiné. 2276. Id., *Histoire des Plantes du Dauphiné.* Grenoble, 1786—1788. 4 vols. 8vo.
Virgil En., Dryden's Trans. 1784. *Dryden's Translation of Virgil,* Carey's ed. Lond. 1806. 3 vols. 8vo.
Visiani Pl. Dalm., ex Bot. Zeit. 587. *Plantæ Dalmaticæ nunc primum editæ a R. De Visiani, M.D., in the Botanische Zeitung.* 1830.
Vitman. Sp. Pl. 2318. Vitmann, *Summa Plantarum quæ hactenus, &c.* Mediol. 1789—1792. 6 vols. 8vo.
Viv. Ann. Bot. 583. Dom. Viviani, *Annali di Botanica.* Genova, 1802—1804. 2 vols. 8vo.
Viv. Fragm. 543. Id., *Floræ Italicæ Fragmenta.* Genæ, 1808. 4to.
Volkamer's Continuation der Nurembergischen Hesperidum. 1337. Nürnberg, 1708. Folio.
Voy. from Egypt. 526. See *Delil. Descr. Egypt.*
Voyage à la Louisiane. 2484. Dupratz.
Voyage aux Iles Canaries. 2263. See *Feuillec H.*
Voyage dans les Petits Cantons, et dans les Alpes Rhétiques. 2276. See *Kasthofer's Voyage dans, &c.*
Voyage de Crète. 638. Siebuhr.
Voyage de Syrie et du Mont Liban. 2409. Par Jean De la Roque. Paris, 1722. 2 vols. 12mo.
Voyage to Crète. 320. See *Voyage de Crète.*
Voyage to the Levant. 2410. *Voyage to the Levant, &c.* By Cornelius Bruyn. London, 1702. fol.

W.

- Wen.* 1074. See *Willd. Enum.*
W. Sp. 238. See *Willd. Sp. Pl.*
Wade's Salices. 1498. *Salices.* By Dr. Wade. Dublin, 1811. 8vo.
Wahl. Fl. Succ. 1138. G. Wahlenberg, *Flora Suecica.* Upsal, 1824—1826. 2 vols. 8vo.

- Wahl. Helv.* 339. Id., De Vegetatione et Climate in Helvetia Septentrionali, intra Rhenum et Arolam. Turici, 1813. 8vo.
- Wahlb. Fl. Carp.* 600. Id., Flora Carpatorum principalium, exhibens Plantas in Montibus Carpaticis, inter Flumina Waagum et Dunajetz, crescentes; cum Tractatu de Altitudine, &c., horum Montium. Götting. 1814. 8vo.
- Wahlb. Fl. Lapp.* 1514. Id., Flora Lapponica, in Itineribus 1800—1810 denuo investigata. Bero-
lini, 1812. 8vo.
- Waldst. et Kit. Pl. Hung.* 372. See *Waldst. et Kit. Pl. Rar. Hung.*
- Waldst. et Kit. Pl. Rar. Hung.* 552. Waldstein et Kitaibel, Descriptiones et Icones Plantarum rariorum Hungariæ. Vienna, 1802—1812. 3 vols. folio.
- Wall. Asiat. Res.* 390. Nath. Wallich, Asiatic Researches; or, Transactions of the Society instituted at Bengal. 1788. 4to.
- Wall. Cat.* 872. Wallich's MS. Catalogue of the Plants contained in the Herbarium of the East India Company, now in the Possession of the Linnean Society.
- Wall. Fl. Ind.* 500. Id., Tentamen Floræ Nepalensis illustrata. Calcutta, 1824. Folio.
- Wall. MSS.* 306. See *Wall. Cat.*
- Wall. Pl. As. Rar.* 245. See *Wall. Pl. Asiat. Rar.*
- Wall. Pl. Asiat. Rar.* 390. Id., Plantæ Asiaticæ rarioræ; or, Descriptions and Figures of a select Number of unpublished Plants of the East Indies. London, 1830—1832. 3 vols. royal fol. 300 plates.
- Wallr. Sched.* 739. F. W. Wallroth, Schedulæ critica de Plantis Floræ Halensis selectis, &c. Halæ, 1822. 8vo.
- Wallroth Monog.* 813. F. G. Wallroth, Rosæ Plantarum Generis Historia succincta, &c. Nord-
husæ, 1828. 8vo.
- Walpole's Memoirs.* 2203. Walpole's (R.) Memoirs relating to European and Asiatic Turkey, &c. London, 1818. 4to.
- Walsh's History of Dublin.* 117. Whitelaw and Walsh's History of the City of Dublin, &c., with numerous plates and maps. Lond., 1818. 2 vols. 4to.
- Wall. Car.* 655. See *Wall. Fl. Car.*
- Wall. Fl. Car.* 240. Thom. Walter, Flora Caroliniana. Londini, 1788. 8vo.
- Wall. Hort.* 1291. A. P. Walthier, Designatio Plantarum Horti ejus. Lipsiæ, 1755. 8vo.
- Wanderings.* 918. Wanderings in South America, &c., in the years 1812, 1816, 1820, and 1824. By Charles Waterton, Esq. London, 1825, 4to; 2d ed. 1829, 8vo; 3d ed. 1836, small 8vo.
- Wang. Am.* 820. See *Wangenh. Amer.*
- Wang. Beytr.* 691. Wangelheim (F. A. G.), Anpflanzung nord-amerikanischer Holzarten. Göt-
tinge, 1787. 3 vols. folio.
- Wang. Forst.* 1879. See *Wang. Beytr.*
- Wangenh. Amer.* 691. See *Wang. Beytr.*
- Wats. Dend.* 470. See *Watson's Dend. Brit.*
- Wats. Dend. Brit.* 1417. See *Watson's Dend. Brit.*
- Watson's Dend. Brit.* 1421. P. W. Watson, Dendrologia Britannica; or, Trees and Shrubs that will live in the open Air of Britain. London, 1825. 2 vols. 8vo, pl. 172.
- Watson's Outlines.* 3019. Outlines of the Geographical Distribution of British Plants. By H. C. Watson. London, 1832.
- Webb and Bertholet's Histoire Naturelle des Iles Canaries.* 1332. Histoire Naturelle des Isles Canaries. Par MM. P. Barker Webb et Sabin Bertholet. Paris, 1826—1828. 4to and fol., many plates.
- Weibe and Nees's Rubi Germanici.* 735. A. Weibe et C. G. Nees ab Esenbeck, Rubi Germanici. Bonn, 1822. Fol., pl. 53.
- Wendl. Eric.* 1081. J. C. Wendlend, Ericarum Icones et Descriptiones. 1798 et seq. 4to.
- Wendl. Hort.* Id., Hortus Herrenhusanus. 1798—1801. 4 fasc. fol.
- Wendl. Obs.* 298. Wendlend (J. C.), Botanische Beobachtungen nebst einigen neuen Gattungen und Arten. Hannover, 1798. Fol.
- Wern. Soc. Trans.* 1137. See *Wernerian Trans.*
- Wernerian Trans.* 1704. Memoirs of the Wernerian Natural History Society. Edinburgh, 1811, &c. 8vo.
- West of England Journ. of Science and Lit.* 1120. Bristol, 1855. 8vo.
- Western Times.* 1837. A provincial newspaper.
- Whecl. Itin.* 399. See *Wheeler's Journey.*
- Wheeler's Journey.* 400. A Journey into Greece in the Company of Dr. Spor of Lyons. By Sir G. Wheeler. London, 1682. Folio.
- Whit. Craven.* 1225. See *Whittaker's History of Craven.*
- Whittaker's History of Craven.* 1225. History and Antiquities of the Deanery of Craven, in the County of York. By the Rev. T. Whittaker. London, 1805. Folio.
- Wibel Wirtb.* 763. A. G. E. C. Wibel, Primitiæ Floræ Wertheimensis. Jenæ, 1799. 8vo.
- Wijón's Garcilasso.* 1654. Works of Garcilasso de Vega, translated into English verse, with an Essay on Spanish Poetry. By J. H. Wiffen. With a portrait and wood-cuts. London, 1823. 8vo.
- Wight's Husbandry of Scotland.* 92. The present State of the Husbandry in Scotland. By Andrew Wight. Edinburgh, 1778—1790. 6 vols. 8vo.
- Wikström Diss. de Daphnæ.* 1511. J. E. Wikström, Dissertatio Botanica de Daphne. Stockholm, 1820. 4to.
- Willd. Ab.* 254. See *Abbildung der Deutschen Holzarten.*
- Willd. Abbild.* 42. See *Abbildung der Deutschen Holzarten.*
- Willd. Arb.* 235. See *Willdenow's Baumzucht.*
- Willd. Baum.* 301. See *Willdenow's Baumzucht.*
- Willd. Berl. Baumz.* 1329. Id., Berlinische Baumzucht. Berl. 1811. 8vo.
- Willd. Enum.* 303. See *Willdenow's Enumeration, &c.*
- Willd. Enum. Suppl.* 351. Id., Enumeratio Plantarum Horti Berolinensis. Supplementum posthumum addidit Schlechtendal.
- Willd. Hort. Ber.* See *Willd. Enum.*
- Willd. Holzart.* 354. See *Abbildung der Deutschen Holzarten.*
- Willd. Sp.* 233. See *Willd. Sp. Pl.*
- Willd. Spec.* 423. See *Willd. Sp. Pl.*
- Willd. Sp. Pl.* Id., Linnæi Species Plantarum. Berol. 1797. 5 vols. 8vo.
- Willdenow's Baumzucht.* 146. Id., Berlinische Baumzucht. Berol. 1811. 8vo.
- Willdenow's Enumeration of the Plants of the Berlin Royal Garden.* 1416. Id., Enumeratio Plantarum Horti Berolinensis. Berol. 1809. 2 vols. 8vo.
- Wilson's American Ornithology.* 86. American Ornithology; or, the Natural History of the Birds of the United States. Philadelphia, 1808—1814. 9 vols. 4to, pl. 76.

- Winch Geogr. Distrib.* 769. An Essay on the Geographical Distribution of Plants through the Counties of Northumberland, Cumberland, and Durham. By N. J. Winch. Newcastle. Ed. 2. 1825. Pamph. 8vo.
- Withering's Botany.* 1223. A Systematical Arrangement of British Plants. By W. Withering, M.D. Birmingham, 1776, 2 vols. 8vo; ed. 7. with additions, London, 1830, 4 vols. 8vo.
- Withers's Treat.* 611. See *Withers's Treatise on the Acacia*.
- Withers's Treatise.* 615. See *Withers's Treatise on the Acacia*.
- Withers's Treatise on the Acacia.* 616. A Treatise on the Growth, Qualities, and Uses of the Acacia Tree, &c. By W. Withers, Esq., of Holt, Norfolk. 1836. 8vo. Not published.
- Wolff's Missionary Journal.* 2411. Wolff (J.), Missionary Journal. London, 1827—1829. 3 vols. 8vo.
- Wood's Bowman's Glory.* 2086. The Bowman's Glory, or Archery Revived; giving an Account of the many signal Favours vouchsafed to Archers and Archery by those renowned Monarchs King Henry VIII., James, and Charles I., as by their several gracious Commissions here recited may appear. With a brief Relation of the Manner of Archers marching on several Days of Solemnity. Published by William Wood, Martial to the Regiment of Archers. London, 1682. 12mo.
- Woodhouselee's Life of Lord Kaimcs.* 94. Memoirs of the Life and Writings of the Hon. Henry Home of Kames. By A. F. Tytler, Lord Woodhouselee. Edinb. 1807. 2 vols. 4to.
- Woodlands.* 188. The Woodlands. By William Cobbett. London, 1825. 8vo.
- Woodville's Med. Bot.* 485. Medical Botany; containing Systematic and General Descriptions, with Plates, of all the Medicinal Plants, indigenous and exotic, &c. By W. Woodville, M.D. London, 1790. 3 vols. 4to.
- Woodville's Med. Bot. Suppl.* 393. See *Woodville's Med. Bot.*
- Words. Poems.* 1339. Wordsworth's Poems.
- Works in Prose and Verse.* 94. See *Pennycook's Works in Prose and Verse*.

Y.

- Yates's Account of New Zealand.* 2604. In the London Phil. Mag.
- Yardley Chase.* 1785. See *Cowper's Yardley Chase*.
- Year.* 1223. A poem by Bidlake.
- Young's Annals of Agriculture.* 615. Annals of Agriculture and other useful Arts. Published in Nos. Bury St. Edmunds, 1790—1804. 40 vols. 8vo.
- Young's Essex.* 1760. General View of the Agriculture of the County of Essex. By A. Young. London, 1807. 2 vols. 8vo.
- Young's Tour in Ireland.* 2335. See *Irish Tour*.

Z.

- Zan. Hist.* 1259. J. Zanoni. Rariorum Stirpium Historia: ex parte olim edita; nunc Centum plus Tabulis ampliata. Ed. C. Montio. Bonon. 1742. Fol. pl. 185.
- Zoological Journal.* 2140. Zoological Journal. London, 1825, &c. 8vo.

CONTENTS OF VOL. I.

The Roman numerals refer to the General Table of Contents, p. xvii. to eliii., where the species and varieties, with all their synonymes, will be found systematically arranged; the first column of Arabic figures, to the pages of the text in this volume; and the second, to those of the supplementary matter contained in Vol. IV.

The names of the half-hardy and suffruticose orders and genera are in small type.

	I.	I.	IV.		I.	I.	IV.
<i>Ranunculàcæ.</i>	xvii.	231	2534	<i>Cruciàcæ.</i>	-	xxi.	312
CLEMATIDÆ.	-	xvii.	232 2534	Vèlla	-	-	xxi. 312
Clématis	-	xvii.	232 2534	<i>Cress-Rocket.</i>	-	-	-
<i>Lady's Bower.</i>	-	-	-	Cheiranthus	-	xxi.	313 2538
<i>Traveler's Joy.</i>	-	-	-	Vesicaria	-	xxi.	313
<i>Virgin's Bower.</i>	-	-	-	Alyssum	-	xxi.	313
Atrágene	-	xvii.	246	Iberis	-	xxi.	313 2538
PÆONIÀCÆ.	-	xviii.	249 2535	Lepidium	-	xxi.	313
Pæònia	-	xviii.	249 2535	Sisymbrium	-	xxi.	313
<i>Tree Pæony.</i>	-	-	-	<i>Capparidàcæ.</i>	-	xxi.	313
Xanthorbiza L.	-	xviii.	255	Cápparis	-	xxi.	313
<i>Yellow Root.</i>	-	-	-	<i>Caper Bush.</i>	-	-	-
<i>Winteràcæ.</i>	-	xviii.	256	<i>Cistàcæ.</i>	-	xxi.	316 2538
Illicium	-	xviii.	256	Cistus L.	-	xxi.	317 2538
<i>Aniseed Tree.</i>	-	-	-	<i>Rock Rose.</i>	-	-	-
<i>Magnoliàcæ.</i>	-	xix.	259 2536	Helianthemum Tourn.	-	xxii.	317 2538
Magnòlia	-	xix.	259 2536	<i>Sun Rose.</i>	-	-	-
<i>Umbrella Tree.</i>	-	-	-	Hudsonia	-	xxv.	354
<i>Cucumber Tree.</i>	-	-	-	<i>Polygalàcæ.</i>	-	xxv.	354
<i>Indian Physic.</i>	-	-	-	Polygala L.	-	xxvi.	356 2538
Liriódéndron L.	-	xix.	284 2536	<i>Milkwort.</i>	-	-	-
<i>Tulip Tree.</i>	-	-	-	<i>Pittosporàcæ.</i>	-	xxvi.	356
<i>Mangifera</i>	-	xix.	291	Billardiæ Sm.	-	xxvi.	356
<i>Michelia</i>	-	xix.	291	<i>Apple Berry.</i>	-	-	-
<i>Dillenàcæ.</i>	-	xix.	292	Sillya Lindl.	-	xxvi.	357
Hibbértia	-	xix.	292	Pittosporum Banks	-	xxvi.	358
Anonàcæ.	-	xx.	293 2536	Senecio Dec.	-	xxvi.	359
Asínina Adans.	-	xx.	292 2536	Bursária Cur.	-	xxvi.	359
<i>Custard Apple.</i>	-	-	-	Cheiranthera Lindl.	-	xxvi.	359
<i>Schizandràcæ.</i>	-	xx.	295	<i>Caryophyllàcæ.</i>	-	xxvi.	359
Schizandra	-	xx.	295	Diáanthus	-	xxvi.	359
Sphero-stéma	-	xx.	295	<i>Tree Clove Pink.</i>	-	-	-
Kadsura	-	xx.	295	Silène L.	-	xxvi.	359
<i>Menispermàcæ.</i>	-	xx.	295	Arenária W.	-	xxvi.	359
Menispermum	-	xx.	296	Drypis L.	-	xxvi.	359
<i>Moonseed.</i>	-	-	-	<i>Linàcæ.</i>	-	xxvi.	360
Cócculus	-	xx.	297	Linum L.	-	xxvi.	360
<i>Ash Berberry.</i>	-	-	-	<i>Tree Flax.</i>	-	-	-
Berberàcæ.	-	xx.	298 2536	<i>Malvàcæ.</i>	-	xxvi.	360
Bérberis	-	xx.	298 2536	Lavátéra	-	xxvi.	360
<i>Berberry.</i>	-	-	-	<i>Tree Mallow.</i>	-	-	-
Mahònia Nutt.	-	xxi.	308 2537	Hibiscus	-	xxvi.	361
<i>Ash Berberry.</i>	-	-	-	<i>Althæa Frutex.</i>	-	-	-
				Sida Bonp.	-	xxvi.	363
				<i>Sterculiàcæ.</i>	-	xxv.	363
				Sterculia	-	xxv.	363

	I.	I.	IV.
<i>Tiliàcææ.</i>	-	xxvii.	364 2538
<i>Tilia L.</i>	-	xxvii.	364 2538
<i>Lime Tree.</i>			
<i>Linden Tree.</i>			
<i>Grewia L.</i>	-	xxvii.	376 2540
<i>Ternströmiaçææ.</i>		xxvii.	376 2540
<i>Malachodéndron Cav.</i>		xxvii.	377
<i>Stuártia Cav.</i>	-	xxviii.	378
<i>Gordônia Ellis</i>	-	xxxviii.	387 2540
<i>Loblolly Bay.</i>			
<i>Camelia L.</i>	-	xxviii.	381 2540
<i>Japan Rose.</i>			
<i>Thea L.</i>	-	xxix.	392
<i>Tea Tree.</i>			
<i>Eurya Royle</i>	-	xxix.	395
<i>Aurantiàcææ.</i>		xxix.	395 2540
<i>Citrus</i>		xxix.	395
<i>Orange Tree.</i>			
<i>Limonia Dec.</i>	-	xxix.	395
<i>Hypericàcææ.</i>		xxix.	397 2541
<i>Hypericum L.</i>	-	xxix.	397 2541
<i>St. John's Wort.</i>			
<i>Androsæmum Chois.</i>	-	xxx.	397
<i>Tutsan.</i>			
<i>Aceràcææ.</i>		xxx.	404 2541
<i>Acer</i>	-	xxx.	405 2541
<i>Maple.</i>			
<i>Sycamore.</i>			
<i>Negundo</i>	-	xxx.	460 2543
<i>Box Elder.</i>			
<i>Æsculàcææ.</i>		xxxii.	462 2543
<i>Æsculus</i>	-	xxxii.	462 2543
<i>Horschestnut.</i>			
<i>Pavia</i>	-	xxxii.	469 2543
<i>Buckeye.</i>			
<i>Sapinulàcææ.</i>		xxxiii.	474
<i>Kölreutéria Laxm.</i>	-	xxxiii.	475
<i>Dodonæa L.</i>	-	xxxiii.	476

	I.	I.	IV.
<i>Meliàcææ.</i>	-	xxxiii.	476
<i>Melia</i>		xxxiii.	476
<i>Red Tree, or Indian Libac.</i>			
<i>Vitacææ.</i>		xxxiii.	477 2544
<i>Vitis</i>	-	xxxiii.	477
<i>Grape Vine.</i>			
<i>Ampelopsis</i>	-	xxxiii.	481 2544
<i>Pine-leaved Ivy.</i>			
<i>Virginian Creeper.</i>			
<i>Cissus</i>	-	xxxiii.	483
<i>Geraniàcææ.</i>		xxxiv.	483
<i>Pelargonium</i>	-	xxxiv.	483
<i>Zygophyllacææ.</i>		xxxiv.	484
<i>Meliánthus</i>	-	xxxiv.	484
<i>Honey Flower.</i>			
<i>Zygophyllum</i>	-	xxxiv.	484
<i>Bian Capar.</i>			
<i>Rutàcææ.</i>		xxxiv.	484 2544
<i>Ruta</i>	-	xxxiv.	484 2544
<i>Rue.</i>			
<i>Aplophýllum Andr. Juss.</i>	xxxiv.		487
<i>Xanthoxylàcææ.</i>		xxxiv.	488 2544
<i>Xanthoxylum</i>	-	xxxiv.	488
<i>Toothache Tree, or Prickly Ash.</i>			
<i>Ptelea</i>	-	xxxiv.	489 2544
<i>Shrubby Trefoil.</i>			
<i>Ailántus</i>	-	xxxiv.	490
<i>Ailanto.</i>			
<i>Coriàcææ.</i>		xxxiv.	492 2545
<i>Coriària Niss.</i>	-	xxxiv.	492 2545
<i>Staphylacææ.</i>		xxxiv.	493
<i>Staphylèa</i>	-	xxxiv.	493
<i>Bladder-nut Tree.</i>			

ALPHABETICAL INDEX TO VOL. I.

	I.	I.	IV.		I.	I.	IV.
<i>Acer</i> - - -	xxx.	405	2541	<i>Hypericæcæ</i> - - -	xxix.	397	2541
<i>Aceracæ</i> - - -	xxx.	404	2541	<i>Hypérieum L.</i> - - -	xxix.	397	2541
<i>Æsculacæ</i> - - -	xxxii.	462	2543	<i>Iberis L.</i> - - -	xxi.	315	2538
<i>Æsculus L.</i> - - -	xxxii.	462	2543	Indian Lilac - - -	xxxiii.	476	
Ailanto - - -	xxxiv.	490		Indian Physic - - -	xix.	276	2536
<i>Ailântus</i> - - -	xxxiv.	490		Itcium L. - - -	xviii.	256	
<i>Althæa Frutex</i> - - -	xxvi.	361		<i>Kadsura</i> - - -	xx.	295	
Alyssum - - -	xxi.	513		<i>Kölreutéria Laxm.</i> - - -	xxxiii.	475	
<i>Ampelopsis</i> - - -	xxxiii.	481	2544	Ladies' Bower - - -	xvii.	232	2534
Anonacæ - - -	xx.	292	2536	Lavatera - - -	xxvi.	360	
<i>Androsæmum Chois.</i> - - -	xxx.	397		Lepidium - - -	xxi.	313	
Anised Tree - - -	xviii.	256		Lime Tree - - -	xxvii.	364	2538
<i>Apophýllum And. Juss.</i> - - -	xxxiv.	487		<i>Limonia Dec.</i> - - -	xxix.	395	
Apple Berry - - -	xxvi.	356		Linacæ - - -	xxvi.	360	
<i>Arenaria W.</i> - - -	xxvi.	359		Linum L. - - -	xxvi.	360	
Ash-barbery - - -	xxi.	308	2537	Liriodéndron L. - - -	xix.	284	2536
<i>Asimina Adans.</i> - - -	xx.	292	2536	Loblolly Bay - - -	x.	387	
<i>Atrágene L.</i> - - -	xviii.	246	2540	Magnolia - - -	xix.	260	2536
<i>Aurantiacæ</i> - - -	xxix.	395	2440	<i>Magnoliacæ</i> - - -	xix.	259	2536
Barberry - - -	xv.	298	2536	Mahonia Nutt. - - -	xxi.	308	2537
Bead Tree - - -	xxxiii.	476		Malachodéndron Cav. - - -	xxvii.	377	
Bean Caper - - -	xxxiv.	486		<i>Malracæ</i> - - -	xxvi.	360	
<i>Berberacæ</i> - - -	xx.	298	2536	<i>Manglietia</i> - - -	xix.	291	
<i>Berberis L.</i> - - -	xx.	298	2536	Maple - - -	xxx.	405	2541
Billardiæ Sm. - - -	xxvi.	356		Melia - - -	xxxiii.	476	
Bladder-nut Tree - - -	xxxiv.	493		<i>Meliacæ</i> - - -	xxxiii.	476	
Box Elder - - -	xxxii.	460	2543	Meliântus - - -	xxxiv.	484	
<i>Bursaria Cav.</i> - - -	xxvi.	359		<i>Menisperm'in rea</i> - - -	xx.	296	
Camellia L. - - -	xxviii.	381	2540	<i>Menispermum</i> - - -	xx.	296	
Caper Bush - - -	xxi.	313		Michelia - - -	xix.	291	
<i>Cappariacæ</i> - - -	xxi.	313		Milkwort - - -	xxvi.	356	2538
<i>Capparis L.</i> - - -	xxi.	313		Moonseed - - -	xx.	296	
<i>Caryophyllacæ</i> - - -	xxvi.	359		<i>Negundo</i> - - -	xxx.	460	2543
Cheiranthæra Lindl. - - -	xxvi.	359		Orange Tree - - -	xxix.	395	
Cheiranthus - - -	xxi.	512	2538	<i>Pæonia L.</i> - - -	xviii.	249	2535
Cissus L. - - -	xxxiii.	485		<i>Pæoniacæ</i> - - -	xviii.	249	2335
<i>Cistacæ</i> - - -	xxi.	316	2538	Pæony - - -	xviii.	249	2535
<i>Cistus L.</i> - - -	xxi.	317	2538	<i>Pavia</i> - - -	xxxii.	469	2543
Citrus L. - - -	lxxix.	395		Pelargonium - - -	xxiv.	485	
<i>Clematideæ L.</i> - - -	xvii.	232	2534	<i>Pittosporacæ</i> - - -	xxvi.	356	
<i>Clématis</i> - - -	xvii.	232	2534	<i>Pittosporum Banks</i> - - -	xxvi.	358	
Clove Pink Tree - - -	xxvi.	359		<i>Polygala L.</i> - - -	xxvi.	356	2538
<i>Cocculus Bauh.</i> - - -	xx.	297		<i>Polygalacæ</i> - - -	xxvi.	355	
<i>Coriaria</i> - - -	xxxiv.	492	2545	<i>Ptelea</i> - - -	xxxiv.	489	2544
<i>Coriaria Niss.</i> - - -	xxxiv.	492	2545	<i>Ranunculacæ</i> - - -	xvii.	231	2534
<i>Cruciacæ</i> - - -	xxi.	312	2538	Rock Rose - - -	xxi.	317	2538
Cucumber Tree - - -	xix.	273	2536	Rue - - -	xxxix.	484	2544
Custard Apple - - -	xx.	292	2536	<i>Ruta L.</i> - - -	xxxiv.	484	2544
Dianthus - - -	xxvi.	359		<i>Rutacæ</i> - - -	xxxiv.	484	2544
<i>Dilleniacæ</i> - - -	xix.	292		<i>Sapindacæ</i> - - -	xxxiii.	474	
<i>Dodonia L.</i> - - -	xxxiii.	476		Schizandra M. - - -	xx.	295	
<i>Drypis L.</i> - - -	xxvi.	359		Schizandracæ - - -	xx.	295	
<i>Eurya Koule</i> - - -	xxix.	395		Senécia Dec. - - -	xxvi.	359	
<i>Geraniacæ</i> - - -	xxxiv.	483		Shrubby Trefoil - - -	xxxiv.	489	
<i>Gordonia Ellis</i> - - -	xxviii.	378	2540	<i>Sida Bonp.</i> - - -	xxvi.	365	
Grape Vine - - -	xxxiii.	477		Silene L. - - -	xxvi.	359	
<i>Græwia L.</i> - - -	xxvii.	376	3540	Sisymbrium - - -	xxi.	313	
<i>Helianthemum Tourn.</i> - - -	xxii.	328	2538	<i>Sollya Lindl.</i> - - -	xxvi.	357	
<i>Ilibertia</i> - - -	xix.	292		Sphaerostema - - -	xx.	295	
<i>Hibiscus L.</i> - - -	xxvi.	361		Staphylæa - - -	xxxiv.	493	
Horsechestnut - - -	xxxii.	462	2543	<i>Staphyleacæ</i> - - -	xxxiv.	493	
<i>Hudsonia</i> - - -	xxv.	354		Sterculia - - -	xxvi.	363	
				<i>Sterculiacæ</i> - - -	xxvi.	363	

	I.	I.	IV.		I.	I.	IV.	
St. John's Wort	-	xxiv.	397	2541	Tutsan	-	xxx.	397
Stuartia Cav.	-	xxviii.	378		Umbrella Tree	-	xix.	269
Sun Rose	-	xxvii.	328	2538	Ulla L.	-	xvi.	312
Sycamore	-	xxx.	414	2542	Uscaria	-	xx.	313
Tea Tree	-	xxix.	392		Virginian Creeper	-	xxxiii.	481
Ternstroemia	-	xxvii.	376	2540	Virgin's Bower	-	xvii.	285
Thea L.	-	xxix.	392		Vitaceae	-	xxxiii.	477
Tilia L.	-	xxvii.	364	2538	Vitis	-	xxxiii.	477
Tiliaceae	-	xxvii.	364	2538	Winterbees	-	xviii.	256
Toothache Tree	-	xxxiv.	488		Xanthorrhiza L.	-	xviii.	255
Traveller's Joy	-	xvii.	235	2531	Xanthoxylaceae	-	xxxiv.	488
Tree Flax	-	xxvi.	360		Xanthoxylum L.	-	xxxiv.	488
Tree Willow	-	xxvi.	360		Yellow Root	-	xviii.	255
Tree Peony	-	xviii.	249		Zygophyllaceae	-	xxxiv.	481
Tulip Tree	-	xix.	284		Zygodium L.	-	xxxiv.	481

ARBORETUM ET FRUTICETUM BRITANNICUM.

INTRODUCTION.

THOUGH, from our title, the *Arboretum et Fruticetum Britannicum*, the reader may expect to find chiefly a history and description of the trees and shrubs which endure the open air in Britain, yet we mean to connect this history with that of the trees and shrubs of all similar climates throughout the world, in such a manner as to show what has been done in the way of introducing them, and what may be anticipated from future exertions. The *Arboretum et Fruticetum Britannicum* may, therefore, be considered as a General History of the Trees and Shrubs of Temperate Climates, but more especially of those of Britain.

Trees are not only, in appearance, the most striking and grand objects of the vegetable creation; but, in reality, they are those which contribute the most to human comfort and improvement. If cereal grasses and edible roots are essential for supplying food to sustain human existence, trees are not less so for supplying timber, without which, there could neither be the houses and furniture of civilised life, nor the machines of commerce and refinement. Man may live and be clothed in a savage, and even in a pastoral, state by herbaceous productions alone; but he cannot advance farther: he cannot till the ground, or build houses or ships, he cannot become an agriculturist or a merchant, without the use of trees.

Trees and shrubs also supply an important part of the food of mankind in many countries; besides all the more delicate luxuries of the table, and the noblest of human drinks in every part of the globe. The fruit of the palms, and of other trees of tropical climates, are as essential to the natives of those countries, as the corn and the edible roots of the herbaceous plants of temperate climates are to us. Wine, cider, arrack, and other liquors, are the products of trees and shrubs; as are also our more useful and exquisite fruits, the apple, pear, plum, peach, orange, mango, and many others. Not to insist in detail on the various

uses of trees and shrubs, it may be sufficient to observe, that there is hardly an art or a manufacture, in which timber, or some other ligneous product, is not, in one way or other, employed to produce it.

The use of trees in artificial plantations, in giving shelter or shade to lands exposed to high winds or to a burning sun, and in improving the climate and general appearance of whole tracts of country; in forming avenues to public or private roads, and in ornamenting our parks and pleasure-grounds, is too well known to require to be enlarged on here.

Every one feels that trees are among the grandest and most ornamental objects of natural scenery: what would landscapes be without them? Where would be the charm of hills, plains, valleys, rocks, rivers, cascades, lakes, or islands, without the hanging wood, the widely extended forest, the open grove, the scattered groups, the varied clothing, the shade and intricacy, the contrast, and the variety of form and colour, conferred by trees and shrubs? A tree is a grand object in itself; its bold perpendicular elevation, and its commanding attitude, render it sublime; and this expression is greatly heightened by our knowledge of its age, stability, and duration. The characteristic beauties of the general forms of trees are as various as their species; and equally so are the beauty and variety of the ramifications of their branches, spray, buds, leaves, flowers, and fruit. The changes in the colour of the foliage of trees, at different seasons of the year, alone form a source of ever-varying beauty, and of perpetual enjoyment to the lovers of nature. What can be more interesting than to watch the development of the buds of trees in spring, or the daily changes which take place in the colour of their foliage in autumn? — But to point out here all the various and characteristic beauties of trees, would be to anticipate what we shall have to say hereafter of the different species and varieties enumerated in our Work.

Shrubs, to many of the beauties of trees, frequently add those of herbaceous plants; and produce flowers, unequalled both for beauty and fragrance. What flower, for example, is comparable in beauty of form and colour, in fragrance, and in interesting associations, with the rose? The flower of the honeysuckle has been admired from the most remote antiquity, and forms as frequent an ornament of classic, as the rose does of Gothic, architecture. In British gardens, what could compensate us, in winter, for the arbutus and the laurustinus, or even the common laurel and the common ivy, as ornamental evergreens; for the flowers of the rhododendron, azalea, kalmia, and mezereon, in spring; or for the fruit of the gooseberry, currant, and raspberry, in summer? And what hedge plant, either in Europe or America, equals the common hawthorn? In short,

if trees may be compared to the columns which support the portico of a temple, shrubs may be considered as the statues which surmount its pediment, and as the sculptures which ornament its frieze.

It is not to be wondered at, that trees and shrubs should have excited the attention of mankind in all civilised countries, and that our accumulated experience respecting them should be considerable. The first characteristic instinct of civilised society is, to improve the natural productions by which we are surrounded; and the next is, by commerce to appropriate and establish in our own country the productions of others, while we give our own productions in exchange; and, thus, the tendency of all improvement seems to be to the equalisation of enjoyment, as well as to its increase.

Notwithstanding the use, the grandeur, and the beauty of timber trees, it is a fact, that, compared with herbaceous vegetables, the number of species distributed over the globe is comparatively small. The palms, the banana, the pine-apple, and other plants, popularly or botanically considered as trees or shrubs, though some of them attain a great height and thickness, are, with very few exceptions, of no use as timber. Almost all the timber trees of the world, with the exception of the bamboo, belong to what botanists denominate the dicotyledonous division of vegetables; and, perhaps, there are not a thousand genera of this division on the face of the earth which afford timber trees exceeding 30 ft. in height. The greater part of these genera, supposing such a number to exist, must belong to warm climates; for in the temperate zones, and in the regions of warm countries rendered temperate by their elevation, the number of genera containing timber trees 30 ft. in height, as far as hitherto discovered, does not amount to a hundred. The truth is, that between the tropics the greater number of species are ligneous, while in the temperate regions there are comparatively few, and in the frozen zone scarcely any. It may naturally be expected, therefore, that, in the temperate regions, there should only be a few timber trees which are indigenous to each particular country. In Britain, for example, there are not above a dozen genera of trees, furnishing in all about thirty species, which attain a height exceeding 30 ft.; but there are other countries of similar climates, all over the world, which furnish other genera and species, to what is, at present, an unknown extent; and it is the beautiful work of civilisation, of patriotism, and of adventure, first, to collect these all into our own country, and next, to distribute them into others. While Britain, therefore, not only enjoys the trees of the rest of Europe, of North America, of the mountains of South America, of India, and of China, she distributes her own trees, and those which she has appropriated, to each of these

countries respectively, and, in short, to all parts of the world; thus contributing almost imperceptibly, but yet most powerfully, to the progress and equalisation of civilisation and of happiness.

It must be interesting to the philosopher and the philanthropist, to know the precise position in which we stand relatively to this kind of interchange of natural productions. Much as has been done within the last century, there is reason to believe, from the number of countries unexplored, that this department of the civilisation of the great human family is yet in its infancy. Hence, in a work like the present, which professes to be a general history of the trees already in, or suitable for being introduced into, Great Britain, it seems desirable to commence with a general view of all other countries with reference to those trees which they contain which have been already introduced, or which, though we do not yet possess, we may expect to obtain and establish. This, therefore, will form PART I. of our Work; and we trust it will be found of considerable interest, by directing the attention of botanical collectors, travellers, and persons resident abroad, to specific objects of research.

In carrying this intention into effect, we shall commence by taking a general view of the trees and shrubs which were known to the ancients; we shall next give an enumeration of those which are indigenous to the British Islands; after which we shall treat of the introduction of foreign trees and shrubs into Britain, from the earliest records up to the end of the year 1834, the period at which this Work was commenced.

Having thus discussed the history of the trees and shrubs, native and foreign, of the British Islands, we shall next give a similar view of the indigenous and introduced trees and shrubs of all those other countries which possess, either by geography or altitude, climates in any degree analogous to that of Britain. This part of the Work will be concluded by a chapter on the literature of the trees and shrubs of temperate climates; in which the principal works which have appeared on the subject, both in Europe and America, will be enumerated.

The next division of our Work, PART II., will be devoted to the science of the study of trees. In this part, trees will be considered in all their various relations to nature and art. They will be considered as component parts of the general scenery of a country; in regard to the expression and character of particular kinds; in regard to the mode of delineating them pictorially, and of describing them popularly and botanically. They will also be considered with reference to uncultivated nature, to cultivated nature, and to man. This part will conclude with a summary of particulars to be taken into consideration, in preparing the description and natural and economical history of trees and shrubs, which are to follow as the third part of this Work.

PART III., which will form our next division, and that, indeed, which will comprise by far the greater part of the Work, will be the history and description of the different species and varieties of trees and shrubs, whether native or indigenous, useful or ornamental, at present cultivated in Britain.

We shall add to the perfectly hardy species the names, and short descriptive paragraphs, of some ligneous plants, which have been found by cultivators to be half-hardy in the climate of London; and of others, which, from their native countries and habits, we think not unlikely to prove so. We make this addition to the *Arboretum et Fruticetum Britannicum* for two purposes: in the first place, because, by trying species from all countries in the open air, some hitherto kept in hot-houses or green-houses may be found quite hardy; such having been the case with *Kérria japónica*, *Cydònia japónica*, *Hydránga Horténsia*, *Aúcuba japónica*, and a number of others. We may add, also, that, though the nature of a species cannot be so far altered as to fit an inhabitant of a very hot climate for a cold one, yet that the habits of individuals admit of considerable variation, and that some plants of warm climates are found to adapt themselves much more readily to cold climates than others. Thus, the common passion flower, according to Dr. Walker, when first introduced into the Edinburgh Botanic Garden, lost its leaves during winter; but, in the course of a few years, the same plant retained the greater part of them at that season. The same author relates that plants of the common yew, sent from Paris to Stockholm to plant certain designs by Le Nôtre, laid out there for the king of Sweden, all died, though the yew is a native of the latter country, as well as of France.

Every gardener must have observed that the common weeds which have sprung up in pots, in hot-beds or in hot-houses, when these pots happen to be set out in the open air during winter or spring are killed, or have their leaves injured; whilst the same species, which have sprung up in the open ground, are growing around them in a flourishing condition.

The obvious conclusions from these facts are, that the habits of plants admit of a certain degree of change with regard to the climate which they will bear; that the degree in which this power exists in any plant is only to be ascertained by experiment; and that the only mode of making these experiments is, by trying in the open air plants usually kept under glass. There is reason to believe, from trials already made, that many of the trees and shrubs of Australia, and particularly those of New Zealand and Van Diemen's Land, will ultimately become so habituated to the climate of London, as to live through the winter against a wall, with scarcely any protection.

The second reason why we have included a number of half-hardy, or supposed half-hardy, trees and shrubs in this Work is, that we think there are few scenes in an ornamental garden or pleasure-ground of greater interest to a person having any knowledge of botany, however slight, than a conservative wall; that is, a wall covered with trees and shrubs, natives of foreign climates, which, though they may be killed to the ground during winter, yet exhibit a degree of luxuriance during the summer season, which they never can display in our green-houses or conservatories. Even were all such plants to be killed by frost every winter, and a reserve obliged to be kept in green-houses or pits to supply their place every spring, still, the splendour of their appearance during the summer months, and the novelty of their forms when compared with those of the spring, usually grown in the open air in Britain, would far more than compensate for the trouble incurred. When we take into consideration how easy it is to have such walls flued, and to heat the borders in front of them by small pipes of hot water, the capacities of a conservative wall and border appear great beyond anything we can at present calculate on; and we are persuaded that, were the subject of conservative walls warmly taken up by a spirited and wealthy individual, something would be produced in this way, as superior to our present green-houses and conservatories, as these are to the orangeries and green-houses of the time of Evelyn, or even of Miller. Having thus given our reasons for the introduction of half-hardy species of trees and shrubs into this Work, we shall next submit a few words with respect to our arrangement and treatment of the hardy species.

The general arrangement of the *Arboretum et Fruticetum Britannicum* will be that of the natural system; by which, whether in botany, zoology, or mineralogy, those objects are brought together which resemble one another in the greatest number of particulars. In describing each species, we shall follow the summary of particulars laid down in the last chapter of Part II.; giving in succession the identification of the name by botanists; the synonymes, botanical and vernacular; references to published engravings; the specific character; the popular description; the geography; the history; the poetical and legendary allusions; the properties and uses; the soil and situation; the propagation and culture; the accidents and diseases which the plant is liable to; the insects and parasitic plants which inhabit it; examples of its growth in the British Islands and on the Continent; and, lastly, its price in some of the principal British, Continental, and American nurseries.

In identifying the name of every species, or variety, with that given to it by botanical writers, we have been guided principally by our own examination of the living plant, and comparing it

with existing engravings and descriptions; but we have also in part relied on the identifications of other botanists.*

In giving the synonymes, we have adopted the same rule as in giving the identifications.

Though we have used every exertion in our power to render this Work as correct, in a botanical and technical point of view, as possible, yet what we consider to be its great practical value (and that which will contribute more than any other cause to the main end which we have in view, viz. that of diffusing a taste for planting collections of trees and shrubs,) is, that we have described scarcely any tree or shrub which we have not seen ourselves, in a living state, within ten miles of London. Some exceptions are necessarily introduced; but, whenever this is done, it is either indicated by the paragraph relating to that tree or shrub being in very small type; or, by our indicating in words, or by some other means, that we have not seen the plant.

The shrubs, as far as practicable, we shall illustrate by engravings of botanical specimens, to be given along with the text; and all these will be to one and the same scale, of 2 in. to 1 ft. Engravings of the trees will be chiefly given in 8vo or 4to plates, apart from the text; and each of these plates will contain a pictorial portrait of the tree, and a botanical specimen. The pictorial portraits will be of two kinds: first, portraits of trees which have been planted ten or twelve years within ten miles of London, all to the same scale; and, secondly, portraits of full-grown trees, chiefly within the same limits, all to another scale.† The use of the first class of portraits is, to show, at a single glance, the comparative bulk which different trees attain in a given climate in a given period; as well as to indicate how far different kinds of trees, at this early age, show anything characteristic in their shape. The portraits of the full-grown trees, it is almost unnecessary to observe, are given in order to exhibit their ultimate magnitude and character. By merely glancing over these portraits, a planter will see at once, first, the effect which any given tree, purchasable in British nurseries, will produce at ten years' growth; and, secondly, what its appearance will be when it has arrived at its average size.

It will be found that in this Work we have, in various instances, reduced the number of species, and even, in some cases, of va-

* For example, in the case of the very first species described, *Clématis Flammula*, having convinced ourselves that our plant was that described by De Candolle (*Prod.*, vol. i. p. 2.), we have not hesitated to give the identifications and synonymes quoted by that eminent author; adding, however, the identifications and synonymes of subsequent authors from our own examination of their works. In the plant referred to we have added to the identifications of De Candolle, Hayne's *Dendrologia* and Don's *Miller*.

† See the Explanatory References, which precede the Table of Contents.

rieties; and this, had we been inclined to trust entirely to our own opinion, we might have carried to a much greater extent.

It is well known to the cultivators of trees and shrubs, that there are a great many names of species enumerated in botanical works, our own *Hortus Britannicus* not excepted, as having been introduced into this country, which are not to be found in any nursery, or even botanic garden. These plants may have been introduced and lost; or the names may have been those of plants already in the country, reintroduced under new names. In either case, according to the present mode of compiling botanical catalogues, the introduction of these names in such catalogues (provided the authorities are given with them) is unavoidable, whether the things to which they apply are in existence or not. It is easy to conceive some of the evils which, in a practical point of view, result from this mode of making catalogues; but it is necessary to be at once a practical botanist and a practical gardener, to comprehend the whole of them. One evil is, that, when collectors of trees, for example, order the plants bearing these names from the nurserymen, they either do not receive any plants at all, or they receive something which they do not want, and, probably, something which they already have. Another evil is, that nurserymen, in order to supply the demand for novelties, or to establish or keep up the appearance of having an extensive collection, too frequently introduce names into their catalogues for which they cannot supply plants; or they introduce synonymes without indicating that they are such. The effect of this is, that gentlemen intending to form collections, finding their intentions frustrated, frequently give up the pursuit in disgust.

Imperfect as are the collections, and erroneous as is the nomenclature in public nurseries, it will readily be conceived how difficult it is for a practical gardener to acquire a tolerable knowledge of the trees and shrubs actually in the country; and it is evident that, without this knowledge on his part, it is not to be expected that any but the most common trees and shrubs can be recommended by him to his employers: indeed, no British gardener, who has not passed some time in acquiring a knowledge of his profession in some of our principal botanic gardens, in the Kew Garden, in the garden of the Horticultural Society, in that of Messrs. Loddiges, in the Botanic Garden at Edinburgh, or in some other garden which contains an arboretum, can be said to know the names of one fourth of the trees and shrubs already in the country. The *Arboretum et Fruticetum Britannicum*, we think, will go far to remedy this evil, by enabling both gardeners and their employers to ascertain, not only what trees are in the country, but where they may see them growing. Nurserymen, by referring to these living trees, will not only

have an opportunity of correctly ascertaining the names of such as they already possess, but of supplying themselves with cuttings or plants of such sorts as they may not have in cultivation. The purchasers of trees, by always using the nomenclature of the *Arboretum Britannicum*, and being able to refer from it to the living specimens from which our engravings were taken, will at once insure certainty as to the kinds they obtain; and stimulate the nurserymen to accuracy, in regard to the names of those plants which they possess and propagate, and to the cultivation of a greater number of species and varieties. After the publication of our Work, it will be the fault of the nurseryman alone, if his nursery do not contain plants of all the species and varieties which we have figured and described.

Many persons, when recommended to plant, reply: "Of what use is it to plant at my age? I can never hope to live to see my plants become trees." This sort of answer does not, at first sight, appear surprising, if we suppose it to come from a person of sixty or seventy years of age; but we often hear it even from men of thirty or forty. In either case, such an answer is the result of a vulgar error, founded on mistaken and prejudiced notions. We shall prove its incorrectness by matters of fact. In the year 1830, there were many sorts of trees in the arboretum of Messrs. Loddiges which had been planted exactly ten years, and each of which exceeded 30 ft. in height. Most of these trees have since been cut down for want of room; but we have the names and the measurement of the whole of them. There are, also, at the present time (December, 1834), many trees in the arboretum of the London Horticultural Society's Garden at Chiswick, which have been only ten years planted, and which are between 30 ft. and 40 ft. in height. Why, then, should any one, even of seventy years of age, assign as a reason for declining planting, that he cannot hope to live to see his plants become trees? A tree 30 ft. high, practically speaking, will effect all the general purposes for which trees are planted: it will afford shelter and shade; display individual beauty and character; and confer expression on landscape scenery.

There is one subject which we shall occasionally touch on, in the history of particular species, and also in taking a general view of the trees of each genus, or of each natural order; and that is, the improvement which many species are probably susceptible of by cross-fecundation with other species nearly allied to them, or by procuring new varieties through the selection of remarkable individuals from seedlings raised in the common way. We shall also bear in mind the manner in which curious varieties are procured by the selection of shoots which present those anomalous appearances which gardeners call sports, and which, when propagated by grafting, continue to preserve their peculiarities. It should never be forgotten by cultivators, that all

our most valuable plants, whether in agriculture, horticulture, or floriculture, are more or less indebted for their excellence to art. Our cultivated fruit trees are very different from the same trees in a wild state; and our garden and field herbaceous vegetables so much so, that, in many instances, not even a botanist could recognise the wild and the cultivated plant to be the same species. There is reason to believe that the same means by which we have procured our improved varieties of fruit trees will be equally effective in producing improved varieties of timber trees. A few species, such as the oak, the elm, the magnolia, &c., have had improved varieties raised from seed by accidental crossing, or by the selection of individuals from multitudes of seedlings; and variegated varieties, and varieties with anomalously formed leaves, or with drooping or erect shoots, have been procured from the sports of parts of different plants. But the mode of improvement by cross-fecundation is yet quite in its infancy with respect to timber trees; and to set limits to the extent and beauty of the new varieties which may be produced by it is impossible. There is no reason why we may not have a purple-leaved oak, or elm, or ash, as well as a purple-leaved beech; or a drooping sweet chestnut as well as a drooping ash. The oak is a tree that varies astonishingly by culture; and, when the numerous American varieties that have been introduced into this country shall have once begun to bear seed, there is no end to the fine hybrids that may be originated between them and the European species. In short, we see no difficulty in improving our ornamental trees and shrubs to as great an extent as we have done our fruit trees and shrubs; though we are as yet only procuring new species from foreign countries, which may be considered as the raw material with which we are to operate.

PART IV., which will form the last division of the *Arboretum et Fruticetum Britannicum*, will be devoted to selected lists of the trees and shrubs described, classified according to their different capacities for fulfilling the various purposes for which trees and shrubs are required by the planter and by the landscape-gardener. For the rest we refer to the Table of Contents.

The utility of such a Work as the *Arboretum et Fruticetum Britannicum* to the gardening world, and to the landed proprietor, will not, we think, be questioned. We shall say nothing, therefore, of the influence which it cannot fail to have in promoting a taste for the culture and spread of such foreign trees as we have already in the country; and in exciting a desire for introducing others from different parts of the world, and for originating new varieties by the different means employed by art for that purpose. One remark, however, we may be permitted to make on the use of such a Work as the *Arboretum et Fruticetum Britannicum* to gentlemen of landed property. Every

proprietor of a landed estate is either a planter, or possesses trees already planted. If he is in the former case, he will learn from this Work to combine beauty with utility, by planting, in the outer margins of his natural woods or artificial plantations, and along the open rides in them, and in the hedgerows of his lanes and public roads, trees which are at once highly ornamental and more or less useful — in some cases, perhaps, even more useful — than the common indigenous trees for which they are substituted. If, on the other hand, his estate is already fully planted, he will learn from this Work how he may beautify his plantations by a mode which never yet has been applied in a general way to forest trees; viz., by heading down large trees of the common species, and grafting on them foreign species of the same genus. This is a common practice in orchards of fruit trees; and why it should not be so in parks and pleasure-grounds, along the margins of woods, and in the trees of hedgerows, no other reason can be assigned than that it has not hitherto been generally thought of. Hawthorn hedges are common everywhere; and there are between twenty and thirty beautiful species and varieties of thorn in our nurseries, which might be grafted on them. Why should not proprietors of wealth and taste desire their gardeners to graft some of the rare and beautiful sorts of tree thorns on the common hawthorn bushes, at intervals, so as to form standard trees, in such of their hedges as border public roads? And why should not the scarlet oak and the scarlet acer be grafted on the common species of these genera, along the margins of woods and plantations? Such improvements the more strongly recommend themselves, because, to many, they would involve no extra expense; and, in every case, the effect would be almost immediate. Every gardener can graft and bud; and every landed proprietor can procure stock plants from nurseries, from which he can take the grafts; or he may get scions from botanic gardens, the garden of the London Horticultural Society, that of the Caledonian Horticultural Society, or the Dublin Garden at Glasnevin.

Amateur landscape-gardeners, and architects who lay out the grounds of the houses they have designed, will be enabled, by this Work, to choose the kinds of trees which they think will produce the best effect in their plantations; and, what is of much more consequence, which will produce a certain effect within a given number of years. Indeed, the want of such a Work as the *Arboretum et Fruticetum Britannicum* to professional landscape-gardeners, and a conviction of the great use it would be of to practical gardeners, and to all persons engaged in laying out grounds, or in forming ornamental plantations, first suggested to us the idea of commencing the Work.

In modern landscape-gardening, considered as a fine art, all the more important beauties and effects produced by the artist

may be said to depend on the use which he makes of foreign trees and shrubs. Our reasons for this are grounded on the principle that all art, to be acknowledged as such, must be avowed. This is the case in the fine arts: there is no attempt to conceal art in music, poetry, painting, or sculpture; none in architecture; and none in the geometrical style of landscape-gardening. Why should there be an attempt to conceal art in modern landscape-gardening? Because, we shall be told, it is an art which imitates nature. But, does not landscape-painting also imitate nature; and yet, in it, the work produced is acknowledged to be one of art? Before this point is settled, it is necessary to recur to what is meant by the imitation of nature, and to reflect on the difference between repetition and imitation. In what are called the imitative arts, it will be found that the imitation is always made in such a manner as to produce a totally distinct work from the thing imitated; and never, on any account, so like as to be mistaken for it. In landscape-painting, scenery is represented by colours on a flat surface; in sculpture, forms, which in nature are coloured, are represented in colourless stone. The intention of the artist, in both cases, is not to produce a copy which shall be mistaken for the original, but rather to show the original through the medium of a particular description of art; to reflect nature as in a glass. Now, to render landscape-gardening a fine art, some analogous process must be adopted by the landscape-gardener. In the geometrical style, he has succeeded perfectly, by arranging grounds and trees in artificial surfaces, forms, and lines, so different from nature as to be recognised at once as works of art. A residence thus laid out is clearly distinguished from the woody scenery of the surrounding country; and is satisfactory, because it displays the working of the human mind, and confers distinction on the owner as a man of wealth and taste. A residence laid out in the modern style, with the surface of the ground disposed in imitation of the undulations of nature, and the trees scattered over it in groups and masses, neither in straight lines, nor cut into artificial shapes, might be mistaken for nature, were not the trees planted chiefly of foreign kinds not to be met with in the natural or general scenery of the country. Every thing in modern landscape-gardening, therefore, depends on the use of foreign trees and shrubs; and, when it is once properly understood that no residence in the modern style can have a claim to be considered as laid out in good taste, in which all the trees and shrubs employed are not either foreign ones, or improved varieties of indigenous ones, the grounds of every country seat, from the cottage to the mansion, will become an arboretum, differing only in the number of species which it contains.

Though a taste for trees has existed from the earliest ages, that taste, in this country at least, may still be considered in its

infancy. An English landowner is almost always a great respecter of trees generally, but seldom knows anything of particular sorts: he, therefore, cares very little for their individual beauties, and contents himself with being an indiscriminate admirer of them. Hence the unwillingness of most persons to cut down trees, however improperly they may be placed; or to thin out plantations, however much they may be crowded, and however great may be the injury which the finer foreign sorts are sustaining from the coarser-growing indigenous kinds. This indiscriminate regard for trees, and morbid feeling with reference to cutting them down when they are wrongly placed or too thick, principally results from ignorance of the kinds and of the relative beauty of the different species, and from want of taste in landscape-gardening. When we consider that it is not much above a century since American trees began to be purchasable in the nurseries of this country, this is not to be wondered at; and, more especially, when it is remembered that planters, generally speaking, have few opportunities of seeing specimens of these trees, so as to become acquainted with them, and thus to acquire a taste for this kind of beauty and its pursuit. The public botanic and horticultural gardens, and the private arboreta and collections of foreign trees and shrubs, now establishing throughout the country; and the mode now becoming general among nurserymen, of planting specimen trees in their nurseries; will tend to remedy this defect, by exhibiting living specimens: and our *Arboretum et Fruticetum Britannicum* will, we trust, aid in attaining the same end.

To artists, the *Arboretum et Fruticetum Britannicum* will not be without its use. It is well known that there are but few landscape-painters who possess that kind of knowledge of trees which is necessary to enable them to produce such portraits as would indicate the kind to a gardener or a forester. This defect, on the part of landscape-painters, arises partly from their copying from one another in towns, rather than from nature in the country; but, principally, from their want of what may be technically called botanical knowledge. The correct touch of a tree, to use the language of art, can no more be acquired without studying the mode of foliation of that tree, than the correct mouldings of a Grecian or Gothic cornice can be understood or represented without the study of Grecian or Gothic architecture. It is for this reason that it will always be found that ladies who reside in the country, and have studied botany, if they have a taste for landscape, will imitate the touch of trees better than professional landscape-painters. We assert it as a fact, without the least hesitation, that the majority of British artists (we may say, of all artists whatever) do not even know the means of acquiring a scientific knowledge of the touch of trees; almost the only works which have noticed the subject, and gone beyond the

mere surface, being the *Remarks on Forest Scenery*, by Gilpin; and Kennion's *Essay on Trees in Landscape*. The perusal of the *Arboretum et Fruticetum Britannicum*, and the comparison of the botanical specimens with the touch to which they give rise in the portraits, will enable artists to investigate from our figures, and afterwards from nature, those differences in the points of the shoots, in the clustering and form of the foliage, and between the appearance of the foliage of spring and that of autumn, which give rise to the difference of touch necessary to characterise a species, and to mark the season of the year. Most artists who have studied trees from nature can give the touch of the oak with characteristic expression; and, by the study of the details of other trees, they may attain a touch which shall characterise them with equal force and accuracy. There is no work extant, however, from which an artist can study, correctly and scientifically, the touch of more species of trees than the oak, the ash, the weeping willow, and one or two others. In proof of this we may refer to the plates in Kennion's work above referred to, as one of the latest and best, where the engravings, in the greater number of instances, have not the slightest resemblance to the trees the names of which are written beneath them. How, under these circumstances, is it possible for an artist, who is not a botanist, and who does not reside in the country, to study the touch of trees? By the *Arboretum et Fruticetum Britannicum* he may acquire as much botanical knowledge as will enable him to distinguish with certainty all the different species of trees to be found in this country; and he will see, in the engravings of the botanical specimens as they appear in autumn, the foundations laid in nature for the different descriptions of touch. The London artist, in addition to the botanical knowledge which he may acquire from our work, may have recourse to the specimen trees (all near London) from which our portraits were taken. Artists generally, by becoming botanically acquainted with the trees, will be able to recognise them in their walks, or professional excursions; to study them under various circumstances, and, when they introduce them in their landscapes, to give their characters with fidelity.

Hitherto there has not been a sufficient demand for this kind of skill on the part of the artist; but, as foreign trees become better known by the public generally, it will be necessary for artists to keep their art on a level with the state of knowledge of the times in which they live. As the foreign trees which are every year being introduced into the country advance in size, the truth of this remark will become more and more obvious.

Having now given a general outline of the plan of our Work, and of the manner in which we propose to carry that plan into execution, we shall next proceed with PART I.

PART I.

GENERAL OUTLINE OF THE HISTORY AND GEOGRAPHY OF THE
TREES AND SHRUBS OF TEMPERATE CLIMATES.

THE use of the slight general outline which we propose now submitting to the reader is, partly, to show the consideration in which trees have been held in all ages and countries, but principally to record what has been done in the introduction of foreign trees into Britain; and to point out, from the ligneous productions of other countries having similar climates, what remains to be accomplished. We shall first notice to what extent a love for, and a knowledge of, trees existed among the nations of antiquity; and, next, give a general idea of the indigenous and introduced trees of those countries occupied by the modern nations of Europe. We shall commence with Britain; and shall take, in succession, France, Germany, and the other European countries. Afterwards, we shall give a slight sketch of the trees suited to temperate climates which are natives of Asia, Africa, America and Australia.

CHAP. I.

OF THE KNOWLEDGE OF TREES AND SHRUBS WHICH EXISTED AMONG
THE NATIONS OF ANTIQUITY.

THE first notices which we have of trees are in the Sacred Writings. The tree of knowledge, and the circumstance of our first parents hiding themselves among the trees of the garden of Eden, are familiar to every one. Solomon appears to have collected all kinds of plants, and not only to have had an orchard of fruit trees, and trees bearing spices, but to have included in his grounds what are called barren trees, and among these the cedar. As this tree is a native of a cold and mountainous country at some distance from Judea, it shows that the practice of collecting trees from a distance, and from a different climate, to assemble them in one plantation or arboretum, is of the earliest date. The cedar, indeed, is frequently mentioned in Scripture; and both that and the fir (including, under this name, probably both *Pinus* and *Abies*, for some species of both are natives of Asia Minor and Greece) are said, in the book of Ezekiel, to be frequent in magnificent gardens. Large trees were then used as places for meeting under (as they are, in the East, to this day); and they were then, as now, planted in cemeteries.

Trees are mentioned in the writings of Hesiod and Homer. The garden of Alcinous is said to have contained various sorts of fruit trees: and directions are given in Hesiod for lopping the poplar, and other species, for fuel; and felling the oak, the elm, and other kinds of large trees, for timber.

The principal trees of the *Egyptians*, according to Herodotus, were, the palm, the sycamore fig, the lote tree (*Urtica australis*, according to Mr. Hogg, *Gard. Mag.*, x. 291.), the olive, and the pomegranate. There are, we know, several other trees which are natives of Egypt; but these were probably thought most worthy of being recorded, as producing edible fruit.

The gardens of the *Persians* contained trees; and those in the garden of the younger Cyrus, at Sardis, were all planted with his own hand, in straight lines; the only mode which, at that early period, when scarcely any but indigenous trees were in use by planters, could convey the expression of art and design. In general, the trees which most attracted the attention of the ancients were those which bore edible fruits, produced spices, had a terebinthine odour, or possessed spreading branches to afford shade. Hence the frequent mention of the palm, the fig, the olive, the cinnamon, the camphor, the cypress, the sycamore fig, and the plane.

The only positive source of information respecting the trees known to the nations of antiquity, down to the time of the *Greeks*, is to be found in the works of Theophrastus. Stackhouse, in his edition of Theophrastus's *Historia Plantarum*, has endeavoured to show the modern botanical names for the plants of which Theophrastus has treated. Sprengel had done the same thing in his *Historia Rei Herbariæ*. Stackhouse has added to his own identifications as many of those of Sprengel as are different from, and supplementary to, his own. From both we have selected the following list of the ligneous species. Stackhouse has stated in the preface to his second volume (his work is in two volumes, 1813, 1814), that Sprengel has carefully ascertained 357 of the kinds treated of by Theophrastus, and that he has passed over the rest, which are nearly as many in number, in silence; except remarking the circumstances which make them so ambiguous as to render the identifying of them hopeless. To some of the identifications which have been proposed, doubt appertains; and, in the case of the ligneous species, in the enumeration below, this doubt is expressed by notes of interrogation. It may be observed, that the greater number of these plants, according to Sibthorp's *Flora Græca*, are natives of Greece, and that most of those which are not, will endure the open air, or are cultivated, in that country. The whole of them, with scarcely any exceptions, are in British gardens and hot-houses; and all those which we have marked

with a star, are indigenous to the British Isles. We have thrown the species into the groups indicated by the natural orders, in order to aid the memory of the botanical reader, and to facilitate generalisation.

Ranunculææ. *Clématis orientàlis.*

Capparidéæ. *Cápparis spinòsa.*

Cistinææ. *Cistus créticus* and *salviaefòlius.*

Malvæææ. *Gossýpium arbòreum*; *Málva tomentòsa*; *Hibiscus*, an arboreous species if a hibiscus.

Tiliæææ. *Tília * europæ'a.*

Aurantiæææ. *Citrus Médica* and *Aurántium.*

Acerinææ. *A'cer * campéstre* and ** Pseùdo-Plátanus.*

Ampélidææ. *Vítis vinífera*, 3 kinds, and *índica.*

Rutæææ. *Rùta gravèolens.*

Celastrinææ. ? *Celástrus*, the species was, in habit, a tree of middle size; ** Euónymus europæ'us.*

Rhámneææ. *Rhámnus lyciòides* and ? *Alatérnus*, *Paliùrus aculeàtus*, *Zizyphus vulgàris.*

Terebinthæææ. *Pistàcia Terebínthus* and *Lentíscus*, *Balsamodéndron gileadense*, *Rhús Coriària más*, *C. fœ'mina*, and *Cótinus*, ? *Mangífera índica.*

Leguminòsææ. *Acàcia arábica*, *Sénegal*, *Cátechu*, *myrrhífera Stackhouse*, and *polyacántha*; ? *Tamaríndus índica*; *Cércis Siliquástrum*; *Colùtea arboréscens* and *cruénta*; *Cathartocárpus Fístula*; *Cýtisis Labúrnum* and "Maránta;" *Ceratònia Siliqua*; *Morínga pterygospérma*; *Genísta lusitánica* and ? *Scórpius (Spártium villòsum Flora Græca)*; *Medicàgo arbòrea*; ? *Coronílla E'merus*; *E'benus crética*; *Astrágalus Tragacántha.*

Rosæææ. *Ròsa*, the 5-leaved, the 10-leaved, the 20-leaved, and the 100-leaved; *Rùbus * fruticòsus*, ** cæ'sius*, and ** idæ'us.*

Amygdàleææ. *Amýgdalus commùnis*; ? *Pérsica vulgàris*; *Cérasus * Pàdus*, *durácina* and ? *Laurocérasus*; *Prùnus * in-sítitia* and *doméstica var. Juliána.*

Pomæææ. *Pýrus * commùnis* wild, and cultivated; ** Màlus* wild, and cultivated, ** Aria*, and *crética*; ** Sórbus*, two varieties of; ** ? torminàlis* and ? a variety of; *Méspilus * ? germánica*, ? a variety of; *Cydònia vulgàris*, wild and cultivated; ? *Ame-lánchier vulgàris*; *Cotoneáster * vulgàris*; *Cratægus Pyracántha*, *Azaròlus*, and ** Oxyacántha.*

Granàteææ. *Pùnica Granàtum.*

Rhizophòreææ. *Rhizòphora Mángle.*

Tamariscínieææ. ** Tamarix gállica.*

Myrtæææ. *Mýrtus commùnis.*

Cactæææ. *Opúntia vulgàris.*

? *Grossulæææ.* ? *Ribes * Grossulària.*

Umbelliferaææ. *Bùbon Gálbanum.*

- Araliæceæ.* *Hédera* * *Hèlix*, and varieties of it.
- Caprifoliæceæ.* *Caprifolium* * *Periclymenum*; ? *Lonicera pyrenæica*; *Viburnum* * ? *Lantana*, * ? *Ópus* and ? *Tinus*; *Sambucus* * *nigra*.
- Córneæ.* *Córnus* más and * *sanguínea*.
- Loranthæceæ.* *Loránthus* europæus; * *Viscum álbum*.
- Compósitæ.* *Santolina rosmarinifolia*; *Helichrysum Stœchas*; *Kentrophýllum* (*Onobròma*) arboréscens; *Conyza saxatilis*.
- Vacciniæceæ.* *Vaccínium* * *Vitis idæa*.
- Ericæceæ.* *Erica*; *Arbutus Andrâchne* and * *Uncdo*; ? *Rhododéndron pónticum*.
- Styracíneæ.* *Stýrax officinále*.
- Ebenæceæ.* *Diospýros Lótus*.
- Oleæceæ.* *Olea* europæa, sylvéstris; *Phillyrea latifolia*; ? *Ligústrum vulgàre*; *Ornus* europæa; * ? *Fráxinus excélsior*.
- Apocýneæ.* ? *Nèrium Oleánder*; * *Vínca* màjor, or * *mínor*.
- Cordiæceæ.* *Córdia Sebestèna* and *Mýxa*.
- Solâneæ.* *Cápsicum frutéscentis*; *Lýcium bárbarum* and ? europæum.
- Labiâtæ.* *Lavándula Spîca*: *Rosmarînus officinális*; *Sálvia tríloba*, crética; *Teucrium créticum*; ? *Marrùbium Pseudo-Dictámnus*, *Oríganum Dictámnus*, *Tournefórtii*, and ? *ægyptiâcum*; *Saturèja capitata*; *Thýmus vulgàris*, ? *Mastichîna*.
- Verbenæceæ.* *Vítex Ágnus cástus*.
- Plantagíneæ.* *Plantâgo* ? *Cýnops*.
- Chenopòdeæ.* *Salicórnica fruticòsa*.
- Lauríneæ.* *Laúrus nóbilis*, var. *platyphýlla* and var. *leptophýlla*; *Cinnamònum vèrum*.
- Myristiceæ.* *Myrística*.
- Thymelææ.* *Dáphne Cncòrum* and *seríceæ*.
- Euphorbiæceæ.* *Euphórbia Pithyûsa*, * *Charâcias*, and *Myrsinîtes*; * *Búxus sempervîrens*.
- Urticeæ.* *Ficus Caprificus*, *Cárica*, religiósa, and *Sycómorus Mòrus nigra*.
- Ulmæceæ.* *Ulmus campéstris*, ? and another kind; *Céltis austrális*.
- Piperæceæ.* *Piper nigrum*.
- Juglândææ.* *Júglans règia*.
- Salicíneæ.* *Sàlix* * *álba*, *nigra*; * *Hèlix variegàta* and *babylónica*; *Pópulus* * *nigra*, * *álba*.
- Betulíneæ.* *Bétula* * *álba* and ? *Álnus* * *glutinòsa*; *Álnus oblongàta*.
- Cupuliferaæ.* *Quércus* ? *Flex*, *Sùber*, *coccífera*, *Ballòta*, *Ægilops*, *Æsculus*, *Cérris*, * *Ròbur*, *fagínea Desf.*, and *Tournefórtii*; * *Còrylus Avellána*; *Castànca* * *vésca*; *Fâgus* * *sylvática*; *O'strya vulgàris*.
- Platâneæ.* *Plátanus orientális*.

Coníferæ. Cèdrus Libàni; Làrix europæa; Pinus halepensis, Pínea, marítima, and *sylvéstris; Abies excélsa and Pícea; Cuprèssus sempervirens; Thùja aphýlla; Táxus * baccàta; Juníperus * commùnis, Oxýcedrus, lýcia, and * nàna.

Cycùdeæ. Cýcas revolùta.

Smiláceæ. Sinilax áspera; Rúscus hypophýllum, * aculeàtus ? racemòsus.

Pálma. Phœnix dactylífera, and four varieties; Còcos nucífera; Chamærops; ? Arèca Câtechu.

Gramíneæ. Bambùsa arundinàcea.

It thus appears that the total number of species known to Theophrastus was not less than 170, which belong to 53 groups or natural orders.

The Romans appear to have begun with a knowledge of all the trees possessed by the Greeks; and there are added to them, in their works, almost all the trees of the colder regions of Europe. It is evident that the Romans introduced trees into Italy from other countries; because frequent mention is made, by their agricultural writers, of the platanus, the cedar, the cypress, and other trees, which are not indigenous to Italy; and the cherry, the peach, and other fruits, we are informed, were imported from Persia. The pine, the bay, the plane, and the box appear to have been the favourite trees of gardens: the first, for its refreshing odour; the second, for its beauty, and because it was used in crowning martial heroes; and the third, on account of its shade. Pliny observes, "In old times trees were the very temples of the gods; and, according to that ancient manner, the plain and simple peasants of the country, savouring still of antiquity, do at this day consecrate to one god or other the goodliest and fairest trees that they can meete withall; and verily, we ourselves adore, not with more reverence and devotion, the stately images of gods within our temples (made though they be of glittering gold and beautiful ivory), than the very groves and tufts of trees, wherein we worship the same gods in religious silence. First, the ancient ceremony of dedicating this and that kind of tree to several gods, as proper and peculiar to them, was always observed, and continues to this day. For the great mighty oak, named esculus, is consecrated to Jupiter, the laurel to Apollo, the olive to Minerva, the myrtle to Venus, and the poplar to Hercules." (*Holland's Translation of Pliny's Natural History*, p. 357.)

The Romans cultivated trees for useful purposes, like the moderns. They planted coppice woods, for fuel, fence wood, and props for the vine: they had osier grounds, for producing hoop and basket willows; single rows of elms and poplars, for supporting the vine and they had indigenous forests on the hills

and mountains, for supplying timber for building and other purposes. The larch was a favourite tree among them; and instances are given by Pliny of the enormous size which it attained, of its durability, and its resistance to fire. The positive knowledge of the Romans, with respect to trees, may be found in Pliny's *Natural History*; and an enumeration of the species which that work contains, as far as they can be guessed at by modern botanists, is given by Sprengel in his *Historia Rei Herbaria*, vol. i. It contains so few, in addition to those known to Theophrastus, that it seems unnecessary to introduce it here.

CHAP. II.

OF THE HISTORY AND GEOGRAPHY OF THE TREES AND SHRUBS NOW IN THE BRITISH ISLANDS.

THE trees and shrubs now indigenous to, or cultivated in, the British Islands, including Ireland, the Channel Islands, and the other adjacent isles, naturally form two divisions; viz., those which are of spontaneous growth, and those which have been introduced or originated by the art and industry of man. In order to convey distinct ideas respecting the number and nature of the native woody plants which may be considered as aboriginal and permanent inhabitants of this country, we shall give an enumeration of them, arranged according to the natural orders to which they belong; and, in order to show the progress of the introduction of foreign species, the number introduced, and the individuals to whom we are indebted for such introductions, we shall treat of the native and foreign plants separately.

SECT. I. *Of the Native Trees and Shrubs of the British Isles.*

THE native plants of any island may generally be considered as the same as those of the continent to which that island belongs; and hence we find that there are very few trees and shrubs which are indigenous to the north of France, Belgium, and the north of Germany, which are not also natives of Britain. In countries which have been long civilised, it appears difficult to determine what trees or plants are aboriginal, and what have been introduced; and even in wild countries, the same difficulties may be said to occur, since the seeds of the plants of one country may be, and undoubtedly are, carried to another country by birds and other animals, and may spring up there, mature themselves, and continue their species like aboriginal plants. It is probable that this process has gone on more or less in every country from its

first existence; and thus, that the tendency even of nature, independently of human art, is to equalise the productions of similar climates.

Cæsar, in his *Commentaries*, states that he found the woods of Britain to contain the same trees as those of Gaul, with the exception of the abies and the fagus: "Materia cujusque generis, ut in Gallia est, præter fagum et abietem." This passage has given rise to much controversy, some authors supposing that Cæsar, by the word abietem, meant the wild or Scotch pine, which is a native of Britain. As, however, the Romans designated the silver fir abies, there can be little doubt that this was the tree alluded to by Cæsar; which not only does not grow wild in England, but was not introduced into this country till modern times. This solution of the passage is so simple, that it is difficult to imagine how any mistake can have arisen, particularly as Pliny speaks of the Scotch pine expressly as *Pinus sylvestris* (*Nat. Hist.*, lib. xv. and xvi.) The only reason appears to be, that the Scotch pine was formerly called the Scotch fir; and that the word abies, being considered to signify fir, was, without further examination, supposed to apply to that tree.

It is more difficult to reconcile Cæsar's assertion that he did not find the fagus in Britain, as that name is generally supposed to have been applied by the Romans to the common beech. Belon informs us that, in his time (*Les Obs., &c., en Grèce, en Asie, et autres Pays étrangers*, 1554), on Mount Athos and in Macedonia, the beech was called phega. It is wonderful, therefore, says Ray, that Cæsar should deny the beech to Britain: his doing so can only be satisfactorily accounted for, by supposing that by the word fagus he meant the *Quercus Æsculus*, the phagos of Theophrastus. Mr. Long, in his *Observations on certain Roman Roads, and Towns, in the South of Britain*, p. 36., asserts that the tree Cæsar called fagus was the sweet chestnut, *Fagus Castanea L.* Mr. Long does not state his grounds for this opinion; but should the fagus of the Romans be our chestnut, and their castanea our beech, it would not only explain this difficulty, but do much to reconcile that passage in the *Georgics*, lib. ii. v. 71., where the fruit of the fagus appears preferred to that of the castanea. If we consider that by fagus Cæsar meant our common beech, all that can be concluded from his remark is, that the beech was not, in his time, discoverable in large masses in Kent; where, though it grows naturally, it is only found on the hills and not in the plains. Mr. Whitaker, in his *History of Manchester*, concludes that the Romans introduced the beech, partly from the assertion of Cæsar above alluded to, and partly from the name for the beech in the British language, foighe, faghe, faydh, being obviously derived from fagus. The name in the Anglo-

Saxon is *bece*; *bèche*, Fr. ; and *buche*, Ger. The Scotch pine, Whitaker thinks, was a native of the island before the arrival of the Romans, though unknown to them at their invasion ; and this pine, judging from the resinous quality of that dug up from peat bogs, he considers to have been of a different species from that now grown in England, and “the same assuredly with the Scotch fir of the highlands of Scotland.” Here we know the author to be entirely mistaken ; the species being every where the same, and the quality of the timber differing only in consequence of differences in the soil and situation. Sir Walter Scott fell into a similar error, when he stated, in the *Quarterly Review* (xxxvi. 580.), that our “Scotch fir was brought from Canada, not more than half a century ago,” and that the true species, found in the north of Scotland in immense forests, grows with “huge contorted arms, not altogether unlike the oak.” The conjectures of these two eminent writers only show that, however great may be their authority in other matters, they are not to be depended upon in what concerns trees. The Scotch pine must, unquestionably, have been indigenous in the highlands of Scotland in the time of Cæsar, though, in all probability, not to be met with, or rare, in England, at least in the southern counties.

The Romans, Whitaker observes, first brought among us, as their present names sufficiently show, “the platanus or plane, the *tilia* or *teil*, the *buxus* or *box*, the *ulmus* or *elm*, and the *populus* or *poplar*. The platanus passed from Asia to Sicily, thence into Italy ; and, before the year 79, as Pliny informs us, it had reached the most northerly shore of Gaul. The apple Whitaker conjectures to have been brought into Britain by the first colonies of the natives, and by the *Hædri* of Somersetshire in particular ; hence Glastonbury was distinguished by the title of *Avellonia*, or the apple orchard, previously to the arrival of the Romans. Before the third century, this fruit had spread over the whole island, and so widely, that, according to Solinus, there were large plantations of it in the “*Ultima Thule*.” The Romans added “the pear, the damson, and the cherry, the *arbor persica*, perch, or peach ; *aprica*, or apricot ; and *cydonia*, or quince.” Cherries were introduced from Pontus and Egypt into Italy by Lucullus, who conquered the former country ; and they were carried into Britain within five years of the first settlement of the Romans in the country. Pears abounded in Italy, though it is uncertain at what time the Romans brought them into England. The damson was originally brought from Damascus to Italy, and thence to Britain, as the quince was from Crete, and the peach from Persia : the latter was common in Gaul in the time of Agricola.

The mulberry, the chestnut, the fig, and the sorbus, or true service, were introduced by the Romans. It is singular, that, not far from one of the very few habitats in which the true

service is to be found in a wild state in Britain, viz., Wyre Forest in Worcestershire, the remains of a Roman villa were some years ago discovered (see *Arch. Mag.*, ii. p. 94.). It is not improbable that the tree referred to may be a descendant from a service tree planted in the orchard belonging to the adjoining Roman villa. The chestnut belongs to Sardis in Asia Minor; and it was brought thence to Tarentum and Naples, where it was cultivated with great success in the reign of Vespasian. That the chestnut was in Britain as early as the 12th century is placed beyond dispute by Giraldus Cambrensis, who, in speaking of the trees of Britain which Ireland wants, mentions the chestnut and the beech. Daines Barrington conjectures that the chestnut was probably brought into England from Spain; and Dr. Ducarel, who had a dispute with Barrington on the subject (see *Phil. Trans.*, lix. and lxi.), endeavours to prove that it is a native. Mr. Whitaker thinks, and, in our opinion, with great reason, that the tree was brought into Britain by the Romans. The medlar, according to Pliny, was brought into Italy from Greece, at what period is uncertain; as is also when it was introduced into Britain. The rose was brought from Italy by the Romans, the best being those of Prænestina and Campania. The rosemary and the thyme are also supposed to have been introduced by the Romans. The thyme, in the days of Vespasian, Pliny observes (xxi. 10), so greatly overspread the plains in the province of Narbonne, that many thousands of cattle were brought every year from the distant parts of the country to fatten upon it.

In a paper on the subject of indigenous trees, in the *Archæologia*, by Daines Barrington, he lays down a test by which it may be known what trees ought to be considered as truly indigenous: that they grow in large masses, and spread over a considerable breadth of surface; that such masses never end abruptly, except where there is a sudden change in the soil or the substratum, and, that the trees or shrubs ripen their seeds kindly, and that when these seeds are dropped, they spring up freely. Applying these tests to what are commonly considered native trees, he rejects positively the sweet chestnut, the lime, the English elm, and the box. As doubtful, he reckons the *Acer Pseudo-Platanus*, and the white poplar (*Pópulus álba*), and even the yew, which, he says, is seldom found but in churchyards or in artificial plantations. He also doubts the spindle tree and the privet. A few lime trees, he thinks, such as those in Moor Park in Hertfordshire, and on the river Neath in Glamorganshire, have been introduced by the alien abbots and priors, when they came to visit their religious houses; but the tree was not generally planted till after the time of Le Nôtre, in the reign of Charles I., who introduced it extensively in avenues, as was then

the custom in France. The antiquity of the sweet chestnut at Tortworth, which he had ascertained from Lord Ducie to be much exaggerated, he alleges to be no proof that the tree is indigenous. "The English, or narrow-leaved, elm," he says, "being much esteemed by the Romans, was probably introduced by them. The box," he erroneously (see *Herb.*, 1597, p. 1226.) states, "is not mentioned by Gerard, and," he adds, "the tree is found nowhere in an apparently wild state, except on Box Hill, where it was planted by Lord Arundel, who designed to build a house there, but who relinquished his intention from the want of water, and built one at Albury hard by." The only native evergreen trees and shrubs of Britain would thus appear to be the Scotch pine, the holly, the juniper, the furze, the spurge laurel, the butcher's broom, and the ivy. The furze Dr. Walker supposes not to be aboriginal, but to have been introduced from the mountains of Portugal, where it abounds. His reason is, that it is the only alleged indigenous shrub which flowers during winter; and that during severe winters it is killed to the ground, both in England and Scotland. According to these authors, the only indigenous evergreen trees are the Scotch pine and the holly; so that we are thus reduced to two evergreen trees and four evergreen shrubs; unless we include such under-shrubs as the heath, the *Andrómeda*, the *Arctostaphylos Uva úrsi*, &c., which do not generally attain the height of two feet.

Perhaps it may be thought unreasonable to allege that the lime and the yew are not natives of Britain, since they unquestionably are of countries which lie farther north; viz., the north of Germany and Sweden: but it must be remembered that the summers of those countries are hotter than those of England, in consequence of which, the lime ripens its seeds every year, which it seldom does in Britain. In countries without extremes either of heat or cold, such as the sea coast of Britain and great part of Ireland, many trees will live and thrive without ever producing seeds. Such trees may remain for ages in a country, without being one step nearer naturalisation than the day on which they were introduced. In Hasted's *Kent* it is stated that Sir John Speilman, who introduced the manufacture of paper into England from Germany, in the time of Elizabeth, and to whom Queen Elizabeth granted the manor of Portbridge in Dartford, introduced the lime tree. He is said to have brought over two trees with him in his portmanteau, and to have planted them at Portbridge, near the dwelling-house belonging to the powder mills; where, according to Hasted, they remained till they were cut down a few years previously to the time when he wrote, which was in 1776. (*Beauties of England, &c., Kent*, p. 562.) The lime, however, is represented by Turner as growing to a large size in 1562; so that the trees introduced by Speilman could

not have been the first that were brought into the country. The *Tilia europæa*, or common lime tree of the north of Europe, is stated by Turner and Gerard to be a native of England; but Ray says, that, though it is an inhabitant of Essex, it is never found in that county, or anywhere else, growing wild. The *Tilia parvifolia*, Ray seems to consider as a native.

The box is one of our most interesting "disputed trees;" for, if we are deprived of that and of the yew, neither of which Daines Barrington will allow us, our only evergreen trees will be the Scotch pine and the holly. Ray says that "the box grows wild on Box Hill, hence the name: also at Boxwell, on Cotswold in Gloucestershire, and at Boxley in Kent, where there were woods of this tree, according to Aubrey. It grows plentifully on the chalk hills near Dunstable." Turner says, "it groweth on the mountains in Germany plentifully, wild, without any setting; but in England it groweth not by itself in any place that I know, though there is much of it in England." (*Herbal*, edit. 1551, p. 159.) Parkinson says it is found in many woods, and that it is also planted in orchards. Evelyn considers it a native, as does Lambarde, in his *Perambulations of Kent*, in 1576. Some curious controversial matter on this subject will be found in the *Gentleman's Magazine*, vol. lvii., for 1787. One writer, T. H. White (p. 667), says, "he called at the village of Boxley, and that, from the strictest enquiries, he was thoroughly convinced that Evelyn was wrong in considering the box to grow wild at this village." It has been said that the Earl of Arundel, who died in Italy in 1646, planted the box trees on Box Hill, with a view to building a house there; but this is denied by another writer, S. H., in the same magazine. "The Earl of Arundel," this writer says, "was a very curious man; and, having a house very near, at Dorking, it has been conjectured, but without foundation, that he planted Box Hill. The ground on which the box trees grow," he continues, "was not His Lordship's property;" and this is confirmed by a passage in Manning and Bray's *Surrey*, where that part of the hill which is covered with the trees is proved to have belonged to Sir Matthew Brown, long before the date when they were said to have been planted by the earl. "Various have been the disquisitions," say these authors, "concerning the antiquity of this plantation, which, however, for aught that has hitherto appeared to the contrary, may have been coeval with the soil. Here was formerly also a warren, with its lodge; in a lease of which, from Sir Matthew Brown to Thomas Constable, dated 25th August, 1602, the tenant covenants to use his best endeavours for preserving the yew, box, and all other trees growing thereupon; as also to deliver, half-yearly, an account of what hath been sold, to whom, and at what prices; and in an account rendered to Ambrose,

his son, by his guardian, of the rents and profits for one year, to Michaelmas, 1608, the receipt of box trees cut down upon the sheepwalk on this hill is 50*l.* We have seen also an account of this manor, taken in 1712, in which it is supposed that as much had been cut down within a few years before as amounted to 3000*l.* (*Manning and Bray's Surrey*, i. 560.) At present the only habitat of this tree in England is Box Hill; and though this circumstance cannot be considered as a proof that it is not indigenous, yet, as it is known that it does not ripen its seeds freely in this country, and seldom sows itself, either on Box Hill or anywhere else, when in a neglected state, we may fairly be allowed, when these circumstances are taken into consideration and conjoined with its Roman name, to doubt whether it be a native. It is so beautiful a tree, that its branches, like those of the bay, were probably in early use both in civic festivals and religious ceremonies; and it appears likely that it was not only introduced, but was cultivated, at an early period. At the same time, it must not be forgotten, that, in estimating the probability of a tree or plant being indigenous to a country, we must add to the other considerations mention that of its native habitat. Now the native site of the box is in woods of deciduous trees, where it is well known a plant may propagate itself by seeds, which would not do so on naked exposed situations. Taking this view of the subject, the box may yet be a native.

The English elm (*Ulmus campestris*) seldom ripens its seeds in England, though it does so freely in the neighbourhood of Paris. It can hardly be considered a native. The common sycamore ripens its seeds kindly, and in woods it sows itself, and the seeds spring up freely; but this may be said of various trees and shrubs which we know are foreign to the soil. The white poplar is found so seldom that it can hardly be considered a native. The yew is found in inaccessible acclivities, and other places where it must have been sown by birds, which is also the case with the spindle tree and the privet; therefore, their being natives cannot reasonably be doubted, except on something like positive evidence.

The trees and shrubs which were known to our Saxon ancestors were, the birch, alder, oak, wild or Scotch pine, mountain ash or rowan tree, juniper, elder, sweet gale, dog rose, heath, St. John's wort, and the mistletoe. All these are considered as aboriginal in the country; but, from the length of time that England was under the government of the Romans, it may reasonably be supposed that, in addition to the native trees and shrubs, there were in the country, when it was taken possession of by the Saxons, several which were natives of France, Spain, or Italy. To what extent this was the case cannot now be known; but it is sufficient for our purpose, that, in the present

day, botanists consider all those plants indigenous to a country, which have existed in it beyond the memory of man or the existence of written records, and which propagate themselves freely by seed, without human agency.

The reputed native plants of Britain have been enumerated and described by different botanical authors: but it will be of little practical use in this case, and in the others which will come before us, to quote from any author who wrote previously to the time of Linnæus; and who, of course, could not adopt his admirable system of giving plants specific names composed of two words, instead of short Latin descriptions. The first author who enumerated the plants of England, and applied the Linnæan specific names, was Hudson, in his *Flora Anglica*, published in 1762; and those of Scotland were first described by Lightfoot, in his *Flora Scotica*, in 1775. Those of Ireland were first enumerated by Threlkeld, in 1727, before the Linnæan system was adopted, and there has not yet been any other flora of the country than a list published by Mr. T. Mackay in 1825. Fortunately, however, there are two recent works, the *English Flora* of Sir J. E. Smith, and the *British Flora* of Dr. Hooker, which contain an enumeration and description of all the plants indigenous to the British Isles, and from them we have compiled the following enumeration. In it are included all the plants, considered by botanists as ligneous, which grow in the British Islands, exclusive of varieties.

To such as are considered by many persons as doubtful natives, we have prefixed, not the point of interrogation used to signify botanical doubts, but the letters qu.

Ranunculaceæ. Clématis Vitálba, a deciduous climber.

Berberídeæ. *Bérberis vulgaris*, a deciduous shrub, 10 ft. high.

Cistíneæ. Heliánthemum marifòlium, surrejànum, vulgàre, tomentòsum, and poliifòlium, evergreen prostrate shrubs, from 6 in. to 1 ft. in height.

Tiliàceæ. Tília europæa qu., a deciduous tree, 50 ft.; grandifòlia (syn. platyphýlla) qu., a deciduous tree, 50 ft.; parvifòlia, a deciduous tree, 30 ft.

Hypericíneæ. Androsæmum officinàle, a deciduous under-shrub, 4 ft.; Hypéricum calycìnum qu., an evergreen under-shrub, 1 ft.

Accríneæ. Acer campéstre, a deciduous tree, 20 ft. high; and *A. Pseùdo-Plátanus* qu., a deciduous tree, 50 ft. high.

Celastríneæ. Euónymus europæus, a deciduous tree, from 15 ft. to 20 ft. high.

Ilicíneæ. Ilex Aquifòlium, an evergreen tree, 30 ft. high.

Staphyleàceæ. Staphylèa pinnàta qu., a deciduous shrub, from 10 ft. to 15 ft. high.

Rhámneæ. *Rhámnus cathárticus* mas, c. fœm., deciduous shrubs, from 10 ft. to 15 ft. high; *Frángula*, a deciduous shrub retaining its leaves late, 5 ft. high.

Leguminòsæ. *Cýtisis scopàrius*, an evergreen shrub, of 5 ft.; *Genísta tinctòria*, an evergreen under-shrub, of 18 inches; *G. pilòsa*, a prostrate evergreen shrub; *G. ánglica*, a prostrate deciduous shrub; *Ulex europæ'a*, an evergreen shrub, of 5 ft.; *U. nána*, an evergreen shrub, of 2 ft.; *U. strícta*, and *U. e. fl. plèno*.

Rosàcææ. *Ròsa cinnamòmea* qu., *rubélla*, *spinosíssima*, *involuta*, *Doniàna*, *grácilis*, *Sabini*, *villòsa*, *tomentòsa*, *Sherárdi*, *rubiginòsa*, *micrántha*, *Bórreri*, *cæ'sia*, *sarmentàcea*, *bractéscens*, *dumetòrum*, *Fórsteri*, *hibérnica*, *canina*, and *sýstyla*, all deciduous shrubs, from 3 ft. to 5 ft.; and *R. arvénsis*, a deciduous trailing shrub; *Rèbus fruticòsus*, *plicàtus*, *rhamnifòlius*, *leucòstachys*, *glandulòsus*, *nítidus*, *affinis*, and *corylifòlius*, all evergreen trailers; *R. cæ'sius*, a deciduous trailer; and *R. suberéctus* and *idæ'us*, deciduous under-shrubs, of 3 ft. Some more species, or reputed species, might be added to the evergreen trailers, from Dr. Lindley's *Synopsis* and our *Hortus Britannicus*. *Potentilla fruticòsa*, a deciduous shrub, above 3 ft.; *Cómarum palústre*, a prostrate deciduous under-shrub, of 1 ft.; *Spiræ'a salicifòlia* qu., a deciduous under-shrub of 3 ft.

Pomàcææ. *Pýrus commúnis*, *Málus torminális*, *doméstica*, *aucupària*, and *pinnatífida*, all deciduous trees, of between 20 ft. and 30 ft.; and *P. Aria*, and *Aria intermèdia*, deciduous trees, between 30 ft. and 40 ft.; *Cratæ'gus Oxyacántha*, and *Méspilus germánica* qu., deciduous trees, between 15 ft. and 20 ft.; and *Cotoneáster vulgàris*, a deciduous shrub, 4 ft. high.

Amygdàlææ. *Cérasus Pádus* and *ávium*, and *Prúnus doméstica* qu., deciduous trees between 20 ft. and 30 ft.; *P. insitítia* and *spinòsa*, deciduous shrubs or very low trees, of 10 ft. or 15 ft.

Tamariscíneæ. *Támarix gállica* qu., an evergreen shrub, between 5 ft. and 10 ft. high.

Grossulàcææ. *Ribes rubrum*, *petræ'um*, *alpínium* mas, a. fœm., *nigrum*, *Grossulària* qu., and *U'va críspa* qu., all deciduous under-shrubs, from 1 ft. to 3 ft. in a wild state.

Araliàcææ. *Hédera Hélix*, a prostrate and clinging evergreen shrub.

Caprifoliàcææ. *Caprifòlium itálicum* qu., and *Periclymenum*, deciduous twiners; *Lonicera Xylòsteum* qu., a deciduous shrub, 10 ft. high; *Sambucus nígra*, a deciduous tree, 15 ft. or 20 ft. high; *Vibúrnum Opulus* and *Lantàna*, deciduous shrubs or very low trees, from 10 ft. to 15 ft. high.

Córneæ. *Córnus sanguínea*, a deciduous shrub or very low tree, from 10 ft. to 15 ft. high.

Loranthàcææ. *Víscum álbum* mas., a. fœm., evergreen parasites.

Vacciniææ. *Vaccinium Myrtillus* and *uliginosum*, deciduous shrubs, 1 ft. high; and *Vitis idæa*, an evergreen shrub, under a foot high; *Oxycoccus vulgaris*, a prostrate evergreen shrub.

Ericæææ. *Calluna vulgaris*, a prostrate evergreen shrub, about a foot in height; *Erica Tétralis cinerea*, *ciliaris*, *mediterranea*, *Mackaiàna*, and *vàgans*, and *Dabæcia poliifolia*, *D. p. álba*, and *cærùlea*, and *Andrómeda poliifolia*, evergreen shrubs, under a foot high; *Arbutus Unedo* qu., an evergreen shrub, 15 ft. high; *Arctostáphylos Uva úrsi*, a prostrate evergreen shrub, and *alpina*, a prostrate deciduous shrub; and *Chamælèdon procumbens*, a prostrate evergreen shrub.

Olcæææ. *Ligústrum vulgàre*, a deciduous shrub, 6 ft. high; *Fráxinus excélsior*, a deciduous tree, 80 ft. high; *heterophýlla*, a deciduous tree, 30 ft. high.

Apocýnææ. *Vinca minor* qu., an evergreen prostrate shrub.

Solànææ. *Solànum Dulcamàra*, a deciduous trailer.

Chenopòdææ. *Chenopodium fruticòsum*, an evergreen shrub, between 3 ft. and 4 ft. high; *Atriplex portulacòides*, a prostrate evergreen shrub, under a foot in height.

Thymelæææ. *Dáphne Laurèola*, an evergreen shrub, 2 ft.; *Mezèrcum* qu., a deciduous shrub, 3 ft.

Elæágnææ. *Hippóphæe rhamnòides* mas, rh. fœm., deciduous shrubs or very low trees, 15 ft.

Euphorbiæææ. *Euphòrbia Charàcias* qu., and *amygdalòides*, evergreen fruticulose shrubs, 2 ft. high; and *Búxus semper-virens* qu., an evergreen tree, from 8 ft. to 20 ft. high.

Ulmæææ. *Ulmus campéstris* qu., a deciduous tree, of 80 ft.; *U. suberòsa* qu., *màjor* qu., and *montàna*, deciduous trees, of 40 ft.; and *U. glàbra* qu., a deciduous tree, of 60 ft.

Cupulíferææ. *Quércus Ròbur* and *sessiliflòra*, deciduous trees, of 80 ft.; *Castànea véscà* qu., a deciduous tree, of 60 ft.; *Fàgus sylvàtica*, a deciduous tree, of 70 ft.; *Córylus Avellàna*, a deciduous shrub, of 10 ft.; and *Cárpinus Bétulus*, a deciduous tree, of 35 ft.

Betulíneææ. *Bétula álba*, a deciduous tree, of 40 ft.; and *B. nàna*, a deciduous under-shrub, of 2 ft.

Salicíneææ. *Sàlix frágilis* mas and *frágilis fœmina*, *Russelliàna* mas, *Russell. fœm.*, *álba* mas, a. fœm., *álba var. cærùlea* mas, a. c. fœm., deciduous trees, of 40 ft. high; *S. triándra* mas, t. fœm., *lanceolàta* mas, l. fœm., *pentáudra* mas, p. fœm., *petiolàris* mas, p. fœm., *vitellìna* mas, v. fœm., *decípiens* mas, d. fœm., *rùbra* mas, r. fœm., *cinèrea* mas, c. fœm., *oleæ-fòlia* mas, o. fœm., *hírta* mas, h. fœm., *càprea* mas, c. fœm., *acuminàta* mas, a. fœm., and *viminàlis* mas, v. fœm., all deciduous trees, 20 ft. or 25 ft. high; *Hoffmanniàna* mas, H. fœm., *amygdàlina* mas, a. fœm., *nígricans* mas, n. fœm., *Borreriàna* mas, B. fœm., *nítens* mas, n. fœm., *Davalliàna* mas, D. fœm., *Wulfeniàna* mas, W. fœm., *tétrapla* mas, t. fœm., *bícolor* mas,

b. fœm., tenuifolia mas, t. fœm., malifolia mas, m. fœm., purpurea mas, p. fœm., *Helix* mas, *H.* fœm., *Lambertiàna* mas, *L.* fœm., *Forbyàna* mas., *F.* fœm., *Croweàna* mas, *C.* fœm., *prunifolia* mas, *p.* fœm., *venulosa* mas., *v.* fœm., *carinata* mas, *c.* fœm., *Stuartiàna* mas, *S.* fœm., *arenaria* mas, *a.* fœm., *lanata* mas, *l.* fœm., *argentea* mas, *a.* fœm., *Doniàna* mas, *D.* fœm., *aurata* mas, *a.* fœm., *aquatica* mas, *a.* fœm., *cotinifolia* mas, *c.* fœm., *rupèstris* mas, *r.* fœm., *Andersoniàna* mas, *A.* fœm., *Forsteriàna* mas, *F.* fœm., *sphacelata* mas, *s.* fœm., *Smithiàna* mas, *S.* fœm., and *stipularis* mas, *s.* fœm., all deciduous shrubs, from 3 ft. to 15 ft. high; *phlycæfolia* mas, *p.* fœm., *vacciniifolia* mas, *v.* fœm., *Myrsinites* mas, *M.* fœm., *Dicksoniàna* mas, *D.* fœm., *arbuscula* mas, *a.* fœm., *livida* mas, *l.* fœm., *glauca* mas, *g.* fœm., *fusca* mas, *f.* fœm., *incubacea* mas, *i.* fœm., and *rosmarinifolia* mas, *r.* fœm., all deciduous, from 1 ft. to 3 ft. high; herbacea mas, *h.* fœm., *reticulata* mas, *r.* fœm., *ætida* mas, *f.* fœm., *repens* mas, *r.* fœm., and *prostrata* mas, *p.* fœm., prostrate deciduous shrubs, under a foot in height. Nearly the whole grow in moist ground. *Pópulus álba* qu. mas, *a.* fœm., *trémula* mas, *t.* fœm., *nigra* mas, *n.* fœm.; and *canescens* mas, *c.* fœm., deciduous trees, from 40 ft. to 60 ft. high.

Myricæ. *Myrica Gåle* mas, *G.* fœm., deciduous shrubs, 2 ft. high.

Conifera. *Pinus sylvèstris*, an evergreen tree, from 60 ft. to 80 ft. high; *Táxus baccata* mas, *b.* fœm., and var. *hibèrnica*, evergreen trees, 20 ft. to 30 ft. high; *Juniperus commùnis* mas, *c.* fœm., evergreen shrubs, from 5 ft. to 7 ft. high; *nàna* mas, *n.* fœm., prostrate evergreen shrubs.

Empètrea. *Empetrum nigrum* mas, *n.* fœm., evergreen prostrate shrubs.

Smilæca. *Rúseus aculeatus* mas, *a.* fœm., and var. *laxus*, evergreen shrubs, from 1 ft. to 2 ft. high.

In estimating the heights of these trees and shrubs, we have supposed them to be growing in their natural and ordinary habitats. Under culture, or even in a wild state under favourable circumstances, many of them would grow higher, particularly the roses, the willows, and the fruticulose plants. The number of the latter might have been increased, by adding the carnation, the pink, &c., which, even as indigenous plants, are certainly as much fruticulose as *Euphòrbia Charácias*, or *E. amygdalöides*.

The above enumeration includes 71 genera, and about 200 species, nearly 100 of which are willows, roses, and brambles; and these species are comprised in 37 groups or natural orders. In greater detail, they are:—

27 deciduous trees, from 30 ft. to 60 ft. in height.

28 deciduous trees, from 15 ft. to 30 ft. in height.

1 evergreen tree, from 60 ft. to 80 ft., the Scotch pine.

3 evergreen trees from 15 ft. to 30 ft., the box, the yew, and the holly.

65 deciduous shrubs, and very low trees, from 5 ft. to 18 ft.; including 21 roses and 32 willows.

26 deciduous shrubs, from 1 ft. to 5 ft.; including 6 roses and 10 willows.

5 evergreen shrubs, from 5 ft. to 15 ft.

7 evergreen shrubs, from 1 ft. to 5 ft.

1 evergreen climber, the ivy.

1 deciduous climber, the clematis.

2 deciduous twiners, honeysuckles.

8 evergreen trailers, brambles.

3 deciduous trailers: the *Ròsa arvensis*, the *Solànum Dulcamàra*, and the *Rùbus cæsius*.

13 evergreen shrubs, or fruticulose plants, from 6 in. to 1 ft. in height; such as the *Vaccinium Vitis idæa*, the ericas, *Andrómeda polliifòlia*, &c.

10 deciduous shrubs, or fruticulose plants, from 3 in. to 1 ft. in height; such as *Cómarum palústre*, *Vaccinium Myrtillus*, *Sàlix reticulàta*, *prostràta*, &c.

SECT. II. *Of the Foreign Trees and Shrubs introduced into the British Isles.*

IF wild plants are said to follow those animals to which they supply food, cultivated plants are the followers of man in a state of civilisation. In all cases of taking possession of a new country, the first step of the settlers has been to introduce those vegetables which, in their own country, they knew to be the most productive of human food; because the natural resource of man for subsistence is the ground. In all temperate climates, the plants of necessity may be considered to be the cereal grasses and the edible roots. Trees, with the exception of such as bear edible fruit, are not introduced till a considerable period afterwards; because all new and uncivilised countries abound in forests of timber. It can only be when this timber becomes scarce, or when wealth and taste have increased to such an extent as to create a desire for new trees as objects of curiosity, that the practice takes place of cultivating indigenous trees, or of introducing new ones. Hence we find that, in England, all the timber required for the purposes of construction and fuel was obtained from the native forests and copses, till about the time of Henry VIII. In this reign and the next, Holinshed informs us that plantations of trees began to be made for purposes of utility; and we find, in the same reign, that attention began to be paid to the trees and shrubs of foreign countries, and that some few, even at that early period in the history of

British tree culture, began to be introduced into our gardens, as objects of rarity and value.

The ornamental trees, or the trees of curiosity, that would first be introduced into any country after those that recommended themselves by their fruit or their medicinal virtues, would be such as were generally planted about houses and in gardens, or such as bore conspicuous seeds. Hence the eypress, the bay, the box, the elm, the lime, and the plane, as being domestic shrubs and trees; and the chestnut, the ilex, the walnut, and the pine, as being trees with conspicuous seeds, would, we may suppose, be those that were first brought over by the Romans, or by the heads of religious houses, ambassadors, or travellers.

In tracing the introduction of foreign trees into this country, from the earliest ages to the present time, we shall first collect such notices as we have been able to obtain of the period from the invasion of the country by the Romans, to the end of the 15th century: and, next, take in succession the 16th, 17th, 18th, and 19th centuries.

SUBJECT. 1. *Of the Foreign Trees and Shrubs introduced into Britain by the Romans, and during the Middle Ages, to the End of the 15th Century.*

THERE can be no doubt whatever that the Romans introduced most of our cultivated vegetables and fruits. Some curious proofs of this are occasionally found in the springing up of Italian plants in the neighbourhood of the ruins of Roman villas, where ground, which had long remained in a state of rest, had been turned over in search of antiquities. Though, as far as we know, no trees or shrubs of Italy have sprung up in this manner from dormant seeds; yet there cannot be a doubt but that some of the trees and shrubs of the Romans would be cultivated in the gardens of their governors and generals, most of whom, it is understood, must have been practically acquainted with husbandry. Such trees would not only be interesting to them as reminding them of their native country, but they would serve to decorate and distinguish their residences, and command the admiration of the Roman army and of the natives.

We have seen, in the preceding chapter (p. 22.), that most of our fruit trees, and in all probability the plane, chestnut, walnut, lime, elm, and box, were introduced by the Romans. Many trees and shrubs introduced by the Romans, or by the monks of the middle ages, may have been afterwards lost; because this is, sooner or later, the case with all neglected plants that are placed in a climate which will not enable them to ripen their seeds.

In the 9th century, during the reign of Charlemagne, some exertions appear to have been made in France for the extension

of orchards; but nothing has reached us respecting the barren trees and ornamental shrubs of that period, either in France or England.

In the tenth century, monasteries and other religious establishments began to abound in the country; and the monks and clergy, who were their principal occupants, were generally either natives of foreign countries, or had been educated in Italy. The occupants of monasteries have, in all times, been attached to gardening; and, among the plants which those of Britain probably introduced from Italy, there can be little doubt that fruit trees were included, and probably, also, some trees of ornament, and shrubs. The sweet bay and the arbutus, if they were not introduced by the Romans, were, in all probability, brought over by the monks. It is conjectured by Dr. Walker (*Essays on Nat. Hist.*), that some trees and shrubs were introduced from the Holy Land during the time of the crusades; and one of these, he thinks, was the English elm. In the dispute already noticed (p. 23.), between Daines Barrington and Dr. Ducarel, on the question of the sweet chestnut being indigenous, the latter refers to a record, dated in the time of Henry II., by which the Earl of Hereford grants to Flexby Abbey the title of all his chestnuts in the Forest of Dean. It appears highly probable that the chestnut, being so productive of human food in Italy in the time of the Romans, would be introduced by them, wherever they went, as one of the most useful of trees.

In the beginning of the 13th century, the apple appears to have been cultivated to some extent in Norfolk. In the 6th of King John (1205), Robert de Evermere was found to hold his lordship of Redham and Stokesly, in Norfolk, by petty serjeantry, the paying of 200 pearmaines, and 4 hogsheads (modios) of wine, made of pearmaines, into the exchequer, at the feast of St. Michael yearly. (*Blomfield's Norfolk*, ii. 242. 4to edit., 1810.)

At the beginning of the 15th century, the rose appears to have been not only known, but in extensive cultivation. Sir William Clopton granted to Thomas Smyth a piece of ground called Dokmedwe in Hausted, for the annual payment of a rose, at the nativity of St. John the Baptist, to Sir William and his heirs, in lieu of all services, dated at Hausted, on Sunday next before the Feast of All Saints, 3 Henry IV. (1402). (*Cullum's Hawsted*, p. 117.)

In explanation of this deed, it may first be observed that ancient deeds are often dated on a Sunday, being executed in churches or churchyards, for the greater notoriety: in the second place, the rose was then in much more extensive use in cultivated society than it is now, when its place is partly occupied by the great variety of other flowers now in cultivation. The demand for roses formerly was so great, that bushels of

them were frequently paid by vassals to their lords, both in France and England. The single rose paid as an acknowledgment was the diminutive representation of a bushel of roses; as a single peppercorn, which is still a reserved rent, is of a pound of peppercorns, a payment originally of some worth, descending by degrees to a mere formality. (*Histoire de la Vie privée des François*, ii. 221., and *Cullum's Hawsted*, 117, 118.)

The well-known story of the quarrel in the Temple Gardens, about 1450, which gave rise to the distinctions of the white and red rose in the wars of York and Lancaster, is in unison with the foregoing authorities.

Towards the end of this century, parks for hunting became common in England, and bushes in gardens were clipped; but we have no evidence that in either case foreign trees or shrubs were made use of; unless, with Daines Barrington, we reckon the yew tree as such. The yew is mentioned in these times as subjected to the topiary operations of the gardener; and there appears little doubt that it was then reckoned one of the principal garden shrubs, and almost the only evergreen one. The trees of the parks were, in all probability, wholly indigenous, and were left to propagate themselves, by shedding their seeds among rough herbage; and the extent of surface they covered was allowed to be curtailed by deer and other animals, or to extend itself, according to the abundance or scarcity of pasture.

Of the foreign trees and shrubs of Scotland and Ireland, at this remote period, scarcely any thing is known. James I. is said to have been an amateur of the fine arts, and to have been fond of gardens, and of grafting fruit trees. James III. had gardens in the neighbourhood of Stirling Castle? and the pear trees and chestnuts, which are known to have existed in Scotland at that period, may have been introduced from France, with which country Scotland was then, and for many years afterwards, on intimate terms, or by the Roman clergy. Dr. Walker mentions a sweet chestnut at Finhaven in Forfarshire, which, in 1760, was conjectured to be upwards of 500 years old, and which is supposed to have been the oldest planted tree in Scotland. (*Essays*, p. 29.)

Still less is known of the introduction of foreign trees and shrubs into Ireland. The arbutus is thought by some to be indigenous; and it is certain that in England, in the 15th century, it was called the Irish arbutus. By others, however, it is said to have been introduced into Killarney by the monks of St. Finnian, who founded the abbey of that name on the banks of the lake, in the 6th century.

SUBJECT. 2. *Of the Foreign Trees and Shrubs introduced into Britain in the 16th Century.*

THERE is no record which throws any light on the subject of the introduction of foreign trees into England previously to the time of Henry VIII. Fitzherbert, in 1523, wrote on planting and preserving trees for timber and fuel; and Gooze, who translated Heresbachius in 1578, notices the same subjects. In Turner's *Names of Herbes* in 1548, the trees mentioned are, the almond, the apricot, the pomegranate, *Cistus salviaefolius*, rosemary, thyme, white jasmine, *Spártium júnceum*, the fig, the oriental plane, the elm, the sweet bay, the common black mulberry, the stone pine, the spruce fir, the *Cuprèssus sempervirens*, and the savin. In his *Herbal* of 1562, he adds the peach, the walnut, and the rue. In 1568 he adds the lavender. It appears that foreign trees and shrubs were not altogether neglected in the royal gardens, in the time of Henry VIII.; since, in a survey of the royal palace at Nonsuch, in Surrey, in the succeeding century, there were, in the wilderness, lilacs, lime trees, yews, junipers, and hollies. L'Obel, who published his *Adversaria* in 1570, includes the *Jasminum fruticans*, the *Pistàcia officinàrum*, and the *Genísta Scórpius*, among his woody plants. Tusser, in 1573, mentions the quince and the Damask rose. Grindal, Bishop of London, is said by Fuller to have introduced the German tamarisk, about the year 1560; but, according to Camden and Hakluyt, better authorities, about 1582. Grindal was visited at Fulham by the queen, who complained that the bishop had so surrounded his house with trees, that she could not enjoy the prospect from her chamber windows. Such excellent grapes were produced at Fulham by this prelate, that some were sent every year to the queen. (*Strype's Life of Grindal.*)

Wimbledon House, which was rebuilt by Sir Thomas Cecil in 1588, and surveyed by order of the parliament in 1649, was celebrated for its gardens and trees. In the several gardens, which consisted of mazes, wildernesses, knots, alleys, &c., are mentioned a great variety of fruit trees, and some shrubs, particularly "a faire bay tree," valued at 1*l.*, and "one very faire tree, called the Irish arbutis, very lovely to look upon, and worth 1*l.* 10*s.*" (*Lysons*, i. 397.) Gerard, the first edition of whose *Catalogue* is dated 1596, appears to have had several foreign trees and shrubs in his garden in Holborn; and, among others, *althæa frutex*, the laburnum, the Judas tree, six different kinds of roses, the laurustinus, the *Diospýros Lôtus*, the white mulberry, the nettle tree, the pinaster, the arbor vitæ, the yucca, and several others, as may be seen by the list below.

Gerard mentions having planted *Phyllirea serrata* in the Earl of Essex's garden at Barn Elms. (*Herbal*, edit. 1597, p. 1210.) Gough (*Brit. Topog.*, p. 61,) says, that, before the year 1597, Gerard had 1000 different plants and trees in cultivation. Tradescant is said by Gough to have been contemporary with Gerard, but he appears rather to belong to the 17th century. The only nursery which we read of as existing in the 16th century is that of Corbet, otherwise called Poynter, the father of Bishop Corbet, at Twickenham, mentioned by Sir Hugh Plat and by Ben Jonson. Gerard says that "Richard Poynter was a most cunning and curious grafter and planter of all manner of rare plants at Twickenham." (*Herb.*, 1597, p. 1269.)

It is uncertain whether Raleigh brought over any hardy American trees or shrubs, though it is highly probable that he did so, as he introduced the cherry tree into Ireland, and his manor at Sherborne, in Dorsetshire, is said to have been magnificently embellished with woods and gardens. Coker, author of a *Survey of Dorsetshire*, published in 1732, but which appears to have been written in the time of James I., says that Sir Walter Raleigh built in "the parke" adjoining the old castle "a most fine house, which hee beautified with orchardes, gardens, and groves of much varietic and great delight; soe, that whether that you consider the pleasantness of the seate, the goodnesse of the soyle, or the other delicacies belonging unto it, it rests unparalleled by anie in those parts." (p. 124.)

The park of Sherborne, after the death of Sir Walter Raleigh, came into the possession of the Earls of Digby, one of whom altered the house, and employed Brown to lay out the grounds. The centre part of the former mansion, which was built by Sir Walter Raleigh, still exists, and bears his arms, and the date 1574 over the windows. In the park there is a grove, said to have been planted by Sir Walter, which still retains his name. (*Beauties of England, &c., Dorsetshire*, p. 438.)

We can state nothing respecting the introduction of foreign trees into Scotland or Ireland during this century.

The trees and shrubs introduced into England during the 16th century, and the persons by whom they were introduced, cultivated, or recorded (the names of the latter being included in parentheses), according to the *Hortus Kewensis*, are as follows:—

1548.	<i>Laurus nóbilis</i>	Italy	(Turner)
	<i>Spártium júnceum</i>	S. of Eu.	Lord Cobham
	<i>Amýgdalus commúnis</i>	Barbary	(Turner)
	<i>Púnica Granátum</i>	S. of Eu.	Syon Garden
	<i>Armeniáca vulgáris</i>	Levant	(Turner)
	<i>Jasmínium officinále</i>	East Indies	(Turner)
	<i>Rosmarínus officinális</i>	South of Europe	(Turner)

1548.	<i>Thymus vulgaris</i>	South of Europe	(Turner)
	<i>Hyssopus officinalis</i>	South of Europe	(Turner)
	<i>Artemisia Abrotanum</i>	South of Europe	(Turner)
1548 or before.	<i>Morus nigra</i>	Italy	(Turner)
	<i>Ficus Carica</i>	South of Europe	(Turner)
	<i>Platanus orientalis</i>	Levant	(Turner)
1548 or before.	<i>Pinus Pinea</i>	S. of Europe	Richmond
	<i>Abies excelsa</i>	North of Europe	(Turner)
	<i>Juniperus Sabina</i>	South of Europe	(Turner)
	<i>Cupressus sempervirens</i>	Candia	Syon Garden
1551.	<i>Cistus salviæfolius</i>	S. of Eu.	Syon Garden
1562.	<i>Ruta graveolens</i>	South of Europe	(Turner)
	<i>Persica vulgaris</i>	Persia	(Turner)
	<i>Persica lævis</i>	Persia	(Turner)
	<i>Saturèja montana</i>	South of Europe	(Turner)
	<i>Juglans regia</i>	Persia	(Turner)
	<i>Juniperus tamariscifolia</i>	South of Europe	(Turner)
1568.	<i>Colutea arborescens</i>	France	(Turner)
	<i>Lavandula Spica and latifolia</i>	South of Europe	(Turner)
1569.	<i>Clématis Viticella</i>	Spain	Hugh Morgan
1570.	<i>Pistacia officinarum</i>	Levant	Gray
	<i>Genista sagittalis</i>	Germany	(Turner)
	<i>Genista Scorpius</i>	South of Europe	(Turner)
	<i>Ononis rotundifolia</i>	Switzerland	Hugh Morgan
	<i>Santolina squarrosa</i>	S. of Eu.	Hugh Morgan
	<i>Jasminum fruticans</i>	South of Europe	(Turner)
	<i>Vitex Agnus castus</i>	Sicily	(Turner)
	<i>Ephedra distachya</i>	France	L'Obel
1573.	<i>Rosa damascena</i>	Levant	(Tusser)
	<i>Cydonia vulgaris</i>	Austria	(Tusser)
	<i>Santolina Chamæcyparissus</i>	South of Europe	(Tusser)
1581.	<i>Quercus Ilex</i>	S. of France	Whitehall Gar.
1582.	<i>Tamarix germanica</i>	Germany	Archbp. Grindal
1596.	<i>Clématis pedicellata</i>	Majorca	Gerard
	<i>Clématis cirrhosa</i>	Spain	Gerard
	<i>Clématis Flammula</i>	France	Gerard
	<i>Cistus incanus</i>	South of Europe	Gerard
	<i>Hibiscus syriacus</i>	Syria	Gerard
	<i>Cytisus Laburnum</i>	Continent of Eu.	Gerard
	<i>Cytisus alpinus</i>	Continent of Eu.	Gerard
	<i>Cytisus spinosus</i>	South of Europe	Gerard
	<i>Coronilla Emericus</i>	France	Gerard
	<i>Medicago arborea</i>	Italy	Gerard
	<i>Cercis Siliquastrum</i>	South of Europe	Gerard
	<i>Rosa centifolia</i>	South of Europe	Gerard

1596.	<i>Ròsa lùtea</i>	Germany	Gerard
	<i>Ròsa moschàta</i>	Barbary	Gerard
	<i>Ròsa cinnamòmca</i>	France	Gerard
	<i>Ròsa provinciàlis</i>	France	Gerard
	<i>Ròsa gállica</i>	France	Gerard
	<i>Amelànchier vulgàris</i>	South of Europe	Gerard
	<i>Plantàgo Cynops</i>	South of Europe	Gerard
	<i>Paliùrus aculeàtus</i>	South of Europe	Gerard
	<i>Rhùs Coriària</i>	South of Europe	Gerard
	<i>Lonicera alpígena</i>	Switzerland	Gerard
	<i>Còrnus mäs</i>	Austria	Gerard
	<i>Philadélphus coronàrius</i>	South of Europe	Gerard
	<i>Tendòria fruticòsa</i>	South of Europe	(Miller)
	<i>Sambùcus racemòsa</i>	South of Europe	Gerard
	<i>Vibùrnum Tinus</i>	South of Europe	Gerard
	<i>Vibùrnum T. lùcida</i>	Spain	Gerard
	<i>Viburnum T. strìcta</i>	South of Europe	Gerard
	<i>Artemisia Santònica</i>	Siberia	Gerard
	<i>Diospÿros Lòtus</i>	Italy	Gerard
	<i>Sàlvia tríloba</i>	South of Europe	Gerard
	<i>Phlòmis fruticòsa and lanàta</i>	Spain	Gerard
	<i>Saturèja capitàta</i>	Levant	Gerard
	<i>Mòrus álba</i>	China	Gerard
	<i>Céltis austràlis</i>	South of Europe	Gerard
	<i>Pìnus Pinàster</i>	South of Europe	Gerard
	<i>Thùja occidentàlis</i>	North America	Gerard
	<i>Yucca gloriòsa</i>	North America	Gerard
	<i>Rúscus hypoglòssum</i>	Italy	Gerard
1597.	<i>Fròsa álba</i>	Crimea	Gerard
	<i>Cérasus Chamæcérasus</i>	Austria	Gerard
	<i>Lonicera nìgra</i>	Switzerland	Gerard
	<i>Syrínga vulgàris</i>	Persia, or probably Hungary, of which country it has been lately discovered to be also a native (<i>Bot. Mag.</i> , 3278., and <i>Gard. Mag.</i> , ix. 706.)	Gerard.
	<i>Phillyrea angustifòlia</i> , and the varieties, <i>mèdia</i> , <i>virgàta</i> , <i>péndula</i> , <i>oleafòlia</i> , <i>ligustrifòlia</i> , <i>lævis</i> , <i>ilicifòlia</i> , <i>latifòlia</i> , and <i>obliqua</i>	S. of Eu.	Earl of Essex
	<i>Periploca græca</i>	Syria	Gerard
	<i>Sàlvia officinàlis</i>	South of Europe	Gerard
	<i>Stÿrax officinàle</i>	Italy	Gerard
	<i>Dàphne Gnidium</i>	Spain	Gerard

It will be observed, from the foregoing list, that the date of the first introduction, or rather, that of the first mention made in books, of foreign woody plants in England, is 1548, when

sixteen were introduced. Among these were the sweet bay, the almond, the apricot, the pomegranate, the mulberry, the platanus, the stone pine, the common spruce fir, the cypress, and the savin juniper. The names of the introducers, or first cultivators, are almost entirely unknown, and, indeed, it is probable that most of the plants named at this early period had been in the country many years previously; some of them, as the rosemary, the thyme, the southernwood, the sweet bay, the apricot, &c., possibly from the time of the Romans; or, at all events, from the period of the establishment of religious houses in England. Among these plants, there are only two from ultra-European countries: the almond, from Barbary; and the jasmine, from the East Indies.

From 1551 to 1596, during the reign of Mary and the greater part of that of Elizabeth, twenty-four plants were first recorded, among which were the peach, the nectarine, and the walnut, from Persia; and the damask rose, the quince, and the *Quercus Ilex*. The names of the introducers are not known, with few exceptions; such as that of Hugh Morgan, apothecary to Queen Elizabeth; Gray, a London apothecary, mentioned by L'Obel; L'Obel, a Fleming, who was afterwards botanist to James I.; and Dr. Grindal, who was bishop of London, and afterwards archbishop of York and Canterbury, during the greater part of the reign of Elizabeth. From 1596 to the end of the century, forty-six different species were introduced, and upwards of thirty of these were first recorded by Gerard. Among these were, the English and Scotch laburnums, the *althæa frutex*, the Judas tree; the musk, the yellow and the hundred-leaved roses; the cotoneaster, Christ's thorn, *Córnus más*, the common syringa, the *laurustinus*, the lilac, and the *phillyrea*. Most of these are from the continent and south of Europe; and there are, in this period, also, the *arbor vitæ* and the *yucca*, from North America.

Thus, the total number of foreign woody plants which are known to have been cultivated in Britain during the 16th century is only eighty-four, exclusive of two varieties of the *laurustinus*, and nine of the *phillyrea*.

It is impossible, at this distance of time, to ascertain the names of all the persons to whom we are indebted for the introduction of these plants; but it is certain that the merit of the first cultivation of the greater part of them belongs decidedly to Gerard.

John Gerard, Pulteney informs us, was born at Nantwich in Cheshire, in 1545, educated as a surgeon, and patronised in London by Lord Burleigh, who had at that time the best collection of plants in the kingdom. Gerard superintended this nobleman's garden, which was in the Straud; Gerard himself living in Holborn, where he had a physic garden, considered

to be at that time the most remarkable in England for the number and variety of its productions. This garden appears by the old maps to have been situated on the brow of the hill between what is now Ely Place, and what was formerly the Fleet River, but what is now called Field Lane, the stream being arched over. Gerard appears to have practised as a surgeon and apothecary, supplying his prescriptions from his garden. He was the author of several works, the principal of which are his *Catalogue* and his *Herbal*. The first edition of the former is dedicated to Lord Burleigh, and the second to Sir Walter Raleigh. It enumerates nearly 1100 sorts of plants, of foreign and domestic growth, all of which (as attested by L'Obel) were to be found in his garden in Holborn. Gerard died about the year 1607, highly respected by the college of physicians and by all his contemporaries.

SUBJECT. 3. *Of the Foreign Trees and Shrubs introduced into Britain in the 17th Century.*

TRADESCANT appears to have come to England towards the end of the preceding century. Wood says he was a Dutchman; that he was in the service of Lord Treasurer Salisbury, Lord Wootton, and the Duke of Buckingham; and that, about 1629, he obtained the title of gardener to Charles I. He is said to have travelled over a great part of Europe, and to have gone into Barbary, Greece, Egypt, and other Eastern countries, in quest of plants and natural curiosities. He had a garden at Lambeth, and a museum there; in the former of which he cultivated many plants, and, as appears by a *Catalogue* published by his son, in 1656, some trees and shrubs. Tradescant's garden and museum were probably not commenced till after he had retired from the service of private noblemen, and entered into that of the king, which would give its origin about 1630. Tradescant's son travelled in Virginia, and introduced various new plants from that country. Tradescant, senior, died about 1652. Tradescant's garden was visited, in 1749, by Dr. Mitchell and Dr. (afterwards Sir) William Watson, F.R.S.; but at that distant period they found very few trees. Among these, however, were *Schubertia disticha*, *Robinia Pseud-Acacia*; *Rhamnus catharticus*, about 20 ft. high, and nearly a foot in diameter; an *Aristolochia*, and several mulberry trees. (*Phil. Trans. Abr*, x. 740.) These were but a few of the species of trees cultivated by Tradescant; as appears by the *Catalogue* published by his son, and by the list at the end of this section.

From a memorandum by Dr. Gray, in his copy of the *Horti Regii Hamptonensis*, &c., now in the British Museum, we learn that many of the plants enumerated in that catalogue were

brought from Soesdyke in Holland, the seat of Mr. Bentinck, afterwards Earl of Portland. The gardens of Holland were at that time the richest in Europe.

The great introducer of foreign trees in this century was Dr. Compton, who was the bishop of London from 1675 to 1713, and who may truly be said to have been the father of all that has since been done in this branch of rural improvement. Bishop Compton was the youngest son of Spencer, Earl of Northampton; he was made Bishop of Oxford in 1674, and was translated to the see of London in the following year. He was a zealous protestant and a most excellent man. He lived a retired life at Fulham, attending to his episcopal duties and to his garden.

In the 32d book of Ray's *Historia Plantarum*, written in 1686, in which he treats of plants imperfectly known, there is a chapter on the rare trees and shrubs which he saw in the garden of Bishop Compton at Fulham. Among these are enumerated the tulip tree, the magnolia, the sassafras, the tree angelica (*Aràlia spinòsa*), the hickory, the box elder, the liquidambar, the Constantinople nut, some species of *Crataëgus*, some of *Rhùs*, some of *Córnus*, and some of *A'triplex*. Bishop Compton died in 1713 at the age of 81 years. His garden was visited by Sir William Watson in 1751, 48 years after his death; and he gave the following account of this bishop and his garden to the Royal Society:—"Dr. Henry Compton," he observes, "planted a greater variety of curious exotic plants and trees, than had at that time been collected in any garden in England. This excellent prelate presided over the see of London from the year 1675 to 1713; during which time, by means of a large correspondence with the principal botanists of Europe and America, he introduced into England a great number of plants, but more especially trees, which had never been seen here before, and described by no author; and in the cultivation of these (as we are informed by the late most ingenious Mr. Ray) he agreeably spent such part of his time as could most conveniently be spared from his other more arduous occupations. From this prelate's goodness, in permitting, with freedom, persons curious in botany to visit his garden, and see therein what was to be found nowhere else; and from his zeal in propagating botanical knowledge, by readily communicating to others, as well to foreigners as to our own countrymen, such plants and seeds as he was in possession of, his name is mentioned with the greatest encomiums by the botanical writers of his time; viz., by Hermann, Ray, Plukenet, and others. As this prelate's length of life and continuance in the see of London were remarkable, so we find the botanists, who wrote after Mr. Ray, most frequently mentioning in their works the new accessions of treasures to this

garden; and of this you meet with a great variety of examples in the treatises of Dr. Plukenet, Hermann, and Commelyn. Botanical much more even than other worldly affairs are subject to great fluctuations, and this arises not only from the natural decay of vegetables, and their being injured by the variety of seasons, but also from the genius and disposition of the possessors of them. So, here, upon the death of Bishop Compton, all the green-house plants and more tender exotic trees were, as I am informed by Sir Hans Sloane, given to the ancestor of the present Earl Tylney at Wanstead. And as the successors of this bishop in the see of London were more distinguished for their piety and learning than for their zeal in the promotion of natural knowledge, the curiosities of this garden were not attended to, but left to the management of ignorant persons; so that many of the hardy exotic trees, however valuable, were removed to make way for the more ordinary productions of the kitchen-garden." (*Phil. Trans.*, xlvii. 243.)

Collinson, speaking of Bishop Robinson, Dr. Compton's successor, says, he was a man of "no such taste" as Bishop Compton. "He allowed his gardener to sell what he pleased, and often spoiled what he could not otherwise dispose of. Many fine trees, come to great maturity, were cut down, to make room for produce for the table. Furber of Kensington, and Gray of Fulham, augmented their collections from this source, with plants not otherwise to be procured."

The following are the principal trees and shrubs which Sir William Watson found in the bishop's garden in 1751:—

Acerinææ. *Acer rubrum*, *platanoides*; *Negundo fraxinifolium*.

Hippocastanææ. *Pavia rubra*.

Terebinthaceæ. *Pistacia officinarum*, *Rhus typhina*.

Leguminosææ. *Robinia Pseud-Acacia*, *Gleditschia triacanthos*, *Cytisus alpinus*, *Cercis Siliquastrum*.

Amygdalææ. *Cerasus Laurocerasus*.

Pomacææ. *Mespilus prunifolia*?

Ericacææ. *Arbutus Uredo*.

Ebenacææ. *Diospyros virginiana*.

Oleacææ. *Ornus europæa*, *rotundifolia*; *Syringa persica* var. *laciniata*.

Laurinææ. *Laurus Benzoin*.

Ulmacææ. *Celtis*.

Juglandææ. *Juglans nigra*.

Cupuliferææ. *Quercus Sæber*, *Ilex*, *alba*; *Corylus rostrata*?

Coniferææ. *Cedrus Libani*, *Larix europæa*; *Pinus Pinea*, *Pinaster*; *Abies Picea*; *Cupressus*, the male cypress, the female cypress; *Juniperus virginiana*.

Smilacææ. *Ruscus hypoglössum*, *racemosus*.

These articles belong to 15 orders, or natural groups, and include 34 trees and shrubs.

A survey of the old trees at Fulham Palace was made by Lysons in 1793, and again in 1809, and published in Lysons's *Environs of London*; by which it appears that several of the trees mentioned by Sir William Watson were still in existence, and in a growing state. The girts of the following trees, taken at these two different periods, are here given from Lysons, as taken at 3 ft. from the ground, to which we have added the dimensions of such as are now (January, 1835) still in existence, which we are enabled to do through the kindness of Dr. Blomfield, the present bishop. We saw the trees ourselves in October last, and found most of those below mentioned still in a growing state, with some robinias and others in a state of venerable decay.

	Girt in	Girt in	Girt in	Computed	
	1793.	1809.	1835.	Height in 1793 & 1835	
	ft. in.	ft. in.	ft. in.	ft.	in.
<i>Negundo fraxifolium</i> , or ash-leaved maple, planted in 1688	6 4	7 1½	—	45	
<i>Acer rubrum</i> , scarlet-flowered maple	4 3	—	—	40	
<i>Juglans nigra</i> , black walnut tree	11 2	11 5½	14 6	70	
<i>Quercus alba</i> , white oak	7 11	8 1½	11 5	70	
<i>Quercus Ilex</i> , evergreen oak	8 0	9 1	10 3	50	
<i>Quercus Süber</i> , cork tree	—	8 4	9 5	50	
<i>Cupressus sempervirens</i> , upright cypress	2 3	—	—	30	
<i>Juniperus virginiana</i> , Virginian red cedar	2 5	—	—	20	
<i>Pinus Pinaster</i> , cluster pine	10 0	10 1	11 10	80	

“There were also,” says Mr. Lysons, in 1793, “the *Quercus Süber*, the *Cytisus Laburnum*, the *Robinia Pseud-Acacia*, and the *Pinus Cædrus*, mentioned by Sir William Watson. The cedar of Lebanon was first planted at Fulham in 1683; the largest, of two measured in 1793, was only 7 feet 9 inches in girt.” “Near the porter’s lodge,” he continues, “are some limes of great age, one of which measured, in 1793, 13 feet 3 inches in girt. It is most probable that they were planted by Bishop Compton about the year of the Revolution (1688), when the fashion of planting avenues of limes was introduced into this country from Holland, where they ornamented the Prince of Orange’s palaces.”

“Upon visiting the gardens at Fulham again in 1809,” Lysons observes, “I could not find the *Cupressus sempervirens*, the *Juniperus virginiana*, or the *Acer rubrum*. The following trees still remain, and they will no doubt be regarded with veneration by the botanist, as the parent stocks of their respective races in the kingdom. The *Acer Negundo*, the girt of which, at three feet from the ground, is now

(1809) 7 ft. $1\frac{1}{2}$ in.; the *Juglans nigra*, 11 ft. $5\frac{1}{2}$ in.; the *Pinus Pinaster*, 10 ft. 1 in.; the *Quercus Ilex*, 9 ft. 1 in.; the *Quercus alba*, 8 ft. $1\frac{1}{2}$ in.; the *Quercus Süber*, of which I had not a satisfactory measure in 1793, is now (1809) 8 ft. 4 in. in girt; the largest cedar now measures 8 ft. $8\frac{3}{4}$ in. in girt; another, in a court of the palace, about 7 ft.: it is probable that the latter has been lessened in girt, from having been drawn up by its situation to a remarkable height. The lime tree above mentioned now measures 14 ft. 1 in. in girt. The *Cytisus Laburnum* is an old decayed tree in the close (without the lodge) near the moat, about 3 ft. in girt. There are two of the *Robinia Pseud-Acacia*, one near the porter's lodge, and one on the lawn near the moat; they are both in a state of great decay, and their trunks in such a state as not to admit of measurement."

All the trees mentioned in the above extract, except those contained in the table, the large limes, the remains of the robinia, and one or two others, are decayed or taken down; the grounds having undergone several alterations during the occupancy of Bishop Porteus, between 1800 and 1816. Both Bishop Porteus and the present bishop have added considerably to the collection.

It would be interesting to know the means by which Bishop Compton procured his trees and shrubs from America, and who were the botanical collectors of that day. Several may have existed whose names are now lost. It appears highly probable that most of the American trees and plants at Fulham were introduced by the Rev. John Banister, who was sent by the bishop as a missionary to Virginia. John Banister, according to Dr. Pulteney (*Sketches, &c.*, vol. i.), was one of the first British collectors in North America. He published a *Catalogue* of the plants he observed there, dated 1680. He is mentioned repeatedly by Ray, as having introduced many plants. Banister was one of the early martyrs to natural history, having, in one of his excursions, fallen from a rock and perished. His *Catalogue* will be found in the second volume of Ray's *Historia Plantarum*, and several of his papers are published in the *Philosophical Transactions*. Plukenet, describing the *Azalea viscosa*, says that a drawing of it, by his own hand, was sent by him to Bishop Compton, his patron.

The name of Evelyn is well known, as belonging to this century. His *Sylva* was published in 1664, from which, and from his *Calendarium Hortense*, it appears that the number of species and varieties of trees and shrubs in the London gardens was then extremely limited. In one of the later editions of the *Sylva*, Evelyn mentions the tulip tree as having been introduced by Tradescant. His description of the tree is curious. He says, "they have a poplar in Virginia of a very peculiar-shaped leaf,

as if the point of it were cut off, which grows very well with the curious amongst us to a considerable stature. I conceive it was first brought over by John Tradescant, under the name of the tulip tree (from the likeness of its flowers,) but is not, that I find, taken notice of in any of our herbals. I wish we had more of them." (*Sylva*, edit. 1670.) The tulip tree was at that time known through all the English settlements by the title of poplar. (*Hunter's Evelyn*, i. 207.) Hermann says that he observed in the park of the Duke of Norfolk, five or six miles [Dutch miles] from London [? Deepdene], a tulip tree which had been planted there twenty years before, but which had never flowered or borne fruit. (*Hort. Acad. Lugd. Bat. Cat.* 1687, p. 615.) At Say's Court, Deptford, one of Evelyn's residences, he is said to have had a variety of trees; but Gibson, who visited it in 1691, after Evelyn had left it, found only the Phillyrea and the holly: of the former, Evelyn had four large round and smoothly clipped plants, on naked stems; and of the latter, a hedge, 400 ft. long, 9 ft. high, and 5 ft. in diameter. Evelyn was very proud of this hedge, and mentions it more than once in his writings. It was ruined by Peter the Great, who, having taken the house at Say's Court, to be near the Deptford Dockyards, had himself wheeled through this hedge in a wheelbarrow for amusement! Evelyn planted cedars, pines, silver firs, ilexes, and walnuts at Wooton, some of which we found still remaining there in 1830. Evelyn however, was more anxious to promote the planting of valuable indigenous trees, than to introduce foreign ones.

Gibson, who made a tour through the gardens about London in 1691, which was published from his M.S. many years afterwards in the *Archæologia*, tells us that he found Sir William Temple's garden, at West Sheen, to excel in orange trees and other "greens," as evergreen shrubs were called at that time; Among these "greens" Italian bays, laurustinuses, and striped hollies were included. Sir Henry Capell is said to have had as "curious greens, in his garden at Kew, as any about London." His two lentiscus trees (*Pistácia Lentiscus*) for which he paid 40*l.* to Verspritt, were said to be the best in England. He had four white-striped hollies, about $\frac{1}{2}$ feet above their cases, kept "round and regular," which cost him 5*l.* a tree; and six laurustinuses, with "large, round, equal heads, very flowery and showy." "In the garden of Sir Stephen Fox, at Chiswick which, though only of five years' standing, is brought to great perfection for the time), are two myrtle hedges about 3 ft. high. They are protected in winter with cases of boards painted." Sir Josiah Child's plantations of walnuts and other trees, at Wanstead, are said by Gibson to be "much more worth seeing than his gardens, which are but indifferent." "Captain Foster's

garden at Lambeth," Gibson observes, "has many curiosities in it, and perhaps the finest striped holly hedge in England. He has many myrtles, not the greatest, but cut in the most fanciful shapes that are anywhere to be seen. He has a walk arched over with trelliswork, and covered with vines, which with others running on most of his walls, without prejudice to his lower trees, yield him a deal of wine."

The commercial gardeners at this time (1691) are thus enumerated by Gibson:—London and Wise had the only extensive nursery; Versprit excelled in hollies and "greens." Ricketts and Pearson were small cultivators for sale. The latter had "abundance of cypresses, which, at 3 ft. high, he sold for 4d. apiece; and, being moderate in his prices, and very honest in his dealings, he got much chapmanry." Darby, at Hoxton, is said "to be master of several curious greens that other sale gardens want." Darby is said to have raised many striped hollies by inoculation; and Captain Foster (who appears also to have sold or exchanged his garden productions) to have propagated the same plants by grafting. Darby also kept a book of dried specimens of plants, to show to his customers. Clements, at Mile End, had many curious "greens," and, the year that Gibson visited him (1691), made "white muscadine, and white Frontignac wine," better than any he (Gibson) had elsewhere tasted. It is worthy of remark, that all these "sale gardeners" had greenhouses, and that they piqued themselves principally upon their plants in pots and on their florists' flowers. It is singular that Gibson does not speak of the Bishop of London's garden, though it must have been in its state of greatest perfection at the time he wrote; and also that he barely mentions the nursery of Messrs. London and Wise, which Evelyn informs us, in the preface to his translation of Quintinye's *Complete Gardener*, published in 1701, "far surpassed all the others in England put together."

The Brompton Park Nursery may, indeed, be considered as the first establishment of the kind which became celebrated. It was founded by Messrs. Cooke, Luere, London and Field, in 1681. Luere, or Lukar, was gardener to the Queen Dowager at Somerset House; Field was gardener to the Earl of Bedford, at Bedford House in the Strand; Moses Cook was gardener to the Earl of Essex, at Cashibury, and author of a work entitled *The Manner of raising Forest Trees, &c.*, 4to, 1676. George London was gardener to Bishop Compton, and afterwards chief gardener, first to William and Mary, and afterwards to Queen Anne. Lukar died in 1686; Cooke and Co. succeeded. Cooke retired in 1689, when Henry Wise, who had been an apprentice to Rose, the royal gardener, as London had also been, became the sole proprietor. In 1593-4, he entered

into a new partnership with London. At that time the grounds exceeded 100 acres in extent. This nursery passed, successively, from London and Wise, in 1701, to Swinhoe; in 1714, to Smith and Co.; in 1756, to Jeffries; in 1788, to Jeffries and Gray; at the death of Jeffries, to Gray and Wear; afterwards to Gray, Wear, and Co.; then to Gray, Son, and Brown; and, lastly, to Gray and Son, in whose occupation it still (1835) is. The grounds are now reduced to thirty acres. In the time of London and Wise, it was thus spoken of by Evelyn, in the preface before alluded to:—“The proprietors, Mr. George London, chief gardener to their majesties, and his associate, Mr. Henry Wise, are recommended for their assiduity and industry; they have not made gain the only mark of their pains, but with extraordinary and rare industry endeavoured to improve themselves in the mysteries of their profession; from the great advantages and now long experience they have had, in being employed in most of the celebrated gardens and plantations which this nation abounds in, besides what they have learned abroad, where horticulture is in high reputation.” He adds, “the grounds and gardens of noblemen and persons of quality, which they have planted *ab origine*, and which are still under their care and attention, justify what I have said in their behalf.” Bowack, who wrote an account of the parish of Kensington in 1705, says, “that some affirm that if the stock of these nurseries were valued at one penny per plant, the amount would exceed 40,000*l.*” London and Wise, in 1694, employed twenty men in their nursery and two women. The foreman had 12*s.* a week, the other men had 8*s.*, and the women 4*s.*

Botanic gardens began to be established in England about the middle of this century; and they contributed to the introduction of hardy trees and shrubs, as well as of herbaceous plants and exotics. The oldest botanic gardens in England are those of Oxford and of Chelsea. Evelyn visited the latter in 1685, and mentions, as rarities, a tulip tree and a tea shrub. Many private botanic gardens were also founded during this century, Among these were the gardens of Ray, in Essex; of the Duchess of Beaufort, at Badmington, in Gloucestershire; of Sir Hans Sloane, at Chelsea; of Dr. Uvedale, at Enfield, &c. The catalogues of these gardens, in the libraries of the Linnæan Society and of the British Museum, show that they contained various foreign trees and shrubs. Dr. Uvedale’s garden, Gibson informs us, “chiefly excelled in exotic greens and orange trees, for which he had six or seven houses or roomsteads.” Gibson adds, “that he understood the culture of particular plants, but had no taste for the disposition of his garden.” We learn from Miller, that Dr. Uvedale had a fine cedar tree, which,

in 1788, was 45 ft. 9 in. high, though 9 ft. had been broken off by the wind. Lysons saw this tree in 1809, and found the girt of it, at 3 ft. 10 in. from the ground (not being able to measure lower, on account of a seat which was fixed round it), to be 13 ft. 1 in. Dr. Uvedale was born in 1642; he became master of the grammar school at Enfield about 1670, and died in 1722. He is said to have devoted so much of his time to his garden, as to be threatened with being removed from his situation by the authorities who had appointed him.

Dr. May, the present master of the grammar school at Enfield, says there is a tradition that one of Dr. Uvedale's scholars, who travelled, had a commission from the doctor to bring a plant of the cedar of Lebanon from Mount Lebanon, and that he brought the tree now standing. Dr. May had it measured in 1821, for the *History of Enfield*; and, the tree being in a state of decay, its dimensions at the present time (January, 1835) are much the same as they were then. The tree lost one of its leading branches in November 1794, previously to which its general form was that of an inverted cone. It was then, and is now, 64 ft. 8 in. high; the girt at one foot from the ground, in 1821, was 19 ft. 9 in.; and the girt is now (1835) 15 ft. 8 in., at 3 ft. from the ground; at 6 ft., 14 ft. There is a portrait of the Enfield cedar in Strutt's *Sylva Britannica*, and the measurements, as taken for us, with the kind permission of Dr. May, will be found in detail in the *Gardener's Magazine*, vol. xi.

The trees and shrubs introduced or cultivated by the curators or proprietors of these different gardens, and others which we have mentioned, will be found in the list which concludes this section, in which the names of Dr. Compton, Gerard, L'Obel, Parkinson, Tradescant, Sutherland, Uvedale, and Sir Hans Sloane, will be found frequently to occur.

In Scotland there appears to have been some taste for botany towards the end of this century, as Patrick Murray had a collection of a thousand plants at Livingstone, and Dr. Balfour founded the botanic garden of Edinburgh in 1680. The curator of the botanic garden at Edinburgh, James Sutherland, was an excellent botanist, and by his correspondents introduced many foreign plants into the garden. It is remarkable that in this garden the cedar of Lebanon was introduced in 1683, the same year in which it is mentioned as having been planted by Bishop Compton at Fulham, and in the Chelsea Botanic Garden.

In Ireland, Sir Arthur Rawdon, struck with the collection of plants in the garden of his countryman, Dr. (afterwards Sir) Hans Sloane, of Chelsea, sent a gardener, who had been a collector for Sir Hans Sloane, to Jamaica, who brought back a shipload of plants to Moira, where various hardy foreign trees were introduced, and kept in good order for several years.

The place is now in the possession of Sir Robert Bateson, but we believe it has been long since dismantled.

The trees and shrubs introduced into England in the 17th century, according to the *Hortus Kewensis*, were as follows:—

1603.	<i>Abies Picea</i>	Germany	Serjt. Newdigate
1616.	<i>Salvia grandiflora</i>	South of Europe	L'Obel
1629.	<i>Cistus ladaniferus</i>	Spain	(Parkinson)
	<i>Cistus cyprius</i>	Greece	(Parkinson)
	<i>Æsculus Hippocastanum</i>	Asia	(Parkinson)
	<i>Ampelopsis hederacea</i>	North America	(Parkinson)
	<i>Coriaria myrtifolia</i>	South of Eu.	Parkinson
	<i>Rhamnus Alaternus</i>	South of Eu.	(Parkinson)
	<i>Rhamnus Clusii</i>	South of Eu.	(Parkinson)
	<i>Rhus typhina</i>	North America	Parkinson
	<i>Cytisus sessilifolius</i>	Italy	Parkinson
	<i>Rosa turbinata</i>	Cont. of Eu.	(Parkinson)
	<i>Rosa sempervirens</i>	South of Eu.	(Parkinson)
	<i>Rosa sulphurea</i>	Levant	J. de Franqueville
	<i>Cerasus Laurocerasus</i>	Levant	James Cole
	<i>Cerasus serotina</i>	North America	(Parkinson)
	<i>Crataegus Pyracantha</i>	South of Eu.	(Parkinson)
	<i>Lonicera caerulea</i>	Switzerland	(Parkinson)
	<i>Helichrysum Stæchas</i>	Cont. of Eu.	(Parkinson)
	<i>Diospyros virginiana</i>	North America	(Parkinson)
	<i>Morus rubra</i>	North America	(Parkinson)
	<i>Juglans nigra</i>	North America	(Parkinson)
	<i>Carya alba</i>	North America	(Parkinson)
	<i>Larix europæa</i>	Germany	(Parkinson)
1633.	<i>Elæagnus angustifolia</i>	S. of Europe	Parkinson
	<i>Laurus Sassafras</i>	North America	Wilmot
1636.	<i>Coronilla juncea</i>	France	J. Tradescant, jun.
1640.	<i>Cistus villosus</i>	South of Eu.	(Parkinson)
	<i>Hypêricum hircinum</i>	South of Eu.	(Parkinson)
	<i>Staphylæa trifolia</i>	N. Amer.	J. Tradescant
	<i>Rhus Toxicodendron</i>	North America	(Parkinson)
	<i>Gelsemium sempervirens</i>	North America	Parkinson
	<i>Astragalus Tragacantha</i>	South of Eu.	(Parkinson)
	<i>Astragalus Poterium</i>	Levant	(Parkinson)
	<i>Cytisus triflorus</i>	Spain	(Parkinson)
	<i>Robinia Pseud-Acacia</i>	N. Amer.	J. Tradescant
	<i>Spiræa hypericifolia</i>	North America	(Parkinson)
	<i>Crataegus Azarolus</i>	S. of Eu.	J. Tradescant, jun.
	<i>Syringa persica</i>	Persia	J. Tradescant
	<i>Stæhelina dubia</i>	Continent of Europe	(Park.)
	<i>Dorýcnium réctum</i>	South of Eu.	(Parkinson)
	<i>Artemisia arboréscens</i>	Levant	(Parkinson)

1640.	<i>Plantago aſra</i>	Sicily	(Parkinson)
	<i>Técoma radicans</i>	North America	Parkinson
	<i>Zízüphus vulgàris</i>	South of Eu.	(Parkinson)
	<i>Dàphne Tartonràira</i>	France	(Parkinson)
	<i>A'triplex Hálimus</i>	Spain	(Parkinson)
	<i>Plátanus occidentàlis</i>	N. Amer. J. Tradescant, jun.	
	<i>Schubértia dísticha</i>	N. Amer. J. Tradescant, jun.	
	<i>Rúscus hypophýllum</i>	Italy	Parkinson
	<i>Cístus álbídis</i>	Spain	(Parkinson)
1648.	<i>Vítis laciniòsa</i>	Cont. of Eu.	Oxford Gar.
	<i>Cérasus lusitànica</i>	Portugal	Oxford Garden
	<i>Èrica mediterrànea</i>	Portugal	Oxford Garden
	<i>Smilax áspera</i>	S. of Eu.	Oxford Garden
1656.	<i>Cístus láxus</i>	Spain	John Tradescant
	<i>Cístus críspus</i>	Portugal	John Tradescant
	<i>Cístus populifòlius</i>	Spain	John Tradescant
	<i>Cístus hirsùtus</i>	Portugal	J. Tradescant, jun.
	<i>Cístus corboriènsis</i>	Spain	J. Tradescant, jun.
	<i>Cístus monspeliènsis</i>	Spain	John Tradescant
	<i>Àcer rùbrum</i>	N. Amer. J. Tradescant, jun.	
	<i>Vítis vulpìna</i>	N. Amer. J. Tradescant, jun.	
	<i>Vítis Labrúsca</i>	N. Amer. J. Tradescant, jun.	
	<i>Pistàcia Terebínthus</i>	S. of Eu.	J. Tradescant
	<i>Rhús Còtinus</i>	S. of Eu.	J. Tradescant
	<i>Caprifòlium sempervìrens</i>	N. Amer. J. Tradescant, jun.	
	<i>Céltis occidentàlis</i>	N. Amer.	J. Tradescant
	<i>Jùglans cinèrea</i>	N. Amer. J. Tradescant, jun.	
	<i>Rhododéndron hirsùtum</i>	Switzld.	J. Tradescant, jun.
	<i>Jasminum hùmile</i>	S. of Eu.	J. Tradescant, jun.
1658.	<i>Polýgala Chamæbúxus</i>	Austria	Oxford Garden
1661.	<i>Phlòmis purpùrea</i>	S. of Eu.	Edward Morgan
	<i>Phlòmis ítálica</i>	Italy	Edward Morgan
1663.	<i>Liriodéndron Tulipífera</i>	N. Amer.	Earl of Norfolk
	<i>Pistàcia Lentíscus</i>	S. of Europe	(Evelyn)
1664.	<i>Juníperus virginiana</i>	North America	Evelyn
	<i>Smilax Sarsaparilla</i>	North America	Evelyn
1665.	<i>Córylus Colúrna</i>	Constantinople	John Rea
1680.	<i>Onònis fruticòsa</i>	South of France	(Morrison)
1683.	<i>Àcer platanòides</i>	S. of Eu.	Jas. Sutherland
	<i>Euónymus americànus</i>	N. Amer.	Jas. Sutherland
	<i>Rhámnus infectòrius</i>	S. of Eu.	Jas. Sutherland
	<i>Ròsa alpina</i>	Switzerland	Jas. Sutherland
	<i>Amýgdalus pùmila</i>	China	(Ray)
	<i>Amýgdalus nàna</i>	Russia	Jas. Sutherland
	<i>Cratægus coccínea</i>	N. Amer.	Bishop Compton
	<i>Pýrus Chamæmèspilus</i>	Pyrenees	James Sutherland
	<i>Còrnus sericea</i>	N. Amer.	Bishop Compton

1683.	<i>Ribes reclinatum</i>	Germany	J. Sutherland
	<i>Baccharis halimifolia</i>	N. Amer.	Bishop Compton
	<i>Santolina rosmarinifolia</i>	S. of Eu.	Jas. Sutherland
	<i>Dorycnium hirsutum</i>	S. of Eu.	Jas. Sutherland
	<i>Lithospermum fruticosum</i>	S. of Eu.	Jas. Sutherland
	<i>Laurus Benzoin</i>	N. Amer.	Bishop Compton
	<i>Quercus coccifera</i>	France	Jas. Sutherland
	<i>Liquidambar styraciflua</i>	N. Amer.	Bishop Compton
	<i>Pinus halepensis</i>	Levant	Bishop Compton
	<i>Cedrus Libani</i>	Levant	Chelsea Garden
	<i>Juniperus phoenicea</i>	S. of Eu.	Jas. Sutherland
1688.	<i>Magnolia glauca</i>	N. Amer.	Bishop Compton
	<i>Magnolia longifolia</i>	N. Amer.	Bishop Compton
	<i>Hypericum serpyllifolium</i>	Levant	Hon. C. Howard
	<i>Negundo fraxinifolium</i>	N. Amer.	Bishop Compton
	<i>Rhus copallina</i>	N. Amer.	Bishop Compton
	<i>Aralia spinosa</i>	Virginia	Bishop Compton
1690.	<i>Spiræa opulifolia</i>	N. Amer.	Bishop Compton
1691.	<i>Menispermum canadense</i>	N. Amer.	Bishop Compton
	<i>Cratægus Crus-galli</i>	N. Amer.	Hon. C. Howard
	<i>Quercus coccinea</i>	N. Amer.	Bishop Compton
1692.	<i>Salix babylonica</i>	Levant	Royal Gardens, Hampton Court
	<i>Populus balsamifera</i>	N. Amer.	Royal Gardens, Hampton Court
	<i>Ostrya virginica</i>	N. Amer.	Bishop Compton
1693.	<i>Juniperus lycia</i>	S. of Eu.	Jacob Bobart
1696.	<i>Rubus occidentalis</i>	N. Amer.	Chelsea Garden
	<i>Lycium barbarum</i>	Barbary	Royal Gardens, St. James's
	<i>Abies balsamifera</i>	N. Amer.	Bishop Compton
1697.	<i>Ornus rotundifolia</i>	Italy	Duchess of Beaufort
1699.	<i>Passiflora cærulea</i>	Brazil	Duchess of Beaufort
	<i>Phyllis Nobla</i>	Canaries	Duch. of Beaufort
	<i>Salvia pomifera</i>	Candia	Hon. C. Howard
	<i>Castanea pumila</i>	N. Amer.	Duch. of Beaufort
	<i>Quercus Süber</i>	France	Duchess of Beaufort
	<i>Myrica cerifera</i>	N. Amer.	Duch. of Beaufort
1700.	<i>Ampelopsis bipinnata</i>	North America	
	<i>Gleditschia triacanthos</i>	N. Amer.	Bishop Compton
	<i>Abies alba</i>	N. Amer.	Bishop Compton
	<i>Abies nigra</i>	N. Amer.	Bishop Compton
	<i>Pyrus arbutifolia</i>	N. Amer.	Lord Clarendon
	<i>Pyrus melanocarpa</i>	N. Amer.	
	<i>Rubus odoratus</i>	N. Amer.	Sir Hans Sloane

The total number of woody plants introduced during the 17th century appears to be upwards of 130.

From the commencement of the seventeenth century to 1636, during the reign of James I., and part of that of Charles I., twenty-six plants were introduced; all, except four, in the year 1629. The reason why so many appear in this year is, that it is the date of the first edition of Parkinson's *Paradisi in Sole*, &c., in which they were first enumerated. The only introducers mentioned are, Mr. Serjeant Newdigate; John de Franqueville, a merchant in London, from whose care, Parkinson says, "is sprung the greatest store of rare plants that is now flourishing in this kingdom;" Wilmot, and Parkinson. Among the plants introduced during this period are some of considerable interest; the silver fir by Serjeant Newdigate, the gum cistus, the horsechestnut, the five-leaved ivy, the common laurel, the pyracantha, the red mulberry, the black walnut, and that most important tree, the larch: the introducers of the last eight valuable plants are unknown. Five of the articles are from North America, one from Asia, and the rest from different parts of the continent of Europe. Evelyn states that "at Harefield Park, in the county of Middlesex, belonging to Mr. Serjeant Newdigate, there are two Spanish or silver firs, that being planted there in the year 1603, at two years' growth from the seed, are now (1679) become goodly masts. The biggest of them, from the ground to the upper bough, is 81 ft., though forked on the top; which has not a little impeded its growth. The girt, or circumference, below, is 13 ft.; and the length, so far as it is timber, that is to 6 in. square, is 73 ft.; in the middle it is 17 in. square; amounting by calculation to 146 ft. of good timber. The other tree is indeed not altogether so large, by reason of its standing near the house when it was burned about 40 years since, when one side of the tree was scorched." (*Silva*, edit. 1706.)

In 1640 (still during the reign of Charles I.), twenty-three plants were introduced. The authority is Parkinson's *Herbal, or Theatre of Plants*, published in that year. The introducers were, Parkinson, Tradescant, and Tradescant junior. Among the articles were, the *Robinia Pseud-acacia*, the azarole, the Persian lilac, the occidental plane, and the deciduous cypress. Seven are from North America, and the rest from different parts of Europe.

In the year 1656 (in the time of Cromwell), sixteen plants were introduced, the authority for which is the *Catalogue of Tradescant's Museum*, published in that year. Among the articles are, *Acer rubrum*, the evergreen honeysuckle, the nettle tree, and the grey walnut.

From 1658 to 1683 (Charles II.), nine plants were introduced, by Edward Morgan, John Rea, Bishop Compton, Evelyn, and the Earl of Norfolk. Among these are, the *Pistacia Lentiscus*,

the red cedar, the Constantinople nut, and the tulip tree. Only three of these are from North America.

In 1683, twenty plants were introduced, by James Sutherland, first curator of the botanic garden of Edinburgh, Bishop Compton, and Parkinson. Among these were, the *Acer platanoides*, the American spindle tree, the kermes oak, the dwarf almond, the scarlet thorn, the *Laurus Benzoin*, the liquidambar, the Aleppo pine, and the cedar of Lebanon. The principal authority is Sutherland's *Catalogue of the Plants in the Edinburgh Botanic Garden*, published in 1683.

From the year 1688 to the year 1700 inclusive (James II., and William and Mary), thirty-one species were introduced, by Bishop Compton, the Honourable Charles Howard, the Duchess of Beaufort, Jacob Bobart, son of the first superintendent of the Oxford Botanic Garden, and others. The authorities are to be found in Ray's *Historia Plantarum*, in the *Phytographia* of Plukenet, and in Bobart's *Historia Plantarum Oxoniensis*. The titles of all these catalogues, and several others used as authorities for the dates of the introduction, or rather first record, of plants, are given in the preface to the second edition of the *Hortus Kewensis*.

The botanists to whom the British arboretum was most indebted during the seventeenth century were, Parkinson, Tradescant junior, Ray, and Sutherland; and the principal botanical amateurs were, the Bishop of London and the Duchess of Beaufort. Parkinson was born in 1567, and was contemporary with Gerard and L'Obel. He possessed a rich garden, and was appointed apothecary to James I. He appears to have died somewhere about 1650. John Tradescant junior inherited his father's museum, and published a catalogue of it, entitled *Museum Tradescantianum*, in 1656. He died in 1662, bequeathing the museum to Mr. Ashmole, who lodged in his house, and whose name the museum now, "unjustly," as Pulteney remarks, bears in Oxford, where it is deposited. John Ray was born at Black Notley, near Braintree in Essex, in 1628. His father, though a blacksmith, contrived to give him a college education. At college, he imbued the minds of some of his companions with a taste for plants, and he pursued this taste himself at every leisure opportunity. In 1660 he was ordained deacon and priest, and after this time he made various journeys throughout Britain, and visited the Continent. He was the author of numerous works, the principal of which relating to plants are, his *General History of Plants*, his *Methodus Plantarum*, and his *Synopsis Methodica Stirpium Britannicarum*. He died in 1704, at his birthplace, at the age of 76.

SUBJECT. 4. *Of the Foreign Trees and Shrubs introduced into Britain in the 18th Century.*

A HOST of amateurs, botanists, and commercial gardeners enriched the British arboretum during this century. In the preceding one, the taste for foreign plants was confined to a few, and these not the richest persons in the community; but generally medical men, clergymen, persons holding small situations under government, or tradesmen. In the 18th century, the taste for planting foreign trees extended itself among the wealthy landed proprietors; partly from the influence of the Princess Dowager of Wales, who established the arboretum at Kew, and partly from the display previously made by Archibald Duke of Argyle at Whitton, the Duke of Richmond at Goodwood, and others. Towards the middle of the century, the change introduced in the taste for laying out grounds, by Pope, Addison, and Kent; and the circumstance that Brown, who had been a practical gardener, was extensively employed in remodelling country residences according to this new taste, must have greatly contributed to increase the number of species employed in plantations; and hence we have the collections at Croome, at Syon, and at Claremont. The writings of Miller, Bradley, Switzer, and Linnæus, and the consequent spread of botanical knowledge among the educated classes about the middle of the century or before, must have enlightened practical men to a degree far exceeding that which had ever previously existed.

In order to give a general view of the state of gardening in England in the first half of the 18th century, as far as it respects foreign trees, we shall begin by giving a summary notice, by Collinson, of the chief encouragers of gardening and planting of his time. Peter Collinson was born in London, in 1693: he was a quaker, and a linendraper. He had a country house and garden, first at Peckham in Surrey, and afterwards at Mill Hill, near Hendon in Middlesex. He appears to have taken possession of the latter place, Ridgeway House, sometime previous to 1749. He was a great lover of animated nature in every form; and in one of his letters, published by Sir James Edward Smith, in the *Linneæan Correspondence*, he declares that every living thing called forth his affections. In a note written in 1768, in one of his copies of Miller's *Dictionary*, which was purchased from one of his lineal descendants in January 1835, by A. B. Lambert, Esq., and which, through the kindness of that gentleman, we have just seen, he declares, at the age of 68, that the plants in his garden at Mill Hill furnish his greatest source of happiness. He died in 1768. In the year 1764, he made notes on some blank leaves in a copy of Miller's *Dictionary*, and

again in 1768, in another copy of that work; and the following extract from those made in 1764 is abridged from a communication by A. B. Lambert, Esq., to the *Linneæan Transactions*, vol. x.:—

“The gardeners about London in 1712,” he says, “were remarkable for fine cut greens, and clipped yews in the shapes of birds, dogs, men, ships, &c. Mr. Parkinson, in Lambeth, was much noticed for these things, and he had besides a few myrtles, oleanders, and evergreens. At that time, Mr. Rench, who lived behind the Earl of Peterborough’s at Parson’s Green, was famous for tulip trees: he began the collecting of evergreens, arbutuses, phillyreas, &c.; and from him came the gold and silver hedgehog holly. He gave rewards for accidental varieties of the common holly, and thus obtained the saw-leaved variety, and a variegated holly which bears his name. He and Parkinson died about 1724. Brompton Park and Hunt’s at Putney were fine nurseries. In 1764, Chelsea Garden excelled all the others in Europe for variety of plants. In 1759, there were, in the American grove at Goodwood, two fine great magnolias [*M. grandiflora*], about 20 ft. high, that flowered annually.” Collinson adds, that his tree of this species flowered in 1760, which he had raised from seed 20 years before. “Lord Petre, he continues, “who was the ornament and delight of the age he lived in, removed, in the spring of 1734, twenty-four full-grown elms about 60 ft. high, and 2 ft. in diameter: all grew finely, and now (1764) are not known from the old trees they were planted to match.” In 1738, he planted an avenue of elms 15 or 20 years old, cedars 20 years old, and larches 11 years old. John Clarke, a butcher at Barnes, was famous for raising cedars from seed, from the great tree at Hendon Place; and also for raising plants of the small magnolia [*M. glauca*]. Clarke sold a thousand cedars in 1761, five years old, for 79*l.* 6*s.*, to the Duke of Richmond, which were all planted at Goodwood, and did well. The cedars at Whitton were all raised from seed by the Duke of Argyle in 1725. In 1762, most of the duke’s rare trees and shrubs were removed to Kew, then belonging to the Princess of Wales, and under the direction of Lord Bute. Mr. Vernon, a Turkey merchant at Aleppo, brought the weeping willow from the river Euphrates to his seat at Twickenham Park, where Collinson saw it growing in 1748. This was the original of all the weeping willows in our gardens. [In the *Hortus Kewensis*, the weeping willow is stated to have been cultivated at Hampton Court in 1692.] In 1761, Mr. Sharpe, at South Lodge, in Enfield Chase, invited Mr. Collinson to dine with him, and to see the *Cornus florida* in flower. In 1746, Mr. Collinson received the first double Spanish broom from Mr. Brewer of Nuremberg. In 1756, the famous tulip tree in Lord Peterborough’s garden at Parson’s Green, near Fulham, died. It

was about 70 ft. high, and perhaps 100 years old, being the first tree of the kind that was raised in England. "It had, for many years, the visitation of the curious, to see its flowers, and admire its beauty. It was as straight as an arrow, and died of age, by a gentle decay." (*Abridged from Mr. Collinson's paper, as quoted by Mr. Lambert, in the Linnæan Transactions, vol. x. p. 282.*)

On a blank leaf of another copy of Miller's *Dictionary*, Collinson adds the following names of proprietors of gardens to the above list:—Reynardson, at Hillingdon, near Uxbridge, whose fine collection, he says, was sold to Mr. Robert Walpole; Mr. Parker, near Croydon; Dr. Lumley Lloyd, at Cheam in Surrey, "who gave his house and great collection of plants to the Duke of Bedford;" Sir Harry Trelawney, of Buttshead, near Plymouth, who had a great collection of hardy trees and shrubs; Sir Harry Goodrick, at Ribstone in Yorkshire, who was a great collector and naturaliser of exotic trees; Mr. Charles Dubois, at Mitcham, remarkable for his collection both of house and of hardy plants; and Mr. Blackburne, at Orford, near Warrington in Lancashire [a catalogue of whose garden was published in 1779], who had a great collection, particularly of stove plants, kept in the highest degree of perfection. Collinson also mentions, in one of the memoranda in this volume, that *Tradescant junior* was the first who propagated American plants for sale in England.

In Collinson's garden at Mill Hill, the *Periploca græca*, and numerous other trees and shrubs, as will be seen by the list at the end of this section, flowered for the first time in England. It was kept up some years after Peter Collinson's death, by his son, Michael Collinson. Afterwards it fell into the hands of Richard Anthony Salisbury, Esq., F.R.S. About the end of the century it was purchased by the protestant dissenters, for a foundation grammar school: the house was turned into lodging-rooms for the boys, and Collinson's stable fitted up as a chapel. A new house has since been built.

On examining the grounds which formerly belonged to Ridgeway House, in January, 1835, several trees and shrubs planted in the time of Collinson were found to be still remaining. A *platanus* 40 ft. high, and $1\frac{1}{2}$ ft. in diameter at a foot from the ground; a deciduous *cypress* 48 ft. high, and $1\frac{1}{2}$ ft. in diameter; four *pinasters*, the diameter of the largest of which was 3 ft.; two of *Pinus Cembra* with trunks nearly 2 ft. in diameter, and from 50 to 60 ft. high, which must be the finest specimens of this tree in England; a tulip tree 30 ft. high, diameter 9 in.; and two cedars with clear trunks between 30 and 40 ft. high, and diameters of nearly 4 ft., the branches of which cover a space of 60 ft. in diameter. Near the spot where Collinson's house stood (for it is now pulled down) there is a cedar 60 ft. high, with its lowest branches reclining on the ground, and covering a space of 70 ft. in diameter.

Near it are a very old laburnum, and a sweet chestnut, with a trunk nearly 5 ft. in diameter, and its branches extending 30 ft. on each side. There are a *Quercus Ilex* covering a space of 35 ft. in diameter with its branches; and a weeping willow 50 ft. high; there are a Chinese arbor vitæ 25 ft. high; two red cedars from 30 to 40 ft. high; an upright cypress 40 ft. high, which the present gardener says was planted by Linnæus, but this could not be the case, as Linnæus left England in 1737; a hemlock spruce with two trunks, each 1 ft. in diameter, and 50 ft. high, with branches extending about 30 ft.; two Portugal laurels, each covering a space 40 ft. in diameter; an arbutus 1½ ft. in diameter, with branches extending 20 ft.; a very handsome variegated holly covering a space 18 ft. in diameter; a handsome box tree 15 ft. high; and a cone of laurustinus 20 ft. in diameter at the base; besides several other trees and shrubs evidently as old as the time of Collinson. It is greatly to the credit of the proprietors of the school, that all these fine specimens are carefully preserved, and the name of Collinson respected as it ought to be.

The large cedar referred to, at Hendon, was blown down, at a mature age, on the 1st of January, 1779. Its height was 70 ft., and the diameter of the space covered by its branches 100 ft. The girth of the trunk, at 7 ft. from the ground, was no less than 16 ft.; at 12 ft., it was 20 ft. in circumference; and the limbs varied in girth, from 6 ft. to 12 ft. The gardener, two years before it was blown down, made 50*l.* of the cones. (*Lysons*, vol. ii. p. 395.)

Peterborough House, mentioned by Collinson, is described by Bowack, in his account of Fulham, in 1706, as having 28 acres of ground attached to it, in which was a tulip tree 76 ft. in height, and 5 ft. 9 in. in girth. Swift, in one of his *Letters*, speaks of Lord Peterborough's garden as one of the finest that he had seen about London. The villa is now (1835) the property of T. Sampayo, Esq., and is no way remarkable for its trees and shrubs.

Whitton, a villa and grounds belonging to the Duke of Argyll, near Hounslow, began to be planted when the duke was Earl of Islay, about the year 1720. Collinson informs us that all the cedars at that place were raised from seed in the year 1722. He also mentions that the *Anona (Asimina)* triloba flowered at Whitton every year; and Weston informs us that the duke's oranges, lemons, limes, and citrons, grown on an open wall, and only sheltered by glass during winter, were the finest in England. (*Tracts, &c.*, p. 201.) Archibald Duke of Argyll was grandson to the Duchess of Lauderdale; he was born at Ham House near Richmond in 1682, and died in London in 1761, aged 79, leaving all his real and personal estate in England to Mrs. Elizabeth Anne Williams. A copy of

his will is given in the *Gentleman's Magazine*, vol. xxxi. p. 206., in which the duke gives one year's wages to each of his servants in London, and to Daniel Craft, his gardener at Whitton, and to his housekeeper there, except, says His Grace, "my cook, whose wages are too high." The duke, who succeeded to that title in 1742, collected all the foreign trees and shrubs which, at that time, were to be procured either at home or abroad: many of the former still remain at Whitton, but all those removable were sent to the Princess Dowager of Wales's garden at Kew, in 1762, after the duke's death, his paternal property and dukedom going to a distant relative. The grounds were sold and divided, and now form three distinct villas, which are remarkable for their fine specimens of cedars, Weymouth pines, silver firs, deciduous cypresses, walnuts, hickories, and American oaks.

In February, 1835, through the kindness of the Misses Gostling, the present proprietors of Whitton Place, and the assistance of Mr. Castle, of the Twickenham Botanic Garden, and of Mr. West, gardener at Whitton, a number of the trees were measured for us. Among these are, a Lombardy poplar 115 ft. high, and 19 ft. 8 in. in girth at 2 ft. from the ground; a cedar of Lebanon 71 ft. high and 14 ft. in girth; a silver fir 95 ft. high, and 10 ft. 6 in. in girth; a Weymouth pine 81 ft. high, and 11 ft. 8 in. in girth; a *Quercus Phellos* (the willow oak) 70 ft. high; and a black hickory 60 ft. high. There are a number of other fine trees, larches, firs, pines, oaks, robinias, gleditschias, &c., on this and other portions of the estate, of which notices will be found in the *Gard. Mag.*, vol. xi.

The seat of the Lord Petre so highly enlogised by Collinson was at Thorndon Hall in Essex; and that place still contains some fine old exotic trees. Speaking of Robert Lord Petre, Collinson, in a letter to Linnæus, dated Jan. 18. 1743, observes that "the death of the worthiest of men, the late Lord Petre, has been the greatest loss that botany or gardening ever felt in this island. He spared no pains or expense to procure seeds and plants from all parts of the world, and then was as ambitious to preserve them." After speaking of his stoves as such "as the world never saw, and may never see again," and giving the dimensions of the more remarkable plants grown in them, he says, "the collections of trees, shrubs, and evergreens in his nurseries at his death, I had told over; and they amounted to 219,925, mostly exotic. As this young nobleman was the greatest man in our taste that this age produced, I thought it might not be unacceptable to give you some account of the greatness of his genius; but his skill in all the liberal arts, particularly in architecture, statuary, planning, and designing, planting, and embellishing his large park and gardens, exceeds my talent to set forth." (*Smith's Linnæan Correspondence*, vol. i.

p. 11.) Robert James, eighth Lord Petre, died in 1742, at the age of 29 years. Linnæus has named a genus of plants *Pètrea*, in commemoration of this nobleman.

Goodwood, near the coast of Sussex, enjoys a mild climate; but the soil, which is thin and on chalk, is not favourable to the growth of trees. The park contains a great number of cedars; but there are not many other foreign trees, except ilices, cork trees, acacias, some acers, and oaks. The magnolias mentioned by Collinson, in his notes of 1764, no longer exist; in Collinson's notes of 1768, he states that all the moveable articles were sold at the duke's death. Miller mentions that a great many trees of the true service were planted at Goodwood; but in 1828, and again in 1831, we sought in vain in the woods for a single specimen.

Samuel Reynardson, Esq., resided at an ancient house, called the Cedar House, from the celebrated cedar which grew in the garden. This cedar was planted by Reynardson, who resided at Hillingdon from 1678, till his death in 1721. This tree was probably one of the first that were planted in England. Lightfoot measured it in 1779; it was then 53 ft. high; the diameter of the space covered by the branches measured from east to west 96 ft., and from north to south 89 ft. The girth, close to the ground, was 13 ft. 6 in.; at 7 ft., 12 ft. 6 in., and at 12 ft., 14 ft. 8 in. It was cut down in 1789, in consequence of one of the branches being broken off by a high wind. It produced 460 ft. of timber, $6\frac{2}{3}$ loads of stack wood, and 125 faggots. It was sold to a carpenter for 10*l.*, and he retailed it for 22*l.* 17*s.* (*Laysons*). Reynardson made a curious will, leaving all his property to the vicar of Hillingdon for ever, to build a room to hold his library and museum. He ordered all his plants to be sold for this purpose. The present vicar of Hillingdon is Dr. Hodgson, the very reverend the Dean of Carlisle, to whom we have written twice, to learn in what state this library and museum now are, but without receiving any answer.

Cashiobury, near Watford in Hertfordshire, the seat of Arthur Capel Earl of Essex (the Earl of Essex who patronised Gerard, and had a seat at Barn Elms, was the celebrated and unfortunate Robert Devereux), was noted for its trees in the time of Evelyn; Cooke, His Lordship's gardener, was the author of a work on forest trees, published in 1676, and afterwards a partner in the Brompton Nursery. Evelyn, in his *Diary*, mentions Cooke as being a skilful artist in the mechanical part of gardening, not ignorant in mathematics, and having some pretensions to astrology. At Cashibury there is now (1835) one of the finest specimens of *Quercus tinctoria* in the country. There are also many fine magnolias, *Pinus Cembra*, tulip trees, cypresses, cedars, and other species.

The limited variety of evergreens which existed in the London nurseries at the beginning of this century, according to Collinson, is confirmed by the preface to the first edition of Miller's *Dictionary*, published in 1724, in which it is stated that the catalogue of evergreens kept by nurserymen for sale contained only twelve sorts; viz., alaternus, arbutus, bay, box, holly, juniper, laurel, laurustinus, phillyrea, pyracantha, Italian green privet, and yew. In the eighth and last edition which Miller published of his *Dictionary*, dated March 1. 1768, the number of plants then cultivated in England is said to be more than double those that were known in 1731. It appears difficult to reconcile Collinson's and Miller's relation with the ample list of trees and shrubs published in 1730, in the nurserymen's *Catalogue*, which we are about to give some account of: most probably Collinson and Miller referred only to the ordinary gardens and nurseries; or perhaps the defective state of the catalogues of these was one reason why the larger catalogue about to be noticed was produced. From Bradley's *New Improvements of Planting and Gardening*, published in 1720, the former appears to be the case; for, after treating of the common shrubs, he speaks of American trees, which were commonly kept in pots, and housed during the winter; so that what are now considered hardy plants were then looked upon as inmates of the greenhouse, or of the cold frame or covered pit. (*Bradley, New Improvements, &c.*, p. 87.)

In the year 1730, the Society of Gardeners, consisting of all the principal nurserymen and florists about London, published *A Catalogue of Trees and Shrubs, both Exotic and Domestic, which are propagated for Sale in the Gardens near London*. This catalogue is in folio, ornamented with an elaborate frontispiece, containing a vista in a garden laid out in the ancient style, exhibiting walls, pillars, and arcades of clipped verdure; and the descriptions of the trees and shrubs refer to engraved plates. As this may be considered a work of unquestionable authority, we shall quote, from the preface, a general view there given of the principal encouragers of planting and gardening in England, previously to and at that time.

The work is dedicated to the Earl of Pembroke and Montgomery, F.R.S.; and, after complimenting His Lordship on his various merits, the Society say, "Your Lordship's good taste in, and great encouragement of, planting and gardening, are fully displayed in those noble gardens at Wilton, where are a greater number of the trees here treated of, and in a more flourishing condition, than can be found in any one garden in this kingdom besides." (p. iv.)

There are now (1831) a considerable number of cedars at Wilton of a large size, and some planes, limes, evergreen oaks,

horsechestnuts, red cedars, arbor vitæ, laurels, bays, &c., which probably were planted in or previously to 1730. (See Index, Cedar, &c.)

In the preface (after praising the temperature of the British climate; the Royal Society, Charles II., and William III., and also Malpighius, Grew, Ray, &c., are complimented, as having paved the way for the improvement of gardening. "The profits and innocent delights of this art," they say, "have allured into it many learned and curious persons, nobility and gentry;" and these "have not contented themselves with the narrow compass and mean stock of our former poorly furnished gardens, but they have industriously procured, from abroad, trees, plants, flowers, and fruit, not only from our own plantations in America, but those also of other parts of Europe, nay, even Asia and Africa. Among these generous procurers of plants, &c., we cannot," the Society add, "forbear mentioning the following worthy persons:—

"First, Dr. Compton, late Bishop of London, who was an early introducer of exotic trees and plants, many of which were grown to a considerable size in the open air, in those formerly well-stocked gardens at Fulham, most of which have been since destroyed, to the great regret of many curious persons.

"Much about the same time Samuel Reynardson, Esq., began to furnish his fine gardens at Hillingdon, near Uxbridge, with a great variety of curious plants, which his great correspondence abroad enabled him to procure from divers parts of the world; but, as he kept them for the most part confined to pots and tubs, preserving them in green-houses in winter, never attempting to naturalise them to our climate, so, soon after his death, that valuable collection was dispersed, as at present to be hardly known what he was possessed of.

"In the like manner, also, the curious Dr. Uvedale of Enfield did, by his great correspondence abroad, collect a very valuable parcel of plants and flowers, which he, with great skill and care, maintained for many years; and some of the valuable trees were planted in the full ground, where they are now (1730) remaining; but the bulk of his collection was sold to Sir Robert Walpole, soon after the doctor's death.

"Her Grace the Duchess of Beaufort did also collect a numerous quantity of rare plants into those famous gardens of Badmington, where she preserved and maintained them with great care in wonderful beauty for many years; but this collection also consisted chiefly of the most tender exotic plants.

"The Earl of Pembroke began about the same time to plant those magnificent gardens at Wilton, with all the different varieties of curious exotic trees and shrubs as would endure the cold of our climate in the open air; in which His Lordship hath

so well succeeded, as to have the best collection of those trees, which are advanced to a considerable size, that can be found now growing in any one garden in this kingdom: and it was from examples of this kind that people were encouraged to make further trials of what plants, trees, flowers, and fruits could be brought to thrive in our climate without the trouble and expense of housing in winter; and it has been from repeated trials and experiments of this kind, that the many noble trees, fruits, and flowers now in England have by degrees been naturalised to our coarse climate, to the no small pleasure of all the delighters in the innocent diversions of gardening.

“Nor should we, in mentioning particular persons, forget the many various gentlemen which at present are carrying this spirit of gardening to a considerable height, by introducing many new kinds of plants, flowers, trees, and fruits, and in making many curious experiments concerning their culture and uses; from all whose observations and experiments we may daily expect something new in the business of gardening and vegetation; amongst which persons are, the Earl of Islay, Lord Wilmington, Sir Charles Wager; Sir Harry Goodrick, Bart., in Yorkshire; the Reverend and Honourable Lumley Lloyd, Esq., at Cheam, in Surrey; Henry Trelawney, Esq., at Buttshead, near Plymouth; Henry Marsh, Esq., at Hammersmith; George Dennis, Esq., in Cornwall; Dr. Beeston, of Ipswich; Mr. James Sherard, of Eltham, in Kent; — Topham, of Windsor; Mr. Peter Collinson, with several others, too many to be here enumerated.

“But to none of the before-mentioned persons is England more indebted for introducing trees, plants, flowers, and fruits, than to the learned and ingenious Charles Dubois, Esq., of Mitcham, who has not only been very industrious to procure plants from abroad, but also as generous in communicating whatever his garden would afford, as also many useful observations relating both to their culture and uses, to all delighters in planting and gardening; and it is to him that we are greatly indebted for many valuable trees and plants which enrich this catalogue.

“And after mentioning the particular persons to whom England is thus indebted for introducing so many valuable trees, plants, flowers, and fruits, we cannot in justice omit to mention the establishing of the public botanic garden at Chelsea, by the worshipful company of Apothecaries of London, not only for medical instruction, but also for introducing still a greater variety of trees and plants.” Afterwards, when speaking of botanic gardens, it is observed by the Society, that there is only one in England which deserves the name, and that is, “that of Chelsea, maintained at the expense of the worshipful company of Apothecaries.” (p. viii.) The reader is next taught what he is to expect; viz., “an exact catalogue of the several sorts of trees

and shrubs, which will endure to be planted in the open air in England, which are to be found in the several nurseries near London," arranged in alphabetical order, and with short descriptions. The preface is signed by the twenty "gardeners and nurserymen" composing the Society, among which are Fairchild of Hoxton, Furber of Kensington, Miller of the Physic Garden, Chelsea, Gray of Fulham, and F. and S. Hunt of Putney.

Some of the patrons of gardening in the above enumeration have been already mentioned, and of the others we know but little. Spencer Compton, speaker of the House of Commons in 1714, and afterwards Earl of Wilmington, was a near relative of Bishop Compton. He died in 1743. Lewis Kennedy, one of the founders of the Hammersmith Nursery, was gardener to him in 1739. Sir Charles Wager had a residence at Parson's Green, where he introduced the scarlet maple (which was then called Wager's maple) in 1725. A *Magnolia grandiflora* flowered in his garden in 1737. He died in 1743. Collinson says that a tulip tree, which had been raised from a seed which he gave Sir Charles Wager, flowered for the first time when it was thirty years old, in 1756; and Lysons mentions a cedar of remarkable growth, which grew near the house, in Sir Charles's garden. (*Environs*, &c., ii. 829.) The grounds at Mitcham, which belonged to Mr. Dubois, are now (Jan. 1835) the property of Mr. Blake, an auctioneer at Croydon. Dubois's house has been long since pulled down; but another has been built, which is occupied by Mrs. Beckford. In the grounds a number of the trees planted by Mr. Dubois still remain. Among these are a very large weeping willow; a nettle tree, with branches covering a space 50 ft. in diameter, and with a trunk 6 ft. 8 in. in circumference. The extremities of the branches hang down nearly to the ground; and on Jan. 10, 1835, when we had the tree examined, the spray was still covered with dark purple berries, rather larger than those of the common hawthorn. There is a pinaster, with a clear trunk about 40 ft. high; the girth, about 3 ft. from the ground, 9 ft.; and the total height 60 ft. The cracks in the bark of this tree are from 6 in. to 8 in. deep. There is a very old, large, and handsome mulberry tree, the branches of which cover a space of 60 ft. in diameter; it bears abundantly every year. Besides these, there are very large and old Scotch pines; a large old stone pine; large *Prunus Mahaleb*; a fine *Ptèlea trifoliata*; a stag's horn sumach, with a trunk 6 ft. in girth; an old *Bignonia radicans*; a large arbutus, and some other fine specimens. Dubois died in 1740, aged 83 years.

The following is an abridged list of the above-mentioned catalogue of the Society of Gardeners, with the modern names, as given in our *Hortus Britannicus*, as far as we have been able to ascertain them:—

1. *Ranunculàcææ*. *Clématis Vitalba*, V. simple-leaved, ? cirrhòsa, ? flòrida; *Viticèlla*, blue; V., blue, double: V., purple; *Viórna*.

2. *Magnoliàcææ*. *Liriodéndron Tulipífera*, also "the laurel leav'd tulip tree."

3. *Berberidææ*. *Bérberis vulgàris*, v. white-fruited, v. seedless-fruited, canadénsis.

4. *Cistinææ*. *Cistus ladaníferus albiflòrus*, l. maculàtus, ? latifòlius, ? cýprius, ? incànus, sp.

5. *Malvæcææ*. *Lavàtera O'lbia*. triloba, "Althæa, frutescens *Bryoniæ folia*. C.B.P. 316." *Hibiscus syriacus*, red flowered, s., purple-flowered; s., white-flowered; s., striped-flowered; s., striped-leaved.

6. *Tiliàcææ*. *Tília europæa*; e., variegated-leaved; platyphýlla, parvifòlia, pubéscens.

7. *Hypericinéææ*. *Hypéricum hircènum*, ? canariènsis.

8. *Acerinéææ*. *Acer Pseùdo-Plátanus*; *Pseùdo-Pl.* variegated-leaved; campéstre, *platanoides*; *pl.* variegated-leaved; *rùbrum* ("the Virginian flowering maple," &c.) ; and another sort of "the Virginian flowering maple." *Negúndo fraxinifòlium*.

9. *Hippocastànææ*. *Æsculus Hippocástanum*; H., with leaves variegated with yellow; H., with leaves variegated with white; *Pàvia rùbra*.

10. *Vites*. *Vítis vulpìna*, ? *Labrúsca* ("the wild Virginian grape"); and these varieties of *vinífera*, "the parsley-leav'd vine," "the blotch'd-leav'd vine," and "strip'd leav'd vine." *Ampelópsis bipinnata* and *hederàcææ*.

11. *Zygophýllææ*. *Meliánthus mājor* and *mìnor*.

12. *Xanthoxýllææ*. *Ptèlea trifoliàta*, *Cnèdum tricóccum*.

13. *Rutàcææ*. *Rúta gravèolens*, ? *angustifòlia*; ? an., var. "the [silver]strip'd narrow-leav'd;" *chalepénsis*.

14. *Ilicinéææ*. *Ilex Aquifòlium*, 33 varieties of, whose characteristics are given; *vomitòria*, "South Sea thea tree;" sp. ("Aquifolium; Carolinianum, angustifolium, spinis raris brevissimis." *Carolina holly*, with smooth leaves.

15. *Staphyleàcææ*. *Staphylèa pinnàta* and *trifòlia*.

16. *Celastrínææ*. *Euónymus europæus* and *latifòlius*.

17. *Rhámneææ*. *Rhámnus Alatérnus*; *A.*, blotched-leaved; *Clùsi*; *C.*, gold-edged-leaved; *C.*, silver-edged-leaved; *Frángula*, *cathárticus*, ? *infectòrius*, sp. ("Rhamnus; spinis oblongis, cortice albo, Monspeliensium. J. B., vol. i. pars 2. 31.") *Palìurus aculeàtus*.

18. *Anacardiàcææ*. *Rhús Cótinus*, *typhìna*, *Toxicodéndron*, *radicans*, ? *vérnix*, *Coriària myrtifòlia*, *Pistàcia*, *Terebínthus* and *officinàrum*.

19. *Leguminòsææ*. *Gledítschia triacánthos*, *Robínia viscòsa*, sp. ("Acacia, Caroliniana, aquatica, *Abruaæ folio*, spinis rarioribus.

Water acacia.") *Pseùd-Acàcia* with rough pods, *Pseud-Acàcia* with smooth pods. *Amórpha*, ? fruticòsa; *Colùtea arborèscens*, *Sutherlandia frutèscens*; *Cýtisus scopàrius*, sp. (? sessilifòlius ? nígricans), alpínus, ? alpínus var., *Labúrnum*, álbus; *Dorýc-nium suffruticòsum*, *Coronilla E'merus*, *E'merus* ? "minor," *U'lex europæ'a* and *nàna*, *Genista ànglica*, *Medicàgo arbòrea*, *Wistària frutèscens*; *Cércis Siliquástrum*, canadénsis, sp. ("Carolina pointed-leav'd Judas tree:" this is given as distinct' from the previous two); *Spártium júnceum*.

20. *Amygdalæ*. *Amýgdalus commúnis*; c., bitter-kerneled; c., tender-shelled; c., white-flowered; *Armeniaca vulgàris*, and a striped leaved variety of it; *Cérasus Pàdus*, *P.* the Cornish variety; the Flanders cluster cherry, the double-flowered cherry tree, the double-flowered cherry with very large flowers, the black cherry or mazzard, the common wild or honey cherry, the wild northern English cherry, the garden cherry with the leaf elegantly striped; *Mahàleb*, *Laurocérasus*, *L.* with the leaf variegated with yellow, *L.* with the leaf variegated with white; *Pérsica vulgàris*, double-flowered; *Amýgdalus nàna* and *pùmila*; *Prúnus spinòsa*, *insitítia*; i., white fruited; and the following varieties of, we assume, domestica: "the damson," "the great violet damson," "the strip'd [leaf] perdrigon plumb," "the strip'd [leaf] muscle plumb," "the cherry plumb," "Christmas plumb or winter creke," and a kind "flore pleno."

21. *Rosàcæ*. *Ròsa canina*, villòsa, sp. allied to villòsa, spinosíssima and two varieties of it, rubiginòsa and var. flore pleno, sp. or var. ("Rosa, rubra, multiplex. C. B. P."), cinnamòmea, c. fl. pl.; lùtea, lùtea punicea, sulphùrea, turbinàta, semper-virens, moschàta; m., double-flowered, "rose without thorns," "the Virginian briar," álba and varities of it; and these ornamental varieties of other species, the single damask, the double damask, the York and Lancaster, the red Belgick, the blush Belgick; the single Provence, the damask Provence, the red Provence, the common Provence, the moss Provence, the Dutch hundred-leaved; the single velvet, the double velvet, the rosa mundi, the marbled, the royal virgin, "the monthly rose," "the strip'd monthly;" *Potentilla fruticòsa*; *Spiræ'a salicifòlia*, *opulifòlia*, *hypericifòlia*; *Rùbus fruticòsus*; fr., white-fruited; fr., variegated-leaved; *cæsius*, *idæus*; i., white-fruited; i., the late red; *odoràtus*, "the upright Pennsylvania bramble, or raspberry," "the Virginian black raspberry."

22. *Pomàcæ*. *Pýrus A'ria*, "the white beam tree with long leaves and small red fruit," "the red chess apple, or English wild service;" *Sórbus* ("the pear-shaped true service"); *S.*, "the round late-ripe service;" *aucupària*, a. "foliis ex luteo variegatis," *torminàlis*, *Màlus*, "the strip'd [leaf] crab tree," "the strip'd [leaf] apple tree," "the paradise apple," "the figg apple,"

“the Virginian crab tree with sweet flowers,” *communis* “with strip’d leaves,” “the double-blossom pear,” “the twice-flowering pear;” *Cydonia vulgaris*, the pear quince, apple quince, and Portugal quince; *Mespilus germanica*; *g.*, the great Dutch; *Crataegus Oxyacantha*; *O. fl. pl.*; *O.*, Glastonbury; *O.*, white-fruited; *Azarolus*, “the Virginian azarol with red fruit;” *Pyracantha*, “the Virginian cockspur,” “the Virginian hawthorn with long sharp thorns,” “the yellow-berried Carolina hawthorn;” *Amelanchier vulgaris*, *Photinia arbutifolia*,? *Cotoneaster vulgaris*.

23. *Granatæe*. *Punica Granatum*. *G.* flore pleno.

24. *Tamariscinæe*. *Tamarix gallica* and *germanica*.

25. *Philadelphæe*. *Philadelphus coronarius*; *c.*, double-flowered; *c.*, variegated-leaved; *c. nanus* (“nunquam florens,” which never flowers).

26. *Passifloræe*. *Passiflora carulea*; *c.*, with yellow blotch’d leaves; “the narrow leav’d passion flower, with lesser and paler colour’d flowers;” “the yellow passion flower,” “the three-leav’d passion flower.”

27. *Crassulæe*. *Sedum populifolium*.

28. *Grossulæe*. *Ribes Grossularia*; eight kinds of goose-berry, among which are “the yellow-leav’d” and “the strip’d-leav’d;” and included in the species *rubrum* these varieties, common red currant, champagne, large red, white, large white, “the strip’d-leav’d white,” “the yellow-strip’d-leav’d currant tree,” “the silver-strip’d currant tree;” *alpinum*: *a.*, with leaf variegated with yellow; *nigrum*; *n.*, variegated-leaved; “*Ribes Americana*, fructu nigro;” “small wild currant.”

29. *Araliæe*. *Aralia spinosa*.

30. *Caprifoliæe*. Kinds of honeysuckle named as follows:—the Italian, yellow Italian, early or French white, late red, Dutch, English long-blowing, oak-leaved, English white, Russian, evergreen, common with striped leaves, striped honeysuckle with hairy indented leaves. *Lonicera Xylosteum*, *alpigena*, and *caerulea*; *Symphoria glomerata*. *Viburnum Lantana*; *L.*, variegated-leaved; *levigatum*, *Opulus*, *O. rosea*, *O. folio variegato*, *Tinus*; *T.*, *hirta*; *T.*, “*foliis ex luteo variegatis*;” *T.*, “small-leav’d;” *lucidum*, *l.* “*foliis ex albo variegatis*.” *Hedera Helix*. *H.*, “yellow-leav’d;” *H.*, “silver-strip’d;” *H.*, “the yellow-strip’d.” *Sambucus nigra*; *n.*, greenish-berried; *n.*, white-berried; *n.*, leaf variegated with yellow; *n.*, “parsley-leav’d;” *racemosa*.

31. *Cornæe*. *Cornus mas sanguinea*; *s.*, striped-leav’d; “the Virginian.”

32. *Compositæ*. *Baccharis halimifolia*.

33. *Vacciniæe*. *Vaccinium uliginosum* and *Myrtillus*.

34. *Ericæe*. *Arbutus Unedo*; *U.*, double-flowered; “the strawberry tree with oblong flowers and egg-shaped fruit.”

35. *Styracineæ*. *Halésia tetráptera*.
36. *Ebenàceæ*. *Diospýros virginíana* and *Lótus*. Perhaps the “*Celtis fructu luteo ampliori*” is a variety of *D. virginiana*.
37. *Oleàceæ*. *Fráxinus excélsior*; e., leaf striped with yellow; ? *carolinàina*, sp. (“the New England ash”); *Órnus europæa* and *rotundifòlia*; *Ligústrum vulgàre* v., leaf yellow-variegated; v., leaf silver-variegated. *Syrínga vulgàris*, these varieties of it, blue-flowered, purple-flowered, white-flowered yellow blotched-leaved, white blotched-leaved; *pérsica*; p., cut-leaved. *Olea europæa*, e. *buxifòlia*, *Phillýrea latifòlia*; l., leaf yellow-variegated; *ligustrifòlia* *angustifòlia*; a., *rosmarinifòlia*: *oleafòlia*.
38. *Jasmíneæ* *Jasminum officinàle*; o., leaf yellow-variegated; o., leaf white-variegated; *frùticans*, *hùmile*.
39. *Bignoniàceæ*. *Técoma radicans*, r. *mìnor*; *Biguònia capreolàta*, *Catálpa syringafòlia*.
40. *Solàneæ*. *Solànum Dulcamàra*; D., white-flowered; D., leaf white-variegated.
41. *Labiàtæ*. *Phlòmis fruticòsa*, ? *lanàta*; *Rosmarínus officinàlis*; o., silver-striped-leaved; o., yellow-striped-leaved; o., *latifòlius*; sp. (“*Rosmarinus*; *Almeriensis*, flore majore, spicato, purpurascente. Tourn. 195. The large flowering rosemary.”) *Sàlvia officinàlis*; o., leaf variegated with green and white; o., leaf red; o., leaf particoloured; sp. (“*Salvia*; *absinthium redolens*. J. B. iii. 307. Wormwood sage”), sp. (“*Salvia*; *minor*, *aurita et non aurita*”), and a variety of the preceding (“*Salvia*; *minor*, *foliis variegatis*”). *Teucrium* ? *frùticans*, sp. (“*Teucrium*; *Bœticum*, calice campanulato. Boerh. Ind. Alt. 181. Spanish tree germander.”) “*Galeopsis*; *hispanica*, *frutescens*, *Teucii folio*. Tourn. The base-horehound tree.”
42. *Verbenàceæ*. *Vítex A'gnus cástus*, A. *latifòlius*.
43. *Chenopòdeæ*. *A'triplex Hálimus*.
44. *Lauríneæ*. *Láurus Benzòin Sássafras*, *nóbilis mas*, and *nóbilis fœm*.
45. *Thymelæèæ*. *Dáphne Laurèola*; L., yellow-variegated; *Mezudèrem*, red-flowered; M., purple-flowered; M., white-flowered; M., white-variegated-leaved.
46. *Elæágneæ*. *Elæágnus angustifòlia*; *Hippóphaë rhamnòides mas*, r. *fœm*.
47. *Euphorbiàceæ*. *Búxus sempervirens*, eight varieties of.
48. *Artocárpeæ*. *Ficus Cárica*, nine varieties of; a tenth kind of *Ficus*. *Morus nigra*, *álba*; a., “the small purplish-white” fruited; “the large-leav'd Virginian mulberry with black shoots,” “the Virginian mulberry with long red fruit.”
49. *Ulmàceæ*. The common elm; the witch hazel, or broad-leaved elm; the small-leaved elm; the smooth-leaved, or witch elm; the Dutch elm; the small-leaved striped elm; the striped witch elm; the yellow-leaved elm; the striped Dutch elm.

“There are some other varieties of these trees in the nurseries near London, which, not having been examined, we shall omit at present.” *Celtis occidentalis*.

50. *Juglâneæ*. *Juglans* *règia*, four varieties of; *nigra*. *Carya*, sp. (“Virginian walnut with long furrow’d fruit”), sp. (“the hickery, or white Virginian walnut”), sp. (“the small white Virginian Walnut, or hickery”).

51. *Salicîneæ*. *Salix* *álba*, *pentándra*, sp. (“the long-leav’d sweet [scented leaf] willow”), *babylónica*, *frágilis*, ? *amygdálina* or ? *triándra*, *vitellina*, ? *càprea*; ? *e.*, round-leaved; *e.*, round-leaved-variegated. *Pópulus* *canéscens*, *álba*; *a.*, variegated-leaved; *trémula*, *nigra*.

52. *Betulîneæ*. *Bétula* *álba*, *Alnus* *glutinòsa* and ? *oblongàta*.

53. *Cupuliferaæ*. *Quercus* *Ilex*, four varieties of; *coccifera*, *Ròbur*; *R.*, leaf white-variegated; *álba*, “evergreen oak with broad leaves like the common oak,” “the scarlet oak,” “the Virginian chestnut-leav’d oak,” “the Virginian willow-leav’d oak,” “the chinquapin oak,” “the Spanish cut-leav’d oak;” *Süber*, “the broad-leav’d cork tree;” *S.*, “the narrow-leav’d cork tree.” *Cárpinus* *Bétulus*; *B.*, striped-leaved; *Ostrya* *vulgàris* and *virginica*; *Castànea* *vésca*; *v.*, leaf elegantly variegated; *pùmila*; *Fàgus* *sylvática*; *s.*, yellow-variegated-leaved: *Córylus* *Avellàna*, and five varieties of it.

54. *Platàneæ*. *Plátanus* *orientàlis*, *occidentàlis*, ? *acerifolia*; *Liquidámbar* *styraciflua*.

55. *Myricææ*. *Myrica* *Gàle*, *cerifera*, *carolinénsis*.

56. *Coniferaæ*. *Cèdrus* *Libàni*, *Làrix* *europæa*: *e.*, with the rudiments of the cone white; *Pinus* *sylvéstris*, *Pináster*, sp. (“*Pinus*; *Americana*, *foliis* *prælongis*, *subinde* *ternis*, *conis* *plurimis* *confertim* *nascentibus*”), *Pinea*, *Stròbus*, *palústris*; *Abies* *excélsa*, *Picea*, sp. or var. (“*Abies*; *minor*, *pectinatis* *foliis*, *Virginiana*, *conis* *parvis* *subrotundis*. *Pluk. Alm. 2. Phyt. tab. 121. f. 1.*”); sp. or var. (“*Abies*; *Piceæ* *foliis*, *brevibus*; *conis* *minimis*. *Rand.*”); *nigra*; *balsamifera*, sp. or var. (“*Abies*; *taxi* *folio*; *fructu* *longissimo*, *deorsum* *inflexo*. *Long-con’d* *Cornish* *fir*”); *Schubértia* *dísticha*, *Cuprèssus* *sempervirens*; ? *s.*, *horizontàlis*; *Thùja* *occidentàlis*, *occidentàlis* with its leaves elegantly variegated; *Táxus* *baccàta*: *b.*, leaf variegated; *b.*, “the broad shining-leav’d yew;” *Juníperus* *commùnis*, *suécica*, *virginiàna*, ? *virginiàna* *hùmilis*; *bermudiàna*, *Sabina*; *S.*, variegated-leaved; ? *S.*, “the berry-bearing or upright savin.”

The introducers of foreign trees and shrubs in the early part of the eighteenth century are much indebted to Mark Catesby, an enthusiastic naturalist, who travelled in North America from 1712 to 1726, when he returned to England, made himself master of the art of etching, and published his splendid work, containing the natural history of Carolina, Florida, and the

Bahama Islands. All the most interesting trees of America are beautifully figured in these volumes; and the appearance of such figures for the first time, in England, must have greatly contributed to induce the wealthy to procure the introduction of the trees they represented into this country. Ryall who wrote the preface to Catesby's *Hortus Americanus Europæus* (which was not published till 1767, nearly twenty years after Catesby's death), observes, "that very little regard was had to the trees and shrubs of America on our first settling in that country; nor indeed, was any considerable step taken about introducing them into England till about the year 1720," chiefly in consequence of Mr. Catesby's exertions. Catesby lived many years at Hoxton, but in the latter part of his life he removed to Fulham, where he occupied a house, and had a garden within the site of what is now the Fulham Nursery, in which some trees remain that were planted with his own hand. *Lysous*, vol. ii. p. 829.) Catesby was born in 1679 and died in 1749. In a notice of his death, in the *Gentleman's Magazine* for 1749 (xx. 30.) he is called the "truly honest, ingenious, and modest, Mr. Mark Catesby."

On a blank leaf of Collinson's copy of Catesby's *Natural History of Carolina* (which, in January, 1835, came into the possession of A. B. Lambert, Esq.) is the following curious memorandum in Mr. Collinson's own handwriting, and signed with his name, "The ingenious author, Mr. Mark Catesby, was born of a gentleman's family at Sudbury in Suffolk. Some of his family being settled in Virginia, and having himself a turn of mind to natural history, he went over there to see his sister and improve his genius. From thence he travelled to Carolina, Bahama Islands, &c., and painted all the subjects from the life. On his return, the subscription being at an end, he was at a great loss how to introduce this valuable work to the world, until he met with a friend (Peter Collinson) to assist and promote his views. He learned to engrave, and coloured all himself, yet it proved so very expensive, that he was many years in accomplishing the work, being himself the principal operator. So noble and so accurate a performance, begun and finished by one hand, is not to be paralleled: but it afforded a subsistence to himself, his wife, and two children, to his death; and his widow subsisted on the sale of it for about two years afterwards, then the work, plates, &c, sold for 400*l*, and about 200*l*. more left by the widow, was divided between the two children, a son and a daughter." At the bottom of the titlepage is written:--- "This edition of this noble work is very valuable, as it was highly finished by the ingenious author, who in gratitude made me this present for a considerable sum of money I lent him without interest, to enable him to publish it for the benefit of

himself and family; else of necessity it must have fallen a prey to the booksellers." Date 1731.

Mr. John Ellis was remarkable for his exertions in devising plans for importing acorns and other nuts and seeds of American trees, which through his means were thus brought over in large quantities. He was a contemporary of Collinson, and, like him, was early in life engaged in merchandise; but he afterwards became agent for West Florida in 1764, and for Dominica in 1770. He had a very extensive correspondence, and was the means of introducing many articles of natural history, besides trees and shrubs. He was the author of *The Natural History of Corallines*, *The Natural History of Zoophytes*, &c., and established the genera *Halesia*, *Gordonia*, *Gardènia*, and others. As a proof of the amiable feeling that subsisted at that time between English and French naturalists, may be mentioned, that, during the war, Duhamel, who was then (1757) at the head of the French marine establishment, promised to Ellis and Collinson to return whatever plants were taken by the French. John Ellis died in 1776, aged 66.

Alexander Garden, M.D., was a Scotch physician, settled at Charleston, in South Carolina, where he married in 1755, and died in 1791, in his 52d year. He sent home a number of American trees and shrubs, including the *Ptèlea*, the fringe tree, several species of *Magnòlia*, *Zàmia integrifòlia*, &c. Linnaeus intended the loblolly bay, called *Lasiánthus* (now *Gordònia Lasiánthus*), to be named after him, which honour Dr. Garden solicited; but, unfortunately, his letter arrived too late by a month, Mr. Ellis having, in the meantime, named it *Gordònia*. Another genus, the *Gardènia*, commemorates the name of this ardent naturalist.

Subsequently to the year 1730, foreign trees and shrubs appear to have been planted in various country seats, and more especially in those laid out in the modern style. Among the earliest of these are included Stowe, and part of the scenery at Blenheim. At the former are some fine old cypresses, cedars, and acacias, planted in Brown's time; and in the latter were, till lately, the oldest deciduous cypresses and Lombardy poplars in England. We believe the very first place in which the Dutch style was made to give way to the English manner was Corby Castle, in Cumberland, which began to display the new taste so early as 1706 (*Warner*); but it does not appear that many foreign trees were planted.

Pains Hill was planted by the Honourable Charles Hamilton, sixth son of the Earl of Abercorn, about the same time that Woburn Farm was laid out, viz. 1735. Mr. Hamilton not only indulged the public with a sight of his improvements at Pains Hill; but allowed strangers the use of low chairs, drawn by

small horses, which were provided at the inns at Cobham, to go over the grounds. In the latter part of his life, Mr. Hamilton retired to Bath, having sold his place to Benjamin Bond Hopkins, Esq., who built the present house, the original one being small. (*Manning and Bray's Surrey*, ii. 768.) Among the trees remaining at Pains Hill are some remarkably fine silver cedars, pinasters, and other pines, American oaks, cork trees, and ilices, a tupelo tree (*Nýssa*), tulip trees, acacias, deciduous cypress, Lombardy and other poplars, &c. Here some of the first rhododendrons and azaleas introduced into England were planted by Mr. Thoburn, who was gardener to Mr. Hamilton, and who afterwards became an eminent nurseryman at Old Brompton. Bowood was laid out about the same time by the first Marquess of Lansdowne (then Earl of Shelburne), who was assisted by Mr. Hamilton of Pains Hill; and, like that place, it was planted with every kind of foreign tree that could be procured at the time. Many of these trees still remain, and have attained a large size: the cedars and tulip trees are remarkably fine. Woburn Farm, which began to be improved by Mr. Southcote in 1735, belongs to this class of places; and also Strathfieldsaye: the former contains one of the largest liquidambar trees in England, a remarkably fine hemlock spruce, very large tulip trees, acacias, hickories, pines, cedars, and cypresses, and a magnificent cut-leaved alder. At Strathfieldsaye are the largest hemlock spruce in England, some remarkably fine scarlet oaks, a large tupelo tree, and many fine pines and firs. Claremont, planted about the same time by Brown, for Lord Clive, contains a great many exotic trees, particularly cedars of large dimensions. There are very large ilices, cork trees, tulip trees, red cedars, a large hemlock spruce, and many other fine specimens of foreign trees. Oatlands, Ashley Park, and more particularly Lord Tankerville's at Walton, were planted soon after this period, and contain many fine specimens.

Upton House, near Stratford in Essex, was planted by Dr. Fothergill about 1762; and, though many of the shrubs were sold at the doctor's death in 1781, the grounds still contain many large and fine specimens. Of these we had the following measured in January, 1835: *Pópulus canadénsis*, 100 ft. high; *P. dilatàta*, 120 ft. high; *Quércus Turneri*, 50 ft. high; *Córylus Colúrna*, with a trunk 5 ft. in circumference, and forming a very handsome tree which bears abundantly every year; *Cuprèssus sempervirens horizontàlis*, 40 ft. high, a fine specimen: two very large cedars, with trunks $9\frac{1}{2}$ ft. in diameter, at 6 ft. from the ground; a large cork tree; *Kölreutèria paniculàta*, 40 ft. high, perhaps the largest in England; a large robinia, &c. Collinson states that the *Arbutus Andráchne* flowered for the first time in this garden, in May, 1766. He adds that the plant was

raised from seeds sent to Dr. Fothergill, by Dr. Russell of Aleppo, in 1756; and that the original plant was sold by auction in August, 1781, after the doctor's death, for 53*l.* 11*s.* There appears to be some mistake in this relation, as Dr. Fothergill did not purchase Upton till 1762, and the *Arbutus Andrachne* was cultivated in 1724. Dr. Fothergill, however, may have raised his plant somewhere else, and removed it to Upton; and, though it was introduced in 1724, it may not have flowered before the period mentioned. Collinson was such a careful observer, that this remark appears due to his memory. Upton House is now in the possession of S. Gurney, Esq.

Purser's Cross was planted by John Ord, Esq., in 1756; and "it is not a little extraordinary," says Lysons, "that this garden should, within the space of little more than fifty years (such have been the effects of good management and a fertile soil), have produced trees which are now the finest of their respective kinds in the kingdom." The following is an account of some of the most remarkable trees at Purser's Cross, as measured by Lysons at three different periods, and for us in Jan. 1835:—

	Girth in							
	1793.		1808.		1809.		1835.	
	ft.	in.	ft.	in.	ft.	in.	ft.	in.
The <i>Sophora japonica</i> , planted in 1756, being then about 2 ft. in height; it flowered for the first time in August, 1807, and has continued to flower almost every year since	8	0	9	4	9	7½	13	0
The ginkgo tree (<i>Salisburia</i>), planted in 1767 (about 37 ft. high in 1809; and in 1835 between 50 ft. and 60 ft.)	2	3	3	6	3	9	5	2
A tree, the seed of which was given to Mr. Ord, by the late Mr. Aiton, as an Illinois nut, and which was sown in 1760 (about 40 ft. high)	2	2	2	10	2	11	4	5
A black walnut tree (<i>Juglans nigra</i>) sown where it now stands, in 1757 (about 64 ft. high)	5	4	6	11	7	3	9	0
A cedar of Lebanon planted in 1756, then two years old (in 1809 about 55 ft. high)	8	8	9	11	9	9	11	10
A willow-leaved oak (sown in 1757)	4	0	5	5	5	7	7	0
The <i>Rhus vernix</i> , or varnish sumach	4	0	4	10	4	10	—	—

Purser's Cross contains a greater number of fine specimens, in a very limited space, than any garden we know of in the neighbourhood of London. In October, 1834, we found there *Magnolia tripetala*, *acuminata*, and other species, of considerable size, *Liriodendron Tulipifera*; *Negundo fraxinifolium*, 40 ft. high; *Asimina triloba*, 10 ft. high, flowering every year; another plant, which died a few years ago, having ripened fruit every year; *Ailantus glandulosa*, 30 ft. high, *Gymnocladus canadensis*, 30 ft. high; *Sophora japonica*, 40 ft. high, which flowers every year; *Robinia* and *Gleditschia*, very fine specimens; *Cratægus*, several species, very large; *Pyrus Sorbus*, very fine specimens; *Celtis*,

Juglans, and *Pópulus*, very large trees; *Quércus* of various species, from 40 ft. to 60 ft. high; *Quércus coccífera* and *gramúntia*, each 30 ft. high, and considered among the finest specimens in the neighbourhood of London; and *Salisbùria adiantifòlia*, nearly 60 ft. high; *Andrómeda arbòrea*, 18 ft. high; and deciduous cypresses, from 70 ft. to 80 ft. high. Purser's Cross is now the property of Lord Ravensworth.

Syon was one of the largest monasteries that were suppressed. It was in Henry VIII.'s hands at his death; and his funeral procession, which is said to have exceeded in magnificence anything of the kind either before or since, was rested a night at Syon on its way to Windsor. King Edward VI. granted Syon to Edward Duke of Somerset, who built the shell of the present mansion. He had a botanic garden there, mentioned by Turner (who was his physician) in his *Herbal*. In 1604, we find Syon House in the possession of Henry Earl of Northumberland, who had laid out 9000*l.* on the house and gardens. The house was afterwards greatly enlarged and improved by Inigo Jones, in 1659. The grounds at Syon are generally understood to have been laid out in their present form by Brown, between 1750 and 1760. They were planted with all the foreign hardy trees and shrubs that could be procured, at that time, in the London nurseries; and the place now contains many very fine old specimens of cedars, pines, planes, gleditschias, robinias, catalpas, and more especially of deciduous cypress.

George William, sixth Earl of Coventry, succeeded to the title, and to the estate of Croome d'Abitot, in the year 1738, being then 17 years of age. He soon afterwards, with the assistance of Brown, began to improve the estate, at that time "a mere bog, and a barren waste" (*Dean's Croome Guide*, 1824, p. 37.), and soon converted it into fertile soil, and planted it with all the useful and ornamental trees and shrubs at that time to be procured in the nurseries. The plants have grown with astonishing vigour, and there is now at Croome an extensive collection of species, containing some of the finest specimens of foreign trees and shrubs in the country.

Numerous gentlemen's seats, planted about this time in every part of England, might be cited as containing fine old specimens of foreign trees and shrubs; but we must limit ourselves to a few which took a lead in this taste. Among these may be mentioned, in addition to those already noticed, Busbridge, near Godalming, in Surrey, in 1751, in the possession of Philip Carteret Webb, Esq., and frequently mentioned by Miller; Mount Edgecombe, Earl of Mount Edgecombe; Mamhead, now belonging to W. Newman, Esq.; Powderham Castle, Earl of Devon; High Clere, Earl of Caernarvon; and Chiswick, Duke of Devonshire. There are, doubtless, many places as much or more worthy of

being quoted than several of those named; but, as we have invited all proprietors and gardeners in the British Isles to send us accounts of their foreign trees and shrubs for this work, and as many of these have done so, we must refer in this place to the paragraph headed *Statistics*, given to each tree and shrub; where, under each county, will be found the names of all those seats most remarkable for foreign trees and shrubs, with the dimensions and other particulars of the plants they contain.

Several botanic gardens were formed during this century, both at home and abroad; and the exchange of seeds and plants which takes place universally among such establishments increased the foreign productions of each respective country. It also became the practice, in the latter part of this century, for private persons and public bodies to send out botanical collectors. Several of these were sent out from the Royal Gardens at Kew, others by the subscriptions of individuals, and some by nurserymen.

Chelsea Garden (already noticed, p. 47.) is said by Collinson to have been, in his time, the richest in plants in Europe. It was brought to the highest degree of eminence during this century by Miller. Its origin is unknown: the first notice of it, in the books of the Apothecaries' Society, is in 1674, when it was proposed to wall it round; and two years afterwards, in 1676, the Society agreed to purchase the plants growing in Mrs. Cape's garden at Westminster. They may probably also have had plants from the garden mentioned in Evelyn's *Diary* for 1658 as "the medical garden at Westminster, well stored with plants, under [Edward] Morgan, a skilful botanist." Piggot is the name of the first curator of the Chelsea Garden, noticed in 1676. Watts, mentioned both by Ray and Evelyn, was an apothecary by profession, but undertook the care of the garden in 1680, at 50*l.* per annum. Miller was appointed to the garden in 1722, at the time Sir Hans Sloane, when applied to for a renewal of the lease of the garden, granted it to the Society in perpetuity, at a rental of 5*l.* per annum, and on condition that specimens of fifty new plants should annually be furnished to the Royal Society, till the number amounted to two thousand, that number, at that time, being supposed likely to exhaust the botanical riches of the whole world. Miller resigned his situation as curator, a short time before his death in 1771, and was succeeded by Forsyth, who left it to become royal gardener at Kensington in 1784, and was succeeded by Fairbairn, who died in the garden in 1814. His situation is now filled by Mr. William Anderson, F.L.S. H.S., &c., who has greatly enriched the garden, and contributed materially to its present high character.

The botanic garden at Kew was established in 1760 by the Princess Dowager of Wales. A catalogue was published in 1768 by Dr. Hill; and a more scientific one by Mr. William Aiton in 1789, a second edition of which appeared in 1810. William Aiton died in 1793, aged 62. He was some time assistant to Philip Miller, at Chelsea, and was recommended to the princess dowager in 1759. In 1783 he was appointed to the care of the pleasure-grounds and kitchen-garden at Kew. The *Aitonia* is named after him. He was succeeded by his son, Wm. Townsend Aiton, the present royal gardener there. Kew is more especially interesting to the planter of trees, from its arboretum having been one of the very first that was formed in Britain; and, though many of the species are now lost, and it does not contain more than a fourth part of what are to be found in the Horticultural Society's garden and in the arboretum of the Messrs. Loddiges, there are still existing there many fine specimens. Dr. James Sherard's botanical garden at Eltham, in which he was assisted by Dillenius, was established in the first years of this century, but declined at Dr. Sherard's death in 1737; and, in 1795, nothing remained of it but a fine cedar of Lebanon close to the house, and a few other trees and shrubs. This cedar measured, at the above period, 9 ft. in circumference, at 3 ft. from the ground; and in 1801 it had increased in circumference $6\frac{1}{4}$ inches. (*Lysons.*) Dr. James Sherard was the brother of Dr. William Sherard, an eminent botanist, and author of several works, who was travelling tutor for many years to several English nobleman, and afterwards British consul at Smyrna, near which he had a fine country house and garden, from which he sent home many seeds and plants. This brother founded the botanical professorship at Oxford, and gave to that establishment his botanical library, and his herbarium. He was the patron of Mark Catesby and of Dr Dillenius.

Mr. William Curtis, author of the *Botanical Magazine*, first established a small botanic garden at Bermondsey. In 1771 he formed one on a more extensive scale at Lambeth Marsh. In 1789 he removed his plants to Brompton, where he died in 1799, aged 53 years. His partner, and successor, Mr. William Salisbury, removed this garden to Cadogan Place, Sloane Street, where an arboretum was planted, and the grounds are now (1835) occupied as a subscription garden and as a nursery.

A private botanic garden was founded at Twickenham about 1789, by William Swainson, the proprietor of some popular vegetable medicines. It contained every tree and shrub that could be procured at the time in the British nurseries, and was kept up in the very first style of order and neatness till Mr. Swainson's death in 1806. It is now the property of Mrs. Canham, and is managed by Mr. Robert Castles, an enthusiastic lover of plants, and an excellent man.

Various other gardens might be mentioned as having contributed to spread a taste for foreign trees and shrubs towards the latter end of this century; and the names of a number of writers on botany and gardening would also deserve commemoration here, were they not already recorded in the historical part of our *Encyclopædia of Gardening*. The reader who thinks we might have extended this part of our work will bear this in mind; and also that it has been our object, throughout this Introduction, to repeat nothing which we have already laid before the public.

The only extensive nursery at the beginning of this century was, as we have before seen (p. 46.), that of Brompton Park, occupied by London and Wise. Those of Gray of Fulham, of Furber of Kensington, of Fairchild of Hoxton, Gordon of Mile End, and Hunt of Putney, became eminent before the middle of the century; and those of Lee and Kennedy, William Malcolm, Russell, Loddiges, and others, were large establishments before the end of it.

Gray commenced his nursery at Fulham early in the 18th century. He received many American trees and shrubs from collectors and resident amateurs in America, and enriched his stock at the sale of Dr. Compton's trees. In 1740, he published a catalogue of his plants, which is said to have been written by Philip Miller. In the preface to Catesby's *Hortus Europæus Americanus*, which is dated 1767, it is said, that "Mr. Gray at Fulham has, for many years, made it his business to raise and cultivate the plants of America, from whence he has annually fresh supplies, in order to furnish the curious with what they want;" and that, "through his industry and skill, a greater variety of American forest trees and shrubs may be seen in his gardens, than in any other place in England." This nursery is now in the possession of Messrs. Whitley and Osborne, and still retains its reputation for American trees and shrubs. It also contains some fine old specimens of the trees planted by Gray. Among these are *Quercus Süber*, *Celtis occidentalis*, *Ailantus glandulosa*, *Laurus Sassafras*, *Kœlreutèria paniculata*, *Diospyros virginiana*, and various others. The first *Magnolia grandiflora* which was brought to England (as generally supposed) was planted in this nursery, and all the old trees of the kind in the country are said to have been propagated from it. The tree died about 1810; but its trunk, which measures 4 ft. 10 in. in circumference, was, till very lately, preserved. The branches extended over a surface 20 ft. in diameter, it was as many feet high, and in the blossoming season, which lasted generally two or three months, it perfumed the whole neighbourhood. It was surrounded by stages from the ground to its summit, on which were placed pots containing layers for

propagation. It was the number of these, and the exhaustion they occasioned, which killed the tree.

Furber, mentioned by Collinson, was a nurseryman at Kensington, and one of those gardeners who formed a society for publishing a work on gardening, of whose *Catalogue* some account is given in p. 60. Miller was secretary of this society, which, as it is said, dissolving through difference of opinion, the papers became Miller's and led to the publication of his *Dictionary*. Furber's grounds are now partly built on, and the remainder forms part of Messrs. William Malcolm and Co.'s nursery.

Thomas Fairchild had a nursery and an excellent vineyard. For the time in which he lived, he was a scientific gardener, and distinguished himself by a paper, in the *Royal Society's Transactions* (vol. xxxiii. p. 127.), "On the different, and sometimes contrary, Motion of the Sap in Plants." He introduced various new trees and shrubs from the Continent of Europe and North America, as will be seen by the list at the end of this section. He was author of the *City Gardener*. He died in 1729, and left funds for a botanical sermon, to be delivered annually on Whitsun Tuesday, at St. Leonard's, Shoreditch. The legacy left by Fairchild produced a guinea a year, but this sum being thought insufficient, a subscription was entered into, the produce of which has raised the annual sum to three guineas. These sermons were preached for many years by Dr. Colin Milne, author of the *Botanical Dictionary*, by whom they were published in 1779. The sermon is now preached annually by the Rev. William Ellis, of Merchant Tailors' School. Some curious details respecting this legacy will be found in Henry Elles's *Account of the Parish of St. Leonard's, Shoreditch*.

James Gordon, nurseryman at Mile End, London, who had previously been gardener to Robert Lord Petre, is thus spoken of in a letter from Ellis to Linnæus, dated April 25, 1758:—

"If you want a correspondent here that is a curious gardener, I shall recommend you to Mr. James Gordon, gardener at Mile End, London. This man was bred under Lord Petre and Dr. Sherard, and knows systematically all the plants he cultivates. He has more knowledge in vegetation than all the gardeners and writers on gardening in England put together; but he is too modest to publish anything. If you send him any thing rare, he will make you a proper return. We have got a rare double jessamine (*Gardènia flórida*) from the Cape, that is not described: this man has raised it from cuttings, when all the other gardeners have failed in the attempt. I have lately got him a curious collection of seeds from the East Indies, many of which are growing, but are quite new to us. He has got the ginkgo (*Salisbúria*), which thrives well, and, when he has in-

creased it, he will dispose of it." (*Smith's Cor.*, vol. i. p. 93.) Gordon commenced his nursery at Mile End in 1757; he disposed of it in 1776, to his sons James and William, and died in 1780. In 1781 we find this nursery in the possession of James Gordon, Thomas Dermer, and Archibald Thompson. James Gordon died in 1794, and Thomas Dermer in 1799, when Archibald Thompson came into possession of the whole. Mr. Thompson died in 1832, and the business is now carried on by his son James. Only a small part of the original ground is now (1835) occupied as a nursery; but in the part that remains there are some fine old specimens. What is believed to be the oldest *salisburia* in England, the ginkgo tree above alluded to, is 5 ft. 5 in. in circumference at 1 ft. from the ground, and 55 ft. high. There are venerable specimens of magnolias, *Laurus Sassafras*, the cork tree and other oaks, the liquidambar both species, ailantus, gymnocladus, and many others, with some of the largest plants of green tea growing in the open air in England. A list of the more remarkable of these trees and shrubs, with their dimensions taken in 1831, will be found in the *Gard. Mag.* for the following year, vol. viii. p. 250.; and subsequent measurements of several of them taken in January, 1835, will be found in the same magazine, vol. xi.

Of the Hunts of Putney we know little, except that their names appear among the authors of the *Catalogue* of the trees and shrubs grown in the London nurseries in the year 1730. The nursery at Putney was little known in our time, and the stock was sold off, and the ground advertised to be let for building on, in December, 1834.

James Lee was born at Selkirk in 1715, and, about twenty years afterwards, walked to London. When he was at Litchfield he was seized with the small-pox, and detained there some time. When he recovered, and came to London, he was employed at Syon, and afterwards at Whitton by the Duke of Argyle. About the year 1760 he entered into partnership with Lewis Kennedy, gardener to Lord Bolton, at Chiswick, and commenced a nursery in what was called the Vineyard, at Hammersmith. At the beginning of the last century, this vineyard produced annually a considerable quantity of Burgundy wine. A thatched house was built in the grounds, the upper part occupied as a dwelling-house and for selling the wine, and underneath were the wine-cellars. Lee was patronised by the Earl of Islay (afterwards Duke of Argyle), the planter of Whitton, who died in 1761; and other noblemen: he corresponded with Linnæus, and composed an *Introduction to Botany*, according to his system, published in 1760, which for many years was in the highest repute. He died in the year 1795, at the age of 80 years; his partner, Kennedy, having died previously. The nursery was carried on

by the sons of the two founders, till 1818, when they dissolved partnership. It then became the sole property of James Lee, the second of that name, who died in 1827, leaving it to his family, and it is now (1835) carried on by his son John. For many years this nursery was deservedly considered the first in the world. Besides an extensive correspondence, and a vigilant attention to procure every new plant as soon as it was introduced by others, Messrs. Lee and Kennedy introduced many plants into the country, through collectors whom they had sent abroad, and through foreign botanists. They maintained a collector in America, who sent home several new oaks; and, in partnership with the Empress Josephine, one at the Cape of Good Hope, who sent home many new ericas, ixias, and other Cape plants. They had also a collector in South America, who sent home the *Fúchsia coccínea*, by which they made a considerable sum of money, selling it for some time at a guinea a plant. They also had the first China rose in 1787, of which they made a large sum. The extent of this nursery has been somewhat curtailed by the approach of London; but it still contains an excellent collection, some fine specimens of magnolias, asiminas, cratæguses, *Pÿrus Sórbus*, and other foreign trees and shrubs, and is conducted with the greatest liberality.

The nursery of Messrs. Malcolm and Co. was established about the middle of the century, first at Kennington, and afterwards at Stockwell, and was at one time one of the most extensive in the neighbourhood of London. The ground has long since been built on; but one of the descendants of the family, as before observed, occupies, with other grounds, part of the nursery which was Furber's, at Kensington.

The nurseries of Russells, at Lewisham; of Bassington, at Kingsland; of Cormack, at New Cross; of Ronalds, at Brentford; and a number of others, some of which are now extinct, and others more eminent than before; were all more or less celebrated during the latter part of this century.

The nursery at Hackney was established about the middle of the century by John Busch (in the *Hortus Kewensis*, erroneously spelt Bush), a German gardener, who, entering into the service of the Empress Catherine in 1771, was succeeded in his nursery, in that year, by Conrad Loddiges, also a native of Germany. This nursery soon became celebrated for the introduction and propagation of American trees and shrubs, particularly magnolias, rhododendrons, and azaleas. It will be noticed more at length in the succeeding section.

The botanical and horticultural authors and garden artists of England during this century, who contributed to the spread of a taste for foreign trees and shrubs by their writings and practice, are numerous. We have already mentioned Brown, to whom

we may add, among artists and authors, Switzer, a seedsman, at "the Flower Pot over-against the Court of Common Pleas in Westminster Hall, or at his garden in Millbank, Westminster," author of *Iconographia Rustica*, and several other works, and the designer of various gardens both in Britain and Ireland; Bradley, a voluminous author; Batty Langley, an architect at Twickenham, who wrote *New Principles of Gardening*, &c., and *The true Method of improving an Estate by Plantations of Timber Trees*, &c.; Dr. John Hill, a voluminous gardening author; Sir William Chambers, who wrote *Dissertations on Oriental Gardening*; Wheeler, a nurseryman at Gloucester, and author of *The Botanist's and Gardener's New Dictionary*; Abercrombie, a well known voluminous author; Weston; Speechley; Dr. A. Hunter, the editor of Evelyn's *Sylva*; Meader, gardener to the Duke of Northumberland; Græfer, gardener to the Earl of Coventry at Croome, and afterwards to the King of Naples at Caserta; and a great number of others. We should have been tempted to submit some notices of these authors and their works, did we not expect a *Chronological and Biographical History* of them from the accurate and learned pen of William Forsyth, Esq. To this gentleman, who has been for many years collecting information respecting trees and shrubs, we are deeply indebted for many corrections and additions to this chapter, and to our work generally.

In order to give a general summary of the trees and shrubs introduced into Britain during the eighteenth century, we shall divide it into periods of ten years; and give in each the names of some of the principal plants introduced, and those of their introducers, according to the *Hortus Kewensis*, and to some farther information on the subject, kindly furnished to us by Mr. Forsyth, and Messrs. Loddiges of Hackney.

From 1701 to 1710 inclusive (Queen Anne), four trees and five shrubs were introduced. Among the trees were, *Fraxinus lentiscifolia* from Aleppo, and *Ptelea trifoliata* from North America, by the Rev. John Banister. *Cratægus parvifolia* was introduced by Bishop Compton and *Ribes oxyacanthoides* by Mr. Reynardson of Hillington. *Bignonia capreolata*, a beautiful climber, and *Colutea cruenta*, a handsome flowering shrub, were introduced during this period.

From 1711 to 1720 (Anne and Geo. I.), three trees and nine shrubs were introduced: among these were, *Pavia rubra*, by Thos. Fairechild; *Pinus Tæda*, and *Ceanothus americanus*, by Bishop Compton; *Cerasus Mahaleb*, from Austria; and *Comptonia asplenifolia*, *Lycium afrum*, and *Iva frutescens*, by the Duchess of Beaufort.

From 1721 to 1730 (Geo. I. and II.), twenty-two trees and twenty shrubs were introduced. Mark Catesby introduced

Gleditschia monosperma, *Catalpa syringifolia*, *Calycanthus floridus*, *Crataegus flava*, and *Amorpha fruticosa*. Mr. Moore introduced *Fraxinus americana*; Robert Furber, *Ostrya vulgaris*, *Pyrus coronaria*, *Platanus acerifolia*, *Quercus alba*, *Rhus radicans*, and *Viburnum laevigatum*. Dr. James Sherard introduced *Arbutus Andrachne Rhus glabra*, and *Rosa caroliniana*. Sir Charles Wager introduced *Acer dasycarpum*; Miller, *Clematis crispa*, *Santolina viridis*, and some others, which are recorded as being cultivated in the Chelsea Garden during this period. Among the species introduced or recorded, but without the name of the introducer, are, *Cerasus virginiana*, *Tilia pubescens*, *Alnus oblongata*, *Carya compressa*, *Cercis canadensis*, *Quercus graminifolia*, *Q. sericea*, and *Q. prinus*, *Pinus palustris*, *Euonymus latifolius*, *Caprifolium gratum*, *Rhus elegans*, *Wisteria frutescens*, and a number of others.

From 1731 to 1740 (Geo. II.), twenty-four trees and forty-five shrubs were introduced. Mr. Stephen Bacon introduced the *Clèthra alnifolia*; Thomas Fairchild, the *Cornus florida*; Miller no fewer than thirty species, including *Crataegus cordata*, *Populus angulata*, *Acer monspessulanum*, *Carpinus orientalis*, *Celtis Tournefortii*, *Platanus cuneata*; *Quercus Ægilops*, *nigra*, *rubra*, and *virens*; *Pinus inops* and *variabilis*; and several others. Collinson, between 1734 and 1739, introduced twenty-six species, among which we find *Magnolia acuminata* in 1736; the first azaleas that were in the country, namely, *Azalea nudiflora*, *viscosa*, and *glauca*; *Kalmia latifolia* and *angustifolia*, *Andromeda mariana* and *racemosa*, *Rhododendron maximum*, *Chionanthus virginica*, *Acer saccharinum*, *Cephalanthus occidentalis*, *Nyssa denticulata*, several species of *Viburnum*, and that beautiful tree, *Larix pendula*. Sir John Colliton had in cultivation the *Magnolia grandiflora* from Carolina in 1734, and the lanceolate-leaved variety in 1737. Dr. James Sherard introduced *Menispermum virginicum*; and Dr. Thomas Dale, *Philadelphus inodorus*.

From 1741 to 1750 (Geo. II.), there were introduced eight trees and twelve shrubs. Sir John Colliton had the *Robinia hispida* in cultivation before 1743. Dr. Amman introduced the *Cytisus austriacus*; Richard Bateman, the *Acacia Julibrissin*; Christopher Gray of Fulham, the *Pyrus angustifolia*. Catesby introduced *Stuartia virginica*; and Archibald Duke of Argyll, the *Pinus Cembra*, *Gymnocladus canadensis*, *Acer montanum*, *Betula papyracea* and *populifolia*, *Crataegus punctata* and *glandulosa*, *Itea virginica*, *Corylus rostrata*, *Amelanchier Botryopium*, *Andromeda calyculata*, and that curious miniature tree, *Dicra palustris*.

From 1751 to 1760 (Geo. II.), twenty-seven trees and forty-seven shrubs were introduced. Peter Collinson introduced

Broussonètia papyrifera from Japan in 1751. Father D'Incarville introduced the *Ailántus glandulòsa* from China, also in 1751; Jas. Gordon of Mile End, the *Ulmus americana* in 1752: that remarkable tree the *Salisbùria adiantifòlia* was cultivated by him in 1754; the parent tree, a male, still exists (see p. 78.), and from it, in all probability, originated all the male trees of the same species, not only in Europe, but in North America; he introduced the *Sophòra japònica* in 1753, and the *Còrnus alternifòlia* in 1760. Archibald Duke of Argyle introduced the *Làrix microcàrpa* and the *Smilax rotundifòlia* in the same year. John Ellis introduced *Halèsia tetràptera* and diptera in 1756 and 1758; Messrs Kennedy and Lee *Euónymus atropurpùreus*; Hugh Duke of Northumberland, *Pinus resinòsa*; Christopher Gray, *Vibúrnum nitidum*. The Duke of Bedford cultivated *Pinus rígida* before 1759; and *Pópulus dilatàta*, the Lombardy poplar, was introduced from Italy by the Earl of Rocheford in 1758. No fewer than fifty articles were introduced or cultivated by Miller during this decade. Among these are, *Acer créticum*, in 1752, probably the small tree still existing in the Chelsea Garden; *A. Opulus*, heterophýllum, and tatáricum; *Dáphne Cneòrum* and *póntica*, *Lonicèra tatárica*, *Magnòlia tripétala*, several species of *Rhámnus*, *Thùja occidentàlis*, *Tília americana*, *Abies rùbra*, *Pinus marítima* and several others, *Bétula lénta*, *Pýrus prunifòlia*; *Cotoneáster tomentòsa*, *Dáphne alpina*, *Liquidámbar imbèrbis*. Among the trees and shrubs recorded in the period, without the name of the introducer, are, *Acer pennsylvánicum*, *Bérberis canadénsis*; *Cérasus caroliniàna*, a beautiful sub-evergreen low tree from Carolina, too much neglected in England; *Ròsa sínica*, *Shephèrdia canadénsis*, *Plánera Richárdi*, and *Oxycóceus macrocàrpus*.

From 1761 to 1770 (Geo. III.), twelve trees and forty shrubs were introduced. Jas. Gordon introduced, or had in cultivation, *Tília álba*, *Bétula excélsa*, *Clématis virginiana*, *Vibúrnum cassinòides* and *Lentàgo*, *Hypéricum alàtum*, and *Euónymus verrucòsus*. John Bartram introduced *Mitchèlla rèpens*; John Busch, *Lèdum palustre*, *Fothergílla ainifòlia*, *Xanthorhiza apiifòlia*; Mr. Bannet, *Lèdum latifòlium*; George William Earl of Coventry, *Kölreutèria paniculàta* from China, *Erica austràlis*, and *Sàlix retùsa* from Italy. John Greening cultivated *Pàvia flàva*; Joseph Brooks, *Erica strícta*; John Cree, *Bumèlia ténax*; Dr. Fothergill, *Pópulus heterophýlla*; Messrs. Kennedy and Lee, *Cratægus elliptica*, *pyrifòlia*, and that fine tree, *Fágu ferugínea*. Sir Joseph Banks introduced *Rhodóra canadénsis* in 1767; John Ord, *Genísta tríquetra*; Peter Collinson, *Alnus serrulàta*, and *Vaccínium virgàtum*; Hugh Duke of Northumberland, *Pópulus græca* and *levigàta*; and Miller, *Sambùcus canadénsis*, *Genísta púrgans* and *Rùbus hispídis*. The

Duchess of Portland introduced *Vaccinium frondosum*. Among the plants respecting which merely the dates at which they were introduced to, or first cultivated in, Britain, are recorded in the *Hortus Kewensis*, are, *Gaultheria procumbens*; *Rhododendron ponticum*, introduced, we are informed, by Conrad Loddiges, who sold the first plant to the Marquess of Rockingham, a noble encourager of botany and gardening; *Andromeda axillaris*, coriacea, and acuminata; *Styrax grandifolium* and *laevigatum*, *Kalmia glauca*, and that delightful shrub, *Chimonanthus fragrans*. The last, we are informed, was first cultivated by the Earl of Coventry at Croome.

From 1771 to 1780 (Geo. III., during the American war), were introduced eight trees and forty-eight shrubs. Mrs. Primmet introduced *Genista lusitánica*; Mons. Richard, *Ulmus pumila*, *Caragana Chamlagu*, and *Caprifolium impléxum*; Sir Joseph Banks, *Salix myrtillóides* from Sweden; Dr. Solander, *Spiræa laevigata* from Siberia; Dr. Hope of Edinburgh, *Pópulus cándicans* and *monilifera*. Messrs. Kennedy and Lee introduced *Aristotèlia Múequi*, *E'phedra monostachya*, *Búddlea globosa*, *Gleditschia hórrida*, *Rhámnus alnifolius*, and others. The celebrated botanist and traveller, Pallas, introduced *Pýrus salicifolia* in 1780, *Diótis ceratóides*, and *Calligonum Pallásia*. Dr. Nicholas Jacquin introduced *Cýtisis capitatus*, and *Drýpis spinosa*; Dr. Pitcairn, *Vaccinium dumosum*; Mr. William Malcolm, *Gordonia pubescens*; Mr. William Young, *Vaccinium staminium*; John Earl of Bute, *Genista germánica*; Hugh Duke of Northumberland, *Caragana spinosa*; Dr. Fothergill, that beautiful tree, *Pýrus spectabilis*, *Búxus baleárica*, and *Clématis flórida*. *Salix incubacea* and *Genista decumbens* were introduced by Drs. Fothergill and Pitcairn about the same time. Francis Masson introduced *Vaccinium Arctostáphylos*. Benjamin Bewick introduced *Vaccinium angustifolium*.

From 1781 to 1790 (Geo. III., intercourse with America being restored), sixteen trees and thirty-five shrubs were introduced, according to the *Hortus Kewensis*. John Bell introduced *Viburnum dáuricum*, *Bétula dáurica*, and *Caragana Altagana*. John Busch introduced *Ribes Diacantha* and *Ulmus incana*; John Græfer, *Pýrus bollwylleriána* and *baccata*, and that valuable evergreen, *Aúcuba japónica* (female). William Forsyth cultivated *Pinus Banksiána* in 1785; William Young, *Fráxinus juglandifolia*; and Daniel Grimwood, *Fráxinus pubescens*. The Hudson's Bay Company introduced *Ulmus undulata*; John Fraser, *Magnòlia auriculata*, *Rhododendron punctatum*, and *Quércus lyrata imbricaria*, and *rotundifolia*; Sir Joseph Banks, *Hydránga Horténsia*, *Magnòlia conspicua*, *Pæónia Moutan*, *Ròsa índica*, *Bérberis sibírica*, and some vacciniums. Gilbert Slater introduced *Ròsa semperflórens* in 1789; and the celebrated Professor Thouin,

the first horticulturist of his day, Nitrària Schóberi in 1788. The following ample list was introduced by Conrad Loddiges during this period ; the names having been supplied to us by the present Messrs. Loddiges, his sons :—*Mòrus* *tatàrica and pennsylvànica ; *Córnus* *circinàta, *Genísta* *sibírica, *Rhododéndron* *Chamæcístus, *Acer* híbridum and trilobàtum, *Prúnus* dasycárpa, *Bérberis* dàurica, *Cratægus* *Oxyacántha* rósea ; *Azàlea* speciòsa crísps, nudiflòra rùbra, and nudiflòra stamínea ; *Bétula* sibírica, *Amýgdalus* sibírica, *Andrómeda* calyculàta var. ventricòsa, *Alnus* pùmila, *Córnus* sibírica, *Ribes* trilòrum, *Caragàna* fèrox, *Ròsa* aciculàris and corymbòsa, *Thùja* plicàta and tatàrica. Of these species introduced by Conrad Loddiges, those marked with a star were received by him from the celebrated botanist and collector for the French government, André Michaux ; almost all the others were received from William Bartram of Kingsessing, Pennsylvania.

From 1791 to 1800 (Geo. III.), were introduced nineteen trees and fifty shrubs. John Bell introduced *Juníperus* dàurica, and *Azàlea* póntica. William Forsyth introduced *Bérberis* ilicifòlia ; Sir George Staunton, *Ròsa* bracteàta ; John Busch, *Caragàna* jubàta, and *Rhododéndron* chrysánthum ; Messrs. Lee and Kennedy, *Ròsa* fèrox ; Messrs. Fraser, *Quércus* tríloba, tinctòria, palústris, and Banísteri. Conrad Loddiges introduced *Atragène* austriaca and americàna, *Cýtisus* *purpùreus, *Andrómeda* *Catesbæ'i, *Aràlia* hispida ; and also, according to Messrs. Loddiges, *Castànea* americàna, *Cýtisus* *supínius, *Juníperus* alpina and sibírica, *Prínos* lanceolàtus and lævigàtus, *Spiræa* canadénsis ; *Vaccínium* *buxifólium, elevàtum, hispídulum, pùmilum, ligústrinum ; *Vítis* vulpina álba, v. nìgra, and v. rùbra. Among the plants recorded in the *Hortus Britannicus*, as having been introduced in this decade, are, *Càrya* porcina and amàra, *Pýrus* spùria, *Magnólia* macrophýlla, *Andrómeda* speciòsa, *Ròsa* suavèolens, *Prúnus* marítima, *Pýrus* auriculàta, *Quércus* microcárpa, and several others.

Of the nearly 500 hardy trees and shrubs introduced during this century, 108 are from the continent of Europe, 300 are from North America, 3 from Chili, 13 from China, 6 from Japan, 2 from the Cape of Good Hope, 33 from Siberia, 2 from Tartary, 1 from Egypt, 2 from Morocco, 1 from Aleppo, 1 from Barbary, and the remaining few chiefly from Asia.

In the early part of the century, the greater number of species appear to have been received by Peter Collinson, from Dr. Garden of Charlestown, John Bartram, Mark Catesby, and other collectors. The progress of introduction was interrupted during the eighth decade of the century (1771 to 1780), owing to the breaking out of the American war ; but it revived with double vigour between 1780 and 1800, during which period by far the

greater number of trees and shrubs introduced were received by Conrad Loddiges, and chiefly from William Bartram, the son of John. The Bartrams, indeed, and André Michaux, were the great collectors of American plants during the 18th century. Michaux sent almost every thing to France, by the government of which he was employed; but the Bartrams were Americans, and corresponded chiefly with the Kew Botanic Garden, and with the London nurserymen and amateurs. A number of trees and shrubs were introduced during the 18th century by John Fraser, but the chief accessions to the British arboretum and fruticetum made by this indefatigable collector were in the succeeding century.

John Bartram, one of the most distinguished of American botanists, was born in Chester County, Pennsylvania, in 1701. His grandfather, of the same name, accompanied William Penn to this country in 1682. John Bartram was a simple farmer; he cultivated the ground for subsistence, while he indulged an insatiable desire for botany. He was self-taught in that science, and in the rudiments of the learned languages, and medicine and surgery. So great, in the end, was his proficiency in his favourite pursuit, that Linnæus pronounced him "the greatest natural botanist in the world." He made excursions, in the intervals of agricultural labour, to Florida and Canada, herborising with intense zeal and delight. At the age of 70, he performed a journey to East Florida, to explore its natural productions; at a period, too, when the toils and dangers of such an expedition far exceeded those of any similar one which could be undertaken at the present time, within the limits of the United States. He first formed a botanic garden in America, for the cultivation of American plants as well as exotics. This garden, which is situated on the banks of the Schuylkill, a few miles from Philadelphia, still bears his name. He contributed much to the gardens of Europe, and corresponded with the most distinguished naturalists of that quarter of the globe. Several foreign societies and academies bestowed their honours upon him, and published communications from him in their *Transactions*. John Bartram died in 1777, in the 76th year of his age. At the time of his death he held the office of American botanist to George III. of England. He was amiable and charitable, and of the strictest probity and temperance. (*Encyc. Amer.*)

William Bartram, fourth son of John Bartram, was born in 1739, at the Botanic Garden, Kingessing, Pennsylvania. [At the age of 16 years he was placed with a respectable merchant of Philadelphia, with whom he continued six years; after which he went to North Carolina, with a view of doing business there as a merchant: but, being ardently attached to the study of botany, he relinquished his mercantile pursuits, and accompanied his

father in a journey into East Florida, to explore the natural productions of that country; after which he settled on the river St. John's, in that region, and finally returned, about the year 1771, to his father's residence. In 1773, at the request of Dr. Fothergill of London, he embarked for Charleston, to examine the natural productions of the Floridas and the western parts of Carolina and Georgia, chiefly in the vegetable kingdom. In this employment he was engaged nearly five years, and made numerous contributions to the natural history of the country through which he travelled. His collections and drawings were forwarded to Dr. Fothergill; and about the year 1790 Bartram published an account of his travels and discoveries in one volume 8vo, with an account of the manners and Customs of the Creeks, Cherokees, and Choctaws. This work soon acquired extensive popularity, and is still frequently consulted. After his return from his travels, he devoted himself to science, and, in 1782, was elected professor of botany in the university of Pennsylvania, which post he declined in consequence of the state of his health. In 1786 he was elected a member of the American Philosophical Society, and was a member of several other learned societies in Europe and America. We are indebted to him for the knowledge of many curious and beautiful plants peculiar to North America, and for the most complete and correct table of American ornithology, before the work of Wilson, who was assisted by him in the commencement of his *American Ornithology*. He wrote an article on the natural history of a plant a few minutes before his death, which happened suddenly, by the rupture of a blood-vessel in the lungs, July 22, 1823, in the 85th year of his age. (*Ibid.*)

In Scotland, as we have seen (p. 48.), very little was done in the way of introducing foreign trees and shrubs during the seventeenth century; though the rudiments of this description of improvement were laid about the end of it, by the establishment of the Edinburgh Botanic Garden. In Nicolson's *Scottish Historical Library*, published in 1702, this garden is stated to have been brought to the highest degree of perfection by its curator, Mr. James Sutherland, "whose extraordinary skill and industry" are said to have greatly advanced this department of natural history in Scotland. In Sibbald's *Scotia Illustrata*, published in 1684, the Edinburgh Botanic Garden is said to contain an arboretum, in which was "every kind of tree and shrub, as well barren as fruit-bearing, the whole disposed in fair order" (p. 66.); and in Sibbald's *Memoria Balfouriana*, published in 1699, this garden is said to be "the greatest ornament of the city of Edinburgh." (p. 73.) The plants of this garden have been twice removed to other situations (first in 1767, and again in 1822), and we believe there is now neither a

tree nor a shrub on the original site. Notwithstanding the example shown by the arboretum in the Botanic Garden, however, the planting of foreign trees and shrubs still appears to have been but little practised in Scotland. A public garden, to contain fruit trees, it appears, was projected for Edinburgh so early as 1662. Maitland informs us that the town council of Edinburgh, "by their act of 15th of March, 1662 (*Coun. Rep.*, vol. xxi. f. 99.), demised to John Thomsone, gardener, for a term of nineteen years, the plot of ground at present called Parliament Close, with the brae or side of the hill, inclosed with a stonern wall; and that "the said Thomsone was to plant a hedge as the eastern boundary." This spot of ground, according to the tenour of the tack, or lease, was to be laid out in walks, and to be planted with trees, herbs, and flowers, exclusive of cabbage, and other common garden stuff. Pursuant to the above agreement, Thomsone, on the 8th of April following, delivered to the council a plan for beautifying the inclosure, which was approved of." Two walks were to be made, a larger and a less one, from east to west; and "their western end, opposite to the Parliament House, was ordered to be planted with plum and cherry trees; and to be bordered with gooseberry, currant, and rose bushes; and flowers to be set along the southern wall or wooden rail at the head of the brae, or brow of the hill; and, at the eastern end, as aforesaid, a hedge." (*Maitland's Hist. of Edin.*, p. 186.) It seems, however, from a subsequent passage, that the plan for this garden was never carried into execution, and that the eastern boundary of the Parliament Close was let soon after for building small shops. Reid, in his *Scots Gardener*, published in 1683, mentions very few trees and shrubs. The most rare of these are, the evergreen oak, the cypress, and the arbutus. He says there are the Indian and Spanish jasmynes, myrtles, olcanders, and orange trees, which some are at great pains in governing; but he adds, "for my part I would rather be in the woods, parks, &c., measuring, planting, and improving." (p. 112.) Those who are curious in trees and other plants, he refers to the catalogue of the "learned and most ingenious Mr. James Sutherland, Physic Gardener at Edinburgh." (p. 114.) It appears by an *Essay on Enclosing, Planting, &c., in Scotland*, published in Edinburgh in 1729, that there was "but a very little stock of trees, either barren, fruit, or hedging quicksets. One who encloses," continues the author, "must get his quicksets from England or Holland, or he must sow Devonshire or French whin seed." (p. 289.)

It is fortunate for the historian of tree culture in Scotland, that such a writer existed before him as the late Dr. Walker, professor of natural history in the university of Edinburgh. This excellent man, whose garden we have seen in our younger

days, at Collington, near Edinburgh, was ardently attached to the study of organised nature from his youth; and, as he mentions in a letter to Lord Kaims, published in Tytler's life of that eminent man, more particularly to plants. Wherever Dr. Walker went, he seems to have paid peculiar attention to trees and plantations; and there are few works which contain sounder information on the subject than his *Economical History of the Hebrides and Highlands of Scotland*, published in 1812, nine years after the author's death, which happened in 1803. The facts, as to trees, given in this history, were collected, Dr. Walker informs us, between the years 1760 and 1786; and, as will appear from the following extracts, they are of very great interest.

"The first trees planted by art in Scotland," says Dr. Walker, "were those of foreign growth, and especially the fruit-bearing trees. Long before the Reformation, various orchard fruits, brought probably from France, were cultivated in the gardens of the religious houses in Scotland. Some of these fruit trees planted, perhaps, but a little before the Reformation, still remain. A few exotic barren trees were likewise propagated, such as the elder and the sycamore, and, at a later date, the beech and the chestnut; but none of our native trees were planted, such as the fir [pine], oak, ash, elm, and birch, till about the beginning of the last century. The first exotic tree of the barren kind planted in Scotland seems to have been the elder. Though a slow-growing and long-lived tree, many generations of it have succeeded each other in that country. Elder trees of a large size and very ancient date still appear; not only about old castles, but about the most considerable and oldest farm-houses. It was very generally planted, and for a very useful and peculiar purpose, the wood of the elder being accounted in old times, preferable to every other sort for the making of arrows. The plane [sycamore, *Acer Pseudo-Platanus*], in point of antiquity, appears to be the next. When it was first introduced is uncertain; but it seems not only to have been planted, but to have been propagated by seeds and suckers, for several generations before any other forest tree was introduced into Scotland. The wood of this tree, in old times, must have been of great value in the hands of the turner; and for that purpose chiefly it seems to have been cultivated. It is better adapted for the wooden bowls, dishes, platters, and other domestic utensils which were universally in use, than the wood of any native tree in the country. These, however, the elder and the sycamore, appear to have been the only two barren trees planted in Scotland, till towards the middle of the seventeenth century."

Perhaps the oldest sycamore in Scotland, and which appears to be at the same time the largest tree of the kind in Britain, is

that at Kippenross, in Perthshire. In 1823, it measured 28 ft. 9 in. in circumference, at a foot from the ground. It appears, from a statement made by the Earl of Mar to Mr. Monteith, that this tree went by the name of "the big tree in Kippenross" in the time of Charles II. (*Monteith's Forester's Guide*, 2d edit. p. 394.) In the grounds of Callendar House, near Falkirk, there are sycamores and other trees of great size, which must have been planted at the commencement of the 18th century, if not in the latter part of the 17th century. Dr. Graham states, on authority which he considers almost approaching to a certainty, that these trees were planted by the Earl of Linlithgow and Callendar, who had accompanied Charles II. in his exile, upon his return from the Continent, after the Restoration. The dimensions of these remarkable trees are given in the appendix to the *General Report of Scotland*, vol. iv. p. 482.

We are not aware of any elder trees in Scotland of any great size or indicating great age. The tree is so completely naturalised there, more especially about houses and places where cultivation has long existed, that, if it were not for Dr. Walker's statement, we should never have supposed it to be otherwise than an indigenous tree.

The sweet chestnut and the beech, producing seeds easily portable from other countries, were in all probability introduced into Scotland by the Romans, and, perhaps, reintroduced by the religious establishments in the middle ages. According to Dr. Walker, "a few chestnuts and beeches were first planted in gardens not long before the middle of the seventeenth century, some of which have remained till our own times. Such was the chestnut at Finhaven; another at Levenside in Dumbartonshire, which was thrown down by the hurricane in 1739; and two or three, which were alive and vigorous, at Kinfauns in Perthshire, in the year 1761. Such was the great beech at Taymouth, overturned by a storm some years ago; the beech at Oxenford; that at Newbattle in Mid Lothian; and another at Ormiston Hall in East Lothian."

The two last-named beech trees, we conclude, from Sir Thos. Dick Lauder's notes respecting them in his edition of *Gilpin's Forest Scenery*, vol. i. p. 266., are decayed; and we believe that the largest beech trees now existing in Scotland are at Ardkinglass in Argyllshire, and one mentioned by Mr. Sang (*Planter's Calendar*, 2d edit. p. 52.) as growing at Panmure in Forfarshire. The latter is 26 ft. 6 in. in circumference, at the surface of the ground; and the former as large, with a longer stem and a finer head. (*Gilpin, &c.*, p. 267.) One of the largest sweet chestnuts in Scotland is at Cairn Salloch in Dumfriesshire; at 2 ft. from the surface of the ground it measures 26 ft. in girth; and it is divided into four large arms, 26 ft., 35 ft. 31 ft., and 28 ft. in

length. There is a very old tree at Riccarton, near Edinburgh, which has been described and figured by Sir Thomas Dick Lauder. The trunk is much injured and decayed; but its boughs and foliage are of luxuriant growth; the branches hang down to the ground, and, in many places, have rooted into it. The trunk is 27 ft. in girth at the surface of the ground, and the branches cover an area of 77 ft. in diameter. (*Ibid.*, p. 268.)

Dr. Walker mentions (p. 213.) some sweet chestnuts which he found, about 1760, in a thriving condition in the Island of Inchmahona, in the Lake of Menteith, in Perthshire, where there was a priory founded by King David I. Dr. Patrick Graham measured some of these trees in 1813, and found the trunks to be 18 ft. in circumference at 6 ft. from the ground. (*General Report of Scotland*, vol. ii. p. 254.) He thinks they were then 300 years old, or upwards, which would carry the date of their planting back to the commencement of the sixteenth century. According to Dr. Walker, as before quoted (p. 34.), the sweet chestnut at Finhaven was both the largest tree of the kind in Scotland, and the first tree planted there by art. "In the year 1760, a great part of the trunk of this remarkable tree, and some of its branches, remained. The measures of this tree were taken before two justices of the peace, in the year 1744. By an attested copy of this measurement, it appeared, at that time, that at half a foot above the ground, it was 42 ft. 8½ in. in circumference. As this chestnut appears, from its dimensions, to have been planted about 500 years ago, it may be presumed to be the oldest planted tree that is extant, or that we have any account of, in Scotland." (*Walker's Essays*, p. 29.) Sir Thomas Dick Lauder states, that, "in the possession of Skene of Carriston, there is a table made of the wood of this tree, having an engraved plate, on which are marked its dimensions. The castle of Finhaven was an ancient seat of the Earls of Crawford." (*Lauders Gilpin*, vol. i. p. 269.)

To the research of Dr. Walker we are indebted for the following list of trees in Scotland, with the name of the places where they were introduced:—

- | | |
|---------------------------------------------------------------------|---------------|
| 1664. <i>Tilia europæa</i> , lime. | Taymouth. |
| 1678. <i>Salix álba</i> , white willow. | Prestonfield. |
| 1682. <i>Abies Picea</i> and <i>excélsa</i> , silver and pitch fir. | Inverary. |
| <i>Acer</i> , maple. | Inverary. |
| 1690, <i>Juglans régia</i> , walnut. | Kinross. |
| 1692. <i>Cárpinus Bétulus</i> , hornbeam. | Drumlanerig. |
| 1695. <i>Cérasus Lusitánica</i> , the Portugal laurel. | Inverary. |
| (Gard. Mag., vol. ii. p. 178.) | |
| 1696. <i>Pópulus nigra</i> , black poplar. | Hamilton. |
| 1705. <i>Cýtisis alpinus</i> , alpine laburnum. | Panmure. |
| 1709. <i>Æsculus Hippocástanum</i> , horsechestnut. | New Posso. |

1710. *Plátanus orientàlis*, Oriental plane. Holyrood House.
 1712. *O'rnus europæa*, flowering ash. Bargally.
 1725. *Pínus Stròbus*, Weymouth pine. Dunkeld.
 1727. *Làrix europæa*, larch. Dunkeld.
 1730. *Quércus I'lex*, evergreen oak. New Hailes.
 1732. *A'bies balsamífera*, balm of Gilcad fir. Arbighland.
 1733. *Taxòdium dístichum*, deciduous cypress. Loudon.
 1734. *Quércus Æ'gilops L.*, Velonia oak. New Hailes.
 1736. *Ulmus campéstris*, English elm. Dalmahoy.
 1738. *A'cer platanòides L.*, Norway maple. Mountstewart.
 1739. *Sálix phloráigna* [q. triándra], Tine- Newhails.
 bark willow.
 1740. *Cèdrus Libàni*, cedar of Lebanon. Hopetoun.
 1743. *Cérasus carolinénsis*, Carolina bird-cherry. Hopetoun.
 1744. *Córylus ? Colúrna*, Hungarian nut. Carmichael.
 1746. *Sàlix amerìna*, Amerina willow. Mellerstane.
 1754. *A'cer sacchàrinum*, sugar maple. New Posso.
 1759. *A'bies canadénsis*, white Newfound- New Posso.
 land spruce, or hemlock spruce.
 1763. *Fráxinus americàna* and sp., white and blue American
 ash.
 Pínus longifòlia, long-leaved American pine.
 A'cer pennsylvànicum, snaked-barked maple.
 Làrix nìgra, American larch.
 Bétula papyrífera, the paper birch.
 1765. *Bétula nìgra L.*, black American birch. Elliock.
 1766. *Pópulus dilatàta*, Lombardy poplar. New Posso.
 1770. *Pópulus balsamífera*, balsam poplar. Leith.

From this period (1770) the intercourse between Scotland and England became so frequent, that the dates of the introduction of foreign trees and shrubs into the two countries may be considered as merged into one.

It would be interesting to know some particulars respecting the tastes and pursuits of the proprietors of the places mentioned in Dr. Walker's list; but at this distant period, we have been able to glean very little suitable to our purpose respecting them.

Taymouth, in the central highlands of Perthshire, is a very old seat of the Campbells. Pennant says the castle was first built by Sir John Campbell, sixth knight of Lochow, who died in 1583. The place, he says, has been much modernised since the days of the founder, and has lost its castellated form, as well as its old name of Balloch Castle. The place has subsequently undergone a great variety of alterations, and at present is remarkable for the extent of its woods and plantations, and for a fine avenue of lime trees. The present proprietor is John Campbell, Marquess of Breadalbane.

Inverary Castle was inhabited by a Colin Campbell before

1480; and is now the property of George William Campbell, Duke of Argyll. It is a magnificent place, from its great extent, with the sea in front, and backed by wooded hills and lofty mountains. The house is an immense quadrangular building, and with the plantations and pleasure-grounds, are said to have cost, within the last half century, upwards of 300,000*l.* The Portugal laurel was introduced here in 1695, and is said to have been brought from Portugal by Duke Archibald; one tree spreads over a circle of 165 ft. in circumference, and is nearly 40 ft. high. In Smith's *Agricultural Report of Argyllshire*, the oldest and largest trees at Inverary are supposed to have been planted by the Marquess of Argyll (frequently mentioned by Evelyn as a great planter), between the years 1650 and 1660. Those of the next largest size and age were raised from seed by Archibald Duke of Argyll in 1746 or 1747. These latter consist chiefly of larches, New England pines, and spruce and silver firs. (*Report, &c.*, p. 156.) The soil and climate at Inverary are said to be remarkably favourable for the growth of trees.

Prestonfield is a well known place in the neighbourhood of Edinburgh, which, in 1783, belonged to Sir Alexander Dick, a great horticulturist as well as agriculturist, and distinguished by having been the first to produce good medicinal rhubarb in Scotland. (See *Wight's Husbandry of Scotland*, vol. iii. pt. ii. p. 443.) Kinross was built and planted, about 1685, by Sir William Bruce, the celebrated architect, for his own residence, and was the first good house of regular architecture in Scotland. It was approached by a fine avenue of trees. Drumlanrig, in Dumfriesshire, was built by the Duke of Queensberry in a commanding situation: it took ten years in building, and was finished in 1689. The duke expended an immense sum in forming terraced gardens, which, according to Gilpin (*Observations, &c., in Scotland*, 1776), served only to deform a very delightful piece of scenery. The duke, he adds, seems to have been aware of his folly, for he is said to have "bundled up all the accounts together, and inscribed them with a grievous curse on any of his posterity who should ever look into them." The property now belongs to the Duke of Buccleugh and Queensberry, who is planting and building there very extensively. (See an account of Drumlanrig, when visited by us in 1831, in the *Gardener's Magazine*, vol. ix. p. 1.)

Hamilton Palace, in Lanarkshire, the ancient seat of the Dukes of Hamilton, was built at different periods; the most ancient part in 1501. The grounds were laid out in the year 1690. The gardens and lawns near the house were planted with foreign trees, especially lime trees, some fine specimens of which still remain. One of the earliest nursery gardens in Scotland appears to have been established at the little village of

Hamilton, close by the palace, being the only garden for the sale of plants mentioned by Reid in his *Scots Gardener*, published in 1683. Among the oaks of Hamilton Park, so famous down to the end of the seventeenth century, there were trees, Nasmyth informs us, which measured 27 feet round the trunk, with wide expansive branches. (*Agriculture of Clydesdale*, p. 144.)

Panmure is the name of an ancient family in Angusshire, whose chief seat is the spacious and hospitable mansion of Brechin Castle, which, from the remotest period of its history, has always been possessed by the Maules, formerly Earls of Panmure. Panmure, another seat of this family, is near Dundee, and was built about 1665. It is a venerable fabric, and is kept by the proprietor, with all its furniture and pictures, in the same state in which it descended from his ancestors. In Dr. Walker's time, Panmure was famous for its laburnums, which were planted towards the end of the seventeenth century, and had attained a great size in 1780. Sang says that a considerable quantity of the laburnums at Panmure and Brechin were cut down in 1809, and sold by public sale at fully 10s. 6d. a foot, chiefly to cabinet-makers.

New Posso, in Peeblesshire, was formerly called Dalwick, Dawick, or Daick. It belonged, in very ancient times, to the chiefs of a very considerable family of the name of Veitch; but, in 1715, it was in the possession of Sir James Nasmyth of Posso, an eminent lawyer, who rebuilt the house and garden, and by some ornamental planting added greatly to the beauty of the place. Pennicuick mentions that, in an old orchard near the house, the herons built their nests upon some pear trees, which were large and old trees in 1715. Armstrong, in 1775, says that New Posso, formerly called Dalwick, "from being a lonely mansion in the bosom of a gloomy mountain, is now the extreme reverse. The vast improvements made by its present possessor have proved not only an ornament to Tweeddale, but a worthy example for emulation in the gentlemen of the county. The botanical and culinary gardens are justly esteemed the most copious in it; and the pleasurable attention with which they are cultivated, is sufficiently expressed on the front of the greenhouse, alluding to its flowers, 'Solomon in all his glory was not arrayed like one of these.'" (*Armstrong*.)

"The name of New Posso," Dr. Pennicuick tells us, "was given to the place by Sir James Nasmyth, grandson of the first possessor of that name, who was sheriff-depute in 1627. The late Sir James Nasmyth of Posso has extended and finished the place, and numerous plantations, with as much taste and elegance as the Dutch mode of gardening by line and rule will admit of. He likewise kept it in high order, and by the superiority of his own external appearance, politeness, knowledge

of the world by travel, and accomplishments, rendered both himself and his seat the models for imitation to the country where he lived. To have every thing about themselves and their houses as like to James Nasmyth and New Posso as possible was then the height of their ambition, about 1775, among the gentry of Tweeddale. A very well written letter, by this Sir J. Nasmyth, on the subject of botany, in answer to one from His Lordship at Blair Drummond, is preserved in Lord Woodhouselee's *Life of Lord Kaims*, and in compliment to him the birch called the *Bétula, Nassmythii* was so named. Many of the fine trees about New Posso have been lately cut down and sold, besides all those at Posso." The above is extracted from Dr. Pennycuick's *Works in Prose and Verse*, which were originally published in 1715, but of which an edition was published in 1814, with notes up to that year. New Posso is at present distinguished for its pine and larch plantations; and, according to Sir Thomas Dick Lauder, the first larches introduced into Scotland were planted at Dawick in the year 1725 (*Lauder's Gilpin's Forest Scenery*, vol. i. p. 148.); though this is doubtful as will hereafter appear. Sir Thomas also mentions a locust tree at Dawick, which, at 3 ft. from the ground, measures 5 ft. 10 in. in girth. The present proprietor of Dawick, or New Posso, is Sir John Nasmyth, grandson of the Sir James celebrated by Dr. Pennycuick.

Holyrood House, where the Oriental plane was first planted in Scotland, is, as every one knows, the royal palace of Edinburgh. The Abbey of Holyrood, according to Maitland (*Hist. of Edin.*), was founded by King David I., in 1128, and consisted of a church and cloister. Maitland speaking of this church and cloister, says:—"After having stood 400 years in the fields, by themselves, King James V., about the year 1528, erected a house to reside in at his coming to Edinburgh, near the south-western corner of the church, with a circular turret at each angle, which is the present tower at the north-western corner of the palace; to which was added, by King Charles II., in the year 1674, all the other parts of the present magnificent royal mansion. The said King James, to accommodate himself with a park, inclosed a large quantity of ground in this neighbourhood with a stonern wall, about three miles in circumference, which probably is no where to be paralleled; for, instead of trees and thickets for cover, which other parks abound with, I could not, after the strictest search, discover one tree therein: in lieu whereof, it is supplied with huge rocks and vast declivities, which furnish the Edinburghers with the best of stones to pave their streets withal; as do the other parts of the said park yield good pasturage, and meadow grounds, with considerable spots of arable land." (*Maitland's Hist. of Edin.*,

fol. 1753, p. 152.) Arnot, in his *History of Edinburgh*, published in 1779, speaking of this park, says:—"In the memory of people not long since dead [Arnot wrote about 1779], the level strip at the foot of the hill [Arthur's seat], which, from the Duke of York having delighted to walk in it, bears the name of 'The Duke's Walk,' was covered with tall oaks; but now there is hardly a single tree in its whole boundaries. Indeed, it is extremely doubtful if, except at the bottom, there ever were any trees on these hills, the height of the ground and barrenness of the soil being very unfavourable to their growth." (*Arnot's Hist. of Edin.*, 4to, Edin., 1779, p. 309.) It is clear, therefore, that the platanus, mentioned by Dr. Walker, was not planted in the park at Holyrood House: but we learn from the same authorities (Maitland and Arnot) that there were two walled gardens attached to the palace; and that "the royal garden at the northern end of the outer court" was "converted into a physic garden," and that it was under the same superintendence, and applied to the same purposes, as the physic garden at the North Loch. There can be no doubt, but it was in the physic garden adjoining the palace, that the platanus mentioned by Dr. Walker was planted; and the planter was probably Sutherland.

Bargally is to us by far the most interesting seat in Scotland, with respect to the introduction of foreign trees and shrubs, and though we have taken the greatest pains to ascertain from what circumstances its proprietor became so much attached to botanical pursuits, as to introduce in a remote part of Scotland, in the 17th century, trees then scarcely known even in its metropolis, and have been in a great measure successful; yet there is still some deficiency in the information we have obtained. Bargally is a small property situated in a glen, the sides of which are covered with natural wood, between Gatehouse in Kirkcudbrightshire, and Newton-Steward. The proprietor's name was Andrew Heron; and he appears, by a family tomb in the grounds, to have died there in 1729. We have applied to about a dozen different persons in Kirkcudbrightshire, whom we deemed most likely to give us information respecting Bargally and its planter, and the following are extracts from the communications we have received, joined to what we have been able to glean from books. One of our correspondents informs us, "that Andrew Heron was a son of Heron of Heron of Kirrauchtrees (Caeruchtred), chief of that name. This Andrew built a cottage, in which he resided, at the upper extremity of the beautiful valley in which the present house of Bargally stands, and planted all the lower part of the valley. The splendid *Quercus Ilex* and the noble beeches, which you saw in 1831, are but miserable relics of the magnificent forest which once rose between Bargally House and the river Palnure.

Andrew Heron married twice, and left a family. He, with one of his wives, was buried in a tomb which lies in front of Bargally House. The date inscribed on this tomb is 1729. Andrew's estate devolved, on his death, to his son, Dr. Andrew Heron; but he was involved in a lawsuit with the Kirauchtree family; and though it was decided in his favour, it ended in greatly injuring his fortune. In consequence of this, he sold Bargally to Hannay of Kirkdale, and retired to a cottage, where he died many years ago at a very advanced age." Another correspondent says, "I was born about two miles from Bargally, and recollect to have seen Dr. Heron, the son of the botanist, when I was very young. I communicated with several people who had lived their whole lifetime near Bargally, and are considerably older than I am, but they all replied that they knew nothing more about old Bargally than what I had stated to them. I recollected that the Herons of Heron [the estate of Heron is in Northumberland, see *A Genealogical History of the Ancient Family of Heron*, imp. 4to, part ii.] of Kirauchtree, and the Herons of Bargally, were originally from the same stock; and, as Lady Heron Maxwell of Springkell is the last of the lineal descendants of the Herons of Kirauchtree, I wrote to Her Ladyship, mentioning that you were engaged on a work that required some information about Andrew Heron of Bargally, and begging that she would tell me what she knew or had heard about him. I received a letter from Her Ladyship yesterday (April 6. 1835), giving me extracts from an old manuscript document in her possession relative to Andrew Heron of Bargally. I enclose a copy, and I hope it will give you all the information you require about that wonderful man. It appears that Andrew Heron was of no profession, simply the 'Laird of Bargally;' but he must have been a person who had travelled a good deal, to induce him to do so much at that early period.

"The old orchard and flower-garden at Bargally have been, to my personal knowledge, a grass field for forty years and upwards; but some of the fine variegated hollies, now large trees, still remain to mark the different divisions of the garden. About thirty years ago, when I was walking over the grass field, which was originally the garden at Bargally, in the month of August, I observed peering through the grass some crocus plants, both white and purple; this surprised me, for I had never seen an autumnal crocus. The gentleman to whom the property then belonged, was also astonished, saying that he had never observed them before. I cut up a few of the roots of the different kinds, with a portion of the turf, and carried them to St. Mary's Isle, and from these roots many plants have been propagated. There are still some curious trees and plants to be seen at Bargally, remaining to sound the praises of old Andrew Heron the

botanist. Bargally was sold by the heirs of Andrew Heron to Mr. William Hannay, the brother of Sir Samuel Hannay of Kirkdale; he was scarce of cash, and cut down the wood of Bargally (including many of the fine trees that had been introduced and planted by Andrew Heron), in the year 1791. I purchased a portion of the trunk of a silver fir, and I made it into a meal chest; the side boards, the bottom, the ends, and top, or lid, of which chest are all out of one board. This chest, is still in my possession, and in use; and it is in depth and breadth, after having been wrought, 2 ft. 2 in.”—*W. M.*

Extract from an old manuscript in the possession of Lady Heron Maxwell of Springkell, relating to Andrew Heron of Bargally:—“Andrew Heron of Bargally was the second son of Andrew Heron of Heron, who settled the lands of Bargally upon him as his patrimony. In 1690 he went to reside at Bargally; in 1693 he built the great dyke for the garden and orchard; and, the next year, he began to collect and fill in a large number of trees, fruit and flowers. His father died in 1695. In that year Andrew Heron employed Mr. Hawkins, an Englishman, to build the stone house. The stone was all got out of a quarry on the east side of the garden; it was finished, watertight and in order, in 1696. In 1697 and 1700 he built the pigeon house and the crews [farm offices]. Andrew Heron of Bargally married, secondly, the relict of John M’Kie of Larg, in April, 1708; and, having lived twenty-one years after his second marriage, hath improved the ground to great advantage, having enclosed all the low ground, and built a new stone house, made large gardens, well stocked with all kinds of fine trees and rare fruits, both stone and core; some portions were stocked with fine flowers, and he had a green-house stocked with oranges, lemons, pomegranates, passion flowers, citron trees, oleanders, myrtles, and many others. The eldest son of Andrew Heron of Bargally was a captain in Lord Monk Kerr’s regiment, and married the daughter of Mr. Vining, a rich merchant at Portsmouth. He left several sons and a daughter: John, bound apprentice to his brother in law, Mr. Reid, a considerable merchant; and Andrew, who is bound apprentice to a surgeon at Bath. He hath also Patrick, Charles, and Benjamin: and of daughters, Jane, who married Mr. Reid, to whom her grandfather, Mr. Vining, gave 3500*l.* in marriage portion, a large fortune at that time.”

Lady Heron Maxwell added, from her own knowledge, the following additional information:—“The first Heron of Bargally was the uncle of my great grandfather, Heron of Heron, who represented the stewartry of Kirkcudbright in parliament at different times; and I am now the last of the direct line of the family of Heron of Heron, and that family held large posses-

sions in Galloway, from father to son, for upwards of 500 years. The only remaining known descendant of Heron of Bargally, in the male line, is Captain Basil Heron of the Royal Artillery, now (1835) on duty at Gibraltar; he married a daughter of Judge Mayne, in Ireland, and has three daughters; he is grandson of Dr. Heron, who sold Bargally, and great-grandson of Andrew Heron the botanist. The male heirs of all the branches of the family of Heron of Heron will be extinct on the death of Captain Basil Heron."

Andrew Heron corresponded with Bradley on gardening subjects; and from this correspondence it appears that he had a curious water-clock in his grounds, that he trained his pear trees in a particular manner, and that he cultivated in his fields what he calls the "true Roman cytissus." (See *Bradley's Treatise on Husbandry and Gardening*, 1726, vol. ii. p. 169.) Mr. Maxwell, writing about the same period to Mr. Hope of Rankeillor, says, "I have of late been with Mr. Heron of Bargally, in whose garden there is a great variety of curiosities to be observed. He is, in my opinion, the most learned and ingenious gentleman, in the article of gardening, I ever conversed with." (*Practical Husbandman*, p. 179.) "The want of money, that great enemy to old timber," another correspondent informs us, "compelled Mr. Hannay, the purchaser of Bargally from Dr. Andrew Heron, who was otherwise a gentleman of fine taste, to cut down a great many of the largest trees, particularly four that grew one at each corner of Heron's tomb."

We visited Bargally in August 24. 1831, and found a number of the trees planted by Andrew Heron still in existence. Having applied to the present proprietor, John Mackie, Esq., for the dimensions of some of these trees, the following is an extract from his letter, dated Bath, March 21. 1835:— "I have now received the measurement of some of the old trees at Bargally, which is as follows, viz.: the circumference of a beech (usually denominated the large beech), at 18 in. from the ground, is 18 ft., and it is upwards of 90 ft. in height. This tree is in the most perfect health, and when in full foliage is truly magnificent. The circumference of an evergreen oak in the garden, at 14 in. from the ground, is 12 ft. and it is above 60 ft. high: this tree is also in a very thriving state, and does not show the least tendency to decay. The circumference of a hop hornbeam (*Ostrya vulgaris*), at 1 ft. from the ground, is 9 ft., and it is above 60 ft. high: this tree is particularly mentioned by Dr. Walker, as having been measured by him in 1780; it was then 4 ft. 1 in. in circumference at 4 feet from the ground, 60 ft. high, and considered the oldest and largest tree of the kind in Scotland. The circumference of a variegated

sycamore is $12\frac{1}{2}$ ft., and it is upwards of 70 ft. high. The girth of a sweet chestnut, at 18 in. from the ground, is 10 ft. 7 in., and it is above 80 ft. high. Mr. M'Nab, my factor, adds, 'Had I measured them at the surface of the ground, they would have been one third more, in consequence of the roots spreading so much as they do.' Mr. Hannay sold the property of Bargally to my father in 1792."

"It is recorded of Mr. Heron, that he went to visit a garden in the neighbourhood of London, and very much astonished the principal gardener, to whom he was a stranger, with the botanical knowledge he displayed; and the gardener having shown him an exotic, which he felt confident Mr. Heron had never seen, he exclaimed, on Mr. Heron's readily naming it, 'Then, Sir, you must either be the devil or Andrew Heron of Bargally;' thereby intimating that Mr. Heron was proverbial, in those days, as a botanist, even with those who had never seen him."

Dr. Walker, in his *Essays* (p. 32.), mentions several firs and pines at Bargally, of large dimensions, which no longer exist. A fir, he says, which was planted in 1697, measured, in 1780, 90 ft. in height. He states that the oldest and largest arbor vitæ in Scotland was at Bargally: it measured, in 1780, 5 ft. 4 in. in girth at 4 ft. from the ground, and was 40 ft. high. He also mentions a flowering ash (*O'rnus europæ'a*), which was cut down in 1780, and 7 ft. of the trunk quartered to make four axles to carts; it was a remarkably handsome tree, 6 ft. 3 in. in circumference at 4 ft. from the ground, and 50 ft. high. Dr. Walker mentions large evergreen oaks, horsechestnuts, and many other species, of extraordinary dimensions. The present proprietor is much attached to this beautiful place, takes the greatest care of the trees, and has lately repaired the tomb of their planter.

We took notes ourselves (in 1831) of several remarkable trees at Bargally, including a large lime tree and a number of beautiful variegated hollies, from 20 ft. to 26 ft. in height, and with trunks from 15 in. to 2 ft. in diameter. Altogether the place is one of very great interest, not only on account of its venerable foreign trees and the tomb of Heron, but from the romantic beauty of the situation, and from the district in which it lies being one of the finest, in point of scenery, in the west of Scotland.

Dunkeld, where, it appears, the Weymouth Pine was first introduced into Scotland, was, in 1727, the property of James Murray, Duke of Athol; the friend and distant relative of John and Archibald, Dukes of Argyll. Dunkeld is celebrated for having been one of the first places where the larch was planted in Scotland; the plants of which, it is said, were sent from

London in pots in the year 1741. Sir Thomas Dick Lauder, as we have seen, however, says the larch was first planted in Scotland, at Sir James Nasmyth's, at Dawick, in 1725. The Rev. James Headrick, in his *Survey of Forfarshire*, gives another account of the introduction of the larch into Scotland. He says, "It is generally supposed that larches were first brought into Scotland by one of the Dukes of Athol; but I saw three larches of extraordinary size and age, in the garden near the mansion house of Lockhart of Lee, on the northern banks of the Clyde, a few miles below Lanark. The stems and branches were so much covered with lichens, that they hardly exhibited any signs of life or vegetation. The account I heard of them was, that they were brought there by the celebrated Lockhart of Lee (who had been ambassador from Cromwell to France), soon after the restoration of Charles II. (about 1660). After Cromwell's death, thinking himself unsafe on account of having served a usurper, he retired some time into the territories of Venice. He there observed the great use the Venetians made of larches in ship-building, in piles for buildings, in the construction of their houses, and for other purposes; and when he returned home he brought a number of larch plants in pots, with a view to try if they could be gradually made to endure the climate of Scotland. He nursed his plants in hot-houses, and in a green-house sheltered from the cold, until they all died, except the three alluded to; these, in desperation, he planted in the warmest and best sheltered part of his garden, where they attained an extraordinary height and girth." (*Headrick's Forfarshire*, p. 374.)

The estate of Dunkeld now contains the most extensive plantations of the larch in the island, spreading over several thousand acres. A copious and most valuable account of these plantations will be found in the *Transactions of the Highland Society*, vol. ix., and an abridgement of that account in our *Encyclopædia of Gardening*, § 6579. edit. 1835. Dunkeld has long been celebrated for its scenery. Dr. Clarke, the traveller, says, the scene that opens before you, after going through the pass, perhaps has not its parallel in Europe. "The grounds of the Duke of Athol," he continues, "I do not hesitate to pronounce without a rival." Gray, the author of the *Elegy*, was "overcome and almost lost" by the beauties of Dunkeld. Gilpin called it the "portal of the Highlands," and Dr. Macculloch has nearly filled a volume on the subject. The house at Dunkeld is a plain large building, erected in 1685, but it has long been in contemplation to remove it, and to build one of superior architectural pretensions. John Murray, the present Duke of Athol, has lately constructed a magnificent public bridge over the Tay at Dunkeld, government assisting His Grace with one sixth part of the expense. The bridge was constructed on dry land, and

the course of the river was afterwards turned to it. It is greatly to the honour of this family, that for a century past their improvements, such as roadmaking, bridge-building, and planting, have been made more with a view to the general benefit of the country than to lodging themselves sumptuously.

New Hailes, near Musselburgh, was a seat of Baron Dalrymple, a celebrated lawyer and improver, and is now the property of Miss Dalrymple.

Arbigland, in Dumfriesshire, was the property of William Craik, Esq., a contemporary of Maxwell and of Fletcher of Saltoun, and one of the original members of the Society for the Improvement of Agriculture in Scotland. He was one of the first to study the works of Tull, and to adopt the drill system. He died in 1798, at the age of 95 years. We visited Arbigland in 1804, and again in 1806, and found the place still celebrated for its old silver firs. A life of this distinguished agriculturist will be found in the *Farmer's Magazine*, vol. xii. p. 145.

Loudon Castle, in Ayrshire, was one of the first places in the West of Scotland, where foreign trees were planted. "John Earl of Loudon," Walker observes, "formed at Loudon Castle, in Ayrshire, the most extensive collection of willows, that has been made in this country, which he interspersed in his extensive plantations. Wherever he went during his long military services, he sent home every valuable sort of tree that he met with. All the willows he found cultivated in England, Ireland, Holland, Flanders, and Germany, as also in America and Portugal, where he commanded, were procured and sent to Loudon. (*Econ. Hist.*, &c., p. 161.) In 1806, and again in 1831, we found a number of fine old trees at Loudon Castle; we recollect, in particular, robinias, gleditschias, American oaks, hickories, walnuts, taxodiums, acers, poplars, and a variety of others. Some are recorded by Dr. Walker as having been remarkably fine specimens in 1780.

Dalmahoy, near Edinburgh, is the property of the Earl of Morton, and there are still a few specimens of old trees there. Mount Steuart, the next place mentioned in the list, is situated in the Island of Bute, and was built in 1718 by James Earl of Bute, father of the celebrated earl of that name, who was minister to George III. The plantations there, according to Dr. Walker, were begun in the same year. Speaking of them in 1780, he says, "They are equal, if not superior, to those of the same age in Ayrshire and Renfrewshire. The Oriental plane grows here almost like a willow; is never hurt in winter, and forms a fine dressed shady tree." The Marquess of Bute's family have planted from 200,000 to 300,000 trees every year since the beginning of the present century. The place contains many remarkably

fine specimens, which will be severally noticed in the course of this work.

Hopetoun House, the property of the Earl of Hopetoun, is still celebrated for its cedars. According to a letter, dated November, 1834, which we received from Mr. Smith, the gardener there, the cedars alluded to by Dr. Walker were brought from London by Archibald Duke of Argyll, and a number of other exotic trees, such as tulip trees, evergreens, oaks, &c., appear to have been planted about the same time. It is remarkable, Mr. Smith observes, that these cedar trees are the fastest-growing trees on the estate. The largest, in 1834, measured nearly $15\frac{1}{2}$ ft. in girth, at a foot from the ground, and was 68 ft. high. The silver fir there was 90 ft. high; the tulip tree 60 ft. high; the Carolina or evergreen birdcherry, mentioned by Dr. Walker, 70 ft. high; the sweet chestnut 75 ft. high; the arbor vitæ 35 ft. high; the common holly 44 ft. high; and the common yew 28 ft. high. On the whole, Hopetoun House is one of the most celebrated places for foreign trees and shrubs in Scotland. (See *Encyc. of Gard.*, § 1225. edit. 1835.)

Carmichael was, we believe, situated in Clydesdale, and belonged to the Earl of Hyndford. Mellerstane, in Berwickshire, was the seat of George Baillie of Jerviswood. The mansion is magnificent, and the grounds extensive. Elliock, in Dumfriesshire, belongs to the Veitch family, some of whom were formerly Lords of Sessions. It has very extensive plantations.

By Leith, where the balsam poplar was first planted, we find, from another passage in Dr. Walker's works, was meant a nursery in Leith Walk; in all probability that of Mr. Richmond, who was the first to establish a nursery there, which about 1780, merged in that of Messrs. Dickson and Co.

It is observed by Dr. Walker, that most of the foregoing trees were only planted in Gardens and pleasure-grounds as objects of rarity or beauty. Planting on a large scale, for profit, was chiefly performed, as may readily be imagined, with indigenous trees. The father of this description of planting in Scotland was, according to the same undoubted authority, Thomas Earl of Haddington, who began to plant Tynningham, near Dunbar, in the year 1705. He enclosed 1000 acres, called Binning Wood, and wrote a *Treatise on Forest Trees*, which was printed in 1733. The earl died at New Hailes near Edinburgh in 1735, and was succeeded by his grandson, to whom he had addressed the letters which compose the treatise. The earl informs us in his treatise, that when he came to live at Tynningham, in the year 1700, there were not above fourteen acres set with trees. The earl's grandfather, he tells us, after the civil wars in the time of Charles I. were over, "tried to raise some trees," and for that purpose planted two rows round the

house and gardens. The author of the treatise tells us that he was "fond of dogs and horses, and had no manner of inclination to plant, till he was obliged to form some enclosures for grazing his horses, as he found the purchase of hay very expensive." After he began, his lady, who "was a great lover of planting, encouraged him to go on, and at last asked leave to go about it herself." The first Marquess of Tweeddale, Lord Rankeilor, Sir William Bruce, his father, and some others, he says, had planted a great deal; yet, he adds, "I will be bold to say, that planting was not well understood in this country till this century began. I think it was the late Earl of Mar, that first introduced the wilderness way of planting amongst us; and very much improved the taste of our gentlemen, who very soon followed his example." (p. 3.) What the earl means by a wilderness, we afterwards learn, is a plantation with straight walks cut through it, in the geometrical style of landscape-gardening; in England, a wilderness plantation is generally understood to be one in which the walks are in irregular directions.

It does not appear, from this treatise, that the earl planted many trees of foreign origin in his woods; but, from the dimensions of some arbor vitæ, evergreen oaks, chestnuts, &c., there can be little doubt that he did not lose sight of such trees in his ornamental plantations near the house. Sang, in the *Planter's Kalendar* (2d edit. p. 551.), mentions a silver fir as having been planted in Binning Wood in 1705. This wood, he says, "reflects great honour on the memory of the lady who planted it;" meaning, no doubt, the Countess of Haddington above mentioned, who is said to have sold her jewels, to enable her to plant Binning Wood. The holly hedges at Tynningham planted by this earl and his successor are unquestionably the finest in Britain. Some notices respecting these hedges are given in the *London Horticultural Society's Transactions*, vol. viii., and in the *Gardener's Magazine*, vol. ii. p. 184. There are in all 2952 yards of holly hedge, in different lengths, of different heights of from 10 to 25 ft., and of widths from 9 to 13 ft.: they are, with the exception of one, regularly clipped every April. The largest single holly at Tynningham, according to the dimensions sent us in January, 1835, was 42 ft. high. The hedges were for the most part planted in 1712. Wight of Ormiston, in his *General Survey of the Agriculture of Scotland*, speaking of Tynningham in 1768, says, these hedges, and the abundance of evergreens, give the place the appearance of summer in the midst of winter.

The great promoter of the planting of foreign trees and shrubs in Scotland, according to Dr. Walker, was Archibald Duke of Argyll; unquestionably, also, as we have seen (p. 57.), the greatest promoter of this kind of planting, in England. The duke communicated this taste to a number of his intimate friends,

both in England and Scotland. Among these, in the latter country, Dr. Walker mentions the Duke of Athol, the Earls of Bute, of Loudon, of Pyndford, and of Panmure; Sir James Nasmyth, Mr. Fletcher of Saltoun, Sir Archibald Grant, and others. By the exertions of these gentlemen, planting became very general in Scotland between the years 1730 and 1750. (*Walker's Hebrides*, vol. i. p. 210.)

Sir Archibald Grant began to plant in 1719. The following is an extract taken from a commonplace book kept by this gentleman, and published in the *Gardener's Magazine*, vol. xi. p. 48. :—"In 1715," Sir Archibald says, "by the indulgence of a very worthy father, I was allowed, though then very young, to begin to enclose and plant, and provide and prepare nurseries. At that time there was not one acre on the whole estate enclosed, nor any timber upon it but a few elms, sycamore, and ash, about a small kitchen-garden adjoining to the house, and some straggling trees at some of the farmyards, with a small copsewood, not enclosed, and dwarfish, and browsed by sheep and cattle."

It is probable that most of the foreign trees and shrubs that were introduced into Scotland previously to the middle of the 18th century, were raised from seeds in the different localities. There could have been few, if any, public tree nurseries in Scotland previously to that period; and the carriage of trees from England must have been extremely tedious and expensive. The Earl of Haddington was in all probability, the originator of nurseries in Scotland, as well as the father of artificial plantations in that country, on a large scale for profit. John Reid, the author of the *Scots Gardener*, published in 1683, mentions Hugh Wood, gardener at Hamilton, dealing in fruit trees and numerous other garden articles, whether English, Dutch, or Scotch, but he makes no mention of forest trees. Sutherland's *Hortus Medicus Edinburgensis*, published in 1683, is stated in the titlepage to be sold by "Mr. Henry Ferguson, seed merchant, at the head of Black Friar's Wynn." That there were plants, trees, &c., sold by the gardeners in Scotland, is obvious from the following advertisement, dated 1721 :—"There is to be sold at John Weir's, gardener at Heriot's Hospital, and at James Weir's, son to the said John, his house at Tolcross, at the end of the West Port, all sorts of garden seeds, fruit and barren trees, and evergreens, as also flowers of the best kinds." Archibald Eagle of Edinburgh was seedsman to the Society of Improvers of Agriculture in Scotland in 1743; and, the Society having been established in 1723, this firm, now Eagle and Henderson, may date from the latter period. They had, however, no nursery for at least half a century afterwards. Dr. Walker seems to indicate that public nurseries for forest trees began to be established in Scotland between the years 1730 and

1760. The most considerable of these, he says, was that of old Mr. Dickson, at Hassendeanburn, in Teviotdale. This nursery, we are informed by the present proprietors, Messrs. Archibald Dickson and Co., was founded in 1729. From it sprang, in 1767, the nursery of Messrs. Dickson, now Dickson and Turnbull, at Perth; and, subsequently, another brother of the Hassendeanburn family, Walter Dickson, began the house of Dickson and Co. of Edinburgh, now Dicksons and Shankley, in connexion with Mr. James Dickson, who was no relative of the family. It thus appears, that Mr. Robert Dickson of Hassendeanburn was the father of commercial forest tree nurseries in Scotland. The three nurseries established by him and his two brothers being still the most extensive in that country. Mr. Archibald Dickson, the present chief of the firm at Hassendeanburn and at Hawick, to whom we are indebted for the above information, states, in his letter of March 24. 1835, that he is now bringing up some of the fifth generation to the trade. The next considerable public establishment of this kind was that of Messrs. Anderson and Leslie of Broughton Park, Edinburgh; and contemporary with this were those of Mr. Richmond of Leith Walk, of Gordon of Fountainbridge, of Boutcher of Comely Bank, of Messrs. Austen of Glasgow, of Thomas Leslie and Co. of Dundee, of Reid of Aberdeen, of Sampson of Kilmarnock, and a number of others. The most scientific nurseryman in Scotland during the 18th century, appears to have been Mr. Boutcher. According to an authority quoted by Sir Henry Steuart, Mr. Boucher was "the honestest and most judicious nurseryman Scotland ever had." He made an attempt to improve Scottish arboriculture about 1760; but, according to Sir Henry, he was "undervalued by the ignorance of his age, and suffered to languish unsupported for years at Comely Garden, and to die at last in obscurity and indigence." (*Planter's Guide*, 2d edit. p. 399.) Boutcher's *Treatise on raising Forest Trees* was the first work on the subject of its time, and Scottish nurserymen have only produced one work on planting superior to it; namely the edition of *Nicol's Planter's Kalendar*, which was edited, and in great part rewritten, by Mr. Sang of Kirkaldy.

The indigenous trees of Ireland are the same as those of Britain, though such as consider the box, the true service, and the common English elm, truly indigenous to England will not accord with this, as these trees are never found in an apparently wild state in Ireland. Those, on the contrary, who consider the *A'rbutus* and *Erica mediterranea* indigenous to Ireland find them wanting in England, and may hence consider that Ireland has more native trees and shrubs than this country. There can be very little doubt that the common yew is an indigenous tree in Ireland, for trunks of it, of large dimensions,

are frequently dug up from bogs. Mr. Mackay has sent us an account received by him from Mr. Charles Hamilton, Honorary Secretary to the Horticultural Society of Ireland, of one dug up in Queen's County, the rings of annual increase of which indicated a growth of 545 years. The greatest diameter of the trunk of this tree was only 19 in.! The growth appeared to be very slow during the last 300 years, for near the circumference there were about 100 rings within the space of an inch. The root and bark were quite sound, and the stem from which the section was taken was about 12 ft. long, and of tolerably even thickness throughout. Mr. Mackay says that he saw a yew tree in the Island of Innisfallen, on the lower lake of Killarney, which must have been as old as that mentioned by Mr. Hamilton; and which, when he measured it about thirty years ago, was nearly double the dimensions. If the Irish yew be a distinct species, Ireland may claim this fine tree as her own. Our own opinion is, that this yew is nothing more than a variety of the common species. The largest specimens of this tree, the *Taxus hibernica* of Mackay, are in a garden at the village of Cumber, near Belfast: they are about 25 ft. in height, and have, at a distance, the appearance of cypresses. They are supposed to have been planted about 50 years, but their history is unknown.

From information procured for us through the kindness of Lord Viscount Ferrard, we find that there is an upright or Irish yew in a garden at Mayland, near Antrim, 130 years old 25 ft. high; the diameter of the space covered by the branches, 10 ft.: and the diameter of the trunk close to the ground, 3 ft. This tree, and three others in the town, are supposed to have been planted by the Refords, when they first settled in Mayland in 1712. "An upright yew, probably the parent of the above trees, and of all others in this country, grew in Mr. Ferguson's garden. It was cut down about 16 or 17 years ago, by the late Mr. Ledlie; and his son, now in Antrim, has several pieces of furniture which were made from it. In the panel 1 ft. broad, of one of these a wardrobe, I can count about 100 annual concentric layers, and as the tree, it is said, was 2 ft. in diameter, this would give 200 years, and 40 or 50 years more might probably be added for the time when scarcely any enlargement took place."—*L. F. Antrim Castle, March 24. 1835.*

If the arbutus be not indigenous to Ireland, it is at least completely naturalised there, being found, as the yew is in England, in places completely inaccessible to a planter, and where the seeds must have been carried by birds. One of the largest specimens stood in Rough Island, on the lower lake of Killarney, in 1805; it was measured in that year by Mr. Mackay, and the trunk found, at a foot from the ground, to be 9½ ft. in

girt. It formed rather an immense bush than a tree, and consisted of four limbs, the branches of which extended from the root to the length of 36 ft. There is one equally large at Powerscourt, Wicklow, which was planted about 90 years ago; and one, of similar size and age, at Newton Mount Kennedy, was blown down in 1804. The *Erica mediterranea* was found growing, by Mr. Mackay, in Cunnemara, on the western coast. (See *Gard. Mag.*, vol. vii. p. 230, and the forthcoming *Irish Flora* of Mr. Mackay.) *Erica mediterranea* has not only been found on the side of Errisbeg mountain, covering a space of three acres, but in the wild district of Erris, in the county of Mayo, in the greatest profusion. It is a distinct variety from the plant of the same name in gardens, and is considered by Dr. Greville to be the same as that found in the Western Pyrenees.

We have not been able to procure much information respecting the dates of the introduction of foreign trees and shrubs into Ireland, though we have looked over numerous books, and entered into an extensive correspondence for that purpose. On the whole, there appears to have been comparatively few foreign trees planted in Ireland previously to the middle of the 18th century; except fruit trees, and probably some ornamental shrubs, as the arbutus, &c., in the gardens of the monastic institutions, and other religious establishments.

A work, entitled *Botanologia Universalis Hibernica*, by J. K'Eogh, A. B., chaplain to Lord Kingston, published in 1735, appears to contain the names of all the foreign trees and shrubs that were in Ireland at that time.

In the preface to this work, the author says: "When I was writing on this subject, I had the advantage daily of viewing the gardens belonging to the Rt. Hon. James Lord Baron of Kingston, wherein were contained near 200 different species of herbs and trees. I was not acquainted with any garden which could show so many. This was no small advantage or conveniency to forward this undertaking." The trees and shrubs enumerated in K'Eogh's work are the following:—

"Abele, about mansion-houses, for shelter; arbutus, wild in Kerry, and is manured in gardens; great bay; box; chestnut, frequently planted in gardens and parks; cypress tree, in gardens, for its pleasant verdure; fig tree; jasminum, planted in gardens; lemon tree, to be seen in the gardens of Mitchelstown belonging to the Rt. Hon. Lord Kingston; medlar tree, in gardens; myrtle tree, it grows in my Lord Kingston's greenhouse, Mitchelstown, and there are also hedges of it in the Lord Inchiquin's gardens at Rostillon; mulberry tree, in gardens.

"Orange trees; of late years they had been transplanted here, which now, by the industry and cultivation of curious gentlemen, are in some gardens brought to perfection. I have seen about

seventy or eighty oranges taken off one tree in the Rt. Hon. the Lord Kingston's garden at Michelstown, as good as any I have seen brought hither from Spain or the West Indies: so you see what a prolific and fertile soil we live in, where the most exotic plants might, by a little care and industry, flourish.

"Peach tree, in gardens; pear tree; pine tree; rose; savin, in gardens, wild in one of the islands of Lough Lane, Kerry; colutea, in gardens, I have seen it flourishing in Mr. Robert Fennell's garden near Mitchelstown; abrotanum; tamarisk, in gardens; vine tree, in some gardens: walnut tree, in walks, parks, and fields."

A nobleman, whose father was one of the greatest planters in Ireland, to whom we were recommended to apply for authentic information, sent us the following statement:—"The gardens of greatest interest in Ireland, as having been the first to introduce exotic trees and shrubs, and as having contained the greatest variety, were those of Lord Moira, at Moira, in Down [noticed p. 48.]; and of Lord Clanbrassill, at Dundalk, in Louth; and Tollymore Park in Down. Sir Robert Bateson, M.P. for the county of Derry, is proprietor of Moira; and the Earl of Roden, of Dundalk and Tollymore. Moira is dismantled, though some of the trees and shrubs may possibly remain. Dundalk is also dismantled, but Tollymore is kept up. Lord Farnham introduced many foreign trees and shrubs to Newton Barry, and may have lately done so at Farnham. (*February, 1835.*)"

Mr. Mackay, the very intelligent curator of the Trinity College Botanic Garden, Dublin, in a letter dated February, 1835, says:—"The late Lord Oriel and the late Earl of Clanbrassill were the persons who introduced by far the greater number of trees into Ireland during the last century. I think they commenced doing so about 1770, or perhaps a few years before that period; the former, Lord Oriel (then Mr. Foster), planted them in his demesne at Collon, in the county of Louth; and the latter, in his fine demesne at Tollymore Park, in the county of Down." Our friend Mr. Murphy, in the *Irish Farm. and Gard. Mag.* (vol. ii. p. 89.), states that Lord Viscount Ferrard, the son of Lord Oriel, possesses more foreign trees and shrubs than any other individual in Ireland. Mr. Mackay also states that John Templeton, Esq., about the same time as the two noblemen above mentioned, introduced many fine American trees and shrubs into his grounds at Malone, near Belfast, where the same family still reside.

The greatest number of species planted in the 18th century, in any one demesne, is at Oriel Temple, and many of these appear to have grown with very great rapidity. A tulip tree, 40 years planted, has attained the height of 43 ft.; an *A'cer*

rùbrum, of the same age, 44 ft.; a *Pàvia flàva*, of the same age 31 ft.; a *Sophòra japónica*, 50 years planted, 35 ft.; an *Aristotèlia Mácqui*, 20 years planted, upwards of 16 ft., though on a strong clayey soil; the *Arbutus Andráchne* seems to grow at the rate of 1 ft. a year; the Portugal laurel, 50 years planted, has attained the height of 35 ft., and its branches cover a space the diameter of which is 45 ft.; the common laurel, of the same age, is 40 ft. high, and its branches cover a space of 36 ft. in diameter; *Pópulus canadénsis*, 40 years planted, is 72 ft. high: *Alnus laciniàta*, 34 years planted, is 44 ft. high; *Quércus palústris*, 50 years planted, is 41 ft. high; and *Q. fastigiàta*, of the same age, is 54 ft. high; *Q. exoniénsis*, 60 years planted, is 67 ft. high; and *Q. Ægilops*, of the same age, is 55 ft. high; the purple beech, 56 years planted, is 54 ft. high; and that beautiful variety of the common beech, *Fàgus sylvàtica pëndula*, at 35 years' growth, is 33 ft. high; the arbor vitæ, 30 years planted, is 30 ft. high; the *Pinus Cémбра*, of the same age, 34 ft. high; the hemlock spruce, 35 years planted, 32 ft. high; the *Cunninghàmia lanceolàta*, in 12 years, 7 ft. high; the *Làrix pëndula*, in 55 years, 62 ft.; the cedar of Lebanon, in 35 years, 33 ft.; and that singularly picturesque, and yet elegant, tree, the cedar of Goa (*Cuprèssus lusitànica*), 32 ft. high, the branches covering a space of 35 ft. in diameter, in 24 years. This cedar was originally brought from the Portuguese settlement at Goa, in the East Indies, to Portugal; and the seeds were brought from that country to Ireland by Lord Ferrard in 1809; and, being sown, produced abundantly. The plants were first kept in a green-house; but, on some of them being transplanted into the open air from want of room, they were found to grow so vigorously, that in three seasons any one branch surpassed in size the entire plant contained in the green-house. A *Rhododéndron pónticum*, at 60 years of age, is 16 ft. high, and the diameter of the space covered by its branches is 38 ft. There was a tree pæony here of extraordinary dimensions. It was 12 ft. high, and was protected during winter by a glass case. About the year 1827 this case was left off, to try to inure the plant to stand without covering. but the winter unfortunately proving severe, it was killed.

At Antrim Castle, also the seat of Lord Ferrard, are some remarkably fine trees and shrubs. There is a yew tree, estimated at 200 years old, which is 35 ft. high; the diameter of the trunk, at 1 ft. from the ground, is 2 ft. 9 in.; and the diameter of the space covered by its branches, 33 ft. There is a Portugal laurel 150 years planted, which is only 18 ft. high: but the diameter of its trunk, at 1 ft. from the ground, is 3 ft. 7 in.; and that of the space covered by its branches, 36 ft. There are an evergreen oak 100 years old and 25 ft. high; and a variegated holly of the same age, 20 ft. high. There are a juniper 18 ft., at 90 years of age: and an arbor vitæ 24 ft. high, at the same age.

At Tollymore Park, in the county of Down, planted by the Earl of Clanbrassill, and now the seat of the Earl of Roden, there are some very fine trees. The soil and situation, the first ridge of the Mourne Mountains, appear to be particularly suited to the larch and silver fir. From a considerable number, of almost equal magnitude, measured for us by desire of Lord Roden, we select one silver fir, planted 60 years ago, which is 84 ft. high; the diameter, at 1 ft. from the ground, 5 ft. 2 in.; at 10 ft., 4 ft.; and at 24 ft., 3 ft. 3 in.; it is beautifully and evenly clothed with branches, the lower tiers of which are pendent to the ground, and the circumference of the space which they cover is 160 ft. The larch of Tollymore Park is in much estimation for its great teracity, and it supplies masts of from 50 ft. to 60 ft. in length. As a selection from a great number, we take one tree, which, at 80 years of age, is 84 ft. high; the diameter, at 1 ft. from the ground, 2 ft. 8 in.; and 10 ft. from the ground, 2 ft. 3 in. another tree, at 60 years of age, is 66 ft. high; the diameter, at 1 ft. from the ground, being 3 ft.; and at 10 ft., 2 ft. 3 in. Among numerous fine specimens of shrubs introduced by the late Lord Clanbrassill, there is a *Rhododéndron póncticum*, which, at 50 years of age, is 10 ft. high, and covers, with its unbroken mass of foliage, a space the circumference of which is 90 ft. The larch at Tollymore Park is grown on the side of a steep hill facing the north, on a stiff gravelly substratum, which corresponds with the natural situation in which the larch is found in Switzerland, as stated by Decandolle in the *Quarterly Journal of Agriculture*, vol. v. p. 403.; and with the situations in the neighbourhood of Dunkeld, where the best larch is grown by the Duke of Athol, as stated in the account of these plantations in the *Transactions of the Highland Society of Scotland*, vol. xi. p. 165. to p. 219. Monteath, the Scotch forester, we are informed by Lord Roden, and also by another correspondent, considers the Tollymore larch as very superior in quality to the generality of the Scotch or Welch larch. Lord Roden states that he uses it for all purposes whatever, and that for forming utensils it is found an excellent substitute for ash. The trees are generally felled at the age of 70 years. The rhododendrons are scattered through the woods; they are found fully as hardy as the common laurel, and many of them have attained a large size. There are many specimens of *Abies excélsa* var. *Clanbrassilliána*, but none of them remarkable.

At Dundalk, also the property of the Earl of Roden, there is a *Magnòlia acuminàta* 27 ft. high; the circumference of the stem, at 1 ft. from the ground, is 5 ft.: and at 3 ft. from the ground, 4 ft. 6 in.; and the branches covering a space measuring 84 ft. in circumference. There is an oak in the park 60 ft. high; the circumference of the trunk, at 1 ft. from the ground, is 15 ft.;

at 5 ft., 10 ft. ; and at 19 ft., 10 ft. ; and the space covered by its branches is nearly 355 ft. in circumference.

At Cypress Grove, near Dublin, Mr. Mackay informs us, the Dowager Lady Clanbrassill resided from 1770 to 1790, during which period she received a number of foreign trees and shrubs from her son. The dimensions of many of these, the present gardener, Mr. Edward Carrol, has obligingly sent us, at the request of Mr. Mackay. The collection is numerous, and some of the specimens have attained a considerable size. *Robínia Pseùd-Acàcia* is 60 ft. high ; *Laúrus nóbilis*, 30 ft. ; *Juglans règia*, 70 ft. ; *Cárpinus Bétulus*, 90 ft. ; *Quércus Cérris*, 70 ft. ; and *Juníperus virginiana*, 40 ft., &c.

At Moira, according to information kindly sent us by the present proprietor, Sir Robert Bateson, there appear to be very few, if any, of the trees existing that were planted by Sir Arthur Rawdon, about the end of the seventeenth century (see p. 48.). A number are of considerable size, but their ages are unknown ; among these are, a lime tree, which is 85 ft. high, the diameter of the space covered by its branches is 60 ft., and that of its trunk, at 1 ft. from the ground, 4 ft. ; a beech, 110 ft. high, the diameter of the space covered by its branches being 80 ft., and that of its trunk, at 1 ft. from the ground, 4 ft. 4 in. ; a variegated platanus, 50 ft. high ; and a *Plátanus acerifòlia*, 70 ft. high ; *Quércus Ilex*, 45 ft. high, with a trunk 4 ft. in diameter at 1 ft. from the ground, and the diameter of the space covered by its branches, 35 ft. ; a broad-leaved elm, 90 ft. high, the trunk 3 ft. 10 in. in diameter, and the diameter of the space covered by its branches, 60 ft. ; *Gledítschia triacánthos*, 55 ft. high ; a sweet chestnut, 40 ft. high ; the silver fir, 90 ft. high ; and the common yew tree, 45 ft. high, with a trunk 4 ft. in diameter, and the space covered by its branches being 39 ft. in diameter.

The late John Templeton, Esq., A.L.S., was a scientific botanist, as well as a skilful cultivator ; he was the author of several articles on botany, and on other branches of natural history, which appeared in different works, and of some valuable papers on acclimatising plants, published in the *Transactions of the Royal Irish Academy*. A very interesting account of his life, by Dr. Drummond, will be found in our *Magazine of Natural History*, vol. i. p. 403. It appears that Mr. Templeton had a country house at Malone, near Belfast, to which he gave the name of Crann-more, that is, Great tree, in honour of the very fine chestnut trees which are in front of the house, and which were probably planted in the 17th century : it had before been called Orange Grove. Mr. Templeton began to cultivate flowers in 1786, and he laid out an experimental garden in 1793. At the time of his death, which happened in 1826, there were, says his biographer, "collected in this garden, from various

parts of the world, many rare and useful plants, which he endeavoured to naturalise in this climate, by placing them in a soil and situation as near as possible to that to which they had been accustomed. By this means there is now growing in his garden in the open air, a wonderful and curious collection of plants from India, China, North and South America, Siberia, &c., which were formerly kept in the green-house, or even hot-house. All the trees at Cranmore, except the chestnuts and oaks, were raised from seed planted by Mr. Templeton himself, and so great a variety of the natives of the forest, has perhaps never before been collected in so small a place." (*Mag. Nat. H.*, i. 405) Mr. Templeton corresponded with all the principal botanists of his time, and discovered several new plants in Ireland; among others the *Ròsa hibernica*, and *Orobánche rùbra*. The dimensions of some of the more remarkable trees and shrubs at Cranmore have been kindly sent to us by Mrs. Templeton. Among these are, a sweet chestnut, 60 ft. high, with a trunk 15 ft. in circumference at one foot from the ground (the tree from which the place takes its name); *Pìnus Cémbra*, 24 ft. high; *Pìnus Banksiàna*, 17 ft. high; *Pìnus Mùgho*, 11 ft. high; and *A`bies canadénsis*, 16 ft. high. There are, an *A`cer rùbrum* 30 ft. high; a liquidambar, 15 ft. high; a Swedish juniper, 18 ft. high; and a *Ptèlea trifoliàta*, 26 ft. high. The first *Rhododéndron máximum* introduced into Ireland is supposed to have been one planted here, which attained a very great size, but died about three years ago. There is one still existing, which is 9½ ft. high, and the circumference of the space covered by its branches is 37 ft.

In the *History of the County of Down*, published in 1745 (p. 60.), speaking of Bangor, the author says, "the gardens are filled with noble evergreens of a great size, cut in various shapes, among which is an evergreen oak, which, though it grows as a shrub in most other places, is here a tall tree, and of considerable girth." At the same place there is now (1835) a very large mulberry tree, which is very uncommon in Ireland. There was also one about the same size at Castle Ward. At Spring Vale, in the same county, is a very large cork tree, which is now in a state of decay.

At Castle Ward, the seat of Viscount Bangor, is a flowering ash (*O`rnus europæa*), 30 ft. high, and 6 ft. 4 in. in girt at 7 ft. from the ground; it flowers frequently, but not every year. There are, also, an evergreen oak, with a trunk 9 ft. 6 in. in girt at 2 ft. from the ground, and 8 ft. 4 in. at 10 ft. from the ground; an arbutus, 5 ft. in girt at 2 ft. from the ground; and a pinaster, 60 ft. high, and 8 ft. 10 in. in circumference at 6 ft. from the ground: these trees are all close to a small bay or arm of the sea. There are, also, a silver fir, 66 ft. high, 8 ft. 5 in. in girt, which it carries up to 30 ft.; a cedar of Lebanon,

at 5 ft., 10 ft. ; and at 19 ft., 10 ft. ; and the space covered by its branches is nearly 355 ft. in circumference.

At Cypress Grove, near Dublin, Mr. Mackay informs us, the Dowager Lady Clanbrassill resided from 1770 to 1790, during which period she received a number of foreign trees and shrubs from her son. The dimensions of many of these, the present gardener, Mr. Edward Carrol, has obligingly sent us, at the request of Mr. Mackay. The collection is numerous, and some of the specimens have attained a considerable size. *Robinia Pseùd-Acacia* is 60 ft. high ; *Laúrus nóbilis*, 30 ft. ; *Jùglans règia*, 70 ft. ; *Cárpinus Bétulus*, 90 ft. ; *Quércus Cérris*, 70 ft. ; and *Juníperus virginiàna*, 40 ft., &c.

At Moira, according to information kindly sent us by the present proprietor, Sir Robert Bateson, there appear to be very few, if any, of the trees existing that were planted by Sir Arthur Rawdon, about the end of the seventeenth century (see p. 48.). A number are of considerable size, but their ages are unknown ; among these are, a lime tree, which is 85 ft. high, the diameter of the space covered by its branches is 60 ft., and that of its trunk, at 1 ft. from the ground, 4 ft. ; a beech, 110 ft. high, the diameter of the space covered by its branches being 80 ft., and that of its trunk, at 1 ft. from the ground, 4 ft. 4 in. ; a variegated platanus, 50 ft. high ; and a *Plátanus acerifolia*, 70 ft. high ; *Quércus I'lex*, 45 ft. high, with a trunk 4 ft. in diameter at 1 ft. from the ground, and the diameter of the space covered by its branches, 35 ft. ; a broad-leaved elm, 90 ft. high, the trunk 3 ft. 10 in. in diameter, and the diameter of the space covered by its branches, 60 ft. ; *Gledítschia triacánthos*, 55 ft. high ; a sweet chestnut, 40 ft. high ; the silver fir, 90 ft. high ; and the common yew tree, 45 ft. high, with a trunk 4 ft. in diameter, and the space covered by its branches being 39 ft. in diameter.

The late John Templeton, Esq., A.L.S., was a scientific botanist, as well as a skilful cultivator ; he was the author of several articles on botany, and on other branches of natural history, which appeared in different works, and of some valuable papers on acclimatising plants, published in the *Transactions of the Royal Irish Academy*. A very interesting account of his life, by Dr. Drummond, will be found in our *Magazine of Natural History*, vol. i. p. 403. It appears that Mr. Templeton had a country house at Malone, near Belfast, to which he gave the name of Crann-more, that is, Great tree, in honour of the very fine chestnut trees which are in front of the house, and which were probably planted in the 17th century : it had before been called Orange Grove. Mr. Templeton began to cultivate flowers in 1786, and he laid out an experimental garden in 1793. At the time of his death, which happened in 1826, there were, says his biographer, "collected in this garden, from various

parts of the world, many rare and useful plants, which he endeavoured to naturalise in this climate, by placing them in a soil and situation as near as possible to that to which they had been accustomed. By this means there is now growing in his garden in the open air, a wonderful and curious collection of plants from India, China, North and South America, Siberia, &c., which were formerly kept in the green-house, or even hot-house. All the trees at Cranmore, except the chestnuts and oaks, were raised from seed planted by Mr. Templeton himself, and so great a variety of the natives of the forest, has perhaps never before been collected in so small a place." (*Mag. Nat. H.*, i. 405.) Mr. Templeton corresponded with all the principal botanists of his time, and discovered several new plants in Ireland; among others the *Ròsa hibernica*, and *Orobánche rubra*. The dimensions of some of the more remarkable trees and shrubs at Cranmore have been kindly sent to us by Mrs. Templeton. Among these are, a sweet chestnut, 60 ft. high, with a trunk 15 ft. in circumference at one foot from the ground (the tree from which the place takes its name); *Pinus Cembra*, 24 ft. high; *Pinus Banksiana*, 17 ft. high; *Pinus Mughò*, 11 ft. high; and *Abies canadensis*, 16 ft. high. There are, an *Acer rubrum* 30 ft. high; a liquidambar, 15 ft. high; a Swedish juniper, 18 ft. high; and a *Ptelea trifoliata*, 26 ft. high. The first *Rhododéndron máximum* introduced into Ireland is supposed to have been one planted here, which attained a very great size, but died about three years ago. There is one still existing, which is 9½ ft. high, and the circumference of the space covered by its branches is 37 ft.

In the *History of the County of Down*, published in 1745 (p. 60.), speaking of Bangor, the author says, "the gardens are filled with noble evergreens of a great size, cut in various shapes, among which is an evergreen oak, which, though it grows as a shrub in most other places, is here a tall tree, and of considerable girth." At the same place there is now (1835) a very large mulberry tree, which is very uncommon in Ireland. There was also one about the same size at Castle Ward. At Spring Vale, in the same county, is a very large cork tree, which is now in a state of decay.

At Castle Ward, the seat of Viscount Bangor, is a flowering ash (*Ornus europæa*), 30 ft. high, and 6 ft. 4 in. in girth at 7 ft. from the ground; it flowers frequently, but not every year. There are, also, an evergreen oak, with a trunk 9 ft. 6 in. in girth at 2 ft. from the ground, and 8 ft. 4 in. at 10 ft. from the ground; an arbutus, 5 ft. in girth at 2 ft. from the ground; and a pinaster, 60 ft. high, and 8 ft. 10 in. in circumference at 6 ft. from the ground: these trees are all close to a small bay or arm of the sea. There are, also, a silver fir, 66 ft. high, 8 ft. 5 in. in girth, which it carries up to 30 ft.; a cedar of Lebanon,

50 ft. high, 5 ft. 3 in. in girt at 8 ft. from the ground; and a sweet chestnut, 10 ft. 3 in. in girt at 3 ft. from the ground. There is a myrtle hedge here at least 120 years old, which grows vigorously. The shrubberies and ornamental planting at Castle Ward were made by Mrs. Ward, the wife of Judge Ward, between 1710 and 1759; and some before that period, as there were some tulip trees of large size cut down some years ago, supposed to be 120 years old — *J.M.R.*

To Mr. Carrol, gardener at Cypress Grove, we are indebted for the measurements of several trees at Howth Castle, near Dublin, the seat of the Earl of Howth. It appears that some foreign trees were planted here even in the 16th century, and particularly an *Ulmus campestris*, which is estimated to be 250 years old. It is only 50 ft. high, but the diameter of the trunk, at 1 ft. from the ground, is 4 ft. 6 in. There is a walnut tree here, considered to be 200 years planted; a *Tamarix gallica*, 100 years; and a tulip tree, 60 years. The tamarisk has a stem 1 ft. 10 in. in diameter at 1 ft. from the ground; it is 20 ft. high, and the diameter of the space covered by its branches is 22 ft. We should suppose it must be the finest specimen of this shrub in existence. The common myrtle stands the open air at Howth, protected by a wall; there is a specimen which has been planted thirty years, which has attained a stem 4 in. in diameter at 1 ft. from the ground.

At Charleville Forest, in King's County, a place where we had the pleasure of staying two or three days in 1811, when consulted professionally by the Earl of Charleville, there is a common lime, supposed to have been planted about seventy years, which is now (1835) 110 ft. high; an *Acer platanoides*, which at 60 years is 68 ft. high; a *Pavia rubra*, 76 ft. high; a common holly, 45 ft. high; a *Robinia Pseud-Acacia*, 50 ft. high; a *Cratægus Azarolus*, 40 ft. high; an *Ulmus campestris*, 85 ft. high; a *Pópulus álba*, 120 ft. high; a *Quercus pedunculata*, planted 60 years, which is 110 ft. high; a *Fagus sylvatica* of the same age and height; a sweet chestnut, 45 years planted, which has attained the height of 85 ft.; a yew tree, 45 years planted, which has attained the height of 50 ft.; an arbor vitæ, planted 25 years, and 20 ft. high; and a *Làrix microcarpa*, 45 years planted, and 94 ft. high. On the whole, there is an excellent collection of trees at Charleville, and they appear to have made extraordinary progress.

At Shelton Abbey, the Earl of Wicklow's, in the county of Wicklow, there are a few remarkably fine specimens of foreign trees and shrubs. A tulip tree, 50 years planted, is 60 ft. high, flowering beautifully every year; a *Robinia Pseud-Acacia*, of the same age, is 65 ft. high, with a trunk $2\frac{1}{2}$ ft. in diameter at 1 ft. from the ground; a Portugal laurel, 40 years planted, is 35 ft. high, has a stem $2\frac{1}{2}$ ft. in diameter at 1 ft. from the ground,

and the diameter of the space covered by its branches is 39 ft. ; a common laurel, 90 years planted, is 45 ft. high. the diameter of its trunk, at 1 ft. from the ground, is 6 ft. [probably from the number of diverging branches proceeding direct from the crown of the root], and the diameter of the space its branches cover is 101 ft. ! A *Záurus nóbilis*, 16 years planted, is 34 ft. high, and the diameter of the space covered by its branches is 25 ft. ; *Cuprèssus sempervirens*, 50 years planted, is 59 ft. high.

At Castle Freke, in the county of Cork, the seat of Lord Carberry, there appear to be some fine specimens. *Rhododéndron pónticum* is 8 ft. high, and the branches cover a space 76 ft. in circumference ; the *Quércus Ilex*, 26 years planted, 36 ft. high ; the Lucombe oak, of the same age, 39 ft. high ; and the sweet chestnut, 44 ft. high ; *Aristotèlia Mácqui*, on light soil over gravel, forms a handsome tree, 26 ft. high, with a trunk $8\frac{1}{2}$ in. in diameter at 1 ft. from the ground.

At Florence Court, the residence of the Earl of Enniskillen, there is a good collection of trees, the dimensions of many of which have been sent us by the gardener there, Mr. Young. The tulip tree, at 38 years' growth, is 35 ft. high ; *A'cer montànum*, at 38 years' growth, 50 ft. high ; the Portugal laurel, at 40 years' growth, is 32 ft. high, and its branches cover a space 22 ft. in diameter ; *Córnus flórida*, 38 years planted, is 16 ft. high, and the branches cover a space of 20 ft. in diameter ; *Sambucus nigra*, at 40 years of age, is 50 ft. high ; the walnut at 50 years of age, is 40 ft. high ; and the Canadian poplar, at 30 years' growth, 70 ft. high ; the scarlet oak, 40 years planted, is 70 ft. high ; and different varieties of *Quércus Cérris*, all planted 38 years, are also 70 ft. high ; the common yew, at 80 years of age, is 30 ft. high, and its branches cover a space of 30 ft. in diameter ; and the Irish yew, sometimes called the Florence Court yew, the original plant being still in existence in the grounds, has attained the height of 26 ft.

At Killruderly House, in the county of Wicklow, the seat of the Earl of Meath, are some remarkably fine evergreen oaks, One of these, by no means larger than the rest, measured for us by Mr. Niven in February, 1835, was 60 ft. high, with a trunk $11\frac{1}{2}$ ft. in circumference at 1 ft. from the ground.

The oldest cedars of Lebanon in Ireland are said to be at Mount Anville Hill, the seat of Counsellor West. K. C. These cedars, we are informed, were brought direct from Mount Lebanon, by an ancestor of Lord Tremblestown. We have tried in vain to get their dimensions, for which reason we suspect they are not very remarkable. The finest, we believe, are at Castletown, Kildare, the seat of Edward Conolly, Esq., M.P. One of these, Colonel Conolly informs us, is 13 ft. in girth at 1 ft. from the ground, and has a clear stem of 30 ft.

50 ft. high, 5 ft. 3 in. in girth at 8 ft. from the ground; and a sweet chestnut, 10 ft. 3 in. in girth at 3 ft. from the ground. There is a myrtle hedge here at least 120 years old, which grows vigorously. The shrubberies and ornamental planting at Castle Ward were made by Mrs. Ward, the wife of Judge Ward, between 1710 and 1759; and some before that period, as there were some tulip trees of large size cut down some years ago, supposed to be 120 years old — *J.M.R.*

To Mr. Carrol, gardener at Cypress Grove, we are indebted for the measurements of several trees at Howth Castle, near Dublin, the seat of the Earl of Howth. It appears that some foreign trees were planted here even in the 16th century, and particularly an *Ulmus campestris*, which is estimated to be 250 years old. It is only 50 ft. high, but the diameter of the trunk, at 1 ft. from the ground, is 4 ft. 6 in. There is a walnut tree here, considered to be 200 years planted; a *Tamarix gallica*, 100 years; and a tulip tree, 60 years. The tamarisk has a stem 1 ft. 10 in. in diameter at 1 ft. from the ground; it is 20 ft. high, and the diameter of the space covered by its branches is 22 ft. We should suppose it must be the finest specimen of this shrub in existence. The common myrtle stands the open air at Howth, protected by a wall; there is a specimen which has been planted thirty years, which has attained a stem 4 in. in diameter at 1 ft. from the ground.

At Charleville Forest, in King's County, a place where we had the pleasure of staying two or three days in 1811, when consulted professionally by the Earl of Charleville, there is a common lime, supposed to have been planted about seventy years, which is now (1835) 110 ft. high; an *Acer platanoides*, which at 60 years is 68 ft. high; a *Pavia rubra*, 76 ft. high; a common holly, 45 ft. high; a *Robinia Pseud-Acacia*, 50 ft. high; a *Crataegus Azarolus*, 40 ft. high; an *Ulmus campestris*, 85 ft. high; a *Populus alba*, 120 ft. high; a *Quercus pedunculata*, planted 60 years, which is 110 ft. high; a *Fagus sylvatica* of the same age and height; a sweet chestnut, 45 years planted, which has attained the height of 85 ft.; a yew tree, 45 years planted, which has attained the height of 50 ft.; an arbor vitæ, planted 25 years, and 20 ft. high; and a *Larix microcarpa*, 45 years planted, and 94 ft. high. On the whole, there is an excellent collection of trees at Charleville, and they appear to have made extraordinary progress.

At Shelton Abbey, the Earl of Wicklow's, in the county of Wicklow, there are a few remarkably fine specimens of foreign trees and shrubs. A tulip tree, 50 years planted, is 60 ft. high, flowering beautifully every year; a *Robinia Pseud-Acacia*, of the same age, is 65 ft. high, with a trunk $2\frac{1}{2}$ ft. in diameter at 1 ft. from the ground; a Portugal laurel, 40 years planted, is 35 ft. high, has a stem $2\frac{1}{2}$ ft. in diameter at 1 ft. from the ground,

and the diameter of the space covered by its branches is 39 ft. ; a common laurel, 90 years planted, is 45 ft. high. the diameter of its trunk, at 1 ft. from the ground, is 6 ft. [probably from the number of diverging branches proceeding direct from the crown of the root], and the diameter of the space its branches cover is 101 ft. ! A *Laúrus nóbilis*, 16 years planted, is 34 ft. high, and the diameter of the space covered by its branches is 25 ft ; *Cuprèssus sempervirens*, 50 years planted, is 59 ft high.

At Castle Freke, in the county of Cork, the seat of Lord Carberry, there appear to be some fine specimens. *Rhododéndron pónticum* is 8 ft. high, and the branches cover a space 76 ft. in circumference ; the *Quércus Flex*, 26 years planted, 36 ft. high ; the Lucombe oak, of the same age, 39 ft. high ; and the sweet chestnut, 44 ft. high ; *Aristotèlia Mácqui*, on light soil over gravel, forms a handsome tree, 26 ft. high, with a trunk 8½ in. in diameter at 1 ft. from the ground.

At Florence Court, the residence of the Earl of Enniskillen, there is a good collection of trees, the dimensions of many of which have been sent us by the gardener there, Mr. Young. The tulip tree, at 38 years' growth, is 35 ft. high ; *A'cer montànum*, at 38 years' growth, 50 ft. high ; the Portugal laurel, at 40 years' growth, is 32 ft. high, and its branches cover a space 22 ft. in diameter ; *Córnus flórida*, 38 years planted, is 16 ft. high, and the branches cover a space of 20 ft. in diameter ; *Sambucus nigra*, at 40 years of age, is 50 ft. high ; the walnut at 50 years of age, is 40 ft. high ; and the Canadian poplar, at 30 years' growth, 70 ft. high ; the scarlet oak, 40 years planted, is 70 ft. high ; and different varieties of *Quércus Cérris*, all planted 38 years, are also 70 ft. high ; the common yew, at 80 years of age, is 30 ft. high, and its branches cover a space of 30 ft. in diameter ; and the Irish yew, sometimes called the Florence Court yew, the original plant being still in existence in the grounds, has attained the height of 26 ft.

At Killrudy House, in the county of Wicklow, the seat of the Earl of Meath, are some remarkably fine evergreen oaks, One of these, by no means larger than the rest, measured for us by Mr. Niven in February, 1835, was 60 ft. high, with a trunk 11½ ft. in circumference at 1 ft. from the ground.

The oldest cedars of Lebanon in Ireland are said to be at Mount Anville Hill, the seat of Counsellor West. K. C. These cedars, we are informed, were brought direct from Mount Lebanon, by an ancestor of Lord Tremblestown. We have tried in vain to get their dimensions, for which reason we suspect they are not very remarkable. The finest, we believe, are at Castletown, Kildare, the seat of Edward Conolly, Esq., M.P. One of these, Colonel Conolly informs us, is 13 ft. in girt at 1 ft. from the ground, and has a clear stem of 30 ft.

The largest old oak tree in Ireland, Sir Robert Bateson informs us, is at his residence, Belvoir Park, near Belfast. It measures about 28 ft. in girt at 6 ft. from the ground; but it is split, and much damaged. It is supposed to be between two and three centuries old. It grows about 50 yards from the banks of the river Lagan, in rather moist soil.

At Hillsborough, the seat of the Marquess of Downshire, in Lady Downshire's garden, a tulip tree carries up the girt of 4 ft. 6 in. to the height of 8 ft., when it branches off. It flowers abundantly, and has flowered for many years past. Close to this tree is a *Magnolia acuminata* 25 ft. high, and 4 ft. 4 in. in girt at 2 ft. from the ground, where it branches; it does not flower every year, but in hot summers very abundantly. There is, also, a cedar 8 ft. 8 in. in circumference at 3½ ft. from the ground, where it begins to branch. It is not tall, and is quite flat at the top. There are several other forest trees, and some shrubs of about the same age, or perhaps older, in the grounds, particularly some very fine evergreen oaks.—*J.M.R.*

No Dr. Walker has yet arisen in Ireland to determine the dates of the introduction of particular species, and all that we have been able to do, therefore, is to place before our readers the foregoing statements. From these it appears that more had been done in Ireland in the way of introducing foreign trees and shrubs, previously to the middle of the 18th century, than is generally imagined; that a good deal has been done since; and that there is every encouragement to proceed, from the extraordinary rapidity of the growth of the trees that have been planted. There are also the greatest inducements, in point of climate, as will appear in our succeeding subsection, when we give a list of what are green-house trees and shrubs in England, but which stand the open air in Ireland.

Nurseries were probably established in Ireland about the time when it became fashionable to plant trees. The oldest we know of is that of Toole and Co. at Cullenswood near Dublin, and at Shank Hill near Bray. In both gardens are some very fine specimens of foreign trees and shrubs. At Cullenswood, *Magnolia grandiflora* has attained the height of 17 ft. in 20 years, and *M. Thompsoniana*, 15 ft. in 6 years; *Arbutus Andrachne*, and *A. hybrida*, 19 ft. in 27 years; *Olea excelsa*, as a standard, 17 ft. in 27 years; *Pittosporum Tobira* as a standard, 10 ft. in 20 years; *Yucca gloriosa*, 8 ft. in 30 years, with a stem a foot in diameter; *Aralia spinosa*, 20 ft. in 20 years; *Eriobotrya japonica*, 20 ft. in 20 years; *Pyrus [Sorbus] nipalensis*, 16 ft. in 7 years; *Laurus nobilis*, 25 ft. in 35 years; and *Pæonia Moultan*, 8 ft. in 20 years. In the Shank Hill Nursery there is an *Arbutus Andrachne*, 20 ft. high, with a head covering a space 30 yards in circumference, 30 years old.

The nursery of Mr. Hodgins at Dunganstown, near Wicklow, was established about 1780, and was well stocked with foreign trees and shrubs. Mr. Niven kindly measured some of the largest of these for us in February, 1835, and a copious list has been sent to us by the proprietor, Mr. Hodgins, through Mr. Mackay. The *Cupressus lusitánica* in this nursery, 54 years planted, is 20 ft. high, with a trunk, at 1 ft. from the ground, 5 ft. in girth, and the branches covering a space the circumference of which is upwards of 120 ft. This is probably the finest in Ireland, next to Lord Ferrard's, mentioned p. 109. There is a hedge of evergreen oaks in this nursery 50 ft. high. There are several large silver firs, with trunks which girth 6 ft. and 7 ft., which have grown to the height of 60 ft.; red cedars 20 ft., and laurels and bays 30 ft. high; there is a Lucombe oak 50 ft. high, with a trunk 18 in. in diameter at 10 ft. from the ground; the cedar of Lebanon, 45 years planted, is from 30 ft. to 35 ft. high; the Portugal laurel is 30 ft. high; the timber of this tree, Mr. Hodgins observes, is better than that of the cherry. There are many pines from 20 ft. to 30 ft. high; aristotelias, 20 ft. high; *Olea excélsa*, 12 ft. high; Norway maple, the wood of which, Mr. Hodgins observes, is as hard as box; and the sugar maple, growing as vigorously as the common sycamore. All these trees, and many others, were planted by the present proprietor, who, Mr. Niven informs us, is a most enthusiastic and successful cultivator, who has done, perhaps, more in Ireland, in the way of cultivating rare trees and shrubs, than any other contemporary; and who, though of an advanced age, is still healthy and vigorous, and derives the greatest enjoyment from the exercise of his profession.

The nursery of Mr. Robertson, at Kilkenny, was founded by the father of the present proprietor about 1765, who introduced most of the foreign trees and shrubs cultivated about that time in the London nurseries. Though most of these were used as stools for propagation, yet a few of them have been allowed to run up as specimens. Among these is a *Córylus Colúrna*, which, at 50 years' growth, is 3 ft. 7 in. in girth at 1 ft. from the ground; it is 25 ft. high, and the diameter of its head is nearly 50 ft. Besides this, an *Ailántus glandulòsa*, a Judas tree, and several others, are worthy of notice; the details of which will be found in the *Gard. Mag.*, vol. xi. p. 210. Most of the other nurseries in Ireland were founded, we believe, in the succeeding century.

The establishment of a Botanic Garden at Glasnevin must have given a considerable stimulus to the introduction of foreign trees and shrubs into Ireland. This garden owes its origin, in 1797, to the late Lord Oriel. The plan of the garden, Mr. Mackay informs us, "was also suggested by his Lordship, but the laying out and arrangement were the work of Mr. Under-

The largest old oak tree in Ireland, Sir Robert Bateson informs us, is at his residence, Belvoir Park, near Belfast. It measures about 28 ft. in girt at 6 ft. from the ground; but it is split, and much damaged. It is supposed to be between two and three centuries old. It grows about 50 yards from the banks of the river Lagan, in rather moist soil.

At Hillsborough, the seat of the Marquess of Downshire, in Lady Downshire's garden, a tulip tree carries up the girt of 4 ft. 6 in. to the height of 8 ft., when it branches off. It flowers abundantly, and has flowered for many years past. Close to this tree is a *Magnòlia acuminàta* 25 ft. high, and 4 ft. 4 in. in girt at 2 ft. from the ground, where it branches; it does not flower every year, but in hot summers very abundantly. There is, also, a cedar 8 ft. 8 in. in circumference at 3½ ft. from the ground, where it begins to branch. It is not tall, and is quite flat at the top. There are several other forest trees, and some shrubs of about the same age, or perhaps older, in the grounds, particularly some very fine evergreen oaks.—*J.M.R.*

No Dr. Walker has yet arisen in Ireland to determine the dates of the introduction of particular species, and all that we have been able to do, therefore, is to place before our readers the foregoing statements. From these it appears that more had been done in Ireland in the way of introducing foreign trees and shrubs, previously to the middle of the 18th century, than is generally imagined; that a good deal has been done since; and that there is every encouragement to proceed, from the extraordinary rapidity of the growth of the trees that have been planted. There are also the greatest inducements, in point of climate, as will appear in our succeeding subsection, when we give a list of what are green-house trees and shrubs in England, but which stand the open air in Ireland.

Nurseries were probably established in Ireland about the time when it became fashionable to plant trees. The oldest we know of is that of Toole and Co. at Cullenswood near Dublin, and at Shank Hill near Bray. In both gardens are some very fine specimens of foreign trees and shrubs. At Cullenswood, *Magnòlia grandiflòra* has attained the height of 17 ft. in 20 years, and *M. Thompsoniàna*, 15 ft. in 6 years; *Arbutus Andràchne*, and *A. hýbrida*, 19 ft. in 27 years; *Olea excélsa*, as a standard, 17 ft. in 27 years; *Pittòsporum Tobira* as a standard, 10 ft. in 20 years; *Yúcca gloriòsa*, 8 ft. in 30 years, with a stem a foot in diameter; *Aràlia spinòsa*, 20 ft. in 20 years; *Eriobótrya japónica*, 20 ft. in 20 years; *Pýrus* [*Sórbus*] *nipalénsis*, 16 ft. in 7 years; *Laúrus nóbilis*, 25 ft. in 35 years; and *Pædnia Moútan*, 8 ft. in 20 years. In the Shank Hill Nursery there is an *Arbutus Andràchne*, 20 ft. high, with a head covering a space 30 yards in circumference, 30 years old.

The nursery of Mr. Hodgins at Dunganstown, near Wicklow, was established about 1780, and was well stocked with foreign trees and shrubs. Mr. Niven kindly measured some of the largest of these for us in February, 1835, and a copious list has been sent to us by the proprietor, Mr. Hodgins, through Mr. Mackay. The *Cuprèssus lusitànica* in this nursery, 54 years planted, is 20 ft. high, with a trunk, at 1 ft. from the ground, 5 ft. in girt, and the branches covering a space the circumference of which is upwards of 120 ft. This is probably the finest in Ireland, next to Lord Ferrard's, mentioned p. 109. There is a hedge of evergreen oaks in this nursery 50 ft. high. There are several large silver firs, with trunks which girt 6 ft. and 7 ft., which have grown to the height of 60 ft.; red cedars 20 ft., and laurels and bays 30 ft. high; there is a Lucombe oak 50 ft. high, with a trunk 18 in. in diameter at 10 ft. from the ground; the cedar of Lebanon, 45 years planted, is from 30 ft. to 35 ft. high; the Portugal laurel is 30 ft. high; the timber of this tree, Mr. Hodgins observes, is better than that of the cherry. There are many pines from 20 ft. to 30 ft. high; aristotelias, 20 ft. high; *O'lea excèlsa*, 12 ft. high; Norway maple, the wood of which, Mr. Hodgins observes, is as hard as box; and the sugar maple, growing as vigorously as the common sycamore. All these trees, and many others, were planted by the present proprietor, who, Mr. Niven informs us, is a most enthusiastic and successful cultivator, who has done, perhaps, more in Ireland, in the way of cultivating rare trees and shrubs, than any other contemporary; and who, though of an advanced age, is still healthy and vigorous, and derives the greatest enjoyment from the exercise of his profession.

The nursery of Mr. Robertson, at Kilkenny, was founded by the father of the present proprietor about 1765, who introduced most of the foreign trees and shrubs cultivated about that time in the London nurseries. Though most of these were used as stools for propagation, yet a few of them have been allowed to run up as specimens. Among these is a *Córylus Colúrna*, which, at 50 years' growth, is 3 ft. 7 in. in girt at 1 ft. from the ground; it is 25 ft. high, and the diameter of its head is nearly 50 ft. Besides this, an *Ailántus glandulòsa*, a Judas tree, and several others, are worthy of notice; the details of which will be found in the *Gard. Mag.*, vol. xi. p. 210. Most of the other nurseries in Ireland were founded, we believe, in the succeeding century.

The establishment of a Botanic Garden at Glasnevin must have given a considerable stimulus to the introduction of foreign trees and shrubs into Ireland. This garden owes its origin, in 1797, to the late Lord Oriel. The plan of the garden, Mr. Mackay informs us, "was also suggested by his Lordship, but the laying out and arrangement were the work of Mr. Under-

wood, the late intelligent curator. The arboretum, which was laid out and planted by him in 1798-9, does him lasting honour." Of the *Pinus Pallasiana*, there are two fine specimens, the largest 40 ft. high, with a trunk 2 ft. in diameter at 1 ft. from the ground, which are probably the best to be met with in Britain or Ireland. What is remarkable in British nurseries, though common in French ones, there is a *Pinus Cémbra*, which was grafted on a Scotch pine about 20 years ago, and now forms a fine tree. The garden contains 30 statute acres; it is described and engraved in Dr. Walsh's *History of Dublin*, and on the whole, is not only the largest in Europe, but the most comprehensive in its plan. Since the death of Mr. Underwood, in 1834, it has been put under the curatorship of Mr. Niven, one of the most scientific gardeners and active-minded men in the profession. Mr. Niven's plan for the improvement and future management of the Glasnevin Garden is intended to include, not merely the extension of the arrangements of plants botanically, but also an agricultural and horticultural selection of hardy fruits and vegetables, for the purpose of following up the important subject of improving, by crossing, the best existing varieties of such fruits and vegetables.

SUBJECT. 5. *Of the Foreign Trees and Shrubs introduced into Britain in the 19th Century.*

DURING that portion of the nineteenth century which has now (1835) elapsed, the taste for foreign trees and shrubs has considerably increased among planters: and the number of new species and varieties that have been introduced, is proportionately greater than at any former period. Botanic gardens and arboretums have also become more general, and the variety introduced into shrubberies and ornamental plantations, though still not so great as it might be, bears some relation to the general improvement. The establishment of the Horticultural Society of London in the early part of the century, has had a material influence in spreading a taste for every department of gardening, not only in Britain, but throughout the civilised world. The interest, however, which belongs to this century, is greatly diminished to the present generation of readers, from the circumstance of the greater part of it being within their recollection. For this reason we shall limit ourselves to giving a short comparative view of the species of trees and shrubs which have been introduced, and a slight notice of the principal arboretums which have been formed; taking, as our authority for the date of the introduction of the trees and shrubs, our *Hortus Britannicus*.

In the first decade of the nineteenth century, viz., from 1801 to 1810 inclusive, ninety-four trees and shrubs were introduced:

eight by Conrad Loddiges; six by Messrs. Lee and Kennedy; three by Fraser; nineteen by Lyon; one by the Kew Garden; one by the London Horticultural Society; one by Don of the Cambridge Botanic Garden; and one by Sir Abraham Hume. Among the most interesting articles introduced during this decade are, *Rosa multiflora*, *Cunninghãmia lanceolata*, *Juniperus excelsa*, *Caprifolium japonicum*, *Rosa Banksia* *Rhododendron catawbiense* (by Fraser), and *Crataegus Aronia*. It is somewhat remarkable, that of such a number of species introduced during this decade, the names of so few of the introducers should be known; but it must be recollected that the means of introducing were, at this period, principally by packets of seeds sent to the nurserymen by foreign correspondents, or by amateurs; and that, as several years must necessarily elapse between the period of introduction, and that of flowering and naming, the name of the collector who sent the seeds, or of the nurseryman who first raised plants from them, is forgotten, or ceases to be of the same interest. The case is different when living plants are brought into the country, and it is, in truth, chiefly of the introducers of such that the names are known.

From 1811 to 1820, three hundred and seventy-four trees and shrubs were introduced, viz., forty-four by Messrs. Loddiges; twelve by Lyon; four by Lee and Kennedy; three by Whitley and Co. among which was *Spiræa bella* in 1820; three by the Horticultural Society (including *Cotoneaster affinis* in 1820); two by Fraser (*Abies Fraseri*, and *Yucca angustifolia* in 1811); one by Don of the Cambridge Botanic Garden: one (the *Ribes sanguineum*, in 1817) by Archibald Menzies, Esq., who sailed round the world with Captain Vancouver; *Genista procumbens* by Schleicher, a botanical collector in Switzerland; one by Knight of the Exotic Nursery, King's Road; and one (*Mahonia fascicularis*) by A. B. Lambert, Esq. Among the most valuable of the species introduced by Loddiges are, *Azalea arborescens*, *A. speciosa*, and *Ribes aureum*, in 1812; *Symphoria racemosa*, *Cytisus ruthenicus*, *Juniperus recurva*, and *Yucca tenuiflora*, in 1817; *Alnus cordifolia* (the most beautiful species of the genus), in 1818; *Armeniaca brigantia*, and *Quercus stellata*, in 1819; *Crataegus melanocarpa*, *C. latifolia*, *C. Olivieriana*, *Fraxinus pannosa*, *F. platycarpa*, *F. lancea*, *Pinus excelsa*, and *Abies Pichta*, in 1820. Among those introduced by Lyon are, *Magnolia pyramidata*, in 1811; and *Andrómèda floribunda*, *Nýssa cándicans*, *Bòrya ligustrina*, *B. porulosa*, *B. acuminata*, *Virgilia lutea*, and *Crataegus apiifolia*, in 1812. Among the fine plants recorded as having been introduced in this decade, without mentioning the names of the introducers, are, *Æsculus glabra* and *pállida*, and *Pàvia híbrida*, in 1812; *Berberis sinensis*, *Cydònia japonica*, and *Dáphne Thymelæa*, in 1815; *Plánera Richárdi*

wood, the late intelligent curator. The arboretum, which was laid out and planted by him in 1798-9, does him lasting honour." Of the *Pinus Pallasiana*, there are two fine specimens, the largest 40 ft. high, with a trunk 2 ft. in diameter at 1 ft. from the ground, which are probably the best to be met with in Britain or Ireland. What is remarkable in British nurseries, though common in French ones, there is a *Pinus Cémbra*, which was grafted on a Scotch pine about 20 years ago, and now forms a fine tree. The garden contains 30 statute acres; it is described and engraved in Dr. Walsh's *History of Dublin*, and on the whole, is not only the largest in Europe, but the most comprehensive in its plan. Since the death of Mr. Underwood, in 1834, it has been put under the curatorship of Mr. Niven, one of the most scientific gardeners and active-minded men in the profession. Mr. Niven's plan for the improvement and future management of the Glasnevin Garden is intended to include, not merely the extension of the arrangements of plants botanically, but also an agricultural and horticultural selection of hardy fruits and vegetables, for the purpose of following up the important subject of improving, by crossing, the best existing varieties of such fruits and vegetables.

SUBJECT. 5. *Of the Foreign Trees and Shrubs introduced into Britain in the 19th Century.*

DURING that portion of the nineteenth century which has now (1835) elapsed, the taste for foreign trees and shrubs has considerably increased among planters: and the number of new species and varieties that have been introduced, is proportionately greater than at any former period. Botanic gardens and arboreta have also become more general, and the variety introduced into shrubberies and ornamental plantations, though still not so great as it might be, bears some relation to the general improvement. The establishment of the Horticultural Society of London in the early part of the century, has had a material influence in spreading a taste for every department of gardening, not only in Britain, but throughout the civilised world. The interest, however, which belongs to this century, is greatly diminished to the present generation of readers, from the circumstance of the greater part of it being within their recollection. For this reason we shall limit ourselves to giving a short comparative view of the species of trees and shrubs which have been introduced, and a slight notice of the principal arboreta which have been formed; taking, as our authority for the date of the introduction of the trees and shrubs, our *Hortus Britannicus*.

In the first decade of the nineteenth century, viz., from 1801 to 1810 inclusive, ninety-four trees and shrubs were introduced:

eight by Conrad Loddiges; six by Messrs. Lee and Kennedy; three by Fraser; nineteen by Lyon; one by the Kew Garden; one by the London Horticultural Society; one by Don of the Cambridge Botanic Garden; and one by Sir Abraham Hume. Among the most interesting articles introduced during this decade are, *Ròsa multiflòra*, *Cunninghàmia lanceolàta*, *Juníperus excélsa*, *Caprifòlium japónicum*, *Ròsa Bânksiæ* *Rhododéndron catawbiénsè* (by Fraser), and *Cratægus Arònia*. It is somewhat remarkable, that of such a number of species introduced during this decade, the names of so few of the introducers should be known; but it must be recollected that the means of introducing were, at this period, principally by packets of seeds sent to the nurserymen by foreign correspondents, or by amateurs; and that, as several years must necessarily elapse between the period of introduction, and that of flowering and naming, the name of the collector who sent the seeds, or of the nurseryman who first raised plants from them, is forgotten, or ceases to be of the same interest. The case is different when living plants are brought into the country, and it is, in truth, chiefly of the introducers of such that the names are known.

From 1811 to 1820, three hundred and seventy-four trees and shrubs were introduced, viz., forty-four by Messrs. Loddiges; twelve by Lyon; four by Lee and Kennedy; three by Whitley and Co. among which was *Spiræa bélla* in 1820; three by the Horticultural Society (including *Cotoneáster affinis* in 1820); two by Fraser (*Abies Fraseri*, and *Yucca angustifòlia* in 1811); one by Don of the Cambridge Botanic Garden: one (the *Ribes sanguineum*, in 1817) by Archibald Menzies, Esq., who sailed round the world with Captain Vancouver; *Genísta procumbens* by Schleicher, a botanical collector in Switzerland; one by Knight of the Exotic Nursery, King's Road; and one (*Mahònia fasciculàris*) by A. B. Lambert, Esq. Among the most valuable of the species introduced by Loddiges are, *Azàlea arboréscens*, *A. speciosa*, and *Ribes aureum*, in 1812; *Symphòria racemòsa*, *Cytisus ruthénicus*, *Juníperus recurva*, and *Yucca tenuiflòra*, in 1817; *Alnus cordifòlia* (the most beautiful species of the genus), in 1818; *Armeniaca brigantiaca*, and *Quercus stellàta*, in 1819; *Cratægus melanocárpa*, *C. latifòlia*, *C. Olivieriàna*, *Fráxinus pannòsa*, *F. platycárpa*, *F. lancea*, *Pinus excélsa*, and *Abies Píchta*, in 1820. Among those introduced by Lyon are, *Magnòlia pyramidàta*, in 1811; and *Andrómeda floribúnda*, *Nýssa cándicans*, *Bòrya ligústrina*, *B. porulòsa*, *B. acuminàta*, *Virgílià lùtea*, and *Cratægus apiifòlia*, in 1812. Among the fine plants recorded as having been introduced in this decade, without mentioning the names of the introducers, are, *Æsculus glàbra* and *pállida*, and *Pàvia híbrida*, in 1812; *Bérberis sinénsis*, *Cydònia japónica*, and *Dáphne Thymelæa*, in 1815; *Plánera Richárdi*

in 1816; *Cratægus prunifolia*, in 1818; *Yucca glaucéscens*, *Ribes caucásicum*, and *Caragána microphýlla*, in 1819; *Pýrus nepalénsis*, *Philadélphus hirsútus*, *Pópulus macrophýlla*, *Tília laxiflora*, *Pinus adúnca*, and *P. uncinàta*, in 1820.

From 1821 to 1830, three hundred and eighteen trees and shrubs were introduced; viz., upwards of sixty by the Horticultural Society; twenty-five by Schleicher (obscure species of willows); twenty-one by Messrs. Loddiges; four by Lord Carnarvon; three by Whitley; one by Malcolm; one by Shepherd of Liverpool; one by Don of Cambridge; one by Low of Clapton; one by Philip Barker Webb, Esq.; one (*Benthàmia fragífera*, in 1825) by J. H. Tremayne, Esq.; one by the late Mr. William Baxter (*Sóllya heterophýlla*, in 1830); one by Bunney; and one (*Ribes speciòsum*, in 1829) by A. B. Lambert, Esq. By far the greater number of the species introduced by the Horticultural Society were sent home by the late unfortunate Douglas, from the north-west coast of North America; among them are, *Bérberis Aquifòlium*, *Gaulthèria Shálton*, *Arctostáphylos tomentòsa*, *Ribes viscosíssimum*, and *Acer macrophýllum*, in 1826; and *Bérberis glumàcea*, *Acer circinàtum*, *Arbutus procèra*, *Ribes níveum*, *inèbrians*, and *divaricàtum*; *Rùbus spectábilis*, *Abies Douglàsii*; *Pinus ponderòsa*, *Lambertiàna*, and *Sabiniàna*; *Amelànchier flórida*, and *Gárrya ellíptica*, in 1827. In this decade the Horticultural Society also introduced the *Cèdrus Deodàra* from Nepal, in 1822; and *Cotoneáster frígida*, and *C. Nummulària*, in 1824. Among those by Loddiges are, *Quércus Taitzin* and *Gledítschia cáspica*, in 1822; *Acer opulifòlium*, *Fráxinus álba*, *epíptera*, *fúscà*, *macrophýlla*, and *quadrangulàris*, all in 1823; and *Fráxinus cinèrea* and *Cotoneáster microphýlla*, in 1825. The greatest number of the valuable trees and shrubs added to the British arboretum, during this century, was introduced by Messrs. Conrad Loddiges and Sons, and the next greatest number by the Horticultural Society. Messrs. Loddiges received their importations chiefly from their foreign correspondents, and more especially from American collectors and nurserymen. The principal British collectors during this period were, Fraser, Lyon, and Douglas. Notices of the first two have been kindly prepared for us by Mr. Forsyth; and of the latter we shall give a short abstract of a biographical memoir which appeared in the *Gard. Mag.*, vol. x. p. 271.

John Fraser was a native of Inverness-shire; he came to London about 1770 (or 1776), married, and settled as a hosier and draper in Paradise Row, Chelsea; but, being of a very active and enterprising turn of mind, and having imbibed a taste for plants in his frequent visits to the physic garden at Chelsea, then under the care of the late Mr. Forsyth, he determined on

proceeding to North America in search of new, rare, and interesting plants.

Accordingly, in 1783 or 1784, he embarked for Charleston, South Carolina, where he made his first collection of many valuable new plants, which he consigned for sale to the care of Mr. Frank Thoburn, nurseryman, at Old Brompton. In the beginning of 1785 he returned to London, and expected to receive the reward of his labours, but was told that all his valuable plants had died, and that those remaining were common, and not very saleable. This created a misunderstanding which led to a lawsuit, which was long and very expensive to both parties.

In the autumn of 1785 he again visited South Carolina, where he made acquaintance with a most valuable friend, Thomas Walter, Esq., an eminent botanist, who had compiled a *Flora Caroliniana*, which MS. Mr. Fraser brought to London, and which was published by him in 1788, 8vo (the original herbarium of Mr. Walter is now in the possession of Mr. J. Fraser). In this second journey he was very successful in bringing home with him many new American plants, seeds, and dried specimens of plants, and various other objects of natural history. These were disposed of principally to the different plant collectors, nurserymen, and others, and he obtained liberal prices for them. Among the plants were several species of pines, oaks, magnolias, azaleas, rhododendrons, &c.; all most valuable and ornamental trees and shrubs, hitherto unknown in the gardens of England. The *Hortus Kewensis* records 16 new plants as having been introduced by Mr. Fraser in 1786, and five more in 1787. He likewise brought home with him, for cultivation, the seeds of a new species of grass, then named *Agróstitis cornucòpia* (now *Trichòdium decúmbens*), an account of which, with a coloured plate, he published in 1787 folio.

In 1790 and 1791 Mr. Fraser made his third and fourth voyages to America, where he extended his researches, and added further to his former collections. In 1791 he introduced the *Thàlia dealbàta*. About 1795 he established himself in a nursery, at Sloane Square, Chelsea, to which place all his subsequent consignments were made.

In April, 1796, he had completed his fifth voyage from America, bringing with him seeds and plants for sale as before. This year he visited Petersburg, taking with him a choice collection of plants, which were purchased, and paid for most liberally, by the Empress Catherine. Upon his return to England, he introduced that fine fruit, the black Tartarian cherry, and also the white Tartarian cherry.

In 1797 and 1798 he repeated his visits to Russia, having been honoured with the commands of the imperial family to

make further additions to their botanical collections; and, in the execution of these commands, he gave such satisfaction, that he was honoured, by special appointment, with the title of Botanical Collector to their Imperial Majesties the Emperor Paul and the Empress Marie, under the sign manual of each, dated Paulowskoe, August, 1798; and in furtherance of this commission he again, with his son John, in 1799, 1800, and 1801, visited the southern states of North America, the Isle of Cuba, the Bahamas, &c. In their passage to the Havannah, from the United States, they were shipwrecked, and saved themselves, with great difficulty, in the Cayos, a small island at the entrance of the Old Channel. In Cuba they had the good fortune to meet with the celebrated travellers Baron von Humboldt and Aimé Bonpland, and from these generous men of science they received every assistance and kind recommendations to the authorities at the Havannah. After an absence of more than two years, Mr. Fraser returned to England with many new and valuable discoveries. (In 1800, the *Hortus Kewensis* records the introduction of thirteen new plants by the Frasers, and in 1801 two more, *Andròmeda cassinifolia* and *Magnòlia cordata*.) He again went to Russia, but, in consequence of the sudden termination of the life of the Emperor Paul, he was unsuccessful, as his services were neither acknowledged nor required by the Emperor Alexander. He made two visits afterwards to the capital of Russia, and to Moscow, in a fruitless attempt to obtain a just remuneration for his arduous and perilous employment.

In the vicinity of Matanzas, in Cuba, they discovered a beautiful species of palm, with silvered leaves (*Còrypha miraguàma Humb. et Bon.*, Nov. Gen. 1. p. 290.), the leaves of which produce a most beautiful and durable material for the manufacture of ladies' hats and bonnets. These were woven by the hand, all in one piece, without sewing, in a new and peculiar manner: a patent was taken out for making them, and the manufacture was patronised by Her Majesty the late Queen Charlotte, and conducted under the management of his sister, Mrs. Christiana Fraser, through whose great perseverance in teaching many young persons the secret of the work, employment was afforded to a number of hands. Subsequently the manufactory proved unsuccessful, from want of capital, more than any other cause.

In 1806, 1807, 1808, 1809, Mr. Fraser made his last excursions to North America, in company with his eldest son. (The *Hortus Kewensis* records nine new plants introduced by them in 1809.) After this, he remained at his nursery in Sloane Square, carrying on the business there, in which, however, he was not successful. Here frequent disappointments, ill-treatment, and other circumstances, all tended to break down

one of the most enterprising, indefatigable, and persevering men that ever embarked in the cause of botany and natural science.

He died at Sloane Square, April 26. 1811, in his 60th year, leaving his wife, who died a few years afterwards, and two sons; John, the eldest, who had been his companion in all his latter voyages to America and Russia, and who is now a respectable nurseryman at Ramsgate, and James Thomas, also living.

Of John Lyon, another botanical collector, very little is known. He is said to have been a natural son of William Lyon, Esq., of Gillogie, Forfarshire, who was afterwards a merchant in London. When he went to America is uncertain; Pursh, who had the management of the gardens of William Hamilton, Esq., at Woodlands, near Philadelphia, informs us that, when he resigned, in 1802, Lyon succeeded him, and remained there till 1805.

During this period Lyon, we are told by the Messrs. Lodiges, sent home several plants and seeds; and the year after he left Mr. Hamilton's service (1806), he brought an extensive collection to England; the plants composing which were partly disposed of by private contract, but were chiefly sold by auction in a garden at Parsons' Green, Fulham. The catalogue of these plants fills 34 closely printed pages, it enumerates 550 lots, and the sale occupied four days. Several of the lots were composed of large quantities of one-year-old seedlings in pots; and ten lots at the end of the sale consisted each of 50 different sorts of seeds. This, it is believed, was by far the greatest collection of American trees and shrubs ever brought to England at one time, by one individual. It contained scarcely any herbaceous plants; and the trees and shrubs were chiefly such as had been already introduced. In the *Hortus Kewensis* fourteen new plants are mentioned as having been introduced by Lyon in 1806, which, doubtless, formed part of the importation of that year.

Mr. Lyon appears to have soon after gone out again, and explored the southern states of North America; viz., the Carolinas, Georgia, and Florida; and, in 1811 and 1812, he again brought over a large collection of plants in cases, which arrived in very fine condition, and were disposed of by public auction at Chelsea. Six plants are mentioned in the *Hortus Kewensis* as having been introduced by Lyon during these years.

Mr. Nuttall separated some of the species of *Andr meda*, and formed of them a new genus, which he named *Lyonia*. "To commemorate the name of the late Mr. John Lyon, an indefatigable collector of North American plants, who fell a victim to a dangerous epidemic amidst those savage and romantic mountains which had so often been the theatre of his labours." (*Gen. of N. American Plants*, Boston, 8vo, 1820, 1. p. 266.) The genus was, however, named before Mr. Lyon's death, as

appears by the catalogue, before referred to, of plants sold in 1806, in which several species of *Lyonia* are mentioned. Mr. Lyon, it is believed, died in 1818.

David Douglas was born at Scone, near Perth, and served his apprenticeship, as a gardener, in the gardens of the Earl of Mansfield. About the year 1817 he removed to Valleyfield, the seat of Sir Robert Preston, Bart., then celebrated for a choice collection of exotics, and shortly afterwards went to the Botanic Garden of Glasgow. Here his fondness for plants attracted the notice of Dr. Hooker, the professor of botany, whom he accompanied in his excursions through the Western Highlands, and assisted in collecting materials for the *Flora Scotica*, with which Dr. Hooker was then engaged. This gentleman recommended him to the late secretary of the Horticultural Society, Joseph Sabine, Esq., as a botanical collector; and in 1823 he was despatched to the United States, where he procured many fine plants, and greatly increased the Society's collection of fruit trees. He returned in the autumn of the same year; and in 1824 an opportunity having offered, through the Hudson's Bay Company, of sending him to explore the botanical riches of the country adjoining the Columbia river, and southwards towards California, he sailed in July for the purpose of prosecuting this mission.

While the vessel touched at Rio de Janeiro, he collected many rare orchideous plants and bulbs. Among the latter was a new species of *Gesneria*, which Mr. Sabine named, in honour of its discoverer, *G. Douglàsii*. He was enraptured with the rich vegetation of a tropical country; he stopped at Rio longer than he anticipated, and left it with regret. In the course of his voyage round Cape Horn he shot many curious birds peculiar to the southern hemisphere, and prepared them for sending home. On Christmas-day he reached the celebrated island of Juan Fernandez, which he describes as "an enchanting spot, very fertile, and delightfully wooded. I sowed a large collection of garden seeds, and expressed a wish they might prosper, and add to the comfort of some future Robinson Crusoe, should one appear." He arrived at Fort Vancouver, on the Columbia, on the 7th of April, 1825. Here an extensive field presented itself to him; and the excellent manner in which he performed his duty to the Horticultural Society cannot be better exemplified than by referring to the vast collections of seeds which from time to time he transmitted home, along with dried specimens, beautifully preserved, and now forming part of the herbarium in the garden of the Society at Chiswick. Of the genus *Pinus* he discovered several species, some of which attain to an enormous size. The *Pinus Lambertiana*, which he named in compliment to Aylmer Bourke Lambert, Esq., vice-president of the Linnæan

Society, is, perhaps, the largest of the whole. One of these, which had been blown down, measured 215 ft. in length, and 57 ft. 9 in. in circumference, at 3 ft. from the ground. The cones of it, which Mr. Douglas sent home, were 16 in. long, and 11 in. in circumference. The kernel of the seed is sweet and pleasant to the taste, and is eaten by the Indians, either roasted, or pounded into coarse cakes for winter store. The resin, which exudes from the trees when they are partly burned, loses its usual flavour, and acquires a sweet taste; in which state it is used by the natives as sugar. Another species, named by Mr. Sabine *Abies Douglassii*, attains nearly the size of the above.

In the spring of 1827 Mr. Douglas traversed the country from Fort Vancouver, across the Rocky Mountains, to Hudson's Bay, where he met Captain (now Sir) John Franklin, Dr. Richardson, and Captain Back, returning from their second overland arctic expedition. With these gentlemen he came to England in the autumn, bringing with him a variety of seeds, as well as specimens of plants and other objects of natural history. Through the kindness of his friend and patron Mr. Sabine, he was introduced to the notice of many of the leading literary and scientific characters in London; and shortly afterwards he was honoured by being elected, free of expense, a Fellow of the Linnæan, Geological, and Zoological Societies; to each of which he contributed several papers, since published in their *Transactions*, evincing much research and acuteness as a naturalist. Some entertaining extracts from his letters to Dr. Hooker were published in *Brewster's Edinburgh Journal* for January, 1827; and a genus of plants belonging to the natural order Primulaceæ was dedicated to him by Professor Lindley, and defined in *Brande's Journal* for January, 1828.

After being in London for two years, Mr. Douglas again sailed for Columbia in the autumn of 1829; where he remained some time, enjoying his favourite pursuit, and adding largely to his former discoveries. His return was expected by the very ship which brought the tidings of his horrible death; an event which was occasioned by his falling into a pit made by the natives of the Sandwich Islands for catching wild bulls, one of the latter being in at the time.

The plants introduced by Mr. Douglas are supposed to be more numerous than those introduced by any other individual whatever; and what greatly adds to their value is, that, being from a temperate region, they will all endure the open air in this country. The number of herbaceous species which he introduced amounts to nearly 100, and of trees and shrubs to 50. The names of the latter compose the following list, which has been kindly communicated to us by Mr. Munro, the head gardener of the London Horticultural Society.

In 1826 and 1827.

Abies Douglàsii.
Acer circinatum.
 macrophýllum.
Amelánchier flórida.
 parvifólia.
Arbutus procéra.
Arctostáphylos tomentósa.
Bérberis Aquifólium.
 glumácea.
Caprifólium ciliósum.
 Douglàsii.
 hispídulum.
Càrya nigro-cathártica.
 **Ceanóthus collinus.*
 *ellípticus.
Gárrya ellíptica.
Gaulthèria Shállon.
 **Laúrus occidentális.*
Pínus Lambertiana.
 ponderósa.
Púrshia tridentàta.
Ribes viscosíssimum.
 aúreum.
 cèreum.
 divaricàtum.
 echinàtum.
 irriguum.
 lacústre.

Ribes níveum.
 petiolàre.
 sanguíneum.
Rùbus nutkànus.
 spectábilis.
Sálvia carnósa [Audibértia in-
 càna.]
Spiræa ariæfólia.
Vaccínium ovàtum.

In 1831.

Abies amábilis.
 grándis.
 Menzièsii.
 nóbilis.
Clématis Douglàsii.
Pínus montícòla.
 mont. var. with red cones.
 Sabiniana.
Pýrus rivulàris.
Ribes glutinósum.
 malváceum.
 speciósum.

In 1832.

Lupínus álbifrons.
Pínus Sabiniana var.

In 1833.

Pínus insígnis.

Of the above specimens, which were all introduced by seeds, the three marked with a * did not vegetate. Some species of *Ròsa* and *Cratægus*, not included in the above list, have vegetated, but are not enumerated, as they have not yet flowered; and consequently have not yet been named or identified.

To enable our readers to take a general view of the various details respecting introductions given in the preceding pages, we shall next endeavour to generalise them; first, numerically; and, secondly, geographically. For the first object, we have had from our *Hortus Britannicus* an enumeration made of the number of species introduced in each decade, from the beginning of the 16th century to the end of the year 1830. We do not give this enumeration as perfectly accurate; because many of the species in our catalogue, as in every other, are doubtful; but it is not of much consequence whether it be perfectly accurate or not; it is sufficiently so to show the ratio of the increase of the introductions, from the earliest periods of which we have any record of them, up to the present time.

There were introduced from the year	to the year	Species.	There were introduced from the year	to the year	Species.
1548	1550	17	1691	1700	24
1551	1560	1	1700	1710	12
1561	1570	18	1711	1720	12
1571	1580	3	1721	1730	44
1581	1590	2	1731	1740	69
1591	1600	48	1741	1750	21
1601	1610	1	1751	1760	77
1611	1620	1	1761	1770	58
1621	1630	22	1771	1780	58
1631	1640	27	1781	1790	49
1641	1650	4	1791	1800	45
1651	1660	17	1801	1810	93
1661	1670	7	1811	1820	361
1671	1680	1	1821	1830	242
1681	1690	27			

The numbers, taken by centuries, are, in the 16th century, 89; in the 17th, 131; in the 18th, 445; and, in the first three decades of the 19th, 699! The total number of foreign trees and shrubs introduced up to the year 1830, appears to be about 1300; or, probably, up to the present moment, including all those species which have not yet flowered, and, consequently, have not yet been recorded in books, about 1400.

The countries from which these 1300 species have been introduced appear, from the *Hortus Britannicus*, to be as under:—

Europe: Greece, Turkey in Europe, and the Levant, 36; Italy, 35; Sicily and other Mediterranean islands, 19; Spain, 69; Portugal, 12; Switzerland, 49; France, 34; Germany, 52; Hungary, 46; Russia, 41; Sweden, 4; Lapland, 4; Spitzbergen, 1; North of Europe, 2; Central Europe, 18; South of Europe, 111: in all, 543. *Asia*: Siberia, 69; Asia Minor, 3; East Indies, 4; Nepal, 54; China, 34; Japan, 11; Persia, 5; Asia, 3: in all, 183. *Africa and the Canary Isles*: Barbary States, 13; Egypt, 3; Cape of Good Hope, 4; Canary Isles, 3: in all, 23. *America*: North America, 528; Mexico, 4; South America, 22; Straits of Magellan, 6: in all, 560. *Australia and Polynesia*: New Holland, 1; Van Diemen's Land, 2; New Zealand, 1: in all, 4.

It would thus appear, that nearly half the foreign trees and shrubs in the country have been introduced during the present century; and that these have been brought chiefly from North America. Among them there are not more than 300 trees which attain a timber-like size, and of these by far the most valuable is the larch. Some of the European acers, the sweet chestnut, some oaks, some poplars, pines, and firs, and the platanus and cedar from Asia, are also valuable as timber trees; but the chief accessions to this class are the acers, oaks, elms, ashes, poplars, birches, pines, and firs of North America. Our principal fruit trees are from Asia, including the common walnut, which is both a fruit and a timber tree; but by far the finest

ornamental trees and shrubs are from North America. Our greatest hopes for future introductions are from the unpenetrated regions of North America, and the mountainous regions of Asia and New Zealand.

We shall conclude this chapter by enumerating some of the principal planters of arboretums, and places where arboretums were planted, during the present century; premising that we do not include in this list any of those places which were commenced during the last century.

Among the planters of arboretums in Great Britain during the nineteenth century, the first place belongs to George, fourth duke of Marlborough. This nobleman, when Marquess of Blandford, resided on the estate of White Knights, near Reading, from the year 1800 till he succeeded his father in 1817. About 1801 he began to collect plants of every description, built numerous hot-houses for the exotics, and occupied a large walled garden with the hardy herbaceous plants, and the more choice trees and shrubs. Soon after, finding this garden too limited, he employed, as an arboretum, a space of several acres, called the Wood; and throughout the park at White Knights he distributed many trees, and a collection, as extensive as could be then procured, of the genus *Cratægus*. About this time magnolias, rhododendrons, azaleas, and other American trees and shrubs, being rare, or newly introduced, bore enormously high prices; but price was never taken into consideration by the Marquess of Blandford. He was never content with only one plant of a rare species, if two or more could be got; and the late Mr. Lee of the Hammersmith Nursery informed us, that he had sold several plants of the same species to the marquess when they were at twenty guineas, and even thirty guineas each. In consequence of a similar mode of proceeding in his transactions generally, the Marquess of Blandford soon found himself involved in debt and lawsuits, which, since 1816, have greatly crippled his exertions. He has still, however, the same taste for plants, and indulges it, as far as his limited resources will permit, in the pleasure-grounds of the palace at Blenheim, where His Grace at present resides. White Knights is now chiefly remarkable for its magnolia wall, which is 145 ft. long and 24 ft. high, entirely covered with twenty-two plants of *Magnòlia grandiflòra*, which flower every year from June till November. They were planted in the year 1800, when the price in the nurseries, for good plants, was five guineas each. In the Wood there are a great number of remarkably fine specimens of all the species of *Magnòlia*, and especially of *M. auriculàta* and *acuminàta*. There are also very fine trees of *Acer rùbrum*, *saccharinum*, and *striatum*; of *Æsculus* and *Pàvia*, of *Arbutus*, of *Kölreutèria*, of *Virgìlia*, of *Còrnus flòrida*, of

Gleditschia, Cércis, Crataegus, and Photinia 15 ft. high; some of the finest trees of Pÿrus nivàlis and bollwylleriana in the country; of Halèsia, Diospÿros, Nÿssa, Gymnócladus, Plánera, Æglans, Stuártia, Láurus, Quércus, Juníperus, Thuja, a remarkably fine CunníngHàmia, and many pines, among which are the greatest number of Pínus Pallasiána to be found together in any grounds in England. Pínus Cémbra has here attained the height of 30 ft. in 35 years; and Làrix péndula that of 50 ft. in the same time. At Blenheim the duke has introduced the finest trees he could procure, in numbers and in masses, as far as he was enabled to do so; and Magnòlia conspícua, of which seldom more than one or two plants are to be found in any one demesne, may be there reckoned by dozens. An account of White Knights, as far as its picturesque beauty extends, will be found in Hoffland's *Description of White Knights*, Lond. 1819, fol.; and of its gardens, in a botanical and horticultural point of view, in the *Gardener's Magazine*, vol. ix. p. 664.; in which work will also be found an account of Blenheim, vol. x. p. 99. The Duke of Marlborough's gardener, from the commencement of the duke's gardening operations at White Knights to the present time, has been Mr. Jones. White Knights, which is now the property of Francis Cholmeley, Esq., has its gardens under the direction of Mr. Ward. The house is at present (1835) unoccupied.

William Beckford, Esq., of Fonthill Abbey, began to plant at that place all the rare trees and shrubs which he could procure, about the same time as the Duke of Marlborough planted White Knights. He paid no attention to house or to herbaceous plants, but, like the duke, he planted the choicest trees and shrubs, in quantities, without any regard to their cost; paying for them, we believe, in ready money. We had the satisfaction of inspecting the grounds at Fonthill twice in 1806, when they were in their highest beauty and keeping; and we spent two days in looking at them again in 1833, when they were in a state of neglect, and when the greater number of the rare trees and shrubs, and in particular the pinetum, thornery, and rosary, were almost obliterated by the growth of common trees and shrubs. There are still some fine magnolias, rhododendrons, and azaleas in the American ground, which have been hardy enough to cope with the native trees which have been planted, or have sprung up fortuitously around them. The scenery of Fonthill has somewhat of a Swiss character, from the hilly ridge on which the Abbey is built, and the prevalence of the pine and fir tribe in the woods; and in it there is an air of melancholy grandeur, unlike that of any other place that we are acquainted with in Britain. A description of Fonthill Abbey, when in its most perfect state, has been given by Britton, in his *Wiltshire*, and a notice of it, as it appeared to us in 1833, will be found in the *Gard. Mag.*, vol. xi. p. 425.

After White Knights and Fonthill Abbey, the following places may be mentioned as subsequently planted, and as containing collections of trees and shrubs more or less extensive. In Bedfordshire, Flitwick House, where an arboretum was planted in 1829. (See *Gard. Mag.*, vol. v. p. 559.) At Woburn Abbey, where a salicetum, or salicetum (as the Duke of Bedford more classically terms it, in his *Salicetum Woburnense*), was planted in 1825, and where an arboretum is now, 1835, commenced. In Berkshire, High Clere, where a number of American trees and shrubs were planted, and a great quantity of fine hybrid rhododendrons and azaleas raised, between 1820 and 1830; and Dropmore, where there is the most complete pinetum in England, the species and varieties amounting, in 1835, to 120. In Cornwall, at Carclew, there is a good collection. In Derbyshire, at Chatsworth, a very complete arboretum was begun in 1834; of which an account and ground plan will be found in the *Gardener's Magazine*, vol. xi. p. 385. In Devonshire, Luscombe near Dawlish (said by Davis, in his *Landscape-Gardener*, to be "an unrivalled production of fine taste"), Endsleigh Cottage, Mamhead, and Bicton, contain good collections. In Essex, Hylands. In Hampshire, Bishop Stoke Vicarage. In Hertfordshire, Cheshunt, which contains a pinetum. In Kent, at Cobham Hall, a very good collection. In Lancashire, Latham House. In Northumberland, Belsay Castle, where there is a pinetum. In Staffordshire, Alton Towers, and Somerford Hall, where there is an excellent arboretum, with sufficient space allowed for the trees to attain their full size. In Suffolk, Barton Hall. In Surrey, Bagshot Park, Milford, where there is a regular arboretum, Oakham Park, Mere Cottage, and Deepdene. In Sussex, West Dean, and Arundel Castle. In Wiltshire, Wardour Castle, where there is a good pinetum, and Boynton.

In Scotland may be mentioned, Hafton, in Argyleshire; Gordon Castle, in Banffshire; Drumlanrig, and Jardine Hall, in Dumfriesshire; Dalhousie Castle, in Edinburghshire; St. Mary's Isle in Kirkcudbrightshire; and Dunrobin, in Sutherlandshire.

In Ireland, Terenure, near Dublin, where there is the most complete arboretum in the country; and Charleville Forest, in Meath, where an arboretum was begun in 1811.

Several public bodies have commenced arboretums during this century. In England the first of these is that of the London Horticultural Society, whose collection, in their garden at Turnham Green, commenced in 1823, may be considered the first in England. It is to be regretted that the space in the garden devoted to this arboretum was originally much too small; and also, that the trees and shrubs were chiefly crowded together in clumps, which have subsequently never been sufficiently thinned out. In consequence of this, the different kinds have

not had an equal chance of displaying themselves, or of attaining that magnitude and character which they ought to have to answer the ends of an arboretum. (See our ideas more at length, and illustrated by figures, in the *Gard. Mag.*, vol. v. p. 346. and fig. 79., and vol. vi. p. 250. and fig. 44.) There is an arboretum in the Liverpool Botanic Garden, in that of Hull, in that of Colchester, in that of Manchester, in that of Birmingham, and one is just commenced in that of Sheffield. The Caledonian Horticultural Society have an arboretum in their experimental garden at Inverleith; that of the Botanic Garden of Edinburgh has been much increased; and there is a good one in the Glasgow Botanic Garden. In Ireland, the Trinity College Botanic Garden was laid out by Mr. Mackay in 1808, and at first contained only three acres. In 1833 two acres more were added, which are principally occupied by ornamental trees on a grass lawn, with surrounding borders for showy herbaceous plants, and trees and shrubs which require the protection of a wall. An arboretum was commenced in the garden of the Cork Institution, soon after the foundation of the Trinity College Botanic Garden; but that institution has been since broken up, and the plants and trees of the garden sold and dispersed. The Belfast Botanical and Horticultural Society established a garden and an arboretum about 1830. The Glasnevin Garden belongs to the preceding century.

The British nurserymen have not been wanting in forming arboretums. Preeminently among them stand the Messrs. Loddiges of Hackney, who have been assiduously collecting trees and shrubs from all parts of the world, since the middle of the last century; and in the year 1818, when these amounted to above 1200 species and varieties, exclusive of azaleas, roses, and willows they were arranged alphabetically on the right hand side of a walk forming a scroll like the Ionic volute, extending over a space of upwards of seven acres, commencing with the letter A, at the outer circumference, and terminating with Z (*Zizyphus*) near the centre. The centre itself forming the eye of the arboretum consists of ten concentric zones, devoted to peat earth plants, commencing in the outer zone with *Andrómēda*, and terminating in the inner one with *Vaccínium*. The collection of willows, which is very extensive, is placed by itself in the circumference; as is also the collection of yuccas. The collection of roses, which exceeds 1500 sorts, is planted on the left hand side of the scroll walk, and their number being about the same, as that of the trees and shrubs which do not require peat earth, they extend to the commencement of the concentric zones. The surrounding boundary walls are covered with half-hardy trees and shrubs. In 1830 this arboretum was in a high degree of perfection, and in the autumn of that year we had sketches taken

of all the trees and shrubs, to the scale of a quarter of an inch to a foot. As the object of Messrs. Loddiges was to include in one place, not only specimens of each particular tree and shrub, but also stools for propagating them, and a stock of young plants for sale, all placed beside the specimen plant, it became necessary to cut down the specimens as soon as they had attained a certain size; and this was accordingly done with many of the timber trees in 1832 and 1833. To us this has been an incalculable loss, because it has prevented us from examining many of the trees in flower; but the stools or young plants still remain, and the collection is increasing every year. So spirited an undertaking cannot be sufficiently appreciated; and it is only to be regretted that the want of a separate piece of ground for containing the stools and the plants for sale, rendered necessary the felling of a collection of specimens such as could be found assembled together nowhere else in the world. Some account of this arboretum, accompanied by a plan, will be found in the *Encyclopædia of Gardening*, edit. of 1835, p. 1217. Such a magnificent example could not be expected to be generally followed by commercial men, but it has rendered more frequent the practice among nurserymen of planting out specimens of choice trees and shrubs; and some have subsequently even formed regular arboretums. Among these the first in the order of time, as well as in completeness, is that of Mr. Donald of the Goldworth Nursery, near Woking, in Surrey, which was commenced in 1831, and of which an account, with a plan, will be found in the *Gard. Mag.*, vol. vii. p. 360. This arboretum, which we viewed in May, 1834, already contains a number of very fine specimens. Mr. Buchanan, jun., of Camberwell, has spared no pains in collecting trees and shrubs; and had got together, in 1834, though in a limited space, a collection which may rank next to that of Mr. Donald. Mr. Miller of the Bristol Nursery has also planted an arboretum; and a nursery has been formed by Messrs. Young and Penny, at Milford, connected with the arboretum of Philip Barker Webb, Esq., which Messrs. Young and Penny are greatly increasing every year, by the addition of new species. This arboretum contains a great number of oaks, acers, and pines. Mr. Rogers of Southampton began an arboretum in 1833; and Mr. Page of the same place is also forming one. More or less has been done in this way, in the nurseries of Messrs. Lee of Hammersmith, Messrs. Osborne of Fulham; Mr. Knight of the Exotic Nursery, King's Road, Chelsea; Messrs. Young of Epsom, Mr. Newman of Chichester, Mr. Cunningham and Mr. Skirving of Liverpool, Messrs. Dickson of Chester, Messrs. Pope of Birmingham, Messrs. Backhouse of York, and a number of others. In Scotland, Mr. Lawson of Edinburgh is most assiduous in collecting trees and shrubs, both at home and

abroad; and he has commenced an arboretum, which already contains a collection of pines and firs not surpassed by any in Britain. An account of this arboretum, which will soon be the first in Scotland, will be found in the *Gard. Mag.*, vol. xi. Messrs. Dickson of Edinburgh, Brown at Perth, and Messrs. Austin of Glasgow, have also a great many choice trees planted out, as have various other nurserymen in that country. In Ireland we have already mentioned the nurseries most celebrated for their fine specimens and extensive collections.

CHAP. III.

OF THE HISTORY AND GEOGRAPHY OF THE TREES AND SHRUBS OF THE CONTINENT OF EUROPE.

THE Continent of Europe has supplied, as we have seen in the preceding chapter, a considerable number of trees and shrubs to the British Arboretum. The different countries which compose it have been so thoroughly explored by botanists, that few farther additions can be expected from them; but it will be, nevertheless, interesting to examine the indigenous ligneous flora of each as compared with that of Britain, and its capacity for receiving additions from the trees and shrubs of other parts of the world. We shall take these countries in the order of France, Holland and the Netherlands, Germany, Scandinavia, Russia and Poland, Switzerland, and Greece, Italy, Spain, and Portugal; and, considering the subject as one of secondary interest to that of the preceding chapter, our observations on it will be brief.

SECT. I. *Of the Indigenous and Foreign Trees and Shrubs of France.*

FRANCE, from its extent, the warmer climate of its southern provinces, and the varied character of its surface, including as it does some of the highest mountains in Europe, and a portion of the shores both of the Atlantic Ocean and the Mediterranean Sea, contains the richest indigenous ligneous flora of any country in Europe. There are few if any trees and shrubs which are indigenous to Britain that are not also indigenous to France; and there are in addition, in the latter country, all the species contained in the following enumeration, taken from Duby and De Candolle's *Botanicon Gallicum*, published in 1828. In this enumeration those orders, genera, or species, marked with a star (*), are either only in cultivation, or known or supposed to be not truly indigenous.

Ranunculácea. *Clématis Flámmula*, F. var. *marítima*, *cirr hòsa* var. *pedicellàta*, *balearica*; *Atragène austriaca*.

Crucifera. *Matthiòla tristis*; *Ibèris Garrexiàna*, *saxátilis*, *semperflorens*.

* *Capparidæa.* *Cápparis spinòsa*.

Cistinæa. *Cistus incanus*, *crispus*, *álbidus*, *salvirefólius*, *corháriensis*, *mons-peliensis*, *Lèdon*, *hirsútus*, *longifólius*, *populifólius*, *laurifólius*, *ladaníferus*; *Helianthemum umbellátum*, *alyssóides*, *alyssóides* var. *rugòsum*, *halimifólius*,

Fumàna, procúbhens, læ'vipes, glutinòsum, g. var. thymifòlium, g. var. juniperinum, organifòlium, celandicum, alpèstre, penicillatum, italicum, canum, lavandulæfòlium, stæchadifòlium, acuminatum, grandiflorum, obscurum, o. var. nummularium, hirtum, pilosum, apenninum, a. var. hispidum, pulverulentum, roseum, *majoranæfòlium*.

Polygaleæ. *Polygala saxatilis*, Chamæbuxus.

Malvæcæ. *Lavatera Olbia*, maritima; **Hibiscus syriacus*.

**Aurantiacæ.* *Citrus Médica*, Limonium, Aurantium, vulgaris: all cultivated.

Acerinææ. *Acer opulifolium*, monspessulanum, platanoides.

**Hippocastanææ.* *Æsculus Hippocastanum*.

**Meliæcæ.* *Melia Azedarach*.

**Vites.* *Vitis vinifera*, laciniosa.

Rutæcæ. *Ruta montana*, graveolens, bracteosa, angustifolia, corsica.

Coriariææ. *Coriaria myrtifolia*.

Celastrinææ. *Euonymus latifolius*.

Rhamnææ. *Zizyphus vulgaris*; *Paliurus aculeatus*; *Rhamnus Alaternus*, Clusii, infectorius, saxatilis, pubescens, pumilus, p. var. rupëstris, alpinus, a. var. corsicus.

Anacardiæcæ. **Pistacia vera*, *Terebinthus*, *Lentiscus*, *L.* var. angustifolia; *Rhus Côtinus*, *Coriaria*, radicans; *Cneorum tricoccum*.

Leguminosææ. *Anagyris foetida*; *Ulex provincialis*; *Spartium junceum*; *Genista candicans*, *linifolia*, *horrida*, *Lobelii*, *Salzmanni*, corsica, c. var. pubescens, *Scorpius*, hispanica, germanica, purgans, cinerea, humifusa, sagittalis, s. var. minor, prostrata, pilosa; *Cytisus Laburnum*, alpinus, sessilifolius, triflorus, spinosus, laniger, supinus, capitatus, argenteus; *Adenocarpus parvifolius*, telonensis; *Ononis arachnoides*, *Natrix*, rotundifolia, fruticosa, arragonensis; *Anthyllis cytisoides*, *Hermannia*, erinacea, *Barba-Jovis*; *Medicago suffruticosa*, s. var. *Benthämii*; *Dorycnium rectum*, hirsutum, h. var. incanum, suffruticosum; *Lotus creticus*; *Psoralea bituminosa*; **Robinia Pseud-Acacia*; *Colutea arborescens*; *Astragalus massiliensis*, aristatus; *Coronilla E'merus*, *juncea*, valentina, glauca; **Ceratonia Siliqua*; *Cercis Siliquastrum*.

Amygdalææ. **Amygdalus communis amara*, **communis dulcis*; **Persica vulgaris*, **laevis*; **Armeniaca vulgaris*, **brigantiaca*; *Prunus spinosa* var. *microcarpa*, domestica var. *pyramidalis*; *Cerasus duracina*, *Juliàna*, *caproniana*, *semperflorens*, *Mahaleb*, **Laurocerasus*.

Rosæcæ. *Spiræa hypericifolia*, h. var. *Plukenettiàna*, h. var. *crenata*; *Rubus tomentosus*, collinus, glandulosus; *Rosa sempervirens*, s. var. *microphylla*, **moschata*, *stylosa*, s. var. *leucochroa*, **indica*, *turbinata*, *gállica pumila*, *gállica officinalis*, *gállica parvifolia*, *lutea*, *lutea* var. *pumicea*, *sulphurea*, *pimpinellifolia*, p. var. *myriacantha*, p. var. *inermis*, *rubrifolia*, r. var. *pinnatifida*, *glandulosa*, *alpina*, a. var. *pyrenæica*, *centifolia*, c. var. *muscosa*, c. var. *pompônia*, *damascena*, *alba*.

Pomæcæ. *Crataegus Pyracantha*, *Azarolus*; *Cotoneaster tomentosa*; *Amelanchier vulgaris*; *Pyrus bollylleriàna*, *salvæfolia*, *amygdaliformis*, *acerba*, *intermedia*, *Chamæspilus*; *Cydonia vulgaris*.

**Granatææ.* *Punica Granatum*.

Tamariscinææ. *Tamarix africana*, germanica.

Philadelphææ. *Philadelphus coronarius*.

Myrtæcæ. *Myrtus communis*.

**Cactææ.* *Opuntia vulgaris*.

Umbelliferaæ. *Bupleurum frutescens*, spinosum, fruticosum.

Caprifoliæcæ. *Sambucus racemosa*; *Viburnum Tinus*; *Caprifolium baleàricum* a var. of *implèxum*, *etruscum*; *Lonicera nigra*, *pyrenæica*, *alpigena*, *cærulea*.

Cornææ. *Cornus mas*.

Lorántheæ. *Viscum Oxycedri*.

Compositææ. *Conyza saxatilis*, *sordida*; *Helichrysum Stæchas*, *angustifolium*; *Bupthalmum maritimum*; *Artemisia arborescens*, *corymbosa*, *arra-*

gonénsis, *Abrótanum*, paniculàta; *Santolìna rosmarinifolia*, víridis, incàna; *Balsamita ageratifolia*; *Stæhelina dùbia*, arboréscens.

Ericàcææ. *Erica scopària*, arbòrea, ramulòsa, multiflòra, mediterrànea; *Rhododéndron ferrugíneum*, hirsútum; *Lèdum palùstre*.

Styràcææ. *Stýrax officinàle*.

**Èbenàcææ*. *Diospýros Lótus*.

Oleàcææ. **O'lea europæ'a*; *Phillyrea angustifolia*, latifolia; *Sýringa vulgàris*, pérsica; ? *argéntea*; *Órnus europæ'a*.

Jasminææ. *Jasminum* * officinàle, fruticans, hùmile.

Apocýneææ. **Nèrium Oleànder*.

Asclepiàdææ. *Gomphocárpus fruticòsus*.

Convolvulàcææ. *Convólulus saxátilis*, s. var. *argénteus*.

Boragínææ. *Lithospérnum fruticòsum*, *oleàfolium*.

Solànææ. **Lýcium bárbarum*, europæ'um; *Solànum* * *Pseùdo-Cápsicum*.

Labiàta. *Rosmarínus officinàlis*; *Sálvia officinàlis*; *Teucrium fruticans*, flàvum, Pòlium, capitátum, flàvicans, *Pseùdo-Hyssòpus*; *Hyssòpus officinàlis*, o. var. *canéscens*; *Phlòmis fruticòsa*, *Lychnitis*; *Lavàndula Stæ'chas brachystàchya*, *S. macrostàchya*, vèra, *Spica*; *Saturýja capitàta*, montàna; *Thýmus vulgàris*, *Zýgis*, créticus, glandulòsus; ? *Oríganum majoranòides*, *Pràsium majus*.

Verbenàcææ. *Vítex Agnus-cástus*.

Globularínææ. *Globulària Alýpum*.

Plumbagínææ. *Státice monopétala*, minùta, pubéscens, fasciculàta.

Plantagínææ. *Plantàgo Cýnops*.

Chenopòdææ. *Camphoròsma monspeliàca*; *Salicòrnia macrostàchya*;

Salsòla prostràta; *A'triplex Hálimus*.

**Laurínææ*. *Laúrus nóbilis*.

Thymelæ'ææ. *Passerina diofca*, nivàlis, *Thomàsü*, hirsùta, h. var. *polygalæfolia*; *Dàphne Gnídium*, *Cneòrum*, oleòides, *Thymelæ'a*, *Tárton-raïra*, alpina.

Santalàcææ. *Osýris álba*.

Elæágneææ. *Elæágnus angustifolia*.

Euphorbiàcææ. *Euphòrbia spinòsa*, dendròides; *Mercuriàlis tomentòsa*.

**Úrticææ*, § *Artocárpeææ*. *Mòrus álba*, nigra; *Ficus Càrica*.

Ulmàcææ. *Ulmus effusa*; *Céltis austràlis*.

**Juglándææ*. *Jùglans règia*.

Betulínææ. *Bétula pubéscens*; *Álnus snavèolens*, víridis, incàna, cordàta, ellíptica.

Salicínææ. *Sàlix cineráscens*, versifolia, *daphnòides*, físsa, monàndra, incàna,

**babylònica*, *cæ'sia*, *pyrenàica*, glàuca, retùsa, *serpyllifolia*, hastàta; *Pópulus virginiàna*, dilatàta.

Cupulifèrææ. *Quérens Cérris*, *Tòza*, pubéscens, apennina, racemòsa, fastigiàta, *Fl'ex*, *Sùber*, coccifera.

**Platàncææ*. *Plátanus orientàlis*, occidentàlis.

Conifèrææ. *Pínus uncinàta*, *Mùgho*, marítima, pumílio, *Pínea*, halepénsis, *Larício*, *Cembra*; *Ábies excélsa*, pectinàta; *Làrix europæ'a*; *Juníperus phænicea*, *Sabina*, *Oxýcedrus*; *E'phedra distàchya*; * *Cuprèssus sempervirens*, * *sempervirens horizontàlis*.

Smilàcææ. *Smilax áspera*, mauritànica; *Rúscus hypoglòssum*.

Asphodèlæææ. *Ásparagus álbus*.

**Pálnæææ*. *Chamæ'rops hùmilis*.

Excluding from the above enumeration the cultivated and doubtful species, there appear to be 346 trees and shrubs indigenous to France, which are not indigenous to Britain; and this number, added to that of the woody species considered as decidedly indigenous to Britain, and supposed to be also indigenous to France, would give a total indigenous ligneous flora to the latter country of 546 species. The number of indigenous timber trees which exceed the height of 30 ft. in Britain appears to be 29; those indigenous in France which exceed that height are, according to the introduction to Michaux's

Arbres de l'Amérique, 30; but, according to the *Botanicon Gallicum*, they are 34. If we add to the indigenous woody plants of France those which are cultivated or doubtful, the total ligneous flora of that country will be above 580. If to this number we add the 528 trees and shrubs of North America (see p. 126.), all of which will grow in France, it will give a total ligneous flora to that country of above 1100 species; which, considering that France possesses in her botanic gardens or nurseries all, or nearly all, the trees cultivated in the open air in Britain, is probably as near the truth as the present state of our catalogues will admit of our arriving at. In the above enumeration of the woody plants of France, we have, as in the case of the enumeration of the woody plants of the British Islands (p. 27.), included all the under-shrubs, and also all those reputed species which we believe to be mere varieties. We have included the under-shrubs, because it is difficult to draw a line of separation between those which might practically be considered as herbaceous plants, though botanically they are suffruticose; and because, in a state of culture, some of these suffruticose plants attain such ample dimensions, and such a ligneous texture, as to assume quite a shrubby character; for example, *Euphórbia Charàcias* in Britain (p. 29.), and *Iberis saxátilis* in France (p. 132.). The first is seldom above 2 ft. high, in its native habitat in woods; and the second is seldom above 6 in. high, on rocks and in gravelly soil: but in dry deep garden ground the euphorbia will, in the course of a few years, form a bush between 3 ft. and 4 ft. high; and the iberis a mass above half that height. We have inserted the names of what we consider only varieties, because we have no doubt that, in most cases, they are plants tolerably distinct; because it is impossible to be quite certain of what are species and what varieties, without comparing them in different stages of their growth, and grown in the same soil, situation, and climate; and because we do not wish to set up our own opinion in this matter as absolute.

In an article by Professor Thouin, published in the *Mémoires d'Agriculture* for the year 1786, it is stated that France then possessed about 84 different species of trees, of which 24 were of the first rank in point of size, or exceeding 100 ft. in height; 16 of the second rank, or exceeding 60 ft. in height; and the remainder of the third rank, or exceeding 30 ft. in height. The names of these trees, and their arrangement according to the heights they attain, will be found in the work last quoted, and also in the *Nouveau Cours Complet d'Agriculture*, edit. 1821, art. *Arbre*. Deleuze states that France contains about 250 species of trees, of which more than three fourths are of foreign origin. (*Annales du Muséum*, tom. iii. p. 191.)

Ample as is the ligneous flora of France, it might be doubled by adding to it the trees and shrubs of Australia, of the mountainous regions of Asia, and of Mexico, Chili, and Peru. We do not speak of the whole of the trees and shrubs of these countries, because the whole are not yet known, but only of those that have been already introduced into Britain, and are treated by us as green-house plants; all of which would succeed in the open air of the southern provinces of France. Were the total number of ligneous species from these countries introduced, the number of trees and shrubs now in France would, in all probability, be quadrupled.

But though the ligneous flora of France is so much more extensive than that of Britain, yet it is far from being so equally spread over the country. Paris is considerably to the south of London, and yet there are above fifty species of evergreen trees and shrubs which are to be found in the open air in the environs of the latter city, which are not to be found in those of the former. We assert this from a comparison between a list of the trees and shrubs now (1835) growing in the *Jardin des Plantes* at Paris, furnished to us by Professor Mirbel, and the list which we have seen in MS. of the trees and shrubs now in the garden of the Horticultural Society of London. No part of France is so far north as Edinburgh; yet, while the cedar of Lebanon attains a large size far to the north of that city, and even in the Highlands of Scotland, it is killed during severe winters at Strasburg and throughout

Lorraine. Nevertheless, the fig and the vine ripen their fruit, and many deciduous foreign trees flower far better in the open air in the neighbourhood of Paris than they do in that of London. There are probably few plants that will endure the open air in the south of France, that might not be kept alive in the open air all the year in the southern extremity of Ireland, or in the neighbourhood of Penzance in Cornwall, though they would, probably, never flower at either of these places. The cause is so well known as to be hardly worth repeating: the summers in France have, proportionately to the latitude, more light and heat than those of Britain, and the winters less heat.

The first foreign trees introduced into France were, in all probability, those fruit-bearing species carried thither by the Romans; among which may doubtless be included the grape, the olive, and the fig, unless these and other fruit trees existed there at a still earlier period. In the progress of civilisation, many ages elapse before barren trees are planted either for timber or ornament. Charlemagne is praised by historians for eradicating the forests, and planting in their stead orchards and vineyards. He left a catalogue of certain plants, among which are some ligneous species, which he desired might be planted in all his gardens; but these, with the exception of the rose, were entirely for medicinal purposes. The earliest positive information that we have been able to obtain, respecting the introduction of foreign trees into France, is from the catalogue of Robin, gardener to Henry IV., which was published in 1610. It contains some few ligneous plants, such as the orange, pomegranate, the usual fruit trees, and a few of the ornamental trees and shrubs which are indigenous to Spain and Italy. Henry IV. was succeeded, in 1610, by Louis XIII.; and the botanic garden of Paris was begun by the latter king, about the year 1626, though the letters patent establishing it were not executed till 1635. Of this garden a catalogue was published by Guy de la Brosse, the first intendant, and who was also physician to the king, in 1636. In the letters patent, Vespasian Robin (son to the Robin who was gardener to Henry IV.) is mentioned as arborist to Louis XIII.; and the first *Robinia Pseud-Acacia* that was brought to Europe from North America was planted by him, in the *Jardin des Plantes*, in 1635. It is still in existence, and is now (1835) 78 ft. high. About 1815 it began to show symptoms of decay, but, the branches being lopped, the trunk has shot out with redoubled vigour. The edict of Louis XIII. also directed that pharmacy and chemistry should be taught in the garden, and illustrated by the demonstration of plants. It is said that the faculty in Paris were strongly opposed to this edict, "and especially desired that chemistry might not be taught." (*DeCuzé's Hist.*, &c. p. 10.) The garden continued gradually increasing in its collection of foreign trees and shrubs, under numerous successive intendants, till 1739, when the celebrated Buffon was appointed intendant; and he, among other improvements, planted, in 1740, an avenue of lime trees, which still exists.

The principal accession to the ligneous flora of France, however, dates from the connexion of that country with North America, which may be said to have taken place about the middle of the 18th century. Of the foreign trees planted in the garden about that time, the following still exist:—*Gleditschia triacanthos* var. *inermis*, 80 ft. high, sent from Canada by M. de la Galissonnière, the friend of Du Hamel, and governor of Canada, in 1748; *Sophora japonica*, the first plant sent to Europe from Japan about the same time, and now 64 ft. high; *Ailantus glandulosa*, brought from China about the same time, 68 ft. high; *Juniperus excelsa*, male, planted by Tournefort, who brought it from the Levant in 1702, 62 ft. high, with a clear trunk of 15 ft.; *Gymnocladus canadensis*, male, 58 ft. high; and a number of others which will be found enumerated in a description of the Paris Botanic Garden, in the *Gardener's Magazine*, vol. xii. The oldest trees at present in the garden, and some of which appear to have been planted soon after its establishment, are, *Acer monspessulanum*, 45 ft. high; *Celtis occidentalis*, 68 ft. high; *Quercus Ilex*, 42 ft. high; *Platanus orientalis*, 74 ft. high; and *Cedrus Libani*, 80 ft. high. This tree is the oldest and largest cedar in France: it was given to

Bernard de Jussieu, when he visited England in 1734, by the benevolent and enlightened Peter Collinson, who had raised some plants (of which he gave Jussieu two) from cones brought from Mount Lebanon. The tree in the Paris garden produces abundance of cones, and is considered the parent of all the cedars in France: it would, no doubt, have attained a greater height, had not the leading shoot been accidentally broken off some years ago (the person who showed it to us in 1815 said by the first shot fired against the Bastile), since when it has increased only in breadth.

Deleuze, who has given a history of the introduction of plants of ornament into France, in the *Annales du Muséum*, tom. viii., states that the taste for foreign trees and shrubs passed from England into France; but that the mode of procuring them from the former country being found too expensive, a plan was devised for importing them direct from America. At the head of this design was the celebrated Du Hamel, who induced his friend, Admiral Galissonnière, to send him several tons of seeds of trees and shrubs, gathered at random in North America. These were sown on a large scale on Du Hamel's estates at Le Monceau and Vriigny, and on those of his brother at Denainvilliers. They succeeded perfectly, and the plants raised were so numerous, that the botanists who afterwards examined them found among them several new species. The brother of Du Hamel the academician, who was the proprietor of Denainvilliers, appears to have had the chief care of these plantations. He also assisted his brother in the preparation of his works, and especially in the *Traité de la Culture des Terres*. The Duke d'Ayen, afterwards Maréchal de Noailles, made an extensive plantation of exotics at St. Germain en Laye, in which flowered, for the first time in France, some American walnuts, and the *Sophora japonica*. This park was open to all amateurs. It was the Maréchal de Noailles who persuaded Louis XV. to establish at Trianon that botanic garden in which Bernard de Jussieu disposed, for the first time, plants in families according to the natural orders of his system. The maréchal was one of the first four honorary members of the Linnæan Society of London. He died in 1793 at the age of 80 years.

The Chevalier Jansen purchased in all the ports of Europe, and in foreign countries, the trees which he hoped he could acclimatise in France; these he planted in his garden at Chaillot, and afterwards distributed among botanists and cultivators. On this spot, in Paris, adjoining the Barrière de Chaillot, may still (1835) be seen superb trees, the seeds of which have produced many others, which have been spread throughout France. That illustrious magistrate and philosopher, Lamoignon de Malesherbes, acclimatized on his estate of Malesherbes a great number of foreign trees and shrubs: he was the first in France to raise fruit trees from seeds on a large scale, in order to obtain new varieties. The celebrated Lemonnier of Montreuil, near Versailles, the friend of André Michaux, encouraged the introduction of trees and shrubs more than any of his contemporaries. He was the first patron of Michaux; and though, as a physician, he was much occupied at court, he employed the greater part of his income, and the whole of his leisure, in procuring rare trees and plants for his garden at Montreuil. There, in a bottom of bog earth, he had a multitude of different species of kalmia, azalea, rhododendron, and other shrubs, among which rose up the superb stems of the Canadian lily. In the shade of spruce firs, of acacias, of tulip trees, and of magnolias, grew the undershrubs of Lapland, of Siberia, and of the Straits of Magellan. His fortune and his garden were much injured during the revolution; but he lived to see the plants which he had introduced become common among his friends everywhere. He died at the age of 84 years.

Through the kindness of M. Vilmorin we are enabled to notice the present state of the different plantations mentioned or alluded to by Deleuze, and of others made by different proprietors about the same period. The plantations of Du Hamel were chiefly cut down, or otherwise destroyed, during the revolution; those of the physician Lemonnier, at Montreuil, were entirely destroyed; those at the Trianon remain, and contain some good specimens of

acacias, deciduous cypresses, pines, and cedars. The dimensions of some of the trees planted by M. Jansen have been sent us by Mr. Blaikie, who now (1835) resides at Chaillot, in a house built in the midst of them: among them are, an *Acer Opalus*, 50 ft. high, with a trunk $1\frac{1}{2}$ ft. in diameter; a *Sophora japonica*, 60 ft. high; and an *Ilex balearica*, 30 ft. high. A great many trees were planted in the great park at Rambouillet, about 1705, chiefly in avenues, after a design made by Le Nôtre, who died a few years before. The majority of the trees are abeles, and they have attained the height of upwards of 100 ft., though many have fallen down from age. Between the years 1787 and 1789 a great many American trees were planted in that part of the grounds at Rambouillet known as the Jardin Anglais, which have thriven well, and many of them have attained considerable size, as will appear from an account of them in the *Gardener's Magazine*, vol. xi. p. 42. and p. 205. At Thury, the property of the learned Vicomte Héricart de Thury (see *Annales d'Hort. de Paris*, tom. xi. p. 298.); at Balaine, near Moulins, the estate of Madame Aglaé Adanson, the daughter of the botanist Adanson, a descendant of Helvetius, and herself the author of *La Maison de Campagne*; at Nerac, on the estate of the Comte de Dijon; and at various other places; are collections of American trees and shrubs planted before the revolution, of which we have received notices from our correspondents, that will be found recorded, when we treat of the trees to which they refer. Near Metz, at Columbière, there are some fine trees of the pine and fir tribe, and many American trees, which were planted about the middle of the 18th century, by the Baron Tschoudi, the father of the baron of that name who was the inventor of herbaceous grafting; and who, after having been many years in the army, has retired to Columbière, and has there an extensive collection of trees and shrubs. At Mereville there are many fine American trees, which were planted by Mr. Blaikie, particularly the ailantus, which grows there to a large size, many specimens having attained the height of 80 ft. in 40 years. At St. Len, the ailantus has also attained a similar height in the same time, with a trunk of $3\frac{1}{2}$ ft. in diameter. One of the oldest magnolias in France is at Maillardière, a property in the neighbourhood of Nantes. An account is given of this tree in the *Nouveau Du Hamel*, tom. ii. p. 220.; and we have also been favoured with its history, communicated by the proprietor, M. le Comte de la Bretesche, to M. Durand de Lançon of Coutance in Normandy, and sent to us by him; and with a description of it by M. Nerrière, a nurseryman at Nantes. The particulars will be found under the head of *Magnolia grandiflora*: it will be sufficient to state here, that, after having sustained many injuries during the century that it has stood at Maillardière, the tree is still in existence, and is now upwards of 30 ft. high.

Historical notices and dimensions of many other large and old foreign trees have been sent us, and they will be found under the heads of their respective genera; but we may remark that there are few large and old trees in France comparatively with what there are in England; not only on account of the great changes which landed property has undergone in France, but because trees in that country are grown principally for timber and fuel, and have at no period been considered so much articles of luxury as they have been and are in England, which is supplied with timber for building from the Baltic, and with fuel from its coal mines.

The knowledge which we in England possess respecting the culture of trees in France may be said to date from the publication of the *Traité des Arbres et Arbustes*, by Du Hamel, in 1755. Du Hamel was contemporary with Miller and Collinson of London, and was in general correspondence with British botanists, to whom, in common with botanists in other parts of the world, he, in the preface to his work, acknowledges his obligations. In the first and second editions (in 2 vols. 4to) of his *Treatise*, he describes 180 genera and nearly 1000 species, without including those small under-shrubs, such as thyme, hyssop, &c., which technically are ligneous plants; and in the third edition, known as the *Nouveau Du Hamel* (in 7 vols. folio), which was

published from 1800 to 1819, nearly 2000 species and varieties are described, of which upwards of 500 are figured.

Du Hamel, in the preface to his work, says that he has treated of shrubs as well as trees, in order to lead to the pursuit of the useful through the medium of the agreeable. "There is reason," he says, "to hope that we shall be better listened to by the rich, when we propose to ornament their mansions with foreign trees, and their parks with thickets of flowering shrubs, than if we were simply to tell them to form plantations on lands unfit for producing corn or grass. If the self-love of the possessors of country seats is flattered by the view of common parks, notwithstanding the revolting uniformity of their thickets, which are only varied by differing in size or in form, is there not reason to hope that they will be much more highly gratified when the thickets in these parks offer that variety which is produced by different kinds of trees and shrubs, and which exhibits beauties suited to every season?" After giving directions for choosing flowering trees and shrubs to form thickets for the early part of spring, for the middle of spring, and for summer, he next shows the superior enjoyment to be derived from the culture of trees, to that which can be derived from the culture of herbaceous plants. "The most beautiful bed of hyacinths or tulips, when the flowers have once faded, leaves nothing but what is withered and unsightly; whilst the flowers of trees and shrubs which generally appear in spring are succeeded by the most vivid green leaves; and even in winter, after these have dropped, the ramification of the branches and spray is beautiful and interesting." (Preface, p. xviii.)

Du Hamel remarks that the greatest difficulty which opposed itself to his plan of rendering foreign trees and shrubs general in France was, that the greater part of them were not to be found for sale in the public nurseries. From this we may conclude that those who did introduce foreign trees and shrubs into France, during the 18th century, received them chiefly from abroad. Of this, indeed, there can be no doubt, since it is attested by a living witness, Mr. Thomas Blaikie, already mentioned, who is a native of Scotland, and has been settled in France as a landscape-gardener since the year 1776. In the *Encyclopædia of Gardening*, edit. 1835, p. 88., will be found a list of gardens and grounds laid out by Mr. Blaikie in France between 1776 and 1794, in which he mentions that for one place (Maison) he "went to England to buy the trees and shrubs; as at that time few trees or shrubs could be found in any nursery near Paris." Mr. Blaikie also laid out several places for the Duke of Orleans, and especially Monçcaux, the trees and shrubs for which were all procured from the Hammersmith Nursery.

The culture of foreign trees and shrubs in French nurseries appears to have commenced about the beginning of the present century, and the principal nurserymen who engaged in that branch were M. Noisette, whose father was gardener to Monsieur, afterwards Louis XVIII., at Brunoy, and M. Cels, who is now dead, and whose nursery is carried on by his son. On this subject, we refer for further details to the historical part of the *Encyclopædia of Gardening*.

Among the principal amateurs who have collected foreign trees and shrubs in France, since the commencement of the present century, may be mentioned, first and principally, the Empress Josephine, who had a collection at Malmaison of all that could be supplied from the London nurseries; the late Baron Pappenheim, who endeavoured to acclimatise many species at Coombe la Ville; Admiral Tchitchagoff, who has a fine collection at Scéaux; Monsieur de Magneville, near Caen, who is noted for his collection of pines; the Duke of Orleans (now King of the French), who has an arboretum at Neuilly, a catalogue of which was published by his gardener, Jacques, in 1833; M. Du Mont de Courset, at Boulogne; M. Soulange-Bodin, at Fromont on the Seine; M. Vilmorin, at Barres; M. le Baron Tsehondi, at Columbière, near Metz; M. le Comte de Montbron, at Clervaux, near Chatelherault; M. Ivoy, in the neighbourhood of Bourdeaux, celebrated for his collection of pines and firs; and General Lemarraiz, formerly aid-de-camp to Napoleon.

This last proprietor has planted in Normandy upwards of 60,000 of the *Pinus Laricio*; and, in the Forest of Fontainebleau, M. de Larminat has grafted 10,000 Scotch pines with scions of this valuable tree; an example well worthy of imitation by the proprietors of newly planted pine woods in Britain.

In the different botanic gardens in France, there are arboreta more or less extensive: the most complete is that of the Paris garden; but those of Metz, Strasburg, Montpellier, and Toulon are also good. In the latter there is a deciduous cypress which, in 35 years, has attained the height of 80 ft., with a trunk 9 ft. in circumference close to the ground. From all these gardens, and several others, we have had lists and dimensions of the trees, which will be found under the different genera.

Some of the nurseries have extensive collections: judging from their sale catalogues, those of Cels, Noisette, and Godefroy appear to be the best in Paris, or its neighbourhood; and those of Audibert of Taraseon, of the Baumans at Bolwyller, and of Jacquemet-Boncoford at Ammonoy, seem to be the most extensive in the provinces. The Bolwyller Nursery, situate near Mulhausen, in Alsace, was established by M. Joseph Baumann (who was formerly gardener to the late Grand-Duchess of Courland), in conjunction with his brother Augustine, about the end of the last century. The establishment of M. Soulange-Bodin at Fromont, in the neighbourhood of Paris, is perhaps the most remarkable in France. It combines the most extensive system of propagation both of hardy and house plants, ligneous and herbaceous, with an institution for the instruction of young men in the science and practice of horticulture. The nature of this establishment, and its extensive collections, will be found at length in the *Annales de l'Institut de Fromont*; in the *Encyclopædia of Gardening*, edit. 1835; and in the *Gardener's Magazine*, vol. ix. p. 141, and in vol. xi.

The individuals who have exercised most influence on the introduction of foreign trees and shrubs into France appear to have been Du Hamel, André Michaux, and Du Mont de Courset.

Henri-Louis Du Hamel du Monceau was born at Paris in 1700, and died in 1782. He was proprietor of several estates, besides that from which he takes his designation. He was appointed inspector of the French navy, and was a member of the *Académie des Sciences*, and a Fellow of the Royal and other Societies in Britain, as well as of several on the Continent. He was the author of a number of works on agriculture, forest trees, fruit trees vegetable physiology, and rural economy, and of the *Elements of Naval Architecture*, all of which appeared between the years 1747 and 1768. His most important work is the *Physique des Arbres*, which contains much of what, in this country at least, has been attributed to subsequent discovery. We allude more particularly to the theory of the ascent of the sap by the wood, and its descent by the bark. Du Hamel is said to have been a man of great modesty, and to have devoted his life to agricultural pursuits, to the mechanical arts, and to his duties as a public officer. He left no child, and his estates went to his nephews. One of these, Fongeroux de Bondaroy, has published an interesting *Mémoire sur les Pins*, inserted in the *Mémoires de l'Académie des Sciences*. Vigny, Du Hamel's principal estate, now belongs to M. Charles de Fongeroux, his grand-nephew, who not only takes the greatest care of the trees left to him by his grand-uncle, but plants extensively himself. Denainvilliers and Monceau now belong to M. de Denainvilliers, the grandson of the brother of Du Hamel du Monceau. There are on these two estates a number of very fine exotic trees, of which the present proprietors take the greatest care. The finest deciduous trees are those that were planted by Du Hamel in some marshy ground at Monceau; and some of them have attained the height of 90 ft.

André Michaux was born in the Park of Versailles, in 1746, and soon evinced a taste for agriculture and botany, which was fostered by his early patron, the court physician, M. Lemonnier. In 1777 he studied botany under Bernard de Jussieu, at Trianon; and in 1779 he was studying in the *Jardin des Plantes*. Soon after this he came to England, and

returned to France with a great number of trees, which were planted in the gardens of M. Lemonnier, and of the Maréchal de Noailles, where they succeeded perfectly. He often used to take from these gardens a packet of grafts, and, going through the woods of Versailles, he would graft them on the trees already there. In 1780, he went to botanise on the mountains of Auvergne with several botanists, among whom were Lamarck and Thouin. Michaux was the most active of all of them; besides his musket, haversack, portfolio, and several specimen boxes, he carried in his pocket seeds of the cedar of Lebanon, which he sowed in favourable situations. Soon afterwards he went to the Pyrenees and travelled in Spain; and, in a short time, accompanied the nephew of the celebrated Rousseau to Persia, the latter being appointed consul to that country in 1782. He went to Aleppo, Bagdad, the Tigris, the Euphrates, Bassora, and many other places, sending home numerous seeds to Thouin, Malesherbes, and others. Persia at that time was a prey to civil wars, and Michaux, plundered of every thing by the Arabs, was supplied with the means of continuing his journey by M. de la Touche, the English consul at Bassora, though France and England were at that time at war; M. de la Touche, his biographer observes, thinking that a naturalist, who travelled for the good of humanity, ought to be protected by every nation. In this part of the world Michaux remained two years, traversing mountains and deserts from the Indian to the Caspian Sea, and proving that the provinces situated between 35° and 45° of latitude in the East have supplied most of our trees, exclusive of those which belong to America. He here verified the fact first noticed by Kæmpfer, that the male flowers of the date will keep during the year, and yet impregnate the female. He sent home sculptured ruins from the palace known as that of Semiramis, near the Tigris, and various other antiques, and objects of natural history. He returned to Paris in June, 1785, and was chosen soon after to go to the United States, to collect seeds of trees and shrubs; to establish an entrepôt for them in the neighbourhood of New York; and to get them sent from that to Rambouillet, which was destined to receive them. He was also commissioned to send home American game. He arrived at New York in October, 1785; established a garden there; traversed New Jersey, Pennsylvania, and Maryland; and, after the first year, he sent home twelve boxes of seeds, and 5000 young trees, together with some Canadian partridges, which afterwards bred at Versailles. In September, 1789, he went to Carolina, making Charleston his depôt; he traversed the Alleghany Mountains, and the whole country north and south, leaving his son at Charleston, in charge of the gardens there. From this place he sent home numerous seeds, and many hundreds of young trees. In April following, he set out to reconnoitre the sources of the Savannah; and there he discovered *Magnolia auriculata*, *Robinia viscosa*, *Azalea n. coccinea*, a *Kalmia*, a *Rhododéndon*, and many oaks and other trees not before known. The manner in which he travelled, his intercourse with the native Indians, and the accidents he met with, are extremely interesting. Whenever he discovered a new plant, it inspired him with such enthusiasm, that he no longer felt fatigue. The discovery of a new *Pavia*, and of the *Pinckneya pibens*, gave him great pleasure. He arrived at New Providence in February, 1799, and returned to Charleston in May of the same year. He afterwards visited the highest mountains of Carolina. The dangers he experienced there convinced him of the necessity of having two guides, because one might perish by the road by a thousand accidents, and it would be impossible for a European to find his way alone through the country. He found in these mountains vast tracts covered with rhododendrons, kalmias, and azaleas, and with forests of trees altogether impenetrable. War, at this time, was declared between France and England; and Michaux was afraid of being forced to leave America. He had been for a long time occupied with the idea of determining the native place of all the American trees; and also at what latitude they begin to grow rare, and where they disappear entirely: in short, he wished to ascertain up to what height they are found on the mountains, and in what soil they prosper best. He con-

sidered the native country of a tree to be that in which it is most numerous, and where it acquires the greatest height and thickness. Thus he fixed on Kentucky as the native country of the tulip tree, because it there forms vast forests, has a trunk commonly 7 ft. or 8 ft. in diameter, and grows 120 ft. high, thriving in a moist clayey soil, but not in one that is frequently inundated. In higher or lower ground, or in a different soil, these trees become smaller and more rare. It was with a view to trace in this manner the botanical topography of North America, that Michaux visited the Floridas, and went as far as Hudson's Bay. He left Charleston in April, 1792; arrived at Quebec in June of the same year; and reached Tadoussac, lat. 52°, in October, 160 leagues from any human habitation. He afterwards planned a journey to Mexico, for the benefit of the United States; but, after very many journeys, he returned to Paris by Amsterdam, where he arrived on the 3d of December 1796, after ten years' absence. He found his friends well, but was grieved beyond measure to learn that the beautiful plantations of Rambouillet, to which he had sent 60,000 young trees, had been destroyed during the revolution, and that but a very small number of the trees was remaining. Seeing that tranquillity was restored, he instantly thought of repairing the loss. After unsuccessfully endeavouring to get sent again to America, he was sent to New Holland. He stopped at the Isle of France, and was very desirous of going to Madagascar; in which island he was attacked by the fever, and he died there in November (an ix.), 1803; aged 57 years.

Michaux not only sent many new trees and shrubs into France, but he sent great quantities of the seeds of the more useful species; such as *Juglans Piccau*, used for making furniture, and which produces the nut oil; *Taxodium distichum* (the deciduous cypress), suitable for planting in very moist soil; *Nyssa caroliniana*, useful for the naves of wheels; *Quercus tinctoria*, for tanning and dyeing; and *Q. virca*, which, he says, grows rapidly on the sandy beach, exposed to the stormy winds of the ocean, where scarcely any other tree can exist, and the wood of which is excellent for ship-building; to these may be added the caryas of Pennsylvania, the tulip trees, and the American ashes, maples, &c., which, in many parts of France, are preferable to the indigenous trees. The administration of the Museum, aware of the services rendered to natural history by Michaux, ordered his bust to be placed on the façade of the green-houses, along with those of Commerson, Dombey, and other travellers who had enriched their collection.

Michaux was too fully occupied in travelling to have much leisure to write; nevertheless, he is the author of *Histoire des Chênes de l'Amérique Septentrionale*, published in 1804; a *North American Flora*; and a *Memoir on the Date Palm*. The particulars of his life, at great length, and proportionately interesting, will be found in the *Annales du Muséum*, tom. iii, p. 191.; from which this notice of his life has been abridged.

F. A. Michaux, the author of *Histoire des Arbres de l'Amérique*, after his father's death, was sent to Charleston, by the French government, to bring over the trees collected in his father's nurseries, and supplies of seeds. During his stay in America, M. Vilmorin informs us that he sent to the *Administration Forestière* larger quantities of acorns and other seeds of foreign trees, than had ever before been sent over from that country. He took that opportunity of visiting Kentucky, the Tennessee, and of penetrating nearly a thousand miles beyond the Alleghany Mountains. On his return to Europe, he published his great work on the trees of North America, and other memoirs on relative subjects; particularly one *Sur la Naturalisation des Arbres Forestières de l'Amérique*, &c. He now resides in the neighbourhood of Paris, and appears to be as enthusiastically devoted to the study of trees and shrubs as his late father. We are much indebted to him for various useful communications having reference to the *Arboretum Britannicum*.

Georges Marie Louis Du Roi, Baron de Courset, author of the *Botaniste Cultivateur*, was the Du Hamel of his time; and, after the revolution, his example and exertions contributed, even more than the influence of the Em-

press Josephine, to spread a taste for exotic trees and shrubs, and the formation of ornamental plantations. He was born in 1746, at the Château de Courset in the Haut Boulonnais. After having received an excellent education, he entered the army at the age of 17 years, and was soon after sent on duty to Languedoc, where the plants of the Pyrenees gave birth to his enthusiastic taste for botany. In 1784 he left the army, and devoted himself wholly to the improvement of his estate at Courset, where, in a short time, he formed by far the richest collection of plants in France, and created an establishment which ranked at that time with the gardens of Malmaison, Kew, &c. In an arid chalky soil, so unproductive as to be called a desert, M. Du Mont created an excellent kitchen-garden, a large orchard, and an ornamental garden devoted to the culture of foreign plants. These gardens will be found described in the *Annales de la Société d'Horticulture de Paris*, tom. xiv.; and in the *Gardener's Magazine*, vol. xii., from our personal inspection. It may be sufficient to state, that, though these gardens do not display fine turf, water, or fine gravel, yet they are of intense interest in point of culture; and that the collection of hardy trees and shrubs, which have attained a considerable size, is not surpassed by any in the neighbourhood of London, in regard to the number of species which it contains. The collection of herbaceous plants is formed into a series of concentric beds. The trees and shrubs are disposed in groups, according to the season of the year at which they flower, as suggested by Du Hamel; but these groups are so thinly planted that room is left for each tree and shrub to acquire its natural size and form. There is an extensive collection of fruit trees, including all the varieties that could be procured in Europe and America. The peat-earth plants are numerous, as are the hot-house and green-house plants. The hot-houses are 200 ft. and the pits 150 ft. in length. In the year 1789 M. Du Mont visited the principal gardens in the neighbourhood of London, and, on his return to his family, was immediately arrested and imprisoned by the government; but he was as promptly set at liberty through the influence with the Committee of Public Safety of his friend, the celebrated Professor Thouin. M. Du Mont published various articles in the public journals of his day; but his principal work is the *Botaniste Cultivateur, or Description, Culture, and Use of the greater Part of the Plants, Foreign and Indigenous, which are cultivated in France and England, arranged according to the Method of Jussieu*, which appeared in five volumes, 8vo, in 1802, and to which two supplementary volumes have since been added. This work has had the same celebrity in France that *Miller's Dictionary* has had in England. M. Du Mont died in June, 1824, at the age of 78 years; his estate is now the property of his daughter, Madame la Baronne Mallet de Coupigny, who has presented the green-house and hot-house plants (with the exception of the pelargoniums) to the *Société d'Agriculture de Boulogne*, but who cultivates the collection of hardy articles, and more especially the trees and shrubs, with the greatest care. The place is visited by gardeners, botanists, and naturalists from every part of the world; and no name in France is mentioned with greater respect than that of the patriarch De Courset.

SECT. II. *Of the Indigenous and Foreign Trees and Shrubs of Holland and the Netherlands.*

THE indigenous trees and shrubs of Belgium and Holland are very few, partly from the limited extent of territory, but chiefly from the great uniformity of the surface, the soil, and the climate. The only *Flora* which has been attempted of Belgium is that of Lejeune and Courtois (reviewed in *Gard. Mag.*, vol. x. p. 449.), of which only a part has been published. Holland can hardly be said to have an indigenous ligneous flora; but into that country foreign trees and shrubs were introduced as soon as they were into any other in Europe. The botanic garden of Leyden, and its earliest catalogues, may be referred to as a proof of this; but for its history, and for various details re-

lating to the subject, we must refer to the *Encyclopædia of Gardening*, edit. 1835, p. 69. M. E. de Wael, the director of the botanic garden at Antwerp, has furnished us with a list of the indigenous trees and shrubs of that neighbourhood, which is even more meagre than we had anticipated; and another correspondent has sent us the following remarks on the subject of the Dutch ligneous flora. "Many causes combine to operate unfavourably on the growth of trees and shrubs in Holland; the numerous heavy winds in the neighbourhood of the sea, and more particularly the north-west wind, destroy the tops of the high-growing trees, break their branches, and, by shaking their trunks, loosen the roots in the soil, or blow the trees down. This is chiefly occasioned by the little depth to which the roots can penetrate into the ground; for, as soon as they reach the water, they are compelled to take a lateral direction, in consequence of which the trees soon become sickly, or are suddenly loosened from the soft, loose, humid soil by the wind. We have here much marsh and fen land. This soil, which is extremely well adapted for supplying turf or peat, is unfavourable to the growth of timber. Should much rain and strong winds occur, the trees on this soil cannot exist long enough to become old, nor even to have good trunks. In order, therefore, to prevent their being blown down, they must, from time to time, be tied or propped up: but the trouble and expense of this operation cause it to be neglected; instead of it the trees are severely lopped, and this, by causing them to throw down a greater quantity of roots into the wet substratum, only renders them more sickly. The truth of this fact may be perceived in the trees in and about most of the Dutch cities.

"When these obstacles do not occur, the trees exhibit a better growth; for the elms on the embankments in Zealand, which have their roots in a good stiff clay soil, and stand high out of the water on the dykes, endure the sea winds without sustaining any injury; besides which, these trees are judiciously pruned, and this, of course, greatly contributes to their large growth and handsome appearance. Whenever the trees are on high ground, and grow in masses, so as to protect one another from the winds, the vegetation is luxuriant, as is the case at the Hague, and in the woods near the Loo. This strikes even a superficial observer at the first glance.

"Most of the country seats in Holland were formerly near Rotterdam, along the Gravenweij, for example; and at Amsterdam, in the neighbourhood of the Diemermeer: but, as all these seats have been demolished, and new ones formed in the high sandy grounds of the provinces of Guelderland and Utrecht, not many remarkable old trees remain in the former district: Those which time might have spared have been cut down in consequence of the removal of the country seats. The *Pópulus canadénsis* [?] *monilifera*] appears every where here in an excellent condition, and grows in places where no other tree will thrive. On the sites of some of the old country seats, especially where the ground is elevated, old tulip trees and catalpas are found, both of which bloom freely. The new country seats are laid out with much taste, in parks on a large scale, and on high grounds, in the provinces of Utrecht and Guelderland; and they are planted with exotic trees and shrubs, which afford very favourable expectations for the future. — *W.*"

But, though Holland and the Netherlands are deficient in an indigenous flora, they are by no means so in collections of plants from other countries. This is ascertained from the early catalogues of the different botanic gardens, and from the magnificent publications on botany and gardening which issued from the press of Leyden, Amsterdam, and other Dutch or Flemish cities, in the 17th century. Great part of the Netherlands, from its moist peaty soil, is particularly adapted for the growth of American trees and shrubs; and these, especially all the more showy-flowering kinds, are in popular cultivation. (See *Gard. Mag.*, vol. vii. p. 279., xi. p. 103. and p. 219.)

The best collections of foreign trees and shrubs, at present in the Low Countries, are in the different botanic gardens; in the garden of the Botanical and Horticultural Society of Ghent, in the garden of the King of the Bel-

gians at Lacken, in the Duc d'Arcmburg's seat at Enghien, and that of Sir Henry T. Oakes near Tournay. The nurseries of Holland are celebrated for their fruit trees, and those of the Netherlands for their magnolias and azaleas, and other peat-earth trees and shrubs. That of M. Parmentier at Enghien has long been remarkable for containing a great many species in a very limited space; and that of M. le Candele at Humbeque, near Brussels, contains the best collection of the genus *Cratægus* in the Low Countries. Some account of this nursery, with notices of its more remarkable trees, will be found in the *Gardener's Magazine*, vol. xi. p. 537.

In the garden at Lacken there are a few fine specimens of foreign trees, particularly a tulip tree, which Mr. McIntosh, the head gardener to the King of the Belgians, informs us, had, in 1834, a clear stem of 20 ft., the diameter of which at the surface of the ground was fully 3 ft., and at the height of 20 ft. about 2 ft.; the head was globular and compact. This tree flowered and ripened seeds every year. When Lacken belonged to France, the palace was occupied by the Empress Josephine, who brought her gardener from Paris to superintend the gardens; and the poor man, while he was gathering the seeds of this tulip tree, fell from it, and broke his neck. The trees and shrubs in the Brussels Botanic Garden have been planted within the last fifteen years: those in the Ghent Botanic Garden are much older; among them is a *Pópulus canadénsis*, 100 ft. high, and upwards of 17 ft. in circumference at 1 ft. from the ground. There are, a *Robínia Pseud-Acácia*, 60 ft. high; a *Catálpa syringa-fólia*, with a trunk between 8 ft. and 9 ft. in circumference; *Vibúrnum Opulus*, 22 ft. high; two tulip trees, 70 ft. high; a *Salisbúria*, 23 ft. high; *Gymnó-cladus*, 70 ft. high; lime trees, 60 ft. high; and *Magnólia auriculáta*, conspícua, and tripétala, from 20 ft. to 25 ft. high. In the grounds of Mr. Herry of Mariakirk is a *Catálpa*, 40 ft. high, with a trunk 6 ft. in circumference at 1 ft. from the ground. In the grounds of Baron le Norman, near the same town, there are, an *Ailántus* (there called the Virginian sumach), 30 years planted, and 45 ft. high; and a *Juníperus virginiana*, 40 years planted, and 30 ft. high. The largest salisburia in Holland is in the botanic garden at Utrecht, its height being nearly 50 ft.

In consequence of the present unfriendly feeling between Holland and Belgium, we have been unable to procure notices of the trees and shrubs of the more remarkable places of either country. We know, however, that there are many fine specimens, and that though the winters are colder than those of England, yet that the summers are warmer, and that the greater part of the deciduous American trees and shrubs thrive there as well as in England. Many of the finest azaleas in our nurseries, and some varieties of magnolia and rhododendron, have been raised from seed in the neighbourhood of Ghent. The winters, however, are unfavourable for evergreens, and but few of these are to be found in any part of the country. In Smith's *Tour on the Continent*, Neill's *Horticultural Tour*, and in various articles in the *Gardener's Magazine*, will be found descriptive sketches of many of the small gardens of Holland and the Netherlands, all more or less remarkable for their American trees and shrubs. Of large places which may be compared with the country seats of England, and which might be supposed to afford many examples of fine old trees, there are comparatively few, as has been already observed above by a correspondent, a native of the country.

SECT. III. *Of the Indigenous and Foreign Trees and Shrubs of Germany, including Hungary.*

THOUGH this portion of Europe is of great extent, yet its ligneous flora is much less varied and numerous than that of France. The reasons are, that it extends in longitude more than in latitude; that it contains few very lofty mountains, and embraces but a small latitudinal portion of the sea shore. It includes Hungary, however, which enjoys a greatly diversified surface, and an extensive range of mountains, with a ligneous flora which has furnished some

fine trees and shrubs to the rest of Europe; for example, several different species of *Pyrus*, and the common and Josikea lilacs. The following enumeration, taken from Roth's *Flora Germanica*, Willdenow's *Baumzucht*, and the *Flora Hungarica* of Waldstein and Kitabel, and kindly revised for us after it was in type by Baron Jacquin of Vienna, indicates those indigenous trees and shrubs which Germany possesses, that are not also indigenous in Great Britain and Ireland; those which are plants of cultivation, or doubtful as natives, being, as in previous lists, indicated by a star.

Ranunculacææ. *Clematis Flammula*, *Atragène austriaca*.

Cistiucæ. *Helianthemum Fumana*, sp. "alpinum foliis Ajugæ" *Ruppini*, vineale.

Polygalacææ. *Polygala Chamæbuxus*.

Tiliacææ. *Tilia pauciflora*, vulgaris, argentea.

Acerincææ. *Acer Pseudo-Platanus*, *platanoides*, austriacum, tatàricum, obtusatum.

Celastrincææ. *Euonymus latifolius*, verrucosus.

Rhamnucææ. *Rhamnus alpinus mas*, a. fœm, saxatilis, infectorius, pumilio, rupèstris.

Staphylacææ. *Staphylèa pinnata*.

Anacardiacææ. *Rhus Coriaria*, *Cotinus*.

Leguminosææ. *Genista radiata*, germanica, sagittalis, procumbens; *Cytisus Laburnum*, alpinus, nigricans, nigricans var. elongatus, hirsutus, supinus, capitatus, austriacus, Weldeni; *Colutea arborescens*, * cruenta; *Coronilla Eimerus*; *Ononis Natrix*.

Rosacææ. *Rosa sempervirens*, fecundissima, gallica, alpina, pendulina, pyrenæica, rubrifolia, alba; *Rubus*, several ligneous species of, described in *Rubi Germanici*, Bonn, 1822; *Spiræa salicifolia*, *chamædrifolia*, *ulmifolia*, mædia *Schmidt*, *oblongifolia*, incana.

Pomacææ. *Mespilus germanica*, *Amelechier vulgaris*; *Pyrus nivâlis*, *bolwylleriâna*, *Chamaemespilus*, *salviaefolia*; *Cydonia vulgaris*, *Crataegus monogyna*.

Tamariscinææ. *Tamarix gallica*, germanica.

Amygdalacææ. *Cerasus Mahaleb*, *Chamaecerasus*.

Philadelphicææ. *Philadelphus coronarius*.

Grossulacææ. *Grossularia Uva crispa*.

Caprifoliacææ. *Lonicera nigra*, alpigena, cærulea, *Xylosteum*; *Sambucus racemosa*.

Cornacææ. *Cornus mas*.

Compositææ. *Helichrysum Stæchas*.

Ericacææ. *Erica herbacea*, *Ledum palustre*, *Andròmeda calyculata*; *Rhododèndron Chamæcistus*, ferrugineum, hirsutum.

Oleacææ. *Syringa vulgaris*, *Josikæa*.

Labiataææ. *Hyssopus officinalis*, *Teucrium montanum*.

Thymelacææ. *Daphne Cneorum*.

Elæagnacææ. *Elæagnus angustifolia*.

Euphorbiacææ. *Euphorbia sylvatica*, *Buxus sempervirens*.

Urticææ. *Urtica alba*.

Ulmacææ. *Ulmus effusa*.

Cupuliferææ. * *Castanea vesca*, *Corylus tubulosa*; *Quercus austriaca*, pubescens.

Betulincææ. *Betula pendula*, pubescens, fruticosa, humilis *Schrank*; *Alnus glutinosa* var. *quercifolia*, incana, ? incana var. minor viridis, ovata, carpatica.

Salicincææ. *Salix Ammanniana*, holosericea, *Hoppeana*, *Jacquiniâna*, hastata, bigemmis, fissa, retusa, lanata, depressa, polyandra *Weigel*, *Meyeriâna*, mollissima, præcox, riparia, serpyllifolia, silesiaca, spathulata, *Starkeana*, uliginosa, undulata, *Weigeliâna*.

Coniferææ. *Pinus pumilio*, nigricans; *Abies excelsa*, *Picea*; *Larix europæa*, *Juniperus Sabina*.

Germany may be fairly considered as possessing the ligneous flora of Britain in addition to her own, though, perhaps, there may be a few obscure species as

exceptions. Supposing this to be the case, the ligneous flora of the British Isles, added to the species above enumerated, will give to Germany a flora of upwards of 360 species of indigenous trees and shrubs.

The introduction of foreign trees and shrubs into Germany, subsequently to the time of the Romans, and to that of the foundation of religious corporations, appears to have commenced with the establishment of botanic gardens. The first tree of note, of the introduction of which we have any record, is the horsechestnut, which, according to Beckmann (*Hist. of Invent., &c.*), was brought to Vienna by the botanist Clusius, somewhere about 1576. In Clusius's *Rariorum Plantarum, &c.*, published in 1601, he states that in 1581 the horsechestnut was considered as a botanical rarity, but that in 1588 there was a tree at Vienna which had been brought there twelve years before, but which had not then produced bloom. M. Bon de Saint-Hilaire (*Mémoire sur les Marrons d'Inde*), however, says that the horsechestnut passed from the mountains of Thibet to England in 1550, and thence to Vienna in 1588. The first plant of *Robinia Pseud-Acacia* was brought to Vienna in 1696; and the remains of it are still living in the courtyard of the palace formerly occupied by Count Fries in the Place Joseph, and now belonging to Baron Sina. The ground on which this tree stands was formerly part of the garden of a convent of nuns, founded by the widow of Charles IX. of France, whose high-steward was the celebrated Augerius, Baron de Barbeck, the friend of Clusius. The oldest foreign trees in Austria are at Schönbrunn, and consist chiefly of tulip trees, platanus, acers, juglans, robinias, and cratægus, planted about the middle of the last century, or earlier. There is a more complete collection, though not quite so old, in the grounds of Prince Lichtenstein at Eisgrub, near Nikolsburg. About the middle of the last century, this nobleman sent M. van der Schott, a German, to North America; who collected there an immense quantity of seeds, which were sown on the prince's estates in Austria, Moravia, and Bohemia, and now form immense forests.

One of the oldest exotic trees in Germany is a *Thuja occidentalis*, near the old castle of Heidelberg, a drawing of which has been sent us by M. Ritter of Pesth, and which must have been planted when the grounds round the castle were laying out by Solomon Caus, as it bears a ticket stating that it was placed there in 1618. Caus began to plant the castle garden in 1615. (*Metzger's Castle of Heidelberg*, p. 60.) This venerable tree is at present about 30 ft. high, with a naked trunk leaning to one side, and a very few branches at top. In the gardens of this castle there are two large yew trees, which were planted in 1650, and some cornelian cherry trees (*Cornus mas*), which were brought from Neuburg on the Danube in 1769. There are also some very old lime trees. The Margraves of Baden have from the earliest ages been much attached to planting and gardening. In the grounds of the ancient grand-ducal palace of Durlach near Carlsruhe, which was the residence of this family for many centuries, and a part of the palace walls of which are supposed to be as old as the time of the Romans, there is an ash 140 ft. high, and 19 ft. in circumference at one foot from the ground. A board fixed to the trunk states that it was 300 years old in 1802. As the ash is not indigenous in the neighbourhood, this ash is, probably, the oldest planted tree in Germany. At Durlach, also, there are the remains of an avenue of chestnuts: the trunks are hollow, but some of them are 120 ft. high and 15 ft. in circumference: they are thought to have been planted about the end of the sixteenth century. The road from Durlach to Carlsruhe is through an avenue of Lombardy poplars, the oldest and the highest in Germany; none of the trees are under 90 ft. high, and many of them are above 120 ft. Nothing of the kind can be more sublime. The worthy old Margrave Charles, the first Grand-Duke of Baden, who died about 1805, and one of his sons yet alive, the Margrave William of Baden, may be reckoned amongst the most zealous promoters of the planting of foreign trees and shrubs; in proof of which, we need only refer to the parks at Carlsruhe, Schwetzingen, Mannheim, and Baden Baden.

By looking into the various catalogues of the German botanic gardens, and particularly into those of Giessen, founded in 1605, and of Altorf, Nuremberg, Rientel, and upwards of twenty others, founded between the commencement of the 17th century and the middle of the 18th century, the dates of the introduction into Germany of a number of trees and shrubs may be found by the curious. It will be sufficient for the purpose of this work, if we commence with the introduction of American trees and shrubs into Germany, which took place shortly after their introduction on a large scale into France; the Argyll of Germany being Prince Lichtenstein of Eisgrub; and the Du Hamel of that country being Baron Otto von Münchhausen of Schwöbbache, near Pymont, in Westphalia, now united to the kingdom of Hanover. This gentleman was the author of a work which obtained great celebrity in Germany in his time, entitled *Der Hausvater* (the Father of a Family). This book, which was printed in 1765, may be compared to the British encyclopædias of domestic economy, except that in it agriculture, gardening, and rural affairs bear a more conspicuous part than housewifery and cookery; it contains a descriptive list of new and desirable trees and shrubs, with directions for their culture, and for their disposition in lines; arguing against clipping them into geometrical figures, as was then the mode. We are assured (see *Gard. Mag.*, vol. ii. p. 386.) that it was the reading of this work, and especially the arguments which it contained in favour of a more natural mode of disposing and managing trees and shrubs in gardens, that gave the Empress Catharine a taste for English gardening; and that it was thus the means of introducing that taste into Russia. As Hanover was at this time closely connected with England, by being under the government of the same monarch, there can be little doubt that the trees planted at Schwöbbache would be procured from the nurseries of this country. Contemporary planters were, in the Hanoverian dominions, Count Veltheim of Harbecke, and Hinuber of Marienwerder near Hanover: also the Duke of Dessau, at Wörlitz, near Dessau, in Anhalt; the Elector of Hesse, at Wilhelmshöhe, near Cassel; Prince Lichtenstein, on his various estates in the Austrian dominions; and the Emperor of Austria, at Schönbrunn, near Vienna. Besides these princes, and Margraves of Baden, already mentioned, the following princes of Germany have distinguished themselves by planting foreign trees: Frederick the Great, and the present King of Prussia, Frederick William IV.; the late King of Saxony, Frederick Augustus IV.; the late King of Wurtemberg, Frederick William; the late Grand-Duke of Weimar; the late Duke Ernest of Saxe-Coburg-Gotha; Prince Prinus of Dahlberg; the Grand-Duke of Frankfurt; and the late Duchess of Courland, at Loebichan in Saxony. We have received Return Papers from all these places, from which we find that some of the foreign trees first planted in them still exist. Schwöbbache is in the possession of the grandson of the author of *Der Hausvater*, and contains a number of very interesting trees. Among these are, a tulip tree, near a pond, 120 years planted, which is 80 ft. high; *Nýssa aquática*, 60 years planted, which forms a magnificent tree 40 ft. high, with a wide-spreading head and branches drooping to the ground: it is in a low moist situation, and its roots, which extend to a great distance, send up innumerable suckers; in the autumn the leaves, before dropping off, become as red as blood. The finest tree of this kind in England is on the Duke of Wellington's grounds at Strathfieldsaye; it is 30 ft. high, and, being rather in a moist situation, will probably one day rival the tree at Schwöbbache, which, in all probability, is the finest specimen of *Nýssa* in Europe. *Córylus arboréscens* (? *C. Colúrna*), at Schwöbbache, 100 years planted, forms a regular-headed tree, with a straight clean trunk 2 ft. in diameter. *Acer eriocarpum*, *saccharinum*, and *Opalus*, have been 80 years planted, and are noble trees; *Aesculus Pávía* and *fláva* are stately trees, and flower freely; *Robínia Pseud-Acácia*, 120 years planted, is a large and most picturesque tree; *Ulmus americana*, 120 years planted, and *Juglans cinerea* and *nigra*, 80 years planted, are noble trees. There are specimens of *Castanea vesca*,

120 years planted; and of *Plátanus orientális*, 100 years planted. *Pínus Stróbus*, 80 years planted, is 100 ft. high. Many of the old trees are in a state of decay, but the present baron still keeps up the collection by planting young ones; and he adds continually to the species, appearing to be as enthusiastically devoted to trees and shrubs, and to gardening generally, as his grandfather. At Harbecke, Count Veltheim's, there are many old foreign trees and shrubs, and a very full collection of young ones.

At Wörlitz there is what is generally considered the fullest collection of old specimens of American trees in Germany; and there they thrive remarkably well on a loamy soil, in a situation damp but not very wet. Many of these trees produce seeds, which are sent to all parts of Germany. These trees, we are informed by M. Schoch, the Duke of Dessau's garden director (and the son of the director of the same name who laid out and planted the garden), were raised on the spot, from seeds brought from England by the Duke Leopold Frederick Francis, who formed the garden between the years 1760 and 1770. A minute and accurate account of all the trees in the garden has been kindly sent us by M. Schoch, with remarks on their different degrees of hardiness, which are very interesting. It appears that the cedar of Lebanon, the common laurel, the Portugal laurel, and even the spurge laurel, require protection during every winter; and that the *Thúja orientális*, the different varieties of common tree box, the *Cratægus Pyracantha*, the common holly, the *Acer erceticum*, the *Negundo*, the *Æsculus Pavia* and *flava*, the *Ailántus glandulosa*, the *Amýgdalus communis*, the *Celtis Tournefortii*, the *Cercis Siliquastrum* and *canadensis*, the *Taxodium distichum*, the *Cytisus Laburnum*, the *Castanea vesca*, the tulip tree, the *Morus alba* and *nigra*, the *Broussonetia*, the *Plátanus orientális*, the *Caragana arborescens*, the *Robinia inermis* and *viscosa*, the *Sophora japonica*, and the *Salix babylonica*, are all killed down to the surface of the ground when the cold is from 20° to 25° of Reaumur, but that they spring up again the following year from the root. Of this list, those which suffer the least are, the holly, the box, the laburnum, the deciduous cypress, and *Robinia inermis*. It is to be observed, that 25° Reaumur, which is exactly 25° below 0 Fahrenheit, is a degree of cold never experienced in any part of either Britain or Ireland, though Wörlitz is about half a degree south of London, and the whole of Scotland is farther north than any part of Germany. Wörlitz has been described and praised by the Prince de Ligne, and, till within the last twenty years, was considered one of the very first places in Germany. A description, at length, of these gardens, translated from an account of them published by the present director, Schoch, will be found in our *Encyclopædia of Gardening*, edit. 1835, p. 188.; and a particular account of the more remarkable trees that they contain is given in the *Transactions of the Prussian Horticultural Society*, vols. iv. and v.

At Schönbrunn, which was planted soon after Schwöbbache, there are several fine specimens of trees, and in particular a *Salisburia*, between 50 ft. and 60 ft. high, which was received from Loddiges' Nursery, and planted there in 1781 (Jacquin *Ueber den Ginkgo*, p. 3.); a *Sophora japonica*, between 80 ft. and 90 ft. high; *Liriodendron Tulipifera*, between 70 ft. and 80 ft. high; *Acer striatum*, between 30 ft. and 40 ft. high, with a trunk 18 in. in diameter; *Æsculus Hippocastanum*, between 90 ft. and 100 ft. high; *Æ. Pavia* and *Æ. flava*, between 30 ft. and 40 ft. high; *Kölreutèria*, 60 ft. high; *Robinia Pseud-Acacia*, 60 ft. high; *Gleditschia triacanthos*, 70 ft. high; *Fraxinus lentiseifolia* and *Ornus europæa*, about 40 ft. high; *Catalpa*, between 30 ft. and 40 ft. high; *Juglans regia*, between 60 ft. and 70 ft. high; *Pópulus dilatata*, upwards of 90 ft. high; *Plátanus orientális*, between 70 ft. and 80 ft. high; *Abies excelsa*, 90 ft. high; *Larix europæa*, 60 ft. high. The cedar of Lebanon, the *Laurus nobilis*, the *Arbutus*, the *Diospyros*, the *Photinia*, the *Aristotelia*, and some other trees, do not stand the open air at Vienna; and some of the magnolias, the *Cercis*, the *Halèsia*, the *Nýssa*, the fig, and several others, though they stand out, require protection.

The modern collections of trees in Germany are too numerous to be mentioned in this work, and we can only, therefore, give the names of those which we have been informed are the most complete.

In Austria there is an excellent collection in the University Botanic Garden of Vienna, under the care of Baron Jacquin, chiefly planted within the last twenty years, but a part much older. Here the macluras, male and female, stand in the open air, but require protection during winter. The salisburia here, and those in several other places near Vienna, flower annually; but they are all male plants. On that in the Botanic Garden, Baron Jacquin has had the female grafted with scions from Geneva, but they have not yet flowered. Laxenburg is more remarkable for native trees than for foreign ones; but there are some very large tulip trees, which ripen their seeds every year, and some very large purple beeches and weeping willows. There is there an *Araucaria excelsa*, protected during winter by a wooden house, which has attained the height of 30 ft. in six or seven years. Baron Jacquin assures us that this is one of the finest and most picturesque specimens of this tree that can be imagined. In the park there are many fine oaks of the growth of several centuries; and a very comprehensive general collection of trees and shrubs, of from ten to forty years' growth. All these have been planted by, and are now under the care of, M. Stephen Rauch, through the exertions of whose son, M. Charles Rauch, now head gardener at Rennweg, we have received much of the information contained in this section; while another son, M. Francis Rauch, now (1835) in London, has drawn from nature the greater part of the botanical specimens by which this work is illustrated. In Austria Proper there are collections at Bruck on the Leytha, on the borders of Hungary; at Dornbach, Prince Schwartzenberg; at Hadersdorf, Baron Loudon (a view of whose mansion we have given in the *Encyclopædia of Gardening*, edit. 1835, p. 136. fig. 87.); and at the nurseries of Rosenthal, and Held in Vienna. On Kopenzel Berg, a hill in the neighbourhood of Vienna, from which there is one of the finest views in Europe, there is a tulip tree 60 years planted, which has an immense globular spreading head, though only 45 ft. high. At Bruck, the trees have been chiefly planted within the last eight or ten years; but there are some older specimens well deserving notice; such as *Acer striatum*, 18 ft. high in 20 years; *Paliurus anstrælis*, 18 ft. high in 30 years; and *Fraxinus atrovirens*, 18 ft. high in 20 years. At Dornbach there is a good collection; but very few trees that have been above 40 years planted. At Hadersdorf we observed, in 1814, some fine cypress trees planted round the tomb of the great Marshal Loudon, but in the Return Paper received they are not mentioned; there is, however, a good collection, the most rapidly growing tree of which appears to be that beautiful species of elm, *Ulmus ciliata*, which, in 20 years, has attained the height of 36 ft. in poor sandy soil. The Vienna nurseries, especially that of Rosenthal, contain good collections planted within the last 20 years.

There are collections at Eisenstadt, and other residences, in Hungary, and in the botanic garden at Pesth. In Bohemia there are collections at Toeplitz, Schönhoff, and other places.

The following enumeration of the evergreen trees and shrubs, foreign and indigenous, which stand the winter at Vienna without protection has been furnished to us by Mr. Francis Rauch:—

Hypericææ. *Hypericum calycinum.*

Hicææ. *Ilex Aquifolium.*

Leguminosææ. *Spârtium juncæum, Cytisus scopârius.*

Pomacææ. *Cratægus Pyracantha var. fructu luteo.*

Araliæææ. *Hedera Helix* and varieties.

Caprifoliæææ. *Caprifolium sempervirens.*

Ericæææ. *Calluna vulgaris.*

Thymelæææ. *Dâphne Lauréola.*

Euphorbiæææ. *Buxus sempervirens* and varieties.

Coniferaææ. *Pinus Banksiana, Cembra, inops, pumilio, Strôbus, rigida,*

sylvéstris, *Tæ'da*, *variábilis*; *A'bies álba*, *balsamífera*, *Píceá*, *canadáénsis*; *Lárix europæa*, *microcárpa*, *péndula*; *Cuprèssus thyóides*; *Thùja austrális*, *cupressóides*, *occidentális*, *orientális*; *Juníperus commúnis*, *Oxýcedrus*, *Sabína* and var., *virginiána*; *Táxus baccàta* and var.

Smilácea. *Rúscus aculeátus*, *andrógynus*.

In Prussia the botanic garden at Berlin contains a very full collection, all planted within the last 20 years, and of which an enumeration, kindly sent us by M. Otto, will be found in the *Gardener's Magazine*, vol. xi. p. 541. In this garden *Magnòlia acuminàta* is from 20 ft. to 30 ft. high; and several species or varieties of American ash trees, such as *F. amer. expànsa*, *F. amer. epíptera*, *F. amer. juglandifòlia*, and several American oaks, are from 20 to 30 years old, and from 25 ft. to 30 ft. high. At Sans Souci there is a collection which has been planted from 10 to 50 years, and in which the tulip tree and the horse-chestnut, in 45 years, have attained the height of 50 ft.; the *Magnòlia acuminàta*, 12 years planted, is only of the height of 6 ft.; *A'cer rubrum*, in 45 years, has attained the height of 38 ft.; and *Ailántus glandulosá*, in 30 years, that of 20 ft. At the Pfauen Insel there is a good collection, from 40 to 50 years planted, among which we observe *Magnòlia acuminàta*, 8 years planted, 18 ft. high; *A'cer eriocárpum*, 40 years planted, 50 ft. high; *Negúndo fraxinifòlium*, 40 years planted, 40 ft. high; *Sophòra japoníca*, 9 years planted, 12 ft. high; and *Plátanus orientális*, 42 years planted, and 55 ft. high. The soil of these three gardens is a deep sand. Prince Püekler Muskau has a collection at Muskau in Silesia, about twenty miles from Dresden; and, according to M. Hofman (*Gard. Mag.*, vol. xii.), it contains some fine tulip trees, and beeches. The public promenades and squares at Breslau are planted with trees, which are placed at a sufficient distance to allow them to attain their full size.

In Bavaria there is an excellent collection in the botanic garden at Munich, and also in the royal gardens at Nymphenburg, and in the royal nurseries. Considering the elevated situation, unfavourable climate, and very indifferent soil, the gardening exertions made at Munich, and the success attending them, surpass those of any other government of Germany. Much of this success is owing to the skill, industry, and enthusiasm, of the late and present garden directors, Charles Skell, and Charles Louis Skell. There is an excellent collection of trees and shrubs around the old castle of Heidelberg, and some specimens of great antiquity there have been already mentioned. (p. 147.)

In Saxony there is a collection in the botanic garden at Dresden, planted since 1815. There is here, in the royal gardens, the largest standard fig tree in Germany; it is 60 ft. high, with a trunk 18 in. in diameter at one foot from the ground. Every year it bears some thousands of figs; but it requires protection by a boarded house during winter. In the royal gardens at Pillnitz are the largest and oldest camellias in Germany; they form bushes about 20 ft. high, the stems 4 in. or 5 in. thick; and they are protected in winter by a wooden house, in the roof of which are small windows. In the garden of Lieutenant Weber, at Dresden, there is an excellent collection of foreign hardy shrubs, as well as some enormously large fig trees, which are known to be above 200 years old. The beautiful road from Worlitz to Dresden is bordered by magnificent oaks, only equalled in Germany by those of the finest parts of the Black Forest.

In Hanover, at Göttingen, there is an excellent collection under the care of our esteemed friend and correspondent M. Fischer, one of the most active and zealous garden directors in Germany; there are, also, the collections at Schwöbber, and the other places already noticed. At Herrenhausen is a rich collection of trees and shrubs, planted in 1834 by M. Wendland. At Haroke, near Helmstadt, there is a very interesting garden laid out in different scenes, which are called Canada, Virginia, &c., from the native countries of the trees planted in them; thus forming a kind of geographical garden. (See (*Gard. Mag.*, vol. xi. p. 647.) Among these trees are a very large *salisburia*, and many large *liriodendrons*. One part of the ground is laid out and

planted with *Pinus Cembra*, so distributed as to resemble a native forest. Baron Hake, at Ohr, near Hamelen, on the river Weser, has formed part of a forest on a hill (Ohr berg) into a park, in which some American trees grow perfectly well: the soil is a sandy loam. There are, also, many large American trees in the garden of Baron Steinberg, at Bruggen, on the road from Hannover to Ernbeck. At Eldagsen, a small village about four miles from Hannover, there are many large trees, which were planted about the same time as those at Schwöbber.

In Wurtemberg there are good collections at the Palace of Rosenstein, and in the royal nurseries, but scarcely any evergreens. The Palace of Solitude (at one time so celebrated for its gardens) is surrounded by a natural forest, which extends many miles in every direction; the trees are chiefly beech, oak, and the trembling poplar, some of which have attained a great size. (See *Encyc. of Gard.*, edit. 1835, p. 158. to p. 169.)

In Baden, at Carlsruhe, there was a very good collection under the care of the late excellent M. Hartweg, author of *Hortus Carlsruhanus*; this collection is still in existence under the care of M. Held, and has since received occasional additions. There is also a very good collection in the celebrated garden of Schwezingen; which, with all the most remarkable gardens of Germany, will be found described at length in the last edition (1835) of our *Encyclopædia of Gardening*. At Donaueschingen (the source of the Danube) there is a tolerable collection, and some specimens of abies, populus, and liriiodendron, of considerable size. In the Black Forest, which surrounds this place, are the largest oaks and silver firs in Germany. In 1828, we spent an entire day examining and admiring these noble trees, many of which we estimated at upwards of 100 ft. in height. All the trees and shrubs enumerated above as enduring the open air at Vienna, without protection during winter, do so at Carlsruhe; with the following additional species, which have been pointed out to us by M. Hartweg, son of the late director of the grand-ducal gardens; a highly educated young gardener, now in the employment of the London Horticultural Society.

Leguminosæ. *Ulex europæa*, nana.

Tamariscinæ. *Tamarix gallica*, germanica.

Araliæcæ. *Hedera canariensis*.

Ericæcæ. *Daboecia* (*Menziëria*) *poliifolia*; *Andrömeda axillaris*, *Catesbæi*, speciosa var. *pulverulenta*; *Gaulthèria procumbens*; *Kalmia angustifolia*, glauca, latifolia; *Rhododendron catawbiense*, caucasicum, maximum, ponticum and varieties, ferrugineum; *Itea virginica*.

Bignoniæcæ. *Bignonia capreolata*.

Myricæcæ. *Myrica cerifera*.

Conifera. *Cedrus Libani*.

Empétræcæ. *Empetrum nigrum*, *Corèma album*.

In Hesse Cassel, the garden of Wilhelmshoe, at Cassel, contains a good collection, chiefly planted within the last 20 years; but some of them have been planted 60 years. From the particulars with which we have been obliged by the director of the garden, M. Claus, we find that the tulip tree here, 60 years planted, has only attained the height of 20 ft.; and the ailantus, 60 years planted, is under 30 ft. The only evergreens of which we have had returns are, the common pines and firs, and *Cupressus thyoides*.

In Nassau, in the Grand-Ducal Botanic Garden at Biebrich, there is a good collection, distributed thinly all round the margin of the garden; and, this garden being of very great length in proportion to its breadth, the space afforded to each tree is such as will enable it to attain a very considerable size. A catalogue of this garden, accompanied by a plan, was published in 1831. We may remark here that the names in this catalogue, as in those of most German catalogues of modern date, generally correspond with the names in the catalogue of Messrs. Loddiges; the reason is, that the collections which have been formed in Germany, during the last 50 years, have, for the

most part, been either procured direct from Hackney, or from German nurserymen who have purchased their foreign trees and shrubs there.

In the dukedom of Mecklenburg, Baron Laffert has a very rich collection of trees and shrubs. Some magnolias grow here in the open air without any kind of protection.

In Anhalt there are few collections besides that of Wörlitz, already noticed (p. 149.); but the cemetery at Dessau, one of the finest in Germany, contains a number of good foreign trees.

In the free townships there is, at Frankfort, a considerable collection in the public garden formed on the ramparts from the plan of M. Zeyer, and planted by M. Rinz. (See *Encyc. of Gard.*, edit. 1835, p. 195.) There are, also, collections of trees, more or less extensive, in the public gardens belonging to the other free towns. On the ramparts of Bremen there is an excellent collection of poplars, of all the different species and varieties that will endure the open air in that part of Germany, of the height of from 60 ft. to 80 ft. This collection was made with great care by the late Professor Mertens.

The principal nurserymen in Germany are, Messrs. Booth of Hamburg, who have an excellent collection of trees in their grounds at Floetbeck; M. Hayen, at Erfurt; M. Schelhaus, at Cassel; M. Scidel, at Dresden; M. Rosenthal, and M. Held, at Vienna; M. Mathieu, at Berlin; and M. Rinz, at Frankfort.

In Germany, as in France, there are very few evergreen trees and shrubs, either in the indigenous or introduced flora, as compared with the ligneous flora of Britain. The cedar of Lebanon requires protection all over Germany, except in the warmest parts of Hungary and Baden; the common laurel, the Portugal laurel, the arbutus, the rhododendron, the kalmia, the laurustinus, the furze, and even the Irish ivy, can only endure the winters in Germany in very favourable situations.

SECT. IV. *Of the Indigenous and Foreign Trees and Shrubs of Scandinavia, including Denmark, Holstein, Sweden, Lapland, Finland, Iceland, Greenland, and the Faroe Islands.*

FROM the northern latitude and severe climate of these countries, it will not be expected that their ligneous flora, either indigenous or introduced, can at all equal that of Britain. The total number of ligneous species enumerated by Retzius, in his *Flora Scandinavia*, edit. 1795, amounts to 133 species, and of that number there appear to be a few, enumerated below, which are not natives of Britain. The most important of these are the *Acer platanoides*, or Norway maple, and the common spruce fir.

Rosaceæ. *Rosa Eglantaria*, cinnamomea, fluviàlis *Retzius*; *Potentilla fruticosa*; *Spiræa salicifolia*.

Pomaceæ. *Crataegus monogyna* *Jacq.*

Leguminosæ. *Genista germanica*, *Coronilla Emericus*.

Salicinceæ. *Salix hermaphrodita*, hastata, myrtilloides, depréssa, lappònum.

Acerinceæ. *Acer platanoides*.

Cistinceæ. *Helianthemum œlândicum*, *Fumana*.

Tamariscinceæ. *Tamarix germanica*.

Ericaceæ. *Phyllodoce taxifolia* (*Menziesia cærulea*); *Andrómèda tetragona*, *hypnoides*, calyculata; *Rhododéndron lappònicum*; *Lèdum grœnlândicum*, not found in the limits of Sweden.

Caprifoliaceæ. *Lonicera Xylósteum*, cærulea; *Linnaea boreàlis*.

Conifereæ. *Abies excelsa*; *Juniperus communis minor*, communis arborescens.

On looking at the *Flora Danica*, *Flora Suecica*, and *Flora Lapponica*, we find the number of ligneous species gradually diminish as we advance northwards, till, in the Faroe Islands, a flora of which has been given by W. C.

Trevelyan, Esq., the total number of ligneous plants consists only of fifteen species, which are all under the height of 3 ft. They are as follow :—

Rosæcæ. Rôsa.

Empétrecæ. *Empetrum nigrum.*

Ericæcæ. *Erica cinerea*, *Calluna vulgaris*, *Chamælèdon procumbens.*

Vacciniæcæ. *Vaccinium Vitis-Idææ*, *uliginosum*, and *Myrtillus.*

Salicincæ. *Salix caprea*, *phylicæfolia*, *hastata*, *lanata*, *arctica*, and *herbæcea.*

Coniferæ. *Juniperus communis.*

The Faroe Islands, Mr. Trevelyan, who resided on them for some years, informs us, are twenty-two in number, and are situated between $61^{\circ} 26'$ and $62^{\circ} 25'$ N. lat., and $6^{\circ} 17'$ and $7^{\circ} 43'$ W. long. Only seventeen of them are inhabited. Most of them may be compared to the summits of mountain ridges, rising out of the ocean to the height of nearly 3000 ft. There is usually deep water close to the land, which often rises in perpendicular cliffs to a height of 1200 ft. and 1500 ft., and, in one instance, to above 2000 ft. The climate is generally mild, but damp. It is not subject to extremes of temperature; the mean of mild years being 49° , and of cool years 42° . The highest temperature during four years was 72° , and the lowest 18° . The only corn cultivated is the Scotch bigg, and that does not always ripen. In the peat bogs occur the remains of birch trees; but these do not now grow in the islands, having probably been extirpated by being used as fuel.

A general view of the arboricultural flora of Sweden, considered geographically and geologically, has been prepared for us by the celebrated botanist Dr. Agardh, formerly professor of botany at Lund; and we have received another for the whole Scandinavian peninsula, by Professor Schouw of Copenhagen; but, as these communications, though excellent in themselves, are somewhat too long for insertion in this work, we have transferred them to the pages of the twelfth volume of the *Gardener's Magazine*; contenting ourselves here with some abridged extracts from them, relative to the introduction of foreign trees into the Scandinavian peninsula.

Foreign trees and shrubs have been introduced into Denmark and Sweden, chiefly in the different botanic gardens, and in the grounds of the royal residences at Copenhagen and Stockholm, and of the wealthy proprietors in the neighbourhood of these capitals, and of the other large towns. There are, however, but few American trees or shrubs to be found as standards in the neighbourhood of either capital. The largest indigenous trees in Denmark are beeches, of which one, in the park of Jägersborg, exceeds 100 ft. in height. The white poplar also grows to the height of 100 ft., and the oak and Scotch pine attain a great size. At Dronninggaard, near Copenhagen, the tulip tree, in 40 years, has attained the height of 80 ft., and also the horse-chestnut. The *Robinia Pseud-Acacia*, at the same place, has, in 40 years, attained the height of 60 ft.; but the *Gleditschia triacanthos*, in the same period, only 16 ft. *Ornus europæa* is 30 ft. high; *Abies Picea*, the silver fir, 100 ft.; and *Abies canadensis*, only 6 ft.; while *Pinus Strôbus* becomes a considerable tree. At the royal gardens of Rosenberg, near Copenhagen, there is an excellent collection, planted for the most part in 1831, 1832, and 1833, a list of which, with their dimensions, has been kindly sent us by the royal gardener there, M. Jens P. Petersen. On looking it over we find that it contains nearly all the species procurable in the London nurseries. Among the hardy trees, however, the cedar of Lebanon is not included, nor the common laurel. In the garden of Christianhome, near Lolland, there are some good trees, the dimensions of which have been sent us by the curator, M. Gentz. Here the tulip tree and the *Acer dasycarpum* are 40 ft. high; the robinia, 36 ft.; the gleditschia, 24 ft.; the *Ornus europæa*, 24 ft.; the walnut, 30 ft.; the Lombardy poplar, 80 ft.; the purple beech, 30 ft.; the platanus, 80 ft.; and the larch, 75 ft.

In Sweden, according to Dr. Agardh, "the central points from which foreign trees and shrubs have spread over the whole country are, Lund, Upsal, Stockholm, and Gottenburg. Some of these introduced trees, such as *Larix*

europæa, *Æsculus Hippocástanum*, some species of *Pópulus*, and *Æcer Pseùdo-Plátanus*, thrive here as well, and are almost as common, as the indigenous trees. Of fruit trees, all that are cultivated north of the European alps grow in Scania; such as peaches, apricots, grapes, almonds, chestnuts, walnuts, and mulberries (*Morus álba* and *nigra*), and they appear to suffer very little from the cold: even figs (*Ficus Cárica*) have lived through some winters. The Japanese shrubs endure the climate of Lund tolerably well, as *Kérria japónica*, and *Broussonétia*, which last had grown to the size of a large tree, one third of a foot in diameter, in the botanic garden at Lund, till accidentally (and not, as it seemed, by the severity of the winter) it died off. But very few evergreens endure our winters; not even the *Aícuba japónica*, or the Portugal or the common laurel; and the holly with great difficulty. The few exotic evergreens that we do possess are, *Búxus sempervirens* and var., *Crataëgus Pyracántha*, *Vínca* sp., and the *Coniferæ*.

"Many of the Swedish noblemen have contributed much to the spreading of foreign trees throughout Scandinavia, by planting them on their estates; as, for example, His Excellency Count Trolle Wachtmeister, His Excellency the Count de la Gardie, Baron Gyllenkrook, and several more, in Scania; also, the late M. Thouse, in West Gotha; His Excellency Count Trolle Bronde, in Upland; M. Wares, in Warmeland; &c. The *Morus álba* thrives well, even as far as Upsal; and, under the protection of our adored Crown Princess Josephine, there is a large plantation of it at Stockholm, for the purpose of breeding and feeding silkworms; and the silk obtained from them is not only abundant in quantity, but the quality of it is excellent. At Stockholm there are several patrons of arboriculture, as regards the cultivation of foreign trees. Some of the most distinguished are, the Counsellor de Pontin, M. Siefwerstrale, and M. Rofenblad; the latter of whom has the richest collection of plants that can be found in any private garden in Scandinavia. There are two public plantations of foreign trees at Stockholm; viz. that of the Forest Institute, directed by M. Ström, and that of the Agricultural Academy; both of which possess a great number of foreign trees.

"As to the height of the trees, I can find no difference between those in Scandinavia and those in Germany, or in any other country north of the European alps. The beeches and oaks are as well grown trees with us as they are in Germany. The sweet chestnut tree and the *Robínia Pseùdo-Acácia* are somewhat smaller, as they have hitherto never attained a greater height here than 50 ft.; but others, as the *Æsculus*, the foreign *tilias*, *populus*, the foreign pines, *juglans*, &c., may be compared with those of Germany. The *Plátanus occidentális* attains a height of 30 ft. The *Plátanus orientális* does not stand in the free ground in our garden. The tulip tree is perfectly hardy. We have not yet tried the cedar of Lebanon in the open air; but we hope to be able to do this at some future time. — *C. Agardh. Lund, Sept. 23. 1835.*"

SECT. V. *Of the Indigenous and Foreign Trees and Shrubs of the Russian Empire.*

THIS immense country, extending in latitude from the Crimea to the Gulf of Bothnia, and in longitude stretching far into Asia, exhibits less variety of surface than might be expected from its great extent. With the exception of its southern and Asiatic provinces, its ligneous flora differs little from that of Germany and of the north of France; but the Crimea, the mountains of Caucasus, the Circassian alps, and the shores of the Caspian and Black Sea, are rich in the productions of warmer climates, and include as indigenous many of the more important trees and shrubs of Switzerland, Italy, and Greece, besides a great number peculiar to themselves. On this account, though the Asiatic portion of the Russian flora has been very imperfectly explored, the number of species that Russia possesses that are not indigenous in Britain is con-

siderable, as appears from the following enumeration, taken from Pallas's *Flora Rossica*, published in 1788.

- Ranunculaceæ.* *Atragène austriaca*, ochoténsis.
Laurinæ. *Laurus nobilis*.
Berberidææ. *Berberis sibirica*.
Philadelphicæ. *Philadelphus coronarius*.
Grossulacææ. *Ribes americanum*, procumbens, *Diacantha*.
Granulææ. *Punica Granatum*.
Elæagnææ. *Elæagnus angustifolia*, orientâlis.
Thymelæææ. *Dôphne alpina L.* (altâica *Pallas*), caucásica, pântica.
Rosacææ. *Rosa alpina*, davûrica, caucásica, parvifolia; *Spiræa chamædri-folia*, *betulifolia*, *trilobata*, *thalictroides*, *crenata*, *alpina*, *salicifolia*, *altâica*, *sorbifolia*.
Pomacææ. *Pyrus salicifolia*, præcox, baccata; *Cratægnus sanguinea*, monogyna rubra, nigra, *Azarolus*, *Pyracantha*; *Méspilus germanica*.
Amygdalææ. *Amygdalus nana*, communis; *Pérsica vulgaris*; *Armeniaca vulgaris*, sibirica; *Cerasus Chamæcerasus*, prostrata, *Mahaleb*, *Laurocerasus*; *Prunus caucásica*.
Leguminosææ. *Cytisus austriacus*, hirsutus; *Halimodéndron argenteum*; *Caragána Altuganu*, frutescens, spinosa, pygmaea; *Colútea cruenta*, *Calôphaca wolgárica*.
Urticææ. *Ficus Carica*.
Ulmacææ. *Ulmus lævis*, pumila; *Celtis australis*.
Cupuliferææ. *Quercus Cerris*.
Betulnææ. *Bétula davûrica*, fruticosa; *Alnus incana*.
Salicinææ. *Salix caspica*, monandra, Gmelini, serotina, sibirica, myrtilloides, arbutifolia, divaricata, lappoonum, lamuginosa, hastata, *rhamnifolia*, *berberifolia*, retusa, ártica; *Pópulus balsamifera*.
Platanææ. *Platanus orientâlis*.
Juglandææ. *Juglans regia*.
Euphorbiacææ. *Buxus sempervirens*.
Rhamnææ. *Rhamnus alpinus*, diûricus, carpinifolius, *Erythrôxyton*, *Erythrôxyton* var. β , ? angustissimus *Dec.*; *Zizyphus vulgaris*, *Paliurus aculeatus*.
Staphyleacææ. *Staphylea pinnata*.
Acerinææ. *Acer platanoides*, Pseudo-*Platanus*, tatáricum.
Vitææ. *Vitis vinifera*.
Anacardiæææ. *Pistacia Terbinthus*, *Rhus Coriaria*, *Cótinus*.
Tamariscinææ. *Tamarix Pallâsii*, germanica.
Nitrariæææ. *Nitraria Schôberi*, sibirica.
Chenopodiæææ. *Salicornia*, 4 species; *Anábasis tatárica*; *Salsola*, 7 species; *Suaeda microphylla*.
Ericacæææ. *Lêdum palustre*; *Andrômeda calyculata*, *lycopodioides*, *hypnoides*, *ericoides*, *tetragona*, *Bryanthus*, *Stelleriana*; *Phyllodoce carulea*; *Azalea pântica*; *Rhododéndron lapponicum*, pânticum, *chrysanthum*, caucásicum, diûricum, camtschaticum.
Vacciniæææ. *Vaccinium Aretostáphylos*.
Caprifoliacæææ. *Lonicera tatárica*, *Nylósteum*, alpigena, caucásica, altâica; *Fibérnum diûricum*, orientâle; *Sambucus racemosa*.
Cornæææ. *Cornus mas*, álba.
Asclepiadæææ. *Periploca græca*.
Ebenææææ. *Diospýros Lotus*.
Jasminæææ. *Jasminum fruticans*, officinále.
Oleacæææ. *Olea europæa*.
Solanææææ. *Lycium tatáricum*.
Verbenacæææ. *Vitex Agnus-cástus*.
Coniferæææ. *Pinus Cembra*; *Abies excelsa*, *Picea*; *Lárix europæa*, *Cuprésus sempervirens*; *Juniperus davûrica*, *lycia*, *Sabina*, *phœnicea*; *Ephedra polygonoides*.
Smilacæææ. *Ruscus hypophyllum*.
Corymbiferæææ. *Pallâsia Pterocæceus L.*

The trees and shrubs which are to be found in the neighbourhood of St. Petersburg, and on the shores of the Gulf of Finland, are as follows:—*Pinus sylvestris*, *Abies excelsa*, *Bétula álba*, *Álnus glutinosa*, *Pópulus trémula*, different *Sálices*, *Juníperus communis*, *Sórbus aucupària*, *Cérasus Pàdus*, *Rhámnus Frángula*, *Tilia europæa*, *Pýrus Málus*, *Acer campéstre*.

The introduction of foreign trees and shrubs into Russia may date from the commencement of the reign of Catharine, or about the year 1768; when, from reading *Der Hausvater*, that empress determined on having the gardens at Tzarsco Celo laid out in the English manner. From the severity of the climate, not many foreign species can endure the winters, either there or any where else, in the neighbourhood of Petersburg; nevertheless, with laudable ambition, many species have been tried at all the imperial residences. The trees and shrubs generally used for planting the Petersburg gardens are of the following genera:—

<p><i>Aquifoliæcæ.</i> Ilex. <i>Leguminosæ.</i> Cytisus, <i>Cara-gána</i>, Genista, Spártium. <i>Rosæcæ.</i> Rôsa. § <i>Potentillæcæ.</i> Potentilla. § <i>Spirææcæ.</i> Spiræa. <i>Amygdalæcæ.</i> Amýgdalus.</p>	<p><i>Pomâcæcæ.</i> Cratægus, Mésplus, Pýrus. <i>Araliæcæcæ.</i> Hédéra. <i>Caprifoliæcæcæ.</i> Sambucus, Lonicera, Viburnum. <i>Córncæcæ.</i> Córnu. <i>Oleæcæcæ.</i> Syringa.</p>	<p><i>Solanæcæ.</i> Lycium. <i>Elæagnæcæ.</i> Hippophæe. <i>Euphorbiæcæcæ.</i> Buxus. <i>Cupuliferæcæ.</i> Fâgus. <i>Amentâcæcæ.</i> Córylus. <i>Taracæcæ.</i> Tâxus. <i>Coniferæcæ.</i> Pinus.]</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

The *Pinus sylvestris* and the *Abies excelsa* attain a considerable size in the elevated light soil at a few miles' distance from St. Petersburg; though in the bog by which that city is surrounded their size is but small. In the Taurida Palace gardens, in the city, the *Ulnus campéstris* has attained the height of 49 ft.; the *Cérasus Pàdus*, and the *Sórbus aucupària*, of 21 ft.; the *Fráxinus excelsior*, of 35 ft.; the *Sàlix fragilis*, 49 ft.; the *Álnus glutinosa*, 56 ft.; the *Bétula álba*, 68 ft.; the *Làrix europæa*, 63 ft.; and, what appears to us remarkable, the *Quércus rubra*, also 63 ft. In the summer gardens of St. Petersburg, planted during the reign of Peter the Great, there is a lime tree 79 ft. high; and a common elm and Norway maple, each 70 ft. high.

On the shore of the Gulf of Finland, opposite the village of Strelna, the small island of Sosnovy Rosha is entirely occupied with tall Scotch pines, from 3 ft. to 5 ft. apart; among which, one has attained the height of 77 ft., and another of 65 ft. In the imperial garden at Strelna is a common English elm, 60 ft. high, the branches of which cover a space of 56 ft. The measurements of a number of trees, grown on the estate of Madame Constantinoff, at Rudets, near St. Petersburg, have been sent us by one of the imperial gardeners, with the following introductory remarks:—"The woods consist principally of pines and firs; the surface of the ground is covered with long moss (*Hýpnum*); the surface stratum is black earth, 6 in. deep; below this a stratum, 4 in. deep, of sand mixed with earth; and under this is clay. The greater number of the trees consists of *Pinus sylvestris*, *Abies excelsa*, and *Bétula*. The pines grow with clean straight stems, of from 50 ft. to 70 ft. high, to where the branches commence, which extend from 14 ft. to 20 ft. more, making the whole height of the tree nearly 100 ft. No care is taken of the woods; the young trees spring from self-sown seed; and the strong plants are suffered to overshadow and destroy the weak ones, till the former at last become large trees. Where the *Pinus sylvestris* grows singly, and has plenty of space to spread its branches, the lower arms are not rubbed off or killed when young, as they are where the tree grows in a thick wood, but they form immense limbs; the consequence is, that the trunk of the tree becomes full of large knots, and, though it is more in diameter, it is less in height; the timber, of course, being of little use but as fuel." Some of the largest specimens of *Pinus sylvestris* in these woods, supposed of 213 years' growth, measured in height 99 ft. and 85 ft.; others, 65 years old, 78 ft. and 64 ft.; one, supposed to be 108 years old, 106 ft.; one, 95 years old, 85 ft.; 120 years old, 99 ft.; 232 years old, 113 ft.; this last tree had a trunk 31 in. in diameter at 1 ft. from the ground; the diameters of the others varied from 26 in. to 12 in., 8 in., and even 7 in. *Bétula álba*, in the same wood, at 40 years old, was 71 ft. high; at 85 years, 85 ft.; and at 75 years, 70 ft. and

64 ft. The *Alnus glutinosa*, at 44 years, was 67 ft. in height. A *Pópulus trémula*, of 78 years' growth, was 74 ft.; and one of 90 years' growth was 71 ft.; the diameter of the trunk being in the latter case 14 in., and in the former 12 in. It may be observed of all these trees, that they have stood very close together, so that the diameter of the trunk is generally very small in comparison with its height.

The following trees and shrubs are found to stand the open air in the neighbourhood of Moscow. By comparing this list with that given above of the trees and shrubs which will stand the open air in the neighbourhood of St. Petersburg, the reader will be able to ascertain what are the very hardiest trees and shrubs of temperate climates, and of high latitudes.

Tilia. *Tilia europæa*. "*Tilia europæa* probably means *T. parvifolia*, which appears the most northern species or variety of *Tilia*. The *Flora Mosquensis* gives *T. parvifolia* 'in sylvis, nemoribusque'; *T. grandifolia*, only 'ad pagos, in hortis'; while *T. europæa* is not named at all. — H. C. Watson. Feb. 6. 1836." *Acer*. *Acer Pseudo-Platanus*. *Hippocastanea*. *Æsculus Hippocastanum*. *Celastrineæ*. *Euonymus europæus* and *verucosus*. *Rhamnææ*. *Rhámnus cathárticus* and *Frángula*. *Leguminosæ*. *Cytisus Labúrnum*, capitatus, ruthénicus, sessilifolius, nigricans; *Caragána frutescens*, acutifolia [?], obtusifolia [?], spinosa. *Amygdalææ*. *Amýgdalus nána*; *Prúnus doméstica*, spinosa; *Cerasus durácina*, ávium, *Pádus*. *Pomacææ*. *Sórbus aucupária*, doméstica, *Aria*; *Cratægus Oxyacántlia*, tomentosa, grandiflora lucida, coccinea ovalifolia, monógyna;

Cotoneáster vulgáris; *Pýrus melanocápa*, communis, *Malus, baccáta*, *prunifolia*, *ovális*. *Caprifoliacææ*. *Córnuş álba*, *sanguinea*, *sericea*; *Sambucus racemosa*, *nigra*; *Fibúrnum O pulus*, *rosea*, *Lantána*. *Oleacææ*. *Fráxinus excelsior*, *álba*, *tomentosa*. *Elæagnææ*. *Hippóphac Rhannióides*, *Elæágnus songárica*. *Ulmæææ*. *Ulmus effúsa*, *campéstris*, *sativa*. *Salicæææ*. *Salix babylónica*, and almost all the other species; all the species of poplar except *dilatáta*. *Betulínæææ*. *Alnus incána*, *glutinosa*; *Bétula álba*, *nána*, *nigra*. *Cupuliferaææ*. *Corylus Avellána*, *cornúta*; *Quercus Iðbur*. "Both the British oaks (*Q. Iðbur* and *sessiliflora*) are included in the *Flora Mosquensis*. — H. C. Watson. Feb. 6. 1836." *Coniferæææ*. *Juniperus Sábina*, *Thuja occidentális*; *Pinus sylvestris*, *Cembra*, *Stróbus*, *montána*; *Abies excelsa*, *álba*, *Pichta*, *canadénsis*, *nigra*, *rúbra*, *balsamifera*; *Lárix communis*.

Deleuze mentions Dimidow as having the richest botanic garden in Russia, and as having sent to the Paris garden many fine trees and shrubs, natives of Siberia. Among these were, *Caragána Allagána*, *C. pygmæa*, and *Halimodéndron argéntum*. The catalogue of Dimidow's collection was published in 1786, soon after which the proprietor died, and the collection was dispersed. The richest collection in 1814 was that at Gorinkí, which suffered greatly when the French were in possession of that part of the country, and has since been destroyed or neglected. The gardens in the neighbourhood of Moscow, though they do not contain a great number of ligneous species, are not without some of very considerable size; more especially the common or Scotch pine, the birch, the white poplar, the ash, the common elm, and the white willow. This last tree, in the south of Russia, is planted in straight lines of apparently interminable lengths, to indicate the road across those immense steppes, over which a traveller may proceed a whole day without seeing any other trace of civilisation than these trees and the post stations. At least we found this to be the case in 1814. M. Fintelman, one of the imperial gardeners at Moscow, visited Britain during the summer of 1835, with a view to add to the collection under his care; and he informed us that the proprietors of gardens in his neighbourhood are most assiduous in the improvement of their grounds, and are most anxious to plant in them every tree and shrub that they think at all likely to endure the climate.

Warsaw, being three degrees further south than Moscow, enjoys a climate better adapted for the introduction of foreign trees and shrubs; and a considerable collection was introduced into the botanic garden there soon after the general peace. On looking over the catalogue of this garden, published by M. Schubert in 1824, we find that the greater number of trees and shrubs which stand the open air in Berlin also do so at Warsaw; but that the Portugal laurel, the common laurel, the laurustinus, the rhododendron, the tree box, the furze, and the broom, are not hardy; and that the ivy and the common holly require protection during severe winters. *Magnolia tripétala* and *acuminata* stand against a wall, with protection, as does also the tulip tree. *Cratægus punctata*, *orientális*, *Crus-gálii*, *pyrifolia*, *coccinea*, *cordáta*, *glandu-*

lōsa, nīgra, tanacetifōlia, *Oxyacantha*, melanocārpa, and prunifōlia, are all tolerably hardy.

Cracow is upwards of two degrees farther south than Warsaw; and it is not much higher above the level of the sea, since the Vistula passes through both cities, and there is not the slightest waterfall on that river between Cracow and its mouth at Dantzic. The greatest cold of the ordinary winters at Cracow is from 13° to 16° Réaumur. (from 2° above zero, to 4° below zero, Fahr.); and the snow seldom lasts longer than from two to three weeks. At Niedzwiedz, about three miles from the city, Count Wodzicki, a gentleman who has never been out of Poland, began, in the year 1814, to plant an arboretum; and he has pursued his plan with such energy, that in 1836 his collection amounted to nearly 200 species and varieties, exclusive of half-hardy species, which he keeps in conservatories, or against walls. An account of this arboretum, as it was in the year 1833, was published in the *Annales de Fromont*, tom. v. p. 177.; and a translation of this account, with some corrections and additions sent us by the author, will be found in the *Gardener's Magazine*, vol. xiv., for 1838. Though Count Wodzicki, as he informs us, was upwards of 61 years of age when he commenced his arboretum, and consequently, in 1836, must have been 83 years old, his passion for trees was then by no means diminished. He was in constant correspondence with Messrs. Booth of Hamburg, M. Soulange-Bodin of Paris, and various commercial cultivators and amateurs; and he spares no expense in procuring every new ligneous plant that is likely to stand the open air at Cracow. A list of the species and varieties which were in the arboretum at Niedzwiedz in September, 1836, with their dimensions and the year in which each was planted, has been kindly sent us by the count; from which it appears that the growth of the hardier species, in that climate, is as rapid as it is in the climate of London. *Acer Pseudo-Platanus*, 24 years planted, is 28 ft. high, with a trunk 22 in. in diameter; *A. platanoides*, of the same age, is 30 ft. high, with a trunk 18 in. in diameter; *A. rubrum*, 20 years planted, is 34 ft. high, with a trunk 13 in. in diameter; and *A. eriocarpum*, of the same age, is 36 ft. high, with a trunk 18 in. in diameter. *Abies rubra*, 25 years planted, is 48 ft. high, with a trunk 17 in. in diameter. *Betula alba pendula*, 24 years planted, is 32 ft. high, with a trunk 18 in. in diameter.

In the Crimea, many foreign trees and shrubs have been introduced into the government garden, and into those of Count Woronzow and some other noblemen.

The introduction of these trees and shrubs may be divided into three periods. In the first period, during the reign of Catharine, the olive, the mulberry, the sweet chestnut, the walnut, the fig tree, the *Diospyros Lotus*, the laurel, the arbutus, and others, were planted in the gardens of individuals, in great part through the influence of Professor Pallas. The second period commenced with the year 1811, when the Duke of Richelieu had the government garden laid out at Nikita, and placed it under the direction of Mr. Steven. This garden soon became celebrated for its collection of trees and shrubs. It contains at present, among other trees, some fine specimens of the genus *Citrus* in the open air, which require only slight protection during winter. The *Magnolia grandiflora* also stands in it in the open air, and flowers every year. The third period commences with the settlement of Count Woronzow at Alpuka, in the year 1823, when he brought with him an English gardener, and an extensive collection of trees and shrubs. Alpuka is finely situated on the sea coast, and it may be worth noticing, that it was admired thirty-six years ago by Prince Potemkin, who caused two cypresses to be planted in the very situation where Count Woronzow is now building a house, under the direction of an English architect.

The winter in this part of Russia lasts but three months, so that oranges, and other trees of the East and West Indies, require only to be protected by a slight roof or tent of boards, covered with leaves or straw. The following species grow freely in the open air:—

Ranunculaceæ: *Clématis flórida* fl. pl., *Pædnia Moutan*. *Magnoliaceæ*: *Magnolia grandiflora*. *Winteraceæ*: *Illicium floridanum*. *Pitosporeæ*: *Pittosporum Tobira*. *Línææ*: *Linum trigynum*. *Camelliææ*: *Camellia japonica*; *Théa Bohéa*, *viridis*. *Ruticææ*: *Corræa álba*. *Leguminosææ*: *Edwardsia microphylla*. *Rosacææ*: *Rubus rosa-folius* fl. pl. *Onagroriææ*: *Fúchia coccinea*. *Salicariææ*: *Lagerstrœmia sudica*. *Myrticææ*: *Myrtus communis*, *Eugénia australis*, *Metrosideros lanceolata*. *Passiflorææ*: *Passiflora filamentosa*. *Cactææ*, or *Opuntiacææ*: *Cactus Tuna*. *Caprifoliacææ*: *Lonicera japonica*, *Fibórnium rugosum*. *Oleacææ*: *Olea fragrans*, *Ligústrum lucidum*. *Jasminææ*: *Jasminum revolutum*. *Apocynææ*: *Nerium Oleander*. *Labiatææ*: *Salvia mexicana*. *Verbenacææ*: *Verbena triphylla*. *Laurinææ*: *Laurus satens*. *Aristolochiææ*: *Aristolochia glauca*. *Cupulliferææ*: *Quercus Súber*, *Ballota*. *Myrticææ*: *Myrica quercifolia*. *Coniferææ*: *Araucaria imbricata*, *Cunninghãmia lanceolata*. *Asphodelææ*: *Phórnium tenax*. *Tulipacææ*: *Yucca aloifolia*, *gloriosa*, *glauca*. *Palmææ*: *Chamaërops humilis*, *Phoenix dactylifera*.

SECT. VI. *Of the Indigenous and Forest Trees and Shrubs of Switzerland.*

SWITZERLAND, from its range of latitude, its lofty mountains, and its hills of every degree of altitude, of every variety of form, and of many different kinds of geological structure, is by far the most interesting country in Europe for the botanist. Here, in consequence of the elevation, we have the plants of the arctic circle on the one hand; while the latitude of the southern extremity, its low level, and proximity to the sea, joined to complete shelter from the north-east and west, are quite suitable for the plants of Italy and Greece. The indigenous ligneous flora of Switzerland, therefore, contains many species not indigenous to Britain. The following enumeration is taken from Suter's *Flora Helvetica*, edit. 1822; and Gaudin's *Flora Helvetica*, 7 vols., recently completed:—

- Ranunculacææ*. *Clématis Flámmula*, *Atrágene austriaca*.
 * *Capparidææ*. * *Cápparis spinosa*. * *Laurinææ*. *Laurus nóbilis*.
Hypericinéææ. *Hypéricum Coris*, *Rícheri*.
Philadelphicææ. *Philadélphus coronarius*.
Grossulacææ. *Ribes reclinatum*. * *Cactææ*. *Opúntia vulgaris*.
 * *Granatææ*. *Púnica Granátum*. *Thymelæææ*. *Dáphne alpina*, *Cneórum*.
Rosacææ. *Rosa Eglantéria*, *cinnamómea*, *montána*, *provinciális gállica*, *rugosa glutinosa*, *pùmila*, *pyrenáica*, *alpina*, *dumetórum*, *rubrifolia*, *spinulifolia*, *collina*, *álba*; *Rubus tomentosus*, *glandulosus*.
Pomacææ. *Pýrus bollwylleriana*, *Chamaemépilus*; *Cydónia vulgaris*, *Ame-lanchier vulgaris*, *Cotonéaster tomentosa*, *Méspilus germanica*; *Crataëgus intermédia*, *monógyna*, * *Azarólus*.
Amygdalæææ. * *Amýgdalus communis*, *Cérasus Mahàleb*.
Leguminosææ. *Genísta radiata*, *sagittális*, *ovata*, *decumbens*, *germanica*; *Onónis Nátrix*, *rotundifolia*; *Colútea arboréscens*, *Astrágalus aristatus*; *Coronilla Émerus*, *glauca*; *Cýtissus alpinus*, *Labúrnum*, *nigricans*, *sessilifolius*, *hirsutus*, *capitatus*.
 * *Urticææ*. *Fícus Cárica*, *Mórus álba*.
Ulmacææ. *U'lmus effúsa*, *Céltis australis*.
Cupulliferææ. *Quercus pubéscens*.
Betulínææ. *Bétula pubéscens*; *Álnus incána*, *viridis*, *glutinosa ineisa*, *glutinosa laciniata*.
Salicinææ. *Salix fissa*, *Villarsiana*, *præ'cox*, *eineráscens*, *styláris*, *myrtillóides*, *arbutifolia*, *retúsa*, *ripária*, *pátula*, *versifolia*, *Lappónum*, *Pontederána*, *Jacquiniána*, *ovata*; * *Pópulus dilatata*.
Euphorbiacæææ. *Euphórbia sylvática*, *Charácias*; *Búxus sempervirens*.
Celastrinæææ. *Euónymus latifolius*.
Rhamnæææ. *Rhámmus saxátilis*, *alpinus*, *pùmilus*.
Staphyleacæææ. * *Staphyléa pinnata*.
Acerinæææ. *Acer Pseudo-Plátanus*, *platanóides*, *opulifolium*.
 * *Vitæææ*. *Vitis vinifera*. *Anacardiæææ*. *Rhús Cótinus*.
Rutacæææ. *Rúta graveolens*, *montána*.
Polygalæææ. *Polýgala Chamæbúxus*.
Cistinæææ. *Cístus salviaefolius*; *Helianthemum Fumána*, *cànum*, *alándicum*, *alpéstre*, *salicifolium*, *pilósum*, *apenninum*, ? II (or ? C.) *calycinum*.
Tamariscinæææ. *Támarix germanica*.

Chenopòdææ. Salsòla prostràta.

Ericæcææ. Erica herbæcæa, arbòrea; Rhododéndron ferrugíneum, hírsútum.

Compòsitæ. Artemísia Abrótannu, Helichrysum Stæ'chas.

Caprifoliæcææ. Lonícera nígra, Xylósteum, alpígena, carúla; Caprifólium étráscum, Sambùcus racemòsa.

Córneæ. Córnu máa.

* *Ebenæcææ.* Diospýros Lótus.

* *Jasmíneæ.* Jasminum officinále.

* *Oleæcææ.* Syringa vulgàris.

Labiàtæ. Hyssòpus officinális, Lavándula Spíca, Teucrium montànum, Rosmarínus officinális, Sálvia officinális.

Conifereæ. Pinus Mùghus, pumíliu, Cembra; Abies Píceæ, excélsa; Lárìx europæ'a, Juníperus Sabína, E'phedra distàchya.

An extremely interesting account of the indigenous and exotic trees of Switzerland, kindly prepared for us by M. Alphonse De Candolle, will be found in the twelfth volume of the *Gardener's Magazine*; and to that we must refer for the geographical distribution of the indigenous species, confining here ourselves to an extract from it respecting the more remarkable native trees, and those which are exotic.

The exotic trees cultivated in Switzerland have been introduced by degrees as ornamental plants; those now most common are, the horsechestnut, the catalpa, the tulip tree, several kinds of limes and maples, and the cedar of Lebanon. Till the beginning of the present century these trees were planted only by proprietors who had some connexions or commercial dealings with other countries, particularly with England. There were scarcely any nurserymen or botanic gardens at Zurich, Bâsle, Geneva, or Berne, but what were reduced to confined spots in the interior of the fortified towns, and which, consequently, could have little influence on the culture of trees in the country.

This state of things has changed greatly within the last twenty years. The new botanic garden of Geneva, planted in 1818, in a less confined situation than the old one, and confided to the care of M. De Candolle, has presented to the public a numerous collection of trees and shrubs, especially of fruit trees, chiefly from the nursery of the Messrs. Baumann at Bollwyler. The sight of this garden, which is constantly open to the public, excited the zeal of proprietors, and soon made them anxious to form plantations of various kinds of trees. This gave encouragement to the nurserymen, and their establishments became, in consequence, greatly extended, and their number increased. The season of peace and security which has followed a long period of war, has produced nearly the same result throughout all Switzerland. The number of country seats has greatly increased, particularly near the towns, and, more especially, in the cantons of Geneva, Vaud, Bâsle, and Berne; and a number of parks and groves have been planted, which now ornament the country. In the environs of Geneva, for example, the sale of trees and shrubs is four or five times as great now, as it was about twenty years ago.

The largest nursery in Switzerland at present is supposed to be that of Messrs. Dailledouze, at Saconex, near Geneva, which may be fairly estimated now to contain nearly 1000 ligneous species, in the botanic sense of the word species; and above 2000 species and varieties, if we count among the number the principal modifications of fruit trees, roses, and azaleas. The other public and private gardens of Switzerland contain few varieties, especially of the hardy kinds, which are not in the nursery of Messrs. Dailledouze. In adding, then, about 200 species to those which are to be found in this nursery, and counting about eighty indigenous ligneous plants, which have not been taken into cultivation, we shall find that the whole ligneous flora of Switzerland amounts to about 1300 species. The result of this approximated calculation shows that, from foreign commerce and improved cultivation, five or six times as many species of trees and shrubs are now grown in Switzerland, as existed there originally.

In the botanic garden, *Zizyphus vulgàris*, the pomegranate, and the fig, live against walls, and ripen their fruits. The kôlreuteria, the cork tree, *Quercus fastigiata*, *Juglans fraxinifolia*, and the *Arando Dônax*, stand at Geneva, even in the open country. *Quercus fastigiata*, in particular, which was introduced by M. De Candolle about 1820, promises to become a great ornament to the Swiss parks. The resemblance which this tree bears to the Lombardy poplar, except that it has an oak leaf, gives it a very extraordinary degree of interest.

If we search in Switzerland for trees remarkable for their antiquity or rarity, we shall find several worthy of being mentioned here. We shall begin with indigenous trees, and then proceed to those which are exotic.

At Fribourg, in the public square, there is a large lime, the branches of which are supported by pieces of wood. This tree was planted on the day when the victory was proclaimed of the Swiss over the Duke of Burgundy, Charles the Rash, in the year 1476; and it is a monument admirably accordant with the then feebleness of the Swiss republics, and the extreme simplicity of their manners. In 1831, the trunk of this tree measured 13 ft. 9 in. in circumference.

The tree of Trons, in the Grisons, is a monument of a similar nature: under the shade of this tree, it is said that the deputies of the country swore to free themselves from the yoke of their lords. This tree is celebrated in all the local poems as being a lime, but the fact is, that it is a sycamore (*Acer Pseudoplatanus*), the trunk of which is now 26 ft. 6 in. in circumference at 1 ft. 6 in. from the ground. We can hardly suppose that it could have been less than 100 years old, when it served as a place of rendezvous for the conspirators, in which case it must be now nearly 500 years old. In the *Bibliothèque Universelle de Genève*, for August, 1831, there is a letter from Colonel Augustus Bontemps, in which it is mentioned, that the probable reason why this sycamore is called a lime in the local poems is, that the German word "ahorn," which signifies a sycamore, is very unpoetical, while that for a lime tree, "linde," is soft and liquid; and this made the former be rejected by the writers of the old ballads.

At Zoffingen there are two lime trees, on the branches of which is placed a plank in such a manner as to enable any one to walk from the one to the other; and thus, people may not only walk, but even dance, upon the foliage of the tree. In the village of Villars-en-Morig, near Fribourg, there is a large lime, which existed there long before the battle of Morat (which the tree at Fribourg commemorates), and which is now of extraordinary dimensions. It was, in 1831, 70 ft. high, and 36 ft. in circumference at 4 ft. from the ground, where it divided into large and perfectly sound branches. It must be nearly 1000 years old. (See De Candolle's *Physiologie Végétale*, p. 987.)

These are certainly the most remarkable trees in Switzerland, because they are all linked, more or less, with the history of the country. They speak to the imaginations of the people, and are connected, not only with the amusements of each generation, but with the victories that, in ancient times, secured the independence of the Swiss.

We shall now mention some trees which are interesting in a botanical point of view. These are almost all found in the neighbourhood of Geneva, where the Messrs. De Candolle, father and son, have taken care to measure them, in order to commence a series of exact observations on the growth of trees. These two naturalists are aware, that, in order to calculate the age and products of old trees, we want data as to their growth after they have passed a century or two of their existence; and they have conceived the idea of making a register of all the numerous measurements that they have taken, designating exactly the local position of the trees. They mean to deposit this register in some public establishment, in order that other botanists may, after them, continue the same kind of observations on the same trees during several centuries.*

* The botanical reader is, no doubt, aware that Professor De Candolle was the first to throw out the idea, that exogenous trees have no definite term affixed to their existence, and, consequently, that there can be no limit to the number of years that a dicotyledonous tree may live. (See *Physiologie Végétale*, vol. iii. p. 957—1022.)

The promenade of the Isle of Barques, at Geneva, at the exit of the Rhone, has several fine hornbeams; the largest of which was, in 1831, 8 ft. in circumference at 3 ft. above the soil. In the same year, a lime tree in the country seat of Vicsseaux, at Chatelaine, had a trunk 13 ft. $\frac{1}{2}$ in. in circumference. Two elms situated at Pré-l'Évêque were, in 1833, at 3 ft. from the ground, 17 ft. in circumference. The largest beeches in the country were situated at the entrance to the Abbey de Pommers sous Salève. One of them was, in 1833, at 2 ft. from the ground, 15 ft. 6 in. in circumference, and the other 15 ft. 4 in.

Among the foreign trees we may notice the horsechestnut of Mr. Charles Martin at Molagnore. It passes for one of the most ancient in the country, and is 13 ft. 3 in. in circumference, with a top which projects very far over the adjacent road. The park at Ferney does not present any remarkable exotic tree; but they show an elm, planted by Voltaire in 1763, of which the trunk, in 1831, was 6 ft. $\frac{1}{2}$ in. in circumference at 4 ft. from the ground. Since that time the tree has been so ill-treated by visitors, who have stripped off portions of its bark as a memorial of the great poet of Ferney, that it has been found necessary to surround it with stakes. The park of Bossière, near Geneva, has some fine trees, and had, a few years since, some *Cytisus alpinus* [Scotch laburnums] which were nearly 40 ft. high. The finest of these trees perished some years ago, but the remainder are still well worth visiting.

Near to Geneva, the country seat which presents the greatest number of old exotic trees is the residence of M. Gaussen, at Bourdigny: it is there that grows the female salisburia, the only old specimen in Europe. M. De Candolle having discovered the sex of this plant, by observing that it bore fruit, about the year 1818, hastened to send cuttings and grafts of it to all the principal gardens in Europe. He published some remarks on it in the *Bibliothèque Universelle*, vol. vii. p. 38. The precise epoch of its being planted is unknown. The former proprietor of Bourdigny, M. Gaussen of Chapeaurouge, was a zealous amateur, who exerted himself to procure foreign seeds, and generally obtained his plants of foreign trees from England. He began his plantations in the year 1767, and he continued planting during 30 years. The female salisburia, when measured in April, 1835, at 1 ft. 10 in. from the ground, was precisely 4 ft. in circumference. The head was depressed, and did not reach higher than 12 ft. or 15 ft.; but it spread out, laterally, to such an extent as to cover a space 25 ft. in diameter.

This tree is perfectly healthy, and produces fruit (*pommes*) every year; which, however, do not contain any fertile seeds, because there is no male tree in the immediate neighbourhood. M. Gaussen has latterly grafted some male branches on his tree, but the grafts have not taken. The only male salisburia which grows in the neighbourhood of Geneva, is three leagues from the female one, at Philosophes, the seat of M. Alexandre Prévost, formerly Swiss consul in England. This tree is 2 ft. 8 in. in circumference, and its habit of growth resembles that of the specimen at Bourdigny. If the grafts do not ultimately succeed, branches of the male plant, in flower, may be brought to fructify the female plant, in the same manner as the castrification of the date palms is effected in Egypt.

In the same country seat belonging to M. Gaussen, there are, also, a cork tree, the trunk of which is above 3 ft. $\frac{1}{2}$ in. in circumference, at 3 ft. from the ground; a female *Negundo fraxinifolium*, which is 3 ft. 3 in. in circumference at $\frac{1}{2}$ ft. from the ground, and at least 40 ft. high; a *Juniperus virginiana*, 3 ft. 2 in. 9 lines in girth; a beautiful chionanthus, some fine platanuses, and other well-grown trees.

The celebrated De Saussure planted several exotic trees on his terrace in the town of Geneva; and persons passing along the Rue de la Corraturie may remark the fine effect produced on this terrace by an old catalpa and a *Juglans nigra*.

The oldest *Taxodium distichum* in the neighbourhood of Geneva is not more than 30 years old; it exists on the grounds of M. Rigot, at Varembe

M. Dumant possesses, on his grounds at Secheron, a *Quercus Ilex*, which is very fine for the country.

At the entrance of the botanic garden there is an old *Ailántus glandulosa*, much older than the garden. It measured, in June, 1833, at the level of the soil, 7 ft. 3 in. in circumference, and was between 45 ft. and 50 ft. high. The unpleasant smell of its flowers is perceived at half a furlong's distance, and its numerous and troublesome suckers rise all round, as far as 40 ft. or 50 ft. from the tree.

The trees contained in the botanic garden itself are not old, as the garden has not been established more than 17 years. Among the rarest and best-grown trees which have been planted from 15 to 17 years, we may mention the following:—A *Photinia serrulata*, spreading into branches from its base, and about 12½ ft. high; a *Magnolia acuminata*, about the same height, with a trunk 7 in. in circumference; a *Kölreuteria*, 15 ft. high, with a trunk 13 in. in circumference; and a tulip tree, 40 ft. high, 3 ft. 5½ in. in circumference; *Acer striatum*, 24 ft. high, and 1 ft. 7½ in. in circumference; *Aesculus flava* and *rubicunda*, 30 ft. high, and 1½ ft. in circumference; *Pavia hibernica*, 20 ft. high, and 14½ in. in circumference; *Cerasus serotina*, 35 ft. high, and 2 ft. 4 in. in circumference; *Mespilus Smithii* Dec., 20 ft. high, and 1 ft. 8 in. in circumference; *Cratægus nigra*, about the same height, and 1 ft. 2½ in. in circumference; *Hippophae rhamnoides*, and *Elaagnus angustifolia*, 12 ft. high, and 1 ft. in circumference; *Plánera crenata*, 35 ft. high, and 1 ft. 9 in. in circumference; *Pópulus angulata*, which sometimes retains its leaves till Christmas, 60 ft. high, and 4 ft. in girth; *Alnus cordata*, 35 ft. high, 1½ ft. in girth; *Quercus alba*, 18 ft. high, and 2 ft. 1 in. in girth; *Juniperus thurifera*, 15 ft. high, and 1 ft. in girth; *Pinus Laricio*, 25 ft. high, and 2 ft. 4 in. in girth; *Pinus Mughus*, 20 ft. high, and 2 ft. 5 in. in girth; and *Larix europæa*, the branches of which hang in a very singular manner, is 30 ft. high, and 2 ft. 5 in. in circumference near to the base of the trunk. All these measurements were taken at such a height from the ground as seemed most likely to give the true dimensions of the trunk, and to avoid the thickness often produced by the graft, or at the base; and these measures were all taken by M. Alphonse De Cándolle, in October, 1835. It is to be wished that a register of similar measurements were opened in every botanic garden, in order to verify the date of the introduction, and the rate of growth, of every species, according to the diverse physical circumstances of each locality. (*A. De C. Nov.*, 1835.)

SECT. VII. *Of the Indigenous and Foreign Trees and Shrubs of Italy, Greece, Spain, Portugal, and the Mediterranean Islands.*

ITALY, having been, during the Roman empire and the dark ages, the centre of civilisation in Europe, would, doubtless, draw from all other countries whatever of their productions was suitable to its climate. Hence the ligneous flora of Italy includes almost all the trees and shrubs indigenous to Greece, Spain, and the Mediterranean islands, which are in any way remarkable for their use or beauty. In the following enumeration, taken from Tenore's *Flora Neapolitana*, Bertoloni's *Flora Italica*, Savi's *Botanicon Etruscum*, Smith's *Prodromus* of Sibthorp's *Flora Græca*, Brotero's *Flora Lusitânica*, Gussone's *Flora Sicula Prodromus*, and Hogg's *Observations on the Classical Plants of Sicily*, we have included all the ligneous plants which are indigenous, or apparently so, in these countries, and which are not included in the indigenous flora of Britain. Those which are believed to be peculiar to any one or two of the countries, have the name of such countries following the name of the plant.

Ranunculaceæ. *Clematis cirrhosa*, *Viticella*, *campaniflora* (Portugal), *Flammula*, *Flammula* var. *rotundifolia* Dec. (Naples).

Berberidæe. *Berberis cretica* (Greece).

Cruciferae. *Iberis sempervirens* (Greece); *Alýssum rupestre* (Naples), *argenteum* (Etruria).

Capparidææ. *Cápparis spinòsa*, Fontanèsü, and rupéstris (Greece).

Cistinææ. *Cistus crispus* (Portugal, Greece), *álbidus* (Portugal, Greece), *salviæfólius*, *hirsütus* (Portugal), *monspeliénsis*, *populifólius* (Portugal), *ladaniferus* (Portugal), *incanus*, *villòsus* (Greece, Etruria), *laurifólius* (Greece), *créticus* (Greece), *parvifórus* (Greece).

Helianthemum Libanòtis (Portugal), *umbellàtum* (Portugal), *umbellàtum* var. (*Cistus verticillàtus Brotero*) (Portugal), *ocymòides* (Portugal), *halimifólius*, *cheiranthòides* (Portugal), *lasiánthum* (Portugal), *involveràtum* (Portugal), *scabròsum* (Portugal), *Fumàna*, *læ'vipes*, *origanifólius* (Portugal), *thymifólius* (Portugal, Greece), *stæchadifólius* (Portugal, Naples), *hispidum* (Portugal), *alpèstre* (Greece), *pilòsum* (Greece), *lavandulæfólius* (Greece, Naples), *racemòsum* (Greece), *hirtum* (Greece), *ellípticum* (Greece), *apenninum* (Greece), *arábicum* (Greece, Etruria), *vírde* (Naples), *itálicum* (Etruria).

Polygálææ. *Polýgala Chamæbúxus* (Naples), *microphýlla* (Portugal).

Caryophýllææ. *Dianthus arbòreus* (Greece), *fruticòsus* (Greece), *Silène fruticòsa* (Greece, Sicily).

Línææ. *Linum arbòreum* (Greece).

Malvæææ. *Lavàtera Olbia* (Greece, Portugal), *triloba* (Portugal).

* *Aurantiæææ.* *Citrus Médica* (* Sicily, * Portugal), *Aurántium* (* Sicily, * Portugal).

Hypericínææ. *Hypericum calycinum* (Greece), *hircinum* (Greece), *empetrifólius* (Greece), *repens* (Greece), *Còris* (Greece, Etruria), *linearifólius* (Portugal).

Acerínææ. *Acer monspessulanum* (Greece, Naples), *créticum* (Greece, Sicily), *obtusifólius* (Greece), *Pseudo-Plátanus* (Naples, Portugal), *O'pulus* (Naples), *neapolitanum* (Naples).

Hippocástànææ. * *Æsculus Hippocástanum* (Greece, * Portugal).

Meliæææ. *Mèlia Azedarách* (Portugal).

* *Vites.* *Vitis vinífera*.

Rutæææ. *Rúta graveolens*, *montàna* (Greece, Naples), *chalepénsis* (Greece), *divaricatà* (Naples), *angustifólia* (Naples), *macrophýlla* (Naples), *tenuifólia* (Portugal), *bracteosa* (Sicily); *Aplophýllum linifólius* (Greece).

Zygophýllææ. *Zygophýllum álbum* (Greece).

Xanthoxýllææ. *Cneòrum tricóccum* (Italy).

Celastrínææ. *Enónymus latifólius* (Greece).

Staphylæææ. *Staphylèa pinnàta*.

Rhámneææ. *Rhámnus infectòrius* (Greece), *oleòides* (Greece, Sicily), *pruinifólius* (Greece), *saxatilis* (Greece), *alpinus* (Greece), *pubéscens* (Greece), *Alatérnus*, *Clùsiü* (Sicily), *lyciòides* (Portugal), ? *buxifólius* (Portugal); *Paliùrus aculeàtus* (Greece, Sicily); *Zizyphus vulgàris*, *Lòtus* (Portugal, Sicily).

Anacardiæææ. *Pistàcia Terebínthus* (Greece, Portugal), *vèra* (Sicily), *Lentiscus*; *Rhüs Coriària*, *Cótinus* (Greece, Sicily), *pentaphýlla* (Sicily), *dioica* (Sicily.)

Leguminòsææ. *Spártium juncèum*; *Genísta Scòrpius* (Greece), *angulàta* (Greece), *hórrida* (Greece), *humifusa* (Greece), *eàndicans*, *sphærocàrpa* (Portugal), *monospérma* (Portugal), *tridentàta* (Portugal), *polygalæfólia* (Portugal), *Bróteri* (Portugal), *lusitànica* (Portugal), *triacánthos* (Portugal), *falcàta* (Portugal), *algarbiénsis* (Portugal), *germánica* (Portugal), *sagittàlis* (Etruria), *radiàta* (Naples, Etruria), *hirsùta* (Naples), *ovàta* (Naples, Etruria), *scariòsa* (Naples, Etruria), *auxàntica* (Naples), *diffùsa* (Naples), *ephedròides* (Sardinia), *Cupàni*; *Cýtisis lániger*, *pónticus* (Greece), *sessilifólius*, *hirsütus*, *triflórus*, *grandiflórus* (Portugal), *pàtens* (Portugal), *Labúrum*, *nigricans* (Etruria), *supinus* (Naples), *argénteus* (Naples), *álbus* (Naples, Portugal), *spinòsus* (Etruria), *capitátus* (Etruria); *Adenocàrpus hispànicus* (Portugal), *parvifólius*; *Stauracánthus aphýllus* (Portugal); *Anthýllis Bárba Jòvis* (Greece, Naples), *Hermànnææ* (Greece); *E'benus crética* (Greece); *Colùtea arborés-cens*; *Coronilla E'merus*, *glàuca* (Greece, Portugal); *Allági Mauròrum* (Greece); *Astrágalus angustifólius* (Greece); *aristátus* (Greece), *créticus*

(Greece), Potèrium (Portugal); Psoràlea bituminosa (Greece, Portugal); *Dorýenium* hirsutum (Greece), réctum (Greece), suffruticosum (Greece); *Ceratonia* Siliqua; *Anagyris* foetida; *Cercis* Siliquastrum; *Medicago* arborea (Greece, Sicily); *Ononis* Natrix (Portugal), hispánica (Portugal); *Lotus* argenteus (Portugal), cretèus (Portugal).

Rosaceæ. *Rosa* glutinosa (Greece, Sicily), sempervirens (Greece, Sicily), scandens *Brotero* (Portugal), heclèliana (Sicily), *Seraphini* (Sicily), pulverulenta (Sicily), gállica (Sicily); *Rubus* tomentosus (Greece, Sicily), hirtus (Sicily); *Spiræa* arenata (Portugal).

Pomaceæ. *Mespilus* germanica; *Cydonia* vulgaris; *Amelanchier* vulgaris; *Pyrus* salicifolia (Greece), erètica (Greece), *Chamæmispilus* (Greece), cuneifolia *Guss.* (Sicily), nebrodènsis *Guss.* (Sicily), præmorsa *Guss.* allied to *aucuparia* (Sicily), acèrba *Dec.* (Sicily); *Crataegus* *Pyracantha*, monogyna, *Azarolus*, *tanacetifolia* (Greece), nigra (Naples), laciniata *Ueria* (Sicily), florentina (Etruria).

Sanguisorbeæ. *Potèrium* spinosum.

Amygdales. *Amygdalus* communis, incana (Greece), nana (Greece); *Persica* vulgaris (Sicily, Portugal); *Armeniaca* vulgaris (Portugal); *Cerasus* lusitánica (Portugal), *Mahaleb* (Greece, Sicily), caproniana (Sicily), *Lauro-cerasus* (Greece), prostrata (Greece).

Granateæ. *Punica* Granatum.

Tamariscinæ. *Tamarix* gállica, africàna (Sicily).

Philadelphicæ. *Philadelphus* coronarius (Naples, Portugal).

Myrtaceæ. *Myrtus* communis, c. var. italica (Sicily), c. var. romana (Sicily), c. var. hætica (Sicily), c. var. lusitánica (Sicily).

Crassulacæ. *Sempervivum* arboreum (Greece, Portugal).

Cactææ. *Opuntia* vulgaris (Portugal, Sicily), maxima (Sicily).

Umbelliferæ. *Bupleurum* fruticosum (Greece, Sicily), *Sibthorpiænum* (Greece).

Araliaceæ. *Hedera* *Helix* chrysoearpa (Sicily).

Caprifoliaceæ. *Caprifolium* étruscum, implèxum (Sicily), canescens (Sicily); *Lonicera* nigra (Greece), *Xylosteum*, alpigena (Greece); *Fiburnia* *Tinus* (Portugal), *T.* hirta (Naples), *T.* lucida (Naples), *T.* virgata (Naples, ? * Sicily); *Sambucus* racemosa.

Cornææ. *Cornus* mas (Greece, Etruria).

Loranthaceæ. *Loranthus* europæus.

Cinchonacæ. *Ernodea* montana (Greece, Sicily).

Compositæ. *Stæhelia* arborescens (Greece), fruticosa (Greece), uniflorulosa (Greece), *Chamaepèice* (Greece); *Artemisia* arborescens (Greece, Portugal); *Melichrysum* *Stæchas* (Greece, Portugal), orientale (Greece, * Portugal); *Conyza* saxatilis, pumila (Greece), candida (Greece, Naples), *limoniifolia* (Greece); *Cineraria* maritima (Greece); *Bupthalmum* maritimum; *Santolina* *rosmarinifolia* (Portugal, Sicily), *Chamaecyparissus* (Portugal); *Calendula* suffruticosa (Portugal).

Ericaceæ. *Erica* arborea, multiflora, manipuliflora (Greece), herbacea (Greece, Etruria), spiculiflora (Greece), scoparia, australis (Portugal), umbellata (Portugal), mediterranea (Portugal), scula *Gussone* (Sicily); *Arbutus* *Andrachne* (Greece).

Styracineæ. *Styrax* officinale (Greece).

Ebenaceæ. *Diospyros* *Lotus* (Greece).

Oleaceæ. *Olea* europæa; *Phillyrea* media, media *ligustrifolia* (Sicily), media *luxifolia* (Sicily), latifolia, latifolia spinosa (Sicily), angustifolia, stricta (Italy), hævis (Naples); *Fontanèsia* *phillyreoides* (Italy, Sicily); *Ornus* europæa (Greece, Italy), rotundifolia (Italy); *Fraxinus* *parvifolia* (Italy), argentea (Italy).

Jasminææ. *Jasminum* fruticans.

Apocynææ. *Nerium* *Oleander*; *Fmea* minor, major.

Asclepiadææ. *Periploca* græca (Greece), angustifolia (Sicily); *Gomphocarpus* fruticosus (Sicily).

Convolvulacæ. *Convolvulus Cneorum, Dorýcnium* (Greece), lanátus (Greece).

Boraginæ. *Lithospermum hispídulum* (Greece), fruticosum (Portugal), rosmarinifólium (Sicily).

Solanæ. *Solanum sodomæum, * Pseudo-Cápsicum* (Portugal), moschátum (Sicily); *Lýcium bárbarum* (Greece), europæum, ? áfrum (Sicily).

Verbenacæ. *Fítex Ágnus-cástus var. latifólia* (Portugal).

Labiátæ. *Teucrium frúticans, brevifólium* (Greece), créticum (Greece), quadrátulum (Greece), *Arduini* (Greece), massiliénse (Greece), flavum, montanum, *Pólium, capitátum* (Greece, Naples), cuneifólium (Greece), alpestre (Greece), spinosum (Naples), *Pseudo-Hyssopus* (Naples); *Saturja nervosa* (Greece), *Thýmra* (Greece), montána (Greece, Etruria), capitata (Greece, Sicily), spinosa (Greece); *Thýmra spicata* (Greece); *Lavándula Spica* (Greece, Etruria), *Stæ'chas, ?* dentata* (Greece), multifida (Portugal); *Sálvia spinosa* (Greece), palestina (Greece); *Beringeria Pseudo-Dictamnus* (Greece); *Phlomis fruticosa* (Greece, Naples), ferruginea (Naples); *Moluccella frutescens* (Greece); *Origanum Dictamnus* (Greece), *Tourneforti* (Greece); *Thýmus vulgaris* (Greece, Portugal), lanceolatus (Greece), *Zýgis* (Greece, Portugal), villosus (Greece), cæspititius (Portugal), *Masticina* (Portugal), *Tragoriganum, micranthus* (Portugal), créticus (Portugal), cephalotus (Portugal); *Acynus graveolens* (Greece); *Prásium majus*; *Rosmarinus officinalis*; *Sálvia officinalis, pomifera* (Greece), calycina (Greece), triloba (Greece, Italy), canariensis (Sicily).

Globularinæ. *Globularia Alýpum.*

Plumbaginæ. *Státice monopctala.*

Plantaginæ. *Plantago Cýnops* (Greece, Sicily), subulata (Sicily), macrolobiza (Sicily), áfra (Sicily).

Amarantacæ. *Achyranthes argentea* (Naples, Sicily).

Chenopodæ. *Salicornia fruticosa, cruciata* (Italy), macrostachya (Sicily); *Anábasis aphylla* (Greece); *Atriplex Halimus, glauca* (Greece, Portugal), græca (Greece); *Camphorosma monspeliaca* (Italy, Sicily); *Salsola vermiculata* (Portugal), *argentina Gussone* (Sicily), oppositifolia (Sicily).

Laurinæ. *Laurus nobilis.*

Thymelææ. *Dáphne dioica* (Greece), *Tarton-raira* (Greece, Naples), argentea (Greece), póntica (Greece), *Gnídium, buxifolia* (Greece), oleoides (Greece), *jasminea* (Greece), sericea (Greece), alpina, collina (Greece), glandulosa *Bertoloni* (? oleoides) (Sicily), *Cneorum* (Etruria); *Passerina hirsuta.*

Santalacæ. *Osyris álba* (Greece, Portugal).

Elæagnæ. *Elæagnus angustifolia* (Greece).

Aristolochiæ. *Aristolochia sempervirens* (Greece), subglauca (Portugal).

Euphorbiacæ. *Euphorbia pumila* (Greece), spinosa (Greece, Naples), dendroides (Greece, Sicily), sylvatica (Portugal, Sicily), *Characias, lanuginosa* (Naples), fruticosa (Sicily), coralloides (Sicily), tanaicensis (Sicily), *Pinea* (Sicily), *Myrsinites* (Sicily), biglandulosa (Sicily); *Báxus sempervirens* (Greece, Portugal); *Mercurialis elliptica* (Portugal), tomentosa (Portugal).

Urticæ. *Ficus Cárica.*

Ulmacæ. *Celtis australis, ? Ulmus Abelicea* (*Fl. Gr. Prod.*).

Cupuliferæ. *Quercus Ballota* (Greece), *Ilex, coccifera, rigida* (Greece), infectoria (Greece), *Ægilops* (Greece), *Esculus* (Greece, Sicily), pubescens (Greece, Portugal), crinita (Greece), racemosa (Portugal), híbrida (Portugal), fruticosa (Portugal), lusitânica (Portugal), rotundifolia (Portugal), *Süher* (Portugal, Sicily), hispânica (Portugal); *Corylus Colurna* (Greece); *Ostrya vulgaris* (Greece); *Castanea vesca* (Sicily).

Betulicæ. *Álnus cordifolia* (Naples).

Salicinæ. *Salix ægyptiaca* (Greece), retusa (Naples), riparia *Tenore* (synonymes, incana *Dec.*, viminális *Villars*) (Naples), salviaefolia (Portugal), atrocinerea (Portugal).

Platanæ. *Platanus orientalis* (Greece, Sicily).

Myricæ. *Myrica Faya* (Portugal).

Coniferae. *Pinus Pinea*, marítima; *Abies Pícea* (Greece); *Cuprèssus sempervirens*, lusitánica (* Portugal); *Juníperus Oxýcedrus*, macrocárpa (Greece), lýcia (Greece), phœnícea (Greece, Portugal), *Sabína* (Greece, Sicily); *Ephedra distácha* (Greece, Portugal).

Empétreae. *Corèma álbum* (Portugal).

Smilácea. *Smilax áspera* (Greece, Sicily), nígra (Greece), excélsa (Greece).

Asphodélea. *Aspáragus acutifólius*, aphýllus (Greece, Sicily), hórridus (Greece, Sicily), verticillátus (Greece), álbus (Portugal, Sicily).

Bromeliácea. *Agave americana* (Portugal, the more southern part of).

Pálmae. *Phœnix dactylífera* (Sicily, * Portugal); *Chamaerops húmilis* (Sicily).

Shortly after this period many foreign trees and shrubs were imported from England into the gardens of Signor del Negro of Genoa, and by him distributed among the amateurs of his neighbourhood. Some account of the gardens in which these trees were planted will be found in the *Encyc. of Gard.* (edit. 1835), made from our personal observations in 1819.

Having thus enumerated the species found in all these countries that are not indigenous to Britain, we shall subjoin some remarks on the trees and shrubs of Italy, of Spain and Portugal, and of Turkey and Modern Greece.

SUBJECT. 1. *Of the Trees and Shrubs of Italy.*

THE introduction of foreign trees and shrubs into Italy, in modern times, may date from the discovery of India by the Portuguese in 1494, or, rather, from their first settlement at Goa in 1510; from the intercourse of France and England with North America in the commencement of the seventeenth century; from the settlement at the Cape of Good Hope by the Dutch in 1650; and, lastly, from the discovery of Australia. From all these countries, but chiefly from the last, a number of trees and shrubs have been brought to Europe; which, though they require the protection of a green-house in England, thrive in the open air in the neighbourhood of Naples, in Sicily, and in warm situations about Genoa. Among Indian plants may be mentioned, as growing freely in the open air in the south of Italy, the orange and lemon, the *Lagerstræmia Índica*, the cotton tree, and the cinnamon tree, which attain the height of small trees; from Syria, the *Acácia Julibrissin*, or silk tree. Among those from North America are, the magnolias, and various shrubs from the southern states, the agave from Mexico, and the palmetto from Louisiana. Among those from the Cape of Good Hope, are all the ligneous *Geraniácea*, many of the heaths, the diosmas, the proteas, the melaleuca, and similar species. From Australia there are many trees in Italy, which have already attained a large size; and there is scarcely a doubt but that nearly all the ligneous flora of that part of the world might be transplanted to Italy, including Sicily, with the most perfect success. As a proof of this, we may refer to the dimensions of certain Australian trees planted at Caserta, in the neighbourhood of Naples, as given in the *Gardener's Magazine*, vol. xi. p. 150. and p. 481. It appears that *Eucalyptus robústa* attains at Caserta, in a very few years, the height of 100 ft.; *Callistemon lophánthus*, and *Acácia heterophýlla*, upwards of 50 ft. The *Magnolia grandiflora* has attained the height of nearly 60 ft.; the camellia 25 ft.; and the melaleuca from 25 ft. to 30 ft. In Sicily, we are informed by Woods, Hogg, and other travellers, the palm and the *Ficus Sycómorus* grow as freely as in Egypt, the sugar cane and the bamboo nearly as well as in the East or West Indies, and the papyrus and the nelumbium succeed in the waters. As the warmest parts of Sicily, therefore, admit of growing the plants of the warmest parts of Africa in the open air, there can, we think, be little hazard in supposing that, between the north of Italy and its southern extremities, the ligneous flora of the whole world might, with a very little assistance from art, be included.

The first introduction of foreign trees and shrubs into Lombardy, we are informed by Signor Manetti, the director and controller of the viceregal gardens at Monza, took place about the year 1770; they were planted by

the brothers Pecinardi, near Cremona. In 1785 great additions were made to the foreign trees and shrubs of the north of Italy, by Count Louis Castiglione, who undertook a voyage to North America, and brought home a great number of seeds, which he sowed at Mozzate, afterwards distributing the plants over all Italy. In 1811 farther additions were made to the foreign trees and shrubs of Lombardy, by M. Villaresi, then director of the gardens of Monza; and in 1814 still further additions were made by the present viceroy. In consequence of these introductions, there are now, in the park and gardens at Monza, many fine specimens of exotic trees. *Magnolia conspicua* flowers every year, and ripens abundance of seed. *M. grandiflora*, at 60 years old, is 36 ft. high; and, though in a very unfavourable situation, viz. a dry soil and a warm sunny exposure, it flowers and seeds freely. There are above 230 of these trees in the plantations of the park, besides numerous plants of all the other species. The tulip tree has attained the height of 70 ft. in 29 years, flowering and seeding every year. *Alantus glandulosa*, 29 years planted, is 60 ft. high; and *Robinia Pseud-Acacia*, of the the same age, is 75 ft. high, with a trunk 2 ft. in diameter, and branches covering a space of 120 ft. in circumference. There are many other fine trees in the grounds at Monza, details respecting which will be found in the *Gardener's Magazine*, vol. xi. p. 639.

In the garden of Count Mellerio, at Brianza, near Milan, the *Melia Azedarach*, 26 years planted, is 40 ft. high, and flowers and seeds freely every year. This beautiful tree is one of the greatest ornaments of the public promenades of the south of Italy; but there are very few parts of Lombardy where it attains so large a size as at Brianza.

In the year 1832, the Abbé Belèse made a tour through the northern part of Italy, chiefly to inspect the gardens; and he noticed, among other trees and shrubs, the following:—Near Milan, at Soma, he saw a cypress of great antiquity, which girted 20 ft., and was 70 ft. high, though it had, for many years, lost its leading shoot; popular tradition says that it was planted previously to the birth of Christ; and the Abbé Belèse's brother assured him, that there was an ancient chronicle in Milan, which proves that this tree existed in the time of Julius Cæsar, B.C. 42. In the botanic garden at Padua, the abbé found two trees of *Magnolia grandiflora*, which had been planted 90 years, soon after the introduction of the tree into Europe; they were 60 ft. high, with trunks 4 ft. in diameter; they were sown by the director of the garden, Farsetti, in 1742. There are in this garden, a salisburia, 60 ft. high; two trees of *Lagerstrœmia indica*, of 40 ft. high, which ripen seed every year; the common red-flowered althea frutex, 50 ft. high, and which, on the 8th of August, 1832, was so covered with blossoms as to resemble one immense flower of the double red camellia. *Quercus Ilex* is here 100 ft. high; *Salix annularis*, 40 ft.; *Lycium japonicum*, 25 ft.; *Acacia farnesiána*, 60 ft., the flowers of which perfumed the air for a great distance round; the date palm, 25 ft.; *Aralia spinosa*, 25 ft.; *Cerasus semperflorens*, bearing fruit and flowers at the same time, 50 ft.; *Vitex Agnus-castus*, 140 years planted, and 35 ft. high; *Tecoma stans*, 30 ft.; *Smilax Sarsaparilla*, 60 ft.; *Nicotiana glauca*, a magnificent tree-like specimen; *Cesalpinia Sappan*, 15 ft.; *Chamærops humilis*, 25 ft.; *Asimina triloba*, 20 ft., and covered with excellent ripe fruit; *Acacia Julibrissin*, 60 ft.; *Liriodendron Tulipifera*, 80 ft.; *Stereulia plataniifolia*, 40 ft.; *Casuarina distyla*, 15 ft.; and a number of others, which will be found recorded in the *Annales de la Société d'Horticulture de Paris*, tom. 12e, p. 68.

In the Isola Bella there are a great many exotic trees and shrubs of very luxuriant growth. Among these are, an immense oleander, numerous trees of *Laurus nobilis*, of great growth; and a hydrangea, 10 ft. in diameter, and 8 ft. high, planted in peat soil, and covered with deep blue flowers. In this, and in other of the Borromeo islands, the *Agave americana* stands the open air, and flowers freely. On the whole, though there are several tropical trees that will not live in the open air in the north of Italy generally, yet

there are portions of it which, from local circumstances, possess so mild a climate, that, with very little art, Lombardy might be made to exhibit specimens of the ligneous vegetation of every part of the globe.

SUBJECT. 2. *Of the Trees and Shrubs of Spain and Portugal.*

FROM the indigenous trees and shrubs of this immense tract of country we may estimate its capacity for growing the trees and shrubs of other climates; and, while those of the North of Europe will find a congenial climate in the mountain ranges, and the elevated region of Madrid, those of the tropics will grow along the shores of the Mediterranean Sea.

There are numerous botanic gardens throughout Spain, and two in Portugal. In these gardens, La Gasca informs us, the trees and shrubs of most parts of the world have, at one time or other, been seen in a flourishing state. In the gardens in the neighbourhood of Madrid are found, the cedar of Lebanon and various other Coniferae, different species of lime, maple, ash, *Aesculus*, *Juglans*, *Morus*, *Crataegus*, *Prunus*, *Pyrus*, and *Cerasus*, *Asimina triloba*, *Magnolia grandiflora*, *Sophora japonica*, the *Calycanthus floridus*, the *Chimonanthus fragrans*, and many others. Capt. S. E. Cook informs us (*Sketches in Spain, &c.*) that the date palm will, in sheltered situations, resist the cold of Madrid; though its fruit only acquires perfect maturity on the coast of Western Andalusia, and in other parts of the shores of the Mediterranean. In the gardens in the neighbourhood of Cadiz the *Musa sapientum* ripens fruit every year; as do the different species of *Capsicum*, *Cactus*, *Cereus*, and *Mesembryanthemum*. All the trees and shrubs of the Cape of Good Hope and Australia grow here as well as in their native countries. In the gardens of Seville, similar exotics thrive freely, and the country is covered with orange, lemon, citron, and lime trees, olives, pomegranates, and algarobas, or carob trees.

We shall pass over the other gardens in Spain, detailed particulars of the more rare foreign trees contained in which, by Sr. La Gasca, will be found in the first and second volumes of the *Gardener's Magazine*, to notice the trees and shrubs of the botanic garden of Lisbon. In this garden the orange, lemon, and citron ripen their fruit in the open air. The *Erythrina picta* grows to the height of 15 ft. in one season, and ripens seed. *Psidium pyriferum* and *poniferum* as standards, set their fruit, and ripen it against a wall. *Coffea arabica* flowers in October, and the berries ripen in the May or June following. *Cárica Papaya* forms a fine umbrageous tree in the open garden, and ripens its fruit; though, being a tree with a succulent or spongy trunk, it is occasionally cut down to the ground. These instances are sufficient to prove, that, with the art of the gardener and the aid of walls, without hot-houses, all the ligneous plants of the world might be grown in the peninsula.

The prominent trees in the forests of Spain are, the *Quercus Robur*, *Q. Cérris*, and its numerous varieties; *Q. Ilex*, with its varieties still more numerous than those of *Q. Cérris*; *Q. Súber*; and *Pinus Pinca*, *Pináster*, *sylvéstris*, *marítima*, &c. The silver fir is also abundant in some native forests, and the *Lárix* in the alpine regions of the northern provinces. The most remarkable of the indigenous trees are the cypresses in the gardens of the palaces which belonged to the Moorish kings; many of these venerable specimens are supposed to be upwards of five centuries old. The prevailing tree about Madrid, as about Paris and London, is the narrow-leaved elm.

The geographical distribution of the indigenous trees of Spain has been given, for the first time, after several years of ardent research and travel, by Capt. S. E. Cook, in his *Sketches of Spain*, published in 1834. Capt. Cook divides Spain geologically into three grand regions. The first is the northern, which includes Galicia, Asturias, &c., and the maritime part of Old Castile. This is the region of humidity and moisture, of the *Quercus Robur*, and *Q. Ilex*, *Ulex europæa*, and *U. c. stricta*; and the *Daboecia polifolia*. The second region includes the Castiles, Estremadura, Aragon, &c.; this is the region of dryness, over which the Merino sheep wander, and in which the olive and the

silkworm are products of culture. "This region," Capt. Cook observes, "contains the vast pine forests of Aragon, of the Sierra de Cuenca, Segura, and the Guadarrama, an l of the central range of Castile. It is characterised by the Spanish ilex; the *Quercus Tōza*; and the *Quercus p̄rasina*, or a species presumed to be so, which is widely spread over its middle elevation; by the white cistus, which grows in prodigious quantities in some of the middle parts; and by the absence of those which are enumerated as marking the divisions on each side of it." The third region lies along the coast of the Mediterranean, and is characterised by a dry and burning summer, and a mild winter. In this region the lemon, the orange, the palm, the sugar-cane, the cotton tree, the *Ceratonia Siliqua*, are the common ligneous plants in cultivation. This region contains no extensive forests, but abundance of orange orchards, olive grounds, and vineyards.

It would occupy too much space, to enter at such length into the ligneous vegetation of each region, as would do justice to the subject, and we must therefore refer the reader to Captain Cook's *Sketches*, or to an extract from them, made with the kind permission of the author, which will be found in the twelfth volume of the *Gard. Mag.* In the third part of this work, when we come to treat of particular species, we shall find much interesting matter, supplied from Captain Cook's volumes, respecting the genera *Pinus*, *Abies*, *Larix*, and *Quercus*.

The most remarkable discovery made by Captain Cook in Spain, and which was made about the same time by Mr. Drummond, the British consul at Morocco, is, that the *alerce*, a timber which is of unparalleled durability, is from the *Thuja articulata*. The roofs of the oldest churches in Spain are of this timber; and some of them, as that of the mosque of Cordova, &c., are known to have existed for nine centuries, the timber, as may be proved by a specimen sent by Captain Cook to the Horticultural Society of London, being still perfectly sound. Captain Cook, also, has collected much new and original information respecting the *Q. Ilex*; and it is remarkable, that the true Spanish evergreen oak (*Q. I. austrālis*), of which acorns can be procured in abundance from Gibraltar, had escaped the notice of both native and foreign botanists, till it was examined by Captain Cook.

SUBJECT. 3. *Of the Trees and Shrubs of Turkey and Modern Greece.*

AFTER having given, in p. 17., the enumeration of the trees and shrubs mentioned by Theophrastus, and in p. 161. those known to modern botanists, it will not be supposed that we can have much to add respecting such a country as Turkey, scarcely, as yet, in the dawn of civilisation; and where, unless the whole surface of the country can be called a garden, there are none but in the cemeteries.

These cemeteries are distinguished by their immense cypresses, and by the occasional appearance in them of the weeping willow. The most common tree in the neighbourhood of Constantinople is the *Quercus Cœrris*, and, next to this, the *Celtis austrālis*, the pinaster, and the stone pine. Other trees, considered interesting in Britain, which abound in the neighbourhood of Constantinople, are the following: *Cercis Siliquastrum*, which is found clothing the shores of the Bosphorus and Mount Libanus; *Ceratonia Siliqua*, *Cupressus sempervirens horizontalis*, *Diospyros Lötus*, *Eleagnus angustifolia*, the wild olive, *Zizyphus vulgaris*, *Palmaris aculeatus*, *Melia Acedarāch*, *Acacia Julibrissin*, *Pistacia Terebinthus*, and *P. Lentiscus*, and *Smilax aspera*, and *S. excelsa*. *S. excelsa* climbs to the tops of the highest trees; and, descending in streaming branches, converts an avenue of trees into two lofty green walls, which, in autumn, are covered with a profusion of rich red berries. There are, also, *Hælera Helix chrysocarpa*; and *Cerasus sativa*, two varieties, one of which is of enormous size, and grows along the northern coast of Asia Minor, whence the original cherry was brought to Europe, and the other is found in the woods in the interior of Asia Minor, and produces an amber-coloured transparent

fruit, of a most delicious flavour. These trees attain the height of 100 ft., with straight trunks of 40 ft. and upwards. *Phoenix dactylifera*, and *Platanus orientalis*, are also frequent. "The Turks," Dr. Walsh observes, "on the birth of a son, plant a platanus, as they do a ypress on the death of one. In the court of the scraglio is a venerable tree of this species, which, tradition says, was planted by Mahomet II., after the taking of Constantinople, to commemorate the birth of his son, Bajazet II.; the trunk of which is 50 ft. in circumference. There is another, of more enormous size, at Buyuk-dere, on the Bosphorus: it stands in a valley, and measures 45 yards in circumference! It, in fact, now consists of fourteen large trees, growing in a circle from the same root, but separating at some distance from the ground. The Turks sometimes encamp here; and the Ben-Bashee pitches his tents in the centre of this tree of trees."

Cistus crispus, *ereticus*, and *salviaefolius* cover all the hills of the Archipelago and Sea of Marmora. All of them have the hypocistus growing on their roots, a succulent parasite of a rich red colour, described by Dioscorides; *Vitex Agnus-castus*, *Nerium Oleander*, *Quercus Ægilops*, *Q. coccifera*, and *Q. Tuuzin* var. *pubescens*, *Pinus Pinaster* var. *maritima*, and *P. Pinea*, *Lavandula Stœchas*, *Ruscus racemosus*, *Kölentéria paniculata*, and a number of others. *Buxus baleariens* grows wild on all the rocky surfaces both of European and of Asiatic Turkey, and the wood is sent to England in large quantities for the use of the wood-engraver, though it is found greatly inferior to that of the *Buxus sempervirens*. This information respecting the trees and shrubs in the neighbourhood of Constantinople is taken from a very interesting paper on the subject, by Dr. Robert Walsh, which will be found in the *Transactions of the Horticultural Society of London* for 1824, and in the *Gardener's Magazine*, vol. i. p. 293.

The ligneous vegetation of the Ionian Islands is given by Pouqueville and Olivier, and resembles that of Greece generally. Whether any foreign species have been introduced, since these islands came under the protection of the British government, we have been unable to ascertain.

CHAP. IV.

OF THE TREES AND SHRUBS OF ASIA, AFRICA, AMERICA, AND AUSTRALIA, WHICH ARE SUITABLE FOR TEMPERATE CLIMATES.

In our notice of the ligneous flora of these countries, we shall confine ourselves entirely to such species as are known, or are supposed, to be suitable for enduring the open air in Britain; and, as in the preceding chapter, we shall chiefly confine ourselves to giving lists compiled from local floras. We shall take the different countries in the usual order of Asia, Africa, America, Australia, and Polynesia. Those who wish more extensive information on this subject, may consult some elaborate communications of M. Mirbel in the *Mémoires du Muséum*, vol. xiv. p. 378.; or, in an English dress, by Dr. Hooker, in Murray's *Encyclopædia of Geography*.

SECT. I. *Of the Trees and Shrubs suitable for Temperate Climates, indigenous or introduced, in Asia.*

ACCORDING to our enumeration (p. 126.), 183 ligneous plants, which endure the open air in Britain, have been introduced from different parts of Asia, and chiefly from Siberia, Nepal, and China. Additions, as we have already observed, when noticing the flora of Asiatic Russia, may still be expected from the former country; and, considering the situation of China, and the character of its surface, when that immense territory comes to be explored by

European botanists, it will, in all probability, afford large additions to our parks and shrubberies. That Himalaya and other mountainous regions of India, temperate from their elevation, have many trees and shrubs in store for Britain, is rendered certain by the excellent *Illustrations of the Botany, &c., of the Himalayan Mountains* of Mr. Royle. That gentleman, who joins to the acuteness of the scientific botanist, the extended views of the general observer, and the knowledge of both Indian and European cultivation, considers that all the plants of regions in India elevated not less than 7000 ft. above the level of the sea, if not found in valleys, are likely to prove hardy in Britain. The following list is of species considered likely to be in this condition, or, at least, most of them. It has been prepared chiefly from Royle's *Illustrations*; but, in cases to which Mr. Royle's work has not yet been extended, or that do not come within its scope, from other sources; these are, Dr. Wallich's manuscript catalogue, to which Mr. Royle has both given the access and indicated the species likely to be fittest for selection in, and our *Hortus Britannicus*. In this list, all those names to which a star (*) is prefixed are, as in preceding lists, supposed not to be indigenous; the dagger (†), before a specific name, indicates that a species of that name from India has already been introduced into Britain; and the point of interrogation (?), put before a specific name, implies a doubt as to the species being capable of enduring the open air in this country.

Ranunculàcææ. Clématis globòsa, † montàna (*Gard. Mag.*, vol. x. p. 564.), † nepalénsis, pubéscens, ritifòlia, Buchananiàna.

Magnoliàcææ. Mr. Royle has adverted to certain species being extant in Nepal that might, perhaps, live in the open air in Devonshire. Some of these are, *Mangliètia insígnis*; *Michèlia lanuginòsa, excélsa, Kisòpa, (?) Doltsòpa*.

Menispermàcææ. Cissámpelos obtécta, † hirsùta; Stauntònia latifòlia, angustifòlia, Brunoniàna; Sphærostèma grandifòrum, and other species. Cócclus (?) laurifòlius.

Berberàcææ. *Bérberis nepalénsis*, † aristàta, † asiàtica, † Wallichianà, † Coriària, kumanaurénsis, floribúnda.

Capparidàcææ. Cáparris (?) obovàta, (?) nepalénsis.

Pitosporàcææ. Senàcia † nepalénsis; Pittósporum eriocárpum.

Terstròmiàcææ. Eùrya acuminàta.

Aurantiàcææ. *Limònia Laurèola*.

Hypericàcææ. *Hypéricum* † cordifòlium, † pátulum, † uràlum, † oblongifòlium.

Aceràcææ. *Acer* † oblongum, cultràtum, caudàtum, sterculiàceum, villòsum; *Negúndo* *fraxinifòlium, † Dobinà'a (?) vulgàris.

Æsculàcææ. Pàvia índica.

Sapindàcææ § *Millingtonièæ*. *Millingtonia simplicifòlia, dilleniacfòlia, púngens*.

Meliàcææ. *Mèlia* † *Azedarách*, * *Buckàyun* (from Europe).

Vitàcææ. *Vitis* parvifòlia, cymòsa, obtécta, macrophýlla, capreolàta; *Ampelòpsis himalayàna*.

Zygophyllàcææ. *Meliánthus himalayànus*.

Rulàcææ. *Rùta* † albiflòra, † * angustifòlia.

Xanthoxylàcææ. *Xanthóxylum* hostile, alàtum.

Coriariàcææ. *Coriària nepalénsis* male, n. female.

Staphylàcææ. *Staphylèa Emòdi*.

Celastràcææ. *Euónymus* tíngens, † echinàtus, vágans, péndulus, frígidus, fimbriàtus, grandiflòrus, † japónicus, † Hamiltoniànus.

Aquifoliàcææ. *Ilex* dipyrèna, excélsa, serràta.

Rhamnàcææ. *Rhàmnus* † virgàtus, † rupéstris *Royle*, purpùreus; *Paliùrus* † virgatus; *Berchèmia flavéscens*; *Hovènia dúlcis*; *Ceanòthus flavéscens*.

Anacardiàcææ. *Rhùs* parviflòra, velutina, *kakrasíngæe*, † vernicífera (syn. *juglandifolia*), † *Bücku-Anèla*, † acuminàta; *Sàbia* parviflòra, campanulàta.

Leguminàcææ. *Caragàna Mooreroftiàna*, brevispina, *Gerardiàna*, polyacántha, spinosíssima, versicolor: one of these has been raised in Edinburgh, or the neighbourhood. *Astrágalus strobilíferus*, polyacánthus, *Grahamiànus*,

múlticeps; *Cytisus flaccidus*; *Colutea* † *nepalensis*; *Piptanthus* † *nepalensis*; *Edwardsia* (?) móllis; *Indigófera heterántha*, † *violæca*, *Gerardiána*; *Desmodium filiaefólium*, † *nütans*, *multiflórum*, *maculátum*, *sequax*; *Acácia* (?) móllis.

Saxifragæcæ. *Astilbe rivularis*.

Rosæcæ § *Chrysobalanæcæ*. *Prinsèpia utilis*.

Rosæcæ § *Potentillæcæ*. *Rubus rotundifólius*, *cóncolor*, *ásper*, *filiaëus*, *paniculatus*, † *pedunculósus*, *ferox*, † *ásper*, † *dístans*, † *micránthus*; *Potentilla rígida*.

Rosæcæ § *Spirææcæ*. *Spiræa callósa Thunb.*, or allied to it, † *chanædrifolia*, † *hypericifolia*, *Lindleyána*, *kanitscháica*, † *nütans*; *Kérria* † * *japónica*.

Rosæcæ § *Amygdalæcæ*. *Amygdalus* † * *commúnis*; *Pérsica* † *vulgáris*, † *lævis*, *saligna*; *Prúnus* † * *doméstica*, *bokhariénsis*, *Aloocha*, * *triflóra*, † *expánsa*; *Armeniaca himalénsis*; *Cérasus tomentósa*, *Púddum*, *undulata*, *capricida*, *cornuta*, *nepalénsis*.

Rosæcæ § *Rosæcæ*. *Rósa Lyéllü*, † *Brunónü*, † *tetrapétala*, *Webbiána*, † *macrophýlla*, *seríceca*, † * *damascèna*, † *microphýlla*, *moscháta* var. *nepalénsis*, *Bánsiæ* var. *múltiplex*.

Rosæcæ § *Póncæcæ*. *Pýrus commúnis*, cultivated varieties of, *sínica*, † *Páshü* (syn. *variolósa*, ? *índica Wall.*), † *lanata*, † *crenata* (syn. *vestita*), † *baccata*, *Máhus*, cultivated varieties of, † *stipulæca*, (*Sórbus*) *foliolósa*; *Cydónia* † * *vulgáris*; *Cotoneáster* † *frígida*, † *acuminata*, † *affinis*, † *microphýlla*; † *Nummularia* † *rotundifolia*; *Eriobótrya* † *elliptica*; *Photínia* † *integrifolia*, *dúbia*; *Cratægus gláuca*, † *crenulata*.

Granæcæ. *Púnica* † *Granátum*.

Tamaricæcæ. *Myricaria bracteata*, *élegans*.

Philadelphæcæ. *Philadelphus tomentosus*; *Deúzia staminea*, *Brunoniána*, *corymbósa*.

Passifloræcæ. *Passiflora* (?) *Leschenaúttü*, (?) *nepalénsis*.

Grossulæcæ. *Ribes* † *glaciæle*, *acuminatum*, *himalénse*, † *Cynósbatü*, *villósum*.

Araliæcæ. *Hédéra Hélix* and var. † *chrysocarpa*. Many other species of *Hédéra* inhabit India, but not any, or but few, of them are likely to thrive in the open air in Britain.

Caprifoliæcæ. *Caprifólium* † *confusum*, † *longifólium*, † *chinénse*, † *japónicum*; *Lonícera diversifolia*, *Webbiána*, *Govaniana*, *angustifolia*, *seríceca*, *obovata*, *elliptica*, *gláuca*, *depréssa*, † *Nylósteum*, *acuminata*, *glabrata*, *lanceolata*; *Abélia triflóra*; † *Leycesteria formosa*; *Sambucus adnata*; *Fibánum cotinifólium*, *Mulláha*, *punctatum*, (?) *erubescens*, *nervosum*, *cordifólium*, *grandifórum*; *Hydránga altissima*, *vestita* (*áspera Don*), *heteromalla*, *Adámia*, † (?) *cyána*.

Cornæcæ. *Córnus* † *oblónca*, *macrophýlla*, *nervosa*; *Benthámia* † *fragifera*.

Vacciniæcæ. *Vaccínium* (*Thibaúdia*) *Sprengéliü*; *Thibaúdia variegata*, *setigera*; *Gaylussáccia* (*Thibaúdia*) *serrata*; *Cavendishia nobilis*.

Myrsinæcæ. *Mýrsine* † *bifaria*, † *semiserrata*, *acuminata*.

Ericæcæ. *Rhododéndron* † *arboreum*, *aristatum*, † *barbatum*, † *cinnamómeum* † *campanulatum*, *formosum*, *lepidotum*, † *anthopogon*, † *setosum*; *Andrómeda fastigiata*, *formosa*, *lanceolata*, *villósa*, *ovalifolia*, *cordata*; *Gaulthéria nummularioides*, *trichophýlla*.

Ebenæcæ § *Styræcæ*. *Sýmplocos racemosa*, *paniculata*.

Oleæcæ. *Olea ferruginea*, *compácta*, *grandiflóra*, *robústa*; *Syrínga Emòdi*; *O'rnus floribúnda*; *Fráxinus xanthoxylóides*.

Jasminæcæ. *Jasminum* † *officinále*, *dispérmum*, *revolútum*, † *pubigerum*, (?) *glandulosum*, (?) *chrysánthemum*, (?) *chrysánthemóides*, (?) *nánum*, (?) *humile*.

Labiæcæ. *Róylea* † *élegans*.

Solanæcæ. *Lýcium* † *europæum*.

Lauræcæ. *Laúrus* (?) *odoratissima*.

Thymelæcææ. *Dáphne* † *cannábina*, *Bhólíca*, *seríceæ* *Don* (syn. *salicifólia Wal.*), *viridiflóra*, *mucronáta*.

Santaláccææ. *Osýris nepalénsis*.

Elaeagnáccææ. *Elæágnus* † *arbórea*, *armáta*; *Hippóphæ* † *salicifólia* (syn. *conférta*).

Euphorbiáccææ. *Búxus emarginatus*; *Pachysándra* † *coriácea*.

Urticáccææ. *Mórus* † * *nigra*, sp. white-fruited, *atropurpúrea*, † *índica*?, † *tatárica*, *parvifólia*, *serráta* (syn. *heterophýlla*), *lævigáta víridis*, † * *mauritiána*, * *scándens*. Some of these names are to be regarded as not established, and as more or less likely to be synonymous with others amongst them. *Broussonétia integrifólia*.

Ulmáccææ. *Ulmus* † *erósa* (syn. *effúsa W.*), † *integrifólia*, *virgáta*. These names are from Dr. Wallich's catalogue. In Mr. Royle's catalogue are three others, but Mr. Royle has advised that they may represent the same species. *Céltis* † *orientális*, (?) *elongáta*, *politória*, *tetráandra*, (?) *cinnanómea*, (?) *móllis*.

Juglandáccææ. *Júglans* † *régia*; *Engelbárdtia Roxburghiána*, *Colebrookiana*.

Salicáccææ. *Sálix Lindleyána*, *obováta*, *polyándra*, † * *babylónica*, (?) *ægyptiáca*, *élegans*, *grísea*, *kamaunénsis*, *erios Stéphya*, *pyrína*. The first five of these names are derived from Mr. Royle's catalogue and work. The rest are from Dr. Wallich's catalogue. In Mr. Royle's catalogue are six other names, but Mr. Royle has told us that they may represent the same species as certain of the names above. *Pópulus cordáta*, *acumináta*, *pyrifórmis*, *ciliáta*, sp.

Cupulíferææ. *Quércus annuláta*, † *dealbáta*, *incána*, *polyántha*, *lanáta*, *floribúnda*, *laxiflóra*, *dentósa*, *semecarpifólia*, *lamellósa*, *lappácea*, *spicáta*, *fenestráta*, *dilatáta*, *lanceafólia*, † *lanuginósa*, † *Phulláta*. These names are from Dr. Wallich's catalogue, and are exclusive of several others that are those of species which Mr. Royle deems likely to be too tender to thrive in Britain. Mr. Royle has in his own catalogue names, distinct from Dr. Wallich's, of seven kinds, of which some may be identical with, some distinct from, those represented by Dr. Wallich's names. *Castánea índica*, *tribulóides*, (?) *microcárpa*, (?) *sphærocárpa*. *Córylus lácera*, ? *fèrox*; in Mr. Royle's catalogue are the names *cashmeriénsis* and *scabérrima*, which may represent the same kinds as the preceding, or distinct ones. *Cárpinus vimínea*, *fágrinea*.

Betuláccææ. *Bétula Bhojpátra* (syn. *útilis*), *acumináta*, *cylindrostéphya*, *nítida*; and, in Mr. Royle's catalogue, three other names of as many kinds, that may be identical with three of those above, or distinct. *Alnus* (?) *nepalénsis*.

Platanáccææ. *Plátanus* † *orientális*.

Myricáccææ. *Myrica sápidá*.

Taxáccææ. *Táxus baccáta*?, † *nuéfífera*?

Coníferææ. * *Píceá Webbiána* (syn. *Abies spectábilis*), *dumósa* (syn. *Brunoniána*); *Pinus* † *excélsa*, † *Gerardiána* (syn. *Nedza*), † *Smithiána* (syn. *Morínda*), *Cèdrus* † *Deodára*; *Cupréssus* † * *sempervirens*, † *torulósa*; *Ephedra Gerardiána*; *Juníperus squamáta*, † * *chinénsis*, * *dimórpha*, † *recurva*, *religiósa*, † *excélsa*, *commúnis*?; *Thúja* † * *orientális*, † *nepalénsis*.

Whoever wishes to become acquainted with the ligneous flora of Himalaya should have recourse to Mr. Royle's *Illustrations*; it will be found to be one of the most scientific and comprehensive works of the kind that have ever been published; embracing, not only the scientific botany and natural history, including the geology, of the district, but the geographical distribution of species and their properties and uses: the whole being generalised with reference to Lower India and Europe.

Of the trees and shrubs of temperate climates, which have been introduced into the different countries of Asia, we profess to know but little. Not many foreign ligneous plants, we suppose, have been added to the ligneous flora of Siberia or Tartary: but the hardy fruit trees and fruit shrubs of Europe have been cultivated for a number of years in the gardens of some of the native princes of India; and many of our ornamental trees and shrubs have been transported to the gardens of our public officers in that country,

and to the botanic gardens established by government. This might be done to a great extent, as Mr. Royle has shown, in the mountainous districts of Himalaya, as well as in other hilly and mountainous regions, both of India and China.

Of that part of the ligneous flora of China which is hardy, very little is known. The following list of some of the ligneous species which inhabit China, and of a few of those which inhabit Japan, has been prepared from these three sources:— 1. Enumeratio Plantarum quas in China boreali collegit Dr. Al. Bunge, anno 1831: this enumeration is published in the *Mémoires présentés à l'Académie des Sciences de St. Petersbourg*, tome ii. livra. 1. et 2., 1833. 2. From Royle's *Illustrations*, in which a sketch of the climate and plants of China is presented, relatively to comparison with the climate and plants of the Himalaya. 3. From our *Hortus Britannicus*. The sign (?) denotes, in this list, as well as in the preceding, doubt of hardihood of the species to which it is applied.

Ranunculacææ. *Clématis intricata*, *Pæonia Moultan* and varieties of it.

Magnoliacææ. *Magnolia conspícua*.

Menispermæcææ. *Menispermum dauricum*, *Stauntonia*, (?) *Kadsura japonica*.

Berberacææ. *Berberis sinensis*.

Sterculiacææ. *Sterculia pyrifórmis*.

Tiliacææ. *Tilia*, (?) *Gréwia parviflora*.

Ternstroemiæcææ. *Thèa viridis*, *Caméllia* (?) japonica.

Hypericacææ. *Hypéricum patulum*, (?) *Ochranthe pallida*.

Accracææ. *Acer truncatum*, palmatum (Japan).

Æsculacææ. *Æsculus chinensis*.

Sapindacææ. *Xanthóceras sorbifolia*, *Kölreutéria paniculata*.

Meliacææ. *Mélia*.

Vitacææ. *Vitis vinifera*, *bryoniæfolia*, *ficifolia*, *humulifolia*, *serianæfolia*, *aconitifolia*; *Ampelópsis*.

Xanthoxylacææ. *Ailantus glandulosa*; *Xanthoxylum* (?) *Avicennæ*, (?) *nitidum*.

Staphyleacææ. *Staphylèa*.

Celastracææ. *Celástrus articulatus*; *Euónymus micranthus*, chinensis.

Aquifoliacææ. *Ilex*, a sp. of, allied to *Aquifolium*.

Rhamnacææ. *Rhámnus Thecæans*; *Zizyphus vulgaris* 1 spinosa, vulgaris 2 incrmis, parvifolia, globosa, (?) sinensis, (?) álbens, (?) nítida: (?) *Hovènia dulcis*.

Anacardiæcææ. (?) *Pistàcia chinensis*; *Rhús Cótinus*, *ailantóides*, (?) *vernicifera* (Japan), (?) *succedanea*.

Leguminacææ. *Sophora japonica* (China and Japan), chinensis; *Indigófera micrantha*, *macrostachya*; *Caragána Chamlágu*, *microphýlla*; *Lespedèza macrocarpa*; *Wistària Consequana*; *Gledítschia chinensis*, *heterophýlla*; *Cércis chinensis*; *Acácia* (?) *macrophýlla*, (?) *Nèmu*.

Rosacææ § *Amygdalææ.* *Amýgdalus communis*, pedunculata, p. múltiplex, p. polýgyna; *Pérsica vulgaris*; *Cérasus chinensis* 2 pluripétala (or flore pleno, syn. *Amýgdalus pumila*), japonica (Japan), j. múltiplex (Japan), serrulata, *salícina*, *Pseudo-Cérasus*; *Prúnus doméstica*?, *trichocarpa*, pauciflora, humilis 1 glabrata, humilis 2 villósula; *Armeniaca vulgaris*.

Rosacææ § *Spiræacææ.* *Spiræa triloba*, dasyantha, *sorbifolia*, callosa; *Kérria japonica pluripétala* (or flore pleno) (Japan).

Rosacææ § *Potentillææ.* *Rúbus purpureus*, *cratægifolius*, (?) *parvifolius*, (?) *reflexus*.

Rosacææ § *Rosææ.* *Ròsa índica*, odorata, longifolia, *Roxbúrhii*, flavescens, nívea, semperflórens, *Lawrenceana*, multiflora, *Grevillei*, *sinica*, *Bánksiæ*, B. flore líteo, microcarpa, bracteata, b. scabricaulis, *pimpinellifolia*, rugosa.

Rosacææ § *Pómææ.* *Pyrus floribunda*, dioica, spectabilis, *betulæfolia*; *Cydònia sinensis*, japonica; *Eriobótrya japonica* (Japan); *Photínia serrulata*; *Cratægus pinnatifida*.

Calycanthaceæ. *Chimonanthus fragrans* (Japan), *f. luteus* (Japan), *f. grandiflorus*.

Granatæcæ. *Punica Granatum*, and the white-flowered and pluripetalous varieties.

Tamaricæcæ. *Tamarix juniperina*, *chinensis*, (?) *indica*.

Philadelphicæcæ. *Deutzia grandiflora*, *parviflora*, *scabra* (Japan).

Grossulacæcæ. *Ribes Cynosbati*.

Hamamelidæcæ. *Hamamelis chinensis*.

Cornacæcæ. *Aucuba japonica* (China and Japan).

Caprifoliæcæ. *Caprifolium chinense*, *longiflorum*, *Loniceræ flexuosa*, *Sambucus racemosa*, *Viburnum* (?) *fragrans*; *Abelia* (?) *chinensis*, (?) *uniflora*; *Hydrangea Hortensia*.

Ericæcæ. *Rhododendron Farreræ*, (?) *leucanthum*; *Azalea* (?) *macrantha*, (?) *Andromeda*.

Ericæcæ § *Vacciniæcæ.* *Vaccinium* (?) *formosum*.

Ebenæcæ. *Diospyros Lötus*, *Schi-tse*.

Oleacæcæ. *Olea*, *Ligustrum lucidum* 1 *floribundum*; *Syringa chinensis*; *Orynus floribunda*.

Jasminæcæ. *Jasminum* (?) *angulare*, *floridum*.

Asclepiadæcæ. *Periploca* (?) *scipium*.

Bignoniæcæ. *Catalpa syringæfolia*.

Thymelæcæ. *Daphne cannabina*, (?) *Passerina Chamædaphne*.

Solanæcæ. *Lycium chinense*, *turbinatum*, *Trewianum*.

Euphorbiæcæ. (?) *Phyllanthus ramiflorus*, *Andrachne chinensis*.

Urticæcæ. *Morus alba* and varieties, *sinensis*, *constantinopolitana*; *Broussonetia papyrifera*.

Ulmæcæ. *Ulmus pumila*, *Celtis chinensis*.

Juglandæcæ. *Juglans regia*.

Salicæcæ. *Salix babylonica*, *Populus*.

Betulæcæ. *Betula*.

Cupuliferæ. *Quercus densifolia*, *chinensis*, and three other species; *Castanea vesca*, the large-leaved chestnut, dwarf chestnuts.

Coniferæ. Pines, fir, larch; *Pinus chinensis*, *Massoniæna*; *Cunninghæmia lanceolata*; *Thuja orientalis*; *Juniperus chinensis*, *glauca*; *Cupressus*, *Salisburia adiantifolia* (Japan).

The northern provinces of China, Mr. Royle observes, are more European in their flora than any parts of the plains of India; and the flora of the mountains has an almost universal identity of genera with that found covering the elevated belt of Himalaya. From these and other remarks we conclude that many species of trees and shrubs in China, now wholly unknown to us, will at some future time be added to the British arboretum.

The Chinese, through the European residents at Canton and other seaports, have become possessed of various of our ornamental ligneous plants. Mr. Reeves (*Gard. Mag.*, vol. xi. p. 437.) mentions that *Magnolia grandiflora* was introduced at Macao by Mr. Livingston, previously to 1830; and *M. acuminata*, *glauca*, and *tripetala*, soon afterwards. The recent discovery of the tea shrub in the province of Assam, through an extent of territory which occupied a month's journey, shows how little of the ligneous flora of that part of the world is yet known. (See *Dr. Wallich*, in *Gard. Mag.*, vol. xi. p. 429.)

SECT. II. *Of the Indigenous and Foreign Trees and Shrubs of Africa.*

THE number of ligneous species which the British arboretum has obtained from Africa, including the Canary Isles, appears to be 23; a number larger than might be expected, considering the tropical situation of this part of the world, and that its mountains are less elevated than those of Asia. Barbary has supplied 13 of these 23 species; because, being situated on the shores of the Mediterranean, its climate is comparatively temperate. More may, per-

haps, be received from the interior of the country, and from the African islands; but, considering that the floras of these islands, and of Egypt and Southern Africa, have been pretty fully explored, our hopes of further additions, fit to endure our climate, are not very sanguine.

The trees and shrubs of temperate climates introduced into Africa must necessarily be very few; and till lately they were limited, perhaps, to a few shrubs in the gardens of the British consuls. Since the introduction of European improvements into Egypt, however, the pacha has established an English garden under the care of an English gardener, Mr. Traill, who is endeavouring to acclimatise the plants and trees both of temperate and tropical climates. Algiers, which came into possession of the French in 1830, is receiving from that nation of naturalists many European plants; as appears in detail in the *Annales de la Société d'Horticulture de Paris* for 1831, and in the *Gardener's Magazine*, vol. xi, p. 632. A nursery has been established by the French authorities, which is said to contain 25,000 trees, bushes, and plants, for the purpose of experiment and naturalisation. It occupies 80 acres, and is under the care of a director and twenty men. Such an establishment may be referred to as one worthy of imitation in colonising a new country.

SECT. III. *Of the Indigenous and Foreign Trees and Shrubs of America.*

By far the greatest and most interesting accessions to the British arbo-retum have been received from North America; but, as some hardy species have also been received from the southern division of that immense country, we shall devote a subsection to each.

SUBSECT. I. *Of the Indigenous and Foreign Trees and Shrubs of North America.*

THE introduction of woody plants from North America into Britain may be said to have commenced with the missionaries sent out by Compton, Bishop of London, about the end of the seventeenth century, and to have continued without interruption ever since. Some species were, doubtless, introduced by Sir Walter Raleigh and others; but the practice of sending out collectors to send home objects of natural history undoubtedly began about the period we have mentioned. We have seen, in preceding parts of this history, that Bannister, Catesby, Garden, John and William Bartram, André Michaux, Fraser, Lyon, and Douglas are the names of the collectors to whom we are chiefly indebted: and that Compton, the Duke of Argyll, Lord Petre, the Duke of Richmond, Ellis, Dr. Uvedale, Dr. Fothergill, and, above all, that most excellent man Peter Collinson, a quaker and linendraper, were the principal amateurs. These gentlemen, and Gray, Gordon, and other nurserymen, in Britain, and Du Hamel, Lemonnier, and Maréchal de Noailles, in France, were the principal persons who encouraged the collectors. Much, also, is due to those American and European authors who have explored the interior of the civilised portion of America, and published the result of their labours. From the *Flora* of Pursh, edit. 1814, we have made the following enumeration of the woody plants of North America not indigenous to Britain.

Ranunculaceæ. *Atrégene americana*; *Clematis virginica*, cordata, holose-ricea *Walteri*, crispa, reticulata, *Viorna*, *Catesbyana*; *Xanthoxiliza apifolia*.

Winteraceæ. *Illicium floridanum*, parviflorum.

Magnoliaceæ. *Magnolia grandiflora* elliptica, grandiflora obovata, grandiflora lanceolata, glauca, longifolia, macrophylla, tripétala, acuminata, cordata, auriculata, pyramidata; *Liriodendron Tulipifera*, T. var. obtusiloba.

Anonaceæ. *Asimina triloba*, parviflora, pygmaea, grandiflora.

Menispermaceæ. *Menispermum canadense*, c. var. lobatum; *Cocculus carolinus*, *Schizandra coccinea*.

Berberidaceæ. *Berberis canadensis*, *Mahonia*, *Aquifolium*, nervosa.

Cistaceæ. *Hudsonia cricoides*.

Malvaceæ. *Hibiscus Mânihot.*

Tiliaceæ. *Tilia glabra*, laxiflora, pubescens, p. var. leptophylla, heterophylla.

Terströmiaceæ. *Gordonia Lasiánthus*, pubescens, *Stuártia virginica*, Malachodéndron ovátum.

Hypericææ. *A'scyrum Crux A'ndrea*, hypericoides, amplexicaule; *Hypéricum Kalmiánum*, frondosum, amœnum, prolificum, nudiflorum, glaucum, densiflorum, galioides, aspalathoides, fasciculatum, tenuifolium.

Acerinææ. *A'cer rubrum*, dasycarpum, barbatum, saccharinum, nigrum, macrophyllum, circumdatum, striatum, montanum; *Negúndo fraxinifolium*.

Hippocastanææ. *Æ'sculus glabra*, pallida; *Pavía*, rubra, hybrida, flava, macrostachya.

Sapindææ. *Sapíndus* ? * *Saponária.*

Vit s. *Vitis Labrusca*, æstivalis, æ. var. sinuata, cordifolia, riparia, rotundifolia, palmata; *Ampelopsis cordata*, *hederacea*, hirsuta, bipinnata.

Xanthoxyleæ. *Xanthoxylum fraxineum*, tricarpum; *Ptelea trifoliata*, t. var. pubescens.

Staphyleææ. *Staphylèa trifolia.*

Celastrinææ. *Celástrus scandens*, bullatus; *Euonymus americanus*, angustifolius, atropurpureus.

Ilicinææ. *Ilex opaca*, laxiflora, *Cassine*, *Dahoón*, angustifolia, a. var. ligustrifolia, vomitoria, prinoides, canadensis, *Myrsinites*; *Prinos verticillatus*, ambiguus, lævigatus, lanceolatus, glaber, coriaceus, c. var. angustifolius.

Rhamnææ. *Rhámnus franguloides*, alnifolius, carolinianus, lanceolatus, minutiflorus; *Ceanothus americanus*, intermedius, sanguineus, microphyllus; *Berchémia volubilis.*

Anacardiæææ. *Rhus typhina*, glabra, elegans, viridiflora, pumila, vernix, copallina, Toxicodéndron, radicans, r. var. microcarpa, aromatica.

Leguminosææ. *Robínia Pseud-Acacia*, viscosa, hispida, macrophylla; *Wistária frutescens*; *Cássia occidentalis*, ligustrina; *Cercis canadensis*, c. var. pubescens; *Virgília lutea*, *Cytisus rhombifolius*; *Amórpha fruticosa*, f. var. emarginata, f. var. angustifolia, microphylla, pubescens, canescens; *Gleditschia triacanthos*, t. var. inermis, brachycarpa, monospermia; *Gymnocladus canadensis.*

Chrysobalanææ. *Chrysobalanus oblongifolius.*

Amygdalæææ. *Cerasus virginiana*, serotina, canadensis, caroliniana, semperflorens, borealis, pennsylvanica, nigra, hyemalis, pygmaea, pubescens, pumila, depressa, chicensa; *Prunus maritima*, domestica var. myrobalana.

Rosæææ. *Rosa parviflora*, nitida, lucida, gemella, *Lyónü*, carolina, rubifolia, lævigata, suavolens, pendulina, lutescens; *Rubus villosus*, strigosus, canadensis, cuneifolius, occidentalis, hispidus, trivialis, flagellaris, inermis, spectabilis, odoratus; *Spiræa salicifolia*, s. var. latifolia, tomentosa, hypericifolia, chamædrifolia, c. var. media, betulafolia, opulifolia, capitata, discolor, sorbifolia; *Púrshia tridentata*, *Potentilla floribunda.*

Pomæææ. *Pyrus coronaria*, angustifolia, microcarpa, americana, arbutifolia, melanocarpa; *Amelánchier Botryápium*, ovalis, sanguinea; *Cratægus apiifolia*, spathulata, turbinata, coccinea, populifolia, pyrifolia, elliptica, glandulosa, flava, parvifolia, punctata rubra, punctata aurea, Crús-gállí splendens, Crús-gállí pyracanthæfolia, Crús-gállí salicifolia.

Calycanthææ. *Calycánthus floridus*, glaucus, lævigatus.

Philadelphæææ. *Philadèlphus inodorus*, *Lewisü*, grandiflorus; *Decumária* bábara, sarmentosa.

Passiflorææ. *Passiflora peltata.*

Cactææ. *Opúntia vulgaris*, *Mammillária vivipara.*

Grossulæææ. *Ribes albinervium*, trifidum, rigens, prostratum, resinosum, viscosissimum, sanguineum, *Menzièsü*, aureum, recurvatum, floridum, laxiflorum, rotundifolium, hirtellum, gracile, triflorum, oxyacanthoides, lacustris, *Cynósati*, speciosum

Araliæææ. *Aràlia hispida*, spinosa, s. var. inermis.

Caprifoliàceæ. *Caprifolium* sempervirens, ciliòsum, flavum, gratum, parviflorum [? dioicium]; *Lonicera* ciliata, ciliata alba, villòsa; *Symphoria* glomerata, racemòsa; *Diervilla* lutea; *Linnaea* borealis; *Viburnum* prunifolium, pyrifolium, Lentago, nudum, obovatum, obovatum panicafolium, *cassinoides*, levigatum, nitidum, dentatum, pubescens, lantanoides, acerifolium, molle, *Oxycoccus*, edule; *Sambucus* canadensis, pubescens; *Hydrangea* arborescens, cordata, nivea, quercifolia.

Córnea. *Cornus* florida, circinata, sericea, asperifolia, stricta, alba, paniculata, paniculata alba, paniculata radiata, alternifolia.

Loranthàceæ. *Viscum* ? flavescens.

Cinchonàceæ. *Pinckneya* pubens, *Chiococca* racemosa, *Cephalanthus* occidentalis.

Compositæ. *Baccharis* angustifolia, glomeruliflora, halimifolia; *Bupthalmum* frutescens; *Tva* imbricata, frutescens.

Vaccinàceæ. *Vaccinium* stamineum, album, diffusum, dumosum, frondosum, frondosum lanceolatum, pallidum, resinosum viridescens, resinosum rubescens, resinosum lutescens, corymbosum, amenum, virgatum, fuscatum, fuscatum angustifolium, *galicans*, *ligustrinum*, tenellum, angustifolium, caespitosum, myrtifolium, crassifolium, nitidum, *Myrsinites*, *M.* lanceolatus, *M.* obtusus, *buxifolium*, ovatum, obtusum; *Oxycoccus* macrocarpus, erythrocarpus, hispidulus.

Ericàceæ. *Andromeda* tetragona, *lypnoides*, *polifolia* angustifolia, *polifolia* latifolia, calyculata, angustifolia, coriacea, axillaris, axillaris longifolia, *Catsbaei*, acuminata, floribunda, mariana, mariana angustifolia, speciosa, speciosa pulverulenta, racemosa, arborea; *Lyonia* ferruginea, rigida, paniculata, frondosa; *Clèthra* alnifolia, tomentosa, scabra, paniculata, acuminata; *Mylocaryum* *ligustrinum*, *Cyrilla* caroliniana; *Arbutus* laurifolia, *Menziesii*, tomentosa; *Gaultheria* procumbens, *Shallon*; *Menziesia* ferruginea, globularis, cymetiformis, cerulea; *Kalmia* hirsuta, glauca, glauca rosmarinifolia, cuneata, angustifolia, angustifolia ovata, latifolia; *Epigaea* repens; *Rhodora* canadensis; *Rhododendron* maximum roseum, m. album, m. purpureum, punctatum, catawbiense; *Azalea* calendulacea flammca, c. crocea, canescens, multiflora coccinea, n. rutilans, n. carnea, n. alba, n. papilionacea, n. partita, n. polyandra, arborescens, bicolor, viscosa, nitida, glauca, hispida; *Ledum* palustre, palustre decumbens, latifolium; *Ammýrsine* *buxifolia*; *Bejaria* racemosa, *Ptea* virginica, *Pyxidanthera* barbulata.

Symplocinèæ. *Symplocos* tinctoria.

Styràceæ. *Styrax* grandifolium, pulverulentum, laevigatum; *Halèsia* tetraptera, díptera.

Sapotèæ. *Bumelia* lycioides, reclinata, lanuginosa, chrysophylloides, serrata.

Ebenàceæ. *Diospyros* virginiana, pubescens.

Oleaèceæ. *Olea* americana; *Chionanthus* virginica, maritima; *Oleus* americana; *Praxinus* sambucifolia, quadrangulata, epiptera, acuminata, caroliniana, platycarpa, pubescens, p. longifolia, p. latifolia, p. subpubescens, juglandifolia; *Catalpa* *springefolia*.

Apocýneæ. *Echites* difformis, *Gelsèmium* sempervirens.

Bignoniàceæ. *Bignonia* crucifera, capreolata; *Tecomia* radicans flammca, radicans coccinea.

Solanèæ. *Lycium* carolinianum.

Labiàtæ. *Salvia* coccinea.

Verbenàceæ. *Callicarpa* americana.

Che nopòdeæ. *Atriplex* *Halimus*, *Diotis* lanata.

Polygonèæ. *Calligonum* canescens.

Laurinèæ. *Laurus* *Catesbyana*, carolinensis glabra, c. pubescens, c. obtusa, *Benzoin*, *Diospyros*, geniculata, *Sassafras*.

Thymelàceæ. *Direa* palustris.

Santalàceæ. *Nyssa* villosa, biflora, candicans, tomentosa, denticulata; *Hamiltonia* oleifera.

Elcàgneæ. *Elcagnus* argentea; *Shepherdia* canadensis, argentea.

Aristolochiææ. *Aristolochia sípho*, tomentosa.

Euphorbiæææ. *Bòrya* porulosa, ligústrina, acuminàta; *Stillíngia ligústrina*, sebifera.

Urticææ. *Mòrus rùbra*.

Ulmæææ. *Ulmus* americana, pèndula, fùlva, alàta; *Plánera* Richárdi, Gmelini; *Céltis occidentális*, crassifolia, pùmila.

Juglândææ. *Juglans* nigra, cinèrea; *Carya* olivæfórmis, sulcàta, álba, tomentosa, amàra, porèina ficifórmis, oboordàta, aquática, myristicæfórmis.

Salicinææ. *Sàlix* cándida, Muhlenbergiana, trístis, recurvata, vestita, Uva úrsi, cordifolia, obovata, planifolia, pedicellaris, fuscata, confèra, myricoides, prinoides, discolor, angustata, longifolia, Houstoniana, falcata, nigra, lucida, rigida, cordata, grisea, ambigua; *Pópulus* balsamifera, cándicans, trépida, monilifera, betulafolia, grandidentata, lævigata, angulata, heterophylla.

Betulínææ. *Bétula* populifolia, excelsa, nigra, papyræa, lénta, pùmila, glandulosa; *Alnus* crispa, serrulata, glauca.

Cupulifercæ. *Quercus* Phellos, Phellos humilis, maritima, sericea, myrtifolia, virens, cinèrea, imbricaria, laurifolia, laurifolia obtusa, agrifolia, heterophylla, aquatica, hemisphaerica, nana, triloba, nigra, tinctoria, discolor, coccinea, ambigua, rubra, Catesbaei, falcata, palustris, Banisteri, obtusiloba, macrocarpa, olivæfórmis, lyrata, álba, álba repanda, *Prinus*, bicolor, montana, Castanea, prinoides; *Castanea* vesca americana, pùmila; *Fágus* ferruginea; *Corylus* americana, rostrata; *Carpinus* americana, *Ostrya* virginica.

Platánææ. *Liquidambar styraciflua*.

Myricææ. *Myrica* cerifera, cerifera pùmila, carolinensis, pennsylvanica; *Comptonia asplenifolia*.

Hamamelidææ. *Hamamelis* virginica, macrophylla; *Fothergilla anifolia*, major, Gardèni.

Conifercæ. *Pinus* inops, resinosa, Banksiana, variabilis, rigida, serotina, pungens, Tæda, palustris, Stròbus; *Abies* balsamifera, Fraseri, taxifolia, canadensis, nigra, rubra, álba; *Larix* pèndula, microcarpa; *Taxodium distichum*; *Cuprèssus thyoides*, *Thùja* occidentalis; *Juniperus* communis deprèssa, virginiana, *Sabina* procumbens, excelsa, barbadensis; *Táxus* baccata.

Cycadææ. *Zamia integrifolia*.

Empétrææ. *Empetrum nigrum*, *Ceratiola ericoides*.

Smilacææ. *Smilax* hastata, hastata lanceolata, bona-nox, quadrangulàris Wálteri, *Sarsaparilla*, ovata, álba, lanceolata, pùbera, Pseudo-china, rotundifolia, caduca, laurifolia, pandurata, circidifolia.

Palmææ. *Sabal* Adansonii; *Chamærops* serrulata, hýstrix, *Palmétto*.

The number of trees and shrubs in the British arboretum, received from North America, is considered to be 528, and they comprise the greater part of the names in the foregoing enumeration. Still, as it is not always certain that the same names in our catalogue are applied to the same things, there may be a number of species described by Pursh which are not yet introduced into Britain. At all events, we have little doubt that, in the unexplored parts of North America, there are many species that will, at no distant day, find way to Europe.

The greater part of the trees and shrubs of Europe, which are remarkable either for beauty or utility, appear in the catalogues of the American nurserymen, more particularly in those of Prince of New York, and of Carr, the successor of Bartram, near Philadelphia. From a MS. which has been kindly sent to us by Dr. Mease, containing the dates of the introductions of a number of European trees into America, we find that some took place as early as the settlers there from this country; and that the introduction of European trees was in an especial degree accelerated by the establishment of tree nurseries. William Hamilton, Esq., of the Woodlands, near Philadelphia, is stated by all the collectors of plants in America, during the last century, to have had the most complete garden in the United States. It is said to have contained not only all the plants of America, but those of Europe and other parts of the world, which were considered of interest either for arts or medicine. In

Bartram's Botanic Garden there appears to have been the best collection in any nursery; and probably, at present, it is only equalled by that of Prince of New York. No nursery in America is superior to Bartram's for fine specimens of trees. The dimensions of some of these, with those of many others, of which accounts have been sent us, will be given when treating of each particular tree, in the third part of this work. (See also *Gard. Mag.*, vol. viii. p. 272.)

Part of the United States, and the Canadas, were visited by two excellent arboricultural observers; Mr. Robert Brown, formerly a nurseryman at Perth, and Mr. James Macnab, the son of the curator of the Edinburgh Botanic Garden, in the autumn of the year 1834; and an interesting notice of the distribution of different species of trees in the countries they passed through has been published in the *Quarterly Journal of Agriculture*, vol. v. p. 594., and will be found also in the twelfth volume of the *Gardener's Magazine*. Before landing at New York, the country appears to the stranger of a very dark and dismal hue, from the quantity of pines and red cedars which clothe the more conspicuous prominences; but, after landing, the whole, from the prevalence of fine trees and shrubs, appears like one vast garden. The stranger is strongly impressed with the beauty and number of trees, which are partly indigenous to the locality or the district, and partly introduced from more southern climates. The diversity of the forms of the trees, and the variety of their foliage, are most remarkable. No remains of ancient forests are observable, as might be supposed, these having been long since cut down for fuel; but forest trees of large size are frequently to be seen, covered to their summits with wild vines. Of these the *Plátanus occidentális*, *liriodendron*, *liquidambar*, *Gledítschia triacánthos*, and the *catalpa* are preeminent. It is worthy of remark, that almost the only foreign trees conspicuous in the artificial scenery of America are, various kinds of fruit trees, the Lombardy poplar, and the weeping willow. The contrast between the regular position and round-tufted heads of the fruit trees and the lance-shaped heads of the poplars, and between both these trees and the wild luxuriance of the indigenous species, is very striking. About 67 miles up the country, on the river Hudson, a limestone district occurs, and on this the lively green of the *arbor vitæ* succeeds to the dark hue of the red cedar. All the uncultivated parts of the surface are covered with this tree, of different sizes, varying from 1 ft. to 20 ft. in height, and always of a pyramidal shape. The woods on both sides of Lake Champlain are very various. The principal trees are, the wild cherry (*Cérasus virginiana*), elms, walnuts, sugar maples, and the aspen poplar. The rocky grounds abound with *arbor vitæ*, and the "appearance of the lofty white, or Weymouth, pine, towering above the deciduous trees, on rising grounds at the base of the hills, of a dark aspect and nearly destitute of branches, was remarkable." The northern extremity of Lake Champlain exhibits the same trees, with the addition of the balm of Gilead fir. The only tree worth notice on the St. Lawrence river was the canoe birch (*Bétula papyracea*). At Montreal our travellers were much "surprised to see the great difference which the Canadian winter produces upon those species of ornamental trees which grace the lawns and cities of the United States. As examples, may be mentioned the *Ailántus glandulosa*, the trees of which were quite small and stunted; *Maclúra aurantiaca* seemed barely alive; and the mulberries were small and unhealthy. The weeping willows here are almost always killed in winter, although in the neighbourhood of New York the stem of this tree is seen averaging from 8 ft. to 15 ft., and sometimes 20 ft. in girth. None of the *catalpas* and *magnolias*, which prove so ornamental in the pleasure-grounds both of New York and Philadelphia, can be made to live here, with the exception of the *M. glauca*, and it is in a very unhealthy condition. *Taxodium distichum* is also much dwarfed, and barely alive. Peaches in this part of the country do not succeed as standards; but several peach trees placed against garden walls possessed well ripened wood, and had every appearance of affording plentiful crops. The principal ornamental tree cultivated in this part of the country, on account of its beauty, is the *Robinia*

glutinosa, which, during the months of June, July, and August, bears a profusion of delicate pink flowers, and does not attain a large size."

In the different islands of Lake Ontario "the hemlock spruce is abundant and of great size, as well as arbor vitæ, walnuts, oaks, sugar maples, and elms." Near Toronto, on the shores of the lake, the weeping willow is healthy and luxuriant, and there are fine specimens of the locust tree, broad-leaved American beech, Canadian and Lombardy poplars, limes, oaks, ashes, elms, white pine, and hemlock spruce. *Pinus resinosa* (the red pine) was observed here for the first time; and it is by no means plentiful, having been found by our travellers only in this tract. The red birch they found a fine-shaped tree, with a trunk about 2 ft. in diameter, and a wide-spreading top like that of the beech tree in Europe. Near the Falls of Niagara is a "very extensive natural forest of sweet chestnuts; and what is very remarkable, the trees are placed at such regular distances that at first one would not hesitate to think that they had been planted by the hand of man. Not a great way from this, we observed a similar forest of large native oaks with precisely the same appearance of regular plantation: yet in both cases the arrangement was wholly the work of nature, the stronger individuals having probably smothered the weaker. In the neighbourhood of the falls, the trees were of very various descriptions, of great size, and more intermixed than we had hitherto seen. The tulip trees were of great height, with stems varying from 8 ft. to 12 ft. in circumference. Platanus trees, oaks, elms, limes, ashes, walnuts, beeches, poplars, and white pines, were all equally large and lofty. The hemlock spruce was scarcely seen, but the arbor vitæ seemed to take its place; for it is, without exception, the most abundant tree in the neighbourhood of the falls, very tall and sometimes tapering to the height of 60 ft. Here, again, the red cedar was observed, with great abundance of dwarf-growing yew (*Taxus canadensis*)."

Between Niagara and Hamilton was the only district in Canada where the *Laurus Sassafras* was seen; the trees were all small, though remarkably healthy. The great natural forests of the country presented chiefly oaks of great height; and, when the ground became in the least degree elevated, white pines bounded. On a flattened, low, moist meadow was an extensive forest of the tamarack, or black American larch (*Larix pendula*), long straggling trees with stems not exceeding 20 in. in circumference. Near New London the specimens of the trees, particularly of the platanus, were very large. Stems were measured of from 15 ft. to 20 ft. in girth, and many of the trees had straight trunks of from 10 ft. to 30 ft. high, before branching. "This tree is always seen largest and in greatest abundance along the moist banks of rivers, where the soil is deep and rich." The white pine, near New London, has a trunk varying from 13 ft. to 18 ft. in circumference; and some trees, which had been blown down, were measured, and found to average 160 ft. in length. The oaks here vary from 10 ft. to 15 ft. in circumference of trunk, with 45 ft. and 5 ft. of straight clear stems. Between New London and Goderich, a distance of 60 miles, the road passes through one continued dense forest. The trees were principally elms, averaging from 10 ft. to 25 ft. in circumference. Mixed with them were beeches, birches, and ashes of ordinary dimensions. On some low swampy ground, there was an extensive forest of hemlock spruce, and on an extended limestone ridge some splendid specimens of arbor vitæ. Horizontal sections of the white pines and hemlock spruce exhibited between 300 and 400 annual layers; oaks, 200; and elms, 300. On the whole, the neighbourhood of Goderich in Canada presented "a much greater and finer collection of large native trees than had before been seen; for, in addition to what have been named, were very large sugar maples, with splendid specimens of the black and white ash, limes, oaks, beeches, birches, cherries, with extensive tracts of balsam poplar (*Populus balsamifera*), and the black American larch, all growing in deep rich soil. On the banks of the Maitland river, many very noble specimens of platanus are seen, with stems varying from 18 ft. to 36 ft. in circumference. It is curious to observe, that

when this tree exceeds 6 ft. in diameter, it seldom has a clear hole of above 10 ft., after which it branches much. Almost the whole of the large trunks are hollow. Along the banks of the lake, extending both ways from Goderich, we observed white American spruces (*Abies álba*); but none of them exceeded 60 ft. in height, and 3 ft. in circumference.

Just before entering the St. Clair river from Lake Huron, lofty white pines are seen towering on both sides; but, on entering the river, they entirely disappear on the Canada side, although, on the United States side, they continue for many miles; and a river, which runs into the St. Clair on that side, has, from the quantity of pines found upon its banks, been named the White Pine River. No very large trees are seen on the banks of the St. Clair, but, on getting back into the country, oaks, elms, limes, and walnuts abound. On the banks of the river, all the way down, many dwarf kinds of shrubby plants adorn its edges. Of these, the different kinds of plums and cherries, with the sweet-scented crab apple, and a variety of thorns, form the chief objects; but what was most gratifying to us, was the great abundance of stag's-horn sumach (*Rhus typhina* var. *elegans*), now loaded with large heads of scarlet fruit.

Crossing Lake Erie to Cleveland, United States, we have, in addition to the ordinary forest scenery, some fine specimens of tulip trees, *Laurus Sassafras*, and cucumber trees (*Magnolia acuminata*). The stems of the latter did not exceed 2 ft. in circumference. Passing through the interior of the country to Pittsburg, the surface was very irregular. The principal native tree on this line of road was the beech, which was seen spread over an extended plain, containing many fine trees. There was also an extensive forest of the larch; and, in similar situations to those before mentioned, bordering all the mixed woods as we proceeded along, was the dogwood tree (*Cornus florida*). These trees, having taken on their autumnal hue (5th September, 1834), were rendered beautiful in consequence. Of this fine tree, none had been seen since leaving New York till now; and from this place it continued to be more or less plentiful, in the different routes which we took, till we reached New York again. It was much talked of by the inhabitants, and praised, on account of its profusion of large white blossoms early in spring, and its dark red leaves and scarlet fruit in the fall, or autumn. Before entering Pittsburg, the sloping wooded banks by the river side presented a rich appearance, from the quantities of rhododendrons, kalmias, azaleas and andromedas, which covered them as underwood. The large trees were much the same as those before mentioned, with the addition of the chestnut-leaved oak and red maple (*Acer rubrum*), of which last some few trees on the banks of the Ohio river, near Pittsburg, measured 12 ft. in circumference. There were likewise some large natural trees of the honey locust.

We left Pittsburg for Philadelphia by way of the Alleghany Mountains. On these the natural grouping of the trees and shrubs was most remarkable. On our first approach to these mountains, we observed the hemlock spruce, of various sizes, covering the banks, having the *Rhododendron catawbiense*, and *Kalmia latifolia* as underwood. Proceeding up the mountain, large tracts of sweet chestnuts are passed through. Above this, extensive tracts of oak; and on the highest summits, all over the mountains, were scraggy trees of the *Pinus rigida*, or pitch pine, with dwarf shrub oaks as underwood. On crossing the different summits, it was curious to see the same arrangement throughout on either side. Having remained for several days on these mountains, our attention was very much taken up with the great natural forests situated on the extended plains between the highest summits. When any one species of a tree is met with, acres of the same are generally seen together. In this way we continued to pass through successive masses of a sort of trees, of the various pines, magnolias, walnuts, poplars, and sour g^{rs}, or tupelo trees (*Nyssa villosa* and *sylvatica*), &c. All of these grow in deep rich soil, with the exception of the pitch pine, which was seen on the mountain tops; and, on poor rocky ground, at different elevations, throughout the mountains, the

Pinus inops, *pungens*, and *Tæ'da* are seen; and on more fertile sheltered situations we find the Canadian Judas tree (*Cercis canadensis*), *Laúrus Benzoin* and *Sassafras*, *Euónymus atro-purpureus*, with nettle trees, witch hazels, and mulberries; also a vast variety of other dwarf-growing trees and shrubs." (*Quart. Journ. of Ag.*, vol. v. p. 605.) These observations of Mr. Brown and Mr. Macnab are well calculated to give an arboriculturist an idea of what he might expect to meet with, in travelling in the northern states of North America. Those who have leisure to pursue the subject, and who wish to form an idea of the scenery of the southern states, may consult Bartram's *Travels*, already referred to p. 86.; Dwight's *Travels in New England*; and some interesting articles communicated by American correspondents to the *Magazine of Natural History*, more particularly in vol. vi. p. 97. and vol. viii. p. 529.

We include Mexico in North America, and have great hopes that some additions may be made to the British arboretum, from the mountainous regions of that extensive country. Some magnificent specimens of *Coniferae* are said to exist in it, particularly an evergreen *taxodium* of enormous height. Dr. Coulter, it is believed, has discovered several new species of pines and firs, which are likely to prove hardy in Britain, as are almost all the resinous trees of other countries.

SUBSECT. 2. *Of the Indigenous and Foreign Trees and Shrubs of South America.*

THE trees and shrubs furnished by South America to the British flora are only 22: nor can it be supposed that that number will be materially increased, the botany of the country having been very generally explored by different botanists. Our hopes are chiefly from the temperate zones of the loftier mountains, and from the shores of the sea and the larger rivers. Mr. Mathews, who has been several years in South America as a natural history collector, speaks of lofty mountain ranges covered with forests, which have never yet been penetrated by civilised man. (See *Gard. Mag.*, vol. xi. p. 636.) The ligneous vegetation, however, of South America appears to be much less varied than that of North America, and to consist chiefly, in the warmer parts of the country, of palms, and in the more temperate regions, of pines, firs, and other *Coniferae*.

With respect to the introduction into South America of trees and shrubs which are foreign to it, we find that the same tendency to equalisation of products has taken place here, as has been the case everywhere else where civilised man has established himself. European and North American trees and shrubs are to be found more or less in the government gardens of Rio de Janeiro, Buenos Ayres, the Caraccas, and other places, and in the gardens of old established European merchants and government officers. The tea shrubs of China, the mulberry, the mango of India, the peach of Persia, the pine-apple of Africa, properly a ligneous plant, and the apple of Europe, are all to be found in the neighbourhood of Rio.

SECT. IV. *Of the Indigenous and Foreign Trees and Shrubs of Australia and Polynesia.*

THE British arboretum has not yet received many additions from these extensive and little known regions; nor can it be conjectured whether they are likely to contain much that is suitable for our purpose, till something more is known respecting the height of the mountains in the interiors of the islands, especially of that of New Zealand. Very few ligneous plants have hitherto been introduced from the latter country; but, if, as there is reason to believe, the interior consists of a chain of mountains with their summits buried in perpetual snow, we may hope to reap a considerable harvest when they have been explored by botanists. Something also may reasonably be expected from Van

Diemen's Land, in addition to the species of eucalyptus, acacia, callistemon, and sida, which live as standards through the ordinary winters in the neighbourhood of London without protection.

A number of the ligneous trees and shrubs of Europe, and some also from other parts of the world, have been taken to New Holland and Van Diemen's Land, by the settlers; and every year packages of plants which stand the open air in England, as well as of house plants, are sent out by the nurserymen. In the botanic garden at Sydney there was in 1828, a collection, which included the fruit trees of every part of the world, as far as they could be collected, and also many of the principal European timber trees and flowering shrubs. An account of this collection by the then curator of the garden, Mr. Charles Fraser, will be found in the *Gardener's Magazine*, vol. v. p. 280. It is there stated that the European trees stood the extreme drought of the year 1827-28 better than those of warmer climates; and, while oranges, limes, shaddocks, guavas, &c., were completely burnt up, apples, pears, &c., stood the shock without any apparent injury. To give an idea of the "capabilities of the climate," Mr. Fraser states that, in "an exposed part of the garden, may be seen growing luxuriantly, in a dense thicket formed by themselves, the following trees; viz., the English ash and elm, *Erythrina Coralodéndrum* in full flower, *Bómbax heptaphýllum*, *Gymnócladus canadénsis*, *Ficus elástica*, *Dalbérgia Sissoo*, *Tectona gróndis*, *Pinus Pináster* and *halepénsis*, *Catálpa syringafólia*, the English lime and sycamore, the mossy-cupped and English oak, *Acácia tamariscína*, *Salisbúria adiantifólia*, the tea and olive, and many others."

The trees of Van Diemen's Land appear to be among the most gigantic of the whole world. Mr. James Backhouse, an English nurseryman who spent some time in Hobart Town and its neighbourhood, and has communicated some interesting information on the vegetation of that country to the *Gardener's Magazine* (see vol. xi. p. 388.), gives the following measurement of ten trees of the *Eucalýptus robústa*, or the stringy-bark tree. They all stood in the neighbourhood of the Emu river, and the circumference of all the trunks were taken at 4 ft. from the ground.

No. 1., 45 ft. in circumference; supposed height 180 ft. The top broken, as is the case with most large-trunked trees; the trunk a little injured by decay, but not hollow. The tree had an exerescence at the base 12 ft. across and 6 ft. high, protruding about 3 ft. No. 2., 37½ ft. in circumference. No. 3., 38 ft. in circumference; distant from No. 2. 80 yards. No. 4. 38 ft. in circumference; distant from No. 3. 56 yards. Nos. 3 and 4. were round trees, upwards of 200 ft. high. No. 5., 28 ft. in circumference. No. 6., 30 ft. in circumference. No. 7., 32 ft. in circumference. No. 8., 55 ft. in circumference; very little injured by decay; and upwards of 200 ft. high. No. 9., 40½ ft. in circumference; sound and tall. No. 10., 48 ft. in circumference; tubercled; tall; some cavities at the base; much of the top gone. A prostrate tree near to No. 1. was 35 ft. in circumference at the base, 22 ft. at 66 ft. 19 ft. at 110 ft. up; there were two large branches at 120 ft.; the general head branched off at 150 ft. the elevation of the tree, traceable by the branches on the ground, 213 ft.

In the *First Additional Supplement to the Encyclopædia of Agriculture* will be found portraits, drawn from nature, of several of the trees mentioned as having been measured by Mr. Backhouse, drawn by Mr. John Thompson, a friend of ours, and an excellent artist, settled at Sydney. The iron-bark tree (*Eucalýptus resinífera*) measured by Mr. Thompson is 200 ft. high, with a clean straight trunk of 130 ft. The most remarkable of these trees in appearance is the grass tree (*Xanthorrhœa arboréscens*).

Mr. Thomas Backhouse has sown the seeds of several species of the trees and shrubs of Mount Wellington and other elevated and exposed situations in Van Diemen's Land, in his nursery at York, and he expresses a hope in a few years to prove their hardiness; and, as they are all evergreens, they will be valuable auxiliaries to our park scenery.

CHAP. V.

OF THE LITERATURE OF THE TREES AND SHRUBS OF TEMPERATE CLIMATES.

A HISTORY of trees and shrubs would be incomplete, without some notice of the literature to which the subject has given rise. In the earlier works on plants, trees and shrubs, as being the more conspicuous division of the vegetable kingdom, occupy a considerable space; and, in modern times, whole works have been exclusively devoted to them. It is only our intention to notice, in a very slight manner, the names of the more remarkable of the works which have been exclusively devoted to the history and description of trees and shrubs, referring, for a chronological enumeration of all the authors who have written on the subject in modern languages, to the second edition of our *Encyclopædia of Gardening*, and to a posthumous work of the late Mr. Forsyth (see *Gard. Mag.*, vol. xi. p. 596.), entitled *Bibliotheca Geoponica*, which will shortly be published.

We have already noticed Aristotle and Theophrastus, as the principal Greek authors who wrote on trees, and Pliny is almost the only Roman one. The information contained in the works of these authors, with some additions from the writings of Cato, Columella, Vitruvius, and others, was used in a new form, on the dawn of literature in the end of the 15th and the beginning of the 16th centuries, in the works on husbandry generally, by Crescentius in Italy (1471), by Fitzherbert in England (1523), Etienne in France (1529), Heresbach in Germany (1578), and Herrera in Spain (1595). The first author who wrote exclusively on trees and shrubs appears to have been Belon, a doctor of medicine of the faculty of Paris, who produced a small quarto volume, entitled *De Arboribus Coniferis, Resiniferis, &c.*, printed at Paris in 1523, and illustrated with a number of engravings on wood. Our copy is the original edition, and consists of thirty-two printed pages, and twenty engravings. Different species of *Juniperus* and *Cupressus*, the *Thuja orientalis*, *Cèdrus Libani*, and several pines and firs, including the *Làrix*, are described and figured; and a number of other plants are mentioned incidentally. Meursius published *De Arborum, Fruticum, et Herbarum, &c.*, in one volume 8vo, at Leyden, in 1600; but, in this work, the medical properties of plants appear to be the main object of the writer. The next work exclusively devoted to the subject of trees is the *Dendrographia* of John Johnston, a Pole, whose work was published in one volume folio, at Frankfort, in 1662. In this work trees and shrubs generally are treated on, and fruit trees at considerable length. It is illustrated with numerous figures, and the object of the author seems to have been to direct attention to the trees which bore edible fruits, or were remarkable for their medical properties. In 1668 the *Dendrologia Naturalis* of Aldrovandus, in one volume folio, appeared at Bon. It is a very thick folio volume, illustrated by numerous engravings, and the medical qualities of the plants are chiefly insisted on. Aldrovandus was born at Bologna in 1557, and died in 1625; he was a great traveller, and one of the most laborious naturalists of the sixteen century.

In England, the first work exclusively devoted to trees and shrubs was Evelyn's *Sylva*, which was published in one volume folio, in 1664. Every one knows the influence which this work had in promoting a taste for planting trees throughout England. It went through several editions during the author's lifetime; and, since his death, an enlarged edition in 2 vols. 4to, with several engravings, edited by Dr. Hunter of York, was published in 1776; and again, with some improvements, in 1786. The first work, after Evelyn's, which was exclusively devoted to trees and shrubs was, the *Descriptive Catalogue of the Trees and Shrubs propagated for Sale in the neighbourhood of London*, by a Society of Gardeners, which we have noticed in p. 60. It

forms a thin volume folio, and appeared in 1730. These are the only works of note, which appeared on the subject of trees exclusively, previously to the time of Linnaeus.

With the exception of nurserymen's catalogues, and some works on planting and managing trees and plantations generally, nothing exclusively devoted to the subject of trees appeared in Britain, till Hanbury published his *Essay on Planting* in 1758: a ponderous folio never in much esteem, and of very little interest. Indeed, the only gardening book in England in which trees and shrubs were described, and treated of botanically as well as horticulturally, previously to the commencement of the nineteenth century, was the *Dictionary* of Miller. The Earl of Haddington, in Scotland, published a *Treatise on Forest Trees*, in 12mo, in 1760; but it can only be considered as a work descriptive of trees and shrubs generally. In 1771, Meader, gardener to the Duke of Northumberland at Syon House, published the *Planter's Guide*, which is little more than a list of trees, with an imaginary engraving showing their comparative heights. A similar list is given at the end of the second volume of Morel's *Théorie des Jardins*, the second edition of which appeared in 1802. In 1772, W. Butcher, a nurseryman at Edinburgh, published a *Treatise on Forest Trees*, already mentioned as a valuable work for the time at which it appeared; and, in 1777, Dr. Anderson, under the name of Agricola, published *Various Thoughts on Planting and Training Timber Trees*. *Planting and Rural Ornament* was published by William Marshall in 1796, in 2 vols. 8vo, one of which is devoted to the description of trees and shrubs, chiefly, as the author acknowledges, taken from Hanbury and Miller. In 1779, Walter Nicol published the *Practical Planter*, and subsequently the *Planter's Calendar*, an edition of which, edited, or rather, rewritten by Mr. Sang, and published in 1812, in 1 vol. 8vo, is the last and the best work on trees and shrubs which has appeared in Scotland.

With the first year of the nineteenth century appeared the *Planter and Forest-Pruner* of William Pontey; but this and the other works on planting of that author belong to the general subject of culture, rather than to the description and history of trees and shrubs. In 1803, Lambert's *Monograph of the genus Pinus* appeared in one volume folio, price twenty guineas; a second volume has since been added; and, in conformity with the spirit of the times, an edition has been published in two volumes 8vo, price 12l. 12s. In 1811, Dr. Wade of Dublin produced a descriptive work on the willow, entitled *Salices*, in one volume 8vo; and, in 1823, Mr. Henry Philips produced, in two volumes 8vo, *Sylva Florifera*, in which the more common ornamental trees and shrubs are treated of in a popular and agreeable manner. Passing over the *Woodlands* of Cobbett, which appeared in 1826, in one volume 8vo, we come to the most scientific work exclusively devoted to trees which has hitherto been published in England, the *Dendrologia Britannica* of P. W. Watson, which was completed in two volumes 8vo, in 1825. The first volume contains 80 plates, and the second 90 plates. The letterpress, with the exception of 72 pages of introductory matter, consists solely of technical descriptions of the figures, arranged in a tabular form under a given number of heads; a very effectual mode of preventing any point, necessary to be attended to in the description of a plant, from escaping the notice of the describer. In this respect, the work is superior to some of its contemporaries, in which the descriptions are sometimes rather disorderly if complete; and are often incomplete, apparently from want of being taken in some fixed and comprehensive order. Mr. Watson was a tradesman in Hull, who afterwards retired from business; and he was one of the principal persons who assisted in founding, and afterwards in laying out and managing, the Hull Botanic Garden, as stated in the introduction to his *Dendrologia*, p. xii. He died, we believe, in 1827. The only work hitherto published in England, which contains a description of all the hardy trees and shrubs in the country, in addition to that of all other plants, ligneous and herbaceous, described by European botanists, is Don's edition of *Miller's Dictionary*, in four volumes 4to, price 11l.

In France, the first really important work on trees, in modern times, is the *Traité des Arbres et Arbustes*, by Du Hamel du Monceau, which was published in Paris, in two volumes 4to, in 1755. In this work the nomenclature of Tournefort is followed, but the names of Linnæus are also given; it is illustrated with numerous figures, partly taken, as the author informs us, from the blocks which were used in the *Commentaries* of Mathioli; and partly engraved on purpose for the work. The first volume contains 368 pages and 275 engravings, and the second 387 pages and 199 engravings. The original edition is not very common, and, when met with in London, sells at from thirty to forty shillings. A new edition of this work was commenced in the year 1800, and it was completed in seven volumes folio in 1819. The letter-press of these volumes was prepared by Mirbel, Loiseleur Deslongchamps, and other botanists; and the drawings were by Redouté, Bessa, &c. The published price of a royal folio copy was 12*l.* 10*s.*, and of a common copy nearly 100*l.* The species are arranged according to the Linnæan system; and the number of engravings of trees and shrubs, including some engravings of fruits, amounts to 498. Both engravings and descriptions are of very unequal merit, and many of the former (at least in our copy, which is a large paper one) are altogether unworthy of the consequence attempted to be given to the work by large type, large paper, and other characteristics of the mode, now gone by both in France and England, of publishing for the few. As a proof of the truth of what we assert, large paper copies may now be purchased in London for between 30*l.* and 40*l.*, and small paper copies for twenty guineas.

In 1809, while the new edition of Du Hamel was slowly publishing in parts, the *Histoire des Arbres et Arbrisseaux*, by M. Desfontaines, appeared in two volumes 8vo, and is still a work of repute. In 1824, *Traité des Arbres Forestières, ou Histoire et Description des Arbres Indigènes, naturalisés, dont le tige a de trente à cent vingt pieds d'élevation*, &c., par M. Jaune St. Hilaire et M. Thouin, appeared in one volume 4to, with coloured plates, price 10*l.* The plates are badly executed, and the work, with the exception of the part written by Thouin, is of a very inferior description,

André Michaux, a notice of whose life has been given, p. 140., published *Histoire des Chênes de l'Amérique*, in one volume folio, in 1801; and his son, F. A. Michaux, published *Histoire des Arbres Forestières de l'Amérique Septentrionale*, in three volumes, large 8vo, in 1812. Of this work there is an English translation entitled the *American Sylva*, which was published in Paris, in 1817, at nine guineas plain, or twelve guineas coloured. F. A. Michaux's work contains 156 plates, including figures of all the oaks described in the *Histoire des Chênes*, and is an excellent work, which still maintains its price both in Paris and London. We ought not to pass unnoticed *Le Botaniste Cultivateur* of Du Mont de Courset, in seven volumes 8vo, which was completed in 1814, and which, though it contains herbaceous and house plants, as well as ligneous hardy plants, is yet more complete in its descriptions of the latter than any other work, except Du Hamel's. There is no French work which brings down the description and history of trees and shrubs to the present time; but, if we were asked what works we would recommend, as making the nearest approach to this, we should say, *Le Botaniste Cultivateur*; *Les Annales de Fromont*; *Le Bon Jardinier*, the edition of which work for the current year contains notices of all the plants newly introduced; and, above all, the excellent *Prodromus* of De Candolle, now in course of publication, and of which four volumes 8vo, price 5*l.*, have already appeared.

In Holland, the only work exclusively devoted to trees and shrubs which, we have heard of, is by Krause, and the title is, *Afbeeldingen der Fraaisste, Meest-wilheemsche Boomen en Heesters*, &c. It appeared at Amsterdam in 1802, in one thick royal 4to volume, the price of which in London is 10*l.* The plates in our copy are executed in a very superior manner, and they are coloured with much more care than those of either Willdenow, Schmidt, or Du Hamel. Some of the German works describing the different kinds of wood were published at Amsterdam, as well as at Leipzig; particularly that

published by Sepps, which appeared in one volume 4to, in 1773, at both places; and at Amsterdam, with a translation of the German descriptions into Dutch. It was published at about 15*l.*, and sells in London for from 8*l.* to 10*l.*

In *Germany*, the first work exclusively devoted to trees and shrubs, which was published in modern times, was the *Harbkesche Baumzucht theils nord-amerikanischen und fremder*, &c., of J. P. Du Roi, which appeared in two volumes 8vo. in 1771-2; to this succeeded the *Osterreichs allgemeine Baumzucht* of Schmidt, the first two volumes of which appeared in 1792, and the third in 1800 price 10*l.* This is an excellent work as far as it goes. The next German work which appeared was the *Berlinische Baumzucht* of C. L. Willdenow, in one volume 8vo, the second edition of which appeared in 1811. In this volume are described all the trees and shrubs which grew at the time in the Berlin Botanic Garden. It is observed by Watson, in his introduction to his *Dendrologia*, p. vii., that, in the *Berlinische Baumzucht*, "the parts of the plant in each description are placed in the same order, and not in the often vague, slovenly, and reiterated way of many botanic writers." In 1810, appeared the first number of *Abbildung der deutschen Holzarten*, &c., by F. Guimpel, C. L. Willdenow, and F. G. Hayne. It was completed with 36 numbers in 1820. It contains 216 coloured plates, and the price in London is 16*l.* In 1819, another work on the trees of Germany appeared, which included all those foreign species which stand the open air in that country. It is entitled *Abbildung der fremden in Deutschland ausdauernden Holzarten*, &c.; by F. Guimpel, F. Otto, and F. G. Hayne. It forms one volume 4to., contains 100 plates, and costs in London 6*l.* The Germans have a species of publication, or rather portable museum, which they call *Holzbibliothek* (Wood Library). A hundred or more sorts of wood, with dried specimens of the leaves, flowers, seed, and winter's wood of each species, are put loose into little cases about the size of 8vo volumes, and these are finished exteriorly like books, and the back lettered with the name of the wood in different languages. There are two libraries of this description, which are more particularly in repute; one published at Munich, and the other at Nuremberg. The former contains 100 sorts of wood, and costs about 100*l.*, and the latter 80 sorts, and costs 80*l.* There are also engravings and descriptions of these woods, at various prices from 10*l.* to 30*l.*; one of the best of these is Hildt's, published at Weimar in 1798, price 15*l.*

We are not aware of any other work of magnitude exclusively devoted to the description and history of trees and shrubs, having been published in any of the other countries of Europe; but in *America*, besides the English translation of Michaux, already mentioned, which appeared there as well as in Paris, in 1817, there is the *Sylva Americana* of D. J. Brown, which was published at Boston in 1832. The cost in London of this work is 1*l.* 1*s.*

It will be recollected, that in this notice we have only given the titles of the principal works devoted *exclusively* to the description and history of trees and shrubs, those which treat on the formation and management of plantations, on forests and woodlands, and on planting trees generally, whether for useful or ornamental purposes, are exceedingly numerous, and will be passed under review in the Introduction to the Fourth Part of this work.

CHAP. VI.

CONCLUSION.

Two considerations may be drawn from the preceding history: the first, respecting the introduction of foreign trees and shrubs; and the second, respecting arboricultural literature.

On comparing the lists which we have given of ligneous plants, found in

the different countries situated in temperate climates, which are not indigenous to Britain, with the catalogues of plants considered as already introduced into this country, it will be found that there are some names of species and varieties, in the lists of almost every country, that are not in British catalogues, and, consequently, not yet introduced. There are a number of names, for example, that are not in our *Hortus Britannicus*. Here, then, is an important use in giving these lists, because they point out to commercial gardeners, and to amateurs and travellers who are botanists, what trees and shrubs it is desirable to enquire after in other countries; and what they should endeavour, if possible, to introduce into their own. It may also be observed, that the same names that are in the lists in preceding chapters frequently occur in British catalogues; yet it is by no means certain that in every case they are applied to the same plants. Here, then, is another point calling for the exertions of the patriotic botanist or traveller; and it may be observed respecting this point, that it will always be the safest side to err on, to reintroduce plants which are already in the country, rather than to remain in any doubt respecting our possessing them. All trees and shrubs liable to great variations, and of great utility or ornament, such as the genera *Quercus*, *Pinus*, *Pyrus*, *Crataegus*, &c., ought to be introduced in all their varieties. We are persuaded that there are, in France and Spain, many interesting varieties of *Pinus*, and of *Quercus Cerris* and *Q. Flex*, which have not yet found their way into British gardens. The *Quercus australis*, recently introduced in consequence of the information given by Captain Cook (see p. 171.), may be cited as a proof of this; and there are, doubtless, hundreds of species and varieties of trees and shrubs in North America, that have never yet been seen in Europe. There is, therefore, ample room for exertion, to those who wish to increase the botanical riches of their country; and more especially in the case of that kind of botanical riches which adds so conspicuously and permanently to its ornament, and to its useful resources. It would, however, be a very contracted view of this subject, to limit our views to the aggrandisement of the collections of trees and shrubs in Britain. The time for believing that the exclusive possession of any benefit contributes to the prosperity or happiness of nations is gone by; and the principles of free and universal exchange and intercourse are found to constitute the surest foundation for the happiness of nations. This is so obviously true in matters of botany and gardening, that it cannot for a moment be doubted.

If it is desirable for us that we should assemble in our country the trees and shrubs of every other similar climate, it must be equally desirable that the inhabitants of every other similar climate should possess all those species for which their climate is adapted; in short, it is desirable for the advancement of civilisation and human refinement, that all the trees and shrubs of temperate climates should be distributed throughout all those climates. This will, no doubt, be the case at some future period, when the civilisation of the whole world is comparatively equalised; and, in the mean time, it may be useful to offer such hints as will contribute towards so desirable a result.

One of the first steps towards the equalisation of the plants of different regions, of similar capacities for growing such plants, is, to determine, with something like precision, what the plants of each region are. This can only be done by assembling living specimens of all of them, or of such a class of them as it may be desirable to equalise, in one garden, and cultivating them there for some time, so as to determine the species and varieties. In old countries of limited extent, such as Britain, which have been examined by botanists for two or three centuries, the establishment of botanic gardens for determining the number of indigenous species of plants may be considered unnecessary; but this is far from being the case with such countries as North America, Upper India, China, Japan, Van Diemen's Land, and a number of others. It would, doubtless, contribute to the spread, all over the world, of the trees and shrubs of North America, if one part of them could be seen in a grand national garden at New York, and another in a garden or arboretum

at Charlestown; or if the whole could be assembled in one grand park and pleasure-ground at Washington. We do not, however, expect this to be done; for, in the progress of civilisation and refinement in every country, there are many much more important points to be attended to than the culture of trees and shrubs.

With a view both to introduction into any particular country, and equalisation in all similar countries, the first thing that ought to be done, after the indigenous plants of the country are clearly determined, is to assemble, in one garden in the given country, all those of all other similar countries, with a view to determining what they are. Thus, in the case of trees and shrubs, all those mentioned in the preceding lists as being foreign, with reference to the trees and shrubs of Britain, ought to be assembled and proved in one grand British national garden; and the same thing ought to be done in a similar garden in every other country. The idea may be thought chimerical; but it is hard to say what will not be done by nations, when they come to cultivate with ardour the arts of peace and of refinement. The cooperation of individuals attached to the same pursuit may, in the mean time, contribute much to the advancement of that pursuit; and hence, though in the neighbourhood of London it might be difficult, or impossible, to induce the metropolis or the government to form an arboretum of 150 acres in extent, so as to exhibit in it all the species of trees and shrubs of temperate climates, that will grow with us in the open air; yet, if the proprietors of a certain number of estates within ten miles of London were to agree each to form a collection of the species or varieties of a particular genus, and to allow these to be examined by botanists, the same result would be obtained, at least as to accuracy of nomenclature, as if the whole of the genera and species were assembled in one national garden. We have explained this cooperative system, at length, in the *Gardener's Magazine*, vol. xi. p. 600., and it is needless to say that it is as applicable to every other country as to England, and to every other large town as to London. The practice might, therefore, be adopted in the neighbourhood of Edinburgh and Dublin; and in that of Exeter, for the south of England; Bristol and Liverpool, for the west; York, for a central situation; Norwich, Lincoln, and Hull, for the east; and Newcastle and Carlisle, for the north.

The number of works which have been written, exclusively devoted to the description and uses of trees and shrubs, shows, in a general point of view, the estimated importance of the subject by authors; and, when we consider the rapidity of the succession of these works within a comparatively limited period, it shows the accumulation of knowledge which is continually being acquired respecting ligneous plants. In this, as in all other branches of natural history, the attention of naturalists was first directed to the objects which more immediately surrounded them in their own country; afterwards it extended to adjoining countries; and lastly, which seems to embrace the most comprehensive view of the subject, to all the other countries in the world which are similarly circumstanced in regard to climate and all that affects the growth of trees and shrubs. Hence, the first books on dendrology were merely local catalogues, enumerating the medical, or such other properties, as, in the age in which these books were produced, excited most attention; while the last are scientific descriptions, with the geography, history, and uses in civilised society, of all the species and varieties enumerated.

The conclusion which we draw with reference to the literature of trees and shrubs is, that, though there are a great many excellent works on the subject already before the public in the English, French, and German languages, yet none of these works embrace the whole subject, and bring it down to the present time; and that, consequently, we are justified in our endeavour to comprise every thing which it is desirable to know, respecting the trees and shrubs which will stand the open air in Britain, in our present Encyclopædia.

PART II.

OF THE SCIENCE OF THE STUDY OF TREES.

THE subject of trees, like every other subject, to be studied in the best manner, must be studied according to some methodical arrangement; and it is the purpose of this part of our work to lay down an outline of that method which we consider the best. By including in such an outline every particular which ought to be taken into consideration in the study of trees, we shall be the less likely to omit any of these particulars in giving the description and history of individual species; and the reader, also, will profit by having his attention previously directed to what this history and description ought to contain.

Trees may be studied both as individual objects, and as connected with other objects. As individual objects, they may be considered pictorially, or as component parts of the general scenery of a country; and botanically, as organised beings. As objects connected with others, trees may be studied with reference to nature, animate and inanimate, which may be considered as their natural history; and with reference to man in a state of civilisation, which may be called their economical history. We shall devote a short chapter to each of these subjects, and in a concluding one give a summary of the whole.

CHAP. I.

OF THE STUDY OF TREES PICTORIALY, OR AS COMPONENT PARTS OF GENERAL SCENERY.

THE study of trees, as component parts of landscape, may be carried on with very little knowledge of either their natural or their economical history; and, indeed, with less knowledge of any other objects or sciences, than most studies. Experience proves that a man may excel as a graphic artist with very little knowledge beyond his art; but, at the same time, it is also found, from experience, that all the greatest artists have been, more or less, philosophers; and hence, though a knowledge of the natural and economical history of trees may not be essential for the artist who studies them pictorially, it will yet be found to render him material assistance.

The subject of this chapter naturally divides itself into the study of the forms of trees and shrubs; the study of their character and expression; and the art of delineating them pictorially.

SECT. I. *Of the Study of the Forms of Trees and Shrubs.*

THE first quality in a tree which will strike a general observer, coming to the study with only a few notions relative to form, will be its bulk, or the space that it occupies in the landscape which meets his eye. This bulk, or magnitude, resolves itself into height and width; and the consideration which immediately follows is, the outline that the tree makes against the sky, or against any other object which appears behind it. The next points that will probably attract notice are, the colour of the tree, and the degree of brilliancy of the lights which appear on its masses. Subsequently, the attention may be drawn to the trunk of the tree: for example, to observe whether it appears to be adequate to the support of the head; whether the head appears equally balanced on it; and whether it stands perpendiculary, or obliquely, to the sur-


face on which the tree grows. The next point is, to observe whether the head is open and airy, or compact; and the last, whether the general form of the tree is regular or irregular. All these particulars are equally applicable to shrubs as to trees; with the exception of those which apply to the trunk, which is almost always wanting in what are pictorially, as well as botanically, considered as shrubs or bushes.

The different points, then, to which attention ought to be directed in the study of trees and shrubs as pictorial forms, are the following:—the height and breadth, or general magnitude, of the tree; the form and outline; the colour, light, and shade; the position of the trunk and branches; the mode of growth; the mode of tufting; the leaves, and the spray and buds.

The height and breadth of trees and shrubs vary according to their kinds, and to the soil, situation, and climate in which they grow. The trees of greatest height, in the temperate regions of the globe, are those of the pine and fir tribe; and they are those, also, which have least breadth in proportion to their height; because their branches, technically called frondose, have not the same tendency as those of other trees to increase in thickness with the age of the tree. The most bulky trees, or those which have the greatest width of head in proportion to their height, are some species of poplars, elms, oaks, &c., and the cedar of Lebanon, which, though it is a tree with frondose branches, yet, when it stands singly, has a head generally broader than its height. The highest shrubs, which grow independently of other objects of support, are, among evergreens, the common laurel, and, among deciduous kinds, the common lilac, the dogwoods, and the Guelder rose. Among the smallest trees, considered as pictorial objects, are the thorns and the spindle tree; and among the smallest shrubs are the heath, the furze, and some of the rhododendrons and azaleas.

The form and outline of trees and shrubs vary chiefly according to their kinds and their age, but partly, also, according to the physical circumstances in which they are placed; such as soil, situation, climate, and, perhaps, above all, proximity to other trees and shrubs. The natural form and outline of a tree can only be ascertained when the tree stands alone. The form which it assumes, when closely surrounded by other objects, will generally be found very different from its natural form; and, therefore, cannot be considered as characteristic of the tree or shrub. Even the climate, or the prevailing soil, or wind, will materially alter the form. The cedar, when planted in masses like the larch or the Scotch pine, produces, like them, a straight trunk, with the branches dying off from the bottom upwards; planted singly, its branches often become like so many trunks: in a sheltered situation its top will be pointed; and, in one exposed, it will become blunt or flattened, like the well-known cedars of the Chelsea Botanic Garden. Nearly the same observations may be made respecting all other trees: even the sturdy oak, in poor soils and cold elevated situations, becomes a bush; in rich soils and low situations it is a lofty tree, with a straight trunk; and in situations exposed to the sea breeze, it has the entire head of the tree leaning to one side, and presents an appearance altogether peculiar. *Fig. 1.* shows the effect of the sea breeze on two oaks growing on the boundary of Camford Heath, near Wimborne, in Dorsetshire; and numerous similar examples might be given from different parts of the island.

However various the outline of trees and shrubs may appear in detail, the general contour may always be reduced to some familiar or recognised form, easily retained in the memory: for example, to that of an oval, or that of an egg placed on its smallest end, which are the most common forms of trees, except in the case of those which have frondose branches; and that of an egg placed on its broadest end, or of a flattened cone, which are the most common forms of shrubs. The Lombardy poplar, the cypress, and some similar trees, may be said to have their heads in the form of an ellipsis: and others such as the common apple, thorn, &c., in that of a globe, or in that of a semi-globe. A few trees, more particularly in their young state, take the form of an inverted cone, such as the planera, as will appear by our figure of that tree at ten years' growth. Within these general forms, the greatest variety of outline



may be produced, by the receding or advancing of the terminations of the branches ; by the forms of these terminations ; by the manner in which they are clustered together ; and by their being open or compact, regular or irregular, densely clothed with foliage, or only loosely covered.

Light and shade. The form of a tree or shrub is generally indicated by its outline against the sky ; but that outline does not convey such a correct idea of the form as where the space enclosed by it exhibits both light and shade, which alone can give it the appearance of substance.

The capacities of different trees for receiving light vary according to the density of the masses of foliage. A compact round-headed tree, like the *Sórbus Aria*, will receive and reflect the light in one large mass ; a tree or shrub, the general form of which is composed of numerous smaller masses, more or less separated from each other, will reflect the light in smaller masses ; and a thin tree, which, in many parts, may be seen through, will not reflect the light from any part in a conspicuous manner. In proportion as the light is reflected from one side of a tree, the other side will appear dark ; and a tree or shrub thinly clothed with foliage will appear to be equally in light and shade throughout. Most light and graceful trees are of this description ; but the want of the beauty produced by light and shade is compensated by the wavy direction of the stem, which gives the idea of gentle motion, and by the variety and intricacy of the tuftings, outline, and disposition of the smaller branches, spray, and leaves.

The colours of trees and shrubs vary not only according to the kinds, but according to the season of the year, the climate, soil, situation, age, and other circumstances. In general, the colours of spring, when the tree is leafing, are more fixed and determinate than those of autumn, when it is losing its leaves ; because, in the former case, it depends chiefly on the nature of the tree, while in the latter it is materially affected by the weather, and also by the soil. The birch, the leaves of which, in most soils, in a wet autumn, will die off of a yellow colour, in a dry autumn will assume a deep red ; and, on a peat bog, birch leaves generally die off of a pale yellow, while on a granitic soil they generally assume a bright red or purple. On the side of the tree next the sun, the leaves of all trees are of a brighter colour than on the shady side. Those trees and shrubs which vary least in colour, whether in summer or autumn, are the evergreens ; and those which vary most are the oaks, the acers, and the thorns.

The trunks of trees vary as much as their heads ; not only naturally, according to the kind of tree, but accidentally, according to the circumstances in which it has been placed by nature, or by the hand of man. In general, a detached tree exhibits a trunk clothed with branches from the ground upwards ; but trees grown in masses generally exhibit naked trunks to a considerable height. In some species, as in the pine and fir tribe, the length of naked trunk is great, in proportion to the size of the head ; in others, as in the oak and

other ramose-headed trees, the length of naked trunk bears a much less proportion to the head. Shrubs are distinguished from trees, not only pictorially, but also botanically, by having no distinct trunk, or naked stem, but, whether large or small, forming one entire bush.

The trunks of trees may farther be considered with reference to the manner in which they rise out of the soil, and to the manner in which the head of the tree is supported by them. Trees which grow in thick woods, where the annual crop of leaves falls down, and rots into mould, have their trunks apparently without bases; but trees which stand in open situations, where the falling leaves are blown away, always rise out of the ground with a conspicuous base, formed by the junction of the trunk with the roots. No circumstance adds more to the effect of a tree, as a stately and durable object in a landscape, than the appearance of its trunk, rising from a secure and widely spreading base; and this appearance is one which may be imitated by art, both in drawing landscapes, and in landscape-gardening, and whether the trees to be introduced are young or old.

The perpendicularity or inclination of the trunks of trees are circumstances well deserving the study of the artist. Wherever trees have grown up fortuitously in groups or masses, the trunks of many of them will be found more or less inclined to the horizon; and their heads will often appear ill balanced when taken singly, though in combinations such trees make the most varied groups. Trees which have been planted singly, and exposed alike on every side, grow up with erect trunks, and form the most stately and well-balanced heads. Such trees always have the appearance of having been planted by art; while the others seem to convey the idea of their having been subjected to the operation of natural causes. Trees of the first kind may be called gardenesque trees, and of the latter, picturesque trees. Trees which grow out of rocks, or on precipices, or on the banks of water, almost always have their trunks inclined to the horizon; as trees on level surfaces, other circumstances being the same, have their trunks at right angles to the horizon.

Though shrubs are without trunks, yet the same general remarks will apply to them. A detached shrub, on a level surface, is clothed to the ground on every side; a shrub which has grown up among other shrubs or trees, or which stands on a slope or hangs over water, will be inclined to one side. Shrubs, however, admit of much less variety in point of attitude than trees.

The branches of trees differ in many particulars. The frondose branches of the pine and the fir tribe never, except under accidental circumstances, attain any great size; on the other hand, the ramose branches of the oak, the chestnut, &c., frequently attain a size which rivals that of the trunk. The branches of some kinds of trees proceed regularly from the stem, in some definite and obvious order of succession; others proceed from the stem apparently in an irregular manner. Some extend a great length horizontally, or at right angles to the stem; and others a great length acutely, or obliquely to it. The heads of some trees, as the cedar of Lebanon, in a detached situation, seem to consist entirely of spreading branches; of others, such as the pinaster, of a few branches at the upper extremity of the trunk. The ramifications of the branches are as various as their mode of insertion in the trunk: some are quite regular, as in the fir tribe; and others are exceedingly various and intricate, as in the oak. The branches of some trees have few branchlets, as in most of the poplars; others have many branchlets, as in the oak, the beech, the birch, &c. The same observations will apply to shrubs, but not in the same degree. In the case of shrubs, numerous stems arise from the root; and these stems, in many cases, produce only branchlets and spray, and, in others, only leaves. In some shrubs, as in the common laurel, the branches trail on the ground; in others, as in the berberry and the lilac, they are numerous and erect. Certain shrubs have stems which twine round, or are otherwise supported by, trees or other objects; such as the ivy, the clematis, &c.; and others rise up, and fall down, as in the case of the common bramble. Some shrubs are, in their details, like a collection of miniature trees; as, for example, the butcher's broom. In

a botanical point of view, the variety which trees and shrubs present, with respect to their trunks and branches, is almost endless: but we have, at present, only to consider them with a view to pictorial effect.

The mode of growth in trees and shrubs varies according to the kinds; and is modified only in a slight degree by situation, soil, and climate. The main stems of almost all trees have a tendency to grow upright; but the branches proceed from these stems in different directions in different trees. The growths, that is, the branchlets and spray produced by the branches, extend themselves horizontally in some cases, as in the oak; upwards in others, as in the beech; and in some downwards, as in the weeping willow and the weeping birch. In some they are rigidly upright and compact, as in the Lombardy poplar; and in others they are upright and sparse (that is, thin), as in the gymnocladus. In young trees and shrubs the mode of growth is more decidedly exhibited than in old trees, because the growth is more rapid; and in detached trees it is more conspicuous than in such as are crowded, because the nature of the tree, in such situations, is more perfectly developed.

In shrubs, the mode of growth is often the principal circumstance by which, in a pictorial point of view, they can be characterised; because they have not sufficient magnitude to admit of great variety of form, or of outline: they are without trunks to admit of variety of attitude; and they are without large branches to admit of marked character. The artist, therefore, in order to render them interesting, must have recourse to their mode of growth, and to their foliage; and for these reasons, also, they ought almost always to be placed in the foreground, both of pictures and of garden landscapes.

The mode of tufting, that is, the mode in which the leaves combine into those masses which appear on the surface of the tree, varies with the kind of tree; and depends principally on the mode of growth, though, to a certain extent, it is influenced by the form of the leaves. The general form of tufts is roundish, or elliptical; the long axis of the ellipsis, or of the roundish figure, being universally in a horizontal direction. One tuft is separated from another by intervals of shade in the body of the tree, and by the background where the tufts compose the marginal outline. All tufts have their margins more or less indented, which depends on the forms and clusters of the leaves.

The leaves of trees and shrubs are almost as various as the species; but, pictorially, their forms are chiefly noticeable only so far as they combine into masses. A single leaf, in a tree viewed pictorially, is at such a distance from the eye as seldom to be seen in its entire form; but something of the general form of the leaves is recognisable in the margins of the tufts of foliage, and more especially in those which are nearest to the eye. The margins of these tufts may be said to be almost always more or less indented or serrated; the serratures differing in magnitude, in depth, in direction, and in being more or less blunt at their extremities.

The spray of trees, that is, the last made shoots on the extremities of the branchlets, is as various as the species of trees. In some it is horizontal, as in the silver fir; in others it is pendent, as in the larch, the weeping willow, and in some species of birch; in others it is rigid, as in the gymnocladus; and in others crooked, as in some of the oaks and thorns; in some it is thick, and unmarked by buds, as in the fig and the walnut; in others it is clothed with leaves throughout the year, as in all the evergreens.

The buds of trees and shrubs may be considered as scarcely perceptible in pictorial point of view; nevertheless, the spray of some trees are without visible buds; and in others, for example, the horsechestnut, the *Bétula nigra*, and the *Quercus sessiliflora*, the buds are very conspicuous. The blossom buds, both of trees and shrubs, are always much larger than the common buds; for which reason, in looking at a tree pictorially, during the spring season, it will be some assistance to the artist to know whether the tree is one which produces large or small flowers; and whether it is of a sufficient age to produce flowers at all. The knowledge of this, and of

other particulars which may be considered botanical, will be of great assistance to an artist, in enabling him to correct his pictorial observations.

These are the chief circumstances, with regard to trees and shrubs, towards which attention ought to be directed, with a view to their pictorial effect, independently of the associations connected with them; and hence, in giving the history of individual species, it would be necessary to test each by all these different properties, with a view to determining its appropriate uses in landscape-gardening, and in ornamental planting.

SECT. II. *Of the Expression and Character of Trees and Shrubs considered pictorially.*

EVERY object in nature that forms a whole has some *expression*. If the nature of the object is unknown to the beholder, the expression which he assigns to it is analogous to that of some object with which he is already familiar; and he uses the same terms to describe its appearance as he would apply to such objects. For example, a tall, erect, regularly clothed tree will be described by the epithets stately, noble, or handsome; another kind of tree, with light airy foliage and a wavy stem, will be called graceful; and so on.

Character is some circumstance added to expression, which renders it more remarkable; and the circumstance which has this effect will generally be found to be the accidental exaggeration of some quality belonging to the natural expression of the object. For example, in the case of handsome regularly clothed trees, supposing a number of them standing together, character would be added to one of them by the extraordinary prolongation and magnitude of one or more of its branches; or by some of its branches having been taken away, so as to expose a portion of the trunk conspicuously, while the remainder continued clothed. Character would also be added to one tree, among a number of the same kind all previously alike, if a portion of this tree were seathed by lightning; or if some circumstance were to occur which threw the trunk over to one side. In either of the latter cases, what is called character would be conveyed by the object displaying, conspicuously, something which did not naturally belong to its species; while, in the former case, character was given by the exaggeration of some quality which was natural to the species.

The expression of trees may be said to be of two kinds: that which proceeds from their organic influence on the eye as forms, without reference to their nature, and altogether apart from moral associations; and that in which moral associations are the principal cause of the expression.

Supposing a person to see a tree or shrub for the first time, and to be totally ignorant of its nature; he could only look upon it in the light of a form; and, in this case, its expression, to him, would depend upon its resemblance to forms which he previously knew, whether geometrical figures, or the figures of other objects. Thus it is, that the first effort which the mind makes to discover beauty in natural forms is, to recognise in them some of the forms of art; and hence, in the infant state of this taste in individuals, the first trees that would be admired would be those the heads of which bore the nearest resemblance to a globe, a cone, or some readily recognisable figure. The next step would be the recognition of some artificial figure, in the trees or shrubs of more regular outlines. To this would succeed the recognition of several figures contained within one general figure; and, lastly, the recognition, among these several figures, of regularity in their arrangement, or of symmetry in their disposition; of variety; of intricacy; and, lastly, of harmony and of character. In this way it will be seen, that a tree or a shrub is capable of exciting many ideas of beauty, considered simply as a form, and altogether apart from considerations of usefulness, of botanical interest, or of moral associations.

A tree, to be regular, or, in other words, to have the expression of *regu-*

larity, must exhibit the same number of tufts of the same size, and at the same distance from each other, on both sides of the trunk, or centre line. Such a tree, as a whole, we may suppose to be a regular globular figure; and it is a property of regularity, that one portion of any figure which is regular, being separated from the rest, will still be regular. If the globular head of the regular tree, therefore, were cut exactly in two, either vertically or horizontally, the separated portions would each be regular figures, that is, semi-globes.

In a *symmetrical tree*, on the other hand, there may be the greatest irregularity in the form and number of the parts, provided, only, that the same quantity of these be distributed on each side of the central vertical line, or trunk, of the tree. In the lower part of the tree, the branches or tuftings may protrude chiefly on the right hand and be large; and, on the upper part of the tree, they may protrude chiefly on the left hand, and be small; or they may be of mixed sizes in both places. Now, it is the characteristic property of a symmetrical figure, that, when it is cut in two, the parts separated, taken singly, are neither regular nor symmetrical; and, consequently, that they will not produce a whole in any way, unless they are reunited as they were before. By imagining such a tree as we have described, separated into two portions by a central vertical line, it will readily occur to the mind, that neither of these portions was either regular, or could in any way form a whole. Whoever wishes to enter on the study of the expression of regularity, symmetry, intricacy, and harmony, in detail, may refer to articles on the subject in the first volume of the *Architectural Magazine*.

The lowest degree of *organic beauty*, in a tree, we may suppose to be the form of a small-sized tree with a lumpish head, like the *Pyrus Aria*; constituting one uniform mass, light on one side, and shaded on the other. A higher degree of beauty will be, where the general form of the mass is that of a cone, or where it is egg-shaped; because these forms contain an additional element of beauty to that constituting a globe, viz. that of elongation. A still higher degree of beauty will be produced, where all the branches of the tree, in the case either of a globe or cone, are distinctly marked and regularly placed on each side of the trunk, as in the case of a fir; and one still higher, where the branches and tufts project irregularly, but are still so balanced, on each side of the centre, as to form, on the whole, a symmetrical figure. Such a figure, where the tufts are numerous, varies in point of size and disposition; and, where the symmetry is perfect, will exhibit the greatest organic beauty of which trees are susceptible.

The association of ideas connected with trees has given rise to what is called their *moral and historical expression*. A tree which is young and growing freely, is said to be in good health, and thriving; and one that is not growing freely, is said to be sickly. A tree with a thick trunk and spreading branches is said to be strong and vigorous; one with a tall and slender trunk, to be light and elegant; one with a bending, or serpentine, wavy-like stem, as we have before observed, to be graceful; a tree with upright growth, to be rigid; and one in which the branches and spray droop, to be mournful, or weeping. In like manner, there are particular associations connected with trees which bear fruit; with those which grow in particular soils or situations, as mountains, marshes, &c.; and with those which are applied to particular purposes, as the oak for ship-building; the pine and fir for house-carpentry; the willow for basket-making; the thorn for hedges; and so on. The historical and geographical associations connected with trees are numerous, and of great interest: for example, the platanus reminds us of the respect paid to this tree in Persia; the sweet bay, of its shoots being used by the Romans to crown their warriors; the vine and the olive, of their unknown antiquity, and the highly prized liquors made from their fruits: and the cedar of Lebanon, of the esteem in which its wood was held in the days of Solomon. A knowledge of all the different associations which belong to each particular kind of

tree, as it must add greatly to the enjoyment derivable from them, ought always to form a part of their individual history.

One of the most common and generally understood expressions of trees is, that which is called their *picturesque beauty*. Much has been written, by Gilpin, Uvedale Price, and others, in order to define the meaning of the word picturesque; and the expression of this beauty has been divided into two kinds: that of the beautiful, or smooth picturesque, suitable for cultivated scenery, and also for painting; and that of the rough picturesque, suitable for wild and forest scenery, and eminently so for graphic representation, whether by the pencil, or the palette. Among trees, whether in a wild or cultivated state, are found both these expressions; and, as an example of the first, we may give young specimens of the willow and the lime, and young trees generally; and of the second, the oak, the cedar, the thorn, and old trees generally. For a tree to be picturesque, it is not necessary that it should stand singly, or be at all symmetrical in point of form, or the general balance of its head; on the contrary, a mutilated tree, or one the trunk of which, from some accidental cause, has grown to one side, may be as picturesque as a tree which grows erect, and is comparatively perfect in all its parts; provided only that it is not a peculiar tree, such as that shown in *fig. 1*, p. 195. The only thing which is essential to picturesque beauty in a tree, is, that it should be capable of readily grouping with another tree, or with any building, object, or animal, so that the combination may form a satisfactory whole. It is evident that this remark applies alike to trees of the rough picturesque, and of the smooth picturesque: since young trees, which belong to the latter kind of beauty, will group together, or with other objects, just as readily as old trees which belong to the former kind of this expression.

The expression of *gardenesque beauty*, in individual trees, differs from the picturesque, in being (whether in the rough or the polished variety of the expression), at all times, regular, or symmetrical. The gardenesque is found exclusively in single trees, which have been planted in favourable situations; not pressed on, during their growth, by any other objects; and allowed to throw out their branches equally on every side, uninjured by cattle or other animals; and, if touched by the hand of the gardener, only to be improved in their regularity and symmetry. A truly gardenesque tree, when fully grown, has always some of its branches depending on the ground, in order to mark it as a tree of the garden or lawn; and not one of the park, where its lower branches would be separated from the ground by that horizontal line formed by the browsing of cattle; or one of the forest, where, from being pressed on by other trees, or, when young, growing up under the shade of an old tree, its form would necessarily belong to some division of the picturesque, or be peculiar; and peculiarity, in trees, as in other objects, as Sir Joshua Reynolds has shown, is deformity rather than beauty.

Architectural and sculptured trees are now no longer in repute: but we see no reason why trees should not be cut into the forms of colonnades, arcades, triumphal arches, and the figures of men and animals, as shrubs are cut into the form of those green walls which, under the name of hedges, separate our fields; and exotic plants are dwarfed, by being grown in pots or boxes; and fruit trees flattened by being spread out against walls. We do not say that arcades of trees, tonsile bowers, and sculptured evergreens, such as were formerly common in French gardens, rank high in the scale of verdant beauty; we merely assert that they are productive of distinct kinds of beauty; and that it is by no means desirable to be exclusive, and limit our notions of what is beauty to that which is highest in the scale, or to those kinds only that happen to be fashionable in our time.

Viewing trees with reference to their beauty as organic forms, and to the interest which they are capable of exciting by calling forth associated ideas, the tree which is considered the most beautiful by man, in any country, will vary according to the knowledge of every individual, and the country in which he

lives. The trees which would most please man in a savage state would be those which had afforded him food or shelter : in a highly refined state, they would be those which afforded him the greatest amount of intellectual enjoyment, including their beauty as organic forms, their beauty as constituting a particular species of a class of organised beings, and their beauty as giving rise to pleasing or interesting associations. Perhaps the most interesting association connected with trees is that of their being employed in ship-building ; because, without ships, mankind must have remained in isolated portions, and could never have been highly civilised. It is probable, therefore, that, in every country where ships are built, and where the trees employed are high in the scale of organic beauty, the most intellectual people of that country will consider such trees as the most beautiful. In Europe and America, the oak is the tree chiefly used in ship-building ; and it is, at the same time, unquestionably fuller of variety and beauty of organic form, and of colour, and light and shade, than any other tree of temperate climates ; the oak, therefore, to the most refined of the inhabitants of these countries, may be considered as the most beautiful of trees.

There are, also, associations of a local nature connected with various species of trees, which, when known, add to the pleasure of the beholder of the particular species : for example, the antiquity of the celebrated chestnut at Tortworth, or of that on Mount Etna, or the celebrity of the platanus at Buyukderé on the banks of the Bosphorus ; or of the elm under which the founder of the state of Pennsylvania signed the first treaty with the Indians ; or of the sycamore of Trons, under which the deputies of the Swiss met in 1424, to swear to free themselves from the yoke of their lords ; lends an interest to every individual of these species. Mount Lebanon is known to every one as the native place of the cedar ; and Wilton is known to many as one of the few places in England where that tree was first raised from seeds brought from that celebrated mountain by Dr. Pococke. An individual, a general observer, but not a botanist, who had never read the history of the cedar, would feel no more interest in a young plant of that species, even if springing from one of these trees, than in a spruce fir. A knowledge of the moral and historical associations connected with trees adds, generally, to the interest of those which are still young. In general, it is thought that such trees can have but a very limited share of beauty ; and that they are chiefly worthy of admiration when they acquire such a size as to invite the painter to delineate them. This opinion can only have arisen from the general ignorance, and consequent want of interest, which prevail respecting trees as organised beings ; from ignorance of their properties in an economical and in a gardening point of view ; and from ignorance of the various associations which are connected with them. The source of interest in objects generally, consists in their positive beauty and utility ; and in their susceptibility of variation, or of changes, in their expression of this beauty and utility. Now, if we compare young trees with old ones in these respects, we think it will not be denied that young trees are objects of much greater interest than old ones. In a picturesque point of view, we allow that the old tree has an advantage ; it has also the advantage in point of shelter and shade ; and, if it were to be cut down, it would produce more timber. But will an old tree prove a source of as much interest to the possessor of it, by its variations, in consequence of its yearly increase in size, as a young tree, provided that possessor has a historical and gardening knowledge of trees ? We think not ; and we would only ask any one who is of a different opinion, whether, if he were to be allowed to have only one tree in his garden, he would prefer a tree of ten years' growth, or a tree that was already full grown ? With the latter tree the mind is carried back to times which, though interesting in some respects, it is desirable should never recur ; with the former, it is carried forward along with all the improvements which are now contemplated, or in progress, in civilised society throughout the world. For our own part, independently of all moral, historical, and economical considerations, so great is the botanical and horticultural

tural interest connected in our minds with young trees, and so delightful to us is the idea of preparing the soil in such a manner as to cause them to grow with extraordinary rapidity, that, if any one were to present us with a timbered estate, the first thing we should do would be, to cut down all the old trees, and to plant young ones.

In treating of trees individually, in the *Arboretum Britannicum*, it will form an important part of their description, to indicate the kind of expression produced by their forms, their attitudes, and their other pictorial qualities; and of their history, to record all those facts respecting each species, which may lead to interesting associations, whenever it occurs, whether it be in a young or an old state.

SECT. III. *Of the Mode of drawing Trees from Nature, in such a Manner as to give the general pictorial Expression of the Species of Tree delineated.*

IN drawing trees from nature, with a view to their introduction into landscape composition, the selection is very different from that made when the intention is to show trees as single objects. Where trees are to be introduced into landscape composition along with buildings, animals, or other trees, the symmetry or beauty of the form of the tree, considered by itself, is a matter of comparatively little importance.

A tree which is mutilated, the branches of which are ill balanced, or imperfectly clothed with tufts of foliage, will group better with other trees or objects, than a tree which is complete in itself. Such trees are perfectly well suited to the landscape-painter; but, except in the case of transplanting very large trees in order to produce immediate effect, they are of no use to the landscape-gardener, the ornamental planter, or the planter with a view to profit or use. To represent a tree mutilated or in any way imperfect, or to represent a group or whole composed of such trees, would be to exhibit what no art of the gardener could produce; and, therefore, what to him is useless, however valuable it might be in a picturesque point of view. Our object, in giving portraits of trees, has reference almost entirely to the gardenesque, to the ornamental, and to the useful. The aim of our portraits, therefore, is natural beauty and expression, with reference to the kind of tree drawn; and not beauty and character with reference to any description of graphic art. It is, in short, the beauty of truth, not local or peculiar truth, or truth with reference to any mode of depicting it; that is, not a portrait of a tree with the peculiarities which it may happen to have at a particular time and place, from peculiar circumstances; or a portrait taken to show the beauties of any particular style of sketching, drawing, or painting. It is not the portrait of a tree which has been overtopped by another tree, been improperly pruned, a part of it scorched by fire, or a part of the leaves destroyed by insects; or a portrait taken to show the picturesque effect of broken lights and shadows, breadth of masses, deep tone of colours, the sharpness of lines printed from copper or steel, or the softness of touches printed from zinc or stone. No: to draw a tree with any of these sorts of peculiarities would be in the same taste as it would be to give, as a specimen of the human being, a portrait of a man mutilated or deformed by accident or disease, or in a grotesque attitude or dress; or, as a specimen of the human face, a portrait of one disfigured with warts or pimples. This would be to portray not merely the individual instead of the species, but the individual under circumstances which had nothing to do with his character or expression, whether moral or graphic, as an individual.

It being agreed, then, that the object in drawing trees for the *Arboretum Britannicum* is to give a faithful portrait of the species, neglecting such circumstances as may be peculiar to the individual, the next point is to determine the season of the year at which the portrait is to be taken. With a view to this object, trees may be divided into three kinds: those the greatest beauty of

which is exhibited when they are in flower or in fruit; those whose greatest beauty is when they are leafing in spring, or just about to lose their leaves in autumn: and evergreens, or those which are clothed with foliage throughout the year. The last two should be drawn in autumn; and those which are most beautiful when they are in flower or in fruit, at the seasons when the flowers or the fruit are in their greatest perfection. For example, the horsechestnut ought to be drawn in June, the laburnum about the same season, and the common apple-tree, the Siberian crab, the quince, and one or two others, in autumn. Some species of the genus *Cratægus* are highly beautiful, both when in flower, in May or June, and when in fruit, from September to December; and these may be drawn at either season. Evergreens may be drawn during autumn and the whole of winter, till they begin to make their shoots in May; from that period they are unsightly for several weeks, while they are losing their old leaves and acquiring new ones; and they are uncharacteristic of the species till the new leaves and shoots have acquired that rigidity which is not produced till after complete maturity. This will be rendered particularly obvious by observing the common spruce fir, the Scotch pine, and the evergreen oak, during the growing season; say, about London, from the middle of May till the middle of June. A young spruce fir tree, drawn in May, would have a touch not unlike that of a horsechestnut; and a pine and an evergreen oak would appear to be trees of quite a different species from what they are. In general, there is a great sameness in the appearance of all trees during the leafing season, from the absence of that rigidity of foliage on the points of the shoots which gives rise to the particular touch of each species. Some deciduous trees are almost as readily known by their appearance in winter, after all the leaves have dropped, as they are in summer. Portraits of some of such trees have been taken during that season; and how very characteristic these winter portraits are, in the case of some species, is rendered obvious by the portraits of the *Gleditschia inermis*, *Ulmus americana*, and others, which will be found in their proper place.

Trees, like other objects, may be represented on paper by colours laid on with a brush or hair pencil, by ink laid on in the same manner, by lines drawn with a pen, or by lines drawn with a black-lead pencil. Whichever of these modes is employed, the object is to give the spectator a correct idea of the tree represented. The style of art in which this is done, whether by the black-lead pencil, the quill and common ink, the hair pencil and Indian ink, or by colours, is a matter of little consequence, provided the delineation be such as to raise up just ideas of the object imitated in the mind of the spectator. Different styles of art may, in this respect, be considered as equivalent to different languages, the object common to all being to convey ideas. As the most convenient and expeditious mode of drawing trees from nature is by the use of the lead pencil, we shall now proceed to give directions for its use. These directions are by no means so full as they might be; but to those who have leisure, and wish to see the subject of drawing trees by the use of the lead pencil treated in the best manner that has hitherto been done, we recommend Harding's *Elementary Art*, published in 1835, a work at once artistical and philosophic.

Previously to proceeding to the place where the tree to be drawn is situated, provide a leaf of drawing paper, or a book of such leaves, of a sufficient size to contain the pictures of the trees of the scale to which it is intended to draw them. In the case of the *Arboretum Britannicum*, we have drawn the young trees, or those which have been ten years planted within ten miles of London, to the scale of a quarter of an inch to a foot; and those which have been fifty years planted within the same distance of London, or are considered as full grown, to the scale of a quarter of an inch to 3 ft. In order to draw trees to these sizes, provide a dozen octavo leaves, and let them be sewed up together at one end, in the form of a small oblong book. Form a parallelogram on the first page, of such dimensions as to include the largest drawing which an octavo page will admit of, and next mark the scale on the boundary of this parallelo-

gram, as is done in *fig. 2*. The parallelogram suited for an octavo page is 4 in. broad, and $7\frac{1}{2}$ in. long; and the divisions are a fourth of an inch each. This parallelogram admits of a drawing of a tree 30 ft. high, with its branches extending over a space 16 ft. in diameter. These dimensions may be considered as the maximum for trees ten years planted, even under the most favourable circumstances; and therefore this size of page, and this scale, are what we have adopted for our *Arboretum Britannicum*. One page being marked as in *fig. 2*., all the other pages in the book, or any number of separate pages, may be marked in a similar manner by pricking through them from the different divisions of the scale.

For the purpose of drawing full-grown trees within the limits of the same sized parallelogram, we assume 90 ft. as the maximum height of the tree, and 48 ft. as the maximum diameter of the space covered by the branches; and, for this purpose, the same division of the inch into four parts will suffice, but with this difference, that each of these parts must now be considered as 3 ft., instead as of 1 ft. *Fig. 2*. is marked in this manner, in the inside of the parallelogram, for old trees; and it is marked in the manner before described, on the outside of the parallelogram, for young trees. In practice, it is unnecessary to mark the figures, as the draughtsman will easily bear in recollection what each division represents. For trees exceeding 90 ft. in height, and 48 ft. in width, the margin may be encroached on, to the extent of half an inch on each side, and at top and bottom, which will afford space for a tree 102 ft. high, and with a head 60 ft. in diameter.

The draughtsman may now be considered as having got his directions, as far as respects paper and scale. Having prepared his pages, he will next procure two black-lead pencils, one hard, and the other hard and black (technically H and HB), together with a foot rule and a chair. The next thing to be done is, to choose the tree and the aspect from which it is to be drawn. In making a choice, the average form, height and character of the species ought to be taken, as far as practicable; and not a specimen remarkable either for its height, or for its singularity or peculiarity; and the point from which it is to be drawn ought, if possible, to be on the south, south-east, or south-west side of it. The rule is, that the sun ought always to be behind the draughtsman, and rather to the left than to the right of him. All other circumstances being the same, therefore, when a tree is to be drawn in the morning, the draughtsman ought to place himself on the south-east side of it, at mid-day on the south side, and in the afternoon on the south-west side. The next point is, the distance from the tree at which the spectator ought to place himself. If he sits, which is, in general, the best mode, though some artists prefer a standing posture when drawing, twice the height of the tree will be a very good distance; but if he stands, and the tree has a very short trunk, say one under 6 ft. in height, it will be advisable for the artist to add to his distance from the tree once, twice, or thrice his own height; otherwise the height of his eye above the lowest branches of the tree might cause the branches to conceal more of the trunk than would be desirable. It may be useful to add, that the principle on which the distance is chosen is that of being able to see the tree as a whole, or as an entire mass of light and shade easily comprehended by the eye fixed in one position, as opposed to that of seeing it in detail, and by changing the position of the eye. Experience has shown that the eye cannot comprehend more with ease than the fourth part of a circle, whether we take this circle as a vertical or as a horizontal plane, or as a solid globe, and imagine the eye in its centre. The principle which directs the position of the sun to be behind the spectator rather than in front of him, and at his left hand rather than at his right, is, that a portion of the tree may be in light, and another portion in shade, in order to show its general form and rotundity, and that the portion in shade may always be, for reasons to be hereafter given, on the right hand. In many cases, it may be necessary to draw the tree from the north side, and, in others, to draw it when the sun does not shine: in both

these cases, the artist must supply the shade, from his knowledge of the manner in which it is supplied by the sun when it shines.

The artist having chosen his tree, and fixed his chair at the proper distance, the next step is, to measure or estimate its height. In the case of young trees, this is easily done by a 10-ft. rod, which, added to the height of a man and the length of his arm stretched above his head, will give 18 ft. ; which will cover the height of most trees of ten years' growth. In the case of old trees, the height may be ascertained by a common quadrant, by a graduated quadrant, or, which in practice, and more especially when trees are crowded together, will be found the best of all modes, by pushing up the side of the trunk a series of rods connected one with the other by small tin tubes. This, and various other modes, will be found described in *Gard. Mag.*, vol. xi. p. 548. ; and the subject will be again adverted to when treating of useful plantations, and felling timber, in Part IV. of this Encyclopædia.

The height of the tree to be drawn being measured, and supposing it to be 19 ft. 7 in., then nineteen divisions and a half of the scale are to be counted down from the top of the parallelogram, and a slight line drawn across, as at *a a*, in *fig. 2*. An estimate is next to be made of the diameter of the space covered by the branches, and also of the extent of the branches on each side of the tree. If the branches extend nearly to an equal distance on each side of the trunk, then all that is necessary is, to make a mark in the centre of the horizontal line *a a*, at *b*, in order to indicate the centre of the trunk. If, on the other hand, the branches extend much more on one side than on the other, then the first step is, to set off the total diameter, so as to reach within equal distances of each side of the page, as at *c c*, in *fig. 2*. ; and supposing the trunk to be one eighth nearer on one side than the other, then the place for its centre may be indicated at *d* on the base line *c c*.

The next step is one of some importance. The artist should go up close to the tree, examine its leaves, and make sketches of an individual leaf, and of a cluster of leaves, both to a larger scale than that to which the tree is to be drawn, and then to the same scale to which the tree is to be drawn. These sketches are merely to be considered as studies made with a view of acquiring what artists call the *touch*, or ultimate character of form, with which the tree is to be clothed. As all the masses of light and shade, and all the various forms which a tree clothed with its leaves presents in nature, result from the various disposition of one form of leaf; so, in a picture, all the imitations of these are formed by the repetition of one character of touch. Sometimes the leaves on the tree, and the touches in the picture, are so crowded as almost to obliterate each other; at other times in both they are more distinct, and the form of the leaf, and the character of the touch, may be more clearly recognised. In densely clothed trees, the form of the leaf, and the character of the touch, are most discernible at the extremities of the branches; in thinly clothed trees they are discernible throughout.

The young artist, however, must not suppose, from all this, that to represent a tree it is only necessary to know the form of its leaf and of its touch; neither must he suppose that, in making out the details of the tufting or subordinate masses of a tree, he is merely to repeat leaf after leaf: on the contrary, having a knowledge of the forms of the leaves when examined singly, and of their clustering as exhibited on the points of the branches in the general outline of the tree when examined singly, and also of the tufting, or subordinate masses, of the tree when examined singly, he must copy from nature, almost without reference to his knowledge of these details; lest, instead of making a picture of the tree as it is in nature, he should portray only his own ideas of how a tree ought to be drawn. We repeat, that he cannot too closely copy nature, and this without reference to any rules; calling to his assistance his technical knowledge of the leaves, of the touch, and of the character of tufting, only where he feels the want of it, to assist him where the appearance of nature may be of doubtful expression. In this way a man writes on any subject, without continually thinking of grammar or syntax; but when he

comes to read over what he has written, and finds some part of it obscure, or of doubtful construction, he is obliged to have recourse to his grammatical knowledge.

One of the many difficulties we have had to contend with, in getting the drawings and engravings of trees prepared for this work, is, the tendency, both of draughtsmen and engravers, to show here and there in their portraits, and sometimes, indeed, throughout the whole portraits, the distinct shapes of the individual leaves. This is just as bad as it would be, in making a drawing of a house, to give the distinct shapes of the bricks. It is true, that the surface of a tree is composed of leaves, as a house is composed of bricks; but our knowledge of these facts is not the result of our looking at the tree or house at a distance as a whole, or as a mere mass of light, shade, and colour, but of knowledge of another kind, quite otherwise acquired. Now, if the artist would only bear constantly in mind, that he is not required to convey, in his picture of the object represented, more knowledge than what a person who knew nothing of its nature might acquire by looking at it from a distance, he could not fail to succeed. The very expression, "Art," implies that the ordinary manner of conveying ideas is not to be adopted; and to show that a tree is composed of leaves, or a house built of bricks, by giving definite figures of the one or the other, is taking a license which robs art of all its charms.


It may be remarked here, that the touch of young trees is in no case so powerfully marked and characteristic in nature as that of old trees, for reasons familiar to every gardener, and which it may be well to notice here for the sake of artists. We have already said that the touch is formed by the clustering of the leaves at the extremities of the shoots. Now, as the terminating shoots of all young trees are chiefly or entirely of one year's growth, they, of course, are long, and terminate in a very few leaves, placed alternately or otherwise, round the shoot or axis, and at some distance, often an inch or more, from each other. Such leaves can never form those striking clusters which are so conspicuous in most old trees; particularly in the oak, the starry touch of which, and especially that of the *Quercus pedunculata*, which is very different from that of *Quercus sessiliflora*, is well known to every artist. The terminating shoots of old trees are generally shoots which grow only an inch or two, or, perhaps, not so much, every year; and, consequently, according to the manner in which trees grow, what is only a single leaf in the young tree of ten years' growth, is, in the spray, or terminal branches, of the old tree, a spur of several years' growth; that is, it is a spur or shoot of half an inch or more in length, protruding from the other shoot, and terminating in a cluster of leaves, perhaps half a dozen or a dozen, all radiating from the same very short axis. These radiating leaves form the touch. Any one may prove this by comparing a young oak tree with an old one. Notwithstanding the great difference between the touch of an old tree and a young tree of the same species, there is a certain distinctive character of touch even in young trees, and much more so in some species than in others; a horsechestnut, for instance, whether young or old, has a very distinct character of touch, from the large size and marked form of its leaves: so have all other trees having large leaves, and most of those having compound leaves, such as the robinias, ashes, elders, &c.

It may not be irrelevant to observe that there is as great a difference between the character of the ramification of an old tree and that of a young one, as there is between the character of their touch. There is a certain degree of sameness in the disposition of the branches of all young trees, from their tendency upwards, and perhaps still more from their being so fully clothed with leaves. Old trees, on the other hand, have generally a majority of their branches in horizontal or very oblique directions, and they are never so fully covered with leaves and spray as is the case with young trees. As a result of what we have stated, the general forms of young trees present a certain degree of sameness; while in old trees of distinct species there is generally a very

distinctive character in the general form, in the trunk, the ramification, the tufting, and the touch. Any one may be convinced of this by observing any particular species, not of very common recurrence, in the arboretum of the Horticultural Society's garden, or in that of Messrs. Loddiges, and observing the same tree of fifty or sixty years of age, at Syon, Purser's Cross, Chiswick, Upton, or any of the places noted for old American trees in the neighbourhood of London. At the same time, while we state this, we must remark that there is still a very great difference in the general form, expression, and character, of even young trees which have been no more than ten years planted. In proof of this, we again refer to the two metropolitan arboretums, and to the engravings of entire young trees, as compared with the full-grown trees, which will be found in this work. We may particularly refer both to the living specimens and to the engravings of the smallest class of trees, such as the thorn, and other *Rosacæ*; which, even in ten years' growth, are remarkably distinct and characteristic, and supply the landscape-gardener with admirable resources for planting small places, as will hereafter appear.

To recur to the subject of the touch, we shall here quote from the *Magazine of Natural History*, vol. i. p. 244., what Mr. Strutt has said on the subject

3




of the touch of the oak, and illustrate it by two engravings from his sketches. "The foliage of the oak," he says, "is particularly suited to the pencil. In those portions which are brought nearer to the sight, the form of the individual leaves (*fig. 3. a*, to the scale of 1 in. to a foot) may here and there be expressed, as shown in the sketch, which also exhibits what is technically called

the touch (*b*, to the scale of the fourth of an inch to a foot), necessary to express its character as it recedes from the eye." As a contrast to the touch of the oak, we shall give that of *Pyrus communis nivālis*, a variety of wild pear, taken at random from the sketch-book of an artist, M. Lejenne, in our employment. In *fig. 4.*, *f* is a single leaf, drawn to a scale of an inch to a foot; *g*, a cluster of leaves to the same scale; *h*, the same cluster of leaves to the scale of a quarter of an inch to a foot; and *i*, the same cluster to a scale of 1 in. to 12 ft. The last is suited for full-grown trees, and the preceding one for young trees drawn to the larger scale.



The artist having made himself acquainted with the touch of the tree, may retire to his seat and commence sketching; unless the tree should have conspicuous flowers or conspicuous fruit, as is the case with the horsechestnut if drawn when it is in flower, or with the laburnum if drawn when it is either in flower or in fruit. In examples of this kind, the artist must use the same means to acquire the touch of the flowers, or that of the fruit, as he has done to acquire the touch of the leaves.

In proceeding to draw the entire tree, the artist will first indicate the out-



lines of the masses, in the slight but accurate manner shown in *fig. 5.*, which is the commencement of a portrait of a young oak: he will then indicate the trunk, and its manner of rising from the ground; as whether perpendicular or inclined, and whether it tapers much or little. All the principal branches of the trees, visible through the leaves, should also be slightly indicated, as shown in the commencement of a portrait of *Cérasus Pâdus*, in *fig. 6.* This being done, the next step is to fill in the details of the leafing, the commencement of which, at the tops of the two trees, is indicated in *figs. 7. and 8.*; and, when this is effected for the entire trees, these two sketches only require the botanical details placed under them, to assume the appearance of the oak (*Quérecus pedunculâta*) and the bird-cherry (*Cérasus Pâdus*) given in their proper places in the series of plates forming our second volume.

The only point which remains to be considered is, that of drawing the botanical specimens. These, in the plates which form our second volume, are all given to one and the same scale; viz. 2 in. to a foot. They ought to be drawn by the artist in a book by themselves, and not on the same page with the portrait of the tree, for various reasons. In the first place, because these specimens require to be drawn at three different seasons: viz. when they are in flower;



6

when they are in fruit; and, in the case of deciduous trees, in winter, when they are in a naked state, to show the appearance of the wood at that season. In the second place, as these require to be drawn with scientific accuracy, they can only be properly done by taking the specimens home, inserting their ends in water, and drawing them with the greatest care before they begin to fade or shrivel. The specimens in flower will naturally, in most cases, be drawn first; and, because the flower is the first in the order of nature, it ought either to be put on the top of the page, or on the left-hand side of it, in order that it may come first in observing or reading. This is the reason why, in our volume of plates, we have, in the case of each young tree, always put the spring or flowering specimen on the left hand,

and the autumn or fruiting specimen on the right hand. For a corresponding reason, we have shaded the entire trees on the right hand rather than the left, because the eye, being first attracted by the light parts of an object, proceeds afterwards to the shade. Where the flowers, when fully expanded, or the fruit or leaves, when fully grown, are less than an inch across, a flower, fruit, or leaf,

7



8



of the natural size is given; and, to distinguish these full-sized specimens from such as are drawn to a scale of 2 in. to a foot, those of the full size are marked with a cross, thus +. Where a tree is of one sex, or has the sexes in different flowers on the same tree, the male flowers are marked by an *m*, and the female flowers by an *f*; and some trees, as in the case of the common ash (*Fraxinus excelsior*), the hermaphrodite flowers by an *h*. In one or two cases, it has been deemed useful to give magnified specimens of flowers or their parts; in which cases the abbreviation *mag.* is added to show this. Where the tree is deciduous, a specimen of the young wood, as it appears in winter, is given to the same scale of 2 in. to a foot. These requisites show that nearly a whole year is required, in order to draw properly the botanical specimens of any one tree.

In the case of full-grown trees, we have in general considered it unnecessary to give more than a sufficient portion of foliage to show the touch of the

tree, because along with the young tree of the same species will always be found the botanical specimens. These full-grown trees have, in every instance, been drawn in the autumn, when the leaves were ready to drop off, at which season alone they have their most forcible character. Some further observations on the subject of drawing trees will be found in the *Gardener's Magazine*, vol. xi. p. 395. to p. 412.; and whoever wishes to become master of the subject will consult the excellent work of Harding already referred to.

CHAP. II.

TREES AND SHRUBS CONSIDERED BOTANICALLY.

THE purpose for which we propose to glance at the study of trees and shrubs, botanically, or as organised beings, is, to explain our reasons for the arrangement which we have adopted in placing them together; for distinguishing between species and varieties; and for the scientific descriptions which we have adopted. It must be evident to the reader, that, before any use can be made of the history of any tree or shrub, means must be devised for distinguishing what particular tree or shrub is meant. From the want of these means, or the ignorance in this branch of knowledge of travellers, many of their remarks on trees, and other organised objects, are of little use: because it is impossible for botanists to ascertain, from their descriptions or names, to what species of tree or shrub these names or descriptions apply. There can be no doubt that the difficulties in this respect are much greater when applied to the whole vegetable kingdom, than when they are limited to trees and shrubs; and more especially when they are limited to the trees and shrubs supposed to be actually growing in Britain. But even among these, which, probably, do not greatly exceed 1500, there is, at present, the greatest uncertainty in the application of names. In genera consisting of many species, there are scarcely two of the London nurseries where the same names are applied to the same things; and what in one nursery is considered as a variety is, in many cases, elevated in other nurseries to the rank of a species. Hence it becomes necessary, in a work like the present, not only to give our reasons for the classification which we have followed, but also for the specific distinctions which we have drawn, and for the kind of descriptions and figures which we have adopted. These reasons will form the subject of three separate sections.

SECT. I. *Of the Classification of Trees and Shrubs.*

MOST authors who have hitherto produced works treating exclusively of trees and shrubs, from Evelyn and Du Hamel to the present time, have arranged them in the order of the alphabet. As we have, on various occasions (see *Encyc. of Gard.*, edit. 1835, pref.), given our objections to this mode of arrangement in any work having pretensions to be scientific, and also shown that all the advantages of an alphabetical arrangement, in the body of a work of any greater extent than a pocket dictionary, may be obtained by an alphabetical index, we shall not farther insist on the subject here; neither is it necessary for us to offer any arguments in favour of the arrangement which we have adopted, which is that of the natural system, now so generally preferred, by botanists and scientific cultivators, before all others. It may suffice to say, in favour of this system, that by grouping together objects which resemble one another in the greatest number of particulars, and which are also most alike in their qualities, every thing which is known respecting the properties, uses, or culture of any one of them, may be inferred, in a great measure, of every individual in the whole group. Hence, in the case

of trees and shrubs, or of any description of natural object arranged in this way, however much the names of the objects may in future be changed, the descriptions of the objects will always be found associated together in the same group, or in groups nearly adjoining. Hence, also, when a plant is received, the name of which is unknown, its nature may be anticipated, by observing its resemblance to some group already known.

It is a common opinion among those who know little of scientific botany, that the natural system is only adapted to those who intend to become professed in the science; and that for practical men, and for amateurs, who merely aspire to a slight degree of knowledge, the Linnæan system is the best. There never was a greater mistake. To become master of the natural system requires, indeed, much study and perseverance; but the possession of such a degree of knowledge of it as shall be of far greater use to the cultivator, to the medical man, to the traveller, and to the amateur, than the most profound knowledge of the Linnæan system, may be easily acquired by any person of ordinary capacity. In fact, every child who is in the habit of seeing a great many trees, shrubs, and plants, though he does not know a single botanical name, may be said to understand, to a certain extent, the natural system; because, to preserve order in his ideas, and to assist his memory, he is obliged to throw all the conspicuous plants that come before him into groups palpably distinct. He would thus form the three grand classes of trees, shrubs, and herbaceous plants; and among the trees he would readily distinguish, and group together in his mind, the broad-leaved and the fir-leaved, the deciduous and the evergreen, the fruit-bearing and the barren. Among herbaceous plants, he would distinguish the grasses as an obviously distinct class; the bulbous flowers as another, and so on. These divisions, so far as they go, are made on the same principles as the natural system; that is, things are brought together, or called by one common name, on account of their general resemblance; that general resemblance comprehending the whole botanical science of the individuals drawing the distinction. All the difference, therefore, between the natural system of the most learned botanist, and that of the most ignorant country labourer, consists in the former having gone more profoundly into the subject; and having his knowledge founded on principles deduced from the facts accumulated by his predecessors, and not merely on personal experience. In short, all sciences not purely abstract are founded on some simple instinct of our nature, which is perceptible in the customs, not only of ignorant persons in civilised society, but of the rudest savages.

We shall not longer occupy our time in contrasting the advantages of a natural arrangement in describing trees and shrubs, either with an alphabetical one, or with the system of Linnæus, or any other artificial system.

SECT. II. *Of the Distinction between Species and Varieties in Trees and Shrubs.*

THIS is an intricate subject; and it is one which we are well aware we shall not be able to treat in a manner that will be satisfactory to all our readers. The reason of this is to be found in the difficulty of determining what are species, or natural and permanent forms; and what forms are accidental, or the result of culture, soil, situation, disease, &c., or of cross fecundation; and because the present disposition of botanists seems to be to multiply species rather than to diminish their number.

When we look into a modern catalogue of plants, we are astonished, and almost confounded, by the great number of specific names which are ranked under one generic name. If we endeavour, by inspecting the names more particularly, to discover any relationship between them, we are utterly at a loss. One name is, perhaps, an adjective denoting colour, or some other property belonging to the plant, another indicates the native country of the species; and a third shows that it has been named in commemoration of

some place, or of some individual. If we look at the column, in such catalogues, which indicates the native country of the species, the difficulty is increased rather than lessened: perhaps a native of the tropics is placed next a plant from the frigid zone. In this, as in similar cases of collecting knowledge, the first step is to accumulate facts, and the second is to generalise on them. Hitherto it would appear, that, as far as regards species and varieties, the great object of botanists has been to increase their number, without much regard to grouping them according to their relationship. It is not for us even to try to remedy this evil in respect to all the species and varieties of plants; but we propose to attempt to do so, in as far as respects the hardy trees and shrubs of Britain. We shall notice, in succession, the subjects of species, races, varieties, and variations; and we shall then offer some remarks on mules, hybrids, and what are called botanical species.

A *species* is defined, by Dr. Lindley, to be "a union of individuals agreeing with each other in all essential characters of vegetation and fructification; capable of reproduction by seed, without change; breeding freely together, and producing perfect seed, from which a fertile progeny can be reared." (*Introd. to Bot.*, p. 365.) This, we believe, is the general definition of a species by botanists; but it evidently requires some modification; for, in the case of many cultivated annual plants, the variety or race is reproduced from seed; and, consequently, if reproduction from seed were considered as a certain test, red, white, woolly-eared, and smooth-chaffed wheat, would be so many distinct species; as would the different varieties of cabbage, turnip, common lupine, &c. In like manner, also, the different varieties of particular species of cultivated fruit trees, might be deemed species; for it is certain that seedlings from such varieties of fruit trees, when no cross fecundation has been effected, always bear a nearer resemblance to the variety which produced the seeds, than to any other variety, or to the original species. The truth we believe to be, that trees and shrubs are subjected to the same law, in regard to the reproduction of varieties from seed, as annual plants; though, from the varieties of the former seldomer falling under our observation, and requiring a longer time to come to maturity, we have not the same opportunity of becoming sufficiently impressed with the identities of their natures as to be able to generalise on them. On examining a number of individual trees or shrubs, raised from seed (say, for example, oaks in an oak wood, or hawthorns in a hedge which has not been cut), we shall not find two individuals exactly alike, either in foliage, in flower, in fruit, in mode of growth, or even in the earliness or lateness of budding, flowering, ripening the fruit, or dropping the leaves. We have no doubt, reasoning from the analogy of the wheat, that, if the acorns or haws of any marked variety in such a wood or hedge as that mentioned were sown, and the plants reared to maturity, they would be found (unless cross fecundation had been accidentally or artificially effected) more like the parent variety than any other in the wood or hedge, just as in the case of seedlings from varieties of wheat, cabbage, or fruit trees.

These may be called cultivated varieties, or, according to De Candolle, *races*; but there are others, which we shall call *accidental varieties* that we are not so certain can be continued by seed. For example, there are weeping varieties of certain trees, such as the common ash; and fastigate varieties of others, such as the Exeter elm, the Cratægn's *Oxyacantha stricta*, and the Lombardy poplar (*P. fastigiata*), which we believe to be only an accidental variety of *P. nigra*: these varieties, we think, would scarcely come true from seed in every, or even in many, cases; though we have no doubt they would in some. Variegated trees and shrubs, we should suppose, would not always come true from seed, any more than variegated annuals or bulbs; but we have no doubt that, as in the two latter cases, a certain proportion of the progeny would be variegated in trees and shrubs, as well as in herbaceous plants. The raising of seedlings from such accidental varieties then, will prove that they are not entitled to rank with cultivated varieties or races.

The difficulty of being able to determine what is only a variety, and what may be ranked as a species, is ably pointed out by Dr. Lindley, in the following passage: — “The manner in which individuals agree in their external characters is the only guide which can be followed in the greater part of plants. We do not often possess the means of ascertaining what the effect of sowing the seed or mixing the pollen of individuals would be; and, consequently, this test, which is the only sure one, is, in practice, seldom capable of being applied. The determination of what is a species, and what a variety, becomes, therefore, wholly dependent upon external characters, the power of duly appreciating which, as indicative of specific difference, is only to be obtained by experience, and is, in all cases, to a certain degree, arbitrary. It is probable that, in the beginning, species only were formed; and that they have, since the creation, sported into varieties, by which the limits of the species themselves have now become greatly confounded. For example, it may be supposed that a rose, or a few species of rose, were originally created. In the course of time these have produced endless varieties, some of which, depending for a long series of ages upon permanent peculiarities of soil or climate, have been in a manner fixed, acquiring a constitution and physiognomy of their own. Such supposed varieties have again intermixed with each other, producing other forms, and so the operation has proceeded. But, as it is impossible, at the present day, to determine which was the original, or originals, from which all the roses of our own time have proceeded, or even whether they were produced in the manner I have assumed; and, as the forms into which they divide are so peculiar as to render a classification of them indispensable to accuracy of language; it has become necessary to give names to certain of those forms which are called species.” (*Ibid.*, p. 366.)

The secret of the great number of names of species which at present form the bulk of names in our catalogues is to be found in what follows from the same author: — “Thus it seems that there are two sorts of species: the one, called natural species, determined by the definition given above; and the other, called botanical species, depending only upon the external characters of the plant. The former have been ascertained to a very limited extent: of the latter nearly the whole of systematic botany consists. In this sense a species may be defined to be ‘an assemblage of individuals agreeing in all the essential characters of vegetation and fructification.’” (*Ibid.*, p. 366.)

The difficulty of determining what is a species, and what is a variety, as far as concerns plants of culture, may here be considered as diminished; but, since it is acknowledged by Dr. Lindley, that nearly the whole of systematic botany consists of what are called “botanical species, depending only upon the external characters of the plant,” the idea of determining, with any thing like absolute certainty, what is a species, at least a botanical species, and what is a variety, seems almost hopeless. The “whole question,” Dr. Lindley observes, “lies with the word essential. What is an essential character of a species? This will generally depend upon a proneness to vary, or to be constant in particular characters, so that one class of characters may be essential in one genus, another class in another genus; and these points can be only determined by experience. Thus, in the genus *Dahlia*, the form of the leaves is found to be subject to great variation; the same species producing, from seed, individuals, the forms of whose leaves vary in a very striking manner: the form of the leaves is, therefore, in *Dahlia*, not a specific character. In like manner, in *Rosa*, the number of prickles, the surface of the fruit, or the surface of the leaves, and their serratures, are found to be generally fluctuating characters, and cannot often be taken as essential to species. The determination of species is, therefore, in all respects, arbitrary, and must depend upon the discretion or experience of the botanist. It may, nevertheless, be remarked, that decided differences in the forms of leaves, in the figure of the stem, in the surface of the different parts, in the inflorescence, in the proportion of parts, or in the form of the sepals and petals, usually constitute good specific differences.” (*Ibid.*, p. 366, 367.)

The subject of species and varieties has, in our opinion, been placed in the clearest light, by Professor De Candolle, in his *Théorie Élémentaire*, and in his *Physiologie Végétale*. In the latter work, this celebrated botanist recognises in plants — species, races, varieties, and variations.

Species. — Under the name of species, that is what we consider aboriginal species in contradistinction to the botanical species of botanists, Professor De Candolle unites all those individuals which bear a sufficient degree of resemblance to each other, to induce us to believe that they might have originated in one being, or one pair of beings. The degree of resemblance which authorises us to unite individuals under the denomination of a species varies much in different families; and it often happens that two individuals which really belong to the same species differ more between themselves in appearance, than others which are of distinct species: thus, the spaniel and the Danish dog are externally more different from each other than the dog and the wolf are; and many of the varieties of our fruit trees offer more apparent differences than are found between many species. (*Physiol. Végét.*, vol. ii. p. 689.)

If all the alleged species and varieties of any tree, shrub, or plant were collected together, and cultivated in the same garden, however numerous were the varieties, and however remote they might appear to be from the original species, it would be practicable, after a series of years, to decide with absolute certainty what were aboriginal or fixed features, and what features were variable. For example, in the case of the apple, notwithstanding the thousands of varieties in cultivation throughout the temperate regions of the world, and the immense difference between some of the varieties (for example, the Alexander or the Hawthornden and the original crab), and even the great difference between the crabs of different parts of Europe, yet in no case is there any danger of one of these varieties being mistaken for a pear. One general character of leaf, flower, and fruit is common to the whole of them, though it may not be easy to define in what this essential character consists, in such a manner as to render it observable to any one who had not seen a great number of varieties of apples and crabs. Again, in the case of the common hawthorn, though some of the varieties have deep red fruit, others pale red fruit, others yellow, and others black fruit; and though some varieties of hawthorn have drooping branches, and others have them rigidly erect and fastigiata; though some have the leaves finely cut, and others obtusely lobed or scarcely lobed at all; though some are polygynous, and some are monogynous; yet there never can be any difficulty, when all these varieties are before us, in determining that they belong to one and the same species. The same observation will apply to the numerous varieties of the cockspur thorn, which now figure in our catalogues as distinct species; and we think that it might be applied to many varieties of the genera *Fraxinus*, *Ulmus*, *Salix*, *Quercus*, *Pinus*, and to various others. Could we bring before us, into one plantation, all those ashes which are natives of America, and watch them for a sufficient number of years, we have no doubt that we should not find it more difficult to assign them to one species, than we do the different varieties of the European ash to the *Fraxinus excelsior*. All the elms of Europe, we are inclined to be of opinion, may be reduced to only three species; and we much question if, on De Candolle's principle of determining what a species is, there would be more than a tithe of the names which are ranked as such under *Salix*, *Quercus*, &c.

Races. — A race in the vegetable, as in the animal kingdom, De Candolle observes, “is such a modification of the species, whether produced by exterior causes, or by cross fecundation, as can be transmitted from one generation to another by seed.” Thus, among all the cultivated vegetables and fruits, both of the garden and of the field, the greater number of sorts may be considered as races, because they may all be continued by seed; the culture given and other circumstances being the same. If the culture were neglected for a series of generations, there can be no doubt that the race would revert to the abori-

ginal species; because a tendency to this has been found to take place both in plants and animals.

Varieties.— A variety differs from a race, in not being susceptible of propagation by seed, at least with any thing like constancy and certainty. For example, the jargonelle pear may be continued by seed; but a jargonelle pear with variegated foliage could not be so propagated with certainty. We allow, however, that, if a great many seeds taken from the fruit of a jargonelle pear with variegated leaves were sown, some of the plants so raised would probably show variegation in their foliage. The same thing, we think, would take place in the case of sowing the seed of variegated hollies, or of fastigate or pendulous-branched plants, but to what extent it is impossible to say. It certainly would not take place to such an extent as to confound varieties with races, or to render it desirable to propagate varieties in this way; and, consequently, varieties are always propagated by some modification of division, such as by cuttings, layers, grafting, &c.

Variations differ from varieties in not being transmittible by any mode of propagation. They are always produced by local circumstances operating on the individual; and the moment these circumstances are changed, the variation disappears. For example, plants grown in the dark will have their leaves white; other plants with hairy leaves, when grown in water, will have their leaves smooth; and the hydrangea, grown in a certain description of soil, will have its flowers blue: but, remove the plant with white leaves to the light, and place the plant grown in water in dry soil, and the hydrangea in common soil, and it will be found that the leaves of the first will become green, and those of the second hairy, and that the flowers of the hydrangea will resume their natural pinkish hue.

Mules and Hybrids.— Some confusion exists as to the use of these terms, when applied to plants. The term mule, we think, ought to be limited to such hybrids as are raised between different aboriginal species, and which it is believed are not susceptible of propagation by seed: such, for example, as the *Pyrus spuria*, which seems to be a hybrid between *Pyrus Sorbus* or *Aria* and *Pyrus Chamæmépilis*. The term hybrid, on the other hand, we think, ought to be limited to the produce, by cross fecundation, of different races and varieties of the same species. Every one knows that this is one of the most important elements of culture, having given rise to the most valuable garden flowers, table fruits, culinary vegetables, and agricultural plants.

Botanical Species.— It will be seen, from the preceding remarks, that we follow De Candolle in denominating what Dr. Lindley and other British botanists distinguish as botanical species, races, or varieties.

It is not to be supposed, however, that we undervalue botanical species, or that we either deny the distinctness of many that exist, or the propriety of having different names for them, and keeping them distinct. On the contrary, to compare plants with men, we consider aboriginal species as mere savages, and botanical species, or, according to De Candolle's classification, races and varieties, as civilised beings. What, then, it may be asked, is our object in endeavouring to show that many of our botanical species are only varieties? We have two objects in view; and both, we think, are very important ones. In the first place, by confounding varieties or garden or botanical species with aboriginal ones, a beginner, ignorant of the extent to which this is done, cannot make a judicious selection; and while, in the case *Práxinos*, for instance (of which there are, in reality, as we think, only three species known, exclusive of *O'rus*), if he wished to select, perhaps, three sorts, he might, instead of selecting the three really distinct species, which would give him a complete idea of the genus, fix on three of the varieties of *F. americana* or of *F. excelsior*, which would only give him an idea of one species. In the second place, we wish to prevent beginners, in the study or cultivation of trees, from puzzling themselves unnecessarily to make out the minute differences which distinguish what are called botanical species; believing, as we do, that it is impossible to make out many of these from the specific charac-

ters given of them in botanical works. The nicety of these distinctions has we know, deterred numbers from the study of practical botany; and has prevented others, who have had the courage to proceed, from ever hoping to attain any satisfactory result. It has also (and this we consider to be the most important part of the evil) prevented many persons from forming collections of trees and shrubs, by inducing them to believe that such collections could never be made anything like complete, without incurring an expense greatly beyond what is really necessary. Instead of this being the case, the number of hardy trees and shrubs is so small, when compared with that of hardy herbaceous plants, or stove or green-house plants, that there cannot be the slightest difficulty in becoming acquainted with all the species, provided these and the varieties are only seen together; and the cost of as complete a collection of species as can be procured in the London nurseries is such as to be within the reach of every planter of the grounds of a villa of a single acre in extent.

The mode by which we propose to attain these objects is very simple. We shall retain the botanical species and varieties in the catalogues, so far as we believe them to exist; but we shall, in every case, place before them the name of the aboriginal species to which they belong: for example, in the case of the genus *Fraxinus*, which, in our *Hortus Britannicus*, appears to consist of 41 species and 12 varieties, we shall rank 30 of the species under the head of *F. americana*, two of them under the head of *F. lentiscifolia*, and the remainder under the head of *F. excelsior*. It may be asked, whether it would not be better at once to make distinct genera of these three species? To which we answer, that it would not; because they are all so obviously of the same general appearance, as evidently to belong to the same family. There would be the same objection to separating the oak family into different genera; though we think it highly probable that there are not a dozen aboriginal species of oak in the world. Every division, or conglomeration, in botany, that can assist the mind to generalise, at the same time assists it in particularising; and it will be found much more easy, after throwing all the races or varieties of *Fraxinus americana* into one group, to distinguish them from each other, than by leaving them as distinct species, and having the trouble of distinguishing them, not only from other races or varieties of *F. americana*, but also from all the races or varieties of *F. excelsior*.

Such are the principles which we have adopted, to guide us in arranging species, races, and varieties, from a perfect conviction of their truth. If we had not had an opportunity of observing, for several years past, the collections of trees and shrubs in the neighbourhood of London, and of studying them at every season of the year, with a view to the production of this work, we should never have been able to arrive at these principles, or to adopt them from others, with any degree of satisfaction to our own minds. We are, however, perfectly satisfied that we are in the right path; and we feel convinced that all practical botanists who have had an opportunity of making similar observations, and who have made them, will approve of our arrangement.

SECT. III. *Of the Mode of describing Trees and Shrubs.*

IT is foreign to the object of this work, to enter any farther into botanical science than becomes necessary to elucidate the reasons which have induced us to depart, in any particular, from general practice. It will readily be conceived, from what has been stated in the preceding section, that we attach no great value to what are called the specific characters of botanical species; that is, of what we shall distinguish as races in some cases, and varieties in others. The reason is, that we do not think it is often practicable to discover a species or race by such characters alone. The specific character of an aboriginal species we consider in a different point of view; for, as we

think all aboriginal species must be decidedly distinct, so we think it practicable to render this distinctness so obvious, in the few words which constitute a specific character, that the name of a plant may be discovered by it. To recur to the genera *Fraxinus* and *Cratægus*, we will ask any botanist, either practical or theoretical, whether, from the specific characters of the botanical species of *F. americana* or of *C. Oxyacantha*, he could discover the individuals to which those names are intended to apply, without having recourse to dried specimens or engravings? We ask the same question with reference to most of the alleged species of *Salix*, *Ulmus*, *Quercus*, *Pinus*, and *Rubus*. We admit that many of these botanical species, or varieties as we consider them, may be made out from lengthened descriptions; but we deny the practicability of doing this, in many cases, from short specific characters. That we may not be misunderstood, we refer more particularly to the genera *Quercus*, *Salix*, *Ulmus*, and *Rubus*; and even to *Tilia*.

But, though we question the utility of specific characters to botanical species as such, we are of opinion that they may be of some use when applied to these species, as being varieties of an aboriginal species, and indicating that they are such. For example, in the case of the specific character of *Fraxinus pubescens*, *caroliniana*, *lanæca*, &c., as absolute species, and to be compared with different botanical species of the same aboriginal species, and also of *F. excelsior*, we think it would be extremely difficult, if not impossible, to apply them; but, if it were known that these botanical species were only varieties of *F. americana*, the difficulties of distinguishing them would be greatly diminished. For this reason we shall, in many cases, adopt the specific characters of botanical species given by botanists, adding to them such descriptive particulars as our own observation has enabled us to supply.

We may here refer to two causes, which have not only contributed to the great imperfection of the specific characters of botanical species; but which have been the means of multiplying the number and descriptions of these species in books, to an extent which, we are persuaded, does not exist in nature. One of the practices to which we allude is, that of describing species from dried specimens only; and the other, that of mistaking varieties for species by collectors. We admit that the first of these practices is unavoidable in the infant state of botanical science; and that it must necessarily be continued, till botanists shall rise up in every country in such numbers, and of such acquirements, as to be able to describe the plants of every country from nature; or till all the species, or all the alleged species, of every genus of plants shall be assembled together in one spot, and what are really aboriginal species shall be determined, after observing them for a series of years. Happily, both these results are in progress of attainment: botanists are beginning to spring up in every civilised country, or to emigrate from old, and settle in newly discovered countries; and, in all the wealthiest governments of Europe, assemblages of plants are being made in botanic gardens. If the directors of these gardens were to cooperate, and each to undertake the collection and the study of one or more genera of hardy plants, we should, at no distant period, be able to say what are really species, and what are not. If botanic gardens were established in every country and climate of the world, and the whole of the directors of these gardens were to act in concert (which concert, being quite distinct from political associations, would not be objected to by any government), in each garden might be assembled all the alleged species or varieties of at least two or three genera, those being selected for which its climate, situation, soil, and extent were most suitable; and, after a few years, the aboriginal species, and the more prominent varieties, might be determined on. In the meantime, this process might be commenced in many of the botanic gardens already established in the temperate regions of the world; and we have already shown (p. 192.) how, in every country in these regions, the determination of species, and their nomenclature, might be effected, as far as respects hardy trees and shrubs.

When the natural system of botany comes to be more generally understood

and cultivated, there will not, we may presume, be that desire, which now seems to exist among botanists, to increase the number of species. Formerly, and more especially among the disciples of Linnæus, the great business of the botanist was to collect, name, and describe plants. These were then the highest departments of the science; but, now, the anatomical, physiological, and chemical studies of plants occupy that station; and the naming and describing of species is considered as comparatively mechanical.

The other cause which has contributed to increase the number of supposed species is, the natural eagerness of botanical collectors, sent abroad in order to discover novelties, to find something new, in order to answer the end for which they were sent out. This is very natural: and where there is a strong desire for, and also an important interest concerned in, obtaining anything, either the thing sought for, or something like it, will be found. Hence the young and ardent collector will seize upon every variation produced by climate, soil, situation, age, or even accident, to add another specimen to his herbarium; which enables the botanist at home to add another name to the number of his species. This we believe to be much more frequently done from practical inexperience, than from any intention to deceive; so different is the appearance which plants present in a wild state and in a state of cultivation, and, often, in one country from what they do in another country; and so difficult is it to judge of an entire tree by a dried specimen, perhaps only a few inches in length. This state of things, in the comparative infancy of botanical science, is perhaps unavoidable; and it is, doubtless, erring on the safe side, to collect and bring home every thing that can be at all considered as distinct, leaving it to cultivators and botanists to determine afterwards whether it is really so. It is proper, however, to notice this state of things, to aid in accounting for the present state of confusion and uncertainty in the names and characters of trees and shrubs; and to show the little faith that is to be placed in botanical descriptions drawn up from dried specimens of any kind, and more especially from those procured by inexperienced collectors. If this may be considered as applicable to plants generally, it is more particularly so in the case of trees and shrubs; which, from the long period which they require to attain maturity, naturally assume very different appearances under different circumstances: and which, therefore, require to be studied, not only in the same locality, but in different localities, for a number of years, before any decided opinion can be pronounced respecting which are species and which are varieties.

It will not, we trust, be supposed, from these observations, that we intend to set ourselves up as a model for imitation, in determining species and describing them; on the contrary, we value the *Arboretum* part of this *Encyclopædia* much more, as containing only the names of such things as we know to be really distinct, and actually in existence in England, than for its pretensions in a purely botanical point of view.

CHAP. III.

TREES AND SHRUBS CONSIDERED WITH REFERENCE TO THEIR USES IN THE ECONOMY OF NATURE AND TO MAN.

THE large proportion which the ligneous vegetation of the earth's surface bears to its herbage, and the immense extent of the forests in comparison with that of the meadows, pastures, or plains, which it contains, seem to indicate that trees and shrubs act an important part in the economy of our globe. In countries uninhabited by man, the influence of forests must be on the climate, on the soil, and on the number of wild animals and herbaceous vegetables. In civilised countries, to these influences must be added the

relation in which trees and shrubs stand to man. It is not our intention to enter farther into these subjects here, than may be necessary to show to what circumstances, in the economical history of trees, we ought chiefly to direct our attention, in composing the history of each particular species. The subject may be divided into two sections.

SECT. I. *Trees and Shrubs considered with Reference to uncultivated Nature.*

It appears highly probable, that the greater part of the surface of our globe has been, at one time, covered with wood; because, among other reasons, coal is found in almost all countries; at all events, it is certain that this has been the case with the greater part of the temperate regions of the world at no very distant period. North America was, till lately, almost entirely covered with trees and shrubs, and presented few naked surfaces, except those of the alluvial deposits on the banks of its larger rivers; and what was so recently the state of America must, we may reasonably suppose, have once, at least, been that of every other part of the world.

The influence which a predominance of forest must have in a country uninhabited by man must have extended to the animals, the herbaceous vegetables, the soil, the waters, and the climate. To wild animals of every kind, especially to those of the more ferocious description, forests have, in all countries, furnished shelter, and, in a great measure, food: birds, insects, and reptiles are the more common inhabitants of forest scenery. Herbaceous plants are, for the most part, destroyed by dense forests; but some kinds, such as epiphytal lichens, mosses, and, in some cases, *Orchidæ*, are encouraged by the thickness of the shade, and the moist heat which prevails among the trunks and branches of the trees. But the great influence of forest scenery in a wild state is on the soil; and, in this point of view, natural forests may be regarded as a provision of nature for preparing the earth's surface for the cultivation of corn, and of the other plants which constitute the food of man, and of domestic animals. It is unnecessary to show how the soil is furnished with that organised matter, on which alone perfect plants can live, by the decay of leaves, and, ultimately, by the decay of trunks and branches. The waters of a country, the rivers and lakes, are necessarily affected by the state of the woods of that country. These woods must, in all cases, act more or less as a sponge in retaining the water which falls on them; and water must thus be supplied more gradually to the rivers, in countries covered with wood, than in countries which are cleared, and regularly drained. The influence of forest scenery in increasing the moisture of the atmosphere, and in preventing a climate from being so hot in summer, and so cold in winter, as it would otherwise be, is well understood, and, in such a slight outline as the present, requires only to be mentioned.

The use of studying the influence of trees in an uncultivated country is, to afford useful hints with reference to the planting or thinning of them in countries which are civilised. That which takes effect on a grand scale, where forests cover many thousand acres, must operate more or less in the same manner where they extend only to hundreds, or even tens, of acres; and, consequently, this influence must be kept in view in the formation of plantations, both useful and ornamental. If the forests and plantations of Britain are no longer of such an extent as to afford a shelter for wolves and hyenas, they still harbour foxes, polecats, snakes, and other noxious animals, and several kinds of carnivorous birds, such as the hawk. The forests in France and Germany still contain wolves and wild boars; and, on most parts of the Continent, the forest is the place of refuge to which man flies for concealment after the commission of crime. (See *Gautieri Dello Influsso dei Boschi*, &c.) If forests in a wild state supply food to birds and insects, in a civilised country birds and insects may be expected to abound more or less wherever there are trees and shrubs to supply them with food and shelter.

The same may be said with reference to different species of reptiles. In Belgium and part of Holland, the caterpillars of some species of moths are so abundant in the woods at a particular season every year, that it is a part of the business of the government police to see that they are destroyed. Hence the advantage of knowing what trees and shrubs are obnoxious to particular insects, and what insects attack trees and shrubs generally. The total destruction of herbaceous plants in dense forests teaches us, that, where we wish the grass or other herbage under trees to thrive, we must plant the trees thinly; and the influence of the decay of leaves, branches, and trees, in adding to the soil, teaches us how barren soil may be improved by trees; and this natural effect has been imitated by trenching down entire plantations of Scotch pine, grown on extremely poor soils in some parts of Scotland. Undrained woods, and especially copse woods, are known to retain the water which falls on them much longer than open groves or plains; and, as increased exhalation and evaporation must be going on from such woods during the period of retention, and increased moisture must be thus produced in the atmosphere, the circumstance may demand consideration in planting extensive shrubberies near dwelling-houses; and, more especially, in planting such as are intended, by frequent digging, always to present a surface of naked loose soil. The influence of trees in modifying both the temperature and moisture of the atmosphere, in civilised countries, and in artificial scenery, is generally known; and this knowledge should not be lost sight of in the disposition of trees and shrubs about a house, more especially in low situations. There is great reason to believe that many country residences in England, naturally healthy, are rendered unhealthy by the superabundance of trees and shrubs, and by the quantity of dug ground close to the house. The insects which infest the rooms of a house are also very much increased by the proximity of wood.

From trees and shrubs in a wild state we can only truly learn their aboriginal natures; because plants, like animals, begin to change their habits as soon as they are taken into cultivation. The fact that this change takes place ought to be borne in view, when speaking of the native soils and situations of different species; because, if it is desirable to improve these species, it may be necessary or advantageous, for that purpose, to place them in a different soil or situation from that in which they are found in a wild state. There are certain soils and situations, however, in which plants are found in a wild state, that can hardly be improved by art; these are peat bogs, or peaty soils, such as are found in North America, and in alpine situations. We mention these particulars merely as a few, among a great number, to which attention ought to be directed in giving the history of particular species of trees and shrubs, and in treating of their introduction into useful or ornamental plantations.

SECT. II. *Trees and Shrubs considered with Reference to Man.*

So various and so important are the uses of trees and shrubs to man, that to say much on the subject here is altogether unnecessary. It must be obvious, that to state what these uses are, in the case of every particular tree and shrub treated of in this work, will form an important part of the information given respecting it. It is not necessary, in every case, to mention how the different kinds of wood are used in particular arts or manufactures; but it is necessary to know, not only the particular sorts of timber, but what modifications of these sorts are best for particular purposes. For example, in the case of ship-building, it is not only necessary to know the different kinds of trees in demand by ship-builders, but the different purposes for which different parts of a tree, and different forms of its trunk or branches, are adapted, and to which they are applied.

Though the timber is the principal part of trees and shrubs which is employed in arts and manufactures, yet, in many cases, the other parts of a tree,

such as the bark, leaves, flowers, fruit, &c., are of importance. Not only are trees used in their different parts after being felled, but, in some cases, a part of their products is gathered yearly; and some sorts, both of trees and shrubs, are in perpetual use in a living state, as fences for separation or enclosure, as avenues for shade, and as belts or screens for shelter. The ornament which trees and shrubs afford to gardens and grounds may also be considered as an important part of their use.

The rearing and culture of trees form an important part of their economical history; and require to be treated of, not only in the history of each individual species, but when treating of trees collectively in plantations. The commencement of the process of rearing is with the gathering of the seed, or the taking off of the cutting, or the forming of the layer; and the termination of the process of culture is with the felling of the tree, or the cutting down of the shrub.

CHAP. IV.

SUMMARY OF PARTICULARS TO BE TAKEN INTO CONSIDERATION IN PREPARING THE DESCRIPTION, AND NATURAL AND ECONOMICAL HISTORY, OF TREES AND SHRUBS.

IN the three preceding chapters we have treated of the science of trees, as pictorial objects, or parts of general scenery; as organised beings, or botanical objects; and as forests or plantations, influencing the physical circumstances of a country, and the condition of man. Our object in those chapters was, to give a general idea of the extensive nature of the study of trees; and in the present chapter we propose to enumerate all the particulars which require to be taken into consideration in giving the specific character, description, history, and uses of each individual species, race, or variety. These particulars, arranged in the order in which they will stand in the succeeding part of this work, are as follows:—

1. *Classification.* We shall adopt the natural system, chiefly for the sake of aiding us in generalising on the genera and species which each order and tribe contains; and we shall refer, for the definitions of these orders and tribes, to the writings of Professor De Candolle and Dr. Lindley, and more especially to the *Prodromus* of De Candolle, and Lindley's *Introduction to the Natural System*, and to his modification of that work in his *Key to Structural, Physiological, and Systematic Botany*, published in 1835.
2. *Genera.* The genera of each order or tribe, with their characters, will be given immediately after the general character of the order or tribe, in the manner of De Candolle, and as adopted by G. Don, in his edition of *Miller's Dictionary*.
3. *Distinctive Characters.* The species, races, or varieties, of each genus, will be enumerated immediately after the enumeration of the genera; and their distinctive characters will be given, with the English name, habit, colour of the flower, and time of flowering, and year of introduction into Britain.
 - A. *Identifications.* These are references to some of the principal works in which the same plant has been described under the same name.
 - B. *Synonyms.* These will be given to as great an extent as can be done with certainty, or apparent advantage. Not only will the scientific synonymes be given, but also those in common or ancient use in this country, and the popular names in other countries.
 - C. *Derivations* will be given, not only of the generic and specific names, but of all the synonymes, where doing so is likely to prove either instructive or interesting.

4. *Engravings.* Engravings of certain species and varieties, to a scale of two inches to a foot, with the flowers and other parts which are less than an inch in diameter of their natural size, are given along with the text.

A. *Engravings of the Trees only* are given in the plates which form a separate volume. The engravings in these plates are of three kinds : first, there is the general figure of the tree, after being ten years planted in the climate of the environs of London, to a scale of one fourth of an inch to a foot. Secondly, on the same plate with the entire tree, are given engravings of botanical specimens in flower, and in fruit, with the winter's wood in the case of deciduous trees, to a scale of 2 in. to 1 ft. ; and, when the flowers or fruits are smaller than an inch across, they are given of their natural size. Thirdly, engravings of full-grown trees of a number of the species are given ; and, on the same plate, a specimen of the leaves to the usual scale of 2 in. to 1 ft. The use of the engravings of the entire trees, of ten years' growth, is to give a palpable idea of the comparative progress of hardy trees during that period, in a given soil and situation ; and the use of the portraits of full-grown trees, all of which are taken from individuals within ten miles of London, is to give a palpable idea of the magnitude and general figure which the particular species assume, when full grown. These full-grown trees are drawn to the scale of one twelfth of an inch to a foot. The trees, whether full grown, or of ten years' growth, and the botanical specimens given along with them, are always referred to as plates ; while the engravings of those species and varieties which are given along with the text are referred to as figures.

B. *Pictorial Signs.* At the commencement of each genus, or sectional division of a genus, will be placed a pictorial sign, or signs, indicating whether the species to be described are trees or shrubs, deciduous or evergreen, climbers, twiners, trailers, or creepers, &c. ; and also to indicate whether they are of the first, second, or third degrees of height, of each of these habits. Before each species and variety will be placed one of the signs in our *Hortus Britannicus*.

5. *Descriptions.*

A. *Descriptive Details.* These will commence with the root, and proceed in the order of stem, leaves, stipules, inflorescence, bracteas, flowers, and fruit.

a. *The Root* will be considered in regard to figure, quality, substance, bark, duration, direction, rootlets, fibres, spongioles, susceptibility of producing buds when made into cuttings, liability to throw up suckers naturally, magnitude relatively to age, soil, native habitat and artificial location ; impulsion, or when it is in most vigorous growth ; and hibernation, or when it is in a state of rest.

b. *The Stem* will be considered in regard to its figure, direction, duration, articulation, surface, bark, ramification, branches, branchlets, twigs, height relatively to age, native habitat, and artificial locality.

c. *The Leaves* will be viewed with reference to their vernal, internal structure, figure, articulation, insertion, circumscription (that is, outline, base, and apex), surface, subface, venation, direction, colour, texture, and duration. *The Petiole* will be considered as to its absence or presence ; and in the latter case its form, surface, texture, length, breadth, variation, duration, &c.

d. *The Stipules* will be considered with reference to position, texture, surface, insertion, figure, magnitude and duration.

e. *The Inflorescence*, or mode in which flowers are disposed upon a plant, will be examined as to its kind and position.

f. *The Bracteas* will be examined under conditions similar to those prescribed for the leaves.

g. *The Flower* will be considered in regard to first appearance, first expansion, colour, magnitude, length of time it continues expanded before it begins to fade, whether the flowering of the whole plant is

simultaneous or continuous, the number of the flowers in proportion to the leaves or surface of the plant, and whether the flowers die off rapidly or slowly. *The Calyx* will be examined as to texture, structure, figure, station relatively to the ovarium and the axis of the flower, surface, size, proportion to the corolla, colour, æstivation, and duration. *The Corolla* will be viewed in its "structure, figure, station with respect to the ovarium and axis of inflorescence and adjacent parts, surface, æstivation, size, colour, proportion to the calyx and stamens, and venation." (*Liubl. Introd.*, p. 141.) *Stamens, Filaments, Anthers, Pollen, Disk, Ovaries, Orules, Styles, and Stigmas*, will all be examined with a view to generic and specific definitions, as well as to general description.

- b. *The Fruit* will be examined as to "texture, form; whether naked or covered with the remains of the floral envelopes; whether sessile or stipitate; mode of dehiscence, if any; number of its valves and cells; situation of the placenta; nature of its axis; number of its seeds" (*Ibid.*, p. 442); magnitude in a wild state, in cultivation; whether generally abundant or not abundant, conspicuous or not conspicuous; colour of the surface; when ripe, length of time in ripening, and duration on the tree.
- i. *The Seed* will be considered scientifically in the generic and specific descriptions, and generally with a view to popular description. Scientifically, it will be examined as to "position with respect to the axis of the fruit, mode of insertion, form, surface; the texture and nature of the testa, arillus, and other appendages, if any; position of the raphe and chalaza. Albumen, its texture, if any. Embryo, its direction, position with respect to the axis of the fruit, to the hilum of the seed, and to the albumen; the proportion it bears to the mass of the latter; the form of its cotyledons and radicle; its mode of germination." (*Ibid.*, p. 442.) Popularly, the seed will be considered as to magnitude, form, integuments, facility or difficulty of separation from its envelopes, peduncles, conspicuousness or inconspicuousness on the tree, duration of the vital principle, &c.

B. *General Descriptions.* After the distinctive characters of a species, race, or variety, have been given, that species, race, or variety, will be described more at length.

- a. *Habit, Bulk, Figure, and Duration.* The entire plant will first be characterised, as whether tree, shrub, undershrub, twiner, climber by tendrils or by elongation, trailer, &c.; its bulk, figure, and duration will then be given, in a natural state in its native habitat, and in an artificial state, more especially in British plantations or gardens.
- b. *Species.* In describing species, two objects will be kept in view; the first to convey a correct idea of the figure of the plant to one who has never seen it, so as to enable him to recognise it; and the second to communicate such ideas respecting its nature, its roots, branches, wood, seeds, &c., as may give a cultivator some notion as to how it may be propagated, and to what purpose its products may be applied.
- c. *Races and Varieties* will be conducted on the same plan, and with a view to the same result as the descriptions of species.
- d. *Mules, or Hybrids.* A hybrid may either be the produce of two aboriginal species, such as the *Pyrus spuria*, in which we shall call it a mule; or it may be the produce of two races or varieties, as are most of our cultivated fruits; in which case, if it reproduces itself from seed it is a race, but if it can only be propagated by division it is a variety.
- e. *Variations.* Variations differ from varieties in not being capable of being continued by propagation. The susceptibility of individuals to vary in their appearance with change of soil, light, or shade, or other circumstances; such as the flowers of the hydrangea becoming blue

in certain soils, &c. ; will be noticed as far as they are known : for, though none of these peculiarities can be continued by propagation, some of them may be produced by culture.

- f. *Impulsion, or Rate of Growth.* The number of feet, or inches, made by shoots of one season's growth, in trees of different ages, will be given ; and the height which the species generally attains in ten, in twenty, in thirty years, and when full grown, in the environs of the metropolis, will be stated as far as has been ascertained.
 - g. *Metamorphoses and Degeneracies.* The doctrine, that all the parts of a plant may be reduced to two (viz. an axis, and a leaf revolving round it), seems to have been hinted at by Linnaeus, but was first brought forward in a conspicuous manner by the poet Göthe. (See De Cand., *Théorie Élémentaire*, p. 105. and *Physiologie Végétale*, p. 771., and Göthe's *Versuch über die Metamorphose der Pflanzen*, 1831.) The doctrine is now generally adopted by botanists ; and double flowers and various other monstrosities and transformations are referred to this head.
 - h. *The Anatomical Structure* of species will be noticed when it is, in any respect, remarkable ; as, when it influences materially the texture or veining of the timber, or the susceptibility of the plant to be united to others by grafting, its fitness for resisting wind, &c.
 - i. *Physiology.* Anything remarkable in the functions of any species will be stated ; together with its bearings on propagation, culture, or duration : such as the kind of sap, whether milky or watery, sugary or alkaline, &c.
 - k. *The Affinities of Species* constitute an important part of their study, with a view to their propagation and culture. Some species may be grafted, not only on every other species of the same genus, as in the case of *Crataegus*, but on every species belonging to the same tribe, such as *Prunus* : other species, which will not unite by grafting to all the species of their own genus, such as *Pyrus communis*, which will not unite to *Pyrus Malus*, will yet unite to *Crataegus* and *Sorbus*. In general, plants which have milky sap will not unite with such as have watery sap, and, indeed, will not unite with other plants at all. Hence, *Acer platanoides*, according to De Candolle (*Physiologie Végétale*, vol. ii. p. 794.), cannot be grafted on any other species of the genus.
 - l. *Resemblances.* Some trees and shrubs bear a resemblance to one another without having any affinity, either organic or physiological ; for example, the different species of *Carya*, *Rhus*, and *Ailantus* ; or the common laurel and the *Magnolia grandiflora*. These resemblances it will be useful to notice, with a view to ornamental plantations.
 - m. *Contemporary Foliation, Flowering, and Defoliation.* For the same purpose as that of indicating resemblances, it will be desirable to note trees and shrubs which come into flower at the same time ; or which either come into leaf, or shed their leaves contemporaneously.
6. *Casualties.* Trees and shrubs are liable to be preyed on by insects, to be injured by vermin and parasitical plants, attacked by diseases, and broken down or destroyed by accidents.
- A. *Insects and Vermin.* The particular species of these which are either peculiar to certain species of trees and shrubs, or liable to attack them, will be described, and occasionally figured ; and the means of protection from their ravages, when known, or of alleviation, or of cure, will be pointed out.
 - B. *Parasitical Plants.* Trees and shrubs are liable to be injured by the growth of lichens, mosses, and other parasites, on their leaves, bark, and wood ; and by *Fungi* on their bark and leaves : among the latter class is the mildew.
 - C. *Diseases.* The diseases to which trees and shrubs are liable, exclusive of the injury done to them by insects and vermin, and by parasitical

plants, are not many; but, still, some, such as the canker, &c., require to be noticed, when the species subject to them come under review.

D. *Accidents.* Some trees and shrubs are more liable than others to be blown down, or have some of their branches broken off by high winds, or by the weight of snow; and these species should be pointed out to the inexperienced planter. Some, also, are said to be less liable to be struck by lightning than others; for example, the beech.

7. *Geographical Distribution.* The different countries will be mentioned where each species is found naturally; and, where practicable, the different localities, soil, elevation and other particulars will be given. It will also be stated, as far as is known, in what countries each species is cultivated, and to what purposes it is applied.

8. *History.* This subject may be included under two heads, viz. retrospective and prospective.

A. *The Retrospective History* of every species or variety will commence with its first discovery, or record by botanists; and its progress will be traced in every country, but more especially in Britain, from that period to the present time. Though the history of some trees and shrubs commences with the time of the Romans, yet that of others is comparatively obscure; and, of some of the finest ornaments of our gardens, little more can be stated than that they are races or varieties, perhaps hybrids, raised by cultivators whose names are unknown.

B. *The Prospective History*, or probable progress, of our knowledge of species may be included under the heads of doubtful species, unnamed species, and expected additions.

a. *Doubtful Species.* In almost every genus, containing several species, there are some of the names which are of doubtful application, which under this head we shall bring together, with their authorities, in order to direct the attention of botanists and cultivators to the subject.

b. *Unnamed Species.* The introduction of new species of trees and shrubs into British gardens is constantly going on, and numbers are also as constantly being raised from seed in the country. In general, neither of these kinds of additions to our ligneous flora can be received into books till they have flowered; when they are named, figured, and recorded in some botanical work. It often happens, however, that the genus to which a new plant belongs is discovered, from the general habit of the plant, long before it has flowered; and in such a case, though the species may not be introduced into botanical catalogues, plants of it may be distributed among cultivators by those who have introduced it, and it may be propagated for sale in the nurseries, under some provisional name. Such species, and also varieties raised in the country from seed, or otherwise procured, deserve notice in a work like the present, and we shall devote this head to the subject.

c. *Expected Additions.* The species of some genera are so numerous, and their geographical distribution is so extensive, that from these circumstances alone we may reasonably anticipate the discovery and the introduction of additional species. Of other genera, many species suitable for our climate, though not yet introduced, have been described by botanists as indigenous in different parts of the temperate hemispheres. It will be useful to direct attention to both these points, with a view of stimulating travellers and others to procure the additional species that are known; and wealthy individuals, or societies or associations, to send out collectors to discover those species which may be yet unknown.

9. *Use.* Trees and shrubs are used in the arts of construction, of machinery, and of fabrication; in the chemical arts of dyeing and colouring; in domestic and rural economy; and in medicine.

A. *The Arts of Construction* are, civil, military, and marine architecture; engineering, carpentry, joinery, cabinet-making, carving, and modelling;

and also cooperage, locksmithry, turnery, mathematical instrument-making, trunk-making, &c.

B. *The Manufacture of Machinery, Instruments, Implements, and Utensils*, comprehends the making of mills, machines, carriages, implements of husbandry, gates, fences, ladders, pumps, water-pipes, gun-stocks, spade-handles, and an almost innumerable number of similar articles.

C. *The Arts of Fabrication* are, weaving, rope-making, mat-making, &c.

D. *The Chemical Arts* include tanning, dyeing, colouring, the expression of oils, the extraction of sugar, the distillation of pyrologeneous acid, of ardent spirits, the fermentation of wine, beer, &c.

E. *The Arts of Domestic Economy* include the preparation of wood for fuel, basket-making, and toy-making; the preparation of walking-sticks, fishing-rods, and other articles used in games, sports, pastimes, recreations, &c., and of chests, desks, and coffins.

F. *The Arts of Rural Economy* comprehend the use of trees and shrubs, in a living state, in agriculture, gardening, and planting; and, also, their use in producing leaves, or stems, to serve as food for domestic animals, fruit for food or drink for man, wood for fencing, draining, &c.

G. *Medicine*. Various parts of trees enter into the materia medica of the medical corporations; while others are used only in empirical practice: both uses will be noticed in a succinct manner.

H. *The Use of Trees by the Priests of particular Religions*, and the ancient uses of some of them as charms, &c., as of the oak and the mistletoe by the Druids, the rowan tree by the believers in witches, &c., will be slightly noticed.

I. *Poetical, Mythological, and Legendary Associations* are connected with various trees and shrubs; and the ideas which these species recall may be considered as a part of their use.

K. *The Picturesque and Decorative Uses of Trees* will, as far as respects planting them, be considered under gardening; but, under this head, will be noticed their suitableness for the landscape-painter; the architect, for architectural ornaments; the house decorator; the decorator of different arts and manufactures, such as those of china, printed tissues, paper-hangings, &c.; and the decorator of theatres, triumphal arches, processions, &c.

10. *Propagation*. In general, all perfect plants may be propagated by all the different modes of propagation known either in nature or art. All perfect plants produce seeds, and may be propagated by them; and they all produce buds, and, for the most part, these buds may be separated from the parent plant, along with a portion of its wood, and inserted in the soil, or in other plants, so as to become plants also. But, as all trees and shrubs are not susceptible of being propagated by all modes with an equal degree of facility, the use of treating of the propagation of individual species is, to point out the methods which are considered most advantageous for each. It is also particularly necessary, to indicate certain modes of propagation as best adapted for certain purposes; as, for example, that of buds, or any other mode of division, for the continuation of varieties, &c.

A. *Natural Propagation* is effected by seeds, by side suckers or root suckers, and by surface stolones or underground stolones.

B. *Artificial Propagation* is effected by seeds, suckers, cuttings, layers, ringing, budding, grafting, and inarching.

a. *By Seeds*. Seeds are employed in artificial, as well as in natural, propagation. The subject embraces their ripening on the tree, their gathering, keeping, preparation for sowing, time of sowing, soil, situation, preparation, and time for transplanting.

b. *By Suckers*. These are of two kinds: side suckers, which rise up close to the stem of the plant, as in the case of the common lilac; and root suckers, which rise up from the roots of the plant, to whatever distance these may extend, as in the case of the common plum, the

elm, and many other trees. The time of separation, the size, the future treatment, and the fitness of plants so produced relatively to those produced by other means of propagation, require to be considered.

- c. *By Division.* Low-growing many-stemmed shrubs, such as the dwarf box, the butcher's broom, &c., and some creepers, such as *Hypéricum calycinum*, &c., are most easily propagated by taking up the entire conglomeration of plants, and separating them.
- d. *By Cuttings.* Cuttings may be taken from the branches, or shoots, and either in summer or winter; they may also, in some cases, be taken from the trunks of trees of large size; they may be taken from roots in many cases; and some evergreens, such as the *Álcúba*, and some deciduous shrubs, such as the *Wistària*, may be propagated by leaves cut off with a bud in their axil. In all these methods, the season, soil, situation, shade, air, temperature, and time for transplanting, require to be taken into consideration.
- e. *By Layers.* These may be made of the winter's wood, or of the summer's shoots, and by a variety of different modes of cutting, twisting, ringing, &c.; in all of which, the season, soil, and situation, and time for detaching and transplanting the layers, require to be treated of.
- f. *By Ringing, and applying a Ball of Earth or Moss.* This is practised in various ways, with or without the aid of a perpetual supply of water; and, as in the preceding cases, the season, locality, and the nature of the subject, with other particulars, require to be mentioned.
- g. *By Budding on other Plants.* Here we have to consider the kind of stock; its age; its influence on the scion; the modes of performing the operation, which are various; the age of the scion from which the buds are taken; the time when the plant is fit for transplanting; and other particulars.
- h. *Budding in the Soil.* Leaves with buds in their axils will, in various cases, both of deciduous and evergreen trees and shrubs, produce plants. Buds, also, without leaves, but with small portions of wood cut from trees, in some cases from the old wood, as in propagating the olive, and in others from the young wood, as in propagating the vine, will produce plants. Buds in the roots may also be so employed; as in the case of many of the *Rosàcèæ*. In general, the buds of the trunks and roots are latent germs, and not visible on the portions that are employed for propagation.
- i. *By Grafting.* With reference to this operation, the kind of stock should be indicated, its age, and its influence on the scion; the mode of performing the operation; the season; the age of the scion; and the time when the subject is fit for transplanting.
- k. *By Inarching.* Here the same considerations require to be taken into view as in grafting; with various additional ones, respecting the mechanical position of the stock, in the case of marching the branches of high trees into stocks in pots.

11. *Culture.* This subject embraces the soil, situation, and exposure; the rearing and culture in the nursery; the choice of plants and planting out; the final culture and management of the plant till it dies, is felled or cut down; and the species adapted to succeed it.

A. *The Soil, Situation, and Exposure.* In general, it may be asserted that the component parts of soils are only of importance relatively to their capacity for retaining, or parting with, moisture; but some plants are absolute in their choice, and will only thrive in particular soils.

B. *Culture in the Nursery.* This, in some cases, will require to be carried on for some time under glass or in pits, or against a wall or with some kind of protection; it may require the plant to be kept in a pot or box, in a shaded or light situation, in a close or airy one, in rows in beds, or singly, &c. The time when the plant will be fit for final transplanting will require to be mentioned; and, also, what is of very

considerable importance, to what size or age the tree or shrub may be kept in the nursery and still be fit to transplant; the number of times which it ought to be transplanted while it remains in the nursery, till it attains that size; its pruning; protection from the weather, from insects, epiphytes, parasites, diseases, &c.

- C. *Choice of Plants, and Planting out.* Some plants are better adapted for transplanting at one age than another; and while some may be taken at once from the seed bed or nursery lines, others should be grown in pots, for more convenient deportation, with all their fibrous roots and spongioles in a living state.
- D. *Culture after final Removal.* This will embrace the treatment of the plant, as a single tree or shrub in a park or lawn; its treatment, as part of a picturesque group, or as part of a gardenesque group; against a wall, as a climber, twiner, trailer, or creeper; collectively, in ornamental plantations, whether gardenesque or picturesque; in useful plantations, whether arranged methodically or planted irregularly; in geometrical plantations; in architectural or sculptural plantations; in avenues, arcades, hedgerows, and hedges.
- E. *Species adapted for Succession.* Natural forests, when they decay by age, are destroyed by fire, or cut down by man, are generally succeeded by a different species of tree from that which before prevailed. It is desirable to imitate this natural process by art, as far as experience and science can direct; and some space will therefore be devoted to the consideration of the subject, in its proper place in our *Encyclopædia of Arboriculture*.
12. *Statistics.* By statistics is to be understood the actual state of any science or art; and the statistics of trees and shrubs may be included under the heads of geographical statistics, and commercial statistics.
- A. *Geographical Statistics.* Under this head we shall include the notices of the age and the dimensions of the trees and shrubs of temperate climates, which we have obtained in consequence of the circulation of the printed forms which we have called Return Papers (see *Gard. Mag.*, vol. x. p. 582.), in Britain, on the Continent, and, as far as we have been able, in North America. The information thus obtained will be useful, as showing the undoubted hardiness of some trees and shrubs; the comparative suitableness of certain soils and climates for particular kinds; those which in general may be considered as most hardy, or of most rapid growth; which attain the largest size, or the greatest age; which are most profitable, or most ornamental, &c.; but, above all, it will show the comparative advances which trees make in a soil prepared, or not prepared, in different parts of Britain. The statistics of trees will be arranged as Domestic and Foreign.
- a. *The Domestic Notices* of the existence of trees and shrubs in certain places, together with notices of their age, rate of growth, &c., will be placed under the heads of—
- a. *In the Environs of London*; that is, within a radius of ten miles from the metropolis.
 - b. *South of London*; that is, in the English counties which are situated wholly, or in the greater part, south of the metropolis
 - c. *North of London*; that is, in the English counties which are situated wholly, or in the greater part, north of the metropolis.
 - d. *Wales*; taking the counties alphabetically.
 - e. *Scotland*; in the same order as in England.
 - f. *Ireland*; also in the same order.
- b. *The Foreign Notices* of the existence and dimensions of trees and shrubs, which we have received, or have collected from books, will be given in the following order:—
- a. *Europe.* 1. France. 2. Belgium and Holland. 3. Germany.

4. Denmark. 5. Sweden and Norway. 6. Russia and Poland.
7. Switzerland. 8. Italy, Greece, Spain, and Portugal.

b. *America*. 1. North America. 2. Mexico. 3. South America.

c. *Asia*. 1. Asia Minor. 2. India. 3. China. 4. Japan.

d. *Australia and Polynesia*. 1. Van Diemen's Land. 2. New South
Wales. 3. New Zealand.

B. *Commercial Statistics*. Trees and shrubs are objects of commerce: in their young state, as plants; and in their more matured state, as timber, fencewood, fuel, bark, leaves, fruit, seeds, &c.

a. *Nursery Commerce, domestic and foreign*. Some trees and shrubs, from being in little demand, are scarcely known out of private gardens, or public botanical establishments; others are cultivated in the nurseries, some very generally, and others only partially. Under the head of Commercial Statistics, we shall notice whether the species is cultivated only in some nurseries, or generally; and we shall give the prices of plants of the smallest size fit for transplanting, and also of seeds when they are to be procured: in London; in the extensive nurseries of Messrs. Baumann at Bollywyller on the Rhine, as a situation central for France, Germany, Switzerland, and Italy; and in New York, as a central situation for North America.

b. *General Commerce, domestic and foreign*. Under this head it will be our object to notice such trees, or their products, as are in general transfer in the internal commerce of the country; and such, also, as are exported or imported. Some woods, as the pine, fir, oak, elm, &c., are in general commerce; and so, also, are some other products, such as oak bark; but the timber of the spindle tree and the laburnum, the inner bark of the holly, and the flowers of the elder bush, enter into the commerce only of particular places. What we shall state respecting either the foreign or domestic commerce of trees and shrubs, will be limited to what relates to the trees and shrubs of temperate climates; that is, to those species which are described in this work.

Such is the *beau idéal* of the desiderata which we intend to keep in view, when describing each species; but we by no means bind ourselves to have, in our descriptions, a separate heading for each of the paragraphs in this Chapter; on the contrary, it will generally be found, that all that we have to say respecting each species will be included in the paragraphs entitled, *Identification, Synonymes, Derivation, Engravings, Specific Character, Varieties, Description, Geography, History, Properties and Uses, Soil and Situation, Propagation and Culture, Accidents and Diseases, and Statistics*.

All the matter included under the first four headings, as being of less interest to the general reader, we have placed in small type, in order that it may occupy but little space, and be easily passed over by those who do not wish to read it. We have also placed in small type the whole of the matter relating to species which have not been seen by us; and also to those which are only half-hardy, and require either to be planted against a conservative wall, or otherwise to receive some kind of protection during the most severe weather in winter. We have done this, though we consider what relates to the species which require some protection, as likely to prove one of the most interesting parts of our work to many gardeners and amateurs (for what would the enjoyments of gardening be, without the elegant cares of exotic culture?), in order that those who take an interest only in hardy trees and shrubs may distinguish, at a glance, what belongs to them.

PART III.

THE ARBORETUM AND FRUTICETUM BRITANNICUM; OR THE DESCRIPTION, HISTORY, PROPERTIES, AND USES, OF THE HARDY TREES AND SHRUBS OF BRITAIN, INDIGENOUS AND FOREIGN.

TREES and shrubs, in common with all other perfect plants, are arranged by botanists in two grand divisions; viz. the Exogenous, or Dicotyledonous, plants, the stems of which increase from without; and the Endogenous, or Monocotyledonous, plants, the stems of which increase from within. The first class includes all the hardy trees and shrubs in Britain, with the exception of shrubs of the genera *Yucca*, *Smilax*, *Ruscus*, and one or two others; and this circumstance, as well as the fact, that the trees and shrubs of Britain are comprised in a very few orders and tribes, has determined us to neglect the great scientific divisions of the natural system, and to adopt only those of the orders and tribes. We proceed, therefore, with the orders of the natural system, much in the same series as that in which they are laid down in De Candolle's *Prodromus*, Don's *Miller's Dictionary*, and in our *Hortus Britannicus*, giving the orders as chapters, and the tribes as sections, and including in our distinctive character of each order, the characteristic of the division to which it belongs: that is to say, whether to Dichlamýdeæ Thalamiðræ, Dichlamýdeæ Calyciðræ, Dichlamýdeæ Corollifiðræ, or Monochlamýdeæ.

CHAP. I.

OF THE HARDY LIGNEOUS PLANTS OF THE ORDER RANUNCULA'CEÆ.

THE term *Ranunculæcæ* is applied to this order, because all the plants contained in it have, more or less, the character of the genus *Ranunculus*. The diagnostic, or distinctive character, of the order is thus given by Dr. Lindley:—"Polypetalous dicotyledons, with hypogynous stamens [that is, stamens under the pistil]; anthers bursting by longitudinal slits; several distinct simple carpella [fruits]; exstipulate leaves, sheathing at their base; solid albumen; and seeds without arillus." (*Introd. to the Nat. Syst.*, p. 6.)

The only ligneous plants belonging to this order are, some species of *Clématis* and *Atragène*, one of *Pæonia*, and the genus *Xanthorhiza*. The stems of the species alluded to, though they are botanically considered as ligneous, yet have very little claim to the appellation in the common sense of the word; and, indeed, with the exception of the stems of *Clématis Vitalba*, *C. Flammula*, and one or two other species of *Clématis* and *Xanthorhiza*, the stems of the plants belonging to this order might be almost called subherbaceous. The species are chiefly natives of Europe and North America; but some are from India, China, and Japan. The *Ranunculæcæ* are considered to indicate a cold damp climate, and to be aerid, caustic, and poisonous, though the root of the peony is said to be antispasmodic. All the plants of the order, with the exception perhaps of a few of the species, seem to be extremely tenacious of life. The tubers of the common *ranunculus* and *anemone*, if kept dry, will vegetate at the end of two, and even three, years; and the seeds of most of the species, more especially those of the *Clematideæ*, may be kept a number of years without impairing their vital powers. The tribes containing ligneous plants are two, *Clematideæ* and *Pæoniæcæ*. The last tribe belongs to a division of the order consisting of what are considered as spurious *Ranunculæcæ*. It includes the ligneous genera *Xanthorhiza* and *Pæonia*, which even a superficial observer may

recognise as differing, in habit and appearance, from the genera *Clématis* and *Atragène*, which are slender-stemmed climbers, while the others are herbaceous-looking undershrubs.

Sect. I. CLEMATIDÆ.

THESE are climbers, characterised by having the aestivation of the calyx valvate or induplicate; with no petals, or with the petals flat; the anther opening outwards; the carpels, or seed-vessels, not opening; one-seeded, terminated by a tail, which is the indurated style. Seed pendulous. Leaves opposite. Deciduous and evergreen climbers. The genera are two; *Clématis* and *Atragène*, which are thus contradistinguished:—

CLEMATIS L. Petals none.

ATRAGÈNE L. Petals several.

GENUS I.



CLEMATIS L. THE CLEMATIS, or VIRGIN'S BOWER. *Lin. Syst. Polyandria Polygynia.*

Identification. The word *Clématis* is said by Donnegan to have been used by Theophrastus, cap. 5. 10., as well as *Atragène*, to designate the *Clématis Vitalba* of Linnaeus. *Clematis* was used by Matthioli, and also by Clusius, who applied it to *C. Viticélla* L. and *C. cirrhosa* L. It has been since generally applied to this family of plants by botanists.

Synonyms. Ladies' Bower *Gerard*; *Clematite*, *Fr.*; *Waldrebe*, *Ger.*; *Clematide*, *Ital.*

Derivations. The word *Clematis*, or *Klematis*, is derived from the Greek word *kléma*, a small branch of a vine; and it is applied to this genus, because most of the plants composing it climb like a vine. The English name of Ladies' Bower was probably adopted from its suitability for covering bowers; and, as the first kind of clematis brought to England (*C. Viticélla*) was introduced in 1569, during the reign of Elizabeth, the name of Virgin's Bower might be intended to convey a compliment to that sovereign, who, as it is well known, liked to be called the Virgin Queen. The German name, *Waldrebe*, is compounded of *wald*, a wood, and *rebe*, the branch of a vine.

Generic Character. *Involucre* none, or situate under the flower, in the form of a calyx. *Calyx* of from four to eight coloured sepals. *Petals* none. *Carpels* numerous, aggregate, terminated by a long, and mostly feathery, tail.—Climbing shrubs, with variously cut opposite leaves. The recent herb of all the species is acrid, and, when applied to the skin, it occasions blisters. (*Don's Mill.*, i. p. 3.) The seed is pendulous, and the carpels are one-seeded; each is terminated by a persistent style, and does not open until ruptured by the germination of the seed.

Description, &c. Root strong; the fibres rather straight, and not very much branched; extended in the soil rather horizontally than perpendicularly. Stem ligneous, not rigid enough to stand erect. Branches the same, and slender. Leaves in decussating pairs; the petiole possessed of a clasping power, the effect of which is the prehension of contiguous plants and objects. The rate of growth in *C. Vitalba* and *C. Flammula* is among the most rapid known in the plants of temperate climates, particularly in the shoots which a well-established vigorous plant throws up, after it has been cut down to the ground. The most ornamental species are *C. Viticélla* and *C. florida*; the most rapidly growing for covering bowers is *C. Vitalba*. The kind most fragrant in its flowers is *C. Flammula*.

Geography, History, Uses, &c. Most of the hardy species of *Clématis* are natives of the middle and south of Europe, and of North America; a few of them are natives of the north of Africa, some of Siberia; there are several in the Himalaya, one in China, and several in Japan. The genus has been known since the days of Theophrastus, and has received various accessions

from the time of Matthioli to the recent introductions from the Himalaya. The acrid properties of the *Clematis* are well known to herbalists. The bark, leaves, and blossoms are used to raise blisters on the skin, or to produce a slight external inflammation: taken internally they are a corrosive poison. The flowers contain a peculiar substance, called clematine, which is similar to gluten; the green leaves, bruised, are applied to ulcers, to produce sloughing. The floricultural use of these plants is to cover bowers, or ornament verandas or trellis-work. The greater number of them ripen their seeds in England, and are easily propagated by them or by layers. They all require support by props of some kind; and all grow freely in any soil that is tolerably dry, but more especially in one that is calcareous. From the acridity of these plants, they are not very liable to be attacked by insects; nevertheless, snails and slugs are occasionally found eating their young herbage. Most of the species and varieties which we shall describe are to be found in the principal botanic gardens of Europe, and have been seen by us in that of the Horticultural Society of London; and the more ornamental of them are cultivated for sale in the principal European and American nurseries.

The ligneous species of *Clematis* are included in four sections; viz. *Flammula*, *Viticella*, *Chiropsis*, and *Anemoniflora*.

§ i. *Flammula* Dec.



Sectional Character. Involucre wanting. Tail of the carpels long, bearded and feathery. Cotyledons distant in the seed. (*Don's Mill.*, i. p. 3.)

1. *CLEMATIS FLAMMULA* L. The inflammatory-juiced Clematis, or sweet-scented Virgin's Bower.

Identification. Lin. Sp., 766.; Willd. Sp., 2. p. 1293.; Hayne Den., p. 119.; Lam. Dict. Encyc., 2. p. 42.; Dec. Prod., 1. p. 2.; Don's Mill., 1. p. 4.

Synonyms. *C. urens* Gerard; *C. maritima* All. Ped.; *C. suaveolens* Satisb. Prod.; *Clematide odorante*, Fr.; *Scharfe walddrebe*, Ger.

Derivation. From *flamma*, to inflame; on account of the blistering qualities of the species.

Engravings. Park. Theat., p. 381. f. 3.; Knorr. His., 2. p. 9.; and our fig. 9.

Specific Character. Leaves pinnate, smooth; with orbicular, oval, oblong or linear, entire or three-lobed, acatish leaflets. (*Don's Mill.*, i. p. 4.) Flowers white. July to Oct. Height 15 ft. 1596.

Varieties and their Synonyms. The following are given by De Candolle; but they are not of much importance in point of general effect:

1. *C. F.* 2 *rotundifolia* Dec., *fragrans* Tenore.—Leaflets almost orbicular.

2. *C. F.* 3 *maritima* Dec.—Leaflets linear.

3. *C. F.* 4 *rubella* Dec.—Leaflets oval, usually emarginate. Sepals four, reddish on the outside.

4. *C. F.* 5 *caespitosa* Dec., *C. caespitosa* Scop., *C. Flammula* Bertol.—Leaflets minute, entire, or cut.

Description. A vigorous-growing plant, the stems of which attain the length of 10 ft. or 15 ft. in a wild state, and from 15 ft. to 30 ft. in a state of culture. The leaves of the entire plant are subject to much variation, from soil, situation, and climate. The shoots of a well-established plant, which has been cut down, grow with great rapidity in the early part of the season, attaching themselves to whatever is near them. The peduncles of the flowers are sometimes simple and sometimes branched. The colour of the sepals is white, slightly pubescent on their exterior margins. The whole plant has a dark green hue; and in autumn it is abundantly covered with flowers, the odour of which is of a honeyed sweetness, exceedingly disagreeable to some persons when near, though at a distance it is not unlike the fragrance of the common hawthorn. The number of the styles varies from five to eight, each

style terminating in a little white feathery process when the seed ripens: the plant at that time appears covered with little tufts of cotton. In its native habitats this plant flowers in July and August; but in Britain it continues in flower from July to October. From the rapidity of its growth, it will in four or five years cover a very large space of wall, roof, or bower. Its herbage is considered less acrid than that of any other of the European species, notwithstanding its name of *Flammula*. (*Dec. Syst.*)

Geography. This well-known species seems confined to the middle and south of Europe and to the north of Africa. It is found in the south of France in hedges, and in waste bushy places; in Greece, Italy, Spain, and Portugal (see p. 132, and p. 164.), and in all these countries, generally in low situations, not far from the sea, and in soil more or less calcareous.

History and Use. *C. Flammula* appears to have been first recorded by Dodonæus, in his *Stirpium Historie Pemptudes*, in 1585; it was recognised by Matthioli and L'Obel, and cultivated by Gerard in 1597; and it is now generally grown in gardens throughout Europe and North America for covering bowers, garden-houses, trellis-work, and naked walls; for which purposes it is well adapted from its rapid growth, its intense fragrance when in flower, and its tufted cottony masses when in seed.

Statistics. Plants may be had in all the European nurseries: about London, of the smallest size, at about 5s. per hundred, or 6d. for a single strong plant; at Bollwyller, at from 6 francs to 8 francs the hundred, or about half a franc a plant; and at New York, for 30 cents per plant.

2. *C. ORIENTALIS* L. The Oriental Clematis.

Identification. Lin. Sp., 765.; Willd. Sp., 2. 1289.; Lam. Dict. Enc., 2. p. 42.; Hayne Dend., 119.; Dec. Prod., 1. p. 3.; Don's Mill., 1. p. 4.

Synonyms. *Flammula scandens apii folio glauco*, Dill. Elth., 144.; *C. flava Moench. Meth.*, 296.; the eastern, or yellow-flowered, Virgin's Bower; *Clematis orientalis*, Fr.; *Morgenländische Waldrebe*, Ger.

Engravings. Dill. Elth., t. 119. f. 145.; and our fig. 10.

Spec. Char. Leaves pinnate; leaflets smooth wedge-shaped, with three toothed pointed lobes. (*Don's Mill.*, i. p. 4.) Flowers greenish yellow, slightly tinged with russet, sweet-scented. Aug. Sept. 1731. Height 15 ft.

Description. The general magnitude of this species resembles that of *C. Flammula*, from which it differs, in its ulterior branches being more persistently ligneous, though the main stem in old plants is seldom seen so thick as that of *C. Flammula*. It is also distinguished from the latter species by throwing up suckers freely, which the other does not. Its leaflets are glaucous, flat, large as compared with those of *C. Flammula*, and it does not produce flowers so profusely as that species; the flowers are yellowish, and not so strongly scented; and the carpels are dissimilar, though still cottony in appearance when the seed is ripe.



Geography, History, &c. *C. orientalis* was discovered by Tournefort in the Levant, and sent by him to the Paris Garden; whence it was sent to Clifford's garden in Holland, and, in 1732, to that of Dr. Sherard at Eltham; when it was first described and figured in the *Hortus Elthameusis*. The plant has been subsequently discovered in Caucasus by Bieberstein, and described by him in his *Flora Taurico-Caucasica*. It is not very generally cultivated, though it is found in several botanic gardens, and may be purchased in some nurseries. Plants of it are in the garden of the London Horticultural Society. Price, in London, 1s. or 1s. 6*d.* a plant; at Bollwyller, ?; and in New York, ?.

‡ 3. *C.* [? o.] GLAU'CA W. The glaucous-leaved Clematis.

Identification. Willd. Arb., 65., and Sp., 2. p. 1290.; Dec. Prod., 1. p. 3.; Don's Mill., 1. p. 4.
Engravings. Willd. Arb., 65. t. 4. f. 1.; Den. Brit., 73.; and our fig. 11.

Spec. Char. Leaves pinnate; leaflets smooth, glaucous, wedge-shaped, with entire bluntish lobes. Peduncles trifid. (*Don's Mill.*, i. p. 4.) Flowers yellow, scentless. July. 1800. Height 10 ft.

Description. The general appearance is the same as that of *C. orientalis*, of which it is probably only a variety; but the whole plant is more decidedly glaucous.

Geography, History, &c. Found in the southern parts of Siberia, in sandy wastes, by Pallas; but when it was introduced into Britain is uncertain. Cultivated in Knight's Exotic Nursery, King's Road, Chelsea, where it flowered in 1822, and whence it was figured by Watson in his *Deudrologia*. There is a plant of it in the London Horticultural Society's Garden, which bears so strong a resemblance to *C. orientalis*, that, supposing them to be correctly named, we have no doubt of their being the same species.



‡ 4. *C.* CHINE'NSIS Retz. The Chinese Clematis.

Identification. Retz. Obs., 2. p. 18.; Dec. Syst., 1. p. 137.; Don's Mill., 1. p. 4.
Synonymes. *C. sinensis* Lour. coch., 1. p. 322.

Spec. Char. Leaves pinnate; leaflets ovate-lanceolate, quite entire. Peduncles few-flowered, longer than the leaves. Ovaries usually four, with almost naked tails. (*Don's Mill.*, i. p. 4.) Flowers [?] purple. 1820. Height 15 ft.

Description. This plant is described in De Candolle's *Systema*, from a dried specimen which he had seen in the Banksian Herbarium. There is a living plant in the Horticultural Society's Garden, which grows vigorously against a wall, producing shoots as long and strong as those of *C. Flammula*; and retaining its leaves till they are blackened by frost. This plant has never flowered in England; but, in its leaves and its general appearance, it seems to resemble *C. orientalis*.

Geography, History, &c. This plant was found in China, in the island called Danes, whence it was received by the Horticultural Society in 1820. It is planted in the garden at Chiswick against a wall, with a southern exposure, and receives some slight protection during winter.

‡ 5. *C.* PANICULA'TA Thuu. The paniced Clematis.

Identification. Thunb. Lin. Soc. Trans., 2. p. 337.; Willd. Sp., 2. p. 1291.; Dec. Prod., 1. p. 3.; Don's Mill., 1. p. 4.

Synonymes. *C. Vitálba japénica* Houtt. Pflanz., 7. p. 309.; *C. crispa* Thunb. Fl. Jap., p. 239.

Engravings. Houtt. Pflanz., 7. p. 309. f. 2.

Spec. Char. Leaves pinnate; leaflets ovate, cordate, acute, entire. Peduncles paniced, many-flowered. (*Don's Mill.*, i. p. 4.) Flowers white, and sweet-scented. 1796.

Description. Described by De Candolle in his *Systema*, from dried specimens, from which it appears that the flowers resemble those of *C. Flammula* in form and colour, and, like them, also, are sweet-scented.

‡ 6. *C.* VITA'LBA L. The White Vine Clematis, or Traveller's Joy.

Identification. Lin. Sp., 776.; Willd. Sp., 2. p. 1292.; Fl. Br. 583.; Hook. Scot., 171.; Lam. Dict. Enc., 2. p. 41.; Hayne Den., p. 120.; Dec. Prod., 1. p. 3.; Smith's Eng. Fl., 3. p. 39.; Don's Mill., 1. p. 4.

Synonymes. Athragenē *Thoph*; *Vitis sylvéstris* Dios.; *C. latifolia* seu *Atragène* Ray; *C. áltera* Math.; *C. tértia* Com.; *Vioria* Ger. and Lob.; *Vitis ngra* Fuch.; *Vitálba* Dod.; the Old Man's

Beard, Bindwith, the common Virgin's Bower, the wild Climber, the great wild Climber; *Clématite brûlante*, *Clématite des Haies*, *l'Herbe aux Gueux*, *la Viorne des Pauvres*, *Fr.*; *Gemeine Waldrebe*, *Ger.*

Derivation. This plant was called *Fitis sylvêstris* (the wood vine) by Dioscorides; and the name of *Vitálba* was given to it by Dodoneus, probably on account of the white appearance of the plant when covered with its seeds in autumn, which whiteness arises from the hairy tails of the carpels. It was called *C. latifolia* by Ray and Baubin, from its broad-leaved variety; *Viórna* by L'Obel and others, from *vina*, a way, and *ornare*, to ornament, in allusion to its ornamental appearance by the way side; and *Fitis ulgra* by Fuchsius, from the dark colour of the bark of its young shoots. Gerard gave it the name of the Traveller's Joy; because of its "decking and adorning the ways and hedges where people travel; and thereupon," he says, "I have named it the traveller's joy." (*Herbal*, by Johnson, p. 886.) The name of Old Man's Beard is very appropriate to the white and hairy appearance of the tails of the carpels; and Bindwith, from the shoots being used instead of those of willows for tying up plants. The French name of *Clématite brûlante* has reference to the acrid properties of the plant; and *Clématite des Haies* to its growing generally in hedges. The name of *l'Herbe aux Gueux* refers to the employment of it by the beggars in France, who use it to make ulcers in their arms and legs, for the purpose of exciting compassion, curing themselves afterwards by the application of the leaves of the beet. *La Viorne des Pauvres* alludes to the same practice, *viorne* being evidently derived from *Viórna*. (*Dict. Gén. des Eaux et Forêts*, t. p. 67.)

Engravings. Jacq. Austr., t. 1. 308.; Eng. Bot., 612.; Willd. Abr., t. 113.; and our fig. 12.

Spec. Char. Leaves pinnate, leaflets ovate-lanceolate, acuminate, cordate at the base, partly cut. Peduncles forked, shorter than the leaves. (*Don's Mill.*, i. p. 4.) Flowers white. From July to September. Britain. Height 20 ft.

Varieties.


1. *C. 2 V. integrâta.* The entire-leaved White Vine Clematis.

2. *C. virginiana* L., to be hereafter described, is considered by some to be only a variety of *C. Vitálba*. It was cultivated under the name of *C. canadensis* by Miller, who says that "it is very like the common sort, but with broader leaves, and rather more tender, the seeds not ripening in England unless the season be very warm."

Description. The stems are woody, more so than those of any other species, angular, climbing to the height of 20 ft. or 30 ft. or upwards, and hanging down from rocky cliffs, ruins, or the branches of trees; or being supported by, and forming tufts on, the upper surface of other shrubs, or low trees, which they often so completely cover as to have the appearance of bushes at a distance.

The footstalks of the leaves are twined about whatever object they approach, and afterwards become hard and persistent, like the tendrils of a vine. The leaves are either quite entire, or unequally cut; sometimes very coarsely so. The panicles are axillary and terminal, many-flowered and downy. The flowers are of a greenish-white colour with little show; but they have a sweet almond-like scent. "The seeds," Smith observes, "have long, wavy, feathery, and silky tails, forming beautiful tufts, most conspicuous in wet weather. [The water on the twigs and branches, which form the background to the carpels, rendering them darker than they are in dry weather.] They retain their vegetative principle for many years, if kept dry." (*Eng. Flora*, iii. p. 39.)

Geography. *C. Vitálba* is found throughout the middle and south of Europe, in the Grecian Archipelago, and in the north of Africa, about Tripoli. One or more varieties of it have been found in North America, and apparently another in Nepal. (See *Dec. Syst.*, i. p. 140.) The topography of this plant in Britain, according to H. C. Watson, extends to Devonshire in the south, and 53° north latitude. According to some, it is found in Scotland;



but Gerard, and also Winch, assert that it is not indigenous in the north of England, which we believe to be the fact. It is common in France and Germany, and is found in the south of Sweden, but not in Denmark.

History. This species appears to have been first recorded by Theophrastus, as *Athragenē* and also as *Klēmatis*; and it has been since frequently mentioned by botanists, under various names, given in our list of synonyms, from the time of Dioscorides to the days of Linnæus.

Properties and Uses. Du Hamel states, that the French gardeners not only use the twigs of this plant instead of withs, for tying up their plants, but that after stripping them of their bark, they make very neat baskets of them (*Traité des Arbres*, §c., 1st edit. 1755, p. 175.); and they also make of them beehives and a variety of other articles of the same kind. The twigs are in the best state for making these articles in winter; and their flexibility is increased by holding them to the fire before using them. Desfontaines says that the young shoots are not corrosive while they are tender and herbaceous, and that in the south of France they feed cattle with them in that state, and eat them pickled in vinegar. It is also said, in the *Dictionnaire Général des Eaux et Forêts* (vol. i. p. 649.), that a very good paper has been made from the feathery part of the seed. Professor Burnet observes that *C. Vitalba* is used in medicine as a rubefacient in case of rheumatism; and that the dried leaves of the plant form a good fodder for cattle, though they [we presume, the matured ones] would poison the animals if they were eaten in a fresh state; hence affording a good example of the rule which predicates the volatile nature of their acridity. (*Outlines of Botany*, vol. ii. p. 338.) The shepherds, in some parts of England and Germany, often cut pieces of the old wood of this plant, which they light at one end, and smoke like a pipe of tobacco. In gardens and plantations the plant is valuable for the rapidity with which it may be made to cover naked walls, unsightly roofs of sheds, or low buildings and arbours; and also to shelter exposed situations, and for a variety of similar purposes. It is interesting both when in flower and when in seed; and the seeds remain on the greatest part of the winter.

Soil and Situation. It is generally found on chalky or calcareous soils, and seldom, if ever, under the dense shade of trees. On the contrary, when it grows up among bushes, it generally spreads over their upper surface, so that its leaves and flowers are fully exposed to the influence of the light and air.

Propagation and Culture. It is readily propagated by seeds, which often remain eighteen months in the soil before they germinate. It will root by layers; but the plant being common in a wild state throughout Europe, it is seldom cultivated in nursery gardens.

♣ 7. *C. VIRGINIANA* L. The Virginian Clematis.

Identification. Lin. Amœn., p. 275.; Willd. Sp., 2. p. 1290.; Lam. Dict., 2. p. 43.; Mich. Fl. B. Amer., 1. p. 318.; Dec. Prod., 1. p. 4.; Don's Mill., 1. p. 5.

Synonymes. *C. canadensis trifolia repens* Tourn.; *C. canadensis* Mill. Dict., No. 5., *Salsb. Prod.*, 371.; *C. cordifolia Moench. Supp.*, 104.; the broad-leaved Canada Virgin's Bower; *Clematis de Virginie, Fr.*; *Virginische Waldrebe, Ger.*


Engravings. Pluk. Mant., 1. 389. f. 4.; Alb. Acad. Ann., 1. p. 79. t. 7.; Den. Brit. (the male plant), t. 74.; En. Pl., f. 7978.; and our fig. 13.

Spec. Char. Flowers panicled, diœcious. Leaves ternate; leaflets cordate, acute, grossly toothed, or lobed. (*Don's Mill.*, i. p. 5.) Flowers greenish white, fragrant. June to August, 1767. Height 15 ft.

Variety.

♣ *C. v. 2 bracteata* Dec. *The bracted Virginian Clematis.* — "Leaflets ovate-lanceolate, entire." *C. bracteata Moench. Supp.*, 103.

Description. The general appearance of this plant is like that of *C. Vitalba*; but it is less robust in all its parts, and less ligneous in its stems and branches; and it is also somewhat more tender. Miller states that, on this last account,



it seldom ripens seeds in England (see p. 236.); but, as it is dioecious, it is possible that he possessed only the male plant.

Geography, History, &c. Found in North America, from Canada to Florida, in hedges, on the margins of woods, and on the grassy banks of rivers. It was cultivated by Miller in 1767; and though it is not so ornamental as most of the other species, it still finds a place in botanic gardens, and may be purchased in some nurseries. Price, in London, 1s. 6d. a plant, and 6d. a packet of seeds; at Bollwyller, 3 francs a plant; and in New York, 25 cents a plant, or 2 dollars a quart of seeds.

♣ 8. *C. TRITERNATA* Dec. The triternate-leaved Clematis.

Identification. Dec. Prod., 1. p. 6.; Don's Mill., 1. p. 6.

Synonymes. *Atragene triternata* Desf. Hort. Par.

Spec. Char. Leaves binate or triternate, smoothish; leaflets oval, cuneated, three-nerved, acutely trifid. (Don's Mill., i. p. 6.) Flowers white? 1806. Height 5 ft.

Description, History, &c. It does not appear that this species has yet flowered in Europe; and hence it cannot be positively determined whether it is a Clematis, or an Atragene. De Candolle appears to have seen a living plant in the Paris Garden; and he notices that it had not there flowered; which is the case, also, with the plant in the garden of the London Horticultural Society. It is a low feeble-growing plant, and might almost be considered as herbaceous.

♣ 9. *C. VIORNA* L. The road-ornamenting Clematis, or leathery-flowered Virgin's Bower.

Identification. Lin. Sp., 765.; Mill. Diet., No. 10.; W. Sp., 2 p. 1288.; Lam. Diet., 2. p. 44., and Michx. Fl. Bor. Amer., 1. p. 318.; Pursh Fl. Bor. Amer., 2. p. 385.; Jacq. f. Ecl., 1. p. 50.; N. Dub. Dec. Prod., 1. p. 7.; Don's Mill., 1. p. 8.

Synonymes. *C. purpurea repens* Ray; *Flammula scandens*, flore violaceo clauso, Dill. Elth.; American Traveller's Joy; the Virginian Climber; the purple Climber; Clematite Viorne, Fr.; Glockenblüthige Waldrehe, Ger.

Derivation. The derivation of *Viorna* has been already given under *C. Vitálba*. Leathery-flowered virgin's bower refers to the remarkably thick texture of the sepals; the German name is a translation of *Viorna*.

Engravings. Dill. Elth., 118. f. 141.; Jacq. fil. Ecl., 1. t. 32.; and our fig. 14.

Spec. Char. Peduncles 1-flowered. Sepals connivent, thick, acumined, reflexed at the apex. Leaves smooth, pinnate; leaflets entire, 3-lobed, alternate, ovate, acute, floral ones entire. (Don's Mill., i. p. 8.) Flowers purple without, and yellow within. June to August. 1730. Height 12 ft. *Variety.* *C. Simsii* is, in all probability, only a variety of this species, as may be possibly, also, *C. reticulata*.

Description, &c. This species is striking in the dissimilarity of its flowers to those of most other species. They may be compared to large pendulous acorns; but the terminal parts of the sepals are curled upward from the terminal part of the acorns, and towards its sides. The species is (in suitable soil) of vigorous growth, and, exclusive of its flowers, assimilates to *C. Viticella*; but its stems and branches are less decidedly ligneous. De Candolle has cited from Barton, that the herb of this species (by which, perhaps, is to be understood the growing parts of it) is intensely acrid. The stems are numerous, slender, and round; the peduncles of the flower are long, deflexed towards the tip, rendering the flowers pendulous, the sepals never open, except at their extreme ends, which are bent back, giving the whole flower a bell shape, but with the mouth of the bell narrower than the body. The sepals are of a greenish purple, or reddish lilac, on the outside, and of a very pale green within. The stamens scarcely emerge from the sepals. The carpels are broad and flat; as they ripen, the tail becomes bent in and plumose, and of a brownish-green colour.



Geography, History, &c. Found in North America, on woody hills in Carolina and Virginia. It was sent to England by Banister, from the latter country, in 1680, and was cultivated by Sherard in 1732; afterwards by Miller; and it is now to be found in the principal botanic gardens, and in many nurseries. As it does not grow to a great height (seldom exceeding 10 ft.), it is most ornamental as a single plant, trained to a rod or to a wire frame. As its branches are not very decidedly ligneous or persistent, but consist mostly of annual shoots from a suffruticose base, and are not much branched, the plant does not exhibit a bushy head. As ligneous branches do not abound to facilitate the propagation of it by layers, seeds are the readier means, and these are sometimes plentifully produced, and grow without difficulty. The sowing of them as soon as ripe is advantageous to their vegetating in the ensuing spring. A plant of this species, with shoots reaching to the height of 10 ft., and studded with its pendulous peculiarly formed flowers (peculiar for a clematis), more or less projected on their rather rigid peduncles, is an interesting object. Plants, in the London nurseries, cost 1s. 6d.; at Bollwyller, 2 francs; and at New York, 50 cents.

10. *C. CYLINDRICA* Sims. The cylindrical-flowered Clematis.

Identification. Sims, in Bot. Mag., t. 1160; Ait., in Hort. Kew., 2d edit., 3. p. 343.; Pursh, in Fl. Bor. Amer., 2. p. 385.; Dec. Prod., l. p. 7.; Don's Mill., l. p. 8.

Synonymes. *C. crispa* Lam., Michx.; *C. Viórna* Andr., in Bot. Rep.; *C. divaricatà* Jacq.; the long-flowered Virgin's Bower; Clematite à longues Fleurs, Fr.

Engravings. Bot. Mag., t. 1160.; Bot. Rep., t. 71.; Jacq. f. Ecl., l. p. 51. t. 33.; and our fig. 15.

Spec. Char. Peduncles 1-flowered. Sepals thin, acuminate, reflexed at the apex, with wavy margins. Leaves slender, pinnate; leaflets stalked, ovate or oblong, middle one sometimes trifid, floral ones entire. (Don's Miller, i. p. 8.) Flowers large, pale purplish blue. July, Aug. 1802. Height 4 ft.

Description. De Candolle has described this in his *Systema* from a dried specimen, and without any acquaintance with it in a living state. He has deemed it related to *C. Viórna*, reticulàta, and crispa, and discriminated it from these. *C. cylíndrica*, he says, differs from *C. Viórna*, in all the segments of its leaves being entire, not usually trifid; in the flowers being blue, and twice the size of those of *C. Viórna* (in this they are of a reddish lilac, pale within); in the sepals being not leathery, but somewhat of the consistence of paper, with the margin waved; the ovaries 12-15, not 25-30. *C. cylíndrica* differs from *C. reticulàta* in its leaves being in consistence papery, not leathery; scarcely veined, not reticulately veined, and in other points. *C. cylíndrica* closely resembles *C. crispa* in habit and mode of flowering, but differs from it in its sepals being waved in the margin, not rolled backwards; in its larger flowers, and especially in its carpels having long bearded tails, and not naked ones. *C. Viórna* and *cylíndrica*, seen together in a living state, are very dissimilar in appearance. *C. Viórna* has vigorous long branches and reddish flowers, which are acorn-like in figure, except that they have a spreading mouth; there is also obvious dissimilarity in the foliage and shoots, *C. cylíndrica* being almost herbaceous.



Geography, History, Use, &c. Found in North America, in Pennsylvania, Carolina, and Virginia. It was discovered by Michaux, and by him sent to Europe, where it may be found in several botanic gardens, and in some nurseries. Plants, in London, cost 2s. 6d. each; at Bollwyller, ?; and at New York, 50 cents.

§ 11. *C. Simsii* Swt. Sims's Clematis.

Identification. Sweet's Hort. Brit., p. 1.; Don's Mill., 1. p. 8.
Synonymes. *C. cordata* Sims; the heart-shape-leaved Clematis.
Engravings. Bot. Mag., 1816, and our fig. 16.

Spec. Char. Peduncles 1-flowered. Leaves pinnate; leaflets cordate, acuminate, entire, ciliated, reticulated. Sepals 4, coriaceous, connivent, lanceolate, reflexed at the apex, curled. (*Don's Mill.*, 1. p. 8.) Flowers lilac. June, August. 1812. Height 8 ft.

Description. The general appearance of this plant is said to give the idea of something between *C. crispa* and *C. Viorna*; and it is said also to bear some resemblance to *C. reticulata*.

Geography, History, &c. It is found in Georgia and Carolina, and was first brought to England in 1812, probably by Lyon, who made a large importation of plants in that year. It appears to have flowered for the first time in England, in Colvill's Nursery, in 1822, whence it was figured by Watson. It is now to be met with in very few collections.



16

§ 12. *C. RETICULATA* Walt. The net-veined-leaved Clematis.

Identification. Walt. Fl. Car., 156.; Gmel. Syst., 873.; Michx. Fl. Bor. Am., 1. p. 318.; Pursh Fl. Bor. Amer., 2. p. 385.; Dec. Prod., 1. p. 7.; Don's Mill., 1. p. 8.

Synonymes. *C. rosea* Abbott; the netted Virgin's Bower; the reticulated Clematis.

Engravings. Dend. Brit., t. 72.; and our fig. 17.

Spec. Char. Peduncles 1-flowered. Sepals connivent. Leaves coriaceous, netted with nerves, smooth, pinnate; leaflets stalked, 3-lobed or entire, ovate. (*Don's Mill.*, 1. p. 8.) Sepals, when expanded, divaricate, and are of a pale purplish red. June, July. 1812. Height 8 ft.

Description. In Don's *Mill.*, the flower of this species is stated to resemble that of *C. Viorna*; but, by the figure in Watson's *Dendr.*, it is quite dissimilar. In *C. Viorna* the sepals do not divaricate, except in their recurved tips: *C. reticulata* is depicted with sepals expanded in the mode of those of *C. Viticella*. A side view of a flower less expanded resembles more the flower of *C. cylindrica*, but the cylindrical portion is shorter. The flowers (sepals) of the two are different in colour. The reticulation of the veins, in the leaves of *C. reticulata*, is the character expressed in the specific epithet. The stems of *C. reticulata* are not truly ligneous. In the new edition of the *Bot. Mag.*, edited by Dr. Hooker, and published in 1833, *C. Simsii* and *C. reticulata* are made synonymous.



17



+

§ ii. *Viticella* Dec.



Derivation. From *viticula*, a small vine; on account of the plants climbing like the *Vitis vinifera* L.

Sect. Char. Involucre wanting. Tail of the pericarp (that is, of the carpel) short, beardless. Leaves ternate, or decomposed.—Stems climbing. (*Don's Mill.*, 1. p. 9.) Deciduous.

§ 13. *C. FLORIDA* Thun. The florid, or showy-flowered, Clematis.

Identification. Thunb Fl. Jap., 210.; Jacq Hort. Schön., 3. p. 57.; Dec. Prod., 1. 8.; Don's Mill., 1. p. 9.

Synonymes. *Atrage indica* Desf.; *Atrage florida* Pers.; Clematis à grandes Fleurs, Fr.; grossbluthige Waldrebe, Ger.

Engravings. Sims's Bot. Mag., t. 874.; Andr Bot. Rep., t. 402.; Jacq Hort. Schön., 3. t. 357.; and our fig. 18.

Spec. Char. Peduncles 1-flowered, longer than the leaves. Leaves ternately decomposed; leaflets ovate, acute, quite entire. Sepals oval-lanceolate,

much pointed. (*Don's Mill.*, i. p. 9.) Flowers white. April to September. 1776. Height 15 ft.

Variety. *C. f. flore pleno*, the double-flowered florid Clematis (*Don's Mill.*, i. p. 9.), is comparatively common in British gardens. It is very handsome, but is not thought so ornamental as the single-flowered variety.

Description. The stem is slender and striated; climbing to the height of 15 feet or upwards where it is trained to a wall with a favourable exposure. It never, however, becomes very woody. The flowers are large, and very handsome either in a single or double state. This species deserves to be recommended for the great size of its greenish-white flowers, especially when they are not double, and the neatness of its foliage. In addition to this, the slenderness of its stems and branches gives such an air of elegance to it, that no lover of plants for their beauty of appearance should be without it, who has a situation in which it will thrive.

Geography. Found in Japan, and introduced into England, about 1776, by Dr. Fothergill. North of London it requires a warm situation; and in Scotland, as well as in France and Germany, it is generally kept in the greenhouse. The best situation for this species is against either a north or south wall; and, where plants can be trained against both, the flowering season will, of course, be continued much longer than if they were planted against one only. In the sunny site, a loamy soil will be best; but, in the northern, heath mould, that is not in a condensed condition, will be most congenial. A mode of pruning plants of this species, by cutting them down to the ground annually, though not generally practised, is said to produce vigorous shoots and fine flowers. This species, in England, seldom ripens seeds, and is therefore generally propagated by layers. Plants, in London, cost 1s. 6d. each; at Bollwyller, ?; and at New York, ?.



‡ 14. *C. VITICÉLLA* L. The Vine-bower Clematis.

Identification. Lin. Sp., 762; Dumont, 4. 422; Dec. Prod., 1. p. 9; Don's Mill., 1. p. 9.

Synonymes. *Viticélla deltoidea Moench*; the red-flowered Lady's Bower, *Gerard*; *Italienische Waldrebe, Ger.*

Engravings. Flor. Græc., t. 516; Curt. Bot. Mag., t. 565; E. of Pl., 7971; and our fig. 19.

Spec. Char. Peduncles 1-flowered, longer than the leaves. Leaves ternately decomposed, lobes or leaflets entire. Sepals obovate, spreading. (*Don's Mill.*, i. p. 9.) Flowers blue or purple. June to September. 1569. Height 15 ft.

Varieties and their Synonymes.

‡ *C. V. 1 carúlea.* The blue-flowered Vine-bower Clematis.

‡ *C. V. 2 purpúrea.* The purple-flowered Vine-bower Clematis.

‡ *C. V. 3 múltiplex* G. Don. The double-flowered Vine-bower Clematis.— Flowers double, blue. *C. pulchélla Pers.* This variety produces more robust, more extended, and fewer shoots than the single-flowered blue or purple varieties; and there is a degree of dissimilarity about it, which might lead distinguishers on minute differences to regard it as of a species distinct from *C. Viticélla*: it is probable that this dissimilarity was the ground of Persoon's naming it *C. pulchélla*.


‡ *C. V. 4 tenuifolia* Dec. The slender-leafted Vine-bower Clematis.— Leaflets oblong-lanceolate. *C. tenuifolia lusitánica Tourne.*

‡ *C. V. 5 baccáta* Dec. The berryed-fruited Vine-bower Clematis.


Description. The stem rises to the height of 10 ft. or 15 ft.; the leaves branch out into many divisions, and the flowers are supported on long

slender peduncles, which render them more obvious; and, as in the case of all the large-flowered species of *Clématis*, are most favourably seen when they are somewhat above the eye. The double-flowered variety is produced by the change of stamens into petals. The single flowers have no petals, but only sepals. *C. Viticélla*, and all its varieties, are tolerably robust and vigorous in their growth, and decidedly ligneous; though plants individually do not endure many years, probably owing to their exhausting the soil in which they grow. Perhaps no mode of disposing plants of this species, for enjoying the effect of their flowers, is preferable to that of planting them so that their branches may be trained over a frame or fence of trellis-work, with both sides free; in which case the clematis will grow and spread so rapidly as to render the fence or hedge in a short time quite a wall of green.

Geography, History, &c. Found in the south of Europe, in hedges and among bushes, particularly in Spain, Portugal, Carniola, &c. It was cultivated in England in 1569, by Mr. Hugh Morgan, being one of the earliest introduced plants on record. This species is perhaps the most beautiful and most estimable of all the kinds of clematis, for the purposes of floral decoration. For the mere covering of bowers and other objects, it is less suited than *C. Vitalba*, *virginiana*, or perhaps even *Flámmula*, as they grow faster, extend farther, and each yields a greater aggregate of herbage, and so covers better; but none of them can vie with *C. Viticélla* and its varieties in beauty; more especially with the single purple and the single blue. The size of the flowers; their being projected on peduncles just long enough to make them obvious beyond the foliage; their being numerous; their conspicuous colour, and their transparency, render their effect extremely beautiful, especially when seen in masses with the sun shining behind them. *C. Viticélla* is more generally cultivated than any of the other species, and may be purchased in all the principal nurseries of Europe. The price, in London, is 25s. a hundred for all the varieties, except the double purple, which is 75s. a hundred; at Bollwyller the species is 80 cents, and the double-flowered variety 2 francs 50 cents; and in New York, ?.



19



20



♣ 15. *C. CAMPANIFLORA* Brot. The bell-flowered Clematis.

Identification. Brot. Flor. Lus., 3. p. 352; Dec. Prod., 1. p. 9;

Don's Mill., 1. p. 9; D. Don, in Sw. Br. Fl.-Gard., 2. s. 217.

Synonymes. *C. viornoides*, received at the Chelsea Botanic Garden by this name from the Berlin Botanic Garden (D. Don, in Sw. Fl.-Gard., 2d ser., t. 217.); *C. viornoides* Schrader, Hort. Brit., No. 28757; *C. parviflora* Dec., according to Sweet.

Engravings. Lod. Bot. Cab., 987.; Sw. Br. Fl.-Gard., 2d series, t. 217.; and our fig. 20.

Spec. Char. Peduncles 1-flowered, somewhat longer than the leaves. Leaves biternately decomposed; leaflets entire, or 3-lobed. Sepals half spreading, dilated at the apex, wavy. (Don's Mill., i. p. 9.) Portugal. Flowers white tinged with purple. June and July. 1810. Height 10 ft.

Description. The habit of growth of this plant is entirely that of *C. Viticélla*, to which it also comes nearest in affinity; but the much smaller flowers, and pointed sepals connivent below, will readily distinguish it. (Sw. Br. Fl.-Gard., 2d series, t. 217.) De Candolle states that this is an intermediate species between *C. Viticélla* and *C. crispa*.

Geography, &c. Found in Portugal in hedges, more especially on the road from Coimbra to Oporto. It appears to have been cultivated in England since 1810. It is a free grower and flowerer, though not so ornamental as *C. Vitiçélla*. It is in the Horticultural Society's Garden, and in the arboretum of Messrs. Loddiges. Price, in London, 1s. 6d.; at Bollwyller, 1 franc 50 cents; at New York, ?.

¶ 16. *C. CRI'SPA L.* The curled-sepaled Clematis.

Identification. Lin. Sp., 765.; Dec. Prod., 1. p. 9.; Don's Mill., 1. p. 9.; Thunb. Fl. Jap., 239.

Synonyme. *C. fôdre crispo Dil. Elth.*

Engravings. Dil. Elth., 1. t. 73. fig. 84.; Bot. Mag., 1982.; E. of Pl., 7975.; and our fig. 21.

Spec. Char. Peduncles 1-flowered, shorter than the leaves. Leaves entire, 3-lobed, or ternate, very acute. Sepals connivent at the base, but reflexed and spreading at the apex. (*Don's Mill.*, i. p. 9.) North America. Flowers purple. July to September. 1726. Height 3 ft.

Description. The flowers of this species are pretty, but perhaps never produced in sufficient quantity to render it highly decorative; though it is very interesting, both in its foliage and in its flowers. The flower is of a pale purple colour; the sepals having their bases approximated so as to form a tube, and their tips spread or reflexed; these are also wavyly crisped with transverse wrinkles. The stems are weak, and do not generally rise higher than 3 ft. or 4 ft.



Geography. Found in Virginia and Carolina, in hedges and among bushes on the banks of rivers. It is also said to be a native of Florida and of Japan. It was cultivated by Miller in 1726, and in the Eltham Garden about the same time. The plants frequently die down to the ground, so that they require to be treated more as herbaceous than ligneous. The species is in most botanic gardens, and in some nurseries. Price, in London, ? 3s. 6d.; at Bollwyller, ?; and in New York, 25 cents.

§ iii. *Cheirópsis* Dec.



Derivation. From *cheir*, the hand, and *opsis*, resemblance; in allusion to the form of the bracteas.

Spec. Char. Involucre in the form of a calyx, from two joined bracteas situated at the top of the peduncle just under the flower. Tails of pericarps bearded.—Climbing or rambling shrubs, with simple or ternate leaves. (*Don's Mill.*, i. p. 9.) The old petioles persistent, and the new leaves and the peduncles produced in clusters from the axils of these. (*Dec. Syst.*, i. 162.) Evergreen.

¶ 17. *C. CIRRHOSA L.* The tendriled-petioled Clematis.

Identification. Lin. Sp., 766.; Willd. Sp., 2. 1827.; Lamarek Dict. Ency., 2. 43.; Dec. Prod., 1. p. 9.; Don's Mill., 1. p. 9.

Synonyms. *Atrágene cirrhôsa Pers. Syn.*, 2. p. 98.; Traveller's Joy of Candia, and Spanish Traveller's Joy, *Johnson's Gerard*; Spanish wild Climber *Parkinson*; the evergreen clematis; Clematite à Vrilles, Clematite toujours verte (*Bon Jard.*), *Fr.*; einfachblättrige (simple-leaved) Waldrebe, *Ger.*

Derivations. The word *cirrhôsa*, which means cirrhose, or tendriled, is applied to this species from the peculiarly grasping and tendril-like action of its petioles, which retain their hold even after the leaflets have fallen. The French word *Vrilles* signifies tendrils; and the German word *einfach* alludes to its comparatively simple leaves.

Engravings. *C. cirrhôsa L.*, Smith's Flor.-Gr., 517.; *C. c. 2 pedicellâta Dec.*, Bot. Mag., t. 1070.; and our fig. 22.

Spec. Char. Peduncles 1-flowered, with an involucre. Leaves ovate, somewhat cordate, toothed, in fascicles. (*Don's Mill.*, i. p. 9.) Evergreen. Flowers whitish. March, April. 1596. Height 10 ft.

Variety and its Synonyms.

¶ *C. c. 2 pedicellâta Dec.* *Pedicelled-flowered tendriled Clematis.*—The chief feature distinctive of this variety from *C. cirrhôsa* is, that the

pedicel between the involucre and the flower is of some length, and causes the flower to seem pedicellated beyond the point of the place of the involucre. *C. balearica* Pers.; *C. pedicellata* Swt. *Hort. Brit.*, p. 2, *Don's Mill.*, 1. p. 9.; *C. cirrhosa* Sims, *Bot. Mag.*, t. 1070.

Description. An elegant evergreen climbing shrub, rising to the height of 8 ft. or 10 ft., and branching freely, so as to become, in two or three years, a very thick bushy plant. The leaves vary from simple to ternate; and, from being entire to being deeply cut. The flowers appear at the end of December, or the beginning of January, and continue till the middle or end of April. They are pendulous and bell-shaped, the mouth being of the breadth of a shilling, or more. Their colour is greenish white, with some purple on the inside. The sepals are downy without, and smooth within. The principal beauties of this species consist in its bright evergreen verdure, and earliness of its flowering in spring; and they may be best obtained by training it against a wall with a southern aspect.



Geography, History, &c. Found in the south of Europe and north of Africa, in hedges and among bushes, particularly in Spain, Majorca, Sicily, Calabria, Algiers, and the islands of the Archipelago. In its native country it is said to climb up and overwhelm the trees; but in England it is a weak plant, not very readily kept. In Loddiges's Nursery it is cultivated in pots and kept in a green-house, or in a cold frame. Miller observes that it stood in the Chelsea Botanic Garden, in the open air, in a dry sheltered situation; and that it flowers better when so treated, than if kept in a house. It is liable to perish, however, in exposed situations. It was first discovered by Clusius in 1565, and is said to have been cultivated by Gerard in 1596; though, as he says that he found it wild in the Isle of Wight and near Waltham Abbey, it was probably some less tender species which he designates by this name. It is not often met with, except in botanic gardens. In Scotland, and in France and Germany, it is kept in the green-house. In London, it costs 1s. 6d.; at Bollwyller, ?; and at New York, 25 cents a plant.

Æ 18. *C. BALEARICA* Rich. The Minorca Clematis.

Identification. Rich. in *Jour. Phys.*, Feb. 1779, 127.; Lamarck's *Dict. Ency.*, 2. p. 43.; Dec. *Prod.*, 1. p. 9.; *Don's Mill.*, 1. p. 9.

Synonymes. *C. calycina* Ait. *Hort. Kew.*, ed. I. vol. 2. p. 98.; *Clematite de Mahon*, Fr. *Engraving.* Sims, *Bot. Mag.*, t. 959.

Spec. Char. Peduncles 1-flowered, with an involucre under the flower. Leaves ternate; leaflets stalked, 3-lobed, deeply toothed. (*Don's Mill.*, i. p. 9.) Evergreen. Flowers whitish. Feb. and March. 1783. Height 10 ft.

Description. Evergreen, and decidedly ligneous; perhaps rather tender. It attains the height of 10 ft. at least; and is pleasing in appearance, both as regards its leaves and flowers. The leaves have their segments narrow, and toothed and lobed, so that they possess fulness of character. The calyxes are larger than those of *C. cirrhosa*; the sepals are whitish, and marked in the inside with a few blotches, not regularly disposed; and, although the flowers are not very showy, they are produced at a season which renders them very grateful. This species, in the green-house, will yield flowers throughout the winter.

Geography, History, &c. Found in Minorca, and first described by L'Héritier. The plant was brought to England by M. Thouin in 1783; and,

after being cultivated in the garden at Kew, it found its way into most of the other botanic gardens, and into some nurseries. It is easily propagated by layers or cuttings; and, when finally planted out, it is the better for having the protection of a wall. North of York, it may be considered a frame or green-house plant, which it is in France and Germany. It is hardy about London, and in the botanic garden of Cambridge.

§ iv. *Anemoniflora.*



Derivation. From the flowers being like those of the *Anemone sylvestris* L.

Sect. Char. Peduncles axillary, 1-flowered, aggregate, not bearing an involucre. Carpel with a feathery tail. Leaves deciduous.

‡ 19. *C. MONTANA* Ham. The Mountain Clematis.

Identification. *C. montana* Ham. MSS.; Dec. Prod., 1. p. 9.; Wall. Pl. As. Rar., 3. p. 12.; Royle, Illustr. Bot. Himalaya, p. 51.

Synonymes. *C. anemoniflora* D. Don; Prod. Fl. Nepal., p. 192.; G. Don's Mill., 1. p. 9.

Engraving. Wall. Pl. Asiat. Rar., 3. p. 12. t. 217.; Swt. Br. Fl.-Gard., 2. s. t. 253.; and our figs. 23. and 24. Fig. 23. is from the plant in the Hort. Soc. Garden, and fig. 24. from a specimen of that at Montreal, Kent.

Spec. Char. Peduncles 1-flowered, not bracteated, several together. Leaves ternately parted, the segments ovate-oblong, acuminate, toothed, the teeth in the mode of incisions. Sepals elliptic-oblong, mucronulate, spreading. Himalayan Mountains (*D. Don*, in *Swt. Br. Fl.-Gard.*, 2d series, t. 253.) Flowers white. May, in England. 1831. Height 15 ft.

Description. A highly ornamental species. The plant is large and branching; the bark thick, ash-coloured, and deciduous. Leaves several together, upon footstalks 1 in. long; their segments, or leafy parts, pale green. Flowers numerous, about the size and form of those of *Anemone sylvestris* L., borne several together, each upon a separate, upright, slender peduncle, about 3 in. long. Sepals 4, 1 in. long, pure white, faintly stained with pink outside at the base. Styles clothed with long white silky hairs; from which it may be inferred that this species will have its fruits terminated with feathery tails, in a state of maturity.

Geography, History, &c. Its native localities are given by Mr. Royle, in his *Illustrations of the Natural History of the Himalayan Mountains*, as "Mussooree, and every where in the Himalayan Mountains, between 5000 ft. and 7000 ft. of elevation" above the level of the sea, where it flowers in April. In the climate of England, Mr. D. Don has stated that it "proves to be quite hardy, and seems to flourish as well as on its native mountains." He received flowering specimens of it in May, 1834, from Montreal, Kent, the seat of Earl Amherst. Dr. Buchanan, whose name was afterwards changed to Hamilton, originally collected specimens of this species at Chitlong, in the valley of Nepal; and from specimens derived from him, in Mr. Lambert's herbarium, it was first described by De Candolle in his *Systema*, vol. i., published in 1818. Plants were soon afterwards raised from seeds in the garden of the



London Horticultural Society, where one plant, on a wall exposed to the east, grows vigorously without any protection, and flowers freely in warm summers. On the whole, it is a very desirable species. It grows best in "a loamy soil, and is readily multiplied by layers." (*D. Don.*)

App. i. *Doubtful Species of Clématis.*

In Sweet's *Hort. Brit.* are enumerated *C. terniflora* Dec., said to be introduced (from what country is uncertain) in 1826; *C. biternata* Dec., from Japan, in 1825; *C. dahurica* Dec., from Dahuria, in 1822; *C. diversifolia* Dec., of uncertain origin; *C. japonica* Dec., from Japan, in 1826; *C. semitriloba* Dec., from Spain, in what year is uncertain; *C. viornoides* Jac. (which we have considered as a synonyme of *C. campaniflora*), in 1826. These species, or names, will be found followed by specific characters in De Candolle's *Prodromus*, and in Don's *Miller*; but, as they are not now to be met with in the gardens about London, we have deemed it not advisable to occupy our pages by describing them.

App. ii. *Anticipated Introductions.*

On recurring to the lists in the historical part of this work, and comparing the names there given with those of the species described in the foregoing pages, it will be found that we are already in possession of all the species of *Clématis* which are considered to be natives of Europe. Of those of Asia, *C. nepalensis*, *pubescens*, *vitifolia*, and *Buchananiana*, according to the list in p. 173, may be expected from the Himalaya. From China, *C. intricata* (mentioned p. 176.) may be expected; and, from North America, there are the names *C. holosericea*, *Walteri*, and *Catesbyana*, which are not in our *Catalogues* as already introduced. There are probably other species in the mountainous regions of Asia, and in China, which will endure the open air in England; and, as the seeds of the genus are light, not bulky, and very tenacious of life, the probability of their growth will amply repay travellers for the trouble of collecting them.

GENUS II.



ATRA'GENE L. THE ATRAGENE. *Lin. Syst.* Polyándria Polygýnia.

Identification. *Lin. Gen.*, p. 615.; *Willd. Baum.*, p. 45.; *Dumont*, 4. p. 426.; *Hayne Dend.*, 118.; *Don's Mill.*, 1. p. 10.

Synonymy. *Clématis Lam. and Dec.*; *Atragene, Fr. and Ger.*

Derivation. The name of *Atragene* appears to be taken from two Greek words; *athros*, pressed, and *genos*, birth; alluding, as it is supposed, to the manner in which the branches press against or clasp the trees that support them. It was first used by Theophrastus, and was by him applied to *Clématis Vitálba L.*

Gen. Char. *Involucre* none. *Sepals* 4, somewhat induplicate in the bud. *Petals* numerous, shorter than the sepals. *Cariopsides* (carpels) terminated by a bearded tail. *Cotyledons* approximate in the seed. — Climbing deciduous shrubs, with variously cut opposite leaves. (*Don's Mill.*, i. p. 10.) Perhaps no genus was ever distinguished from another on slighter differences than those extant between *Clématis* and *Atragene*. These are, the presence of petals in the flowers of the latter genus; though this is scarcely the case in *A. ochoténsis*; and, in the double-flowered variety of *C. Viticélla*, the metamorphosed stamens, which give the flower its fullness, are considered to be petals. Hence De Candolle regards the different species of *Atragene* only as a section of the genus *Clématis* (§ iv. *Atragene, Prod.*, i. p. 9.); but, as

the genus is retained in Don's *Miller*, and is current among British and German cultivators, we have thought it best to preserve it. On the same principle (that of simplification), we prefer retaining the genera *Mahònia*, *Sórbus*, *Málus*, *Ábies*, *Làrix*, *Cèdrus*, and others; though, in a strictly scientific point of view, they may not be valid.

Description. The atragenes differ from the clematides in producing leaves and one flower from the bud contemporaneously; whereas in most clematides the flowers are produced upon wood developed previously to their appearance, and during the same season. Hence the winter buds of *Atrágene* are larger than those of *Clématis*, from their including the flower as well as the leaves of the year succeeding. In atragenes the leaves are less divided than in many of the species of *Clématis*, and they are always divided ternately. All the species of *Atrágene* described in this work have petioles which not only clasp objects like those of *Clématis*, but maintain the prehension effected for more than the season, like the vine. All the species are extremely interesting from the beauty of their blossoms.

Geography, History, &c. The hardy species are found in the south of Europe, one in Siberia, and one in North America. They are not numerous; and it appears to us doubtful, whether, in reality, more than two hardy species have been yet discovered. This genus is comparatively of recent introduction into British gardens; but, as it is ornamental, plants of it may be procured in most nurseries. The culture is the same as in *Clématis*, and generally by layers.

§ 1. ATRÁ'GENE ALPI'NA L. The Alpine Atragene.

Identification. Lin. Sp., 764.; Willd. Baum., 45.; Dumont, p. 426.; Hayne Den., p. 116.; Don's Mill., 1. p. 10.

Synonymes. *Clématis cærúlea Bauh.*; *Atrágene austríaca Scop.* and *Bot. Mag.*; *Atrágene clematides Crantz*; *Clématis alpina Mill. Dict.*, No. 9.; *C. alpina Dec. Prod.*, 1. p. 10.; *Atragène des Alpes, Fr.*; *Alpen Atragene, Ger.*

Engravings. Bot. Rep., t. 180.; Bot. Mag., t. 530.; and our fig. 25.

Spec. Char. Peduncles 1-flowered, longer than the leaves. Leaves biternate; leaflets ovate-lanceolate, acuminate, serrate. Petals somewhat spatulate, blunt. (*Don's Mill.*, i. p. 10.) Austria. Flowers blue. May to July. 1792. Height 8 ft.

Varieties. De Candolle mentions its varying with white flowers; and *A. sibírica* Lin., described below as a species with yellowish white flowers, appears to us nothing more than a variety of *A. alpina*.

Description. The stems are numerous, branching, weak, forming knots at the joints where the leaves and flowers are protruded. One flower on a longish scape springs from between the leaves; the sepals are twice the length of the petals, and are blue on both sides. The petals are of a dirty white colour, and, in general, 12 in number.

Geography, History, &c. Native of the mountains of the south of Europe, from the height of 2400 to 6000 feet, especially on a calcareous soil, in Austria, Carniola, Piedmont, Dauphiné, the Eastern Pyrenees, Hungary, &c. It appears to have been first remarked by Allioni and Jacquin, in Switzerland. The species is very ornamental, and is to be had in most nurseries. The price, in London, is 1s. 6d. a plant; at Bollwyler, ?; and at New York, ?.



25

§ 2. A. SIBI'RIKA L. The Siberian Atragene.

Identification. Sims, Bot. Mag., t. 1951.; Don's Mill., 1. p. 10.

Synonymes. *Atrágene alpina Gmel. Sib.*, 4. p. 194., *Pall. Flor. Ross.*, 2. p. 69.; *Clématis sibírica Mill. Dict.*, No. 12., and *Dec. Prod.*, 1. p. 10.

Engravings. Sims, Bot. Mag., t. 1951.; *Pall. Fl. Ross.*, 2. p. 69. t. 76., and our fig. 26.

Spec. Char. Peduncles 1-flowered, almost equal in length with the leaves. Leaves biternate; leaflets oblong-lanceolate, acuminate, serrated, petals emarginate at the apex. (*Don's Mill.*, i. p. 10.) Siberia. Flowers white. June or July. 1753 Height 12 ft.

Variety. A blue-flowered variety of this species is mentioned, in *Bot. Mag.*, t. 1591.

Description. There is a considerable similarity in this to the last, in foliage and habit of growth; but it is less robust and less branching; its branches are more ligneous-looking, and the segments of the leaves longer. The calyxes of the flower are white, longer, and with the tips rather connivent than spreading. The whiteness of the calyxes seems to be partaken of by the bark and foliage, as these are of a lighter colour than in *A. alpina*, the calyxes of which are of a blue colour. The flowers of *A. sibirica* are longer than those of *A. alpina* and perhaps less numerous.

Geography, History, &c. Found in the mountainous districts of Siberia, as far as the Eastern Ocean, covering the shrubs and underwood, much in the same manner as *Clematis Vitalba* does in England. Pallas observes that it flowers profusely in June, and that in autumn it delights the eye with its clusters of white feathery seeds. The plant is not very common in British gardens, and has been chiefly cultivated by Messrs. Loddiges.

‡ 3. *A. OCHOTENSIS* Pall. The Ochotskoi Atragene.

Identification. Don's Mill., i. p. 10.; Sweet's Hort. Brit., p. 2.

Synonymes. *Atragene violacea* Pall.; *Clematis ochotensis* Poir.; and *Dec. Prod.*, i. p. 10.

Spec. Char. Peduncles 1-flowered. Leaves biternate; leaflets oblong-lanceolate, acuminate, serrated. Petals few, linear. (*Don's Mill.*) Siberia. Flowers violet. From May to July. 1818. Height 12 ft.

Description. The plant bears a strong resemblance to *A. sibirica*, of which it is probably only a variety. The flowers have four ovate mucronate sepals, which are downy in the margin. Petals none, or sometimes with the outer stamens abortive, a little elongated, and linear. Filaments velvety, a little shorter than the sepals. (*Don's Mill.*, i. p. 10.)

Geography, History, &c. Native of Siberia, towards the Ochotskoi Sea; and of Kamtschatka, between Ochotsk and Kantsch. It appears to have been introduced into England in 1818, but we have not seen it in cultivation.

‡ 4. *A. AMERICANA* Sims. The American Atragene.

Identification. Sims, in *Bot. Mag.*, t. 887.; Swt. Hort. Brit.; Don's Mill., i. p. 10.

Synonymes. *Clematis verticillaris* Dec. *Prod.*, i. p. 10.

Engravings. *Bot. Mag.*, 887.; E. of Pl., 7965., and our fig. 27.

Spec. Char. Peduncles 1-flowered; leaves whorled, in fours, ternate; leaflets stalked, cordate, lanceolate, acuminate, entire or somewhat lobed or serrated. Petals acute. (*Don's Mill.*, i. p. 10.) North America. Flowers purplish blue. May to July. 1797. Height 15 ft.

Variety.

‡ *A. a. 2 obliqua* Douglas MS. *The oblique American Atragene.*—Leaflets bluntly serrated. (*Don.*)‡

Description. This species is distinguishable from all the other Clematideæ described in this work, by the peculiarity of its leaves being disposed, not oppositely, in alternately decussating pairs, but in whorls of four. This is an anomalous characteristic, which De Candolle has expressed by his specific epithet *verticillaris*. The flowers are large, of a palish purple, and less showy than those of *A. alpina*.

Geography, History, &c. Found in North America, in shady places, on the sides of rivulets, climbing and creeping among loose rocks; at New York and in Pennsylvania; near the foot of the Blue Mountains; on the eastern declivity of the Rocky Mountains; and at Cape Mendocina, on the north-west coast. Douglas's variety is found on the eastern declivity of the Rocky Mountains, in valleys; and at Cape Mendocina, on the western coast. The species appears

26



27



to have been introduced into England in 1797: it is found in some gardens and nurseries. The price, in London, is 1s. 6d. a plant; at Bollwyller,?; and at New York, 1 dollar.

§ 5. *A. OCCIDENTALIS* Horn. The Western Atragene.

Identification. Sprengel's Syst.; Swt. Hort. Brit.; Don's Mill.

Synonyme. *Clematis occidentalis* Dec. *Prod.*, 1. p. 10.

Spec. Char. Leaves opposite, ternate; leaflets nearly entire, shining; sepals ovate-lanceolate; flowers somewhat bell-shaped. (*Don's Mill.*, i. p. 10.) Native country, and colour of the flower, unknown. 1818.

Description, &c. We have never seen this species, nor is any thing stated in books respecting it, further than what we have given above.

Sect. II. PÆONIA`CEÆ Dec.

THIS section, or tribe, is the only one contained in De Candolle's second of his two divisions of Ranunculaceæ, namely, in Ranunculaceæ spuria, and is identical with that division. It is at once distinguishable from the other division, named Ranunculaceæ vera, by the character of the anthers opening to admit the escape of the pollen on the side next the ovaries. In the other, the anthers open on the side outward to the ovaries. De Candolle has questioned (*Prod.*) whether the Ranunculaceæ spuria, or Pæoniaceæ, may not be a proper order. They differ from Clematideæ in the character named, and, besides, in having the æstivation imbricate, and the carpels from one-seeded to many-seeded. The ligneous species are included in two genera, Pæonia and Xanthorrhiza, and of the characteristics of these the following are contrasting ones:—

PÆONIA L. Sepals 5, persistent. Petals 5 or more, orbicular, without claws. Stamens numerous. Ovaries 2—5, girded by a fleshy disk. Carpels each containing several seeds.

XANTHORHIZA L. Sepals 5, deciduous. Petals 5, truncately 2-lobed, narrowed into a pedicel. Stamens 5—10. Ovaries 5—10, not girded by a fleshy disk. Carpels each containing 1—3 seeds.

GENUS I.



PÆONIA L. THE PEONY. *Lin. Syst.* Polyándria Di-Pentagýnia.

Identification. The term Pæonia was applied by the Greeks to these plants, which have continued to bear that name ever since.

Synonyms. Peony; Piony; Pivoine, *Fr.*; Gichtterrose, and Páonie, *Ger.*; Rosa del Monte, *Span.*; Peonia, *Ital.*

Derivations. The term Pæonia is said to have been given by Hippocrates and Dioscorides, in commemoration of Pæon, the physician who first used it in medicine. Mr. D. Don has stated (*Sv. Br. Fl.-Gard.*, 2d series, 258.) that it is much more probable that it is derived from Pæonia, a mountainous country of Macedonia, where some of the species grow wild. Most of the other European names are mere adaptations of the classical one, except gichtterrose, *Ger.*, which signifies the gouty rose, from the knobby, or gouty, appearance of the roots of the herbaceous species.

Gen. Char. *Calyx* of 5 leafy, unequal, permanent sepals. *Petals* from 5 to 10, somewhat orbicular. *Stamens* numerous. *Disk* fleshy, girding the ovaries. *Carpels* follicular, from 2 to 5, large, many-seeded, terminated with thick bilamellate stigmas. *Seeds* rather globose, shining. *Leaves* biternate or bipinnate. *Flowers* large, rosy, or rosy and white, usually with a strong disagreeable smell. (*Don's Mill.*, i. 65., with adaptation.) Height from 3 ft. to 10 ft.

Description. There is but one ligneous species, *P. Moutan*; but there are several varieties of this: all are undershrubs, which never attain a great height, and the wood of which always retains a herbaceous character, with a large pith. The roots are ramose rather than tuberous.

Geography, History, &c. The ligneous species is found in China and Japan;

in a wild state in the north of China, and on the mountains in the province of Ho-Nan; and it is cultivated in most gardens in both countries. Its first introduction into England was in 1789. The species and varieties are all beautiful; they are in cultivation in first-rate gardens, and are propagated in the principal nurseries of Europe and America.

✻ *P. Mou'TAN* Sims. The Moutan, or Tree, Peony.

Spec. Char. Segments of leaves oval-oblong, glaucous underneath. Carpels 5, villose. (*Don's Mill.*, i. p. 65.) Height 10 ft.

Identification. Anderson, Linn. Trans.; Dec. Prod., l. p. 65; Don's Mill, l. 65.

Synonyms. *Pæonia arborea* Don, *Hort. Can.*; *P. suffruticosa* Bot. Rep.; Pivoine moutan, and Pivoine en Arbre, Fr.; baumartige Gichtterrose, Ger.; Hoa-Ouang, and Pé-Leang-Kin, Chinese.

Derivations. The word moutan has been applied to this species of peony, in China, for above 1100 years. *P. arborea* and *P. suffruticosa* signify the tree and the sub-shrubby peony. The German name signifies the tree-like gouty rose. The Chinese name Hoa-Ouang signifies the king of flowers, alluding to the beauty of the plant; and Pé-Leang-Kin, a hundred ounces of gold, in allusion to the high price which some of the varieties bear in China.

Varieties and their Distinctions.

- ✻ *P. M. 1 papaveracea* Andrews. *The Poppy-flowered Moutan Peony.*—Petals from 8 to 13, white, with a purple spot at the base of each. Capsules altogether enclosed in the urceolus, or disk. (*Don's Mill.*, i. p. 65.)

Identification. Andr. Bot. Rep.; Don's Mill.

Engravings. Andr. Bot. Rep., t. 463; Lodd. Bot. cab., 547; Bot. Mag., 2175, and our fig. 28.

Introduced in 1806. Mr. D. Don has remarked (*Sw. Br. Fl.-Gar.*, 2d ser., 238.) that the *P. M.*

var. *papaveracea* appears to be really the normal form of the species, as the late Mr. George Anderson suggested.

- ✻ *P. M. 2 variegata* D. Don. *The particoloured-petaled Moutan Peony.*—A low-growing bushy kind, branching from the ground, and scarcely woody. Flowers about 6 in. across. Petals white, stained with a deep rose-colour in various parts; the base marked with numerous radiating streaks of violet and purple. Anthers yellow. The Earl of Mount Norris, whose successful culture of the tree peony has been rewarded by the production of several splendid varieties, far excelling any of those imported from China, has been so fortunate as to raise this fine variety also, which is remarkable for its dwarf and almost herbaceous habit. It was raised from seeds of the *P. M. papaveracea*, which the Earl of Mount Norris supposes had been accidentally fertilised by some of the herbaceous species. All the varieties raised at Arley were from *P. M. papaveracea*, and not from *P. M. Banksii*, as the gardener had inadvertently stated. (*D. Don*, in *Sw. Fl.-Gar.*, 2d ser., t. 238.; *G. M.*, vol. x. p. 284.)

- ✻ *P. M. 3 Banksii* Andrews. *Banks's Moutan Peony.*—Flowers double. Petals slightly tinged with blush, becoming nearly white at the edges, marked at the base with purplish red. In the centre of the flower are some elongated petals, which sometimes appear to rise from amongst the germens. (*Don's Mill.*, i. 65) Cultivated in 1794.

Identification. Anders. Lin. Trans.; Don's Mill.

Engravings. Bot. Rep., t. 418; Bot. Reg., 379; Bot. Mag., t. 1154; and our fig. 29.

- ✻ *P. M. 4 Humei* Ker. Sir A. Hume's *Moutan Peony.*—Flowers double. Petals of the same colour as those of *P. M. Banksii*, with a bunch of long petals rising from the middle of the flower. (*Don's Mill.*, i. p. 63.) Cultivated in Britain as early as 1817.

Engravings. Bot. Reg., 579.

28



29

- ♣ *P. M. 5 ròsea* Dec. *The rosy-petaled Moutan Peony.*—Flowers semi-double. Petals rose-coloured. Segments of leaves with very blunt fissures at the apex. (*Don's Mill.*, i. 65.) Petals large, of a very deep pink. Cultivated in 1794.

Identification. Dec. Prod., 1. 65.; Don's Mill., 1. 65.

This variety is much esteemed, and is at present comparatively high-priced. In *G. M.*, xi. 79., a *P. M. ròsea Courtois* is mentioned, which Dr. Courtois of Belgium has stated to be distinct from the variety known in Britain by that name; and, also, that the latter is identical with *P. M. rùbra Courtois*.

- ♣ *P. M. 6 ròsea semiplèna.* *The semidouble rosy-flowered Moutan Peony.*—This is described, in our *Hortus Britannicus*, as introduced from China in 1794; as attaining the height of 2 ft., and flowering from April to June; and as having red flowers.

- ♣ *P. M. 7 ròsea plèna* Hort. Trans. *The double-rosy-flowered Moutan Peony.*—Flowers very double, of a fine deep pink, nearly scentless. Petals jagged. (*Don's Mill.*, i. p. 65.) Introduced from China in 1804. It flowers from April to June. (*Hort. Brit.*)

Identification. Hort. Trans., 6. p. 477.

Synonymes. *P. suffruticòsa* Andr. Bot. Rep.

Engravings. Andr. Bot. Rep., t. 373.; Bonpl. Pl. Rar., t. 23.

- ♣ *P. M. 8 Rawèsii* Hort. Trans. *Rawes's Moutan Peony.*—Flowers single, pale, slightly tinged with pink. The foliage much resembles that of a herbaceous peony. (*Don's Mill.*, i. 65.) First cultivated in 1820.

Identification. Hort. Trans., 6. 479.; Don Mill., 1. 65.

- ♣ *P. M. 9 càrnea plèna* Hort. Trans. *The flesh-coloured double-flowered Moutan Peony.*—Flowers very double, of a delicate purplish pink, with a rich purple rayed spot at the top of each. It is very like *P. M.* var. *Báncsü*, but has not the central elongated petals of that variety. (*Don's Mill.*, i. 65., with adaptation.)

Identification. Hort. Trans., 6. p. 481.; Don's Mill., 1. 65.

- ♣ *P. M. 10 àlbida plèna* Hort. Trans. *The whitish double-flowered Moutan Peony.*—Flowers double, very pale, suffused with purple. (*Don's Mill.*, i. 65.)

Identification. Hort. Trans., vol. 6. p. 482.; Don's Mill., 1. p. 65.

- ♣ *P. M. 11 Anneslèi* Hort. Trans. *Annesley's* (the family name of the Earl of Mount Norris) *Moutan Peony.*—Flowers small, almost single, of a rich purplish pink. Petals usually 9, obovate, slightly jagged at the margins, of a darker colour at their bases. (*Don's Mill.*, i. 65.)

Identification. Hort. Trans., 6. p. 482.; Don's Mill., 1. 65.

Engravings. Hort. Trans., 6. t. 7.

- ♣ *P. M. 12 làccera* Lindl. *The cut-petaled Moutan Peony.*—This very beautiful variety is strikingly different from the others in the bright rosy red of the petals, the innermost of which are very much cut and gashed, curled up, and distinctly bordered with a narrow edge of light carmine, which sets them off to great advantage, and gives the whole flower a peculiarly rich and finished appearance. Raised in 1831, from seed, by Mr. William Hyland, gardener to the Earl of Sandwich, at Hinchbrook, near Huntingdon. (*Bot. Reg.*, July, 1835; *Gard. Mag.*, vol. xi. p. 418.)

Engraving. Bot. Reg., 1771.

Expected Varieties. Those mentioned under *P. M. 2 variegàta*, above, as raised by the Earl of Mount Norris.—A *yellow-flowered variety* is mentioned by Mr. Main (*G. M.*, ii. p. 423.), as having been imported by Gilbert Slater Esq., in 1794, and which died.—The Chinese pretend to have a *black-flowered variety*, and a *double blue* one, which is only possessed by

the emperor, and which is said to have from 100 to 1000 petals.—It is recommended to those who wish to import plants from Canton, to furnish themselves with good Chinese drawings of the varieties they wish to obtain; and to send these to an agent or collector. Mr. Main also recommends adding to the drawing the Chinese character for the variety, if it can be procured. The word *moutan* is written in Chinese as in *fig. 30*.

Description. The *Pæonia Moutan*, in a sheltered situation, will attain the height of from 6 ft. to 10 ft. in ten years; and no plant can be a more gorgeous ornament of the garden than such a bush, abounding as it does in leaves, striking from their branched character and numerous segments, and in very magnificent flowers, of extraordinary size: both leaves and flowers being produced early in the spring.

Geography, History, &c. *P. Moutan* appears to have been first observed by Europeans in the gardens at Canton, where, however, it is neither indigenous nor propagated; but the plants are brought by the commercial gardeners of the provinces of Nan-Kin and Ho-Nan, where it is a native, and sold by them to the merchants and others possessing gardens at Canton. From Chinese drawings, and from the extravagant praises bestowed upon this plant in the *Mémoires sur la Chine*, published by the missionaries, an ardent desire was excited, in Sir Joseph Banks and others, to import plants into England; and, previously to 1786, Sir Joseph Banks engaged Mr. Duncan, a medical gentleman attached to the East India Company's service, to procure a plant for the royal garden at Kew, where it was first received, through Mr. Duncan's exertions, in 1787. (See *Gard. Mag.*, ii. p. 424.) Various plants have since been imported by different individuals, mostly of the common sort (*P. M. Banksii*), but including also some of the varieties which have been enumerated above. In 1803, the *P. Moutan* was introduced into France, and various other parts of the continent of Europe, having been sent from the Hammersmith Nursery to the garden of Malmaison.

Soil and Situation. On its first importation this plant was grown in sandy peat; but it has since been found to thrive best in deep rich loamy soil. An open situation is preferable, both on account of maturing the wood and leaves, and for displaying the flowers to advantage; but the plant must be sheltered from the north and east, or other cold spring winds, unless it is intended to cover it, when it is in flower, with a movable glass, or canvass, case. The protection given to this plant is necessary, not so much to prevent it from being injured during winter (for it will bear the winters of Paris without any protection, if the wood has been properly ripened), as to protect the tender leaves and flowers when they first appear, in April and May, from being blackened by the frost. Mr. Curtis finds, at Glazenwood in Essex, that by giving the plants no protection whatever, they flower somewhat later than if protected; and this slight retardation, in his part of the country, is sufficient, in most years, to prevent the flowers from being injured by frost. In severe weather, however, when they are in flower, he protects them with poles and mats. (*Curt. Bot. Mag.*, vol. i. p. 26., 1833, new edit.)

Properties and Uses. The whole plant possesses narcotic and poisonous qualities, which are common to the genus. No use, however, is made of the ligneous species, but as an ornamental flowering shrub, as which, it is needless to say, it holds the very first rank.

Propagation and Culture. The Chinese are said to propagate this plant by seeds, in order to procure new varieties; but they also multiply it by parting the roots, and by layers and cuttings; and it is said that "they generally inoculate the buds of different varieties upon the several branches of the same colour. When the time of flowering approaches, they carefully remove all the superfluous buds, in order to strengthen those which they intend to expand; and these they also protect from the scorching heat of the sun." (*Kamp. Amœn.*

30

Exotica, p. 862.) In Britain the tree peony is propagated by seeds, by division of the root, by grafting, by summer budding, by layers, by cuttings, and by single winter buds.

Seeds. These are occasionally ripened by plants bearing single or semi-double flowers. Mr. Curtis finds that seeds are produced by *P. M. papaveræa* in abundance. In England, plants were first raised in this way from *P. M. papaveræa*, at Arley Hall, the seat of the Earl of Mount Norris. They have since been raised at several other places; and, among these, at Glazenwood, where the seeds are sown as soon as gathered in the autumn, and while some of them vegetate in the following spring, many of them remain a year or more before they come up. At Hinchbrook, it was found that the seeds did not germinate till eighteen months after sowing. (*Bot. Reg.*, 1771.; *Gard. Mag.*, xi. p. 418.)

Division of the Root. This operation is so simple, that it is unnecessary to describe it. When the wounds made are large, it may be advisable to seal them, or to cover them with grafting wax. It is necessary, for the success of each divided portion of the main root, that some fibrous roots should be attached to it when taken off.

Grafting. This operation is performed on the roots of herbaceous peonies, at any time from the beginning of September to the middle of March. Select some good tubers of *P. officinalis*, or of any other hardy herbaceous kind, and take off cuttings of *P. Moûtan papaveræa*, or any other tree peony that it is desired to increase. Then slit the tuber from the crown downwards about two inches; form the scion like a wedge; insert it into the slit of the tuber, and make the barks fit on one side as exactly as possible; then bind them well together with strands of good bast matting; over which put one turn of brass wire, to prevent the parts from separating after the strands of bast have decayed. Put the tubers into pots deep enough to allow the mould to cover the top of each tuber; set them in a cold frame or pit; keep them close, rather dry, and defended from the sun during the first month, and from frost during winter. When they have perfected one season's growth, turn them out, and treat them like established plants. (*Gard. Mag.*, iii. p. 293.) We are not aware of any ligneous variety of peony having been grafted on any other variety of ligneous peony.

Budding. This mode, it is stated by Kæmpfer, as quoted above, is practised by the Chinese, which is the more remarkable, as grafting is said to have been unknown to that people till lately. We do not doubt its practicability, though we have never seen it tried.

By Layers. These are made of wood of the preceding year's growth, either in autumn or spring; and tongued and pegged firmly under the soil, between 2 in. and 3 in. They will throw down roots the first year; but it is generally found desirable, to let them remain two years before separating them from the stool. When they are taken off, they should be potted, and kept in pots till wanted for final planting.

By Ringing, by Buds, and by Cuttings. The following directions for these modes of propagation are taken from Maund's *Botanic Garden*:—"In February select any of the stems of the *Pæonia Moûtan*, or all may be used; and, at the distance of half an inch from the centre of each bud, both above and below it, cut out entirely round the stem a small ring of the bark, rather more than the sixteenth of an inch wide, in the manner of common ringing, as practised on fruit trees. Thus every bud will occupy one inch of the stem, where the direct continuation of the bark is obstructed, both above and below, by the rings which have been cut out of it. The stems, so prepared, are then to be laid horizontally about 3 in. beneath the soil, leaving only the leading bud at the end of each branch above the surface. In six months every bud will have made a vigorous shoot, and, in general, will have two radical fibres at its base. In August, remove the soil from above the layers; and, having raised the newly made roots, carefully separate each young shoot from the main layer, by passing a small knife from one ring to the other, cutting out about one third part of the old stem. The young plants should then be immediately potted, to

remain till they are required for planting out in their final situations. After thus gathering the first crop of young plants, the old layers should be again covered with good soil, and left as before; and, in the following summer, a second and greater crop of plants will be produced than in the first season; and, what is most remarkable, they will issue from various parts of the stem, where no trace of a bud was previously indicated. Again, if a stem be detached from the parent plant, and treated as described above, and then laid in soil in a pine-pit or stove, it will shoot almost as freely as if connected with the original root."

Cuttings. "In another experiment, cuttings of about an inch in length were made of the *Pæonia Moutan*, in the manner of vine cuttings, having one bud on each, and about half of the stem behind the bud slit up, and the pith removed. These were put 3 in. deep in pots of soil, and plunged into an exhausted bark-bed, having a temperature of about 60°. In the space of two months, these cuttings made young shoots through the soil, and grew freely." (*Bot. Gard.*)

Nursery Culture. Stools are planted in the London nurseries, either in cold-pits, or in the open ground, to be slightly protected during winter; and, when these have been two or three years established, they throw up abundance of shoots every year, which are laid down, either with or without the ringing process described above, in autumn, or early in spring, and taken off in about a year or two years afterwards. As the tree peony does not transplant well, from the length of its descending roots, and the paucity of their fibres, plants should always be kept in pots, by which means, instead of suffering from removal, they will be improved by it.

Final Culture. The plant has always the best effect when placed singly on a lawn, or in a border, in such a manner as that it may become an orbicular bush, free on all sides. As it is of slow growth, it requires little or no winter pruning, except for the removal of dead or injured wood; but in spring, it is desirable to thin out the embryo blossoms, as soon as they can be distinguished, so as to leave no more than what the plant can bring to a high degree of perfection; and so as not to injure any part of the embryo foliage. The plant has been tried against a wall; but in such a situation the heat appears too great for it in summer. It succeeds well in a conservatory or in a cold-pit. South of London, and in most parts of Ireland, it thrives perfectly well in the open air; though in the latter country it requires a slight roof to be thrown over it while it is in flower.

Accidents, Diseases, &c. The shoots are liable to be killed back by the winter's frost, especially after a wet summer, when they have not ripened thoroughly. In spring, the leaves and blossoms are liable to be blackened, when the plants are not protected by a slight covering stretched over them horizontally during frosty nights. A piece of thin muslin, canvass bunting, or woollen netting, stretched over a wooden frame, formed like an hood, would be quite sufficient for this purpose, and the trouble of covering the plant would not be great. The bitter quality of the leaves prevents them from being attacked by many insects; though the blossom buds are sometimes punctured by them.

Statistics. One of the largest tree peonies within ten miles of London stood, till lately, in the grounds at Spring Grove, where it was planted by Sir Joseph Banks. It was 6 ft. or 8 ft. high, and formed a bush 8 ft. or 10 ft. in diameter in 1825. South of London, there are equally large plants at Rook's Nest, near Godstone, Surrey, which were planted in 1818. North of London, the largest plant in the country (*P. M. papaveracea*) is at the seat of Sir Abraham Hume, at Wormleybury, in Hertfordshire. It is 7 ft. high, and forms a bush 14 ft. in diameter, after having been planted thirty years. It stands the winter, in general, very well; but, if the flower-buds swell too early in February, it becomes advisable to cover the plant slightly with a mat. In the year 1835, this plant perfected 320 flowers; but it has been known to bear three times that number. In most parts of Scotland, the tree peony will grow with protection, and near the sea coast nearly as well as in England. The largest

plants are at Hopeton House, and in Dalkeith Park. In Ireland, the plant attains a large size with little or no protection, as will appear by the notice of one 12 ft. high at Lord Ferrard's, already given in p. 109.

Commercial Statistics. Soon after the tree peonies were imported from China, they began to be propagated in the nurseries, and the price was, at first, ten guineas a plant. In 1820 the price had fallen to six guineas, and in 1825 to two guineas. Price in 1836, in London, from 3s. 6d. to 10s.; at Bollwyller, where it is a conservatory plant, from 10 francs to 30 francs; and in New York, ?.

GENUS II.



XANTHORHI'ZA L. THE YELLOW ROOT. *Lin. Syst.* Polyándria Mono-Tri-gýnia.

Gen. Char. *Calyx* of 5 deciduous sepals. *Petals* 5. *Carpels* 2—3-seeded, but usually solitary from abortion. (*Don's Mill.*, i. p. 65.) There is only one species known.

æ X. APIFO'LIA L'Hérit. The Parsley-leaved Yellow Root.

Identification. *Lin. Gen.*, 1581.; *Lam. Ill.*, 854.; *Dec. Prod.*, 1. p. 65.; *Don's Mill.*, 1. p. 65.

Synonymcs. *Xanthorhiza apiifolia*; *Zanthorise à Feuilles de Persil*, *Fr.*; *Sellerie-blättrige Gelbwurz*, *Ger.*

Derivation. From the Greek words *xanthos*, yellow, and *rhiza*, a root, applied from the deep yellow colour of the roots. The French name needs no explanation; and the German is a literal translation of the English one.

Engravings. *Lam. Ill.*, t. 854.; *L'Hér. Stirp. Nov.*, p. 79. t. 38.; *Bot. Mag.*, 1736.; *E. of Pl.*, 3993.; and our *fig.* 31.

Description. A small shrub with yellow creeping roots, throwing up numerous suckers; with irregularly pinnate leaves, branched racemes, and small purplish flowers, which are usually unisexual from abortion, rising from the scaly buds. (*Don's Mill.*, adapted.) The flowers appear early in May, and continue a month or upwards before they drop off. We have never heard of its ripening seeds in Europe. Nevertheless, this may have often occurred, and been overlooked, from the inconspicuousness of the shrub, and the smallness of its fruit.



Geography, History, &c. Found on shady banks of rivers from Virginia to Georgia, where it flowers in May. It was first described by L'Héritier, and introduced in England, about 1776, by John Bush. It is so readily propagated by suckers, of which it throws up a great number, that it is in most botanic gardens, though, from having no great show in a shrubbery, it is seldom met with in private collections. To us it appears a plant of very great interest, from the uncommon colour of its elegant panicles of flowers, their early appearance in spring, and the finely divided light green leaves, which succeed them. The plant is of so limited a habit of growth, that under no circumstances do we suppose that it would rise higher than 3 ft. or 4 ft., which height it attains in three or four years; but it spreads by its suckers, which, though they are numerous, do not come up at a great distance from the stem. As the shoots are not liable to be killed by frost, and never require pruning, it forms a desirable bush for a shady moist situation. Its roots afford abundance of deep yellow juice; but they have hitherto not been used in dyeing; probably, because there are already so many plants, which grow more rapidly, that afford a yellow colour. Plants are so seldom asked for in the nurseries, that the name of this shrub is seldom to be found in the catalogues. Price, in London, 1s. 6d.; at Bollwyller, ?; and in New York, 25 cents.

CHAP. II.

OF THE HARDY LIGNEOUS PLANTS OF THE ORDER WINTERACEÆ.

This order, which was formerly included in Magnoliaceæ, has been separated from it by Mr. Brown. It is named Winteraceæ, because the *Drimys Winteri* Forst., previously *Wintera aromatica* Murr. (named in honour of Captain Winter, who sailed with Sir Francis Drake), is the type of the order. The diagnostics are, calyx of 2—6 deciduous sepals, and 2 to many petals; the sepals and petals, when more than two, disposed ternarily; carpels whorled, very rarely solitary from abortion; leaves full of pellucid dots. *Illicium* is the only genus of this order which contains species that will stand out in the open air.

GENUS I.



ILLICIUM L. THE ILLICIUM, or ANISEED TREE. *Lin. Syst.* Polyándria
Polygýnia.

Identification. Lin. Gen., 611.; Lam. Ill.; Dec. Prod., 1. p. 77.; Don's Mill., 1. p. 78.

Synonyms. Badiane, or Anis étoilé, Fr.; Sternanis, Ger.

Derivations. The generic name, *Illicium*, is formed from the Latin word *illicio*, to allure, on account of the agreeable aromatic smell of all the species. It is called the Aniseed Tree, from its smell bearing a strong resemblance to that of aniseed. Badiane appears to be an aboriginal French word; Anis étoilé, and Sternanis, signify literally the starry anise, and may allude to the starry disposition of the parts of the flower and of the capsules.

Gen. Char. Calyx of 3-6 petal-like sepals. Carpels stellately disposed, capsular, opening on the upper side, 1-seeded. (*Don's Mill.*, i. p. 79.)

Description. The species are evergreen shrubs, with smooth, shining, oblong, stalked, leathery leaves; generally attaining the height, in their native countries, of from 5 ft. to 12 ft.; and, in this country, of from 3 ft. to 8 ft. in the open air, and more in a conservatory. They are nearly hardy.

Geography, History, &c. These shrubs are found in the southern states of North America, and in China and Japan. One species has been known in Britain since 1766; but the others are of more recent introduction. They are all spicy and aromatic; and, are employed, in their native countries, in the same manner as anise and coriander seeds are in Europe. Northward of London, as well as on the Continent, they are placed in the green-house during winter, or planted in the conservatory. Wherever they are planted, in common with most evergreen trees and shrubs having broad shining leaves, they prefer the shade to the sun; and, consequently, if they are planted against a wall, it ought to be one facing the south-east, or the south west, and never against one full south. They are all slow growers, and, to produce any effect, should never be planted near trees or shrubs which grow rapidly. They may all be planted in the open ground, in warm sheltered situations; but they require protection during winter. They are all easily propagated by cuttings of the ripened wood planted in sand, and covered with a glass, or by layers; and they all grow in a light loamy soil, or in a mixture of loam and sandy peat. One or more of the species is in most botanic gardens; and small plants may be procured, in some of the principal London nurseries, at from 2s. 6d. to 5s. each. The species are *I. floridanum*, *I. anisatum*, and *I. parviflorum*.

• *I. ILLICIUM FLORIDANUM* Ellis. The Florida Illicium.

Identification. Lin. Mant., 395.; Willd. Sp., 2. p. 1294.; Dec. Prod., 1. p. 77.; Don's Mill., 1. p. 79.

Synonyms. The Florida Aniseed Tree, red-flowered Anise-seed tree *Mor. Hist.*; Badiane de la Floride, Fr.; unächter spürnis Sternanis, Ger.

Engravings. Lam. Ill., t. 493. f. 1.; Curt. Bot. Mag., 439.; Lodd. Bot. Cab., t. 209.; E. of Pl., 7901.; and our fig. 52.

Spec. Char. Petals 27—30, dark purple, outer ones oblong, inner ones lanceolate. (*Don's Mill.*, i. p. 79.) West Florida. Evergreen. Flowers of a dark reddish purple. April to June. 1766. Height 6 ft.

Description. A compact, many-stemmed, bushy, evergreen, slow-growing shrub, attaining, in the neighbourhood of London, the height of 6 ft. or 8 ft. or upwards, and flowering every year. The leaves are oblong-lanceolate, quite entire, pointed at both extremities, smooth, shining, and, in common with the whole plant, have a rich reddish hue. The flowers are numerous, solitary, and terminal; and bear some general resemblance to those of *Calycánthus floridus*.



Geography, History, &c. Found in West Florida, on the banks of the river Mississippi, and in marshy places near the town of Pensacola, by Bartram, in 1766. At first this plant was kept in stoves and green-houses; but some specimens were planted out in the open air, by Mr. Gordon, in his nursery at Mile End, where they remained above forty years, sometimes without any protection at all, and at others with only a mat thrown over them, or a slight covering of peas haulm. They grew in a deep, dry, sandy soil, in a warm situation, sheltered from the north-east and east, and open to the south. They were sold, with some other fine specimens, in 1834 and 1835. The properties of this species are of a very decided kind. The leaves and the entire plant are strongly impregnated with a spicy aromatic taste and smell, approaching to that of the seeds of the anise or coriander. The leaves, when bruised, smell strongly of anise, and may be distilled like the seeds of that plant. Though not used in medicine, there is little doubt, Professor Burnet observes, that the bark would answer the same purposes as that of canella, or saffaras. (*Outlines, &c.*, ii. p. 836.) The soil in which this plant is generally grown is a light sandy loam; but Dumont observes that, after trying it in loam, both in the open air and under glass, he found the leaves assume a yellow hue. He then tried pure heath soil, and soon perceived that the leaves had resumed their deep green colour, and the entire plant had begun to grow vigorously. The manner in which the plant is propagated in the London nurseries is, generally, by forming stools of it in a cold-pit, and laying down the shoots, which require two years to root sufficiently to admit of their being separated from the parent plant; but it is sometimes propagated by cuttings both of the young and of the old wood. As soon as the layers are taken off, they are potted, and kept in a green-house or frame till wanted for final transplanting. The situation of such a shrub in a garden should be in a select spot, where it can only be compared with slow-growing plants like itself. Being an evergreen, and finishing its growth early in summer, it would take little harm by being covered with mats or fronds for several months, provided these were supported so as not to touch it, and they were occasionally opened on fine days, on the south side, to admit the sun and air, in order to dry up the damps generated within; a covering of spruce fir branches would also form a very suitable protection. For a small conservatory where there is no means of heating, but from which the extreme cold is excluded, by covering the glasses with mats in severe weather, this, and other species of Illi'cium, would prove very suitable shrubs. Plants may be obtained, in the principal London nurseries, at from 2s. 6d. to 5s. each; in the Bollwyller Nursery, at 4 francs; and in New York, at 1 dollar. They are always sold in pots, being so grown in order that they may be protected by a frame during winter.

❖ 2. ILLI'CIUM ANISA'TUM L. The Anise Illi'cium.

Identification. Lin. Sp., 664.; Gært. Fruct., 1. p. 338.; Dec. Prod., 1. p. 77.; Don's Mill., 1. p. 79.
Synonyms. The Chinese Aniseed tree; Badiane de la Chine, and Anis étoilé, Fr.; ächter (true) Sternanis, Ger.

Engravings. Kämpf. Amœn., t. 881.; Gært. Fruct., 1. p. 338. t. 69.; Lam. Ill., t. 493. f. 2.

Spec. Char. Petals 27—30, yellowish, outer ones oblong, inner ones linear

awl-shaped. (*Don's Mill.*, i. p. 79). China. Flowers yellow. April till June. 1790. Height 10 ft.

Description. A large shrub or low tree, with a thick branchy stem, attaining the height of 20 ft. or upwards in its native country, but seldom appearing of half that height, even in conservatories, in England. The leaves are large, somewhat like those of the common laurel but smaller, lanceolate, thinly distributed on the lower parts of the shoots, but closer together near their points. The wood is hard, and finely scented, but fragile; and the bark is smooth, and russet-coloured. The fruit is composed of from 9 to 12 capsules (carpels) united at their base, and spread out at their extremities in the form of a star. Every carpel contains one oval seed, lance-shaped, and of a russet colour, which encloses a whitish kernel, somewhat oily, and agreeable to the taste, which is similar to that of the anise seed, but is more pungent. The shoots grow at about the same rate as those of the preceding species, and the plant attains its ultimate size of 10 ft. or 12 ft. in twenty or thirty years.

Geography and History. It is found wild in China and Japan, where it is cultivated in gardens as a sacred tree, and also for its carminative and stomachic qualities. It was first described by Clusius, and afterwards seen in Japan by Kämpfer, who has given us a great many interesting particulars respecting it. It was introduced into England in 1790, and soon after into the garden of Courset, near Boulogne-sur-Mer; and it is now occasionally met with in the green-houses of the principal botanic gardens of the middle and north of Europe, and in the open ground of those in Italy.

Properties and Uses. The whole plant is stomachic and carminative, and it is used in the East both medicinally and in cooking. The Chinese chew it after dinner to promote digestion, and as a sweetener of the breath: the same practice prevails in Japan. In some parts of the East Indies the leaves are mixed with tea and sherbet, and the capsules are imported from China into Europe, under the name of Chinese anise, for the purpose of flavouring dishes, and making the liquor called *anisette de Bordeaux*. The Chinese make an infusion of the seeds with the roots of ninsin (*Sium Ninsi* L.), and drink it, when they are fatigued, to recruit their strength and refresh their spirits. They also mix it both with coffee and tea, to improve the flavour. The capsules and seeds are infused in water, and fermented, so as to produce a vinous liquor, very much esteemed, and which the Dutch import under the name of anise arrack. Kämpfer states that a branch of this plant, though not poisonous in itself, if put into a decoction of the poisonous fish called by the Dutch *opblaser* (a species of *Tétrodon*), increases the violence of the poison and makes it occasion death almost instantaneously. (*Kämpf. Amœn.*, fasc. v. p. 883.) The wood, which is called anise wood, is employed in cabinet-work: it is very hard and durable, but is rather brittle while being worked. In China, the watchmen powder the bark, and with it fill long narrow wooden tubes, which are graduated on the outside at regular distances. The powder is then lighted at the farther extremity of the tube, and, as it burns regularly and slowly, it is always the same time in burning a given distance; and the watchman, when he sees by the graduated scale that the flame has reached a certain point, rings a bell, thus forming of it a kind of pyrotechnic clock. The Japanese and Chinese consider the tree as sacred: they burn the bark as incense on their altars, and with the branches decorate the tombs of their friends. (*Thouin, Diet. Agric.*; *Roy., Cours d'Agric.*)

Soil and Situation as in the preceding species, of which this, as Thunberg conjectures, is, in all probability, only a variety. It is, perhaps, rather more tender, and, therefore, should be placed in the warmest situation that can be found, but where it will, at the same time, not be too much exposed to the direct influence of the sun during the hottest months.

The Propagation and Culture are the same as of the preceding species. Plants are to be met with, in the principal London nurseries, at from 2s. 6d. to 5s. each; and at Bollwyller, where the price is 3 francs: in New York, ?.

■ 3. ILLI'CIUM PARVIFLO'RUM Michx. The small-flowered Illicium.

Identification. Mich. Fl. Bor. Amer., 1. p. 326.; Ph. Fl. Amer. Sep^{t.} 2., p. 380.; Dec. Prod., 1. p. 77.; and Don's Mill., 1. p. 79.

Synonymes. Illicium anisatum Bartram; the small-flowered Aniseed Tree; Badiane à petites Fleurs, Fr.; kleinbluthiger Sternanis, Ger.

Engravings. Vent. Cels, t. 22.; Lois. Herb. Amat., t, 330.

Spec. Char. Petals 9—12, yellowish, ovate-roundish. Sepals 3, ovate, somewhat ciliated. (*Don's Mill.*, 1. p. 79.) West Florida. Flowers yellowish white. From May to June. 1790. Height 8 ft.

Description. This species is more upright and free-growing than either of the other kinds. The twigs and stems, which are numerous, are quite green when young; but as they become old they assume a greyish hue. The leaves are of the same consistency as those of *I. floridanum*, and of nearly the same size, but longer, and paler on the under side, with very short petioles. The flowers are smaller, and not so open as those of the other species; and their colour is of a yellowish white. *I. parviflorum* is a low evergreen shrub, or rather bush, highly aromatic in every part when in a growing state; but the scent soon evaporates from every part of the plant after it is gathered, except from the fruit, the scent of which becomes even more fragrant, and more penetrating, when it is dry. In England, this plant seldom produces annual shoots of more than 5 in. or 6 in. in length; and it will probably attain the height of 5 ft. in ten years, and its full size in about twice that period. In Florida it forms a compact evergreen bush from 8 ft. to 10 ft. high.

Geography, History, &c. This shrub was first discovered by Bartram, and afterwards by Michaux, in Florida, on the bank of the lake Georgia; and it was sent by the latter to France, where it was cultivated in Cels's Nursery and in the *Jardin des Plantes*, in 1789. It was kept the first winter in the conservatory, and flowered in the course of the following summer. (*Nouv. Duh.*) It was introduced into England in 1790. The bark has exactly the flavour of the sassafras root, and the dried fruit is used for scenting linen. The propagation, culture, and price, in the London nurseries, are the same as those of the preceding species; at Bollwyller, ?; and in New York, 1 dollar.

CHAP. III.

OF THE HARDY TREES AND SHRUBS BELONGING TO THE ORDER MAGNOLIA' CÆÆ.

THE term *Magnoliææ* is applied to this order, because the genus *Magnolia* is considered as its type. The diagnostic of the order, as given by De Candolle, was made to comprehend some tribes which have since been separated from it; but, as the hardy species still belonging to *Magnoliææ* are all included under the genera *Magnolia* and *Liriodendron*, a sufficient character for them will be: a calyx of 3 deciduous sepals; a corolla of 3—12 petals, disposed in threes; anthers adnate, elongated; carpels numerous, disposed along a spiked axis; leaves destitute of pellucid dots, stipulate when young, the stipules convolute, and enclosing the unexpanded leaves. Evergreen and deciduous trees and shrubs. (*Don's Mill.*, adapted.) The differential characters of the genera *Magnolia* and *Liriodendron* are as follows:—

MAGNO'LIA L. Carpel dehiscent; that is, opening to admit the escape of the seed.

LIRIODE'NDRON L. Carpel indehiscent; that is, not opening to admit the escape of the seed.

GENUS I.

**MAGNOLIA L. THE MAGNOLIA.** *Lin. Syst. Polyándria Polygýnia.*

Identification. *Lin. Gen.*, 690; *Gart. Fruct.*, 1. p. 343; *Dec. Prod.*, 1. p. 79; *Don's Mill.*, 1. p. 82.

Synonymes. *Magnolie, Fr. and Ital.*; *Bieberbaum Hart.*, ; and *Magnolie Willd., Ger.*

Derivations. The name *Magnolia* was given to this genus by Linnæus, in honour of Pierre Magnol, professor of medicine, and prefect of the botanic garden at Montpellier. He was author of *Botanicum Mompeliense*, published in 1676, and of other works. He died in 1715. The German name *Bieberbaum*, beaver-wood, is applied generically by Hartweg in the *Hortus Carlsruhensis*; but, in America, beaver-wood appears to be applied only to *M. glauca*.

Gen. Char. *Calyx* of 3 deciduous sepals, that resemble petals. *Corolla* of from 6—9 petals. *Stamens* numerous. *Pistils* numerous. *Carpels* disposed compactly in spikes, opening by the external angle, 1—2-seeded, permanent. *Seeds* baccate, somewhat cordate, pendulous, hanging out beyond the carpels by a very long umbilical thread. — Trees and shrubs with large entire leaves; and solitary, terminal, large, odoriferous flowers. Trees chiefly deciduous, but partly evergreen. (*Don's Mill.*, adapted.)

Description. One of the species is a lofty evergreen tree; but the others are deciduous, and partly trees and partly shrubs. The flowers of most of the species are white; but in some they are of a greenish yellow, and in others they are tinged with purple. The seeds are mostly of a scarlet colour. The roots are branched, and yet but sparingly supplied with fibres.

Geography. The native country of most of the hardy magnolias is North America; but there are some hardy species found in China and Japan, and, perhaps, also, in the Himalaya. No species of *Magnoliaceæ* has hitherto been found in Europe, Africa, South America, or Australia; and the geographical range of this order in America and Asia is comparatively limited. The first magnolias were imported into Europe about the year 1730, and into France and England nearly about the same time. From that period to the present new species have been introduced at intervals, and some may be expected from the mountainous regions of India. Most of them are to be found in all the botanic gardens, and in the principal European nurseries. All the magnolias are highly ornamental; and though this is their principal use in Europe, yet in America they are valued for their medical properties. The bark of all the species is bitter; and it is used in America for the same purposes as the Jesuits' bark is in Europe. The magnolias may be cultivated in most parts of Britain, and of the middle and southern states of Europe; but, north of London and Paris, some of the species require protection during winter, or to be kept in the green-house. A deep sandy soil, and a situation sheltered from the north and east, will suit most of the species; though some, as *M. glauca*, for example, thrive best in a moist peaty soil. Few of the species ripen seeds in England, but most of them do so in France. From these seeds, or from such as are imported, all the American species are most frequently raised; but the species from Asia are increased by layers, as are the American species very frequently. In France, all the deciduous magnolias endure the open air, even in the northern provinces, without protection during winter, because, in consequence of the warm summers, the wood is thoroughly ripened. In the neighbourhood of Paris, all the species, even those of Asia, ripen seeds; though the evergreen magnolia requires protection there during winter. In Holland and Belgium, the deciduous magnolias thrive nearly as well as in France, and some of them ripen their seeds. In Italy, the magnolias suffer from the heat, except when planted in moist situations, and among other trees. In the north of Germany, in Russia, and in the greater part of Sweden, most of the magnolias are green-house plants.

The hardy species of this genus are included in two sections, *Magnoliæstrum* and *Gwillimia*.

§ i. *Magnoliástrum*.

Derivation. *Magnolia*; and *astrum*, from *ad instar*, an affixed particle, signifying likeness.

Sect. Char. American species, with one spathe-like bractea enclosing the flower-bud; ovaries approximate; anthers bursting outwards. (*Don's Mill.*, i. p. 83.)

† 1. *MAGNOLIA GRANDIFLORA* L. The large-flowered Magnolia.

Identification. Lin. Sp., 755.; Lam. Ill., 490.; Mich. Arb., 3. p. 71.; Du Ham. Arb. Nouv. 2. p. 219.; Dec. Prod., 1. p. 80.; Dou's Mill., 1. p. 82.

Synonymes. Laurel-leaved Magnolia, the large-flowered evergreen Magnolia, the Laurel Bay, big Laurel, the large Magnolia; Laurier tulipier, Fr., in Louisiana; Magnolie à grandes Fleurs, Fr.; grossblumiger Magnolie, or Bieberbaum (Beaver-wood Tree), Ger.

Engravings. Mill. Ic., 2. t. 172.; Mich. Arb., 3. p. 71. t. 1.; Du Ham. Arb., 2. p. 219. t. 65.; E. of Pl., 7904.; and our Pl. I. in Vol. II.

Spec. Char. Evergreen. Leaves oval-oblong, coriaceous, upper surface shining, under surface rusty. Flowers erect, 9—12 petals, expanding. (*Don's Mill.*, i. 82.) North America. Flowers white. June to Sept. 1737.

Varieties. The aboriginal varieties, that is, those which have been found in a wild state in North America, are few, or perhaps only the *M. g. obovata*; because it is uncertain whether the original Exmouth magnolia, which has been so extensively propagated, was an imported plant, or one raised from seed on the spot. The principal varieties cultivated in the London and Paris nurseries are the following:—

† *M. g. 2 obovata* Ait. *The obovate-leaved large-flowered Magnolia.*—Leaves obovate-oblong. Flowers expanded. (*Hort. Kew.*, vol. iii. p. 329.) This seems to be the only variety found in a wild state. In British gardens it is a magnificent plant, the broad ends of its leaves forming a conspicuous feature, and distinguishing it readily from the original species, the leaves of which are pointed; but it does not flower freely.

† *M. g. 3 rotundifolia* Swt. *The round-leaved large-flowered Magnolia.* (*Swt. Hort. Brit.*, p. 14.)—Leaves roundish. Not a very distinct or handsome variety, and not a free flowerer.

† *M. g. 4 exoniensis* Hort. *The Exmouth large-flowered Magnolia.*—The leaves are oblong-elliptical, generally rusty underneath. Flowers somewhat contracted. *M. g. stricta* in some gardens. Figured in Lodd. *Bot. Cab.*, 1814, and in our Second Volume. This is the most distinct of all the varieties of the species, and, on account of its flowering early and freely, the one best deserving of general culture. It forms a tall, fastigate, elegant bush, or tree, and has attained the height of 30 ft., as a standard, at various places in the south of England.

† *M. g. 5 ferruginæa* Sims. *The rusty-under-surfaced-leaved large-flowered Magnolia;* or the rusty-leaved evergreen Magnolia.—This differs from the preceding in having rather broader leaves and larger flowers, and in forming a broader and more compact tree or bush.

† *M. g. 6 lanceolata* Ait. *Bot. Mag.* t. 1952. *The lanceolate-leaved large-flowered Magnolia.*—This differs from the preceding varieties, in not having the leaves rusty underneath; in not being quite so fastigate in general form as the former, nor so broad and bushy as the latter.

† *M. g. 7 elliptica* Ait. *The elliptic-leaved large-flowered Magnolia.*—Leaves oblong-elliptical. Flowers contracted as in the three preceding varieties, from which it differs only in the form of the leaves.

† *M. g. 8 angustifolia* Hort. *The narrow-leaved large-flowered Magnolia.*—Leaves lanceolate, pointed at both extremities, wavy. A very distinct variety, introduced from Paris about 1825, which has not yet flowered in England. The best specimens of this variety are in the London Horticultural Society's garden; in the arboretum of Messrs. Loddiges; and in Lee's Nursery.

† *M. g. 8 præcox* Hort. *The early-flowering large-flowered Magnolia.*—Leaves oval-oblong. Flowers fully expanded. This is an early

variety, introduced from Paris about (?) 1830; and there is a plant of it in Lee's Nursery, and another in the garden of the London Horticultural Society, named *M. g. præcox André*. The flowers are as large as those of any of the varieties, and they are produced from the end of May till the approach of winter.

Other Varieties. In consequence of the great demand for this species in the nurseries, many slight variations have been noticed by cultivators, and named as distinct. In the garden of the London Horticultural Society are the following names:—*M. g. vèra*, *M. g. latifolia*, *M. g. exoniensis var.*, *M. g. rubiginosa*, and some others.

Selection of Varieties. *M. g. obovata* deserves the preference for the magnificence of its foliage; and *M. g. exoniensis*, because it flowers early and freely; and because, from the fastigate form of the tree, it is less liable to be injured by a heavy fall of snow; it seems also to grow faster than any of the other varieties. Where the tree is to be trained against a wall, *M. g. præcox* deserves the preference on account of the largeness of its flowers, and because they appear early, and continue during the whole summer. *M. g. angustifolia* deserves culture on account of its foliage, which is quite distinct from that of all the other varieties. The species sold in the nurseries as the common broad-leaved *Magnolia grandiflora* is frequently raised from American, French, or Italian, seeds; and, hence, the plants, though they grow freely, do not flower for 20 or 30 years after being planted out. For this reason, when it is desired to have plants of the *Magnolia grandiflora* which will flower early, those plants which have been raised by layers from flowering trees ought to have the preference; or the Exmouth variety should be made choice of, because it is always raised from layers.

Description. In its native country, the *M. grandiflora* is a tree varying from 60 ft. to 100 ft., or upwards, in height. According to Bartram, its head forms a perfect cone, placed on a straight clean trunk, resembling a beautiful column; and, from its dark green foliage, "silvered over with milk-white flowers," it is seen at a great distance. In Europe, except in some situations in Spain and Italy, and a few in the south of England, the *M. grandiflora* is chiefly to be considered as a wall tree. There are standards of it, in the neighbourhood of London, of 20 or 30 years' growth; but these are not common; and, on account of the snow breaking down their branches, they require protection during winter. The leaves vary from 9 in. to 1 ft. in length, and from 3 in. to 4 in. or more in breadth; they are always smooth and shining, and perfectly entire on the edges. They bear a strong resemblance to those of the common laurel (*Cerasus Laurocerasus*) both in form and colour; and to those of the orange tree in colour and glossiness, but not in size. In most of the varieties, they are of a rusty brown underneath; and one takes its name from this appearance, though it varies in intensity so much with soil and situation, as to seem rather a variation, than a variety which may be continued by propagation. In America, the flowers appear in May; in England, seldom before June; and they continue in some varieties till they are destroyed by frost. The flowers are produced on the summits of the last year's shoots, and are from 6 in. to 8 in., or even 10 in., across. It is remarked by Collinson, of the flowers of this species of *Magnolia*, that, unlike those of all the other species (unless we except *M. glauca*, when it is planted in moist situations), they are produced throughout the whole season; whereas those of all the other species are produced comparatively at once, and last only a short time. The odour of the flowers is exceedingly sweet, and overpowering to some when near, though it is agreeable at a distance. The seed of the species has in only one or two cases been ripened in England; but it ripens occasionally in Italy and in the neighbourhood of Paris (*Boupl. Malmaison*, p. 54.); and young plants are frequently raised from seed brought from North America.

Geography. Found in the forests of America, from North Carolina to Louisiana. Of all the trees of North America, east of the Mississippi, it is observed by Michaux, and in the *Sylva Americana* of Brown, the big laurel

is the most remarkable for the majesty of its form, the magnificence of its foliage, and the beauty of its flowers. It is first seen in the lower part of North Carolina, near the river Neuse, in the latitude of 35°; proceeding from this point, it is found in the maritime parts of the southern states and of the Floridas, and as far up the Mississippi as Natchez, 300 miles above New Orleans, which embraces an extent of 2000 miles of territory. The French of Louisiana call it *lawrier tulipier*. It grows only in cool and shady places, where the soil, composed of brown mould, is loose, deep, and fertile. These tracts lie contiguous to the great swamps, which are found on the borders of the rivers, and in the midst of the pine barrens, or form themselves a part of these swamps; but they are never seen in the long and narrow marshes called branch swamps, which traverse the barrens in every direction, and in which the miry soil is shallow, with a bed of white quartzose sand beneath. (*Syl. Amer.*, p. 211.)

History. The precise date of the introduction of *M. grandiflora* into Britain is uncertain. In the *Hortus Kewensis* it is stated, on the authority of Catesby, that it was cultivated before 1737 by Sir John Colliton, at Exeter; and, as far as is known, the tree there, which is the parent of all those varieties bearing the name of *M. g. exoniensis*, was the first which was raised or planted in England, and, in all probability, in Europe. This tree, a notice of which will be found in the *Gardener's Magazine*, vol. xi. p. 70., was cut down, through mistake, about the year 1794. It had previously been much disfigured from the great number of layers that had been taken from it; and, though the trunk was 18 in. in diameter, its height was not more than 5 ft. It had been surrounded by a scaffolding for many years, on which tubs were placed to receive the branches laid down for propagation. The tree seems to have been rented by different gardeners, who at first sold the layers at five guineas each; but the price gradually fell to half a guinea. From the source of supply being in this part of England, it will readily be conceived that the largest specimens are in Devonshire. None of these, however, of which we have been able to obtain dimensions, exceed 30 ft. in height. Collinson mentions that on returning to Goodwood, after nine years' absence, he found two plants of *Magnolia grandiflora* in flower (see p. 55.); and that he had a plant, raised by himself from seed, which flowered for the first time in 1760, when twenty years old.

The history of the introduction of the *M. grandiflora* into France is thus given by M. Merlet de la Boulaye, professor of botany at Angers, in the new edition of Du Hamel:—"There is at Maillaudière, distant about 5 miles from the town of Nantes, a fine magnolia, which was brought from the banks of the Mississippi in 1732, and planted in a poor soil. It grew there more than thirty years without any care having been taken of it; as the marine officer who brought it died soon after he had planted it, and his heirs did not trouble themselves about a tree which had as yet produced them nothing, and appeared to them merely a variety of the *Cérasus Laurocérasus*. M. Bonami, a physician of Nantes, and professor of botany there, who published a *Flora* of this district, recognised, in 1758, this beautiful tree to be the *Magnolia grandiflora* of Linnæus; and, at the meeting of the states of Bretagne, which was held at Nantes in September, 1760, he presented to the Princess of Rohan-Chabot a fine branch of this magnolia in flower, which became a subject of conversation and interest to all there assembled. Louis XV. possessed several small plants of the *Magnolia grandiflora* in his garden of the Petit Trianon, but they did not thrive; and, having heard of a magnolia 35 or 40 ft. high, which every year was covered with fine flowers of a delicious perfume, he sent two of his gardeners to ascertain if it was possible to transport this fine tree to Versailles; and, above all, should they do so, if it would be certain to grow. They saw the tree; and, being of opinion that it would not survive removal, it was suffered to remain in its place. This magnolia was, at that time, from 35 ft. to 40 ft. high; but, during the troubles of the civil war of La Vendée, it was mutilated, and lost most of its branches. Afterwards, the burning of the house near which it was planted having damaged its fine head, it was treated as an orange tree injured by the frost; that is to say,

the branches were cut off close to the trunk. It shot out vigorously; but the young shoots, not having had time to ripen, were destroyed by the frost. Notwithstanding this check, it again recovered, and afterwards became a fine tree, between 25 ft. and 30 ft. high, with a large, well proportioned head, and a trunk 4 ft. in circumference, the lower branches sweeping the ground; and the whole tree producing annually from 350 to 400 large, elegant, and sweet-scented flowers. The seeds, however, never arrive at perfect maturity; although the fruit attains its full size, and remains upon the tree till the following spring." (*Nouv. DuR.*, i. p. 220.) This tree, as we have noticed in p. 138., still exists, and is now upwards of 30 ft. high, and 100 years of age.

M. grandiflora, soon after its introduction into France and England, would doubtless find its way into the botanic gardens of Germany. In Italy, as already noticed in p. 169., the first planted trees were in the botanic garden at Padua, where, in 90 years, they have attained the height of 60 ft., with trunks 4 ft. in diameter; and in the botanic garden at Pisa, as we are informed in the *Nouveau Du Hamel*, it produced perfect seeds, from which plants were raised by M. Marnier, on his estate at Rois, near Besançon. The tree has been introduced into the botanic gardens of Spain; also, it is believed, into those of South America and India; and, as stated in p. 176., into the gardens of China.

Properties and Uses. The medical properties of the genus have been already mentioned. In Europe, it can only be considered in the light of an ornamental tree, or rather, perhaps, shrub; as, generally speaking, it can only be cultivated with success when trained against a wall.

Soil and Situation. In Europe, a deep sandy loam, dry at bottom, and enriched with vegetable mould or heath soil, seems to suit all the varieties of this species. The situation in the colder parts of Europe may be exposed to the direct influence of the mid-day sun; but, in the south of France, and in Italy, the tree always thrives best when in the shade of other trees; and in these countries, also, it requires a moister soil than in England. Where the tree is to be treated as a standard, the situation should be sheltered from the points from which the highest winds are expected, but it should be open to the south or south-east, to admit abundance of light and warmth. In general, where the fig tree will grow as a standard, and survive the winter without protection, there the *Magnolia grandiflora* may be planted, and treated as a standard also. The best situations are, the south-east margin of a shrubbery or wood, a sheltered place on a lawn, or an open glade in a plantation. Perhaps the finest situation for displaying the flowers of this tree, as a standard, would be a sloping bank of sandy soil facing the south-east. Here it might be mixed with a few of the deciduous magnolias, and particularly with *M. conspicua* and *Soulangeana*, which, as they flower before their leaves come out, would be set off to great advantage by the evergreen leaves of *M. grandiflora*. When this species is to be trained against a wall, any aspect may be chosen, except, perhaps, the north-east. To display the flowers to the greatest advantage, to a spectator walking in a direction nearly parallel to the wall, the ground plan of the latter should be curvilinear, by which means a direct or front view of a considerable portion would be brought before the spectator. In general, a segment of a circle will be sufficient for a short wall; and two or more segments, forming an ogee, or a serpentine line, for a longer wall. In a very exposed situation, a magnolia wall, 20 ft. or 30 ft. high, might form three parts of a round tower, open at top; the lateral opening facing the south, and the trees planted inside the tower. All the trees might easily be protected by throwing a slight roof of boards over the tower during winter. If the tower were not more than 20 ft. or 30 ft. in diameter, the walls, if built of brick and cement, need not be thicker than 9 in., even if made 30 ft. high. Magnolias might be grown in the inside of such a tower, and camellias on the outside. The wall may be of any height, from 10 ft. to 20 ft., or even 50 ft. A wall covered with evergreen magnolias, interspersed with a few plants of *M. conspicua*, forms a beautiful medium of connexion between a conservatory and a flower-garden or shrubbery; or it forms an admirable northern boundary to a winter


walk, or a winter flower-garden. The finest magnolia wall in England is that at White Knights, near Reading; it is 145 ft. long, and 24 ft. high. The upper part of it is formed of trellis-work, which projects with a curve, the tangent to which forms an angle of, perhaps, 45° with the face of the wall. On this trellis the upper branches of the trees terminate, and, by their projection, protect all those beneath them from perpendicular rains or snows. Such protection, however, is altogether unnecessary, as the magnolias against walls, in Messrs. Loddiges's arboretum, and in various nurseries and gardens about London, abundantly testify. Indeed, it cannot fail to be ultimately injurious, not only by keeping off perpendicular rains, but by excluding the direct influence of the sun's rays from the upper part of the tree. As this species of magnolia does not flower till June or July, its blossoms are in no danger of being injured by frost; and, therefore, it requires no projecting coping, or covering of any sort, during winter, at least about London. Where danger is anticipated from severe frost, attention should be directed to protecting the roots, and especially the collar and the stem, for 2 ft. or 3 ft. above the ground.

Propagation, in the London nurseries, is generally effected by forming stools either in warm situations in the open air, to be protected during winter, or in cold-pits. The shoots are laid down in autumn, and require two years to become sufficiently rooted for separation; they are then potted, and kept in pits or under glass during winter, and set in the open air, in a shady place, during summer, till wanted for final planting. *M. grandiflora* is also occasionally raised from American seeds; but, as plants so originated are much longer in coming into flower than plants raised from layers (as we have before observed, p. 262.), they are not in demand.

Choice of Plants. In no case whatever would we recommend purchasing any species of magnolia not grown in a pot; because plants so grown may be sent to any distance without injury to the roots, which are few and succulent, and easily damaged by exposure to the air and light.

Planting, Culture, and Management. In planting, the ball should be carefully broken by the hand, and the roots spread out in every direction, and covered with heath mould, or a mixture of leaf mould and sandy loam. The soil ought to be made firm to the fibrous roots, not by treading, but by abundant watering, and, if the plant be large, by fixing with water; that is, while the earth is being carefully put about the roots by one man, another should pour water over it from a pot held 6 ft. or 8 ft. above it, so that the weight of the water may wash the soil into every crevice formed by the roots, and consolidate it there. Shading will be advisable for some weeks, or even months, after planting: to a standard, this may be given by placing a cone of wicker-work over the tree by day, and taking it off at night; or by sticking a few spruce fir or other twiggy branches in the soil round it, or, at least, on the south side. Against a wall or trellis, it may be sheltered by an old net during day, which is to be taken off at night, or by any other convenient means. If the magnolia be intended to form a handsome tree as a standard, it should not only have a sufficient depth of suitable soil, and a dry sheltered situation open to the south, but it should be pruned to a single stem for at least 3 ft. or 4 ft. from the ground, to direct the growth to the head. If the plant does not grow freely after it has been three or four years planted, it ought to be bent down to the ground, and kept in that position till it throws up one strong shoot from the collar. The old stem should then be cut away, leaving only the new shoot; and this shoot, which will probably extend to 3 ft. or 4 ft. the first season, will soon form a handsome tree. The greater part of the magnolias which are planted as standards, as far as we have been able to observe, are treated in such a manner that they can never be expected to become any thing else than mere bushes. The soil is not properly prepared; or, if prepared, a sufficient quantity of it is not brought together: because, to admit of this species growing to a tree, the subsoil ought to be prepared by art as well as the soil. The plants, too, are generally turned out of their pots without breaking the ball, and spreading the roots; and it has been ascertained, that the *Magnolia grandiflora*, and various other exotic trees, when

treated in this manner, will remain for several years before the roots strike into the adjoining soil. If the Exmouth variety of this species be made choice of, layers will produce flowers in a year or two after being separated from the parent plant, if kept in pots; but, when they are planted out, and grow freely, so as to make shoots 2 ft. or 3 ft. every season, they will probably not flower for three or four years. Whether the tree be against a wall or trellis, or treated as a standard, all the pruning it will require, after it has begun to grow freely, will be, to cut out the stumps from which the flowers or the strobiles have dropped off, and any dead or decaying wood, and any branches which cross and rub on each other. For a few years after being planted as a standard, it may be advisable to form a small cone of thatch, litter, leaves, or spruce fir branches, round the stem, as practised by M. Boursault in Paris, and exhibited in fig. 33. M. Boursault found that, by this kind of protection, he could grow the magnolia, and various other exotics, as standards, to a size which had never before been seen in Paris (See *Gard. Mag.*, ii. p. 63.) Magnolias against a wall require very little protection, even when young; and this can easily be given by mulching the ground at the roots, and covering their branches with a mat, or with the fronds of the spruce fir.



Statistics. Magnolia grandiflora in the Environs of London. At Syon, 50 years planted, 25 ft. high; at Chiswick, 30 years planted, 15 ft. high; in a garden at Isleworth, 20 ft. high; at Fulham palace, *M. g. exoniensis* 8 years planted, 15 ft. high: these are all standards. There are numerous instances of this species, or its different varieties, planted against walls in the neighbourhood of London, attaining the height of 20 ft., reaching above the wall, and extending 15 ft. or 20 ft. on each side of the main stem. Among the most remarkable may be cited, the magnolias in the botanic garden at Kew, those at Purser's Cross, and at Harringay; at which last place there is one, 20 years planted, which is 20 ft. high.

Magnolia grandiflora South of London. As standards, the largest are at Powderham Castle, and at Coombe, near Plymouth; at both places upwards of 30 ft. high: at Saltram, 60 years planted, 25 ft. high: at Killerton, 18 years planted, the Exmouth variety has attained the height of 23 ft., and flowers nine months in the year. At Eastwell Park, in Kent, 6 trees of *M. g. obovata*, 20 years planted, have attained the height of 25 ft. Examples of *Magnolia grandiflora* against a wall, growing vigorously, and flowering freely, might be given by hundreds. The most remarkable are those at White Knights, see p. 217, and p. 265. The wall was planted in 1800, with twenty-two plants, which cost six guineas each. They were placed in a prepared border, 12 ft. wide, and 6 ft. deep, the soil being a mixture of sand, vegetable mould, and loam; and the subsoil a retentive loam. The trunks of the trees, in 1835, were from 5 in. to 7 in. in diameter; and the plants produce flowers every year, from the beginning of June till they are checked by frost. At Sandown Place, in Surrey, there is a *Magnolia grandiflora* trained against a house, 40 ft. high; at Farnham Castle, there is one against a wall, 20 ft. high; at Leigh Court, in Somersetshire, there is one against a wall, 20 ft. high; at Cowdray, in Sussex, one 24 ft. high; and at Bowood, in Wiltshire, one 25 ft. high.

Magnolia grandiflora North of London. In Hertfordshire, at Hatfield, against a wall, there is a tree 20 ft. high; and, at Wormleybury, one 30 years planted, 24 ft. high; at Golden Grove, in Pembroke-shire, one 12 ft. high, as a standard; at Whitley Abbey, in Warwickshire, there is one 17 ft. high, as a standard; and in Worcestershire, at Croome, against a wall, one 35 ft. high.

Magnolia grandiflora in Scotland. In the Edinburgh Botanic Garden, 12 ft. high, against a wall; at Dalkeith Palace, 10 ft. high, against a wall; in the Glasgow Botanic Garden, only a few feet high, and requiring protection during the winter. At Thainston, and other places in Aberdeenshire, from 6 to 10 ft. high.

Magnolia grandiflora in Ireland. In the Collingswood Nursery, near Dublin, 20 years planted, and 17 ft. high; against a wall, flowering freely every year. In the Trinity College Botanic Garden, 22 years planted, and 10 ft. high. In Ireland generally, the *Magnolia grandiflora* grows much better than in Scotland, or in the north of England; but the dimensions which have been sent to us are all of young trees; for it has not been long the custom in Ireland to plant any trees against walls, except those bearing fruit.

Magnolia grandiflora in Foreign Countries. In France, the best collection of varieties of this species was formerly at M. Boursault's (*Gard. Mag.*, ii. p. 63); and is now at Admiral Tchitchagoff's, at Scéaux, where the highest standard plants, about 15 years planted, are 20 ft. high. The largest trees in France appear to be at Maillardière, near Nantes, where, besides the parent tree already mentioned (p. 263), there are others, varying from 30 ft. to 45 ft. in height, which have been planted from 50 to 80 years. In the Botanic Garden at Toulon there is a tree, 18 years planted, which is 20 ft. high. In Belgium and Holland, the *M. grandiflora* is trained against a wall, and protected during winter, or treated as a conservatory plant, as it is in the greater part of Germany, Denmark, Sweden, and Russia, except in the Crimea, where, as we have seen (p. 179), there are some large specimens, as standards, in the open air. In Italy, there are a greater number of fine specimens of this tree than there are in any other country in Europe; as may be seen by referring to p. 168, and p. 169. The highest trees appear to be those in the botanic garden at Padua, and in the English garden at Caserta, which have attained the height of 60 ft. The tree of this species in the botanic garden at Pisa, as we have seen (p. 169.), ripened seeds many years ago.

Commercial Statistics. Price, in the London nurseries, of young plants in pots, *M. grandiflora*, 3s. 6d.; *M. g. obovata*, or *obtusifolia*, 7s. 6d.; *M. g. exoniensis*, 5s.; at Bollwyller, where this species is a green-house plant, from 3 francs to 25 francs; and in New York, 1 dollar, and the seeds 9 dollars per quart.

‡ 2. *M. GLAÚCA* L. The glaucous-leaved Magnolia.

Identification.—Lin. Sp., 2. p. 755; Willd. Arb., p. 230; Dec. Prod., 1. p. 80; Don's Mill., 1. p. 82. *Synonymes.* *M. fragrans* *Saltb.*; Swamp Sassafras, Beaver-wood, white Bay, small Magnolia, Swamp Magnolia; Magnolie glauque, Arbre de Castor, *Fr.*; grauer Bieberbaum, *Ger.* *Derivations.* It is named Swamp Sassafras on account of its growing in boggy places, and resembling in qualities the *Laúrus Sassafras*; and Beaver-wood, because the root is eaten as a great dainty by the beavers, and these animals are caught by means of it. It also grows in the swamps, which they inhabit; and Michaux tells us that it is felled by them for constructing their dens and houses, in preference to any other tree, on account of the softness of the wood. *Engravings.* Lodd. Bot. Cab., t. 215; Sims, Bot. Mag., 2164; Nouv. Duh., 2. p. 223; E. of Pl., 7905; and our plate of this species in Vol. II.

Spec. Char. Almost deciduous. Leaves elliptical, obtuse, under surface glaucous. Flower 9—12-petaled, contracted. Petals ovate, concave. (*Dou's Mill.*, i. p. 82.) North America. Flowers white, scented. June to September. 1688. Height, in England, 20 ft.

Varieties. The only aboriginal varieties, if varieties they can be called (for they appear to be only variations), are, that which retains its leaves during the greater part of the year, which appears to depend upon the moisture of the soil in which the plant grows, and which is called, in the nurseries, *M. gláuca sempervirens*; that which assumes more of the tree character, and is called *M. gláuca arborea*; and a third, noticed by Pursh, which has the under surface of the leaves somewhat silvery, and is accordingly named *M. g. argentea*.

‡ *M. gláuca* 2 *Thompsoniana* *Thomp.* *Thompson's glaucous-leaved Magnolia.* *M. gláuca* var. *a* major (*Bot. Mag.*, new edit. p. 36.)—This is a supposed hybrid between *M. gláuca* and *M. tripétala*; though it may possibly be only a long-leaved arboreous variety of *M. gláuca*. It was noticed in a pot of seedlings by Mr. Thompson, in his nursery at Mile End, and by him kept distinct, and propagated under the above name. We should say that it was only a variety or race of the aboriginal species, enlarged in all its parts.

‡ *M. gláuca* 3 *longifolia* *Hort.*, the long-leaved glaucous *Magnolia*, is a variety produced, as it is supposed, between *M. tripétala* and *M. gláuca*. It was originated in Belgium, and imported into this country by Mr. Knight of the Exotic Nursery, King's Road, Chelsea. It is in appearance intermediate between the two species; and seems to correspond with the *M. longifolia* of Pursh. The leaves are elliptical, acute at both ends, longer than those of *M. gláuca Thompsoniana*, and resembling those of *M. tripétala*, but thicker, smaller, and glaucous underneath. It is a very handsome tree, and, supposing it to be that described by Pursh, it is found in Florida and Georgia. The flowers are sweet-scented, and resemble those of *M. gláuca Thompsoniana*, but are smaller.

Other Varieties. *M. gláuca Gordoniana*, and *M. gláuca Burchelliana* are variations or varieties, the names of which are found in some nurserymen's catalogues; the former having been raised by Mr. Gordon, at Mile End, and the other by Mr. Burchell, at Fulham: but they do not appear to us at all distinct. There are also semi-double and double-flowered varieties in some nurserymen's catalogues, but the only distinct varieties are *M. gláuca Thompsoniana*, and *M. gláuca longifolia*.

Description. A low tree, nearly evergreen, with slender stem, covered with a smooth whitish bark. The wood is white and spongy. The leaves are smooth, of a bluish green on their upper surface, and whitish or glaucous and a little hairy underneath. The flowers are produced in May or June, at the extremity of the last year's shoots. They have six concave white petals, and

an agreeable odour. The spike of fruits is an inch or more in length, conical, an inch in diameter in the widest part, and of a reddish brown colour when ripe. This tree begins to flower, in the United States, in May, June, and July; when the perfume of the flowers may be perceived at a considerable distance. A few of these flowers, shut up in a room over night, communicate to the air a heavy and almost insupportable odour. (*Bigelow's Med. Bot.*, vol. ii. p. 68.) When the plant is in a soil supplied with moisture during the summer, it continues to produce flowers till the autumn, and retains part of its leaves all the winter: in dry situations the leaves drop off. Seeds are frequently ripened in England: they are of a bright scarlet, and they hang down by slender white threads, as in all the other American species. The young shoots are from 1 ft. to 18 in. in length, and the plant, in ordinary circumstances, will attain the height of 12 ft. in ten years: when full-grown, it is seldom, either in its native country or in England, more than 18 ft. or 20 ft. high; which height in England it arrives at in twenty years.

Geography. Found in low situations near the sea in North America, from Massachusetts to Louisiana; more especially in New Jersey and Carolina, where it seldom grows to above 12 ft. high. It has the most extensive range, especially near the sea, of any of the genus. Its most northern boundary appears to be in a sheltered swamp in Manchester, Cape Ann, about thirty miles north of Boston. It here attains to but small size, and is frequently killed to the ground by severe winters. It is common in the middle and southern states; and Michaux informs us that it is one of the most abundant trees in the morasses of Florida and Lower Louisiana. According to this author, however, it is not usually met with far in the interior, or to the west of the mountains. Its common names are various, and change with almost every district. It is naturally a tenant of deep boggy swamps, and is somewhat irregular in its growth. It acquires more symmetry of form when cultivated in an upland soil, although its transplantation is difficult. To insure its successful cultivation in a dry soil, the tree should be raised from the seed. (*Bigelow's Med. Bot.*, vol. ii. p. 68.)

History. This species was introduced into England by Banister, who sent it to Bishop Compton, at Fulham, in 1688. It was soon afterwards generally propagated by American seeds, and became known throughout Europe many years before any of the other species.

Soil and Situation. The soil ought to be a deep sand, or a sandy peat, kept moist, more especially in summer. The situation should be sheltered and shaded by larger trees, but not overtopped by them.

Properties and Uses. The bark is aromatic and pungent, apparently more so than that of most of the other species. When distilled, it has a peculiar flavour, and an empyreumatic smell. In a dry state it affords a little resin. The aroma is volatile, and probably an essential oil or variety of camphor. The bark, seeds, and canes, are employed in tincture, with very good success, in chronic rheumatism. The inhabitants of the marshy countries of America have used the bark, like that of the cinchona, in the case of intermittent and remittent fevers. The inhabitants of the countries where the plant is indigenous cure coughs and other pectoral diseases by putting the fruit into brandy, and administering the decoction every morning. The wood is employed for making joiners' planes. The flowers, in a dried state, may be used in drawingrooms for *pot pourri*, as a substitute for those of the lily of the valley. In Europe, the plant can only be considered as ornamental.

Propagation and Culture. Plants are generally originated from seeds imported from America; but *M. glauca Thompsoniana*, and the other varieties, are propagated by layers, which require two years to root properly; or by inarching. The seeds should be sown in pots of bog earth about the beginning of March, and placed in gentle heat under glass. In a year they will be fit to transplant into small pots; and every year they should be shifted into others of a larger size, till wanted for final planting out.

Statistics. *Magnolia glauca* in Great Britain. This species is by no means uncommon, but always as a standard, and, when in a thriving state, in moist peat soil. The largest tree in the neighbourhood of London is that at Syon, figured in our Second Volume. There are in Thompson's Nursery, at Mile End, various specimens from 17 ft. to 20 ft. in height; and one of the var. *Thompsoniana* still higher. There is one at Kew, 30 years planted, which is 20 ft. high; one of the same age and height at White Knights, and another in the Killerton Nursery; one at Cobham Hall, Kent, 25 years planted, and 25 ft. high. At Woburn Farm, Chertsey, there was formerly a row of trees of this species 20 ft. high, and nearly a century old, which were cut down when the new house was built. (*J. M.*, in *Genl. Mag.*, new series, vol. iii. p. 226.) At Alton Towers, in Staffordshire, the tree has attained the height of 12 ft. in 10 years; and at Croome, in Worcestershire, 15 ft. in 25 years. At Cownan House, in Cromarty, in lat. 55° 35'', and 161 ft. above the level of the sea, young plants were growing freely in 1835. At Oriel Temple, near Dublin, trees 10 years planted have attained the height of 11 ft.

Magnolia glauca in foreign countries. In France this species is not very abundant, but with the great heat of the summers, and the general dryness of the air; but, in some shaded moist situations at Versailles and the Petit Trianon, it has attained the height of 15 ft. There are trees of most of the varieties, of from 10 ft. to 12 ft. in height, at Scéaux and at Fromont, and small plants in many of the botanic gardens both of France and Germany. In Belgium there are trees upwards of 15 ft. high in various private gardens, and of a smaller size in the botanic garden at Ghent, and in the grounds of the palace of Lâcken. In the north of Germany, and in Sweden and Russia, it is a green-house plant; and, though it is to be found in the south of Europe in most of the botanic gardens and best private collections, yet, owing to the heat and dryness of the air, it does not thrive in these countries. *M. glauca*, and all its varieties, will be found at Monza.

Commercial Statistics. Plants for sale, whether seedlings or layers, are generally kept in pots. The price, in the London nurseries, is 2s. 6d. each; at Bollwyller, 10 francs; and in New York, 12 dollars per hundred, or 1½ cents each, and 2 dollars 75 cents per quart of seed. In London the seed is sold by Charlwood for 1s. 6d. an ounce.

‡ 3. MAGNOLIA TRIPE'TALA L. The three-petaled Magnolia.

Identification. Lin. Sp., 2. p. 756.; Willd. Baum., 231.; Pursh, 2. p. 381.; Michx., 3. p. 90.

Synonymes. *M. umbrellâ Lam.*, *Nouv. Duh.*, *Dcc. Prod.*, *Don's Mill.*; *M. frondôsa Salisb.*; the Umbrella Tree; Umbrella Magnolia; Elkwood; Magnolie Parasol and Arbre Parasol, *Fr.*; dreiblättriger Bieberbaum, dreiblättrige Magnolie, *Ger.*

Derivations. This species is called the Umbrella Tree, according to Michaux, because its leaves, which are thin, oval, entire, and acuminate at both extremities, 18 in. or 20 in. long, and 7 in. or 8 in. broad, are often disposed in rays at the extremity of vigorous shoots; and these display a surface of 2½ ft. in diameter, in the form of an umbrella. The tree is called Elkwood in the mountains of Virginia, probably from the resemblance which the points of the shoots bear to the horns of the elk. The French names merely signify Umbrella Tree, and the German ones the three-petaled Beaver Tree, or Magnolia.

Engravings. Michx. Arb., 3. t. 5.; Lodd. Bot. Cab., t. 418.; Otto., t. 18.; *Nouv. Duh.*; and the plate of this species in our Vol. II.

Spec. Char. Deciduous. Leaves lanceolate, spreading, adult ones smooth, younger ones pubescent underneath. Petals 9—12, exterior ones pendent. (*Don's Mill.*, i. p. 83.) North America. Flowers white. In May and June. 1752. Height 30 ft.

Description. This tree, both in America and Europe, is remarkable for the largeness of its leaves, and its flowers. The wood is spongy, brittle, with a large pith, soft, porous, and of very little use. The bark upon the trunk is grey, smooth, and polished; and, if cut while green, it exhales a disagreeable odour. In Britain the tree sends up various shoots from the root, to replace the stems, which are seldom of long duration; so that a plant that has stood thirty or forty years in one spot has had its stems several times renewed during that period. In America it seldom exceeds the height of 35 ft.; and, in England, 36 ft. is the greatest height that it has yet attained. The trunk, in both countries, is from 5 in. to 6 in. in diameter. The stem is seldom erect

but generally inclined, branching, and rising from the root in twos or threes. The leaves are 18 in. or 20 in. long, and 7 in. or 8 in. broad. The flowers are 7 in. or 8 in. in diameter, with large white flaccid petals; they are placed on the extremities of the last year's shoots, have a languid luxurious appearance, and a sweet but heavy odour. The fruit, which is conical, is 5 in. or 6 in. long, and about 2 in. in diameter. It ripens in America about the beginning of October; and in England in fine seasons, about the end of the same month. It is of a beautiful rose colour, and contains usually from 50 to 60 seeds, which should be sown immediately after they are gathered, as otherwise they become rancid and lose their vital qualities; though, if enveloped in moist moss or earth, they may be preserved for several months. This species is very hardy, and can withstand the most rigorous winters, when the summer has been sufficiently hot to ripen the wood thoroughly. As it is a short-lived tree, and consequently flowers early, there is not the same objection to raising plants of it from seed, as there is to raising plants in that manner of *M. grandiflora*, which is a long-lived species.

Geography. The umbrella tree, according to Michaux, is first seen in the northern part of the state of New York, and it extends on wooded mountains to Carolina and Georgia, as well as Virginia. Though met with over a great extent of country, it appears only in situations perfectly adapted to its growth, which are always shady, and, where the soil is deep, strong, and fertile. In the lower parts of South Carolina and Georgia, it is found only near the alluvial flats which lie along the banks of the rivers, and there it is accompanied by the *Magnolia grandiflora*, but never by the *Magnolia glauca*, which is confined to situations where, according to Michaux, the soil is black, shallow, and often miry.

History. This species of *Magnolia* was brought to England about 1752, and soon after it passed into France, and was cultivated on the Continent generally. In France and Italy it seeds freely; and even in England, at Deepdene in Surrey, self-sown seeds have produced plants. It may now be considered as the commonest of all the magnolias; because, though in point of beauty it is not so popular as *M. grandiflora* or *M. glauca*, yet, as a peat soil is not essential to it, it is more easily preserved.

Soil and Situation. The soil should be a deep, rich, sandy loam, and the situation sheltered and shaded. A situation exposed to the sun is injurious; and, trained against a wall, the plant suffers extremely. A sheltered glade, in a shrubbery or wood, where it is sufficiently distant from other trees not to be injured by the roots, is the most desirable site.

Propagation and Culture. In the nurseries it is almost always propagated by seeds, but sometimes also by layers. In either case the plants are kept in pots until required for final transplanting.

Statistics. Magnolia tripétala in Great Britain. The largest plants in the neighbourhood of London are at Purser's Cross and at Syon; and they are about 30 ft. high. There is one in the Mile End Nursery 20 ft. high. The largest in England are at Cobham Hall, in Kent, and about 36 ft. high. At Walton House, in Surrey, there are plants 30 ft. high; and at Bowood, in Wiltshire, one 13 ft. high. At Golden Grove, in Pembrokeshire, there is a tree 23 ft. high; and at Croome, in Worcestershire, one 20 ft. high. The tree does not thrive about Edinburgh; and in the north of Scotland it is trained against a wall. In the Perth Nursery there is one, trained against a wall, 16 ft. high; and at Gordon Castle, one 14 ft. high. In Ireland, in the Glasnevin Botanic Garden, there is a standard tree 14 ft. high; and at Pakenham Hall, in the province of Leinster, there is a standard tree which in 10 years has attained the height of 10 ft.

Magnolia tripétala in Foreign Countries. In the Ghent Botanic Garden there is a standard 25 ft. high; and one at Sécaux, near Paris, which has attained the same height in 12 years. At Schwöbber, in Hanover, there is a tree 25 years planted, which is 30 ft. high. The species is in most of the Continental botanic gardens, in France, Belgium, Holland, and the south of Germany, as a

standard in the open air; and in the north of Germany, and in Denmark and Sweden, as a green-house plant. It is also in botanic gardens of the south of Italy, but it does not thrive there. Some of the finest trees of this species in Italy are at Monza.

‡ 4. *M. MACROPHYLLA* *Mich.* The long-leaved Magnolia.

Identification. Mich. Bor. Amer., 1. p. 327; Dec. Prod., 1. p. 80; Hayne Dend., p. 117; Don's Mill., 1. p. 83.

Synonymes. Large-leaved Umbrella Tree, *Amer.*; Magnolia Michauxii Hort.; Magnolier à grandes Feuilles, Magnolier baumier, *Fr.*; grosseblättrige Bieberbaum, *Ger.*

Engravings. Bot. Mag., 2189; E. of PL., 7915; and our plate in Vol. II.

Spec. Char. Deciduous. Leaves very large, oblong-obovate, somewhat panduriform, cordate at the base, under surface whitish, glaucous. Petals 6—9, ovate. (*Don's Mill.*, 1. p. 83.) A tree of the middle size. North America. Flowers white. June and July. 1800. Height 35 ft.

Description. The general appearance of this tree, Michaux observes, greatly resembles that of *M. tripétala*. The terminal arrangement of the leaves is the same, and it is remarkable that the two trees are almost always found together. In point of size, it exceeds the *M. tripétala*, both in its leaves and general height; but it is seldom found higher than 35 ft., which exceeds the height of the other by a sixth part only. The body of the tree is covered with a smooth and very white bark, by which, in the winter, when stripped of its leaves, it is readily distinguished from *M. tripétala*. At this season, also, it may be distinguished by its buds, which are compressed and covered with a soft and silvery down; whereas in *M. tripétala* they are prominent and rounded at the end. The leaves, in its native country, are 35 in. long, and 9 in. or 10 in. broad; and in vigorous plants, in England, they sometimes even exceed these dimensions. They are borne on petioles, short in comparison with the size of the leaves, and are of an oblong oval shape, pointed at the extremity, and cordiform at the base: their colour is light green above, and glaucous beneath. The flowers are white, and larger than those of any other species of magnolia; for, when fully blown, they are sometimes 8 in. or 9 in. in diameter: they are composed of six petals, longer and broader than those of the umbrella tree. Within the flower, near the bottom of the petals, is a purple spot, 7 or 8 lines in diameter. The flowers diffuse a fragrant odour, and their beauty is heightened by the luxuriant foliage which surrounds them. The fruit is about 4 in. long, nearly cylindrical, and of a vivid rose-colour when arrived at maturity. In the arrangement of the carpels and of the seeds, the fruit resemble those of *M. tripétala* and *M. acuminata*; it should be remarked, however, that it is destitute of the appendages visible on that of the last-mentioned species, especially when it is dry. The seeds of the large-leaved umbrella tree require, in order to preserve their power of germination, the same attention as those of the preceding species. (*Michx.*) The stipules, in this species, and the manner in which they envelope the unexpanded leaves, are interesting subjects of observation, more especially when the leaves are emerging from the bud. The stipules are large, and placed mainly upon petioles of the leaves; yet the office of the stipules borne by the petiole of any leaf is not to envelope and protect that leaf, but the leaf next inward to it. The outermost wrappers of the leaf-buds are (as examination will show) stipules upon the rudiments of petioles. Young plants of this species grow very slowly till they are thoroughly established, which will require, in general, two years. The year's shoots may then be from 1 ft. to 2 ft.; so that in ten years a plant may attain the height of 12 ft. or 15 ft. It may be considered a short-lived tree, and, like all such, it comes into flower when young. The largest tree of this species, in England, is 28 ft. 6 in. high.

Geography. This is the rarest of the American species of magnolia. It was generally confounded by the native collectors with the *Magnolia tripétala*, till separated from it by Michaux, by whom it was discovered in 1789, in the mountainous regions of North Carolina, 10 miles south of Lincoln town, and 250 miles from Charleston. Extensive researches made in quest of it, in the

upper part of the southern states, and east of the Alleghanies, have been unsuccessful. In Tennessee it is found sparingly at intervals of 10 or 50 miles. It appears to delight in cool sheltered situations, where the soil is deep and fertile; and, as already observed, it is constantly attended by the *M. tripétala*.

History. It was discovered by the elder Michaux, in 1789, but was not introduced into England till imported by the Messrs. Loddiges in 1800. It has rarely, if ever, been propagated in this country by inarching or layers, and very seldom from seeds; and, hence, the plant is very sparingly distributed. In France, it seems to have been introduced about the same time as in England; and it seems to prosper better in the climate of Paris, as there, in the nursery of M. Godefroy, it has ripened seeds, from which, in 1827, young plants were raised. In Britain, young plants are constantly imported from the New York and Philadelphia nurseries.

Soil and Situation. The most suitable situation for this species is one perfectly sheltered on every side, and slightly shaded from the mid-day sun. The soil should be a deep dry sand; at all events, those trees in England which have attained the largest size, stand in soil of this description; our comparatively moist winters rendering such a soil advantageous, by preventing the excess of moisture from rotting the roots, or damping off the plant when young, at the surface of the ground.

Propagation and Culture. Neither this species nor *M. tripétala* can be readily grafted or inarched on each other, or on any other species, as far as experience has hitherto gone in Britain; probably from the large proportion which the pith bears to the ligneous part in young shoots; nevertheless, according to Bonpland, it has been in one or two cases successfully effected in France. M. Soulange-Bodin, having been unsuccessful in various attempts to inarch *M. macrophýlla* on *M. tripétala*, thought of trying it on *M. auriculata*; but it only lived a very short time, and then died. It will root by layers with great difficulty; and plants so raised, from their want of vigour, will probably not be of long duration. The only mode worthy of general adoption is, to raise it from seed; and, as these are produced in abundance at Fromont, at the nursery of M. Godefroy, at Ville d'Avry, and at other places in France, there is no necessity for having recourse to any other method. If any species of magnolia, the young shoots of which are so abundant in pith, and the entire plant so liable to die down to the ground and shoot up again, as *M. tripétala* and *M. macrophýlla*, is to be grafted at all, the operation ought, as it appears to us, to be performed on the root, which, as in the case of all ligneous plants, is without pith.

Statistics. The largest *Magnolia macrophýlla* in England, is that at Arley Hall, the seat of the Earl of Mount Norris. This fine tree is a standard, 28 ft. 6 in. high, with a trunk 6 in. in diameter at a foot from the ground, and a head 17 ft. in diameter. The next largest standard is that at the Duke of Devonshire's villa at Chiswick, which, in 1835, was 22 ft. high. At Harringay there is a tree against a wall, which, in 1835, was 22 ft. high, and, like that at the Duke of Devonshire's, it flowers abundantly every year. It was planted in 1814, and has never received the slightest protection. In the grounds of a villa at Kensington Gore, adjoining the Brompton Nursery, is a standard tree 18 ft. high, which flowers every year. At White Knights there is a standard tree, 20 ft. high, that has been 30 years planted. At Southill, in Bedfordshire, there is a standard, 22 years planted, which is 12 ft. high. The tree stands in the open air in the neighbourhood of Edinburgh and of Dublin; but there are no remarkable specimens. In France, the largest *M. macrophýlla* is at Fromont, which, in 1835, measured 24 ft., and the branches covered a space of 15 ft. in diameter. It has flowered every year since 1826, and it ripens seeds in October, from which many young plants have been raised. There are plants of this species, of considerable size, at Ville d'Avry, and in several of the other Parisian nurseries. In Germany, the species is a green-house plant, and in the south of Europe it has not yet been generally tried. There are plants of it in the collection at Monza. Plants should always be purchased in pots. The price, in London, of two-years-old seedlings, is 15s.; at Bollwyler, 2; and at New York, 1 dollar.

‡ 5. *M. ACUMINATA* L. The pointed-leaved Magnolia.

Identification. Lin. Sp., 756; Dec. Prod., 1. p. 80.; Willd. Baum., p. 230.; Hayne Dend., p. 117.; Don's Mill., 1. p. 83.

Synonymes. *M. rustica*, and *M. pennsylvanica*, of some; the blue Magnolia, *Eng.*; the Cucumber Tree, *U. S.*; Magnolier acuminé, Magnolier à Feuilles pointées, *Fr.*; zugespitzter Bieberbaum, *Ger.*

Derivations. This species is called the Cucumber Tree, in America, from its fruit resembling a small cucumber. The other names are translations of the botanic one.

Engravings. Mich. Arb., 3. p. 82. t. 3.; Sims, Bot. Mag., 247.; Hayne, t. 17.; E. of Pl., 7913.; and our plate in Vol. II.

Spec. Char. Deciduous. Leaves oval, acuminate, under-surface pubescent. Flowers 6—9-petaled. (*Don's Mill.*, p. 83.) A deciduous tree. North America. Flowers yellowish. May to July. 1736. Height from 30 ft. to 40 ft.

Varieties.

‡ *M. a. 2 Candolli Savi.* *De Candolle's acuminate-leaved Magnolia.*—Leaves ovate, oblong, acute. Flowers greenish. Figured in Savi's *Bibl. Ital.*, p. 224.

‡ *M. a. 3 maxima* Lodd.—Leaves much larger than those of the original species. Introduced by Messrs. Loddiges, and cultivated in different nurseries.

Other Varieties. The *Magnolia acuminata* being frequently raised from seed, and the seedlings varying much in the size of their leaves, and in the presence or absence of pubescence, both on the leaves and wood, it would be easy to select several varieties apparently as distinct as those above mentioned. In the Goldworth Nursery, Woking, Surrey, are some which appear remarkably distinct.

Description. A deciduous tree, in its native country, from 60 ft. to 80 ft in height, with a straight trunk, from 3 ft. to 4 ft. in diameter; numerous branches, and regularly distributed shoots. The leaves are from 6 in. to 7 in. long, and from 3 in. to 4 in. broad, upon old trees, but double that size upon young vigorous-growing plants. Michaux describes them as oval, entire, and very acuminate; but, in the seedlings raised in British nurseries, they are found sometimes ovate, nearly orbiculate, and cordate-acuminate. The flowers, which are 5 in. or 6 in. in diameter, are bluish, and sometimes white, with a tint of yellow. They have but a feeble odour; though, as they are large and numerous, they have a fine effect in the midst of the superb foliage. Plants raised from seeds do not usually produce flowers till they are eight or ten years old, when the tree will probably be from 15 ft. to 20 ft. in height; but plants raised from layers produce flowers in two or three years. The fruit is about 3 in. long, and nearly 1 in. in diameter. It is nearly cylindrical, and often a little larger at the summit than at the base: it is convex on one side, and concave on the other; and, when green, it nearly resembles a young cucumber. The fruit is rose-coloured; and, as in the case of the other species, the seeds, before they drop, remain suspended for some time by long white threads. The wood of this tree is of a fine grain, and of an orange colour.

Geography. The most northerly point at which this tree is found is Niagara, near the Falls, in lat. 43°. It abounds along the whole mountainous tract of the Alleghanies, to their termination in Georgia, over a distance of 900 miles. It is also common on the Cumberland Mountains, which divide the state of Tennessee. The situations peculiarly adapted to its growth, according to Michaux, are the declivities of mountains, narrow valleys, and the banks of torrents, where the atmosphere is constantly moist, and where the soil is deep and fertile. "At the distance of 40 or 50 miles from these mountains, either eastward or westward, the cucumber tree is met with only accidentally upon the steep banks of rivers, where the atmosphere is constantly refreshed by the evaporation from their surface. We may conclude that this tree is a stranger to all the regions north of the river Hudson, and to all the Atlantic parts of the United States, to the distance of 100, 150, and 200 miles from the sea; the nature of the soil, and the extreme heat of the climate in summer, being utterly uncongenial to its growth. It is also

rare in the parts of Kentucky and West Tennessee which are most remote from the mountains, where the face of the country is less uneven." (*Mitchaux.*)

History. *M. acuminata* was first discovered by John Bartram, and was sent by him to Mr. Peter Collinson, in 1736. Being readily propagated by layers, and very hardy, it was soon spread extensively through European gardens; and there are now numerous trees of it in Britain, France, and in the north of Italy, from 40 ft. to 60 ft. in height.

Properties and Uses. The tree being comparatively rare in its native country, its timber is not in general use. Where it can be obtained, it is employed in joinery and cabinet-making; and, from its size and lightness, large trunks are selected for scooping out into canoes. The inhabitants of the Alleghanics gather the fruits about midsummer, when they are half ripe, and steep them in whisky: a glass or two of this liquor, which is extremely bitter, is considered to be a preventive against the autumnal fevers. In Europe, the tree can only be considered as ornamental; though its fruit might no doubt be applied in the same manner as in America.

Soil and Situation. A free, deep, and rather moist soil answers best for this species; but, as it is much hardier than any of the other species in this section, it will grow in almost any soil that is moderately free, and not overcharged with moisture. To attain a large size, it requires a sheltered situation, and a deep rich soil; but it will grow in exposed sites, and even flower there freely.

Propagation and Culture. It is generally propagated in the London nurseries by layers, the plants so produced flowering much sooner than seedlings; but the latter, as they make far more durable plants, should always be preferred when this species is used as a stock to graft or inarch others on. It is so used very generally, not only for *M. auriculata* and *cordata*, but for *M. conspicua* and *Soulangiana*. The plants are, in some nurseries, grown in the free soil; but it is always preferable to rear them in pots; because, in that case, they are not checked by transplanting, and at least a year is gained in their growth.

Statistics. Magnolia acuminata in Great Britain. The largest tree stood in the garden of Lord Petre, at Thorndon Hall in Essex; but it was cut down some years ago, and its exact dimensions we have not been able to ascertain; though we have seen a section of the trunk which exceeded 27 in. in diameter: there is one still standing in the same park, which is 37 ft. high, with a trunk 7 ft. 2 in. in circumference. In Thompson's Nursery at Mile End, and in the arboretum at Kew, there were formerly trees between 30 ft. and 40 ft. high. At Syon there is one 49 ft. high, see our plate in Vol. II.; and at White Knights there are many trees of this species from 20 ft. to 35 ft. high, all planted within the last 30 years. At Cobham Hall, in Kent, there is a tree 17 years planted, which is 30 ft. high; and there is one of the same height at Eastwell Park, in the same county. North of London there are some hundreds of trees of which we have received the dimensions. They grow in various soils, clayey loam, sand, prepared soil, &c.; and, in 10 years, generally acquire the height of 15 ft., and in 20 years, of 30 ft. In Scotland, the tree is usually trained against a wall; but in the neighbourhood of Dublin there is a tree, 15 ft. high, growing as a standard in the Glasnevin Garden, and another, equally high, at Cypress Grove. At Oriel Temple there is a tree 35 years planted, which is 17 ft. high; and one, not quite so old, at Dundalk, 27 ft. high, with a trunk 20 in. in diameter, and a head 85 ft. in circumference. In England these trees flower freely every year, but not quite so much so in Ireland, owing to the wood not ripening so thoroughly.

Magnolia acuminata in Foreign Countries. There are various trees of this species, in the neighbourhood of Paris, from 20 ft. to 30 ft. high; and in Belgium there are a number still higher. In Germany, the largest tree that we have had an account of is at Schwöbber, where it has attained the height of 25 ft. It stands in the open air at Berlin, Dresden, and Vienna, but never acquires a timber-like size as a standard. In the Berlin Botanic Garden it is 25 ft. high against a wall. In Italy, there are trees of this species in the Eng-

lish garden at Caserta, and in most of the botanic gardens; but, as already observed, the deciduous American magnolias do not thrive in the south of Europe, except in particular localities. In North America, there is a tree of this species in Bartram's Botanic Garden, Philadelphia, 80 ft. high, which supplies a great part of the seeds sent yearly to Europe.

Commercial Statistics. The price of plants, about London, is 5s. each, and of seeds 2s. 6d. an ounce; at Bollwyller, from 5 francs to 10 francs each plant; in New York, plants are 25 dollars a hundred, or 30 cents each, and seeds are 9 dollars a quart.

‡ 6. M. (? ACU.) CORDATA *Mr.* The heart-leaved Magnolia.

Identification. Mich. Bor. Amer., 1. p. 328.; Dec. Prod., 1. p. 80.; Hayne Dend., p. 118.; Don's Mill., 1. p. 83.

Synonymes. The heart-leaved Cucumber Tree, *Amer.*; Magnolier à Feuilles en Cœur, *Fr.*; herzblättriger Bieberbaum, *Ger.*

Engravings. Bot. Cab., 474; and our plate in Vol. II.

Spec. Char. Deciduous. Leaves heart-shaped, somewhat ovate or cordate, acute, under surface tomentose, upper surface smooth. Petals 6—9, oblong. (*Don's Mill.*, i. p. 83.) A middle-sized tree. North America. Flowers white and purple, scented. June and July. 1800.

Description. This tree, in its native country, attains the height of 40 ft. or 50 ft., with a trunk 12 in. or 15 in. in diameter, straight, and covered with a rough and deeply furrowed bark. Its leaves, which are borne upon petioles, are from 4 in. to 6 in. in length, and from 3 in. to 5 in. wide, smooth and entire. The flowers, which appear in April, are yellow, with the interior of the petal longitudinally marked with several reddish lines. They are from 3 in. to 4 in. in diameter, and are succeeded by fruit about 3 in. long, and nearly 1 in. in thickness, of a similar form to those of the preceding species. The wood is light and soft, and is used in joinery and cabinet-making, where it can be found; but the tree is not common in America. In Britain, the tree attains the height of 20 ft. or 30 ft., and flowers freely.

Geography. Found on the banks of the river Savannah, in Upper Georgia, and on those of the streams which traverse the back parts of South Carolina. The nearest point to the sea at which the younger Michaux found it, was in the plantation of Goodrest, 12 miles from Augusta, along the sides of Horn Creek. The tree is rare in Upper Georgia, never making its appearance in forests, but only in isolated situations, along the banks of rivers.

History, &c. This tree appears to have been discovered by the elder Michaux. It was brought to England in 1801 by Mr. Lyon; and the original tree, not 15 ft. high, still exists in the nursery of Messrs. Loddiges. This tree agrees in very few particulars with Michaux's description, and, taken together with the various and very opposite appearances assumed by the seedlings of *M. acuminata*, convinces us that *M. cordata* is nothing more than a variety of *M. acuminata*. The soil and situation may be considered the same as in the preceding species; but, as this race or variety seems, in its native country, to inhabit higher and drier localities than *M. acuminata*, it may probably be placed in still more exposed situations than that species in Britain.

Statistics. Though this species is by no means uncommon in British gardens, we are not aware of many large specimens of it. The highest we know of is at Claremont, where it has attained the height of 27 ft. in sandy loam on clay. At Luscombe, in Devonshire, there is a tree 8 years planted, which has attained the height of 14 ft.; and at West Dean, in Sussex, is one 9 years planted, which is 13 ft. high. At High Clere, in a situation upwards of 500 ft. above the level of the sea, a plant 12 years planted is 12 ft. high. In the Perth Nursery, one 8 years planted is 15 ft. high against a wall. There are trees as standards at Oriel Temple, Terenure, Charleville, and various other places in Ireland. At Paris, there are trees at Sécaux, and in most of the nurseries and botanic gardens. Plants, both seedlings and layers, are not infrequent in the nurseries. The price, in London, is from 7s. 6d. to 21s.; at Bollwyller, 6 francs; in New York, 1 dollar.

Y 7. *M. auriculata* Lam. The auricled-leaved Magnolia.

Identification. Willd. Sp., 2. p. 1258.; Dec. Prod., 1. p. 80.; Hayne Dend., 117.; Don's Mill., 1. p. 83.

Synonyms. *M. Fraseri* Walt.; *M. auriculàris* Salisb.; Indian Physic, and long-leaved Cucumber Tree, Amer.; Magnolier auriculé, Fr.; geöhrtet (eared) Bieberbaum, Ger.

Engravings. Bot. Mag., 1206.; E. of Pl. 7916.; and our plate in Vol. II.

Spec. Char. Deciduous. Leaves smooth, under surface somewhat glaucous, spatulately obovate, cordate at the base, with blunt approximate auricles. Sepals 3, spreading. Petals 9, oblong. (*Don's Mill.*, i. p. 83.) A tree of the middle size. North America. Yellowish-white flowers. April and May. 1786.

Description. This tree attains the height of from 40 ft. to 45 ft., with a straight trunk 12 in. or 15 in. in diameter, often undivided for half its length; the branches spread widely, and ramify but sparingly; and this circumstance, Michaux observes, gives the tree a very peculiar air, so that it may readily be known at a distance, even in winter. The leaves are of a light green colour, of a fine texture, 8 in. or 9 in. long, and from 4 in. to 6 in. broad: on young and vigorous trees they are often one third, or even one half, larger. They are smooth on both surfaces, acuminate at the summit, widest near the top, and narrowest towards the bottom. The base is divided into rounded lobes, whence is derived the specific name of *auriculata*. The flowers are 3 in. or 4 in. in diameter, of a milky white, of an agreeable odour, and are situated at the extremity of the young shoots, which are of a purplish red, dotted with white. The fruit is oval, 3 in. or 4 in. long, and, like that of *Magnolia tripetala*, of a beautiful rose colour when ripe. They differ from those of the other species by a little inferiority of size, and by a small appendage which terminates the carpels. Each carpel contains one or two seeds. The wood is soft, spongy, very light, and unfit for use. The bark is grey, and always smooth, even on the oldest trees. When the epidermis is removed, the cellular integument, by contact with the air, instantly changes from white to yellow. The bark has an agreeable aromatic odour, and an infusion of it in some spirituous liquor is employed as an excellent sudorific in rheumatic affections. (*Michaux.*) In England, annual shoots of young plants are from 1 ft. to 2 ft. or more in length; and the height which the tree usually attains in 10 years is from 10 ft. to 15 ft. The highest tree within ten miles of London is at Messrs. Loddiges's, where it has attained the height of 30 ft. The following extract from *Bartram's Travels* will be read with interest. "This exalted peak I named Mount Magnolia, from a new and beautiful species of that celebrated family of flowering trees, which here, at the cascades of Falling Creek, grows in a high degree of perfection. I had, indeed, noticed this curious tree several times before, particularly on the high ridges betwixt Sinica and Keowe, and on ascending the first mountain after leaving Keowe, where I observed it in flower: but here it flourishes and commands our attention. This tree (or perhaps rather shrub) rises 18 ft. to 30 ft. in height. There are usually many stems from a root, or source, which lean a little, or slightly diverge from each other, in this respect imitating the *Magnolia tripetala*; the crooked wreathing branches arising and subdividing from the main stem without order or uniformity; their extremities turn upwards, producing a very large rosaceous, perfectly white, double, or polypetalous, flower, which is of a most fragrant scent. This fine flower sits in the centre of a radius of very large leaves, which are of a singular figure, somewhat lanceolate, but broad towards their extremities, terminating with an acuminate point, and backwards they attenuate, and become very narrow towards their bases, terminating that way with two long narrow ears, or lappels, one on each side of the insertion of the petiole. The leaves have only short footstalks, sitting very near each other, at the extremities of the floriferous branches, from which they spread themselves after a regular order, like the spokes of a wheel; their margins touching, or lightly lapping upon, each other, form an expansive umbrella, superbly crowned or crested with the fragrant flower, representing a white plume. The blossom is succeeded by a

very large crimson cone, or strobile, containing a great number of scarlet berries, which, when ripe, spring from their cells, and are, for a time, suspended by a white silky web or thread. The leaves of those trees which grow in a rich humid soil, when fully expanded and at maturity, are frequently above 2 ft. in length, and 6 in. or 8 in. where broadest. I discovered, in the maritime parts of Georgia, particularly on the banks of the Alatamaha, another new species of *Magnolia* [*M. aur. pyramidata*], whose leaves were nearly of the figure of those of this tree; but they were much less in size, not more than 6 in. or 7 in. in length, and the strobile very small, oblong, sharp-pointed, and of a fine deep crimson colour; but I never saw the flower. These trees grow straight and erect, 30 ft. or more in height, and of a sharp conical form, much resembling the cucumber tree [*M. acuminata*] in figure." (*Bartram's Travels*, p. 338.)

Geography. This species appears to be confined to a particular part of the Alleghanies, nearly 300 miles from the sea. It is found on the steep banks of the rivers which rise in these lofty mountains. It appears to be very sparingly distributed; the distance of 150 miles occurring, in some cases, between the spots where it is to be met with. Michaux says, "I have no where found it so abundant as on the steepest part of the lofty mountains of North Carolina, particularly those which are called by the inhabitants, Great Father Mountains, and Black and Iron Mountains." The soil of these mountains is brown, deep, and of an excellent quality; and the tree is found to multiply so fast from seed, that a thousand plants might be collected in a single day. The atmosphere, in such situations, is continually charged with moisture, from the number of torrents which rush down from the summits.

History. This tree was discovered by Bartram, from whom it was first received in England by Messrs. Loddiges, in 1786. It was, probably, soon afterwards sent to France; because we find Madame Lemonnier, the widow of Michaux's patron and friend, describing a tree of this species in her garden in 1800, which was 9 ft. high, and had already flowered. As it is of difficult propagation, it is not very generally distributed; but it is found in the principal botanic gardens of the middle of Europe, and in first-rate nurseries.

Soil, Situation, &c. The soil, as we learn from Michaux, ought to be free and deep; and the situation low, sheltered, and moist, rather than dry. As seeds are not very easily procured, the common mode of propagation is by layers, or by inarching on *M. acuminata*. In both modes, two years are required before the plants can be separated from the parent stock. If the demand for plants were adequate, abundance of seeds might, no doubt, be procured from America.

Statistics. The parent tree is at Messrs. Loddiges. In the Exotic Nursery, King's Road, there is a tree of this species 16 ft. high; and in the Hammer-smith Nursery there is one 18 ft. high. At Sherwood, one 18 years planted is only 12 ft. high; at White Knights, one 33 years planted is 26 ft. high; and in the Killerton Nursery, Devonshire, one 8 years planted is 14 ft. high. At Cobham Hall, in Kent, one 17 years planted is 25 ft. high; in Knap Hill Nursery there is a tree 20 ft. high; and at Barton, in Suffolk, one only 10 years planted which is 19½ ft. high. There are trees of this species in Scotland and Ireland, but they are chiefly trained against walls. There are several in the gardens about Paris, and some at Secaux, which have attained the height of 20 ft. In the botanic garden at Rouen the height is 10 ft. In the botanic garden at Ghent there is a specimen 22 ft. high. In Germany, *M. auriculata* is either trained against a wall, or treated as a green-house plant. The price of a single plant, about London, is from 7s. 6d. to 21s.; at Bollwyller, 10 francs; in New York, 1 dollar.

‡ S. M. (? AUR.) PYRAMIDA`TA Bartr. The pyramidal-headed *Magnolia*.

Identification. Mich. Bor. Amer., 1. p. 328.; Dec. Prod., 1. p. 80.; Don's Mill., 1. p. 83.

Engravings. Bot. Reg., t. 497.; E. of Pl. 7917.; and our plate in Vol. II.

Spec. Char. Deciduous. Leaves smooth, the same colour on both surfaces, spatulate, obovate, cordate at the base. Auricles spreading. Sepals 3,

spreading. Petals 9, lanceolate, pointed. (*Don's Mill.*, i. p. 83.) A low tree. North America. White flowers. May and June. 1811. Height 20 ft.

Description, History, &c. This tree, which is found in the western parts of Carolina and Georgia, resembles the preceding species in every particular (except size) so closely, that we have not the least doubt of its being only a variety of it, or, at least, its bearing the same relation to that species which *M. cordata* does to *M. acuminata*. It has been found in only two or three localities. One of these is on the banks of the Alatamaha river in Georgia, 40 miles south of Savannah, where it was discovered by M. le Conte. (See *Gard. Mag.*, vol. viii. p. 288.) It was brought to England in 1818 by Mr. Lyon; and the original tree still exists in Messrs. Loddiges's nursery. It is extremely difficult to propagate (which is done by inarching on *M. auriculata*); and it is, in consequence, very sparingly distributed over the country. Plants, about London, cost 21s. each; at Bollwyller, 15 francs; in New York, ?

§ ii. *Gwillimia* Rott. in *Dec. Syst.*



Derivation. General *Gwillim*, some time governor of Madras. (*Don's Mill.*, i. p. 83.)

Sect. Char. Asiatic species, generally with two opposite spathe-like bractees enclosing the flower-bud. Anthers bursting inwards. Ovaries somewhat distant. Perhaps the species of this section, with one bractea, should have been given among the *michelias*. It is, however, evident, that none of them are true *magnolias*. (*Don's Mill.*, i. p. 83.)

‡ 9. *M. CONSPICUA* *Salisb.* The *Yulan*, or conspicuous-flowered *Magnolia*.

Identification. *Dec. Prod.*, i. p. 81; *Don's Mill.* i. p. 83.

Synonymes. *M. præcia* *Correa*; *M. Yulan* *Desf.*; *Yu lan*, *Chinese*; the Lily-flowered *Magnolia*; *Magnolier Yulans*, *Fr.*; *Yulans Bieberbaum*, *Ger.*

Derivations. The epithet *præcia* was given to this *magnolia* by M. Correa, because it produces its flowers before its leaves. *Yu lan* signifies the lily tree.

Engravings. *Bot. Mag.*, 1621; *Otto and Hayne*, t. 72; *E. of Pl.*, 7907; our *fig.* 34. and the plate of this species in Vol. II.

Spec. Char. Leaves obovate, abruptly acuminate, younger ones pubescent, expanding after the flowers. Flowers erect, 6—9-petaled. Styles erect. (*Don's Mill.*, i. p. 93.) A middle-sized tree in China. Flowers white. Feb. to April. Introduced in 1789. Height from 30 ft. to 40 ft.

Varieties.

‡ *M. c.* 2 *Soulangeana*. *Soulange's* conspicuous-flowered *Magnolia*.

Synonymes. *M. Soulangeana* *An. Hort. Soc. Par.*, *Swt. Fl.-Gard.*, *Don's Mill.*; *Magnolier de Soulange*, *Fr.*

Engraving. *M. Soulangeana* *Swt. Brit. Fl.-Gard.*, t. 260.

Description. The leaves, wood, and general habit of the tree bear so close a resemblance to those of *M. conspicua*, that, when the plant is not in flower, it is almost impossible to distinguish it from that species. The flowers resemble in form those of *M. purpurea* var. *gracilis* or of *M. purpurea*, and the petals are slightly tinged with purple. It was raised at Fromont, near Paris, from the seeds of a plant of *M. conspicua*, which stood near one of *M. purpurea*, in front of the château of M. Soulange-Bodin; the flowers of the former of which had been accidentally fecundated by the pollen of the latter.

‡ *M. c.* 3 *Alexandrina* *Hort.* The Empress *Alexandrina's* conspicuous-flowered *Magnolia*.—This variety so closely resembles the preceding one, as not to be distinguishable from it otherwise than by its flowering somewhat earlier. It was originated at Paris a few years after the preceding variety, and sent to London by the Parisian nurserymen in 1831.


‡ *M. c.* 4 *speciosa* *Hort.* The showy conspicuous-flowered *Magnolia*.

— This is another variety similarly originated, and scarcely, if at all, distinguishable from the last.

‡ *M. c. 5 citriodora* Hort. *The Lemon-scented conspicuous-flowered Magnolia.* — This variety was raised by M. Parmentier of Enghien, but is little known; and, we believe, wholly without merit as a new variety. There are plants of it in the Exotic Nursery, King's Road, Chelsea, which Mr. Knight considers as differing very little from *M. conspicua*.

Other Varieties of M. conspicua. This species, as will hereafter appear, has ripened seeds in various places; and, as it fertilises readily with *M. purpurea* and *M. gracilis*, many new varieties may be expected when the attention of cultivators is more especially directed to the subject. *M. c. Norbértii* is a seedling variety, of which there is a plant in the garden of the Horticultural Society, which has not yet flowered. The plants raised from seed of *M. c. Soulangeana* at Fromont may be productive of something new, as may those raised by Mr. Curtis at Glazenwood, and by Mr. Ward at White Knights. If Signor Manetti succeeds in raising plants from the seeds of *M. c. Soulangeana*, which have ripened at Monza, he also may introduce some new varieties.

Description. This is a very showy tree, distinguishable from all the other magnolias of both sections, by its flowers being of a milk white, and expanding



[Scale, 1/2 in. to 1 ft.]

before any of the leaves. The tree assumes a regular conical shape, with a grey bark and numerous branches and twigs, which generally have a vertical, rather than a horizontal, direction; so that a large tree of this species would probably be more fastigate than any of the others. The young shoots are from 1 ft. to 18 in. in length, and the tree, in ten years, will attain the height of from 10 ft. to 15 ft., flowering the second or third year after grafting. The size of the full-grown tree, in its native country, is said to be from 40 ft. to 50 ft.; the highest which we know of in England is at Eastwell Park, in Kent, which, in 1825, appeared to be upwards of 30 ft.

Geography and History. This tree is said to be a native of the southern provinces of China; and to be extensively cultivated there in the gardens of the emperor, and in those of all eminent persons who can afford to procure it. It began to be cultivated in China in the year 627; and from that time it has always held the very first rank, as an ornamental tree, in their gardens. It is not only planted in the open ground, and allowed to attain its full size, but dwarfs are kept in pots and boxes, and forced throughout the winter, so as to keep up a perpetual supply of bloom in the apartments of the imperial palace. So highly is this tree valued, that a plant in flower, presented to the emperor, is thought a handsome present, even from the governor of a province. In very severe winters, the trunks of the trees in the open air are sometimes wrapped round with straw ropes; but it never requires any other protection, even in the


climate of Pekin. The tree was introduced into England by Sir Joseph Banks in 1789; but it was many years before it attracted much attention, being considered as requiring a green-house or conservatory. So little was it known in 1807, that it is not enumerated among the magnolias described in Martyn's *Müller's Dictionary*, published in that year. Within the last twelve years, it has been discovered to be nearly as hardy as the American species, and it is now most extensively cultivated in the nurseries, both in Britain and on the Continent, and finds a place in every collection. It flowers freely every year, as a standard, in the neighbourhood of London, when the wood has been properly ripened during the preceding summer: and, at White Knights, in England; at Fromont, and various other places, in France; and at Monza, in Italy, it has ripened seeds from which young plants have been raised. Some of the flowers having been fecundated with the pollen of *M. purpurea* or *gracilis*, some hybrid varieties have been produced, of which the most beautiful is *M. c. Soulangeana*.

Properties and Uses. Besides its value as an ornamental plant, the Chinese pickle the flower-buds, after having removed the calyx, and use them for flavouring rice. Medicinally, the seeds are taken in powder, in colds and inflammations of the chest. It is also regarded as stomachic; and water, in which it has been steeped, is used for bathing the eyes when inflamed, and for clearing them of gum. The Chinese poets call the tree the symbol of candour and beauty. (*N. Duh.*, i. p. 225.)

Soil and Situation. A rich sandy loam seems to suit this species best; but it will grow in any deep free soil, properly drained, and moderately enriched. The situation, when it is to be treated as a standard, ought to be sufficiently open to admit of ripening the wood in autumn, and yet not so warm as to urge forward the flower-buds prematurely in spring, as they are very liable to be injured by frost; from which, however, they may be protected by a very slight covering (during nights and frosty days) of gauze or bunting, stretched over the tree horizontally and supported by posts. Against a wall, the tree shows itself in its greatest beauty; and there it can easily be protected, by a projecting coping, from the severest weather ever experienced in the neighbourhood of London. In warm situations, sloping to the south or south-east, the tree has a fine effect planted in front of a bank of evergreens; and, indeed, wherever it is planted, evergreens should be placed near it, and, if possible, so as to form a back ground, on account of the flowers expanding before the tree is furnished with any leaves.

Propagation and Culture. The species and all the varieties are propagated by layers, or by inarching on the *Magnolia purpurea*, or on the *M. acuminata*. When grafted on *M. purpurea*, the tree is comparatively dwarfed, by which it is rendered very convenient for use as a shrub, or for growing in pots; but, when it is intended to form a tree, it should either be grafted on *M. acuminata*, or raised from layers or seeds. It generally requires two years before the plants can be separated from the parent stock. Some plants of this species have been raised from seed ripened in Europe; and we have no doubt that, when this magnificent tree becomes better known and more generally in demand, it will be raised in this way as extensively as *M. acuminata* and *M. glauca* are at present.

Statistics. An original imported plant, against a wall at Wormleybury, measured, in April, 1835, 27 ft. high, covered a space laterally of 24 ft., and had on it, at that time, 5000 flowers! In Lee's Nursery there are several plants above 20 ft. high, as standards, which flower magnificently every year. At Harringay there is a tree of *M. conspicua* 25 ft. high, against a wall; and in the same garden there is one of *M. c. Soulangeana*, 6 years inarched upon a strong plant of *M. cordata*, which is 20 ft. 6 in. high, the branches extending over a space 16 ft. in diameter, and the diameter of the trunk, at a foot from the ground, being 1 ft. 1 in. In the Kensington Nursery there is a tree of the same height (*fig. 35.*), of which we had a drawing made in the first week in April, 1827, when it was covered with 1100 blossoms. There is a large



[Scale, $\frac{1}{2}$ in. to 1 ft.]

tree at Cashiobury; and several at White Knights, one of which, 16 ft. high, has ripened seeds. At Farnham Castle, in Hampshire, one 10 years planted is 16 ft. high; and at Cobham Hall, in Kent, one 16 years planted is 20 ft. high. The largest *M. c. Soulangeana* in the neighbourhood of London is in Brown's Nursery at Slough, where it has attained the height of 15 ft. in less than ten years, and is thought by Mr. Brown to be rather more hardy than *M. conspícua*. At High Clere, on an elevated exposed situation, *M. c. Soulangeana* has attained the height of 7 ft. 6 in. in 4 years, as a standard. *M. conspícua* grows freely, against a wall, in all the low districts of Scotland; and, as a standard, in the neighbourhood of Dublin. In the neighbourhood of Paris, the largest plant of *M. conspícua* is at Fromont. It measured, in 1835, 40 ft. high; and the circumference of the trunk, at 2 ft. from the ground, was more than 2 ft.; and the diameter of the space covered by the branches is 24 ft.; it flowers magnificently every year, at the end of March and be-

ginning of April, and the odour of the flowers is perceived at a great distance. This is the tree, the ripened seeds of which produced *M. c. Soulangeana*. The original plant of *M. c. Soulangeana*, at Fromont, is not more than 12 ft. high, and, though it has flowered every year for several years past, it did not ripen seeds till 1834. These seeds have been sown; and *M. Soulange-Bodin* informs us that he expects some interesting new sorts from them. In the botanic garden at Ghent there is a standard tree of *M. conspœua* 22 ft. high. In Germany, *M. conspœua* is trained against a wall, or kept in the conservatory; and in Italy, and the South of Europe, it has not been long planted; though both the species, and the var. *M. c. Soulangeana*, are at Monza, where, as before noticed, *M. conspœua* has ripened seeds; as has also, as Signor Manetti informs us, *M. c. Soulangeana*.

Commercial Statistics. The price of *M. conspœua* in the London nurseries is from 2s. 6d. to 5s. a plant; at Bollwyller, 10 francs; in New York, ?.

♁ 10. *M. PURPUREA* *Bot. Mag.* The purple-flowered Magnolia.

Identification. Sims, in *Bot. Mag.*

Synonymes. *M. obovata* Thun.; *M. discolor* Vent.; *M. denudata* Lam.; the obovate-leaved Magnolia; *Magnolie discoloré* Bon. Jard., and *Magnolie bicoloré* Dun., Fr.; rothe Bieberbaum, Ger.

Engravings. Sal. Par., t. 87.; *Bot. Mag.*, t. 390.; *E. of Pl.*, 7908.; and our fig. 36.

Spec. Char. Deciduous. Leaves obovate, acute, reticulately veined; almost smooth. Flowers erect, of 3 sepals and 6 obovate petals; styles very short. (*Don's Mill.*, 1. p. 84.) A shrub from Japan, with flowers purple on the outside, and white within. March, April, and May. 1790.

Varieties. In De Candolle's *Prodromus*, and in Don's *Miller*, three varieties are described: *M. p. denudata* Lam., distinguished by the flowering branches being without leaves; *M. p. discolor* Vent., which is said to be rather more tender than the species; and *M. p. liliflora* Lam., the petals of which are white on both sides. These varieties were originally described by Kæmpfer; but, as far as we know, none of them are in British gardens. Several plants of this species having been raised from seed ripened in this country, the plants may exhibit slight shades of difference, as has been the case with certain seedlings raised in the Brentford Nursery; but, as far as we have observed, none of these are worth keeping distinct. The only variety which we consider truly distinct is *M. p. gracilis*, considered as a species by Salisbury and other botanists, but which, we are convinced, is nothing more than a race, or a variety.

Description. A deciduous shrub, attaining, in the gardens about London, the height of from 6 ft. to 10 ft. in as many years, and seldom growing much higher as a bush. The stems are numerous, but not much branched; the leaves are large, of a very dark green; and the plant produces a profusion of flowers, which do not expand fully till a day or two before they drop off; and which, unless the weather is warm, do not expand at all, but wither on the plant, and disfigure it. The flowers are large, more or less purple (according to the season, but never wholly dark purple) without, and always white within. The bark, when bruised, has an aromatic odour. It is a very ornamental species, and no garden ought to be without it.

Geography, History, &c. Found wild in Japan; and cultivated there, and in China, in gardens. It was discovered by Thunberg, and imported by him into England, in 1790. It has since been generally distributed throughout the botanic and first-rate private gardens of Europe. About London and Paris, it is not only propagated for

36



sale as a flowering shrub, but as a stock for grafting other species on, even of the tree kinds: such as *M. conspicua*, *cordata*, and others. In the north of France, and in Germany, it is generally treated as a green-house plant.

Soil and Situation. This species is generally considered as requiring a mixture of heath soil, or sandy peat, with loam; but in many gardens about London it succeeds perfectly both in sand and clay; the latter soil being rendered free by sand, leaf mould, or manure, and drainage. The situation, when the plant is treated as a bush, ought to be open, in order that the wood may be ripened; and the plant should be detached, in order that it may be covered with foliage and blossoms on every side. North of London, in most situations, it requires a wall, and few plants are more deserving of one. Against a wall, it will reach the height of 15 ft. or 20 ft.

Propagation and Culture. In the London nurseries, it is generally propagated by layers; but it will also strike by cuttings, both of the ripened and the herbaceous wood. The stools are generally formed in pits; or, if in the open ground, they are generally covered with mats during winter. Seeds have been ripened both in England and France; and from these plants have been raised in some few nurseries. The plants, whether raised from layers, cuttings, or seed, should always be kept in pots till wanted for final planting.

Statistics. There are fine plants of this species, trained against walls, at Harringay, Wormleybury, White Knights, and numerous other places. The largest bushes in the neighbourhood of London are in the Mile End Nursery; and there are very handsome specimens in the Hammersmith Nursery. There is one 20 ft. high in the garden of the Rev. J. Mitford, at Benwell in Suffolk, which, we believe, is the largest in England. Price, in London, from 1s. 6d. to 2s. 6d. a plant; at Bollwyller, 5 francs; in New York, ?.

♯ 11. *M. (? P.) GRACILIS.* The slender-growing purple-flowered Magnolia.

Synonymes. *Magnolia Kobus* Kæmpf., Dec., and Don.; *M. glauca* var. *a* Thun. *Fl. Jap.*; *M. tomentosa* Thun. in *Linn. Soc.*; Sidi Kobusi, *Japanese*; *Magnolier grêle*, Fr.; *filziger Bieberbaum*, Ger.

Derivations. Kobus, or Sidi Kobusi, is the name of the plant in Japan. It is named *gracilis* from its slender habit of growth; and *tomentosa* from the comparatively downy surface of the leaves. It was called *M. glauca* by Thunberg, because he originally supposed it to be a variety of the *M. glauca* of North America. The French and German names signify the same as the specific one.

Engravings. Kæmpf. *lc.*, t. 42.; Par. *Lor.*, 87.; E. of Pl., 7909.

Description. In its native country it is a small tree with rough bark; but in England it is a somewhat delicate shrub, with slender stems and branches, growing rather more erect and fastigate than *M. purpurea*. The leaves are of the same form, but a little longer, and always of a decidedly paler green. The young leaves are pubescent underneath, as are the young shoots. In England, though this kind cannot be considered as a tree, yet it has a different habit of growth from *M. purpurea*; and, instead of, like it, forming a broad spreading bush, it is a narrow, upright, slender, fastigate-growing one. The bark has the same odour as that of *M. purpurea*. The two main points of difference between it and *M. purpurea* are, the paler green, and somewhat narrower shape, of the leaves; and the longer and more slender form of the flower, the points of the petals of which are slightly turned back; while the flower of *M. purpurea* is more cup-shaped, and the petals at the points are rather turned inwards. The petals of *M. gracilis* are exteriorly of an entirely dark purple, whereas those of *M. purpurea* melt off into white at their upper extremities.

Geography, History, &c. It is a native of Japan, where it is said by Kæmpfer to attain the size of a cherry tree. It was brought to England in 1804, but is not very common, probably from its being very generally confounded with *M. purpurea*. Its management, in all respects, is the same as that of the species, except that it is, perhaps, somewhat more tender. The largest plant that we know of is in the conservatory at White Knights, where it forms a narrow bush about 10 ft. high. It is kept in the conservatory there, not on account of its tenderness, but because it was considered to be a conservatory plant when it was first planted; and it is now much too

large to be removed with safety. There are bushes of this variety in the open border, both in the Hammersmith Nursery and at Mile End, between 3 ft. and 4 ft. high, and 2 ft. and 3 ft. broad, which flower freely every year, without any protection whatever. Price, in the London nurseries, 5s.; at Bollwyller, ?; and in New York, ?.

App. i. *Half-hardy Magnolias.*

Magnolia fuscata Andr., figured in *Bot. Mag.*, t. 1008., and introduced from China in 1789, is common in conservatories. At Claremont it forms a large evergreen bush, 15 ft. high; and at Taplow Court, a bush 10 ft. high. It flowers in April, May, and June: its fragrance is much more grateful than that of the other magnolias, and not at all oppressive. The plant has been tried against a conservatory wall; and also, in the Goldworth Nursery, in the open ground, as a bush; and in both situations, when once firmly established, it is found to endure moderate winters with a little protection. On account of its being an evergreen, and from the fragrance of its flowers, which are of a dark brownish red or purple, tinged with yellow in the inside of the petals, it is a very desirable plant. There are two other species, natives of China (*M. ananclifolia* and *M. pumila*), also evergreens, and treated as green-house or stove plants; but we have not heard of either of them having been tried out. Possibly, they may prove nearly as hardy as *M. fuscata*.

App. ii. *Additional Magnolias.*

It is highly probable that there are other species of the genus *Magnolia*, in the mountainous regions of India, and in China, which will endure the open air in Britain, though none of these have yet been described by botanists, with the exception of some by Dr. Wallich, which are now considered to belong to *Michelia*. Some expected additions of genera closely allied to *Magnolia* will be noticed in the concluding section of this chapter. Possibly, by cross fecundation, some mules might be produced, between the species mentioned in the preceding paragraph and the hardy species. If the refreshing fragrance of *M. fuscata* could be thrown into the flowers of *M. grandiflora*, or of any of the other species which continue flowering for a long time, the result would be a desirable acquisition. We recommend the subject to the attention of ingenious cultivators.

GENUS II.



LIRIODE'NDRON *L.* THE TULIP TREE.

Gen. Char. *Carpels* 1-2-seeded, disposed in spikes, indehiscent, deciduous, drawn out into a wing at the apex. *Calyx* of 3 deciduous sepals. *Corolla* of 6 petals, conniving into a bell-shaped flower. (*Don's Mill.*, i. p. 86.) — There is only one species; a deciduous tree of the first rank. North America. Flowers yellowish, variegated with green, red, and orange. June. 1688. Height, in England, 70 ft.

♂ I. LIRIODE'NDRON TULIPÍFERA *L.* The Tulip-bearing Liriodendron, or Tulip Tree.

Identification. *Lin. Sp.*, 755.; *Hayne Dend.*, 115.; *Dec. Prod.*, 1. p. 82.; *Don's Mill.*, 1. p. 86.

Synonymes. The Poplar, White Wood, Canoe Wood, the Tulip Tree, *Amer.*; Virginian Poplar, Tulip-bearing Lily Tree, Saddle Tree, *Eng.*; Tulipier de Virginie, *Fr.*; Virginscher Tuliperbaum, *Ger.*

Derivations. This tree is called Liriodendron, from *leirion*, a lily, and *dendron*, a tree; from the flowers resembling those of a lily, though more correctly those of a tulip, as the specific name implies. It is called Poplar, from its general resemblance to trees of that genus; White Wood, from the colour of its timber; Canoe Wood, from the use to which it is applied by the native Indians; Tulip Tree, from its tulip-like flowers; and Saddle Tree, from the form of its leaves. The French and German names are literal translations of the words Virginian tulip tree.

Engravings. *Bot. Mag.*, 27.; *Dub.*, tom. 3. t. 18.; *Willd. Ab.*, t. 29.; *Krause*, t. 51.; *E. of Pl.*, 7905.; and our plate in Vol. II.

Spec. Char. Leaves smooth, truncate at the top; 4-lobed, resembling a saddle in shape. Flowers large, solitary, terminal, variegated with green, yellow, and orange colour; furnished with two deciduous bracteas under flowers. (*Don's Mill.*, i. p. 86.)

Varieties.

- ✕ *L. T. 2 obtusiloba* Michx., *integrifolia* Hort., the obtuse-lobed, or entire-leaved, Tulip Tree, Yellow Wood, or Yellow Poplar, has the leaves with blunter lobes than the original species, but is in no other respect different from it. See the plate of this tree in our Second Volume.
- ✕ *L. T. 3 acutifolia* Michx. has the leaves smaller and more acutely cut than either the preceding variety or the species. We are not aware of any plants of it existing in British gardens.
- ✕ *L. T. 4 flava* Hort. has yellow flowers. As most of the tulip trees in Europe have been raised from seeds, it is probable that their flowers vary in degrees of yellowness; but we are not aware that any variety, with the flowers entirely yellow, is propagated in British nurseries. It is, however, in the catalogue of the garden at Courset, and in some of the Paris catalogues.

Description. This tree, in the Atlantic states of North America, according to Michaux, especially at a considerable distance from the sea, is often seen from 70 ft. to 100 ft. in height, with a trunk the diameter of which varies from 18 in. to 3 ft. The elder Michaux found in Kentucky, on the road from Beardstone to Louisville, tulip trees which appeared to be 15 ft. or 16 ft. in circumference; and, three miles and a half from Louisville, he measured one which, at 5 ft. from the ground, was 22 ft. 6 in. in circumference, the height of which he found to be from 120 ft. to 140 ft. Of all the deciduous trees of North America, the tulip tree, next to the button-wood (*Platanus occidentalis*), attains the amplest dimensions; while the perfect straightness and uniform diameter of its trunk for upwards of 40 ft., the regular distribution of its branches, and the richness of its foliage, give it a decided superiority to that tree, and entitle it to be considered one of the most magnificent trees of the temperate zones. In the development of its leaves the tulip tree differs from most other trees. The leaf-buds, in general, are composed of scales closely imbricated, which, in the spring, are distended by the growth of the minute bundle of leaves that they enclose, till they finally fall off. The terminal bud of each shoot swells considerably before it gives birth to the leaf: it forms an oval envelope, which contains the young leaf, and which produces it to the light only when it appears to have acquired sufficient force to endure the influences of the atmosphere. Within this envelope is found another, which, after the first leaf is put forth, swells, bursts, and gives birth to a second. On young and vigorous trees five or six leaves issue, successively, in this manner, from one bud. Till the leaf has acquired its growth, it retains the two scales which composed its envelope, and which are now called stipules. In the spring, when the weather is warm and humid, the growth of the leaves is very rapid: they are 6 in. or 8 in. broad, borne on long petioles, alternate, somewhat fleshy, smooth, and of a pleasing green colour. They are divided into three lobes; of which the middle one is horizontally notched at its summit, and the two lower ones are rounded at the base. This conformation is peculiar to the tulip tree, and renders it easily distinguishable in the summer. The flowers, which are large, brilliant, and, on detached trees, very numerous, are variegated with different colours, among which yellow predominates; they have an agreeable odour, and, surrounded by the luxuriant foliage, they produce a fine effect. In the spring they are gathered by women and children in the neighbourhood of New York, and sold in the market of that city. The fruit is composed of a great number of thin narrow scales, attached to a common axis, and forming a conical spike 2 in. or 3 in. in length. Each fruit contains 60 or 70 carpels; of which never more than a third, and in some seasons not

more than seven or eight in the whole number, are productive. It is also observed, that, during ten years after it begins to yield fruit, almost all the seeds are unproductive; and that, on large trees, the seeds from the highest branches are the best. The bark, till the trunk exceeds 7 in. or 8 in. in diameter, is smooth and even: it afterwards begins to crack, and the depth of the furrow, and the thickness of the bark, are proportioned to the size, and to the age of the tree. The heart, or perfect wood, of the tulip tree is yellow, approaching to a lemon colour; and its sap, or alburnum, is white. (*Michaux.*) In Europe, though the tulip tree does not attain the same magnitude that it does in situations favourable to it, in its native country, it still forms a magnificent tree; in some cases, both in Britain and in the middle of the European continent, reaching the height of 90 ft. or 100 ft., flowering freely, and sometimes ripening seed. The annual shoots of young plants, in the neighbourhood of London, are from 18 in. to 2 ft. in length; and the tree will, in favourable circumstances, attain the height of from 15 ft. to 20 ft. in ten years; seldom, however, flowering till it is upwards of twenty years old. The height, in England, frequently exceeds 70 ft.; and it has ripened seeds here, occasionally, from which young plants have been raised. It ripens its fruit very generally in France; though it is observed, in the *Nouveau Du Hamel*, that these seeds do not vegetate so freely as those which are imported from America.

Geography. The southern extremity of Lake Champlain, in latitude 45°, according to Michaux, may be considered as the northern, and the Connecticut river, in the longitude of 72°, as the eastern, limit of the tulip tree. It is found beyond the Hudson, which flows two degrees farther west; and below 43° of latitude it is frequently met with, and fully developed. Its expansion is not here repressed, as in Vermont, and in the upper part of the Continent, by the excessive cold, and by a mountainous surface unfavourable to its growth. It abounds in the middle states, in the upper parts of the Carolinas and of Georgia; and is found still more abundantly in the western country, particularly in Kentucky. Its comparative rareness in the maritime parts of the Carolinas and of Georgia, in the Floridas, and in Lower Louisiana, is owing less to the heat of the summer than to the nature of the soil; which, in some parts, is too dry, as in the pine barrens, and in others too wet, as in the swamps which border the rivers. Even in the middle and western states, the tulip tree is less abundant than the oaks, the walnuts, the ashes, and the beeches, because it delights only in deep, loamy, and extremely fertile soils, such as are found in the rich bottoms that lie along the rivers, and on the borders of the great swamps that are enclosed in the forests. In the Atlantic states, especially at a considerable distance from the sea, tulip trees are often seen 70 ft., 80 ft., and 100 ft. in height, with trunks from 18 in. to 3 ft. in diameter; but the western states seem to be the natural soil of this magnificent tree, and there it displays its most powerful vegetation. It is commonly found mingled with other trees, such as the hickories, the black walnut and butter nut, the Kentucky coffee tree (*Gymnocladus*), and the wild cherry tree: but it sometimes constitutes, alone, pretty large tracts of the forest; as was observed by the elder Michaux on the road from Beardstone to Louisville. In no other part of the United States did he find tulip trees so lofty, and of so great a diameter. (*Michaux.*) The artificial geography of this tree may be said to embrace the middle region of Europe, from Berlin and Warsaw, on the north, to the shores of the Mediterranean and Naples, on the south; Ireland, on the west; and the Crimea, on the east.

History. When the tulip tree was first introduced into England is uncertain; but it was cultivated by Compton, at Fulham, in 1688. It was, however, at that time, wholly unknown as a timber tree. Evelyn, speaking of it, says, "They have a poplar in Virginia of a very peculiar-shaped leaf, which grows well with the curious amongst us to a considerable stature. I conceive it was first brought over by John Tradescant, under the name of the tulip tree, from

the likeness of its flower; but it is not, that I find, taken much notice of in any of our herbals. I wish we had more of them; but they are difficult to elevate at first." (*Hunter's Evelyn*, i. p. 207.) According to Miller, Mr. Darley, at Hoxton, and Mr. Fairchild, were the first who raised this tree in any quantity from seeds; and from their nurseries it is probable that the numerous old trees which are spread all over the country were procured. The first notice which we have of the tulip tree on the Continent is in the *Catalogue of the Leyden Garden*, published in 1731. From the number of tulip trees existing in France, the south of Germany, and Italy, there can be little doubt that it spread as rapidly in those countries as it did in Britain. Public avenues are planted of it in Italy, and as far north as Strasburg and Metz. It stands the open air at Vienna, and attains a large size there; but it will not endure the open air north of Warsaw, or at Moscow, without protection. The first tulip tree which flowered in England was one in the gardens of the Earl of Peterborough, at Parson's Green, near Fulham. This is understood to have been the first tree which was planted in the open ground: previously, they had, like most other American trees in those times, been grown in pots, and housed every winter. This tree at Parson's Green, Miller says, convinced gardeners of their mistake, by the great progress which it made; so that afterwards there were a great many planted in open ground, which, more especially those on a moist soil, speedily attained a large size. Some at Waltham Abbey, and at Wilton, are referred to by him as among the oldest and largest.

Properties and Uses. The timber of the tulip tree, though classed among light woods, is yet, Michaux observes, much heavier than that of the common poplar; its grain is equally fine, but more compact; and the wood is easily wrought, and polishes well. It is found strong and stiff enough for uses that require great solidity. The heart-wood, when separated from the sap, and perfectly seasoned, long resists the influence of the air, and is said to be rarely attacked by insects. Its greatest defect, when employed in wide boards, and exposed to the weather, is, that it is liable to shrink and warp, by the alternations of dryness and moisture: but this defect is, in a great measure, compensated for by its other properties, and may be, in part, owing to its not being allowed sufficient time to be properly seasoned. The nature of the soil has so striking an influence upon the colour, and upon the quality of the tulip wood, that the mechanics who employ it have made the remark; and have distinguished it by the names of the white poplar, and the yellow poplar. The external appearances which mark these varieties are so equivocal, that they can only ascertain to which of them a tree belongs by cutting it. It is known, in general, that the white poplar grows in dry, gravelly, and elevated places; it is recognised, too, by its branchy summit, and by the small proportion which the light yellow heart-wood bears to the sap-wood. The grain, also, is coarser and harder, and the wood decays more speedily; hence it is always neglected, when the other variety can be obtained. The yellow poplar possesses every quality requisite to fit it for a great variety of uses. At New York and Philadelphia, and in the adjacent country, it is often employed in the construction of houses, for rafters and for the joists of the upper stories, for which purposes it is esteemed on account of its lightness and strength. In the other middle states, in the upper parts of the Carolinas, and, above all, in the western states, it is more generally used in building, and is considered as the best substitute for the pine, the red cedar, and the cypress. Wherever it abounds, it serves for the interior work of houses, and sometimes for the exterior covering. The panels of doors and of wainscots, and the mouldings of chimneypieces, are made of this wood. In the states of Ohio and Kentucky, on the banks of the Miami river, and in the upper part of North Carolina, shingles of it, about 15 in. long, are preferred for covering roofs; because they are the most durable, and because they are not liable to split from the effect either of intense frost, or of ardent sunshine. In all the large towns of the United States, tulip tree, or, as they are there called, poplar, boards, which are

often 2 ft. or 3 ft. wide, are exclusively used for the panels of carriages. When perfectly dry, they take the paint well, and admit of a brilliant polish. The vicinity of Boston does not produce this tree, and the coachmakers there procure it from New York and Philadelphia; it is also sent for the same use to Charleston, S. C., where the tulip trees are few in number, and inferior in size. The seats of the Windsor chairs which are fabricated at New York, Philadelphia, and Baltimore, and in many other towns, are always of this wood: a large quantity of it is consumed in this way, and also in the manufacture of trunks which are covered with skins, and of bedsteads which are stained in imitation of mahogany. The circular board and wings of winnowing-machines are made of this wood, as it is easily wrought in the lathe, and is very light; it is also much used for wooden bowls, and for the heads of hair brooms, or sweeping-brushes. The farmers use it for the eating and drinking troughs of cattle: these troughs are formed of a single piece; and, exposed to the weather, they last as long as those made of chestnut and butternut (*Carya*). In Kentucky, the wood of the tulip tree is sometimes employed for the rails of rural fences. It is found useful, also, in the construction of wooden bridges, as it unites lightness with strength and durability. The Indians who inhabited the middle states, and those who still remain in the western country, prefer this tree for their canoes; which, consisting of a single trunk, are very light and strong, and sometimes carry twenty persons. The wood of the tulip tree affords excellent charcoal, which, in America, is employed by the smiths in the districts that furnish no fossil coal. In the lumber yards of New York, Philadelphia, and Baltimore, a great quantity of this wood is found, in forms convenient for the uses above enumerated. In America it is very cheap; being sold at half the price of black walnut, wild cherry, and curled maple. In all the country watered by the river Monongahela, between 39° and 40° of latitude, the tulip tree is so abundant, that large rafts, composed wholly of its logs, are made to float down the stream. Michaux remarks that, when a poplar is felled, the chips of the heart-wood that are left upon the ground, particularly those which are half buried in the leaves, suffer, at the end of three or four weeks, a remarkable change; the lower part becomes of a dark blue, and they exhale a fetid ammoniacal odour; though the live part of the bark of the trunk, the branches, and still more of the roots, has an agreeable smell, and a very bitter taste; and, even under the same circumstances as the heart wood, it neither acquires the blue colour, nor the disagreeable smell.

Medical Properties. In Virginia, some of the inhabitants of the country steep the bark of the roots of this tree, with an equal portion of dogwood bark, in brandy, during eight days; and this tincture is considered a cure for intermittent fevers. Poplar bark, reduced to powder, and given in substance to horses, appears to be a pretty certain remedy for worms. The *American Museum* for December, 1792, contains details of the valuable properties ascribed to this bark, by Dr. Young of Philadelphia; from which it appears that it is nearly equal to quinquina, being a powerful tonic and antiseptic: the aromatic principle appears to reside in a resinous part of its substance, which stimulates the intestinal canal, and which operates as a gentle cathartic. In many instances, the stomach cannot support it, unless each dose is accompanied by a few drops of laudanum. In Paris, a spirituous liquor is made from the fresh bark of poplar roots, with the addition of a sufficient quantity of sugar to render it agreeable to the taste. (*Michaux.*)

The Uses of the Tulip Tree in Europe are limited almost entirely to those of ornament; for, though there are numerous trees which would produce excellent timber if cut down, we have never heard of any having been felled for this purpose, or, indeed, for any other. Every possessor of a tulip tree, in Europe, values it far higher for its beauty in a living state, than for its products, or the artificial applications of them. On the Continent, where trees ripen seeds, they may be considered as affording some profit from that source.

Soil and Situation. In its native country, according to Michaux, the tulip tree delights only in deep, loamy, and extremely fertile soils, such as are found

in the alluvial plains on the margins of rivers, and on the borders of swamps. Like almost all other trees, however, it will grow on soils of different descriptions, and have its timber and other properties more or less affected by the circumstances in which it is placed. In deep rich soil, the wood is yellow and heavy; in dry gravelly soil, on an elevated situation, it is white, light, coarse, hard, and decays more readily than the wood which has grown on rich soil. According to Du Hamel, it neither thrives in France on a dry and gravelly soil, nor on one with the subsoil of clay or marl. The most rapid-growing young tulip trees which we have heard of in England were planted in a deep sandy loam, in a rather moist climate, in the West Riding of Yorkshire; and the progress of these has been at the rate of 16 ft. in 10 years, from the seed. (See *Gard. Mag.*, vol. xi. p. 259.) The situation most favourable for the tulip tree is one which, while it is sheltered from high winds, is, at the same time, sufficiently exposed to the light and air to admit of the maturation of its leaves on every side, and the perfect ripening of its wood, without which it can neither resist the severe frosts of winter, nor form blossom-buds. If it were desired to grow the tree for the purpose of forming straight clean timber, it ought to be placed in a close plantation, where one plant would draw up another.

Propagation and Culture. The species is seldom, if ever, propagated otherwise than by seeds, which come up best in heath soil, very fine mould, or sandy loam, in a shady situation, kept rather moist; but the varieties are, of course, multiplied by layers, budding, grafting, or inarching. When the seeds are sown in autumn, they generally come up the following spring; but, sown in spring or the beginning of summer, they generally remain a year in the ground. Formerly, nurserymen used to raise them on heat; by which means spring-sown seeds came up the succeeding summer. In France, and, occasionally, in England, the obtuse-lobed variety is raised by layers or inarching; but, in either case, it requires two or three years before the plants can be separated from the parent stock. The tulip tree, like the magnolias, having roots furnished with but few fibres, does not transplant readily; and, therefore, the plants ought either to be kept in pots, or, if in the free ground, transplanted in the nursery every year; or, if neither of these modes be practicable, removed to their final situation, when not more than two, or at most three, years old. The tree is, like the magnolias, not very patient of the knife, either in a young or in an old state; and, from the bitter qualities of the leaves, it does not seem to be much attacked by insects.

Geographical Statistics. *Liriodendron Tulipifera* in the *Environs of London.* The largest tulip tree that we have seen in the neighbourhood of London is at Syon, where, in about 70 years, it has attained the height of 76 ft. The trunk, at 1 ft. from the ground, measures 2 ft. 6 in. in diameter; and the diameter of the space covered by the branches is 46 ft. (See the plate of this tree in our Vol. II.) The next largest tree is at Mount Grove, Hampstead, the residence of T. N. Longman, Esq., 80 years planted, 70 ft. high, the diameter of the trunk is 3 ft. 10 in., and that of the head 49 ft. The oldest tree, estimated at 150 years, is at Fulham Palace: it is 55 ft. high; the trunk, at 1 ft. from the ground, is 3 ft. in diameter; and the head, which is in a decaying state, is 25 ft. in diameter. In the arboretum at Kew, there is a tree 60 years planted, which is 70 ft. high; the diameter of the trunk is 2 ft. 8 in.; and that of the space covered by the branches is 35 ft. All these trees flower freely every year, and, in fine seasons, ripen some seeds.

Liriodendron Tulipifera South of London. The dimensions of a great number of specimens have been sent us, from which we select the following, partly to show the rate of growth, and partly to show the ultimate magnitude. In Berkshire, at High Clere, 14 years planted and 28 ft. high, in an exposed situation, 500 ft. above the level of the sea. In Cornwall, at Port Elliot, 80 years planted and 60 ft. high; and at Carlew, 40 years planted and 60 ft. high. In Devonshire, at Killerton, 70 years planted and 63 ft. high; and in the Killerton Nursery, 22 years planted and 37 ft. 6 in. high; at Luscombe, 21 years

planted and 37 ft. high; at Endsleigh, 18 years planted and 35 ft. high. In Hampshire, at Farnham Castle, 55 years planted and 40 ft. high, on chalk. In the Isle of Jersey, in Saunders's Nursery, 10 years planted and 18 ft. high, on strong clay. In Somersetshire, at Hestercombe, 96 ft. high; with a trunk nearly 3 ft. in diameter; a magnificent tree, which ripens seeds every year. In Sussex, at Cowdray, 40 ft. high. In Surrey, at Claremont, 70 ft. high; in sandy loam on clay; at Oakham Park, 28 years planted and 32 ft. high; at Melbourne, 70 ft. high. In Wiltshire, at Longleat, 70 ft. high; at Corsham, 66 ft. high; and at Wardour Castle, 50 years planted and 60 ft. high.

Liriodendron Tulipifera North of London. In Bedfordshire, at Southill, 22 years planted and 38 ft. high. In Herefordshire, at Stoke Edith, 20 years planted and 36 ft. high. In Lancashire, at Latham House, 60 years planted and 43 ft. high. In Leicestershire, at Elvaston Castle, 34 years planted and 45 ft. high. In Northamptonshire, at Wakefield Lodge, 9 years planted and 15 ft. high. In Pembrokeshire, at Golden Grove, 70 years planted and 60 ft. high. In Radnorshire, at Maeslough Castle, 50 years planted and 73 ft. high; the trunk 2 ft. in diameter, and the diameter of the tree 36 ft.; the soil a deep yellow loam. In Staffordshire, at Trentham Hall, 50 ft. high, with the trunk 2 ft. in diameter. In Suffolk, at Culford, 8 years planted and 14 ft. high; at Livermere, 30 years planted and 26 ft. high, on strong clay in a northern exposure; at Wolverton Hall, 60 ft. high, and the trunk 2 ft. 8 in. in diameter. In Warwickshire, at Combe Abbey, 50 years planted and 40 ft. high. In Worcester-shire, at Kinlet, 50 years planted and 60 ft. high; the diameter of the trunk 2 ft. 7 in., and of the head 60 ft., containing 35 cubic feet of timber; the soil a sandy loam, and the situation sheltered. The lower part of this tree always comes into leaf before the upper part has the least appearance of doing so; the cause of which is, that the lower part is sheltered by high ground, while the upper part is exposed to a strong west wind: it flowers freely, and has a splendid appearance at that season, and also in autumn, before it sheds its yellow leaves. In the same county, at Croome, 70 years planted, 75 ft. high; the trunk 2 ft. 6 in. in diameter, and the diameter of the head 35 ft. on strong loam; at Hagley, 12 years planted and 15 ft. high, on a sandy loam. In Yorkshire, in the Hull Botanic Garden, 20 years planted and 30 ft. high, in strong loam on clay; at Ripley Castle, 10 years planted and 16 ft. high; at Knedlington, 10 years from the seed, sown on the spot, 14 ft. to 16½ ft. high; the trunk from 2½ in. to 4 in. in diameter, and the diameter of the head 5 ft. or 6 ft.; at Grimstone, 12 years planted and 35 ft. high; the diameter of the stem 7½ in., and of the head 15 ft.; the soil a deep free loam, and the situation sheltered.

Liriodendron Tulipifera in Scotland. Near Edinburgh, at Gosford House, 20 years planted and 20 ft. high; and at Hopeton House, two trees, 86 years planted and 60 ft. high, with trunks 2 ft. 4 in. in diameter, and heads 30 ft. in diameter. There is a tree at Tynningham, 72 years planted, 34 ft. high, and the diameter of the trunk 2 ft. 3 in.; at the Hirsell, a low tree, 100 years planted, 13 ft., the trunk 4 ft., and the branches 33 ft., in diameter; at St. Mary's Isle, 60 years planted and 50 ft. high; at Cassinacrie, 55 ft. high; in the Glasgow Botanic Garden, 14 years planted and 13 ft. high, the young shoots occasionally cut down in winter, especially if the preceding summer has been such as not to ripen them fully; at Roseneath Castle, 55 ft. high. In Fifeshire, at Dinbristle Park, 40 ft. high. In Perthshire, at Annat Gardens, 27 years planted and 20 ft. high; at Gerthy, 2 trees, 40 ft. high, which flower occasionally. In Ross and Cromarty, at Coul, 10 years planted and 10 ft. high, and the diameter of the trunk 2 in.; the situation 160 ft. above the level of the sea, and in north latitude 55° 35'. In Sutherlandshire, at Dmrobin Castle, 20 years planted and 10½ ft. high; the diameter of the trunk 2¾ in., and of the head 11 ft.

Liriodendron Tulipifera in Ireland. Near Dublin, in the Glasnevin Garden, 30 years planted and 20 ft. high; at Cypress Grove, 35 ft. high, flowering freely every year in strong loam or clay; at Howth Castle, 36 ft. high; at Terenure, 9 years planted and 11 ft. high; in Cullingswood Nursery, 24 years planted and 26 ft. high. In Munster, at Castle Freke, 35 ft. high. In Leinster,

at Oriel Temple, 40 years planted and 43 ft. high; at Charleville Forest, 45 years planted and 54 ft. high; at Shelton Abbey, 50 years planted and 60 ft. high. In Ulster, at Florence Court, 38 years planted and 45 ft. high; the diameter of the trunk $2\frac{1}{2}$ ft., and of the head 30 ft. In Connaught, at Mackree Castle, 37 ft. high.

Liriodendron Tulipifera in Foreign Countries. In France, in the Toulon Botanic Garden, 48 years planted and 40 ft. high, the trunk 3 ft. in diameter, in calcareous soil; at Mereville, 30 years planted and 60 ft. high, in a free moist soil; near Nantes, 40 years planted and 50 ft. high. In Holland and the Netherlands, in the Ghent Botanic Garden, 70 ft. high; in the grounds of the palace of Läckén, near Brussels, there is a tree which ripens seeds every year, noticed in p. 145. In Prussia, at Harbeke, 10 years planted and 14 ft. high; at Sans Souci, Potsdam, 42 years planted 50 ft. high; in the Berlin Botanic Garden, 18 years planted and 40 ft. high; the shoots sometimes injured by the frost. At Schwöbber, near Hanover, 120 years planted, 80 ft. high; the diameter of the trunk 2 ft., and of the head 30 ft.; in alluvial soil near water. In Saxony, at Wörlitz, 60 years planted and 30 ft. high. At Munich, in the public garden, 20 years planted and 20 ft. high; in a private garden near the city, 36 years planted and 50 ft. high, flowering freely every year. In Cassel, at Willhelmsöhe, 60 years planted and 20 ft. high. In Austria, in the University Botanic Garden, 20 years planted and 24 ft. high; at Laxenburg, 40 years planted and 30 ft. high; at Kopenzel, near Vienna, 60 years planted and 45 ft. high; at Brüek on the Leytha, 40 years planted and 51 ft. high. In Italy the tree abounds, and attains the height of 70 ft. or 80 ft., flowers freely, and ripens seeds, as may be seen by referring to p. 169.

Commercial Statistics. Plants are abundant in all the European and American nurseries. In London, seedlings are 12s. a hundred; transplanted plants, 2 ft. high, 50s.; and those from 3 ft. to 4 ft. high, 75s.; and seeds are 1s. 6d. a quart. At Bollwyller, one year's seedlings are 20 francs a hundred; two years' seedlings, 35 francs; plants in pots, 1 franc 5 cents each; and plants from 6 ft. to 9 ft. high, from 2 francs to 3 francs 50 cents; and the entire-leaved variety is charged 4 francs. In New York, plants are 20 cents each, and seeds 4 dollars and 50 cents a bushel.

App. i. *Expected Additions to the Order Magnoliacæ.*

In our list (p. 173.) of the Magnoliacæ of the Himalaya, which might probably endure the open air in England, are included *Manglietia insignis*, the *Magnolia insignis* of Dr. Wallich, which grows on the mountains of Nepal; *Michèlia lanuginosa*, *excelsa*, Kisòpa, and Doltsòpa; all of which, being found in elevated regions in the Himalaya, Mr. Royle conjectures would stand the open air in Devonshire, and, with a little protection, in the climate of London. *Michèlia Doltsòpa* is one of the finest trees in Nepal, yielding a fragrant wood much used there for house-building. (*Don's Prod.*, 226.) *Michèlia excelsa*, according to Dr. Wallich, produces a valuable timber of a fine texture, at first greenish, but soon changing into a fine yellow. We have already observed (p. 173.) that there are probably various species of Magnoliacæ in China and Japan, not yet introduced, which would prove hardy, and the introduction of which would amply repay patriotic travellers and European residents in those countries.

As many of the species of *Magnolia* seem to admit of cross-fecundation, it is possible that the same thing may be practicable, to a greater or less extent, between the genera composing the order. The tulip tree, rendered sub-evergreen, would be an interesting object; as would a variety of it with fastigiate branches, like the Lombardy poplar; or one as truly pendulous as the weeping ash; or one with dark leaves, like those of the purple beech. No doubt, a variegation might be produced in the leaves both of the tulip tree and of the magnolia.

CHAP. IV.



OF THE HALF-HARDY LIGNEOUS PLANTS OF THE ORDER DILLENIACEÆ.

We introduce this order chiefly for the sake of recommending some species of *Hibbertia* as half-hardy climbers. All the *Hibbertias* are either natives of the Cape of Good Hope, or of Australia; and, probably, the whole of them might live against a wall with protection. They grow freely either in sandy loam mixed with leaf mould, or in sand and peat; and they are readily increased by cuttings, either of the young or of the ripened wood.

1. *Hibbertia volubilis* Bot. Rep., t. 126., and our fig. 38., the twining *Hibbertia*, was introduced from the Cape of Good Hope in 1790, and has been long known, in green-houses and conservatories, as *Dillenia volubilis*, or *Dillenia scandens*. The flowers are about the size of those of *Hypericum calycinum*: they are of a bright yellow, and are produced all the season, from the beginning of May to the end of October. The plant is a vigorous grower; and, in conservatories, will extend to the height of 8 ft. or 10 ft. in about as many years.



2. *Hibbertia dentata* R. Br., Bot. Reg., t. 282., and our fig. 37., the toothed-leaved *Hibbertia*, was introduced from New Holland, where it grows on the Blue Mountains, in 1814. It is a vigorous-growing twiner, like the preceding species, with narrower leaves and rather smaller bright yellow flowers, which are produced from February or March till August. It has been tried in the open air in several places, and found to stand the winters



of London with very little protection; sometimes, when neglected, being killed down to the ground, but shooting up again the following spring. One in our garden at Bayswater has stood since 1831; and one in the garden at Bicton, near Exeter, since 1833.

3. *Hibbertia grossulariifolia* Sal, Bot. Mag., t. 1218. The Gooseberry-leaved *Hibbertia*.—This is an elegant trailer, from New Holland, which has been in cultivation since 1816. The leaves are nearly round, beautifully notched; and the flowers are on peduncles opposite the leaves, and of a bright yellow. It is rather a procumbent than a climbing plant; but thrives well against a wall, or on rockwork, during the summer months.

Other Species of *Hibbertia*, from New Holland, are in cultivation in Britain; and upwards of a dozen, which have been described, remain to be introduced; all of which, there can be no doubt, will stand our British winters with little protection, and produce a fine show of their brilliant yellow flowers during the summer months.

CHAP. V.

OF THE HARDY LIGNEOUS PLANTS OF THE ORDER ANONACEÆ.

The characteristics of this order assimilate most to those of *Magnoliaceæ*, and those distinctive of it from that order are: anthers with an enlarged four-cornered connectivum, which is sometimes nectariferous; albumen pierced by the substance of the seed-coat; leaves without stipules, conduplicate in the bud; properties aromatic. (*Lindl. Introd. to N. S.*, and *Don's Mill.*) The leaves of *Magnoliaceæ* are involute in the bud; and, perhaps, they are generally less obviously feather-nerved than those of *Anonaceæ*. The hardy species of this order are included in the genus *Asimina* Adans., formerly *Anona* L., and are natives of North America.

GENUS I.

*ASIMINA* Adans. THE ASIMINA. *Lin. Syst. Polyándria Polygýnia.*

Identification. Adans. Fam., 2. p. 365.; Dec. Prod., 1. p. 87.; Don's Mill., 1. p. 91.

Synonymics. *Annona* L.; *Orchidocarpum* Mx.; *Porcêluc* sp. Pers.; Custard Apple; Asiminer, and Anone, Fr.; Flaschenbaum, Ger.

Derivations. *Asimina* is Latinised from a word of Canadian origin, the meaning of which is not known. *Orchidocarpum* was, it is probable, intended to express a likeness between the figure of the fruit, and that of some species of *Orchis*. *Porcèlia* is a name given by Ruiz, in honour of Antonio Porcel, a Spanish promoter of botany. *Anona* is a South American word, that signifies a mess, or dish of food, to be eaten with a spoon. Linnæus, in applying this word, says, "anona [spelling it with two ns] and guanabannus are barbarous words; that the sound, however, may be kept, I name it Annona, on account of the fruit which is so grateful to the natives." (*Hort. Cliff.*) This word, guanabannus, is a synonyme to *Anona muricàta*, a stove plant. The German name, Flaschenbaum, flask tree, is given from the shape of the fruit.

Gen. Char. *Calyx* 3-parted. *Petals* 6, spreading, ovate-oblong, inner ones smallest. *Anthers* numerous, nearly sessile. *Ovaries* many, but for the most part only 3, ovate or oblong. *Carpels* the same number as the ovaries, baccate, sessile. *Seeds* many, disposed in a single or double row. (*Don's Mill.*, i. p. 91.)—Low trees or shrubs, deciduous, with white or purplish flowers, and fruit about the size of small plums. Rather tender, and difficult of culture.

Description. The plants, in their native countries, are shrubs or low trees, varying from 2 ft. to 30 ft. in height. In this country they are, for the most part, shrubs; though there are specimens of *A. triloba*, near London, in the Hammersmith Nursery, and at Purser's Cross, 10 ft. high. All the species require peat soil, and they are only propagated from imported seeds.

Geography, History, &c. The species are found in Virginia, Georgia, Carolina, and Florida; generally in shady places on the margins of woods, and almost always in sandy soil. They have been introduced at different periods, from 1736 to 1820.

§ 1. *A. PARVIFLO'RA* Dun. The small-flowered *Asimina*.

Identification. Dun. Mon. Anon.; Dec. Prod., i. p. 87.; Don's Mill., i. p. 91.

Synonymes. *Porcèlia parviflora* Pers.; *Orchidocarpum parviflorum* Mx.

Engravings. Dunal Monog., p. 82. t. 9.

Spec. Char. Leaves cuneate-obovate, mucronate; under surface, as well as branches, covered with brown pubescence. Flowers sessile; outer petals scarcely twice as long as the calyx. (*Don's Mill.*, i. p. 91.) This is a deciduous shrub, from 2 ft. to 4 ft. in height. The flowers have the outside of the calyx and corolla clothed with brownish tomentum; and the inside of the petals is of a dark purple colour. The berries are aggregate, sessile, fleshy, of the size of a small plum.

Geography, History, &c. Found in Virginia, Georgia, and Carolina, in shady woods near rivers and lakes; flowering in April and May. It was introduced into England in 1806, but is little known either in botanic gardens or nurseries. In New York plants are 1 dollar each.

§ 2. *A. TRI'LOBA* Dun. The three-lobed-calyxed *Asimina*.

Identification. Dun. Monog.; Dec. Prod., i. 87.; Don's Mill., i. 91.

Synonymes. *Annona triloba* L., and Mx. in Arb.; *Porcèlia triloba*, Pers.; *Orchidocarpum arietinum* Mx. Bor. Am.; the Papaw, Amer.; *Asiminier* de Virginie, and *Annone à trois Lobes*, Fr.; dreylappiger (three-lobed) Flaschenbaum, Ger.

Engravings. Mill. Icon., i. t. 35.; Du Ham., 2. t. 25.; Mx. Arb., 3. t. 9.; E. of Pl., 7927.; and our fig. 39.

Spec. Char., &c. Leaves oblong-cuneated, acuminate, and as well as the branches, smoothish. Flowers on short peduncles; outer petals roundish-ovate, four times longer than the calyx. (*Don's Mill.*, i. p. 91.) A small tree, densely clothed with long leaves, lying over one another in such a manner as to give a peculiarly imbricated appearance to the entire plant. The flowers are campanulate and drooping, and appear before the leaves; the outer petals are purple, and vary in colour in different plants; in some being very dark, and in others light, inclining to yellow. The berries are large, yellow, ovate, oblong, and eatable. They contain a yellow pulp, of a sweet luscious taste, in the middle of which lie, in two rows, 12 seeds, divided by as many thin membranes. All parts of the tree have a rank, if not a fetid, smell; and the fruit is relished by few persons except negroes. The fruit ripens in America in the beginning of August, and is about 3 in. long and 1½ in. thick, oval, irregular, and swelling into inequalities.

Geography, History, &c. Michaux did not observe this tree north of the Schuylkill river; and it appears to be unknown, or extremely rare, in the low and maritime part of the southern states. It is not uncommon in the bottoms of the rivers which stretch along the middle states; but it is most abundant in the rich valleys intersected by the western waters; where, at intervals, it forms thickets exclusively occupying several acres. In Kentucky and the western part of Tennessee, it is sometimes seen, also, in forests where the soil is luxuriantly fertile; of which its presence is an infallible proof. In these forests it attains the height of 30 ft., with a trunk 6 in. or 8 in. in diameter, though it generally stops short of half this height. (*Michaux.*)

This species was imported to England, under the name of *Anona triloba*, by Peter Collinson, in 1736; and it has since become known in the principal botanic gardens of Europe, and procurable in first-rate nurseries. Miller mentions that the largest plant he had seen was in the Duke of Argyll's garden at Whitton. (See p. 57.) The largest tree that we have heard of is that already mentioned, at Purser's Cross; where, some years ago, a tree of about the same size, since dead, ripened fruit. The plants are always raised from seeds; and they seldom produce shoots exceeding 5 in. or 6 in. in length: hence a plant, in ten years, does not reach above 3 ft. or 4 ft. in height; and will not flower till of 15 or 20 years' growth.

Properties, Uses, &c. The fruit in America is never brought into the markets, and is sought in the woods only by children. A spirituous liquor has been made from it, but it is of little worth. The wood is spongy, extremely soft, destitute of strength, and applicable to no use in the mechanical arts. In England, it may be considered as a curious, slow-growing, deciduous shrub, well deserving a place in gardens, but which ought always to be isolated, and at some distance from rapid-growing plants. Relatively to growth, it may be placed near *Dica palustris*, some of the daphnes, or *Illicium parviflorum*. Plants, in the London nurseries, are 2s. 6d. each, and seeds 1s. an ounce; at Bollwyller, 5 francs a plant; and in New York, 40 cents.

3. *A. PYGMÆA* *Dun.* The Pygmy Asimina.

Identification. *Dun. Monog.*; *Dec. Prod.*, l. 87.; *Don's Mill.*, l. 92.

Synonymes. *Annona pygmæa* *Bartr.*; *Orchidocarpum pygmæum* *Mr. Fl. Bor. Amer.*; *Porcilia pygmæa* *Pers.*

Engravings. *Bartr. Trav.*, p. 21. t. 1.; *E. of Pl.*, 7932; and our fig. 40.

Spec. Char., &c. Stem suffruticose. Leaves oblong-linear, cuneate, blunt, and, as well as the branches, smooth. Flowers on short peduncles. Outer petals obovate-oblong, much larger than the calyx. (*Don's Mill.*, i. p. 92.) A little shrub, hardly 1 ft. high, with twiggly branches, and long, cuneate, narrow leaves. Outer petals much larger than the inner ones, and all white. Found in Georgia, Florida, and Carolina, in sandy fields. Flowers white. April and June.

Geography, History, &c. We have never seen this species and know nothing more of its history than what is above stated. By the catalogues it appears to have been introduced into England in 1812, and, probably, is since lost. In New York, plants are 1 dollar 50 cents each.

4. *A. GRANDIFLORA* *Dun.* The large-flowered Asimina.

Identification. *Dun. Mon.*; *Dec. Prod.*, l. p. 87.; *Don's Mill.*, l. p. 92.

Synonymes. *Annona grandiflora* *Bartr.*; *A. obovata*, *Willd.*; *Orchidocarpum grandiflorum* *Mr. Fl. Bor. Amer.*; *Porcilia grandiflora* *Pers.*; *Asiminer à grandes Fleurs* *Bon. Jard.*

Engravings. *Dun. Mon.*, t. 11.; *Bartr. Trav.*, t. 2.

Spec. Char. Leaves cuneate-obovate, obtuse; under surface, as well as the branches, clothed with brown pubescence. Flowers sessile; outer petals

39



40



obovate, much larger than the calyx. (*Don's Mill.*, i. p. 92.) A small smooth-branched shrub, with white flowers, very large for the size of the plant; the outer petals are larger than the inner ones; the berries are smooth, and oblong-obovate. Height 2 ft.

Geography, History, &c. Native of Georgia and Florida, in sandy woods and shady places; and brought to England in 1820. It is still rare, or, rather, scarcely to be met with. It may ultimately turn out that these four alleged species are only varieties of one species, modified by local circumstances. At all events, one of them (*A. triflora*) is quite sufficient in a general collection, to give a correct idea of the genus.

CHAP. VI.

OF THE HALF-HARDY LIGNEOUS PLANTS OF THE ORDER SCHIZANDRA`CÆÆ.

SOME of the genera of this order have been referred to Menispermæcæ, and some to Anonæcæ; we introduce it here, in order to notice a beautiful ligneous climber, *Schizandra*.

Identification. *Don's Mill.*, i. p. 101.; *Blum. Bijdr. Fl. Ind. ex Schlecht. in Linnæa*, i. p. 497. obs. *Synonymes.* Part of Menispermæcæ and part of Anonæcæ with Dec.; Anonæcæ § *Schizandra*; *Lindley's Key*, p. 46.

GENUS I.



SCHIZANDRA *Michx.* THE SCHIZANDRA. *Lin. Syst. Monæ'cia Pentândria.*

Identification. *Michx. Fl. Bor. Amer.*, 2. p. 18.; *Dec. Syst.*, 1. p. 548.; *Don's Mill.*, 1. p. 101.

Deriv'ion. From *schizō*, to cut, and *andrōs*, a man; stamens cleft.

Gen.'char. Flowers monoecious. *Sepals* 9, in a ternary order. *Petals* nôbe. *Male flowers* with 5 others, which are joined at the apex; *female* ones with an indefinite number of ovaries. *Berries* disposed in spikes along an elongated receptacle. (*Don's Mill.*, i. p. 101.)—A deciduous climber.

± 1. SCHIZANDRA COCCI'NEA *Michx.* The scarlet-flowered Schizandra.

Engravings. *Michx. Flor. Bor. Amer.*, 2. t. 47.; *Sims, Bot. Mag.*, t. 1413.; *Encyc. of Pl.*, 13259.; *Don's Mill.*, f. 26.; and our fig. 41.

Spec. Char., Description, &c. Leaves alternate, oval-lanceolate, pointed at both ends, rarely toothed, of a beautiful green, smooth above and pale beneath, petiolated. Flowers scarlet, disposed in spikes in the axils of the leaves. A climbing, deciduous, half-hardy shrub, found in shady woods in Georgia and Florida, and also in Carolina. It flowers in June and July, and was introduced into England in 1806. It is generally treated as a green-house plant; but it stood out through the winters of 1832, 1833, 1834, and 1835, in the garden of the Horticultural Society, trained against a wall, and very slightly protected. It forms a most desirable ornament in the summer season, and should have a place against every conservative wall. It prefers a light sandy soil, and is easily propagated by ripened cuttings, in a pot of sand, placed under a hand-glass. Price, in London, 5s.; at Bollwyller, ?; and in New York, 75 cents.

41



App. i. Anticipated Additions to the Hardy Species of Schizandra`cææ.

Sphaerostema grandiflorum, and other species from Nepal, commonly included under Menispermæcæ (see p. 173.), but properly belonging to this order, may possibly be found half-hardy; as may *Kadsura japonica*, which, as the name implies, is a native of Japan.

CHAP. VII.

OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE ORDER
MENISPERMA'CEÆ.

DISTINCTIVE Characteristics. Thalamiflorous (*H. B.*). Sexes, in most, dioecious; in the rest, monœcious or polygamous. Sepals and petals similar; the latter not present in some. Stamens monadelphous, or rarely free; equal in number with the petals, and opposite to them, rarely double that number, or fewer. Ovaries, in some, numerous, each one-styled, all somewhat connected at the base; or, in others, only one, crowned with many styles, and many-celled, and, therefore, consisting of many carpels grown together, very rarely, one-celled, and this, most likely, by abortion. Fruit, in most, baccate or drupaceous, one-seeded or many-seeded, oblique or lunulate, compressed, with the seeds of the same form. Embryo curved or peripheric. Albumen none, or very sparing and fleshy. (*Don's Mill.*, i. p. 102., with adaptation.)—Climbing or twining shrubs, mostly natives within the tropics, with alternate, stalked, usually cordate or peltate, simple, rarely compound leaves, always with the middle nerve terminating in an awn or point; destitute of stipules. Flowers small; in most species, in axillary racemes. (*Don's Mill.*, i. p. 102.) The species in British gardens are included in the genera *Menispermum* and *Cœculus*, and are natives of North America and Dahuria. They are all of the easiest culture, and are propagated by dividing the root, or by cuttings.

GENUS I.

MENISPERMUM L. THE MOONSEED. *Lin. Syst.* Diœ'cia Dodecándria.

Identification. Tourn.; Dec. Prod., l. p. 102.; *Don's Mill.*, l. p. 112.

Synonymes. Ménisperme, *Fr.*; Mondsaaime, *Ger.*

Derivation. From *mène*, the moon, and *spërma*, a seed; from the seeds being crescent-shaped.

Gen. Char. *Sepals and petals* disposed in a quaternary order, in two or three series. *Male flowers* with 16 to 20 stamens; *female flowers* with 2 to 4 ovaries. *Drupe* baccate, roundish-kidney-shaped, 1-seeded.—Climbing shrubs, with alternate, peltate or cordate, smooth leaves. Peduncles axillary or supra-axillary. Male and female peduncles rather dissimilar. Flowers small, greenish-white. (*Don's Mill.*, i. p. 112.)

♁ I. M. CANADÉ'NSE. The Canadian Moonseed.

Identification. *Lin. Sp.*; Dec. Prod., l. p. 102.; *Don's Mill.*, l. p. 112.

Synonymes. *M. canadense* var. α *Lamarck*; *M. angulatum Moench*; Ménisperme du Canada, *Bon.*

Jard.; Canadischer Mondsaaime, *Ger.*

Engravings. Schkuhr. II., 3. t. 337.; *Lam. Dict.*, t. 824.; and our *fig.* 42.

Spec. Char. Leaves peltate, smoothish, somewhat cordate, roundish-angular; angles bluish, terminal one abruptly awned, mucronate. Racemes solitary, compound. Petals 8. (*Don's Mill.*, i. p. 112.) A twining shrub, with thick woody roots, and numerous very slender shoots, which rise to the height of 12 ft. or 14 ft., but which, though ligneous, never attain any considerable diameter, and are not of many years' duration. The stem twines in a direction contrary to the sun's apparent motion, and is smooth and even, having more the appearance of a herbaceous plant, than of a shrub. and the berries black.



The flowers are small,

Variety.

‡ *M. c. 2 lobatum* Dec. *The lobed-leaved Canadian Moonseed. M. virginicum* L. — This variety is distinguished by the angles of the leaves being acutish, and the flowers of a greenish white. Figured in *Dill. Elth.*, t. 178. fig. 219.

Geography, History, &c. Found in North America, among bushes, on the banks of rivers, and on fertile declivities, from Canada to Carolina; and also indigenous in Siberia. It was cultivated, in 1713, by Bishop Compton; and is not unfrequent in British botanic gardens, and in our principal nurseries. It will grow in any free, deep, and rather moist soil; and, as it sends up numerous shoots from its thick woody roots, it is easily propagated by dividing them, or by layers made in autumn, which will root in one year. Both the male and female plants are in Lee's Nursery; and the male of the variety *M. c. lobatum* is in the garden of the London Horticultural Society. Price, of plants in the London nurseries, 1s. 6d. each; at Bollwyller, 90 cents; in New York, 25 cents.

‡ 2. MENISPERMUM DAÛRICUM Dec. *The Daürian Moonseed.*


Identification. Dec. Prod., 1. p. 102.; Don's Mill., 1. p. 112. *Synonymæ.* *Trilophus Ampelisàgia* Fisch.; *M. canadense* var. β Lam. *Engravings.* Deless. Icon., 1. t. 100. and our fig. 43.

Spec. Char. Leaves peltate, smooth, cordate, angular; angles acute, terminal one acuminate hardly mucronate. Racemes in pairs, capitate. (*Don's Mill.*, 1. p. 112.) A twining shrub, resembling *M. canadense*, but smaller in all its parts, and, probably, only a variety of that species. Flowers yellowish. June and July. 1818. Found in Daüria, on rocky hills near the river Chilca, and said to be introduced into England in 1818; but we have never seen it.



‡ 3. *M. SMILA'CINUM* Dec. *The Smilax-like Moonseed.*

Identification. Dec. Syst., 1. p. 541.; Don's Mill., 1. p. 112. *Synonymæ.* *Cissampelos smilacina* Lin. *Engravings.* Jacq. Icon., t. 629.; Catesb. Carol., 1. t. 51.; and our fig. 44.



Spec. Char. Leaves peltate, smoothish, cordate-roundish, bluntly angular, under surface glaucous. Racemes simple. Petals 4. (*Don's Mill.*, 1. p. 112.) A climbing shrub, with slender stalks, and leaves resembling those of the common ivy. The flowers, which appear in July and August, are white, and the berries are red, about the size of small peas, and grow in clusters. Found in Carolina by Catesby, and first described by him. It was introduced into Britain in 1776, by Dr. Hope, then professor of botany at Edinburgh. The plant is rather scarce in British gardens; and, when it is met with, it is generally in a greenhouse; though there can be little doubt of its being half-hardy.

GENUS II.



‡ COCCULUS Bauh. *THE COCCULUS. Lin. Syst. Dia`cia Hexàndria.*

Identification. Bauh. Pin., 511.; Dec. Prod., 1. p. 96.; Don's Mill., 1. p. 104. *Synonymes.* *Menispermum* L.; *Weudlândia* Willd.; *Andróphilax* Wendl. *Derivation.* From *coccus*, the systematic name of cochineal, which is applied to this genus on account of the greater number of the species bearing scarlet berries.

Gen. Char. *Sepals* and *petals* disposed in a ternary order, in 2, very rarely in 3 series. *Male flowers* with 6 free stamens opposite the petals; *female* ones with 3 or 6 carpels. *Drupe* baccate, 1 to 6, usually obliquely reniform, somewhat flattened, 1-seeded. *Cotyledons* distant. (*Don's Mill.*, 1. p. 104.)

Description. This is a genus of climbing or twining shrubs, with peltate, cordate, ovate or oblong, entire, rarely lobed, leaves. Peduncles axillary,

rarely lateral; those bearing male flowers are usually many-flowered; but those bearing female flowers are few-flowered, either free from bracteas, or furnished with very small ones. The berries of many of the species of this genus are often made into a paste, and used in their native countries to intoxicate fish and birds, &c., in order to take them; and it is said that brewers use them to give their ale and porter an intoxicating quality. (*Don's Mill.*, i. p. 104.) The species are chiefly tropical, and only one that is hardy has yet been introduced into the British gardens.

3. *COCCULUS CAROLINENSIS* Dec. The Carolina Cocculus.

Identification. Dec. Prod., 1. p. 98.; *Don's Mill.*, 1. p. 107.

Synonymes. *Menispermum carolinum* *Lin.*; *Wendlandia populifolia* *Willd.*, *Pursh.*, and *Dill.*; *Andropilax scandens* *Wendl.*; *Baumgärtia scandens* *Moench.*; *Ménisperme de la Caroline*, *Fr.*; *Carolinischer Mondsaame*, *Ger.*

Engravings. *Dill. Elth.*, 223. t. 178. f. 219.; *Wendl. Obs.*, 3. t. 16.; and our *fig.* 45.

Spec. Char., &c. Leaves cordate or ovate, entire, obtuse, and somewhat 3-lobed; under surface velvety pubescent. Male racemes floriferous from the base, female ones 3-flowered. (*Don's Mill.*, i. p. 107.) A twining shrub, a native of Carolina, Georgia, and Florida, in woods and hedges, as the black bryony is in England. The flowers are diœcious, but, according to *Wendland*, often hermaphrodite. Though ligneous in its native country, in cold countries it is often herbaceous or subherbaceous. The flowers, which appear in June and July, are greenish; and the berries, when ripe, are of a red colour. It was introduced into England in 1759, and is not uncommon in botanic gardens and the principal nurseries. Price, in London, 1s. 6d. a plant; at *Bollwyler*, ?; in New York, 1 dollar.



App. i. *Anticipated Menispermæcæ.*

In p. 175. are enumerated some genera and species belonging to this order which are natives of the Himalaya; and in p. 176. some that are natives of China and Japan, which, it is considered, would be found half-hardy in our gardens.

CHAP. VIII.

OF THE HARDY LIGNEOUS PLANTS OF THE ORDER *BERBERACEÆ.*

This order is distinguished from other thalamiflorous ones by the following traits. Sepals usually 6, in two whorls, deciduous, and furnished with petal-like scales on the outside. The petals are equal in number with the sepals; and the stamens equal in number with the petals, and opposite to them. The anthers "open by reflexed valves; that is to say, the face of each cell of the anther peels off except at the point, where it adheres as if it were hinged there;" a structure so remarkable, *Dr. Lindley* observes, as to be "found in no European plants except *Berbericææ*"

and the laurel tribe." (*Penny Cyc.*, vol. iv. p. 259.) The genera containing the hardy species are two, *Berberis* and *Mahonia*. They are shrubs, or low trees, inhabitants of Europe, Asia, and of North and South America; but they are not met with in the interior of Africa, or the South Sea Islands. They are usually found in the temperate zones; but some of them inhabit high mountains within the tropics. The seeds are very tenacious of life, and, being small, and easily conveyed from one country to another, a number of new species have recently been introduced from Nepal and South America. The wood of some of the species is used for dyeing yellow; and the more common have been admitted into the *materia medica*, from the days of Galen to the present time, on account of their bitter and astringent properties. All the species are ornamental, and those of them which are evergreen eminently so. They are all readily propagated by seeds, which most of them ripen in England, and also by side suckers and root suckers, which almost all the species produce in abundance. The fruit is generally edible, and abounds in the malic acid. The genera and species of this order have recently been arranged and described, in a masterly manner, by Dr. Lindley, in the *Penny Cyclopædia*; from which article, from *Don's Miller*, and from our own observations, we have drawn up this chapter. The distinctive characters of the two genera of *Berberis* and *Mahonia* are as under:—

BERBERIS. Sepals 6, furnished on the outside with 3 scales. Petals 6, with 2 glands on the inside of each at the base. Stamens toothless. Berries 2—3-seeded. (*Don's Mill.*, i. p. 114.) Flowers in simple, mostly pendulous, racemes; in some species solitary. Leaves undivided.

MAHONIA. Sepals 6, furnished on the outside with 3 scales. Petals 6, without glands on the inside. Stamens furnished with a tooth on each side, at the top of the filament. Berries 3—9-seeded. (*Ibid.*) Flowers in erect racemes, that are disposed several together in a panicle. Leaves pinnate.

GENUS I.



BERBERIS Lin. THE BERBERRY. *Lin. Syst.* Hexándria Monogýnia.

Identification. *Lin. Gen.*, 442; *Dec. Prod.*, i. p. 105; *Don's Mill.*, i. p. 114.

Synonyms. Piperidge Bush; E'pine vinette, *Fr.*; Berberitze, *Ger.*

Derivations. *Berberis* is the Arabic word used for this plant by Avrrhoeos and other writers on medicine; but some persons derive the name from the Greek word *berberi*, signifying a shell, from the leaves of the common species having a hollow surface. Bochart says that the word *Berberis* is derived from the Phœnician word *barar*, which signifies shining like a shell, from their shining leaves. Gerard says that the word *Berberis* is a corruption of *amyrberis*, the name given to the plant by Avicenna. Du Hamel says that *Berberis* is derived from an Indian word signifying mother of pearl. Piperidge bush, or piprage tree, Gerard says, is Dr. Turner's name for the plant, and it is still given to it in Cambridgeshire. E'pine vinette signifies the acid, or sorrel, thorn, from the taste of the fruit and leaves.

Gen. Char. Sepals 6, guarded on the outside by 3 scales. Petals 6, with 2 glands on the inside of each. Stamens toothless. Berries 2—3-seeded. Seeds 2, rarely 3, laterally inserted at the base of the berries, erect, oblong, with a crustaceous coat and fleshy albumen. Cotyledons leafy, elliptical. Radicle long, capitate at the tip. (*Don's Mill.*, i. p. 114.) *B. heterophýlla* *Juss.* has toothed stamens.

Description. The species are all shrubs of from 2 ft. or 3 ft. to 18 ft. or 20 ft. in height, in a wild state; some of them attaining the height of 30 ft. in gardens. They all throw up numerous side suckers, and the stronger-growing species, if these were carefully removed, might be formed into very handsome small trees. In all the species the flowers are yellow. The fruit is generally red; but in some species it is black or dark purple, and in some varieties of the species it is white or yellow: it is always acid, and more or less

astringent. "The spines of the common berberry are a curious state of leaf, in which the parenchyma is displaced, and the ribs have become indurated. They, as well as all the simple leaves of ordinary appearance, are articulated with the petiole, and are therefore compound leaves reduced to a single foliole; whence the supposed genus *Mahonia* does not differ essentially from *Berberis* in foliage any more than in fructification." (*Lindley, Introd. to N. S.*, p. 31.) The species are generally thorny, and most of them flower freely in spring, bearing fruit abundantly in autumn.

The irritability of the stamens of the genus *Berberis*, and more particularly of those of the common berberry, of *B. canadensis*, *B. sinensis*, and, perhaps, of all the species the flowers of which expand, is a very remarkable property, which was first discovered by Kölreuter; probably from observing that the stamens were put in motion by the proboscis of insects extracting honey from the flowers. Sir James Edward Smith has given a copious account of this phenomenon in the *Phil. Trans.*, vol. lxxviii. p. 158., and the last, and most clear and concise, description of it will be found to be that by Dr. Lindley, under the article *Berberis*, in the *Penny Cyclopaedia*. The stamens, "when the filament is touched on the inside with the point of a pin, or any other hard instrument, bend forward towards the pistil, touch the stigma with the anther, remain curved for a short time, and then partially recover their erect position. This is best seen in warm dry weather. After heavy rain, the phenomenon can scarcely be observed, owing, in all probability, to the springs of the filaments having been already set in motion by the dashing of the rain upon them, or to the flowers having been forcibly struck against each other. The cause of this curious action, like that of all other vital phenomena, is unknown. All that has been ascertained concerning it is this, that the irritability of the filament is affected differently by different noxious substances. It has been found by Messrs. Macaire and Mareet, that, if a berberry is poisoned with any corrosive agent, such as arsenic or corrosive sublimate, the filaments become rigid and brittle, and lose their irritability; while, on the other hand, if the poisoning be effected by any narcotic, such as prussic acid, opium, or belladonna, the irritability is destroyed by the filaments becoming so relaxed and flaccid, that they can be easily bent in any direction. It is difficult to draw from this curious fact any other inference than this, viz. that in plants, as well as in animals, there is something analogous to a nervous principle, which is more highly developed in some plants, or in some organs, than in others." (*Pen. Cyc.*, iv. p. 260.) According to Sir J. E. Smith, the purpose of this remarkable function in the stamens of the berberry is evident. "In the original position of the stamens the anthers are sheltered from rain by the concavity of the petals. Thus, probably, they remain till some insect comes to extract honey from the base of the flowers, and, thrusting itself between the filaments, unavoidably touches them in the most irritable part; and thus the impregnation of the germs is performed." (*Phil. Trans.*, lxxviii. p. 158.) All the species are easily propagated by seed, which most of them produce in abundance; those which do not are readily increased by the removal of their side suckers, or by layers. They will grow in any soil, though they mostly prefer one that is calcareous.

Though the species of this genus are commonly treated as shrubs, and these shrubs, from their numerous side suckers, have, in general, a rough, inelegant appearance; yet there are some of them which may be formed into the most beautiful and durable small trees that can be introduced into gardenesque scenery. The common berberry, when pruned up to a single stem to the height of 8 ft. or 10 ft., and all suckers from the root, and all side buds from the stem removed the moment they appear, will form a fine orbicular head with the extremities of the branches drooping; and this pendulous appearance will increase with the age of the tree. Such a tree, covered, as it will be every year, with yellow blossoms in the beginning of summer, and with bright scarlet fruit in autumn, may rank in beauty and value with the low trees of the genera *Crataegus*, *Cotoneaster*, and *Amelanchier*.

A. *Leaves thin, deciduous. Flowers solitary.*

❖ 1. *B. SIBIRICA* Pall. The Siberian Berberry.

Identification. Pall. Fl. Ross., 2. p. 42.; Dec. Prod., 1. p. 108.; Don's Mill., 1. p. 117.; Pen. Cyc., 4. p. 260.

Synonymes. *B. altaica* Pall.; Vinettier de Sibérie, Fr.

Engravings. Pall. Fl. Ross., 2. t. 67.; and our fig. 46.

Spec. Char., &c. Spines 3—7-parted. Leaves lanceolate-obovate, ciliately serrated. Peduncles 1-flowered, shorter than the leaves. (*Don's Mill.*, i. 117.)—A small shrub, found in rocky places, on the hills and lower mountains of Altai, Siberia, &c., and introduced into England by Pallas in 1790. Pallas states that the priests of the Mongols, who also act as physicians, being taught by the Tunguti, use the bark of the trunk and the yellow pulpy matter of the root for various diseases; and that a decoction of the young twigs is sometimes applied with a pencil to the eyes as a charm. In British gardens this species is a low scrubby bush, seldom exceeding 2 ft. in height. Price, in London, 10s. 6d. each.

46



B. *Leaves thin, mostly deciduous. Flowers in Racemes.*

❖ 2. *B. VULGARIS* L. The common Berberry.

Identification. Lin. Sp., 472.; Dec. Prod., 1. p. 105.; Don's Mill., 1. p. 115.

Synonymes. Pimperidge Tree, Dr. Turner; Épine vinette, Fr.; gemeine Berberitze, Ger.

Engravings. Eng. Bot., t. 49.; Willd. Baum., t. 39.; E. of Pl., 4922.

Spec. Char. Spines 3-parted. Leaves somewhat obovate, ciliately serrated. Racemes many-flowered, pendulous. Petals entire. (*Don's Mill.*, i. p. 115.)

Varieties. These are numerous. Those recognised by De Candolle and G.

Don are as follows:—

- ❖ *B. v. 2 lutea.* The yellow-fruited common Berberry.—Fruit yellow.
- ❖ *B. v. 3 alba.* The white-fruited common Berberry.—Fruit white.
- ❖ *B. v. 4 violacea.* The violet-coloured-fruited common Berberry.—Fruit violaceous.
- ❖ *B. v. 5 purpurea.* The purple-fruited common Berberry.—Fruit purple, leaves narrow, hardly ciliated. *B. innominata* Kiehm.
- ❖ *B. v. 6 nigra.* The black-fruited common Berberry.—Fruit black; leaves oblong, ciliately serrated, serratures few. The fruit of this plant is said by Tournefort, who found it on the banks of the Euphrates, to be of delicious flavour.
- ❖ *B. v. 7 dulcis.* The sweet-fruited common Berberry.—Fruit red, somewhat less acid than that of the common berberry. Leaves of a bright shining green. Native of Austria.
- ❖ *B. v. 8 asperma.* The seedless Berberry.—Fruit destitute of seeds. Miller, and also Du Hamel, both say that suckers taken from this variety commonly produce fruit with seeds; that, as the tree grows older, the seeds become fewer, and that it is the age of the plant that at last causes the fruit to be seedless; in that case this plant must be considered more a variation than a variety. *B. v. asperma* is said by Du Hamel to produce the best fruit for preserving; and it is from it that the delicious *Confitures d'Épine vinette*, for which Rouen is so celebrated, are made. (*Nor. Dub.*, iv. p. 13.) Price, 2s. 6d. each.

All these varieties are in the garden of the London Horticultural Society.

Other Varieties. Dr. Lindley says, that "there is in the *Catalogues* a Canadian berberry, which appears to be nothing more than a common berberry, brought from North America; and also *Berberis daurica* and *altaica*, neither of which merits to be distinguished from *B. vulgaris*."


Description. In a wild state, the common berberry is seldom found higher than 4 ft. or 5 ft., but in a state of culture it may be grown to nearly 30 ft.

high. The stems are upright, and much branched towards the top; smooth, slightly grooved, covered with a whitish or ash-coloured bark, which is yellow within, and they have a large white pith. The main stem soon becomes so surrounded by side suckers as to be concealed by them; so that, even where the height of the plant is that of a tree, its character is still that of a bush. The blossoms are, in general, abundant, and produce a fine appearance in April and May; their smell is offensive when near, but not disagreeable at a short distance. The tree will live for two or three centuries, without increasing much in size. The wood is hard and brittle, of a yellow colour, but little used except for dyeing. The rate of growth, when the plant is young, is rapid; and, in consequence, in five or six years it will attain the height of 7 ft. or 8 ft.; but it grows slowly afterwards, unless the suckers are removed from it as they are produced. It is seldom seen above 10 ft. high; but there are examples of trees of it 30 ft. high, probably of 30 years' growth.

Geography and History. Found wild in most parts of Europe, and in many parts of Asia and America; in the warmer parts of those last countries, on mountains; in the colder parts of Europe in plains, as in Norway, near Christiania. The berberry is found on Mount Lebanon, and on Mount Etna; in which last situation it becomes a low shrub, in the last zone of vegetation, at the height of 7500 French feet above the level of the sea. In England it is found in indigenous woods and hedges, more especially on calcareous soils. It is so common in the hedges of Saffron Walden, in Essex, where corn grows frequently quite up to the hedge, that Professor Martyn refers to this circumstance, as a proof that the prejudice respecting its originating the mildew on wheat is unfounded. It is indigenous in Scotland and Ireland, but not very common in those countries. The plant is mentioned by Pliny; and, among moderns, seems first to have been recorded by Bauhin in his *Pinar.*, and subsequently by all the writers on plants, under different names, till the time of Ray, who first called it *Berberis*; which name was afterwards adopted by Linnæus, and by all the botanists since his time.

Properties and Uses. The inner bark both of the stems and roots affords a yellow dye. The leaves are agreeably acid, and, according to Gerard, were used in his time "to season meat with, and instead of a salad, like sorrel." The berries are so acid, that birds seldom touch them. They are not eaten raw, but are excellent when preserved with sugar in syrup, or candied. They are also made into jelly and rob, both of which are not only delicious to the taste, but extremely wholesome; and they are pickled in vinegar, when green, as a substitute for capers. In some countries in the north of Europe, the berries are used instead of lemon for flavouring punch, &c.; and when fermented it produces an acid wine, from which tartar is procured by evaporation. They are also in general use for garnishing dishes. Medicinally, the berries, leaves, and roots are powerfully acid and astringent; the bark is purgative and tonic; and the berries, when bruised and steeped in water, make a refreshing drink in fevers. The astringent principle is so abundant in the bark, that it is used for tanning leather in Poland; and it dyes it of a fine yellow at the same time. A decoction of the bark is said to make a good gargle to strengthen the throat and gums. The plant is cultivated in gardens as a fruit tree or fruit shrub; and the variety, or rather variation, in which the seeds are said to be wanting, and that in which the fruit is sweet, are recommended in preference. The plant makes an excellent hedge; but there exists a prejudice against it among agriculturists, from its supposed influence in producing blight, or mildew, on the corn adjoining it. This opinion, though totally unfounded, is of unknown antiquity. It appears to have been first considered as an erroneous prejudice by Du Hamel, who assures us that it is totally void of foundation; and Broussonet and other botanists subsequently proved the fact; but the most scientific refutation of the error was given by Dr. Greville, in his *Scottish Cryptogamic Flora*. In that excellent work Dr. Greville has shown that the mildew which attacks the berberry (*Æcidium Berberidis* Pers., fig. 47.) is quite different from any of the *Fungi* which are found on

corn. The berberry mildew, when magnified, is found to consist of a number of small orange cups, with a white film over each. When ripe these films burst, and the tops of the cups assume a ragged uneven appearance, in which state they look like white *Fungi*. The cups are filled with innumerable little cases, containing seeds, or sporules, and these constitute the bright orange powder that is seen on the leaves and flowers of the common berberry. "Among the many beautiful objects that are to be met with in the lower and more imperfect tribes of plants," Dr. Lindley observes, "it is difficult to find one more worthy of an attentive examination than the *Æcidium Berberidis*." The blight on corn is generally a species of *Urædo*, and does not correspond in botanical characters with the *Æcidium*.



47

Propagation and Culture. The original species is propagated in the nurseries by seeds, and the varieties by suckers. For ordinary purposes, no plant requires less culture; but, to produce large fruit, it should be planted in a deep, well manured, somewhat calcareous soil, and be constantly freed from side suckers. The racemes of the blossoms, also, should be thinned out, in order by reducing the number of bunches of fruit, to increase its size. When the berberry is intended to become an ornamental tree, it should be trained with a straight stem to the height of 8 ft. or 10 ft., and then suffered to branch out, thinning out the shoots where necessary, and destroying every sucker as it appears. So treated, it forms a singularly beautiful small tree, of great duration.

Diseases, &c. The common berberry is very subject to the mildew, *Æcidium Berberidis* Pers., before described.

Statistics. The berberry is to be found in all European gardens that pretend to have a complete collection of fruit trees, and in most shrubberies. The original species is to be procured in all nurseries, and the varieties in some of them. Seedlings of the species, in the London nurseries, are 8s. a thousand; and transplanted plants 25s. a thousand: at Bollwyller, the varieties are a franc each; plants of the species, 50 cents each: in New York, ?

3. *B. CANADENSIS* Mill. The Canadian Berberry.

Identification. Pursh's Fl. Amer., Sept., 1. p. 219.; Nutt. Gen. Amer., 1. 210.; Dec. Prod., 1. p. 106.; Don's Mill., 1. p. 115.

Synonymes. *B. vulgaris* Mx. Fl. Bor. Amer. 1. p. 205.; *B. vulgaris* var. *canadensis* Martyn's Mill., No. 1.

Engravings. Hayne Abbild., t. 63.; and our fig. 48. after that author.

Spec. Char., &c. Spines 3-parted. Leaves obovate-oblong, remotely serrated, upper ones nearly entire. Racemes many-flowered, nodding. (*Don's Mill.*, i. p. 115.) A shrub, or low tree, with yellow flowers, from April to June. Cultivated in 1759. Height 5 ft. It is found in North America, on fertile hills and among rocks, especially in the Alleghany Mountains, from Canada to Carolina, and also in Tennessee. The berries are said by Pursh to be more fleshy and less acid than those of *B. vulgaris*. The same opinion prevails in the United States as in England, as to the berberry producing mildew on wheat. From the appearance of this alleged species in the garden of the London Horticultural Society, we are induced to consider it only a variety of *B. vulgaris*. De Candolle thinks it for the most part intermediate between *B. vulgaris* and *B. chinensis*. Plants, in London, are 1s. 6d. each; at Bollwyller, 1 franc; and in New York, 15 cents.



48

4. *B. EMARGINATA* Willd. The emarginated-petaled Berberry.

Identification. Willd. Enum., 1. p. 395.; Dec. Prod., 1. p. 105.; Don's Mill., 1. p. 115.

Synonymes. Ausgerandete (serrated) Berberitze, Ger.

Engravings. Hayne Abbild., t. 62.; and our fig. 49.

Spec. Char., &c. Spines 3-parted. Leaves lanceolate-obovate, ciliate serrated.

Racemes scarcely pendulous, shorter than the leaves; petals emarginate. (*Don's Mill.*, i. p. 115.) A shrub closely resembling *B. vulgaris*, of which it is, doubtless, only a variety; but it is one half smaller in all its parts, and has the petals emarginate. It is found wild in Siberia, and was introduced into England in 1820. In the garden of the London Horticultural Society it has attained the height of 7 ft. in 10 years. Price, in the London nurseries, 2s. a plant; at Bollwyller, 2 francs; and in New York, ?.



5. *B. IBÉRICA* Stev. The Iberian Berberry.

Identification. Stev. and Fisch. in Litt.; *Don's Mill.*, i. p. 115.; and Lindl. in Pen. Cyc., 4. p. 261.
Synonymes. *B. vulgaris* ? v. *ibérica* *Dec. Syst.*, 2. p. 6.; *B. sinensis* *Wat.*
Engravings. *Dend. Brit.*, t. 26.; and *E. of Pl.*, 4928., as *B. sinensis*; and our fig. 50.

Spec. Char., &c. Spines simple, and 3-parted; leaves obovate-oblong, quite entire. Racemes many-flowered; petals entire. (*Don's Mill.*, i. p. 115.) A shrub closely resembling the common berberry, but, according to Dr. Lindley, readily distinguished from it by its smaller leaves, and its almost upright racemes. The berries are dark purple. It is a native of Iberia, whence it was brought to England in 1790. Height 5 ft.

6. *B. SINENSIS* Desf. The Chinese Berberry.

Identification. Desf. *Catal. Hort. P.*, 150.; *Dec. Prod.*, 1. p. 106.; *Don's Mill.*, 1. p. 115.
Synonymes. *B. vulgaris* *Thunb. Jap.*, 1. p. 146.

Spec. Char., &c. Spines 3-parted. Leaves oblong, obtuse, entire, or the lower ones a little toothed. Racemes many-flowered, nodding. (*Don's Mill.*, i. p. 115.) A shrub closely resembling *B. vulgaris*, but seldom growing more than 4 ft. or 5 ft. high. The berries are oval, of a deep red colour (*Dec.*), or, according to Dr. Lindley, of a dirty red; 1-2-seeded. It is a native of China, where it was found during Lord Macartney's embassy, between Peking and Gehol; and it was introduced into England in 1800. There are plants of it in the garden of the London Horticultural Society, and in the arboretum of Messrs. Loddiges. Dr. Lindley observes that it is more common in French than in English gardens, and that it most resembles *B. ibérica*.



7. *B. CRÉTICA* L. The Cretan Berberry.

Identification. *Lin. Sp.*, 472.; *Dec. Prod.*, 1. p. 106.; *Don's Mill.*, 1. p. 115.
Synonymes. *B. crética buxifolia* *Tourn.*; *Vinettie* de Crète, *Fr.*; *Cretische Berberitze*, *Ger.*
Engravings. *Fl. Græc.*, t. 242.; *Candian Berberry*.

Spec. Char., &c. Spines 3—5-parted. Leaves oval-oblong, entire, or somewhat serrated. Racemes 3—8-flowered, rather shorter than the leaves. (*Don's Mill.*, i. p. 115.) A low shrub, seldom exceeding 3 ft. or 4 ft. in height, with numerous suckers, forming a compact bush, densely covered with leaves intermixed with spines. The leaves are produced without any obvious order, and in their shape they resemble those of the narrow-leaved variety of the common box. The berries are ovate, black, 2-seeded, more astringent than acid; stigma on a very short style. It is a native of Crete, or Candia, of Cyprus, and also of Japan; and it has been cultivated in England since 1759; but, being a plant of no great show, it is not very common in gardens or nurseries. There are plants of this species in the arboretum of Messrs. Loddiges, and also in the arboretum of Messrs. Buchanan and Oldroyd, at Camberwell. It is also in the garden of the Horticultural Society. Dr. Lindley observes of it, that "it is a dwarf scrubby bush, looking like a starved specimen of the common berberry." Price, in London, 1s. 6d. a plant; at Bolwyller, 1 franc 50 cents; in New York, ?.

Variety.

♣ *B. c. 2 serratifolia* Poir. *The serrated-leaved Cretan Berberry.*—Leaves ciliate-serrated.

♣ *B. CRATÆGINA* Dec. *The Cratægus-like Berberry.*

Identification. Dec. Syst., 2. p. 9.; Don's Mill., 1. p. 116.

Spec. Char., &c. Spines simple. Leaves oblong, reticulated, hardly serrated. Racemes many-flowered, crowded, spreading, scarcely longer than the leaves. (*Don's Mill.*, 1. p. 116.) Native of Asia Minor. "Allied to *Berberis crética* and *sinensis*" (*Dec.*): said to be like *B. vulgaris* by *G. Don*. "Described by De Candolle from specimens collected in Asia Minor. Young plants, of what is said to be this species, are in the gardens, but they have not yet flowered." (*Lindl.*) Where we meet with many doubts, we are always disposed to simplify; and, from the geography of this species, we think it highly probable that it will turn out to be a mere variety of *B. vulgaris*.

C. Leaves leathery, evergreen or sub-evergreen. Flowers solitary or in Clusters.

♣ *B. DU'LCIS.* *The sweet-fruited Berberry.*

Identification. Swt. Brit. Fl.-Gard.; Lindl. Pen. Cyc., 4. p. 261.

Engravings. Swt. Brit. Fl. Gard. 2d ser., t. 100.; and our fig. 51.

Spec. Char., &c. Spines long, slender, simple, or 3-parted.

Leaves obovate obtuse, with or without a bristly point, quite entire, glaucous on the under side. Flowers solitary, on slender stalks, twice as long as the leaves. (*Lindl., Pen. Cyc.*, 4. p. 216.) This species, or alleged species, is not recognised by De Candolle or George Don: it is said by Dr. Lindley to be "a native of the south-western part of South America, from the Straits of Magellan to Valdivia, where it forms a small evergreen bush. The species has been some years in this country; but it is at present very rare." (*Pen Cyc.*, 5. p. 261.) There are plants in the Hort. Soc. Garden between 2 ft. and 3 ft. high; and it is to be procured at Low's Nursery, Clapton, where it flowered in March, 1831. It was sent to Mr. Low by Mr. Anderson, the collector attached to Capt. King's expedition, from the Straits of Magellan. In *Sweet's Flower-Garden*, where it is figured, it is said that, in its native country, "the fruit is used, both green and ripe, as we use gooseberries, for making pies and tarts, and preserves, for which it is most excellent. The berries are round and black, being about the size of a black currant, and are produced in great abundance. The flowers are very handsome, being of a bright yellow, and nodding: they make a very elegant appearance." (*Swt. Fl.-Gard.*, 2d s., i. t. 100.) It is quite hardy, and evergreen; but there is a deciduous variety, also possessed by Mr. Low, which, by some, is supposed to be a distinct species.



♣ *B. HETEROPHYLLA* Juss. *The various-leaved Berberry.*

Identification. Juss. in Poir. Dict., 8. p. 622.; Dec. Prod., 1. p. 108.; Don's Mill., 1. p. 117.; Lindl., Pen. Cyc., 4. p. 261.

Synonyms. *B. bicifolia* Forst.; *B. triscupidata* Smith.

Engraving. Hook Exot. Fl., 1. t. 14.; and our fig. 52.

Spec. Char., &c. Spines 3-parted. Leaves ovate-lanceolate, glabrous, some of them entire, others furnished with 3 pungent teeth. Pedicels solitary, 1-flowered, hardly longer than the leaves. Filaments toothed. (*Don's Mill.*, 1. p. 117.) This species Dr. Hooker describes as a shrub about 3 ft. in height, much branched, and the older branches covered with dark wrinkled

bark. The leaves clustered, and of two kinds; the old ones terminated with a sharp spinose point, and having a lateral spinule on each side, above the middle, and the younger ones being pale green, unarmed, and having their margins entire and softish. The old leaves are also quite rigid, dark green, and shining. The flowers are solitary, about the size of a pea, and of an orange-yellow colour. (*Exot. Fl.*, i. t. 14.) Dr. Hooker also observes, that this species "departs from the generic character of *De Candolle*, inasmuch as the calyx has no scales at its base, nor are the filaments destitute of teeth, for there are two most distinct ones just beneath the anther." This shrub is a native of the Straits of Magellan, where it was discovered by Commerson; but when, and by whom, it was introduced in our gardens, Dr. Hooker informs us, is not known. According to Dr. Lindley, it is "an inelegant bush, about 3 ft. high, bare of leaves, and having nothing but its rarity to recommend it." It is the *B. bicifolia* of English gardens.



☞ 11. *B. EMPETRIFOLIA* Lam. The Empetrum-leaved Berberry.

Identification. Lam. Ill., t. 253; Dec. Prod., 1. p. 107; Don's Mill., 1. p. 117; Pen. Cyc., 4. p. 261.
Engraving. Lam. Ill., t. 253. fig. 4.

Spec. Char., &c. Spines 3-parted. Leaves linear, quite entire, with revolute margins. Pedicels 1—2, 1-flowered. (*Don's Mill.*, i. p. 117.) According to Dr. Lindley, the leaves are collected in bundles in the axils of the spines, and the pedicels of the flowers are about as long as the leaves. "A very curious and pretty plant, found wild from the Cordilleras of Chili to the southern point of the American Continent, in subalpine woods. In general aspect it is much more like a heath than a berberry, seldom exceeding 2 ft. in height. It has been some years in the Horticultural Society's Garden, and is in Young's Nursery at Epsom, and in the Fulham Nursery, but is to be found in few others. It flowers in December, and is said, in *Sweet's Brit. Fl.-Gard.*, 2d series, t. 100., to have flowered at Low's Nursery, Clapton.

D. *Leaves leathery, evergreen or sub-evergreen. Flowers in Racemes.*

☞ 12. *B. FLORIBUNDA* Wall. The many-flowered Berberry.

Identification. Wall. MSS; Don's Mill., 1. p. 115; Pen. Cyc., 4. p. 261.

Synonyms. "Out of accidental variations of this species, and its mode of leafing and flowering, the spurious species called *B. alluinis* and *B. ceratophylla* have been constituted. By Dr. Wallich, in his distribution of the herbarium of the East India Company, *B. floribunda* has been mistaken for *B. aristata*." (*Pen. Cyc.*, 4. p. 261.)

Spec. Char., &c. Spines 3-parted and very stiff. Leaves oblong or oblong-lanceolate, nearly entire, or toothed in various degrees, sometimes very deeply and coarsely veined; flowers in long, loose, slender racemes. (*Pen. Cyc.*, iv. p. 261.) This species is supposed to grow about 10 ft. high in Nepal, and, as Dr. Lindley observes, is, "apparently, extremely common in the whole of the north of India, where it forms a tall bush, varying considerably in the form and size of the leaves, and in the degree in which they are toothed, but always well marked by its slender, pendulous, or erect racemes of flowers, which are much longer than the leaves, and in no degree corymbose. It is to be found occasionally in the more choice collections in this country." (*Pen. Cyc.*, iv. p. 261.)

☞ 13. *B. ASIATICA* Roxb. The Asiatic Berberry.

Identification. Roxb. in Dec. Syst., 2. p. 13; Dec. Prod., 1. p. 107; Don's Mill., 1. p. 116; Pen. Cyc., 4. p. 261.

Synonyms. *B. tinctoria* Lech.; the Raisin Berberry *Pen. Cyc.*
Engraving. Deless. Icon. sel., 2. t. 1.

Spec. Char., &c. Spines trifid, or simple. Leaves oval, cuneated or elliptical, mucronate, smooth, under surface glaucous, entire or spinulosely toothed.

Racemes short, many-flowered, corymbose, shorter than the leaves. Pedicels elongated, 1-flowered. Berries oval. (*Don's Mill.*, i. p. 116.) A vigorous-growing shrub, with numerous luxuriant side suckers, approaching in vigour those of *B. aristata*; but the leaves resembling those of *B. heterophylla*. It is a native of the East Indies and Nepal, and was introduced into England in 1820. *B. asiatica*, Royle observes, "is found on the Neelgherries, and was called, by M. Lechenault de la Tour, *B. tinctoria*, from the use to which it has been applied; and it has been proved by the experiments of M. Vauquelin to be inferior to few woods for dyeing a yellow colour." (*Royle's Illust.*, p. 63.) According to Dr. Lindley, the fruit is round, covered over with a thick bloom, and has altogether the appearance of the finest raisins. In the garden of the London Horticultural Society the plants of this species are about 7 ft. high, and they flower and fruit freely. They are easily distinguished from *B. aristata*, by their very short racemes. Plants 3s. 6d. each.

14. *B. DEALBATA* Lindl. The whitened-leaved Berberry.

Identification. Bot. Reg., t. 1750; Pen. Cyc., 4. p. 261.
Synonyme. *B. glauca* Hort.
Engraving. Bot. Reg., t. 1750; and our fig. 53.


Spec. Char., &c. Spines scarcely any. Leaves roundish, coarsely toothed, rather glaucous, white beneath. Racemes very short and compact, pendulous. (*Pen. Cyc.*, iv. p. 261.) A native of Mexico, whence it was introduced into England by the London Horticultural Society in 1830. "It is a tall slender evergreen bush, with deep brown branches, and scarcely any spines. The flowers, which appear in December, are yellow, and the fruit red. The leaves are sometimes wedge-shaped and 3-toothed, but more frequently are nearly round, with two or three spiny teeth on every side." (*Pen. Cyc.*, iv. p. 261.) A curious and beautiful species, well deserving of cultivation. There are plants in the Fulham Nursery 21s. each.



15. *B. ARISTATA* Dec. The bristled-tooth-leaved Berberry.

Identification. Hook. Exot. Flor., 2. t. 98; Dec. Prod., 1. p. 106; Don's Mill., 1. p. 115.
Synonymes. *B. Chitria* Buch; *B. angustifolia* Roab.; *B. sinensis* Desf.
Engravings. Hook. Exot. Flor., t. 98; Bot. Reg., t. 729; and our fig. 54.

Spec. Char., &c. Lower spines 3-parted, simple; leaves obovate-acute, tapering much to the base, ending in a mucro (prickly point) at the apex, membranous, smooth on both sides, serrated, with 4 or 5 bristly teeth. Racemes nodding, many-flowered, longer than the leaves. Berries oblong. (*Don's Mill.*, i. p. 115. adapted.) A robust shrub, very distinct from any of the preceding species or varieties, growing with extraordinary vigour, and capable of being formed into a very handsome small tree. It is a native of Nepal, and is found on mountains at from 5000 ft. to 8000 ft. of elevation, flowering there in May. The root and wood are of a dark yellow colour, and form the yellow wood of Persian authors; they are used as a dye, and, being bitter and a little astringent, they, as well as the bark, are employed in medicine. (*Royle's Illust.*, p. 63.) The plant was introduced into England in 1820, and is already in several gardens. In Nepal, the fruit of this species is dried, like grapes for forming raisins, in the



sun. After being once established, plants of this species grow with extraordinary rapidity till they attain the height of 8 ft. or 10 ft., after which they continue throwing up suckers, and some of these which we have measured in the Fulham Nursery, and in the Goldworth arboretum, were 9 ft. long, and, at the lower end, three fourths of an inch in diameter. It is a most desirable plant, and calculated to produce a splendid effect, both when in flower and when in fruit, upon an open lawn. As a rapid grower, it ought not to be planted near slow-growing shrubs or trees. Price of plants, in the London nurseries, 1s. each; at Bollwyller, ?; in New York, ?.

Varieties. Mr. Royle has the following remarks. "Under *B. aristata*, I conceive two species have been included, or at least two such very distinct varieties as to require particular notice. These are distinguished by the natives, apt to confound things together, by the names of *kushmul* and *chitra*. The former growing at as low elevations as 3000 feet. and therefore easily acclimated in the plains of India, has the leaves and branches pale-coloured, and more thorny; the flowers more numerous, racemes erect, appearing earlier in the season, and having less pleasant-tasted fruit: while *chitra*, which I conceive to be the true *B. aristata*, and have not found below 5000 feet of elevation, has brownish-coloured branches, smooth, shining, almost entire leaves, each flower much larger than those of *kushmul*, though less numerous than those on each of the drooping racemes." (*Illustr. &c.*, p. 64.)

App. i. *Additional Species of Bérberis.*

B. Wallichiana Dec., *synon. atro-viridis*. A native of the higher parts of Nepal, and, according to Dr. Lindley, "exceedingly well worth procuring, on account of its deep-green evergreen leaves," which will, in all probability, prove hardy. It is figured in Wallich's *Plant. Asiat. Rar.*, t. 243. — *B. kunawurënsis*, a native of the Himalaya, is also a very desirable species. — *B. actinacantha* is mentioned by Dr. Lindley as a very common plant, between Valparaiso and Santiago, which might be easily introduced: besides which, he says, there are other evergreen South American species of great beauty. "Some from the south of Chili particularly were found by Mr. Bridges near Valdivia, with shining holly-like leaves, long racemes of orange-coloured flowers, and young branches covered with rusty down." (*Penny Cyc.*, iv. p. 261.) — *B. buxifolia* Lam. Ill., t. 253. f. 3., and our fig. 55., a small twisted shrub, with bluish purple berries, a native of the Straits of Magellan, would be a desirable acquisition, as it is doubtless as hardy as *B. empetrifolia*, p. 306. In the garden of the Horticultural Society there is a berberry raised from seeds received from M. Ledebour, under the name of *B. filicifolia*, which has not yet flowered, but which is entirely different from the *B. heterophylla*, *synon. B. filicifolia* of the nurseries. Seeds of berberries from distant countries, the same authority observes, "would certainly reach England in safety, if mixed with tenacious earth and rammed into a box." The species both of *Bérberis* and *Mahonia* are so eminently beautiful, that too much can scarcely be said in their favour.



GENUS II.



MAHONIA Nutt. THE MAHONIA, or ASH BERRBERRY. *Lin. Syst.* Hexándria Monogýnia.

Derivation. Named by Nuttall in honour of Bernard M^{rs} Mahon a seedsman at Philadelphia, the author of the *American Gardener's Calendar*, and an ardent lover of botanical science.

Identification. Nutt. Gen. Amer., l. p. 307.; Dec. Prod., l. p. 108.; Don's Mill., p. 117.

Synonymes. *Bérberis* of authors; *Odostrëmon Raf.*; *Ash Berberry Pen. Cyc.*

Gen. Char. *Sepals* 6, guarded on the outside by three scales. *Petals* 6, without glands on the inside. *Stamens* furnished with a tooth on each side at top of the filament. *Berries* 3—9-seeded. (*Don's Mill.*, i. p. 118.) — The species are elegant evergreen shrubs with yellow flowers and pinnate leaves. The latter resemble pretty much those of the ash, and hence, doubtless, the name of ash berberry. Natives of the north-west coast of America, and also of Nepal, and perhaps Japan. Though some botanists think that the

characters ascribed to this genus, and those ascribed to *Berberis*, as exhibited in p. 229., are not sufficient to keep them separate as genera; yet the habits of the species of one, as to the mode of growth, foliage, and inflorescence, are so distinct from those of the other, as to induce us to adopt *Mahonia*. The species in British gardens are all of comparatively slow growth, and admit but of slow multiplication by layers, and scarcely at all by cuttings. Some of them, however, seed freely, and are readily propagated in that way. Four species have been introduced, and they are described by Dr. Lindley, in the *Penny Cyclopædia*, as being included in a section of the genus *Berberis*.

■ 1. *M. FASCICULARIS* Dec. The crowded-racemed Mahonia, or Ash Berberry.

Identification. Hook. Fl. Bor. Amer., 1. p. 28.; Dec. Prod., 1. p. 108., and Syst., 2. p. 19.; Don's Mill., 1. p. 118.

Synonymes. *Berberis pinnata* Lag., Bot. Reg., t. 702., and Bot. Mag., 2d edit. vol. 1. t. 88.; *E. fascicularis* Pen. Cyc. In the same work it is stated that *Mahonia diversifolia* is the same as this species; though it is figured and described by Sweet, as a species from Monte Video: see *Swt. Br. Fl.-Gar.*, 2d series, t. 56.

Engravings. Ker. Bot. Reg., t. 702.; Kth. Nov. Sp. Amer., 5. p. 71. t. 434.; Bot. Mag., t. 2396.; and our fig. 56.

Spec. Char., &c. Leaves

of 3—6 pairs with an odd one, the lowest pair near the base of the petiole.

Leaflets ovate-lanceolate, rather distant, one-nerved,

spiny-toothed, with 4 or 5 teeth on each side.

Racemes nearly erect, much crowded. Filaments

bidentate. (*Don's Mill.*, i. p. 118., adapted.)

A very handsome tall evergreen shrub, which

attains, in British gardens, especially if against a wall, the

height of 8 or 10 ft. and produces its

yellow flowers in abundance, from the middle of March to the middle of May.

“Perhaps the most showy of all the family.” (*Bot. Mag.*, 2d edit. vol. i. p. 48.) It is found in the mountainous parts of California and Mexico. It is readily distinguished at a distance from the other mahonias, by the glaucous green and subdued tone of colour of its leaves; those of all the others being of a darker green, and more or less shining.

The plant is rather too tender to be treated as a bush, unless some slight protection be given to it during very severe frosts; but it will grow freely against a wall with scarcely any protection.

There is a fine specimen of it in the London Horticultural Society's Garden, more than 8 ft. high. In the nurseries, plants are at present rather scarce, and cost from 5s. to 7s. each.

■ 2. *M. AQUIFOLIUM* Nutt. The Holly-leaved Mahonia, or Ash Berberry.

Identification. Nutt. Gen. Amer., 1. p. 212.; Dec. Prod., 1. p. 108.; Don's Mill., 1. p. 118.

Synonyme. *Berberis Aquifolium* Ph. and Pen. Cyc.

Engravings. Parsh. Fl. Amer. Sept., 1. t. 4.; Bot. Reg., t. 1425.; and our fig. 57.

Varieties. One variety, *M. A. nutkana* Dec., is mentioned by De Candolle (*Prod.*, i. p. 108.); and another, found at the junction of the Portage river with the Columbia, by G. Dou. (*Don's Mill.*, i. p. 118.)



Spec. Char., &c. Leaves in 4 pairs of leaflets with an odd one, the lower pair distant from the base of the petiole; leaflets ovate, approximate, cordate at the base, one-nerved, spiny-toothed, with 9 or 6 teeth on each side. Racemes erect, and much crowded. Filaments bidentate. (*Don's Mill.*, adapted.) One of the handsomest of hardy evergreen shrubs, attaining the height of 6 ft. in 6 years, quite hardy, producing a profusion of bunches of yellow flowers during April and May. It is a native of the north-west coast of America, from New Albion to Nootka Sound, growing in rich vegetable soil among rocks, or in woods, where it forms a thick and rich under-growth. It was introduced into England in 1823, and is to be found in all good collections. According to Dr. Lindley, it is "perhaps the handsomest hardy evergreen we yet possess. Its foliage is of a rich, deep, shining green, becoming purple in the winter; it bears fruit in some abundance, which consists of clusters of roundish black berries, having their surface covered with a rich violet bloom. It most resembles *M. fasciculàris*, from which its large shining leaves at once distinguish it." (*Penny Cyc.*, iv. p. 262.) This species is propagated very slowly by layers, and, for some years, plants were sold in the nurseries at ten guineas each. Lately, however, a number of ripe seeds have been produced in England, or imported from America through the Hudson's Bay Company; and from these, many young plants have been raised, in the Epsom and other nurseries; so that small plants may now be obtained for 5s. each, and in a few years they will, no doubt, not cost half that sum. In Prince's *Catalogue* for 1825, the price is stated as 25 dollars (3l. 5s.) each.



57

3. *M. NERVO'SA* Nutt. The nerved-leaved Mahonia, or Ash Berberry.

Identification. Nutt. Gen. Amer., 1. p. 212; Don's Mill., 1. p. 118.

Synonymes. *Berberis nervosa* Ph.; *Mahonia glumàcea* Dec.; *Berberis glumàcea* Pen. Cyc.

Engravings. Pursh. Fl. Amer., t. t. 5.; Bot. Reg., t. 1426.; and our fig. 58.

Spec. Char., &c. Leaves of 5—6 pairs, with an odd one, the lower pair distant from the petiole; leaflets ovate, acuminate, and remotely spiny-toothed, somewhat 3—5-nerved, with 12 or 14 teeth on each side; racemes elongated; filaments bidentate. (*Don's Mill.*, i. p. 118.) An evergreen undershrub, in its native habitats seldom exceeding the height of 3 ft., and producing its yellow flowers in October, succeeded by roundish fruit, of a glaucous-purple colour, and having an insipid taste. The plant is found in shady pine woods, on the



58

north-west coast of North America, along the river Columbia. According to Dr. Lindley, the stem of this species does not grow more than 6 in. or 8 in. high, and is, in fact, shorter than its leaves. The petioles of the leaves, he says, "are jointed at every pair of leaflets, in the manner of a bamboo stem." The plant is hardy, and will thrive in a shady border of peat soil. It was introduced into England in 1822, and may be seen in the London Horticultural Society's garden, but it is not yet extensively distributed. In London, plants cost 10s. 6d. each.

■ 4. *M. REPENS* G. Don. The creeping-rooted Mahonia, or Ash Berberry.

Identification G. Don. in Loud. Hort. Brit., No. 28182.; and in Don's Mill., i. p. 118.

Synonymes. *Berberis Aquifolium*, Lindl. Bot. Reg., t. 1176.; *Berberis repens* Pen. Cyc., iv. p. 262.

Engravings. Bot. Reg., t. 1176.; and our fig. 59.

Spec. Char., &c. Leaflets 2—3 pairs, with an odd one, roundish-ovate, opaque, spiny-toothed. Racemes diffuse. Root creeping. Filaments bidentate. (*Don's Mill.*, i. p. 118.)

A small branched evergreen shrub, seldom rising higher than 2 ft., with the leaves somewhat glaucous on both surfaces. The racemes of flowers are terminal, numerous, fasciated, diffuse, rising from scaly buds. The plant, in British gardens, produces a profusion of rich yellow flowers in April and May, but these have not yet been succeeded by fruit. Found wild on the east side of the Rocky Mountains of the west coast of North America, and perfectly hardy in British gardens. It is propagated by layers or suckers, but does not strike readily; and it has, in consequence, been but sparingly distributed. Price of plants, in London, 10s. 6d. each.



App. i. *Additional Species of Mahonia.*

Mahonia nepalensis Dec., *Berberis nepalensis* in the list in p. 173., is an evergreen Nepal shrub, from 4 ft. to 6 ft. high, of great beauty, which, if it could be introduced, would probably be found as hardy as the American species. — *M. acanthifolia*, if different from the foregoing, would also be very desirable. It is said to be a fine pinnated-leaved plant, with round black fruit, found on the Neelgherry Mountains of India, at the elevation of 8000 ft. *M. nepalensis* grows at the height of 5000 ft. and 6000 ft., and attains, in shady situations, an elevation of 12 ft. It is also found on the Neelgherries, in 11° of north latitude. (*Royle's Illust.*) In the *Penny Cyclopædia*, it is suggested that *M. nepalensis* "ought to be obtained from India at any cost, as it would in all probability succeed in this climate. — *M. tragacanthoides*, with not more than one or two pairs of leaflets, found along the banks of the river Kur, near Teflis; and *M. caraganæfolia*, a Chinese plant very like the last, having the points of the leaflets hardened into spines; well merit introduction." (iv. p. 262.) A plant is mentioned by Thunberg, under the name of *Ilex japonica*, which appears to be a *Mahonia*. It is found in the island of Nippon in Japan, and, as it would very likely prove hardy, ought by all means to be procured.

CHAP. IX.

OF THE HARDY LIGNEOUS SPECIES OF THE ORDER CRUCIACEÆ.

DISTINCTIVE Characteristics. Thalamiflorous. (*II. B.*) The order Cruciacæ is readily recognised by the cruciform arrangement of the petals, which are always four, in conjunction with tetradynamous stamens, and the fruit a silique or silicle. Though there are several species, which, technically considered, are ligneous plants, such as *Alýssum saxátile*, *Ibèris sempervirens*, *Cheiránthus Chèrri*, and some others; yet, in a popular point of view, the only shrub included in the order is the *Vélla Pseùdo-Cýtisis*.

GENUS I.



VÉLLA L. THE VELLA. *Lin. Syst.* Tetradynàmia Silicnòsa.

Derivation. The word *Vélla* is Latinised from the word *vetlar*, the Celtic name of the cress.

Gen. Char. *Stamens* the 4 longer in 2 pairs, the 2 of each pair grown together.

Style ovate, flat, tongue-shaped, at the tip of the silicle. *Silicle* ovate, compressed, its valves concave. *Partition* elliptic. *Cotyledons* folded, the embryo root disposed in the sinus of the fold. (*Dec. Syst.*)

1. *VÉLLA PSEU'DO-CÝTISIS L.* False Cytisus, or shrubby, Cress-Rocket.

Identification. *Lin. Sp.*, 895.; *Dec. Prod.*, l. p. 223.; *Don's Mill.*, l. p. 254

Synonyms. *Vélla integrifolia Sal.*; *Faux-cytise, Fr.*; *strauchartige (shrubby) Velle, Ger.*

Engravings *Cav. Ic.*, l. 42.; and our *fig.* 60.


Spec. Char., &c. Petals yellow, with long dark purple claws. Larger stamens perfectly connate by pairs. (*Don's Mill.*, l. p. 254.) A low evergreen shrub, seldom exceeding 4 ft. in height, with glaucous green leaves, and bright yellow flowers, which appear in the beginning of April, and continue till the middle of May. It is a native of Spain, on gypsaceous hills about Aranjuez, where it was first observed by Minuart, and, afterwards, by Cavanilles. It was cultivated by Miller in 1759, as a greenhouse plant; but is found sufficiently hardy to stand the open air with a slight protection. It has stood for several



years in the garden of the London Horticultural Society, planted on rockwork, where the dry soil renders protection unnecessary. It has also stood for a number of years in the open garden in the Hammersmith Nursery, and for five years in our garden at Bayswater, where it appears to be as hardy as the common azalea. It is a desirable shrub, on account of the early period at which it flowers; and also because it is a free flowerer. On a mound of rockwork it would form a most ornamental bush, and might be associated with the dwarf furze and *Nitrària Schóberi*. It is easily propagated by cuttings of the young wood, planted in sand under a hand-glass. Price, in London, 1s. 6d.

App. I. *Other ligneous or suffruticose Cruciferae.*

Those who wish to include in their collections all the hardy plants of Cruciferae, cultivated in the gardens, which are botanically considered as ligneous, will find them enumerated in our *Hortus Britannicus*. The principal are, *Cheiranthus Cheiri*, and several varieties, more especially *C. C. fruticulosus*, the wild wallflower; *Vesicaria utriculata*; *Alyssum argenteum*, *A. saxatile* (fig. 61.), and *A. gemonense*; *Iberis sempervirens* (fig. 62.), *I. saxatilis*, and *I. corifolia*; *Lepidium subulatum* and *L. suffruticosum*. *Sisymbrium Millefolium*, from the Canaries, might also, if it is possible, stand out; and it is very interesting, from its finely cut leaves, a character which is comparatively rare in cruciaceous plants. All the ligneous plants of this order are particularly adapted for rock-work; and, like all low-growing woody plants, even when grown in a common border, each ought to be elevated on a small mound or hillock of stones, of such a size as that, the plant after three or four years' growth, might hang down over it on every side, so as completely to conceal the stones.



CHAP. X.

OF THE HALF-HARDY LIGNEOUS PLANTS OF THE ORDER CAPPARIDA'CEÆ.

DISTINCTIVE Characteristics. Thalamiflorous. (*H. B.*) Sepals 4. Petals 4, cruciformly disposed. Stamens often numerous; if few never tetradynamous, or scarcely ever. Ovary stalked upon the receptacle. Fruit either pod-shaped or baccate, 1-celled, very rarely 1-seeded, most frequently with many seeds attached to two narrow simple parietal placentæ. Seeds kidney-shaped. Properties stimulant or tonic. (*Lindley's Introd. to N. S., and Key.*)

The only genus in this order, which contains any half-hardy ligneous plant, is *Capparis*.

GENUS I.



CA'PPARIS L. THE CAPER BUSH. *Lin. Syst.* Polyándria Monogýnia.

Derivation. From *kabir*, the Arabic name of the common caper.

Identification. *Lin. Gen.*, 643.; *Dec. Prod.*, 1. p. 245.; *Don's Mill.*, 1. p. 278.

Gen. Char. *Calyx* 4-parted. *Torus* small. *Fruit* a silique, somewhat baccate, upon a slender stalk.

✱ *I. C. SPINOSA* L. The spined, or common, Caper Bush.

Identification. Lin. Sp., 720; Dec. Prod., l. p. 245; Don's Mill., l. p. 278.

Synonymes. The caper having been in extensive cultivation, and used for so long a period, has names in most of the languages of the civilised world, and these, though too numerous to be given here, will be found in Newman's *Dictionary*, or in the index of synonymes to the *Encyclopædia of Plants*. In French, the name of the caper is Caprier or Tapenier; in Italian, Capriolo or Cappero; and in German, Kaperstrauch.


Engravings. Blackw. Herb., t. 417; Smith's Spic., 18. t. 12; Fl. Gr., 4^o6; Bot. Mag., t. 291; and our fig. 63.

Spec. Char., &c. Stipules spinose, hooked. Leaves ovate, roundish, deciduous.

Pediceles solitary, 1-flowered. (*Don's Mill.*, i. 278.)

Variety. There is said to be a variety without spines, and with ovate leaves, the latter more or less hoary.

Description, &c. The common caper plant is a well-known shrub, trailing and rambling like the bramble, found wild on the rocks in the south of Italy, in the Grecian Islands, and in various parts of Asia Minor. The stems are woody, and covered with a white bark, round, smooth, and branching. The leaves are oval or roundish, succulent, glaucous green, and deciduous. The stipules, which are two, at the base of the footstalks, are transformed into spines. The flowers are white, numerous, axillary, solitary, large, handsome, and without smell. The petals are much larger than the sepals; spreading, obovate, waved, tender and flaccid; white, faintly tinged with red or lilac. The stamens are about 60, of the same length as the petals, sometimes a little longer. The fruit is an oblong-oval coriaceous capsule. The root is long and woody, and covered with a thick bark.



Geography and History. Found wild in the southern countries of Europe, in the Levant, in Sicily, and in the south of Spain, on rocks, walls, and dry places. It is mentioned by Theophrastus as a wild plant, and incapable of being cultivated; but, in the south of France, it has been grown for the flower buds from time immemorial. It was cultivated by Gerard in 1596, who tells us that he put the seeds into the brick walls of his garden, "which did spring and grow green;" and Bradley states that, he sowed some seeds which he procured from Italy on the garden walls of Camden House, near Kensington, about the year 1716. Mr. Miller mentions an old plant growing there (probably the same that Bradley sowed), which resisted the cold many years, and annually produced many flowers, but the young shoots were frequently killed to the stump during the winter. This plant died about the year 1816. In general, the caper bush is grown in green-houses or stoves, but even in them it is not very frequent, and is seldom seen in flower. The largest plant that we know of in England is in the bed of a conservatory at Troughton Hall, near Manchester; its shoots cover a space between 20ft. and 30ft. long, and 4ft. or 5ft. wide, and it is profusely covered with blossoms every year.

Properties and Uses. The flower buds are well known throughout Europe and America as a pickle, and in the south of Italy the fruit is prepared in the same way as the buds. Their properties are, acridity, bitterness, and aroma. The buds are gathered daily, from the middle of May, when they begin to appear, till the end of autumn. They are taken when about half the size which they would attain when just about to expand. They are then thrown into a cask, among salt and vinegar, in which they remain till the end of the season, when they are taken out and sorted, and put into other casks with fresh vinegar, when they are fit for sale. Covered with vinegar, caper buds will keep many years. It is said that in order to increase the green colour, it is customary to put filings of copper in the first pickle. Bosc states that, in order

to effect the same object, they use sieves formed of copper wires, when separating the large buds from the small ones, previously to placing them in fresh vinegar; the consequence is, he says, that capers are always more or less poisonous. (*N. Cours d'Agr.*, tom. iii. 413.) The substitutes for capers are, the green fruits of the nasturtium (*Tropæolum majus*), and the unripe pods of the *Euphórbia Láthyris*.

Soil and Situation. A very dry soil, somewhat calcareous, and a situation fully exposed to the mid-day sun, are essential. It should either be planted against a wall, or on the south side of dry elevated rockwork; and, in either case, it will require some protection during winter. In the *Nouveau Du Hamel* it is stated that it will not grow at all if placed in the shade. In the neighbourhood of Paris, it is grown in light soil, on a stratum of broken limestone, and protected during winter with straw or leaves. There is a plant against the wall in the London Horticultural Society's garden, which in 1835 had stood there 8 years with very little protection. There is a large and vigorous plant of it in the botanic garden at Cambridge, planted in the open air, but in front of a stove, and near the furnace, which produces strong shoots, and flowers abundantly every year.

Propagation and Culture. In France, where ripe seeds can be procured, it is raised from them; but they require to be sown immediately after they are gathered. About Marseilles, where it is cultivated extensively in the fields, it is multiplied chiefly by cuttings; but partly also by division of the root. (See *Statistics*.)

Statistics. The caper is cultivated for its fruits and buds on both shores of the Mediterranean; and in Greece, and even in Egypt, the buds are gathered for sale from wild plants. In France, the only caper plantations are in the neighbourhood of Marseilles and Toulon, and these have existed from the time that Marseilles was founded by a colony from Greece. The plants are there grown in open fields, planted at 10 ft. apart in quincunx. They attain the height of 4 ft. or 5 ft., and the bush covers a space of about the same diameter. Every autumn all the shoots are cut off within 5 in. or 6 in. of the root; and, over the stools so formed, a little heap of earth is thrown up, of from 6 in. to 8 in. in thickness. In spring this earth is spread out, and the ground is hoed or ploughed; and this is the whole culture which the plant receives. As soon as the plants begin to flower, which, about Marseilles, is early in May, women and children are employed to gather the buds, and they continue doing so throughout the season, till the commencement of frost in November. Every day's gathering is thrown into a eask in the evening, and every addition of capers is followed by an addition of vinegar, with a little salt in it, so as to keep the buds always covered with liquor to the depth of 2 in. When a new plantation is to be made, the shoots cut off in the autumn are formed into cuttings of about a foot in length, which are immediately planted in a nursery, and covered with straw, to protect them from frost. They remain there two years, and afterwards are transplanted to their final situation, where two, and sometimes three, plants are always placed together to provide for deficiencies from deaths. Sometimes new plantations are formed by dividing the roots of old plants, and this operation is always performed in spring. The culture of the caper has been tried, with a view to commercial objects, in the neighbourhood of Paris, but without much success; not so much on account of the severity of the frosts there, as owing to the humidity both of the situation and of the climate. In Spain, on the shores of the Mediterranean, the caper is planted on the face of terrace walls on the sides of hills. Bose observes that the gathering of the caper buds by women and children is a "cruel torment" to them, on account of the numerous spines which cover the branches; and he adds that he has heard of a variety, which, however, he says, is not known in France, which is without spines, and which it would be very desirable to substitute for the other in general cultivation. In the south of France, every one who has a garden grows his own capers; and cottagers sometimes plant them in their garden walls, in order to sell the

produce. The caper is cultivated extensively in the neighbourhood of Tunis, and exported both to America and Europe. In commerce, the buds are of three different qualities, the nonpareil, the capucine, and the capotte. McCulloch says, the best capers imported into Britain are from Toulon; some small salt capers come from Majorca, and a few flat ones from about Lyons. In the year 1832, 6213 lbs. were entered for home consumption. (*Com. Dict.*)

The caper plant has, we believe, been introduced into Australia, and it is highly probable that it would thrive particularly well in that dry and warm climate; as it would, doubtless, in the Himalaya, and in other parts of India. For these reasons, we have departed from the rule we laid down, p. 230., which would have obliged us to print our account of this species, as being only half-hardy, in small type.

* 2. *C. FONTANÆ* Dec. Desfontaines's Caper Bush.

Identification. Dec. Prod., l. p. 245.; Don's Mill., l. p. 279.

Synonymes. *C. ovata* Desf. *Fl. Atl.*, l. p. 404.; Caprier oval, *Fr.*

Engraving. *Bocc. Sic.*, t. 42.

Spec. Char., &c. Stipules spinose, hooked. Leaves ovate, cordate at the base, acutish at the tip. (*Don's Mill.*, l. p. 279.) Flowers dull white. Fruit club-shaped. A deciduous bush, closely resembling *C. spinosa*, of which it is, in all probability, only a variety. It was found in Mauritania, near Oran, in fissures of rocks, by M. Desfontaines, and it is also to be met with in Sicily, Italy, Spain, and the states of Barbary. In the *Nouveau Du Hamel* it is stated that it differs from *C. spinosa* in nothing but the forms of the leaves, which are oval-acuminate, while those of the other are round. It appears to have been introduced into England in 1800, but we have not seen it. As it is, doubtless, equally hardy with the other, it well merits a place against a conservative wall.

From the habits common to the genus *Capparis*, and more especially from the principal part of the plant which contains the vital power being under ground, it is not improbable that all the greenhouse species might stand against a conservative wall with very little protection. One only is introduced, namely *C. ægyptia* Lam., from Egypt; but there are described by De Candolle, and by G. Don: *C. nepalensis* Dec., from Nepal; *C. nummularia* Dec., *C. quiniflora* Dec., and *C. umbellata* R. Br., from New Holland; *C. canescens* Banks, from New South Wales; *C. heteracantha* Dec., and *C. leucophylla* Dec., from between Bagdad and Aleppo; *C. volkamæriæ* Dec., *C. citrifolia* Lam., *C. cluytiefolia* Burch., *C. oleoides* Burch., *C. coriacea* Burch., *C. albitrunca* Burch., which is a tree 16 ft. high, *C. punctata* Burch., and *C. racemosa* Dec., all from the Cape of Good Hope; and *C. saligna* Vahl, from Santa Cruz.

CHAP. XI.

OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE ORDER CISTACEÆ.

DISTINCTIVE Characteristics. Thalamiflorous. Sepals 5, incompletely whorled, two of them being exterior. Petals 5, crumpled in æstivation, very fugitive. Stamens numerous. Fruit capsular, usually 3-valved or 5-valved, occasionally 10-valved; either 1-celled, with parietal placenta in the middle of the valves; or imperfectly 5-celled or 10-celled, with dissepiments proceeding from the middle of the valves, and touching each other in the centre. Embryo inverted. Properties balsamic. (*Lindl. Introd. to N. S.*, and *Key.*)

Description, History, &c. The species are all low ornamental shrubs, sub-evergreen or evergreen, most of them trailers, and only a few of them attaining the height of 5 ft. or 6 ft. They are natives of the south of Europe and north of Africa, but are scarcely known in America or Asia. One or more of the species of the Cistaceæ have been known from the days of Hippocrates. Linnæus included the whole of what were known in his time under two genera, *Cistus* and *Hudsonia*; but a new arrangement was published by Professor De Candolle (*Prod.* i.), in 1824, which he had adopted from Dunal, and this was followed by Sweet, in 1830, in his *Cistaceæ*; and by G. Don, in 1831, in his edition of Miller's *Dictionary*. This arrangement we shall adopt

in the present chapter, though we are convinced that most of the species described are mere varieties, some of them of the most fugitive kind. Our own opinion is, that all the different alleged species of the genera *Cistus*, *Heliánthemum*, and *Hudsonia* are, properly, only races or varieties of three or four original forms. The Cistaceæ have no medical properties; but the resinous balsamic substance called ladanum or labdanum is produced from *C. créticus*, *C. ladaniferus*, *C. laurifolius*, and one or two other species. (See *Mag. Nat. Hist.*, vol. ii. p. 408.) Some of the species which inhabit Turkey and Greece are liable, in those countries, to be injured by the growth of the hypocistis on their roots. The hypocistis is the *Cýtinus Hypocistis L.*, *Gynándria Octándria L.*, *Aristolochiæ Juss.*, and *Cytíneæ R. Br.* It is nearly allied to *Nepénthes* and *Aristolochia*; and is a succulent parasite of a rich red colour, bearing a distant resemblance in size and form to the *Orobánche*. It has been known from the days of Theophrastus, but, as far as we know, has never been seen in a living state in Britain. It is figured in *Du Ham.*, i. t. 68.; and in Gerard's *Herbal*, p. 1275. The use of the Cistaceæ in gardens is for ornamenting rockwork, or for keeping in pits during the winter, and planting out in flower-borders in spring; as, from the tenderness of the finer species, they are unfit for a permanent place in a shrubbery or arboretum. Most of the larger-growing kinds require some protection during winter; but they will all grow freely in any soil that is dry; and they are readily propagated by seeds, which, in fine seasons, they produce in abundance, or by cuttings; the plants, in both cases, flowering the second year. In the London nurseries the plants are generally kept in pots; and the price of the commoner sorts is from 1s. 6d. to 2s. 6d. a plant; at Bollwyller, where they are mostly green-house plants, it is 1 franc 50 cents; and in New York, ?.

The hardy ligneous species are included in three genera; which are thus contradistinguished by De Candolle and G. Don:—

CISTUS. Calyx of 5 sepals, 2 outer ones unequal or absent. Capsule covered by the calyx, 10—5-celled, from having a dissepiment in the middle of each valve.

HELIAÑTHEMUM. Calyx of 3 equal sepals, or of 5 unequal sepals. Capsule triquetrous, 1-celled, 3-valved, with a narrow dissepiment, or a placental nerve in the middle of each valve.

HUDSONIA. Calyx of 5 equal sepals. Capsule 1-celled, 3-valved, 1—3-seeded.

GENUS I.



CISTUS L. THE CISTUS, or ROCK ROSE. *Lin. Syst.* Polyándria Monogýnia.

Derivation. From the Greek word *kisté*, a box or capsule, or the Anglo-Saxon, *cist*, a hollow vessel; on account of the shape of its capsules. In Martyn's *Miller*, the name is said to be derived from that of the youth *Cistus*, whose story is to be found in *Cassianus Bassus*. Others derive it from *kis*, a worm or weevil.

Identification. Tourn., *Lin.*, Dec., G. Don.

Synonymes. Holly Rose *Gerard*; Gum Cistus; Ciste, *Fr.*; Cisten Rose, *Ger.*

Gen. Char. Calyx of 5 sepals; sepals disposed in a double series; 2 outer ones unequal, sometimes wanting. *Petals* 5, equal, somewhat cuneated, caducous. *Stamens* numerous, usually exerted from the glandular disk. *Style* filiform. *Stigma* capitate. *Capsule* covered by the calyx, 5- or 10-valved, with a seminiferous partition in the middle of each valve, therefore 5- or 10-celled. *Seeds* ovate, angular. *Embryo* filiform, spiral.—Elegant, erect shrubs or subshrubs, with opposite, exstipulate, entire or somewhat toothed leaves, and axillary, 1- or many-flowered peduncles. Flowers large, beautiful, resembling a single rose, red or white. (*Don's Mill.*, i. p. 298.)

§ i. *Erythrocistus*, Dec. i. p. 264.

Derivation. From *erythros*, red, and *cistus*; because the flowers of all the species in this section are red or purple.

Sect. Char. Outer sepals narrowest, and usually smallest; inner ones concave at the base, with scarious margins. Petals rose-coloured, red, or purple, with a yellow spot at the base of each. Capsule 5-celled, from having 5 seminiferous partitions, one in the middle of each valve. (*Dec. Prod.*, i. p. 264.; *Don's Mill.*, i. p. 298.) Low shrubs, evergreen, sub-evergreen, or deciduous, generally with large showy flowers.

A. *Peduncles 1-flowered, axillary or terminal, solitary or umbellate. Style cylindrical, generally longer than the Stamens. Stigma capitate, 5-furrowed.* (*Ibid.*)

α 1. *CISTUS PURPUREUS* Lam. The purple-flowered Cistus, or Rock Rose.

Identification. Lam. Diet., 2. p. 14.; Ker, in Bot. Reg., t. 408.; Dec. Prod., l. p. 264.; Swt. Cist., t. 17.; Don's Mill., l. p. 298.

Synonymes. *C. créticus Hort. Kew.*; the purple Gum Cistus, the purple Shrubby Cistus; *Ciste pourpre, Fr.*; *purpurrothe Cisten Rose, Ger.*

Engravings. Bot. Reg., t. 408.; Swt. Cist., t. 17.; and our fig. 64.

Spec. Char., &c. Leaves oblong-lanceolate, obtuse or acute, and more or less rugose; reticulately veined, with undulated margins. Petioles short, hairy, connected at the base, and sheathing the stem. Flowers terminal, from 1 to 6, on short peduncles. Bractees sessile, leaf-like, pubescent, broad and concave at the base, where they are connected, and terminating in acute points. Pedicels short, and with the calyx hairy; calyx of 5 sepals. Petals 5 or 6, obovate or wedge-shaped; very much imbricate, more or less crumpled. Stamens numerous, filaments smooth. Style very short; and stigma large, capitate, 5-lobed, papillose. (*Swt. Cist.*, 17.) A shrub about 3 ft. or 4 ft. high, and much branched; the branches are erect, and clothed with a brownish pubescence. The flowers are very large and handsome, of a bright reddish purple, with a yellow spot at the base, above which is a large dark velvet mark, surrounded with red, and slightly branched. The petals are imbricate, and much crumpled. It is a native of the Levant; but when it was introduced into England is uncertain: it seems to have been cultivated by Gerard under the name of *Cistus más angustifolius*, "with flowers of a purple colour, in shape like unto a single-flower briar rose, having leaves very like those of sage, wrinkled somewhat like unto a cloth new dried before it be smooth." It is rather tender; but, if planted near or against a wall, requires no other protection. It flowers abundantly in June and July, and is very ornamental. It grows very fast, and is easily propagated by cuttings.



64

α 2. *CISTUS HETEROPHYLLUS* Desf. The various-leaved Cistus, or Rock Rose.

Identification. Desf. Atl., l. p. 411. t. 104.; Dec. Prod., l. p. 264.;

Swt. Cist., t. 6.; Don's Mill., l. p. 298.

Synonymes. The Gum Cistus of Algiers; *Ciste hétérophylle, Fr.*

Engravings. Desf. Atl., l. t. 104.; Swt. Cist., t. 6.; and our fig. 65.

Spec. Char., &c. Leaves ovate-lanceolate; petioles very short, hairy, and sheathing at the base; margins of the leaves revolute, green on both sides. Peduncles hairy, one-flowered, with two leafy bractees about the middle of them. Flowers large, terminal. Calyx of 5 hairy sepals. Petals 5 or 6, imbricate, obovate, with roundish points. A stiff upright woody shrub, with short rigid branches, thickly clothed, as well as the other parts of the plant, with a hairy pubescence. The



65

flowers are of a bluish rose colour, with a bright yellow spot at the base; and the petals are imbricate, and much crumpled. The leaves are very small, and the whole plant has the appearance of a miniature tree. It is a native of uncultivated hills in Algiers; but by whom it was discovered, and when brought to the country, are unknown. It is rather tender, and requires protection during winter. It does best trained against a wall, where it has a very brilliant appearance in June and July, when it is covered with flowers. The seeds sometimes ripen in this country; and, when they do, they afford the best means of propagating the plant, as it does not strike freely from cuttings. It requires a light rich soil, and does best in a mixture of sandy loam and peat. (*G. Don. Sweet.*)

■ 3. *C. PARVIFLORUS Lam.* The small-flowered Cistus, or Rock Rose.

Identification. Lam. Dict., 2. p. 14; Dec. Prod., 1. p. 264; Swt. Cist., t. 14; and Don's Mill., 1. p. 298. *Engravings.* Swt. Cist., t. 14. Smith's Fl. Græc., t. 495.

Spec. Char., &c. Leaves ovate, acute, somewhat tomentose, drawn out into the footstalks at the base, and somewhat connate. Peduncles 1-flowered, 3 or 4 together, almost terminal. (*Don's Mill.*, i. p. 298.) A shrub about 3 ft. in height; a native of Crete. Mr. Sweet thinks it was probably introduced by Dr. Sibthorp, having been found in our collections ever since the doctor's return from that country. The petals are small, purplish, or pale rose-coloured, and distinct or separated from each other. It flowers in June and July, and sometimes ripens seeds, from which, or from cuttings, it is readily propagated. Plants were in the Chelsea Botanic Garden, and in the Fulham Nursery, in 1826. (*Sweet.*)

■ 4. *C. COMPLICATUS Lam.* The complicated Cistus, or Rock Rose.

Identification. Lam. Dict., 2. p. 14; Don's Mill., 1. p. 298.

Spec. Char., &c. Leaves roundish-ovate, bluntish, approximate, clothed with white tomentum; under surface reticulated; footstalks dilated at the base, with pilose margins, channelled above, and sheathing at the base. Peduncles short, 1-flowered, three or four together, somewhat terminal. (*Don's Mill.*, i. p. 298.) A shrub, from the Levant, and from the mountains of Valencia, in Spain, growing to the height of 3 ft., and producing small rose-coloured or purplish flowers in June and July. Introduced into England in 1818.

■ 5. *C. VILLOSUS Lam.* The villous Cistus, or hairy Rock Rose.

Identification. Lam. Dict., 2. p. 12; Lin. Sp., 736; Willd. Sp., p. 1181; Hort. Kew., 2d. edit., 5. p. 303; Dec. Prod., 1. p. 264; Don's Mill., 1. p. 298.

Synonymes. *C. salviifolius Hort.*; *C. undulatus Macrch.*; *Cistus más mājor folio rotundiore Duh.*; *C. creticus Hort. Lam.*; the shrubby *Cistus Mart. Mill.*; *Ciste velu, Fr.*; Raube Cisten Rose, *Ger. Engravings.* Duh. Ar., 1. t. 64; Swt., t. 55; Willd., p. 2. 1181.

Spec. Char., &c. Leaves roundish-ovate, wrinkled, tomentose, and hairy, stalked; footstalks furrowed, connate at the base. Peduncles 1-flowered, 1 or 3 together. Sepals villous. (*Don's Mill.*, i. p. 298.) A shrub 3 ft. high. A native of the south of Europe and the north of Africa, which has been in the English and French gardens for the last two hundred years. It is, as Mr. Sweet observes, one of the commonest species in all the nurseries about London, where it is sold under several names, and generally for *C. salviæfolius*; which, however, is a white-flowered species, though it resembles the present plant in habit. This shrub forms a "snug compact bush," and continues in flower for a long time. The flowers vary in colour from a pale lilac to a dark purple, and even very much on the same plant at different times. In severe winters it requires a little protection; and it will generally be found safe to keep a reserve of young plants in pots, in a pit or cold-frame.

Variety.

■ *C. v.* 2 *rotundifolius.* The round-leaved villous Cistus, or Rock Rose. *C. rotundifolius Sweet*; *C. villosus β virescens Dec.* (*Swt. Cist.* t. 75.)—Leaves more obtuse than in the species.

■ 6. *C. CRÆTICUS L.* The Cretan Cistus, or Rock Rose.

Identification. Lin. Sp., 1. p. 738; Jacq. Icon. rar., 1. t. 95; Smith's Fl. Græc., 495; Buxb. Cent. 3. p. 34. t. 64. f. 1.; Dec. Prod., 1. p. 264; Swt. Cist., t. 112.

Synonymes. *Lédon Diosc.*; *Ciste de Crète, Fr.*; *Cretische Cisten Rose, Ger.*

Engravings. Jacq. Icon. rar., 1. t. 95.; Fl. Gr., t. 495.; Buxb. Cent., 3. p. 34. t. 64. f. 1.; Swt. Cist., t. 112.; and our fig. 66.

Varieties. *C. c. 2 crispatus* Dec. has the leaves waved or curled; and *C. c. 3 touricus* Dec. has the leaves flat, and very villous, on the under surface.

Spec. Char., &c. Leaves spatulate-ovate, tomentosely hairy, wrinkled, tapered into the short footstalk, waved on the margin. Peduncles 1-flowered. Sepals villous. (*Don's Mill.*, i. p. 298.) This species, Sweet observes, resembles *C. villösus* and *C. undulatus* in appearance, and is often confused with those species in collections. In the nurseries, *C. purpureus* is very often sold for it; but the fine yellow spots at the base of its petals readily distinguish it from that species. It is a shrub, a native of Crete, Syria, and Greece, growing to the height of 2 ft., and generally requiring protection in the gardens about London; which as it does not often receive, it is, in consequence, scarce. The gum ladanum is the produce of this species. Dioscorides tells us that in his time



66

the gum that exuded from the glands of the leaves was obtained by driving goats in among the shrubs, or by these animals naturally browsing upon them, when the substance adhered to their hair and beards, whence it was afterwards combed. This resin being at present collected to supply an extended commerce, a peculiar instrument is employed for the purpose, which is figured and described by Tournefort, and which is a kind of rake with a double row of long leathern straps. (See *Mag. Nat. Hist.*, vol. iii. fig. 21.) The following is the description of the mode of gathering the gum given by Sieber in his *Voyage to Crete*: — “It was in the heat of the day, and not a breath of wind stirring; circumstances necessary to the gathering of ladanum. Seven or eight country fellows, in their shirts and drawers, were brushing the plants with their whips; the straps whereof, by rubbing against the leaves of the shrub, licked up a sort of odoriferous glue, sticking on the leaves; this is a part of the nutritious juice of the plant, which sweats through the texture of the leaves like a fatty dew, in shining drops, as clear as turpentine. When the whips are sufficiently laden with this grease, they take a knife and scrape it clean off the straps, and make it up into a mass or cakes of different sizes: this is what comes to us under the name of *ladanum*, or *labdanum*. A man who is diligent will gather three pounds in a day, or more, which they sell for a crown on the spot. This sort of work is rather unpleasant than laborious, because it must be done in the sultry time of the day, and in the deadest calm; and yet the purest *ladanum* cannot be obtained free from filth, because the winds of the preceding day have blown dust upon the shrubs.” (*Sieber's Crete*, as quoted in Murray's *Encyc. of Geog.*, p. 835.) Formerly ladanum was a good deal used in pharmacy, but at present it is comparatively neglected. In the west of Europe, a considerable quantity of it, however, is annually collected in Crete, and sent to Constantinople, where it is chewed by the Turks, and used in various preparations of laudanum, and for fumigating churches and mosques.

67

n. 7. *C. INCA'NUS* L. The hoary Cistus, or Rock Rose.

Identification. *Län. Sp.*, 737; *Smith's Fl. Græc.*, 494; *Don's Mill.*, I. p. 298.; *Cist.*, t. 44.

Synonymes. *C. albidus Hort.*; *C. cymösus Dec.*; *Ciste cotonneux, Fr.*; *be-staubte Cisten Rose, Ger.*

Engravings. *Bot. Mag.*, t. 13; *Swt. Cist.*, t. 44; and our *fig.* 67.

Spec. Char., &c. Leaves spatulate, tomentose, wrinkled, somewhat 3-nerved, sessile, somewhat connate at the base, upper ones narrower. Peduncles 1—3-flowered. (*Don's Mill.*, i. p. 298.) A shrub, a native of Spain and France, about Narbonne, and which has been in our gardens since the time of Gerard. It grows to the height of 3 ft., forming a hoary bush, with reddish purple flowers, having the petals emarginate,



and flowering in July and August. It will endure our mildest winters in the open air; but in severe frosty weather it will require to be protected by glass, or by some slight covering. Plants of this species were in the Hammersmith Nursery in 1826.

Variety.

- *C. i. 2 canescens.* *The canescent-leaved Cistus, or hoary Rock Rose.* *C. canescens* Swt. (*Don's Mill.*, i. p. 298.); *Cistus mās* Clus.; *C. incānus* var. β Dec. (*Swt. Cist.* t. 45.)—Leaves oblong-linear, bluntish, tomentose, hoary, waved, rather 3-nerved, sessile, somewhat connate at the base. Peduncles terminal, 1-flowered, or somewhat cymose. Sepals ovate, acute, nerved, clothed with starry pubescence. Petals obovate, distinct. (*Don's Mill.*, i. p. 298.) Native of the south of Europe. A shrub growing to the height of 2 ft. in British gardens, and greatly resembling the preceding species; the general colour and surface of the plant being the same, and also the colour of its flowers. Mr. Sweet says that he has no doubt of its being perfectly distinct; which it may be, and yet be only a variety. It is tender, and requires protection like the species.

- 8. *C. UNDULATUS* Dec. *The waved-leaved Cistus, or Rock Rose.*

Identification. Dec. Prod., 1. p. 264.; *Don's Mill.*, 1. p. 299.

Synonymes. *C. crēticus* Swt., t. 63., afterwards corrected to *C. undulatus*; perhaps *C. crispus* var. *Don.*

Engraving. Swt. Cist., t. 63., under the name of *C. crēticus*.

Spec. Char., &c. Leaves sessile, linear-oblong, acute, with waved margins, 3-nerved at the base. Peduncles solitary, each furnished with a bractea. Sepals taper-pointed, villous. (*Don's Mill.*, i. p. 299.) A shrub 2 ft. high, cultivated in collections, but of which the native country is unknown. It has purple flowers, which appear in June and July; and is probably a hybrid. It is rather tender, and not very frequently to be met with. It was in the Hammersmith Nursery, when Mr. Sweet's drawing was made, in 1827.

- 9. *C. CRISPUS* L. *The curled-leaved Cistus, or Rock Rose.*

Identification. Lin. Sp., 738.; Swt. Cist., 22.; *Don's Mill.*, 1. p. 299.

Synonymes. *Ciste crépu*, Fr.; krause Cisten Rose, Ger.

Engravings. Cav. Icon., 2. t. 174.; Swt. Cist., 22.

Spec. Char., &c. Leaves sessile, linear-lanceolate, undulately curled, 3-nerved, wrinkled, pubescent. Flowers almost sessile, 3 or 4 together, somewhat umbellate. (*Don's Mill.*, i. p. 299.) Native of the south of France, Spain, and Portugal, and introduced into England in 1656. It is a shrub, growing to the height of 2 ft., and producing showy purple, or reddish purple, flowers in July and August. The leaves are ribbed, or nerved, and covered with hairs, much undulated at the edges, and of a whitish green. They vary considerably in size, as well as in form. It is a very distinct sort, and forms a very pretty bush, which will stand the severity of our winters without protection. Cuttings of the young wood, Mr. Sweet observes, planted under hand-glasses in autumn will strike root readily; but they will not strike so freely in summer. Plants of this kind were in the Fulham Nursery in 1826.

- 10. *C. ALBIDUS* L. *The white-leaved Cistus, or Rock Rose.*

Identification. Lin. Sp., 737. *Don's Mill.*, 1. p. 299.

Synonymes. *Ciste blanche*, Fr.; weissliche Cisten Rose, Ger.

Engraving. Swt. Cist., t. 31.

Spec. Char., &c. Leaves sessile, oblong-elliptical, hoary-tomentose, somewhat 3-nerved. Flowers 3 or 8, terminal, somewhat umbellate. Outer sepals largest. (*Don's Mill.*, i. p. 299.) Native of France, about Narbonne, Spain, and Portugal. In British gardens, a shrub growing to the height of 2 ft., erect, much branched, and thickly crowded with white hoary leaves. The flowers, which are of a pale purple, a bright lilac, or a pale rose colour, terminate the branch in a sort of umbellate corymb, and appear in

July and August. It is one of the most desirable species of the genus, being quite hardy, having flowers of the largest size, and thriving in almost any soil or situation not too moist. Cuttings put in in autumn are soon rooted, and the plant ripens seeds plentifully in ordinary seasons.

α. 11. *C. CANDIDISSIMUS* *Dun.* The whitest-leaved *Cistus*, or *Rock Rose*.

Identification. *Dun. ined. Dec. Prod.*, 1. p. 264; *Swt. Cist.*, 3.; *Don's Mill.*, 1. p. 299.

Engravings. *Swt. Cist.*, t. 3.

Spec. Char., &c. Leaves ovate-elliptical, acute, densely clothed with hoary tomentum, 3-nerved; footstalks short and sheathing at the base, with pilose margins. Peduncles solitary, 1-flowered, shorter than the leaves. Outer sepals one half shorter than the rest. (*Don's Mill.*, 1. p. 299.) Native of the Grand Canary Island, in elevated pine forests. This is a noble species, growing to the height of 4 ft. and upwards, with fine poplar-like leaves, and large pale rose-coloured flowers, with distinct petals, which appear in July and August. It was introduced into the Botanic Garden at Chelsea in 1815, where it flowered soon afterwards, and plants existed there in 1825. It forms a shrub, not sufficiently hardy to stand through the winter, about London, in the open air, as a bush; but, with dry litter laid about its roots, and a slight covering of mats in the most severe weather, it may be preserved.

α. 12. *C. VAGINATUS* *Ait.* The sheathed-petioled *Cistus*, or *Rock Rose*.

Identification. *Hort. Kew.*, 3. p. 304; *Jacq. Hort. Sch.*, 3. p. 17.; *Swt. Cist.*, t. 9.; *Don's Mill.*, 1. p. 299.

Synonymes. *Cistus symphytifolius Lam.*; *Ciste à Feuilles de Consoude, Fr.*; *schidenartige Cisten Rose, Ger.*

Engravings. *Jacq. Hort. Sch.*, 3. p. 17. t. 282.; *Bot. Reg.*, t. 225.; *Swt. Cist.*, 1. 9.

Spec. Char., &c. Leaves lanceolate, acute, 3-nerved, hairy, under surface reticulated; footstalks furrowed, dilated, and sheathing at the base, with pilose margins. Peduncles 3-flowered, axillary or terminal, long, bracteate at the base. (*Don's Mill.*, i. p. 299.) Native of the Island of Teneriffe. Introduced in 1779. A splendid-flowered species, easily distinguished by its panicle flowers, and large, imbricate, obovate, crumpled petals. It grows to the height of 4 ft.; and, being rather tender, it is generally kept in green-houses or pits. Its flowers are light rose colour, darker without and pale within: their general appearance is that of an apple blossom on a large scale; and they continue appearing from April to June. Plants were in the Kensington Nursery in 1826.

α. 13. *C. SERICEUS* *Vahl.* The silky-leaved *Cistus*, or *Rock Rose*.

Identification. *Vahl. Symb.*, 1. p. 37.; *Don's Mill.*, 1. p. 299.

Engravings. *Barrel. Icon.*, 1315.

Spec. Char., &c. Leaves ovate, tomentose, 3-nerved; lower ones on footstalks, upper ones sessile. Peduncles hairy. (*Don's Mill.*, i. p. 299.) Native of Spain, producing its purple flowers in June and July. It was in cultivation in 1826; and is said to grow to the height of 3 ft.

α. 14. *C. HYBRIDUS* *Vahl.* The hybrid *Cistus*, or *Rock Rose*.

Identification. *Vahl. Symb.*, 1. p. 37.; *Don's Mill.*, 1. p. 299.

Spec. Char., &c. Leaves ovate, hoary, on footstalks. Branches beset with yellow scales. Peduncles elongated, subracemose, hairy. Outer sepals caducous. (*Don's Mill.*, i. p. 299.) Native of Spain, where it produces its purple flowers in June and July. It is said to grow to the height of 3 ft., but has not yet been brought to Britain.

B. *Peduncles cymose. Style almost wanting. Stigma capitate, shorter than the Stamens.*

α. 15. *C. CYMOSUS* *Dun.* The cymose-flowered *Cistus*, or *Rock Rose*.

Identification. *Dun. ined. Dec. Prod.*, i. p. 265.; *Don's Mill.*, 1. p. 299.

Synonymes. *Cistus incanus Sib. Fl. Gr.*, and at one time in the garden of Cels.

Engravings. *Fl. Gr.*, t. 494., as *C. incanus*; *Swt. Cist.*, t. 90.

Spec. Char., &c. Leaves broad-ovate, twisted at the top, acutish; under surface wrinkled and hoary. Footstalks dilated at the base, and somewhat sheathing, furrowed above. Peduncles cymose, 5- or 10-flowered, hoary, axillary, or terminal. (*Don's Mill.*, i. p. 299.) A very handsome, small, bushy shrub, growing in British gardens to the height of 3 ft. It is a native of the Levant, and has been some years in cultivation in British gardens; but how long is uncertain. Sweet says it is often mistaken for *C. incanus*,

but that it is nearer related to *C. villòsus*. It is somewhat tender, like all the other species from the Levant; and, in the climate of London, it requires a green-house, a cold-frame, or other protection, during winter.

§ ii. *Ledònia* Dec.

Derivation. From *lèdon*, a name given by Dioscorides to the plant that produces the ladanum; but which is supposed by some to be *Cistus Lèdon* (*Dec. Prod.*, 1. p. 265.), and by others to be *C. créticus*. (See p. 320.)

Sect. Char. Sepals 5, 2 outer ones largest, and very much pointed, or wanting. Petals white or whitish, with a yellow or purple mark at the base of each. Stamens numerous, longer than the pistil. Stigma almost sessile, large, capitate. Capsules 5- or 10-celled, from being furnished with 5 or 10 seminiferous partitions, one in the middle of each valve. Sub-evergreen shrubs or subshrubs. Leaves usually covered with clammy gluten. (*Don's Mill.*, i. p. 299.) This section includes some of the finest species of the genus; such as *C. cýprius*, *C. ladaníferus*, *C. laurifolius*, &c.: almost all of them are evergreen, and many of them form bushes from 4 ft. to 6 ft. in height, or more, which, when covered with flowers, are among the most ornamental objects that can be introduced into a shrubbery or flower-garden.

a. *Peduncles 1-flowered or many-flowered, cymose. Sepals 5, outer ones usually cordate at the Base, and pointed at the Apex. Capsules 3-celled.*

a. *Peduncles naked at the Base, usually bearing beneath their Middle two opposite small Leaves.*

ii. 16. *C. SALVIEFO`LIUS* L. The Sage-leaved *Cistus*, or *Rock Rose*.

Identification. *Lin. Spec.*, 38.; *Cav. Icon.*, 2. p. 31.; *Jacq. Coll.*, 2. 120.; *Swt. Cist.*, t. 54.; *Smith's Fl. Græca*, t. 497.; *Don's Mill.*, 1. p. 299.

Synonymes. *Cistus foemina Clus. Hist.*, 1. p. 70.; *Ciste à Feuilles de Sauge, Fr.*; *Salbey-blättrige Cisten Rose, Ger.*

Engravings. *Cav. Icon.*, 2. t. 137.; *Jacq. Coll.*, 2. t. 8.; *Swt. Cist.*, t. 54.; *Smith's Fl. Græc.*, t. 497.

Spec. Char., &c. Leaves stalked, ovate, obtuse, wrinkled; under surface tomentose. Peduncles long, white from tomentum, 1-flowered, articulated above, solitary or ternary. (*Don's Mill.*, i. p. 299.) Native of the south of France, Italy, Greece, Spain, and Portugal. Flowers white. A branchy shrub in British gardens, producing white flowers in July and August, and cultivated since the year 1548. The leaves are of a whitish or pale green colour; and, like every other part of the plant, are covered with numerous short hairs. It is readily distinguished from other species, by its solitary, 1-flowered, jointed peduncles, and its obtuse leaves. It is a very hardy species; and, in sheltered situations, it will endure the winter without any protection. The sage-like leaves and neat flowers of this plant give it a very pleasing appearance.

Varieties.

ii. *C. s. 2 creticiusculus* Dec. has the stem rather more erect than the species; and *C. s. 3 ochroleucus* Dec. has the flowers cream-coloured.

ii. 17. *C. OBTUSIFO`LIUS* *Sict.* The obtuse-leaved *Cistus*, or *Rock Rose*.

Identification. *Swt. Cist.*, 1. 42.; *Don's Mill.*, 1. p. 299.

Engraving. *Swt. Cist.*, t. 42.

Spec. Char., &c. Leaves almost sessile, tapering to the base, ovate-oblong, obtuse, wrinkled, clothed with starry pubescence; margins somewhat denticulated. Peduncles terminal, cymose, many-flowered. Outer sepals broadly cordate, acute. Petals obcordate, imbricated. (*Don's Mill.*, i. p. 299.) Native of Crete. Petals white, with a yellow spot at the base of each. This is a dwarf shrub, seldom growing higher than a foot and a half. Being a native of Crete, it requires shelter in severe frost. Mr. Sweet thinks that it has been in our gardens since the time of Dr. Sibthorp, without being noticed as distinct from *C. salviæfolius*, of which, we have no doubt in our own minds, it is only a variety. Plants were in the Fulham Nursery in 1826.

■ 18. *C. CUPANI'ANUS* Presl. Cupani's Cistus, or Rock Rose.

Identification. Presl ex Spreng. Syst. Append., p. 206; Don's Mill., i. p. 299.
Engraving. Swt. Cist., t. 70.

Spec. Char., &c. Stem erect. Leaves stalked, cordate-ovate, 3-nerved, reticulately veined; upper surface scabrous, under surface covered with fasciated hairs; margin fringed. Peduncles pilose, 2- or 3-flowered. Sepals villous, acuminate. Petals imbricated. (*Don's Mill.*, i. p. 299.) Native of Sicily, but when introduced into England is uncertain. It grows to the height of 2 ft., and produces white flowers, with a spot of yellow at the base of each petal, in June and July. Sweet says that it is very nearly hardy, requiring protection only during the severest frosts; and, in sheltered situations, requiring no protection at all. It is nearly related, he says, to *C. salviaefolius*, and to *C. corbariensis*; but is of stronger growth than either of these. Plants were in the Fulham Nursery in 1827. The heart-shaped leaves of this species render it easily distinguishable from the others that have white flowers.

Variety.

- *C. C. 2 acutifolius.* The acute-leaved Cupani's Cistus, or Rock Rose. (*Swt. Cist.*, t. 78.; *Don's Mill.*, i. p. 299.) *C. acutifolius* Swt.; *C. salviaefolius* β humifusus Dec. Prod., i. p. 265., *Swt. Cist.*, t. 78.—Leaves cordate-ovate, 3-nerved, reticulately veined, pubescent on both surfaces. Branches twiggy, diffuse, rather prostrate. Peduncles tomentose, generally 3-flowered. Sepals cordate, acute, shining, rather pilose, ciliated. Petals obovate, imbricated at the base. (*Don's Mill.*, i. p. 299.) Cultivated in Colvill's Nursery, Chelsea, in 1827; and, according to Sweet's figure, so closely resembling the preceding sort, that we have no doubt of their belonging to one species or race.

■ 19. *C. CORBARIENSIS* Pourr. The Corbières Cistus, or Rock Rose.

Identification. Dec. Prod., i. p. 265.; Swt. Cist., t. 8.; Don's Mill., i. p. 299.
Synonymes. *C. salviaefolius* β Dec. Prod. Fl. Fr., 4. p. 813.; *C. populifolius* minor, in some nurseries; *C. hybridus* Pourr., not of Vahl.
Engraving. Swt. Cist., t. 8.

Spec. Char., &c. Leaves stalked, somewhat cordate, ovate, acuminate with fringed margins, wrinkled on both surfaces, and very glutinous. Peduncles long, 1—3-flowered. (*Don's Mill.*, i. p. 299.) Found in the south of France, on the mountains of Corbières, and also in Spain; and cultivated in the English gardens in the year 1656. It is a shrub, 2½ ft. high, handsome, and producing its flowers in May and June. According to Sweet, it is one of the hardiest species of the genus, thriving well in the common garden soil, and in any situation where it is not too moist. It continues in bloom for about two months; and every day during that period the plant is covered with a profusion of handsome white flowers, the margins of which are tinged with rose colour. The rose-coloured buds are also very pretty before the flowers expand. Plants were in the Hammersmith Nursery in 1825.

■ 20. *C. FLORENTINUS* Lam. The Florentine Cistus, or Rock Rose.

Identification. Lam. Diet., 2. p. 17.; Swt. Cist., t. 59.; Don's Mill., i. p. 300.
Engraving. Swt. Cist., t. 59.

Spec. Char. Leaves narrow-lanceolate, wrinkled, reticulated on the under surface, almost sessile. Peduncles villous, generally 3-flowered. (*Don's Mill.*, i. p. 300.) A native of Italy, whence it was brought to England in 1825. It is a shrub 3 ft. high, producing its white flowers in June and July. It is a rare and very distinct species, Mr. Sweet observes; and forms a very pretty upright bush, requiring some protection during winter. Plants of it were in the Fulham Nursery in 1826.

■ 21. *C. MONSPELIE'NSIS* L. The Montpellier Cistus, or Rock Rose.

Identification. Lin. Sp., 737.; Lam. Ill., t. 477.; Cav. Icon., 2. t. 137.; Jacq. Coll., 2. t. 8.; Swt. Cist., t. 27.; Smith's Fl. Græc., t. 493.
Synonymes. Ciste de Montpellier, Fr.; Französische Cisten Rose, Ger.
Engravings. Lam. Ill., t. 477. fig. 4.; Cav. Icon., 2. t. 137.; Jacq. Coll., 2. t. 8.; Swt. Cist., t. 27.; Smith's Fl. Græc., t. 493.

Spec. Char., &c. Leaves linear-lanceolate, sessile, 3-nerved, clammy, villous on both surfaces. Peduncles pilose, cymose, somewhat secund. (*Don's Mill.*, i. p. 300.) A native of the south of France, Spain, and Portugal; and, at one time, abundant at Montpellier, as the name implies. A shrub which grows to the height of 4 ft., and has been in cultivation since the time of Gerard, in 1656. The leaves are lanceolate, wrinkled, and reflexed at the points; and the flowers are middle-sized, white, with the petals broadly cuneate. This sort is rather scarce in British collections, though it is of the easiest culture, and is a handsome-growing plant, and an abundant flowerer. It requires a warm border, and some protection in severe weather.

■ 22. *C. PLATYSE'PALUS* Swt. The broad-sepaled Cistus, or *Rock Rose*.

Identification. Swt. Cist., t. 47; Don's Mill., i. p. 300.

Engraving. Swt. Cist., t. 47.

Spec. Char., &c. Leaves oblong-lanceolate, sessile, wrinkled, 3-nerved, villous on both surfaces. Peduncles cymose, and are, as well as the calyxes, villous. Sepals acuminate, outer ones broadly cordate. Petals obcordate, distinct. (*Don's Mill.*, i. p. 300.) A shrub growing to the height of 3 ft. or 4 ft.; a native of Crete; and producing its white flowers in June and July. It is uncertain when it was introduced; but plants of it were in the Fulham Nursery in 1826. Sweet says that it is frequently confused with *C. monspeliensis*, though no two plants need be more distinct, and it is much nearer related to *C. hirsutus*.

■ 23. *C. LE'DON* Lam. The Ledon Cistus, or *Rock Rose*.

Identification. Lam. Dict., 2. p. 17; Duh. Arb., 1. p. 168, t. 66. Don's Mill., 1. p. 300.

Synonymes. *C. undulatus* Link? *C. ladaniferus monspeliensium* Bauh.; Ciste de Montpellier qui donne du ladanum, Ciste de Ledon, Fr.

Engraving. Duh. Arb., 1. p. 168, t. 66.

Spec. Char., &c. Leaves connate, oblong-lanceolate, nerved; upper surface smooth, shining, under surface silky, villous. Flowers corymbosely cymose. Peduncles and calyx clothed with silky hairs. (*Don's Mill.*, i. p. 300.) Found wild in the south of France, and figured and described by Du Hamel in 1755. He says ladanum is obtained from this species, and from others that are easily known by their having the smell of that gum. In catalogues, this species is marked as having been introduced in 1730; but Mr. Sweet remarks, in 1829, that he had not seen it about London for the last 10 years.

■ 24. *C. HIRSU'TUS* Lam. The hairy Cistus, or *Rock Rose*.

Identification. Lam. Dict., 2. p. 17; Clus. Hist., 1. p. 78; Don's Mill., 1. p. 300.

Engraving. Swt. Cist., t. 19.

Spec. Char., &c. Leaves sessile, oblong, blunt, hairy. Peduncles short, 1-flowered, or cymosely many-flowered. Capsules small, covered by the large, hairy, pyramidal calyx. (*Don's Mill.*, i. p. 300.) A shrub growing to the height of 4 ft.; a native of France and the mountains of Spain; producing its white flowers in June and July; and cultivated in England in the year 1656. It forms a pretty little branching bush, readily distinguished from all other species of this section, by its very large pyramidal calyx, and small capsules. It bears our winters, in the open borders, without protection, except when the frost is very severe. It produces a great profusion of flowers, which continue to expand in succession for a considerable time. Plants were in the Fulham Nursery in 1826.

■ 25. *C. SIDERITIS* Presl. The Ironwort-like Cistus, or *Rock Rose*.

Identification. Spreng. Syst. Append., p. 204; Don's Mill., 1. p. 300.

Spec. Char., &c. Decumbent. Leaves stalked, obovate, a little wrinkled, covered with hoary tomentum on the under surface. Peduncles elongated, 1—2-flowered, articulated above; and are, as well as the calyx, tomentose. Flowers nodding before expansion. (*Don's Mill.*, i. p. 300.) A decumbent shrub, with white flowers from June to August, found on rocks in Sicily, and not yet introduced into British gardens.

■ 26. *C. LA'XUS* Ait. The loose-flowering Cistus, or *Rock Rose*.

Identification. Ait. Hort. Kew., 3. p. 305; Don's Mill., 1. p. 300.

Synonymes. ? *C. capensis* Lin. Sp. 736; schlaffe Cisten Rose, Ger.

Engraving. Swt. Cist., t. 12.

Spec. Char., &c. Leaves on short footstalks, ovate-lanceolate, acuminate, with wavy somewhat toothed margins, smoothish, upper ones hairy. Flowers cymose. Peduncles and calyx hairy. (*Don's Mill.*, i. p. 300.) A shrub 3 ft. high, with white flowers and imbricate petals. Found wild

in Spain and Portugal, and introduced into England in 1656. It is erect, and not much branched; quite hardy, or requiring only very slight protection in very severe frost. Dumont thinks it probable that it is only a variety of *C. lædon*.

☞ 27. *C. OBLONGIFOLIUS* Swt. The oblong-leaved Cistus, or Rock Rose.

Identification. Swt. Cist., t. 87.; Don's Mill., 1. p. 300.

Engraving. Swt. Cist., t. 67.

Spec. Char., &c. Erect. Branches hispid, villous. Leaves on short footstalks, oblong-lanceolate, obtuse, pubescent, and waved at the margins; under surface veiny. Peduncles cymose. Petals concave, imbricated. (*Don's Mill.*, i. p. 300.) A native of Spain, attaining the height of 4 ft., and producing its white flowers in June and July. When it was introduced into England is uncertain; but Sweet found plants which appeared to him to be of this species in Colvill's Nursery in 1826. It forms a strong handsome evergreen shrub, which is quite hardy, and of the easiest culture.

☞ 28. *C. ASPERIFOLIUS* Swt. The rough-leaved Cistus, or Rock Rose.

Identification. Swt. Cist., t. 87. Don's Mill., 1. p. 300.

Engraving. Swt. Cist., t. 67.

Spec. Char., &c. Leaves almost sessile, ovate-lanceolate, acute, 3-nerved, wrinkled, smoothish, with wavy margins, somewhat denticulated, ciliated, netted with veins beneath, with the nerves and veins rough. Flowers cymose. Peduncles and calyxes hairy. Petals imbricate. (*Don's Mill.*, i. p. 300.) A shrub growing to the height of 2 ft., and producing its large white flowers from May to August. Its native country is uncertain; and Sweet says it may probably be a garden production, and, if so, intermediate between *C. læxus* and *C. oblongifolius*. It forms a handsome, strong, upright, evergreen shrub, and is quite hardy, producing its flowers all the summer, and till late in autumn, in the open border. There were plants of it in Colvill's Nursery in 1826.

☞ 29. *C. PSILOSEPALUS* Swt. The glabrous-sepaed Cistus, or Rock Rose.

Identification. Swt. Cist., t. 33.; Don's Mill., 1. p. 300.

Engraving. Swt. Cist., t. 33.

Spec. Char., &c. Leaves on short footstalks, oblong-lanceolate, 3-nerved, acute, with undulated margins, which are somewhat denticulated and ciliated, rather hairy. Flowers somewhat cymose. Peduncles hairy, tomentose. Sepals with long points, glabrous, shining, and with ciliated edges. Petals broad, cuneated, imbricated. (*Don's Mill.*, i. p. 300.) A shrub growing from 2 ft. to 3 ft. high, with white flowers in July and August. Apparently a garden production. Plants of it were in the Hammersmith Nursery in 1826. It approaches near to *C. longifolius*, but is very different, according to Sweet, from that species. The plants are bushy, and the shoots are terminated by large cymes of white flowers, which open in succession, and produce a contrast with the dark green leaves with which the plants are clothed.

b. *Peduncles with small, concave, coriaceous, yellowish, decussate, caducous Bracteoles at the Base; and with two larger opposite ones beneath the Middle.*

☞ 30. *C. LONGIFOLIUS* Lam. The long-leaved Cistus, or Rock Rose

Identification. Lam. Diet., 2. p. 16.; Don's Mill., 1. p. 300.

Synonyms. *Cistus nigricans* Pourr. *Act. Toul.*, 3. p. 311.; *C. populifolius* var. *longifolius* Dumont.

Spec. Char., &c. Leaves on short peduncles, oblong-lanceolate, with waved and pubescent margins; under surface veiny. Peduncles cymose. (*Don's Mill.*, i. p. 300.) A shrub growing to the height of 4 ft., and producing its white flowers in July and August. It does not appear to have been introduced into England. Dumont de Courset, in his *Botaniste Cultivateur*, says that this species is only a variety of *C. populifolius*; seeds of the latter having, with him, produced the former.

☞ 31. *C. POPULIFOLIUS* L. The Poplar-leaved Cistus, or Rock Rose.

Identification. Lin. Sp., 736.; Don's Mill., 1. p. 300.

Synonyms. Ciste à Feuilles de Peuplier, Fr.; Pappel-blättrige Cisten Rose, Ger.

Variety.

☞ *C. p. 2 minor* Dec. Prod., 1. p. 266.—Peduncles and calyx smoothish, shining, clammy. *Lædon latifolium*, n., *Cus. Hist.*, i. p. 78.; *Cistus populifolius* Cav. *Icon.*, 3. 215., *Swt. Cist.*, 23.

Spec. Char., &c. Leaves stalked, cordate, acuminate, wrinkled, smooth. Flowers cymose. Peduncles bracteate. Bracteas oblong. Sepals acuminate, clammy. (*Don's Mill.*, i. p. 300.) A shrub of vigorous growth, attaining the height of from 3 ft. to 5 ft. in British gardens, into which it was introduced in 1656. It has large dark green cordate leaves, with undulate margins, and white flowers, with distinct petals, which it produces from May to July. It is one of the most robust of the species, and is found wild both in the south of France and in Spain and Portugal. It is of the easiest culture, but requires a little protection in very severe winters. Plants were in the Hammersmith Nursery in 1826.

■ 32. *C. LATIFO'LIIUS* Swt. The broad-leaved Cistus, or Rock Rose.

Identification. Swt. Cist., 15; *Don's Mill.*, i. p. 300.

Synonymes. Cistus populifolius var. α major *Dec. Prod.*, 1. p. 266.

Engraving. Swt. Cist., t. 16.

Spec. Char., &c. Leaves stalked, broadly cordate, acute; with curled, waved, denticulated, ciliated margins. Peduncles bracteate, somewhat cymose, pilose. Sepals broad, cordate, villous. Petals imbricated. (*Don's Mill.*, i. p. 300.) A shrub from Barbary, with white flowers from May to July, growing to the height of 3 ft. or 4 ft., cultivated in British gardens since 1656. It has broadly cordate leaves, with reflexed points, and imbricate, obcordate petals. It is a most ornamental plant, robust in its growth, but rather tender. Plants of it were in the Hammersmith Nursery in 1826. We have no doubt that this and the two preceding sorts are nothing more than varieties of the same race.

B. *Peduncles bracteate with caducous decussate Bracteas, lower ones smaller; 1-flowered, axillary and solitary, or terminal and umbellate. Calyx of 3 Sepals. Capsules 5—10-celled*

a. *Stigma large, sessile.*

■ 33. *C. LAURIFO'LIIUS* Lin. The Laurel-leaved Cistus, or Rock Rose.

Identification. Lin. Sp., 736; Swt. Cist., t. 52; *Clus. Hist.*, 1. p. 78. f. 1.; *Don's Mill.*, 1. p. 300.

Synonymes. Ciste à Feuilles de Laurier, *Fr.*; Lorbeer-blättrige Cisten Rose, *Ger.*

Engravings. Swt. Cist., t. 52; *Clus. Hist.*, 1. p. 78. f. 1.

Spec. Char., &c. Leaves stalked, ovate-lanceolate, 3-nerved, upper surface glabrous, under surface tomentose. Footstalks dilated, and connate at the base. Capsules 5-celled. (*Don's Mill.*, i. p. 300.) A shrub growing in British gardens to the height of 4 ft. or 5 ft., and producing large white flowers in July or August. It is a native of the south of France and Spain, and was introduced into England in 1771. It is a robust bush, with large green laurel-like leaves: it produces an abundance of flowers, which, with their light red bracteas, are very ornamental before they expand, resembling, at a distance, the bursting buds of roses. It requires no protection, and may be raised from seeds, which it ripens in abundance, and also by cuttings, which, however, do not strike so freely as in some of the other species.

■ 34. *C. CY'PRIUS* Lam. The Gum Cistus, or Cyprus Rock Rose.

Identification. Lam. Diet., 2. p. 16; *Don's Mill.*, 1. p. 300.

Synonymes. C. ladaniferus *Bot. Mag.*, t. 112; Cistus stenophyllus *Link. Enam.*, 2. p. 74. ?; C. salicifolius of some.

Engraving. Swt. Cist., t. 39.

Spec. Char., &c. Leaves stalked, oblong-lanceolate, upper surface glabrous, under surface clothed with hoary tomentum. Peduncles generally many-flowered. Petals spotted. Capsules 5-celled. (*Don's Mill.*, i. p. 300.) A splendid shrub, growing in British gardens to the height of 5 ft. or 6 ft.; introduced from the Island of Cyprus in 1800. Its flowers are large, $2\frac{1}{2}$ in. or 3 in. across; white, imbricated, each petal having a dark, rich, brownish crimson spot at the base. It is one of the handsomest species of the genus, and one more generally in demand than any other. In the nurseries, it is generally found under the name of *C. ladaniferus*, or that of *C. salicifolius*. Young plants require protection; but when they get to the height of 3 ft. or 4 ft. they are tolerably hardy. Young cuttings, Sweet observes, planted under hand-glasses in autumn, will strike root; but the best way is to raise them from layers or from seed. There is a plant of this species at Minard, in Argyllshire, 7 ft. 9 in. high, with a head 12 ft. in diameter, which is clothed

with flowers every year. In the London nurseries, where this species is generally called *C. ladaniferus*, or by its English name of gum cistus, plants are from 1s. to 1s. 6d. each.

♣ 35. *C. LADANIFERUS* L. The Ladanum-bearing Gum Cistus, or Rock Rose.

Identification. Lin. Sp., 757; Don's Mill., 1. p. 300.

Synonymes. Ciste ladanifère, Fr.; Ladanum Cisten Rose, Ger.

Varieties. C. l. 1. *albiflorus* Dec. Prod., i. p. 266.; Swt. Cist., t. 94.; Lèdon, i., *Clus. Hist.*, i. p. 78. ic.; and C. l. 2. *maculatus* Dec. Prod., l. c., Swt. Cist., C. l. 3. *plenifolius* *Ait. Hort. Kew.*, iii. p. 305., are varieties of this species.

Spec. Char., &c. Leaves almost sessile, connate at the base, linear-lanceolate, 3-nerved, upper surface glabrous, under surface tomentose. Capsule 10-celled. Petals imbricate. (*Don's Mill.*, i. p. 301.) A shrub 4 ft. high, a native of Spain and Portugal, upon hills; introduced into England in 1629, and producing large white flowers in June and July. The leaves are lanceolate, and nearly sessile, of a deep green; the flowers terminating the branches, solitary, white, and large; each flower being from 1½ in. to 2 in. broad. The plant requires a little protection during winter, and was to be had in Colvill's Nursery in 1826. This species, as well as *C. Lèdon* and *C. créticus*, and doubtless various others, produces the resinous exudation known as gum ladanum, the mode of gathering which is described in p. 320.

b. *Stigma capitate, small.* Style cylindrical, equalling the Stamens in Length.

♣ 36 *C. CLUSII* Dunal. Clusius's Cistus, or Rock Rose.

Identification. Dunal. *inced.*, Dec. Prod., l. p. 266.; Swt. Cist., 32.; Don's Mill., 1. p. 301.

Synonymes. Cistus Libanotis *β* *Lam. Dict.*, 2. p. 18.; *Desf. Atl.*, 1. p. 412.; *C. undulatus* Link; Lèdon, vii., *Clus. Hist.*, 1. p. 80. ic.

Engraving. Swt. Cist., t. 32.

Spec. Char., &c. Erect. Leaves somewhat 3-nerved, linear, with revolute margins, under surface canescent. Flowers somewhat capitate. Calyx 3—5-sepaled, pilose. Sepals ovate, acute. Capsules 5-celled. (*Don's Mill.*, i. p. 301.) A shrub 2 ft. high, from Spain and Barbary, in 1810. The leaves and flowers are smaller than those of any of the other sorts here described. The plant forms a handsome and compact bush, and stands the winter well in a dry situation. It approaches the nearest to *C. monspeliensis*. Plants were in the Fulham Nursery in 1826.

GENUS II.



HELIANTHEMUM. THE HELIANTHEMUM, or SUN ROSE. Lin. Syst. Polyándria Monogýnia.

Identification. Tourn. Inst., 248. t. 128.; Gært. Fr., 1. p. 371. t. 76.; Dec. Fl. Fr., 4. p. 815.; Prod., 1. p. 265.; Don's Mill., 1. p. 301.

Synonymes. Cisti species of Lin.; Helianthe Sonnen Gurtel, Ger.

Derivation. From *helios*, the sun, and *anthe*, a flower; because the flowers open with the rising of the sun in the morning, and the petals fall off with the setting of the sun in the evening. The flowers of Helianthemum, as well as of Cistus, only last for a few hours when the sun shines; and if the weather is dull, and the sun does not make its appearance, the flowers do not open, but remain unexpanded. Should this continue for several days together, they will decay in the bud.

Gen. Char. Calyx of 3—5 sepals; when 3, these are equal, and disposed in a single series; but, when 5, they are unequal, and disposed in a double one; the two outer sepals are usually smaller than the inner ones, very rarely larger. Petals 5, usually regularly denticulated at the top. Stigma capitate. Style sometimes almost wanting, sometimes straight, sometimes oblique, and sometimes bent at the base. Ovary triquetrous. Capsule 3-valved; valves with a narrow dissepiment, or a semiferous nerve in the middle of each. Seeds angular, smooth. Albumen mealy. Embryo uncinately inflexed—Erect or trailing herbs, subshrubs, or shrubs. Leaves opposite and alternate, with or without stipules, 3-nerved or feather-nerved. Pedicels usually furnished with bractees at the base

sometimes opposite the bractæas, or opposite the leaves, sometimes solitary, sometimes umbellate, and sometimes racemose; and, before the flowers expand, the racemes at the top are bent or twisted backwards, and become gradually erect as the flowers expand. (*Dec. Prod.* and *G. Don.*) Flowers yellow, red, or white. All the species are ornamental; those which form erect bushy undershrubs are suitable for warm dry borders; and those which are trailers of the lowest size, for growing on rockwork, for ornamenting old walls or ruins, or for growing in pots. This genus has been arranged by De Candolle in three divisions, which have been subdivided into sections. The divisions are as follows:—

- I. Style straight, erect, almost wanting, or shorter than the stamens. Stigma capitate. § i. ii. and iii., *Halimium*, *Lecheoides*, and *Tuberaria*, belong to this division.
- II. Style straight, erect, equal with or longer than the stamens. § iv. and v., *Maculãria* and *Brachypetalum*, belong to this section; but, as they are entirely herbaceous, we omit them.
- III. Style bent at the base; to which belong § vi. vii. viii. and ix., *Eriocarpum*, *Fumãna*, *Pseïdo-Cistus*, and *Euheliãthemum*.

§ i. *Halimium* Dec. Prod., 1. p. 267.

Derivation. From *halimos*, marine; alluding to the habitation of the plants by the sea-side. (*Don's Mill.*, i. p. 301.)

Sect. Char. Calyx usually of 3 equal sepals; rarely of 5 unequal sepals, but when this is the case the two outer ones are the smallest. Petals rarely white, usually yellow, wedged-shaped, truncate, and marked at the base with a dark bloody or intense yellow spot. Style straight, short or almost wanting. Stigma capitate, somewhat 3-lobed. Seeds few, blackish, minutely mucicated, and somewhat angular. Erect shrubs. Leaves opposite, 3-nerved, without stipules, pilose, or tomentose. Peduncles 1—3-flowered, axillary, solitary, or umbellate, or rarely panicled. (*Dec.* and *G. Don.*)

A. *Style short, straight.*

1. H. LIBANO'TIS Willd. The Rosemary-leaved Helianthemum, or *Sun Rose*.

Identification. Willd. Enum., 570; Don's Mill., i. p. 301.

Synonyms. *Cistus Libanôtis* Lin. Sp., 739; *Brot. Fl. Lus.*, 2. p. 261., but not of Lam. or Desf.; *H. rosmarinifolium* Lag. in *Litt.*, but not of Pursh; *Barrct. Icon.*, 294; *Lêdon*, viii., *Ctus. Hist.*, i. p. 80. ic; *Ciste à Feuilles de Rosmarin*, Fr.

Spec. Char., &c. Erect, smoothish, branched. Leaves sessile, linear, with revolute margins; upper surface brownish green, under surface somewhat canescent. Bractæas oblong-linear, shorter than the peduncles. Peduncles solitary, 1-flowered. Calyx of 3 sepals, smooth, shining, ovate-acuminate. (*Don's Mill.*, i. p. 301.) A shrub growing to the height of 1 ft., a native of Portugal and Mauritania, and introduced into England in 1752. It produces yellowish-white flowers in June and July, and is of the easiest culture.

2. H. UMBELLA'TUM Mill. The umbellate-flowered Helianthemum, or *Sun Rose*.

Identification. Mill. Dict., No. 5; Don's Mill., i. p. 302.

Synonyms. *Cistus umbellatus* Lin. Sp., 739; *Cistus Libanôtis* of some.

Spec. Char., &c. Suffruticose, branched, younger branches tomentosely pilose, clammy. Leaves sessile, linear-oblong, with revolute margins, clammy; under surface tomentose. Peduncles 1-flowered, disposed in whorled racemes, terminal, umbellate. Calyx 3-sepaled, villous. (*Don's Mill.*, i. p. 302.) An undershrub 1 ft. high, with white flowers from June to August. Introduced into England in 1731. This is a very handsome and very distinct sort, well adapted for ornamenting rockwork, or for growing in pots. Sweet says it is generally sold in the nurseries under the name of *Cistus Libanôtis*. Plants were in the Hammersmith Nursery in 1825.

Varieties.

- *H. u. 2 erectum* Dec. (Swt. Cist., 5.), *Cistus umbellatus* Lam., *Cistus verticillatus* Brot., has the stem erect, and the leaves on the under surface clothed with greenish tomentum.
- *H. u. 3 subdecumbens* Dec., *Cistus umbellatus* var. *a* Lam., has the stem somewhat decumbent. Leaves somewhat ciliated, and clothed with white tomentum on the under surface.
- *3. H. OCYMOÏDES Pers.* The Basil-like Helianthemum, or *Sun Rose*.

Identification. Pers. Syn., 2. p. 76.; Don's Mill., 1. p. 302.

Synonymes. *Cistus ocyroides* Lam. Dict., 2. p. 18.; *Cistus sampsucifolius* Cav. Icon., 1. p. 65. t. 96.

Engraving. Clus. Hist., 1. p. 72. ic.

Varieties. Two are mentioned by De Candolle; one of them with the peduncles glabrous, and the other with the branches and peduncles rather hairy.

Spec. Char., &c. Erect, branched. Branches hoary. Cauline leaves obovate, or ovate-oblong, 3-nerved, almost sessile, green; those of the branches are stalked, and keeled on the back, reflexed at the top, hoary on both surfaces. Peduncles long, branched, paniculate. Pedicels opposite, somewhat umbellate. Calyx of 3 much-pointed sepals. (*Don's Mill.*, i. p. 302.) A shrub growing to the height of 3 ft., a native of Spain and Portugal; but when introduced into England is uncertain. According to Sweet, this species is very often confused in the collections with *H. algarvense*, though it is very different. It is a very handsome undershrub, and, being rather tender, requires the protection of a wall, or a covering of mats during severe frosty weather.

B. *Style almost none. Stigma large.*

- *4. H. ALYSSOÏDES Vent.* The Alyssum-like Helianthemum, or *Sun Rose*.

Identification. Vent. Choix., t. 20.; Lam. Dict., 2. p. 20.; Dec. Fl. Fr., 4. p. 818.; Don's Mill., i. p. 302.

Synonymes. *Cistus alyssoides* α Lam. Dict., 2. p. 20., Dec. Fl. Fr., 4. p. 818., Don's Mill., 1. p. 302.

Engraving. Vent. Choix., t. 20.

Spec. Char., &c. Erect, much branched, diffuse, spreading. Branches hoary, tomentose hairy at the top. Leaves sessile, tapering towards the base, oblong-ovate, bluntish, covered with short hairs; younger ones rather hoary, adult ones green. Peduncles terminal, solitary, or umbellate, 1—2-flowered, longer than the leaves. Calyx 3-sepaled, acuminate, hairy. (*Don's Mill.*, i. p. 302.) A shrub, native of Spain and the west of France, growing to the height of 3 ft., and producing its yellow flowers from June to August. It is remarked of the flower buds, that they are of an intense purple colour at the apex. It does not appear to be in cultivation in British gardens.

- *5. H. RUGOSUM Dum.* The wrinkled-leaved Helianthemum, or *Sun Rose*.

Identification. Dunal. ined. Dec. Prod., 1. p. 268.; Swt. Cist., t. 65.; Don's Mill., 1. p. 302.

Engraving. Swt. Cist., t. 65.

Spec. Char., &c. Erect. Branches rather hairy, clothed with leprous tomentum, scabrous, of a brownish grey colour. Leaves sessile, tapering into the footstalks at the base, obovate-oblong, bluntish, rather oblique, with the margins somewhat denticulated, and a little curled, tomentose on both surfaces, wrinkled, under surface hoary. Peduncles terminal, axillary, or umbellate, 1—3-flowered, hairy, shorter than the leaves. (*Don's Mill.*, i. p. 302.) A shrub 3 ft. high, a native of Spain, introduced in 1800, and producing its fine yellow flowers from June to August. Its stem is shrubby, erect, or a little flexuose; the branches spreading, more or less hairy; the young shoots reddish; and the flowers terminal, with broadly ovate petals of a golden yellow, each with a large dark spot near the base. It is distinguished from all other sorts by the stiff bristle-like purple hairs of its calyx. It is rather tender, but it will amply repay protection, either against a wall or in a pit. Plants were in the Bristol Nursery in 1827.

- *6. H. MICROPHYLLUM Sut.* The small-leaved Helianthemum, or *Sun Rose*.

Identification. Swt. Cist., t. 96.; Don's Mill., 1. p. 302.

Synonymes. *H. rugosum* β *microphyllum* Dec. Prod., 1. p. 268.; *H. alyssoides* β *microphyllum* Dec. Fl. Fr. Suppl., p. 622.

Engraving. Swt. Cist., t. 96.

Spec. Char., &c. Much branched, erect. Branches blackish grey, hairy, tomentose at the apex. Leaves almost sessile, obtuse, keeled, tapering to the base, obscurely greyish, tomentose. Flowers terminal, panicle. Pedicels 1—3-flowered, very short. Calyx of 3 sepals, very hairy. Petals cuneated, distinct. (*Don's Mill.*, i. p. 302.) A suffruticose bush, seldom growing higher than 2 ft.; found in the west of France, near Bourdeaux, and introduced into England in 1800. The leaves are small, nearly sessile, very glaucous. The petals are of a bright yellow; and the plant flowers from June to August, or later. It is commonly kept in a green-house or pit, in pots; but it will stand in the open air, if planted near a wall or fence, protected during severe weather by mats. Plants of it were in the Chelsea Botanic Garden in 1828.

■ 7. *H. SCABRO'SUM Pers.* The rough Helianthemum, or *Sun Rose*.

Identification. Pers. Ench., 2. p. 76.; Brot. Fl. Lus., 2. p. 265.; Don's Mill., i. p. 302.

Synonymæ. *Cistus scabrosus Art. Hort. Kew.*, 2. p. 236.

Engravings. Swt. Cist., t. 81.; and our fig. 68.

68

Spec. Char., &c. Erectish. Branches pilosely tomentose, scabrous, canescent. Leaves sessile, tapering to the base, oblong-ovate, acutish, roughish, 3-nerved, with waved revolute margins; upper surface green; under surface clothed with grey tomentum. Peduncles terminal, 1—2-flowered, shorter than the leaves. Calyx 3-sepaled, hairy. Petals distinct. (*Don's Mill.*, i. p. 302.) A shrub, from the north of Portugal, growing to the height of 3 ft.; cultivated in British gardens since 1775; and producing its fine yellow flowers from June to August. It forms a handsome little bush, and requires a sheltered situation. Plants of it were in the Bristol Nursery in 1827.



■ 8. *H. ALGARVE'NSE Dum.* The Algarve Helianthemum, or *Sun Rose*.

Identification. Dum. ined.; Dec. Prod., 1. p. 268.; Swt. Cist., t. 40.; Don's Mill., 1. p. 302.

Synonymæ. *Cistus algarvensis Bot. Mag.*, t. 627.; *H. algarvense Tourn. Inst.*, 250.?

Engraving. Swt. Cist., t. 40.

Spec. Char., &c. Stem branched. Leaves sessile, ovate-lanceolate, obtuse; hoary on the under surface; upper surface green, pilose. Peduncles somewhat panicle, pilose. Calyx 3-sepaled, acute, hairy. (*Don's Mill.*, i. p. 302.) A shrub 3 ft. high, found in Portugal, in the Algarves; introduced into England in 1800; and producing its fine deep-yellow flowers from June to August. The petals are crenated, with a dark base, similar in colour to those of *Calliopsis bicolor*. According to Sweet, this species is nearly allied to *H. ocymoides*. It is rather tender, and requires protection during winter. Plants of it, at 1s. 6d. each, may be obtained in all the London nurseries.

■ 9. *H. FORMO'SUM Dum.* The beautiful Helianthemum, or *Sun Rose*.

Identification. Dunal. ined. Dec. Prod., 1. p. 268.; Don's Mill., 1. p. 302.

Synonymæ. *Cistus formosus Bot. Mag.*, t. 264.

Spec. Char., &c. Erect. Branches tomentously villous, canescent. Leaves on short footstalks, obovate-lanceolate, tomentously villous, younger ones hoary. Peduncles and calyxes villous. Calyx 3-sepaled. (*Don's Mill.*, i. p. 302.) A shrub growing to the height of 4 ft., introduced in 1780, and producing its fine large flowers, with yellow dark-spotted petals, from May to July. The stem is erect and much branched, the leaves lanceolate and 3-nerved, and the flowers the largest of the genus. It is a most desirable plant to keep in pots, and turn out into borders in the spring; or, by keeping it against a wall and protecting it with mats, it will produce a fine show

of flowers from the beginning of May to the end of July. It ripens seeds in abundance; but, as Sweet remarks, "the colour of the flowers varies considerably on different plants," and therefore the seeds should always be saved from those of the brightest colours. Plants are to be procured in all the London nurseries.

■ 10. *H. ATRIPLICIFOLIUM* Willd. The Orache-leaved Helianthemum, or Sun Rose.

Identification. Willd. Enum., 569; Don's Mill., 1. p. 302.

Synonymy. *Cistus atriplicifolius* Lam. Dict., 2. p. 19.

Engraving. Barrel. Icon., t. 292.

Spec. Char., &c. Erect. Branches white from leprous tomentum. Leaves stalked, broad-ovate, bluntnish, waved at the base, covered with leprous tomentum on both surfaces. Peduncles racemose, hairy. Calyx hairy, 3-sepaled, rarely 5-sepaled, with the two outer ones very minute. (Don's Mill., i. p. 302.) A shrub growing to the height of 3 ft., with leaves like those of *Atriplex Halimus*; introduced from Spain in 1826, and producing its yellow flowers in June and July. Mr. Sweet observes, in 1829, that this species, he believes, has "quite disappeared from our collections."

■ 11. *H. LASIANTHUM* Pers. The hairy-flowered Helianthemum, or Sun Rose.

Identification. Pers. Ench., 2. p. 76; Don's Mill., i. p. 302.

Synonymy. *Cistus lasianthus* Lam. Dict., 2. p. 19.

Engraving. ? Barrel. Icon., t. 289.

Spec. Char., &c. Stem suffruticose, and much branched. Branches dark-cinereous, at top tomentose hairy. Leaves almost sessile, ovate-oblong, often blunt, keeled, of an obscure greyish colour, tomentose. Peduncles 1—2-flowered, hairy, very short. Calyx usually 3-sepaled, very hairy. (Don's Mill., i. p. 302.) A shrub 3 ft. high, from Spain, introduced in 1826, and producing its yellow flowers in June and July.

■ 12. *H. INVOLUCRATUM* Pers. The involucreted-flowered Helianthemum, or Sun Rose.

Identification. Pers. Ench., 2. p. 76; Don's Mill., 1. p. 302.

Synonymy. *Cistus involucratus* Lam. Dict., 2. p. 20.

Spec. Char., &c. Branched, erect. Branches rather greyish, tomentose. Lower leaves stalked, somewhat ovate, small, hoary-tomentose; upper ones oblong-lanceolate, sessile, greenish, and roughish. Peduncles very short, surrounded by the leaves. Calyx 5-sepaled, inner ones hoary-tomentose, outer ones linear, smoothish, and greenish. (Don's Mill., i. p. 302.) An erect shrub, 2 ft. high, from Spain and Portugal, in 1826, which produces its yellow flowers in June and July.

■ 13. *H. CHEIRANTHOIDES* Pers. The Wallflower-like Helianthemum, or Sun Rose.

Identification. Pers. Ench., 2. p. 76; Don's Mill., 1. p. 303.

Synonymy. *Cistus cheiranthoides* Lam. Dict., 2. p. 19; ? *Cistus elongatus* Fahl. Symb., 1. p. 38.

Cistus halimifolius, ii., *Ctus. Hist.*, 1. p. 71.

Engraving. Swt. Cist., t. 107.

Spec. Char., &c. Erect, branched; younger branches villously tomentose, hoary. Leaves tomentose, hoary, oblong-lanceolate, tapering into the foot-stalks. Peduncles very short, 2-flowered. Calyx somewhat villous, 5-sepaled, outer sepals very minute. (Don's Mill., i. p. 303.) An erect handsome bushy shrub, growing to the height of 3 ft., and producing yellow flowers, without dark spots on the petals, in July and August. This very handsome species, Sweet observes, was in the Bristol Nursery in 1828.

■ 14. *H. CANDIDUM* Swt. The white-leaved Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 25; Don's Mill., 1. p. 303.

Engraving. Swt. Cist., t. 25.

Spec. Char., &c. Erect. Branches leprously white. Leaves, which are obovate-lanceolate, equally white on both surfaces, and tapering to the base; somewhat stalked, upper surface pilose, under surface scabrous from papillæ, rather 3-nerved; floral ones opposite, sessile, and green on both surfaces. Peduncles long and rather panicle, glabrous or with a few scattered hairs. Calyxes with 3 or 5 acute sepals, villous. Petals imbricated. (Don's Mill., i. p. 203.) A shrubby, erect, much branched plant, from Spain, with bright yellow flowers from June till August. When it was introduced is uncertain, but in 1826 it was in the Fulham Nursery, under the name of *H. algarvensis*. It is somewhat tender, but a highly ornamental species, the flowers resembling those of *Calliopsis bicolor*.

n. 15. *H. HALIMIFOLIUM* Willd. The Sea-Purslane-leaved Helianthemum, or Sun Rose.

Identification. Willd. Enum., 569; Swt. Cist., t. 4; Don's Mill., i, p. 303.

Synonymes. *Cistus halimifolius* Lin. Sp., 738; *Cistus folio Halimi*, l., *Chus. Hist.*, 1. p. 71.

Engraving. Swt. Cist., t. 4.

Spec. Char., &c. Erect, branched. Branches leprously white at the top, as well as the leaves, on both surfaces. Leaves on very short footstalks, ovate-oblong, tapering to the base. Peduncles long, branched, somewhat panicled, leprously white. Calyx leprous, 5-sepaled, two outer ones very narrow, linear. (*Don's Mill.*, i. p. 303.) An erect bush, growing to the height of 3 ft.; found in Spain and Portugal by the sea-side, and cultivated in England since 1656; producing beautiful yellow flowers, spotless, or each marked with a small dark bloody spot at the base, in July and August. It is somewhat tender during winter, but grows freely during summer, and ripens abundance of seeds. Plants of it were in Colvill's Nursery in 1826. De Candolle notices a variety with obtuse leaves.

§ ii. *Lecheïdes* Dec. Prod., i. p. 269.

Derivation. From *Lechè*, and *eidōs*, appearance; plants with the habit of some species of *Lechè*.

Sect. Char. Calyx 5-sepaled, 2 outer sepals narrow, linear, 3 inner ones acute, with scarious margins. Petals yellow. Style almost wanting, or very short, erect. Stigma large, capitate. Ovary triangular. Capsule smooth, shining, 3-valved, 1-celled. Seeds rufescent, small. Stems herbaceous or suffruticose, ascendant or erect, usually dichotomous. Lower leaves opposite, cauline ones alternate, feather-nerved, on short footstalks or sessile, without stipules. (*Don's Mill.*, i. p. 303.) The species included in this section have generally very small flowers; and, as remarked below, their flowers are frequently apetalous.

A. *Peduncles many-flowered. Flowers small, crowded.*

n. 16. *H. CORYMBO'SUM* Michx. The corymbose-flowered Helianthemum, or Sun Rose.

Identification. Michx. Fl. Bor. Amer., 1. p. 307; Don's Mill., 1. p. 303.

Spec. Char., &c. Suffruticose, branched, erect. Branches dichotomous, rather pubescent, somewhat tomentously cinereous at the top. Cauline leaves alternate, oblong-lanceolate, bluntnish; under surface clothed with woolly tomentum; upper leaves with revolute margins. Corymbs fastigate, crowded. Calyx tomentously hairy, canescent; outer sepals linear, blunt; inner ones ovate acute, somewhat shorter than the capsule. (*Don's Mill.*, i. p. 303.) This species is one of the few belonging to the order which are natives of America, having been found by Michaux in New Jersey and Georgia. It grows to the height of about 1 ft., and produces its (♀) yellow flowers in July and August. We are not aware of its having been introduced into England.

n. 17. *H. GLOMERA'TUM* Lag. The glomerate-flowered Helianthemum, or Sun Rose.

Identification. Lag. in Litt.; Swt. Cist., t. 110; Don's Mill., 1. p. 303.

Synonymes. *Cistus glomeratus* Lag. *Gen. et Spec.*, p. 16; the cluster-flowered Helianthemum.

Engravings. Swt. Cist., t. 110; and our fig. 69.

Spec. Char., &c. Suffruticose, erect, somewhat dichotomous. Branches rather tomentously cinereous. Leaves lanceolate-oblong, tapering to the base; under surface hoary. Racemes axillary or terminal, many-flowered, smaller than the leaves. Flowers glomerate. (*Don's Mill.*, i. p. 303.) Found wild near Acapulco and Cimupan, in New Spain. Seeds of it were brought to England from Mexico, by Mr. Bullock, in 1823. It is a low but erect undershrub, scarcely reaching 1 ft. in height, and producing very small or apetalous flowers, in July and August. Mr. Sweet remarks that all the species belonging to this section (*Lecheïdes*), when they flower in the spring and early in the summer, produce flowers with petals; whereas, when they



flower in autumn, they are apetalous. The species in question (*H. glomeratum*), he says, had, in his garden, its shoots killed back a good way in winter, so that it did not flower till the autumn, and was, therefore, apetalous, like *fig.* 69. Had the plant been protected during the winter, it would, most probably, he says, have produced plenty of perfect flowers in spring, as is the case with *H. polygalæfolium*, *H. brasiliense*, *H. carolinianum*, and the other American species.

B. Peduncles 1-flowered, bractless, situated on the Branches.

α. 18. *H. BRASILIËNSE Pers.* The Brazilian Helianthemum, or Sun Rose.

Identification. Pers. Eneb., 2. p. 77.; Swt. Cist., t. 43.; Don's Mill., 1. p. 303.

Synonyms. *Cistus brasiliënsis Lam. Dict.*, 2. p. 22.; *Cistus alternifolius Vahl. Symb.*, 1. p. 58.

Engravings. Swt. Cist., t. 43.; and our *fig.* 70.

Spec. Char., &c. Suffruticose. Branchlets simple, hairy. Leaves ovate-oblong, acute, sessile, hairy. Peduncles and calyx hairy, caescent. Peduncles solitary, 1-flowered, inner sepals ovate, acuminate. (*Don's Mill.*, 1. p. 303.) A low shrub, found on the mountains of Brazil, introduced in 1823, and producing bright yellow flowers in June and July. It scarcely reaches 1 ft. in height, and might almost be considered as herbaceous. It is rather tender, and is best preserved when grown in pots, and taken into a frame or green-house during the winter. Plants of it were in the Botanic Garden at Chelsea, in 1826, where it flowered in the spring of that year, with petals, as in *fig.* 70., and in the autumn of the same year, without petals.



Other Species belonging to this Division of Lecheïodes.

H. polygalæfolium Swt. Cist., t. 11., from Brazil, in 1823, with white flowers, and growing to the height of half a foot. *H. astylum* Moc. and Sesse, a native of New Spain, has not been introduced; and *H. tripetalum* and *H. obcordatum* Moc. and Sesse, both from Mexico, are also but little known to botanists, and not in cultivation in British gardens.

§ iii. *Tuberaria Dec. Prod.*, 1. p. 270.

Derivation, unknown.

Spec. Char. Calyx 5-sepaled, 2 outer sepals smaller or larger, usually spreading. Petals yellow, often marked with a dark purple spot at the base of each, entire, denticulated, serrated. Stamens numerous, much longer than the pistil. Style straight, almost wanting. Stigma capitate. Capsule 3-valved. Seeds minute, yellowish. Roots woody or herbaceous. Stems erect or ascendent. (*Don's Mill.*, 1. p. 304.) There is only one ligneous species in this section.

α. 19. *H. LIGNOSUM Swt.* The woody Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 46.; Swt. Hort. Brit., p. 469. No. 88.; Don's Mill., 1. p. 304.

Engravings. Swt. Cist., t. 46.; and our *fig.* 71.

Spec. Char., &c. Stem tetragonal, shrubby, clothed with rough scaly bark. Branches ascending, covered with hispid hairs. Leaves ovate-oblong, ending in the petiole, 3-nerved, also beset with hispid hairs, caescent; under surface nerved, upper surface furrowed; floral leaves sessile, glabrous, oblong-lanceolate, uppermost ones alternate. Pedicels few, furnished with bractes at the base, rather paniced, about the length of the calyx. Petals

obovate, distinct, spreading. (*Don's Mill.*, i. p. 304.) A shrub about a foot high, a native of the south of Europe, producing its yellow flowers in July and August. Introduced in 1809, by Mr. George Don, in whose father's garden, at Forfar, it was cultivated for many years; but whence he obtained the seeds Mr. Don is uncertain. It is a very curious species, and merits a place in collections of the genus.



§ iv. *Eriocarpum* Dec. Prod., i. p. 273.

Derivation. From *erion*, wool, and *karpos*, a fruit; because the capsules are pilose.

Sect. Char. Calyx of 5 sepals. Sepals beset with silky hairs on the outside, or rather tomentose, shining on the inside; the 2 outer ones minute, linear, the 3 inner ones ovate, furnished with 4—5 stripes. Petals a little longer than the calyx. Style erectish, bent at the base. Ovary pilose, or villous. Capsule pilose. Seeds numerous, rufescent, small. Subshrubs with round branches, younger ones clothed with cinereous pubescence. Leaves opposite and alternate, bluntish; under surface cano-cinereous. Stipules linear, shorter than the footstalks. Racemes secund, small, opposite the leaves. Flowers crowded, small, sessile, or larger on short pedicels. (*Don's Mill.*, i. p. 306.) Erect and trailing undershrubs.

α. 20. *H. LIPII* Pers. Lippi's Helianthemum, or *Sun Rose*.

Identification. Pers. Ench., 2. p. 78.; Vahl. Symb., 1. p. 39.; Don's Mill., 1. p. 306.

Synonymy. *Cistus Lippii* Lin. Mant., 245.

Spec. Char., &c. Stem erect, pubescent, whitish, somewhat bifid, or dichotomous. Leaves opposite and alternate, on short footstalks, elliptic-lanceolate, or linear, oblong, obtuse, rather scabrous, glaucescent; under surface canescent; stipules narrow, erect, of the length of the footstalks. Racemes short. Flowers sessile, crowded, bracteate at the base. Bractees very minute. (*Don's Mill.*, i. p. 306.) A shrub, a native of Egypt, brought to England in 1820, growing to the height of 1 ft., and producing its yellow flowers in June or July.

α. 21. *H. SESSILIFLORUM* Pers. The sessile-flowered Helianthemum, or *Sun Rose*.

Identification. Pers. Syn., 2. p. 78.; Don's Mill., 1. p. 306.

Synonymy. *Cistus sessiliflorus* Desf. *Fl. Atl.*, 1. p. 427. t. 106.

Engraving. Desf. *Fl. Atl.*, 1. t. 106.

Spec. Char., &c. Erect, much branched. Branches pubescent. Leaves opposite and alternate, linear, clothed with very short cinereous tomentum, with revolute margins. Stipules linear, small. Racemes short. Flowers sessile, furnished with minute bractees. (*Don's Mill.*, i. p. 306.) A shrub, growing from 1 ft. to 2 ft. high, in the north of Africa, on arid hills. It produces its yellow flowers in July and August, but has not yet been introduced into England.

α. 22. *H. KAIRIUM* Del. The Cairo Helianthemum, or *Sun Rose*.

Identification. Del. *Fl. Ægypt.*, t. 31. f. 2.; Don's Mill., 1. p. 307.

Synonymy. *Cistus stipulatus* β. *Forsk. Ægypt.*, 101.

Engraving. Del. *Fl. Ægypt.*, 93. t. 31. f. 2.

Spec. Char., &c. Stem much branched, twisted at the base. Branches ascendant. Lower leaves opposite, the rest alternate, obovate, with revolute margins, hoary, stipulate; under surface nerved. Racemes secund. Flowers on short pedicels. Pedicels and calyxes villous. Sepals acute. Capsule oblong, villous. (*Don's Mill.*, i. p. 307.) A shrub 1 ft. high, brought from Egypt in 1820, and producing its yellow flowers in June and July.

§ v. *Fumana* Dec.

Derivation unknown.

Sect. Char. Calyx twisted at the apex before expansion, 5-sepaled; 2 outer sepals narrow, small; 3 inner ones ovate, acuminate, 4—5-veined, with scarious margins. Petals yellow, small, almost twice the length of the sepals. Stamens few. Style straight, rather longer than the stamens; when in flower oblique, after flowering erectish. Stigma capitate, fringed, somewhat 3-lobed. Capsule 3-valved, open, spreading. Seeds few, blackish or rufescent, angular. Stems suffruticose. Leaves linear, sessile,

or subsessile, narrow. Pedicels 1-flowered, drooping before the expansion of the flower; when in flower erect, but afterwards reflexed.

A. *Leaves alternate, without Stipules.*

23. II. *Fumàna* Mill. The *Fumana Helianthemum*, or *Sun Rose*.

Identification. Mill. Dict., No. 6.; Don's Mill., 1. p. 307.

Synonymes. *Cistus Fumàna* Lin. Sp., 740., Jacq. Aust., t. 252.; *Cistus humilis*, seu *Chamaecistus Ericæ folio luteus erectior* Bauh. Pin., 465., Magn. Bot., 69.

Engraving. Jacq. Aust., t. 252.; Swt. Cist., t. 16.; and our fig. 72.

Varieties. De Candolle distinguishes three forms of this species: *H. F. major*, *H. F. minor*, and *H. F. virgata*.

Spec. Char., &c. Stem branched, twisted, rather diffuse, erectish; lower branches procumbent. Leaves alternate, linear, with pilose, roughish, rather involute margins; lower leaves short, crowded, upper ones scattered and longer. Peduncles solitary, 1-flowered, rarely rameal, usually almost opposite the leaves, or terminal, longer than the leaves. Capsules open, naked. (*Don's Mill.*, i. p. 307.) A shrub 1 ft. high, a native of the south of France, of Switzerland, Italy, Spain, and Portugal; and cultivated in England since the year 1752. It flowers in June and July, and is distinguished by its heath-like leaves and ovate petals. It is a very desirable species for growing in pots, or for the south side of rockwork. Sweet says it is often confused with *H. procumbens*, but is readily distinguished when both are growing together. The true *H. Fumàna* was in the Chelsea Botanic Garden in 1825.



24. II. *PROCU'MBENS* Dun. The procumbent *Heath-like Helianthemum*, or *Sun Rose*.

Identification. Dun. ined., and Dec. Prod., 1. p. 275.; Swt. Cist., t. 68.; Don's Mill., 1. p. 317.

Synonymes. *Cistus humilis*, sive *Chamaecistus Ericæ folio humilior*, Magn. Bot., p. 69.

Engravings. Barrel. Icon., t. 445.; Swt. Cist., t. 68.

Spec. Char., &c. Stem procumbent, branched. Branches elongated, younger ones hoary. Leaves alternate, linear, rather lax, with the margins pilose, as well as the under surface; strigose. Peduncles almost axillary, shorter than the leaves. Capsules open, bearing the seeds. (*Don's Mill.*, i. p. 307.) A native of the south of France, Italy, and Tauria; and cultivated in the Chelsea Botanic Garden in 1825, where it produced its small yellow flowers from June to August. Uses and culture as in the preceding species.

B. *Leaves alternate, stipulate.*

25. II. *ARA'BICUM* Pers. The *Arabian Helianthemum*, or *Sun Rose*.

Identification. Pers. Ench., 2. p. 80.; Don's Mill., 1. p. 307.

Synonymes. *Cistus arabicus* Lin. Sp., 745., Smith's Fl. Græc., t. 503.; *Cistus ferrugineus* Lam. Dict., 2. p. 25.; *Cistus Savi* Bertol.; *H. viscidulum* Stev.

Engravings. Swt. Cist., t. 97.; Smith's Fl. Græc., t. 503.

Spec. Char., &c. Stem hairy, ascendent. Branches twiggy, leaves alternate, linear-oblong, hairy, almost sessile. Peduncles solitary, 1-flowered, almost opposite the leaves, rameal or terminal. Calyxes hairy. (*Don's Mill.*, i. p. 307.) A native of Arabia, Spain, and Italy, where it is a trailing shrub, seldom exceeding half a foot in height; though it acquires double that height in British gardens, where it produces its yellow flowers in June and July. It was introduced before 1826, as plants of it flowered in that year in the Chelsea Botanic Garden.

26. II. *LÆ'VIPES* Willd. The smooth-peduncled *Helianthemum*, or *Sun Rose*.

Identification. Willd. Enum., 570.; Don's Mill., 1. p. 307.

Synonymes. *Cistus lævipes* Lin. Sp., 739., Jacq. Hort. Schönb., t. 158., Cav. Icon., 2. p. 56. t. 173., Ger. Gallo-Pror., p. 294. No. 6. t. 14.

Engravings. Swt. Cist., t. 24.; Jacq. Schönb., t. 158.; Cav. Icon., 2. t. 173.; Jacq. Ger. Gallo-Pror., t. 14.

Spec. Char., &c. Stem ascendent. Leaves stipulate, setaceous, glaucous, smoothish. Buds leafy, axillary. Stipules long, filiform. Peduncles long, disposed in secund racemes. Pedicels glabrous, and bracteate at the base. Calyxes hairy. (*Don's Mill.*, i. p. 307.) A native of the south of Provence, Spain, and Dalmatia, on rocks exposed to the sun, where it forms an under-shrub 1½ ft. in height, and produces its yellow flowers from June to August. It is an elegant little plant, but rather tender during winter. Plants of it were in Mr. Colvill's Nursery in 1825.

C. Leaves opposite and alternate, furnished with Stipules.

27. *H. LÆVE Pers.* The smooth Helianthemum, or *Sun Rose*.

Identification. Pers. Ench., 2. p. 78.; *Don's Mill.*, i. p. 307.

Synonymes. *Cistus lævis Cav. Icon.*, 2. p. 35. t. 145. f. 1.

Engravings. Cav. Icon., 2. t. 145. f. 1.

Spec. Char., &c. Stem erectish, glabrous, branched. Branches erect. Leaves linear, sessile, glabrous, with revolute margins, keeled, opposite; upper ones alternate, stipulate. Stipules long, awl-shaped. Peduncles solitary, 1-flowered, subterminal. Calyxes smooth. (*Don's Mill.*, i. p. 307.) Found wild on the hills of Spain, and raised in England from seeds sent to the Chelsea Botanic Garden in 1826. It grows 1 ft. high, and produces its yellow flowers in June and July.

28. *H. VIRIDE Tenore.* The green-leaved Helianthemum, or *Sun Rose*.

Identification. Tenor. Prod. Fl. Neap., p. 31.; *Don's Mill.*, i. p. 308.

Spec. Char., &c. Stem ascendent, glabrous. Leaves opposite, linear, with revolute margins, glabrous, green, somewhat mucronate, stipulate. Stipules awl-shaped, much smaller than the leaves. Peduncles racemose, beset with clammy hairs, as well as the calyx. (*Don's Mill.*, i. p. 308.) Introduced from Sicily in 1825, where it is a shrub 1 ft. in height, producing yellow flowers in June and July.

29. *H. JUNIPERINUM Lag.* The Juniper-like Helianthemum, or *Sun Rose*.

Identification. Lag. in Litt., and Dec. Prod., i. p. 275.; *Don's Mill.*, i. p. 308.

Synonymes. *Cistus juniperipes Durand, Gouan Fl. Monsp.*, 263. ?; *Cistus mauritanicus Thib. ined.*, *Barrel. Icon.*, t. 443.

Engraving. Barrel. Icon., t. 443.

Spec. Char. &c. Stem ascendent, branched. Leaves linear awl-shaped, ciliated, mucronate, flat, with rather revolute margins, opposite; upper leaves alternate. Stipules awl-shaped, upper ones longest. Peduncles racemose, and are, as well as the calyxes, clothed with clammy hairs. (*Don's Mill.*, i. p. 308.) A shrub 1 ft. in height, introduced from the south of Europe in 1800, and producing its yellow flowers from June to August.

30. *H. BARRELIERI Tenore.* Barrelier's Helianthemum, or *Sun Rose*.

Identification. Tenor. Prod. Fl. Neap., p. 31.; *Don's Mill.*, i. p. 308.

Engravings. Barrel. Icon. rar., 416.; *Bot. Mag.*, t. 2371.

Spec. Char., &c. Stem erect. Branches villously pubescent. Leaves linear-oblong, narrowed at the base, pubescent, with revolute and ciliated margins, opposite; upper leaves alternate. Stipules linear awl-shaped, mucronate, erect. Peduncles racemose, few-flowered, and are, as well as the calyxes, beset with clammy hairs. (*Don's Mill.*, i. p. 308.) Found wild in Italy and Spain, where it grows 1 ft. high. It was introduced in 1820, and produces yellow flowers from June to August. It is rather tender.

31. *H. THYMIFOLIUM Pers.* The Thyme-leaved Helianthemum, or *Sun Rose*.

Identification. Pers. Ench., 2. p. 79.; *Don's Mill.*, i. p. 308.

Synonymes. *Cistus thymifolius Lin. Sp.*, 743., *Smith's Fl. Græc.*, t. 500., *Barrel. Icon. rar.*, t. 444.; *H. glutinosum β, Fl. Fr.* 4. p. 821.

Engravings. Sat. Cist., t. 102.; *Sm. Fl. Gr.*, t. 500.; *Barrel. Icon. rar.*, t. 444.; *Fl. Fr.*, 4. p. 821., under *H. glutinosum β*.

Spec. Char., &c. Stems procumbent. Branches pubescent. Leaves almost linear, very short, pubescent, opposite, upper ones alternate. Stipules mucronate, erect. Peduncles few-flowered, clothed with clammy hairs. (*Don's Mill.*, i. p. 308.) A native of Spain, which has been in cultivation in British gardens since 1658. It is a pretty little dwarf thyme-looking plant, and produces abundance of small bright yellow flowers from June to August; and it often ripens seeds. It is one of the most desirable of the Cistaceæ for being kept in pots.

32. *H. GLUTINOSUM Pers.* The clammy Helianthemum, or *Sun Rose*.

Identification. Pers. Ench., 2. p. 79.; *Don's Mill.*, i. p. 308.

Synonymic. *Cistus glutinosus Lin. Mant.*, 246.

Engravings. Swt. Cist., t. 83.; *Barrel. Icon. rar.*, 512. t. 415. ?; *Cav. Icon.*, 2. t. 145. f. 2.

Spec. Char., &c. Stem ascendent. Branches clothed with clammy hairs, somewhat cinereous. Leaves almost linear, with revolute margins, villous,

clammy, somewhat cinereous, opposite, upper ones alternate. Lower stipules minute, the rest long and loose. Peduncles and calyxes villous, clammy. Petals distinct. (*Don's Mill.*, i. p. 308.) Found within the south of France and Spain, and introduced in 1790. It is a slender plant, growing about 1 ft. in height, and producing very small pale yellow flowers from May to September. It is a distinct sort; and, though not very ornamental, it is valuable on account of its flowering the whole summer. Its flowers are very fugacious, the petals expanding in the morning, and dropping before the middle of the day.

§ vi. *Pseudo-Cistus* Dec. Prod., i. p. 276.

Derivation. From *pseudēs*, false, and *Cistus*; false cistus.

Sect. Char. Calyx of 5 sepals; outer sepals narrow, minute; inner ones 4-veined. Petals yellow, small, scarcely twice the length of the sepals. Style twisted at the base, and bent inwards at the apex, usually shorter than the stamens, rarely longer. Stigma capitate, 3-lobed. Capsule small. Seeds few, rather rufescent. Perennial herbs or subshrubs. Leaves stalked, feather-nerved, opposite, usually without stipules, rarely with stipules at the summits of the branches. Flowers secund, racemose, or panicled. Pedicels bracteate at the base, recurved before flowering, when in flower erect, but afterwards reflexed. Bracteas sessile, linear-lanceolate. (*Don's Mill.*, i. p. 308.) Evergreen undershrubs, bushes, or trailers, of the smallest size.

2. 33. II. MO'LLIS *Pers.* The soft-leaved Helianthemum, or Sun Rose.

Identification. *Pers. Ench.*, 2. p. 76.; *Don's Mill.*, 1. p. 308.

Synonymes. *Cistus mollis Cav. Icon.*, 3. p. 31. t. 262. f. 2.

Engravings. *Cav. Icon.*, t. 262. f. 2.

Spec. Char., &c. Suffruticose. Branches almost simple, pilose. Leaves roundish-ovate, obtuse, stalked, hairy, tomentose on both surfaces, soft. Racemes simple, and arc, as well as the calyxes, hairy, tomentose, cinereous. (*Don's Mill.*, i. p. 308.) A native of Spain, whence it was introduced in 1817; grows 1 ft. high, and produces its yellow flowers from June to August.

2. 34. II. ORIGANIFOLIUM *Pers.* The Marjoram-leaved Helianthemum, or Sun Rose.

Identification. *Pers. Ench.*, 2. p. 76.; *Don's Mill.*, 1. p. 308.

Synonymes. *Cistus origanifolius Lam. Dict.*, 2. p. 20., *Cav. Icon.*, 3. p. 31. t. 262. f. 1.

Engraving. *Cav. Icon.*, 3. t. 262. f. 1.

Spec. Char., &c. Stem suffruticose, di-tri-chotomous. Leaves stalked, ovate, pilose on both surfaces. Racemes short, terminal. Petals scarcely longer than the calyx. (*Don's Mill.*, i. p. 308.) A trailer, a native of Spain, whence it was introduced in 1795. The flowers are exceedingly small; but they are produced in abundance in June and July.

2. 35. II. DICHO'TOMUM *Dunal.* The dichotomous-branched Helianthemum, or Sun Rose.

Identification. *Dunal ined.*, and *Dec. Prod.*, 1. p. 276.; *Don's Mill.*, 1. p. 308.

Synonymes. *Cistus dichotomus Cav. Icon.*, 33. p. 2. t. 263. f. 1.

Engraving. *Cav. Icon.*, 3. p. 32. t. 263. f. 1.

Spec. Char., &c. Suffruticose. Branches dichotomous, smoothish. Leaves minute, ovate, acute, glabrous, with revolute margins, on short footstalks. Racemes slender, few-flowered. (*Don's Mill.*, i. p. 308.) A native of Spain, whence it was introduced in 1826. It is a neat little prostrate shrub, with small leaves, having the appearance of those of *Thymus Piperella*; and exceedingly small flowers, hardly the size of those of *Spérgula nodosa*, but of a deep yellow. They appear in the beginning of June, and continue till the end of August.

2. 36. II. OELANDICUM *Dec.* The Oeland Helianthemum, or Sun Rose.

Identification. *Dec. Fl. Fr.*, 4. p. 817.; *Don's Mill.*, 1. p. 308.

Synonymes. *Cistus oelandicus Lin. Sp.*, 741.; *Chamaecistus*, ii., *Clus. Hist.*, p. 73. ic.

Engraving. *Swt. Cist.*, t. 85.

Spec. Char., &c. Stem suffruticose, procumbent, branched. Leaves lanceolate-elliptical, bluntish, green on both surfaces, usually glabrous, sometimes ciliated, stalked; upper leaves sessile. Racemes simple, few-flowered.

Calyx somewhat globose-ovate. (*Don's Mill.*, i. p. 308.) Found in the Alps of France, Switzerland, and Austria; introduced in 1816; and, in our gardens, a low trailing shrub, producing yellow flowers from June to August. It is nearly related to *H. alpestre*, but has narrower leaves and smaller flowers, and is of weaker growth. Plants were in the Chelsea Botanic Garden in 1828.

2. 37. *H. PULCHÉLLUM* *Swt.* The neat *Helianthemum*, or *Sun Rose*.

Identification. *Swt. Cist.*, t. 74.; *Don's Mill.*, i. p. 308.

Synonymes. *H. alpestre Spreng. Syst.*, 2. p. 590.

Engraving. *Swt. Cist.*, t. 74.

Spec. Char., &c. Stem suffruticose, procumbent, branched. Branches clothed with hoary tomentum. Leaves roundish or ovate, obtuse, upper surface green, beset with hispid hairs; under surface clothed with hoary tomentum, with the margins a little revolute. Racemes simple. Calyxes pilose, hoary. Petals imbricate. (*Don's Mill.*, i. p. 308.) A native of Germany, introduced in 1820, and producing its yellow flowers from June to August. A neat little prostrate shrub, hardy, and of the easiest culture. Plants were in the Roehampton Nursery in 1828.

2. 38. *H. ALPE'STRE* *Dunal.* The Alpine *Helianthemum*, or *Sun Rose*.

Identification. *Dunal ined.*, and *Dec. Prod.*, 1. p. 276.; *Don's Mill.*, i. p. 308.

Synonymes. *Cistus alpestris Crantz Austr.*, p. 103. t. 6. f. 1.; *Wahl. Hede.*, p. 103.; *Cistus celandicus Jacq. Austr.*, t. 399.

Engravings. *Crantz Austr.*, t. 103. t. 6. f. 1.; *Jacq. Austr.*, t. 399.

Varieties. *De Candolle* records *H. a. glabratum*, *H. a. elongatum*, and *H. a. canescens*.

Spec. Char., &c. Stem suffruticose, procumbent, branched. Branches pilosely hairy. Leaves greenish on both surfaces, oblong-elliptical, rather glabrous, or with hairs in fascicles, stalked; upper leaves almost sessile. Pedicels and calyxes pilosely hairy. Hairs cinereous. (*Don's Mill.*, i. p. 308.) A native of Germany, Switzerland, France, and Italy, on rocks. It was introduced into England in 1818, and produces its yellow flowers, large for the size of the leaves, in July and August. It is an elegant little plant, and quite hardy. It was in Mr. Colvill's Nursery, Chelsea, in 1824.

2. 39. *H. PENICILLA'TUM* *Thib.* The pencilled *Helianthemum*, or *Sun Rose*.

Identification. *Thib. ined.*, and *Dec. Prod.*, 1. p. 277.; *Don's Mill.*, i. p. 309.

Synonymes. *Cistus echioides Lam. Dict.*, 2. p. 21.; *Cistus anglicus Lin. Mant.*, 245.?

Spec. Char., &c. Suffruticose. Branches procumbent, long, hispid. Leaves green, with the leaves on both surfaces hispid, as well as the margins; lower leaves stalked, ovate, smaller; upper ones linear-oblong, almost sessile. Racemes simple, and are, as well as the calyxes, hispid. Flowers minute. (*Don's Mill.*, i. p. 309.) A native of France and Spain, and introduced in 1826. It is a trailing plant, with the habit of *Echinopsernum Lappula*. Introduced in 1826, and producing its yellow flowers from June to August.

2. 40. *H. OBOVA'TUM* *Dunal.* The obovate-leaved *Helianthemum*, or *Sun Rose*.

Identification. *Dunal ined.*, and *Dec. Prod.*, 1. p. 277.; *Don's Mill.*, i. p. 309.

Synonymes. *Cistus italicus Lin. Sp.*, 740.

Spec. Char., &c. Suffruticose. Branches spreading, somewhat dichotomous, clothed with cinereous tomentum towards the apex. Leaves obovate, or oblong-obovate, green on both surfaces, ciliated, pilosely strigose; lower leaves minute. Racemes simple, 3-flowered. Bractees green. Calyxes pilose, cinereous. (*Don's Mill.*, i. p. 309.) Found in Spain, near Aranjuez, where it produces its yellow flowers in June and July. It was introduced in 1826; and, in British gardens, is a trailing plant, about 1 ft. in height.

2. 41. *H. ITA'LICUM* *Pers.* The Italian *Helianthemum*, or *Sun Rose*.

Identification. *Pers. Ench.*, 2. p. 76.; *Don's Mill.*, i. p. 309.

Synonymes. *Cistus italicus Lin. Sp.*, 740.; *Cistus marifolius Bieb. Fl. Taur. Cauc.*, 2. p. 8.

Engraving. *Barrel. Icon. rar.*, 510. t. 566.

Varieties. *De Candolle* distinguishes three forms of this species, viz. *H. i. strigosum*, *H. i. candidissimum*, and *H. i. albidum*.

Spec. Char., &c. Suffruticose. Branches simple, erect, long, pilosely tomentose. Leaves pilosely hispid; hairs strigose, appressed; lower leaves ovate, smaller; upper ones lanceolate, oblong or oblong-linear. Racemes simple, and are, as well as the calyxes, pilosely hispid, caescent. (*Don's Mill.*, i. p. 309.) A native of the Mediterranean, on dry hilly surfaces. Introduced in 1799, and producing its yellow flowers from July to September. It is a glaucous-looking trailing plant, seldom exceeding 1 ft. in height.

2. 42. *H. VINEALE Pers.* The Vineyard Helianthemum, or Sun Rose.

Identification. Pers. Ench., 2. p. 77.; Don's Mill., 1. p. 309.

Synonymes. *Cistus vinealis Willd. Sp.*, 2. p. 1195.

Engraving. Swt. Cist., t. 77.

Spec. Char., &c. Suffruticose, procumbent. Branches ascendent, pilosely tomentose, caescent. Leaves ovate-oblong; upper surface green, strigosely pilose, under surface tomentose, hoary. Racemes simple, few-flowered, and are, as well as the calyxes, pilosely tomentose, caescent. (*Don's Mill.*, i. p. 309.) A native of the south of Germany, Switzerland, France, and Spain; and introduced in 1817. It is a trailing plant, growing to about 1 ft. in height, and producing its pale yellow flowers in June and July. Sweet says that it is covered with flowers during a great part of the summer. After flowering, most of its leaves become quite silvery on both sides. It requires very little protection. Plants were in the Chelsea Botanic Garden, raised from seeds received from Germany, in 1828.

2. 43. *H. CANUM Dunal.* The hoary Helianthemum, or Sun Rose.

Identification. Dunal ined., and Dec. Prod., 1. p. 277.; Don's Mill., 1. p. 309.

Synonymes. *Cistus canus Lin. Sp.*, 740.; *Chamaecistus*, iii., *Clus. Hist.*, p. 74.

Engravings. Swt. Cist., t. 56.; Jacq. Aust., t. 277.; All. Ped., No. 1664. t. 45.

Spec. Char., &c. Stem suffruticose, procumbent, branched, ascendent, pilosely tomentose, hoary. Leaves obovate, ovate, ovate-oblong, or elliptical, pilose; upper surface green, under surface somewhat tomentose, hoary. Racemes simple. Pedicels and calyxes pilose, caescent. Petals distinct. (*Don's Mill.*, i. p. 309.) Found wild in the south of France and Germany, and probably also in Britain. It is a procumbent shrub, which has been cultivated in British gardens since 1772, and produces its yellow flowers in June and July. It is nearly related to *H. alpestre*, and *H. vineale*; and, according to Sweet, is sometimes mistaken for *H. marifolium*. Plants were in Colvill's Nursery, at Chelsea, in 1827.

2. 44. *H. MARIFOLIUM Dec.* The Maroon-leaved Helianthemum, or Sun Rose.

Identification. Dec. Fl. Fr., 4. p. 817.; Don's Mill., 1. p. 309.

Synonymes. *Cistus marifolius Lin. Sp.*, 741., but not of Bieb. Fl. Taur. Cauc., *Burcel. Icon. var.*, 521. t. 441.

Engraving. Barrel. Icon. rar., 521. t. 441.

Spec. Char., &c. Suffruticose, procumbent. Leaves without stipules, stalked, ovate-cordate, or ovate, acutish; upper surface green, pilose, under surface hoary. Racemes solitary, simple, few-flowered, terminal. (*Don's Mill.*, i. p. 309.) Found wild in the south of France, and in Spain and Italy; and introduced in 1817. It is a trailing plant, seldom exceeding 1 ft. in height, and producing its yellow flowers in June and July.

2. 45. *H. CRASSIFOLIUM Pers.* The thick-leaved Helianthemum, or Sun Rose.

Identification. Pers. Ench., 2. p. 77.; Don's Mill., 1. p. 309.

Synonymes. *Cistus glaucus Desf. Atl.*, 1. p. 418., but not of Cav.; *H. Sézte* Lag. in Litt.

Spec. Char., &c. Stem suffruticose, erect, rather glabrous. Leaves somewhat fleshy, on short foot-stalks; lower leaves ovate, acute, without stipules; upper ones oblong-linear, stipulate. Racemes short, rather umbellate. Calyxes pilose at the base. (*Don's Mill.*, i. p. 309.) Found wild in Barbary and Spain, and introduced into England in 1818. It grows to 1 ft. in height, and produces its yellow flowers from May to July. It was called *H. Sézte*, on account of its being found in Valencia, where it is called *sezte* by the common people.

2. 46. *H. PANICULATUM Dunal.* The paniced-flowered Helianthemum, or Sun Rose.

Identification. Dunal ined., and Dec. Prod., 1. p. 278.

Synonymes. *Cistus marifolius Herb. Thib.*; *Cistus mummularius var. Lag. in Litt.*; *H. sp. nova Schouw, in Litt.*

Spec. Char., &c. Suffruticose, procumbent. Branches ascendent and erect; floriferous branches long; upper part stipulate. Leaves stalked, ovate, bluntish, rarely roundish; upper surface green, under surface hoary. Racemes opposite and ternary; paniced. (*Don's Mill.*, i. p. 309.) A native of the mountains of Spain and Sicily, and introduced here in 1826. It is a procumbent plant, with very small flowers of a yellow colour, which are produced in abundance from June to August.

§ vii. *Euhelianthemum Dec. Prod.*, i. p. 278.

Derivation. From *eus*, genuine, *helios*, the sun, and *antheion*, a flower; that is to say, genuine species of helianthemum.

Sect. Char. Calyx of 5 sepals, rather twisted at the top before expansion;

outer sepals usually spreading, much smaller than the inner ones, which are usually 2- or 4-ribbed, furrowed, with scarious margins, with the inner surface shining, and with the angles generally pilose. Petals 2, 3, or 4 times longer than the calyx. Stamens numerous. Style bent at the base, but somewhat club-shaped at the apex. Stigma simple. Capsule covered by the calyx, 3-valved, 1-celled, opening at the apex. Seeds few, convex on the outside, and angular on the inside. Subshrubs, with the stems branched from the base; branches numerous, erect or procumbent, but generally ascendent. Leaves opposite, on short footstalks; lower ones smallest, usually with revolute margins, stipulate. Stipules linear-lanceolate. Racemes terminal, second, simple, curved backwards before flowering; after flowering erect, elongated. Pedicels laterally bracteate at the base, drooping before flowering; when in flower erect, after flowering recurved or reflexed. (*Don's Mill.*, i. p. 310.) Evergreen undershrubs, bushes, and trailers, of the smallest size; natives of Britain, and the southern part of the European continent. Many of the sorts are hybrids originated in British gardens.

A. *Petals yellow.*

- ♁ 47. *H. LAVANDULÆFOLIUM* Dec. The Lavender-leaved Helianthemum, or *Sun Rose*.

Identification. Dec. Fl. Fr., 4. p. 820.; *Don's Mill.*, 1. p. 310.

Synonyme. *Cistus lavandulæfolius* Lam. *Dict.*, 2. p. 25.

Engraving. Barrel. Icon., t. 288.

Varieties. De Candolle notices *H. l. syriacum*, the *Cistus syriacus* of Jacquin; and Persoon records *H. l. Thibaudi*, the *Cistus racemösus* of Cavanilles.

Spec. Char. &c. Stem suffruticose, erect, branched. Branches long, terete, canescent. Leaves oblong-linear, with revolute margins; under surface tomentose, hoary; younger leaves canescent on both surfaces. Stipules and bractees linear, acute, ciliated. Racemes 1—3-flowered, terminal. Flowers crowded. Calyxes glaucous. Sepals ciliated, outer ones minute; these become reflexed after flowering; inner sepals 2-nerved, oblique, acute. (*Don's Mill.*, i. p. 310.) A bush about 1 ft. in height, a native of the south of France, and found also in Barbary, Spain, and Syria, in dry places; producing its yellow flowers in June and July. It was introduced into England in 1739, and, probably, is now lost, or confounded with some other sort.

- ♁ 48. *H. STÆCHADIFOLIUM* Pers. The French-Lavender-leaved Helianthemum, or *Sun Rose*.

Identification. Pers. Ench., 2. p. 79. *Don's Mill.*, 1. p. 310.

Synonyme. *Cistus stæchadifolius* Brot. *Fl. Lus.*, 2. p. 270.

Spec. Char. &c. Stem erect. Branches hoary, tomentose. Leaves oblong-linear, bluntnish, somewhat tomentose on both surfaces; under surface hoary, upper surface greenish grey, with revolute margins. Stipules rather villous, linear-lanceolate. Racemes revolute before flowering. Flowers crowded. Calyxes villous. Outer sepals ciliated, green, inner ones acuminate, hoary. (*Don's Mill.*, i. p. 310.) A native of Spain and Corsica. An upright bush, introduced in 1816, and producing its yellow flowers in June and July.

- ♁ 49. *H. CRO'CEUM* Pers. The Saffron-coloured-flowered Helianthemum, or *Sun Rose*.

Identification. Pers. Ench., 2. p. 79.; *Don's Mill.*, 1. p. 310.

Synonyme. *Cistus crœceus* Desf. *Fl. Atl.*, 1. p. 422. t. 110.

Engravings. Swt. *Cist.*, t. 53.; Desf. *Fl. Atl.*, 1. t. 110.

Varieties. De Candolle (*Prod.*, i. p. 279.) records three forms of this species: one with the stipules longer than the footstalks of the leaves; another, with the stipules setaceous, shorter than the footstalks of the leaves; and the third with procumbent branches, shorter leaves, and racemes few-flowered.

Spec. Char. &c. Stem shrubby, branched, somewhat procumbent. Branches simple, erect, hoary-tomentose. Leaves rather tomentose; under surface canescent, upper surface glaucous with revolute margins; lower leaves almost round; middle ones elliptical, obtuse; upper ones, lanceolate, acutish. Stipules and bractees erect, linear, oblong, villous, rather greenish. Calyxes yellowish-glaucous, minutely pubescent. (*Don's Mill.*, i. p. 310.) A procumbent plant, with large dark yellow flowers, found in Spain and Barbary, and, according to Sweet, cultivated in the Chelsea Botanic Garden in 1826, where, growing on rockwork, it covered more than a yard in diameter, and made a grand appearance in June and July, when it was covered with flowers. It is one of the most ornamental species of the genus, and no collection ought to be without it, if it were only for planting out in borders in the summer season.

2. 50. *H. ANDERSONI* Swt. Anderson's Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 89.; Don's Mill., i. p. 310.

Engraving. Swt. Cist., t. 89.

Spec. Char., &c. Stem suffruticose, proeminent, branched. Branches ascending, canescently tomentose. Leaves oblong-lanceolate, acutish, rather tomentose, grey above, and canescent beneath, with the margins a little revolute. Stipules linear, awl-shaped, ciliated, a little longer than the petioles. Calyx tomentose. Petals imbricate. (*Don's Mill.*, i. p. 310.) A pretty and curious plant, produced from the seed of *H. erœcum*, that was fertilised by the pollen of *H. pulverulentum*, in the Chelsea Botanic Garden, in the year 1828. The seeds were self-sown in that year; in the year following, the plants produced flowers, from May to the end of November. "The flowers were very variable, some being of a bright yellow, and others, on the same plant, and sometimes on the same branch, of a pale straw colour." (*Sweet.*) It is a remarkably fast grower, an abundant flowerer, seeds freely, and is quite hardy. Sweet says, "We have named it in compliment to our respected friend Mr. William Anderson, the curator of the garden, to whom we are obliged for the opportunity of making drawings of many rare species, which we have not seen in any other collection."

2. 51. *H. NUDICAULE* Dunal. The naked-stemmed Helianthemum, or Sun Rose.

Identification. Dunal. ined., and Dec. Prod., i. p. 279.; Don's Mill., i. p. 310.

Spec. Char., &c. Stem shrubby, branched. Branches smooth at bottom, but hoary-villous at top. Leaves oblong lanceolate, with revolute margins, tomentose on both surfaces; under surface hoary, upper surface yellowish green. Stipules linear, longer than the petioles. Calyxes profoundly sulcate, hardly pubescent, with elevated pilose nerves. (*Don's Mill.*, i. p. 310.) A native of Spain, and found on mountains in the kingdom of Valentia. It was introduced in 1826, producing its yellow flowers in June and July. De Candolle doubts whether it is not only a variety of *H. erœcum*.

2. 52. *H. GLAUCUM* Pers. The glaucous Helianthemum, or Sun Rose.

Identification. Pers. Euch., 2. p. 78.; Dec. Prod., i. p. 279.; Don's Mill., i. p. 311.

Synonymes. *Cistus glaucus* Cav. *Icon.*, 3. p. 31. t. 261., but not of Desf.

Engraving. Swt. Cist., t. 111.

Varieties. Two forms of this species are mentioned by De Candolle, viz.: *H. g. acutiusculum*, the upper leaves of which are oblong, and rather acuminate, with the upper surface glaucescent; and *H. g. obtusiusculum*, the upper leaves of which are oblong elliptical, bluntish, the upper surface roughish and green.

Spec. Char., &c. Stem suffruticose, branched. Branches ascendant, hoary-tomentose, hispid at the top. Leaves ciliated on their margins, scarcely revolute, tomentose on both surfaces; under surface hoary, upper surface greenish glaucous. Lower leaves round, the rest elliptic or lanceolate-oblong. Stipules and bractæas pubescent, green. Pedicels and calyxes beset with white hairs. (*Don's Mill.*, i. p. 311.) A suberect glaucous plant, from Spain and Italy, in 1815; and producing its small pale yellow flowers from June till August. It is a very desirable plant to keep in pots, for turning out in the borders, or on rockwork, in spring, as it is rather tender, and liable to damp off in the free soil, unless the situation is very dry and warm. Plants were in the garden of Robert Barclay, Esq., at Bury Hill, near Dorking, in 1829.

2. 53. *H. TOMENTOSUM* Dunal. The tomentose Helianthemum, or Sun Rose.

Identification. Dunal. ined., and Dec. Prod., i. p. 279.; Don's Mill., i. p. 311.; Smith's Eng. Bot., 2208.; ? Scop. Carn., t. 24.

Engraving. ? Scop. Carn., t. 24.

Spec. Char., &c. Stem suffruticose, branched. Branches elongated, ascendent, somewhat canescent. Leaves lanceolate-oblong, usually with revolute margins; under surface hoary-tomentose, upper surface smoothish, green. Calyxes furrowed, with elevated pilose nerves. (*Don's Mill.*, i. p. 311.) A trailing plant, 1 ft. in height, found on the mountains of Scotland, and also in Spain and France, producing its yellow flowers, with imbricate petals, in July.

▮ 54. *H. BARBA'TUM* Pers. The bearded-racemed Helianthemum, or Sun-Rose.

Identification. Pers. Ench., 2. p. 79.; Swt. Cist., t. 73.; Don's Mill., 1. p. 311.

Synonymes. *Cistus barbatus* Lam. Dict., 2. p. 24.

Engraving. Swt. Cist., t. 73.

Spec. Char., &c. Stem suffruticose, erect, much branched. Branches clothed with fasciated hairs. Leaves hairy, green on both surfaces; lower ones roundish-ovate, upper ones elliptical. Stipules oblong, ciliated, hairy, longer than the footstalks of the leaves. Racemes long, hairy, bearded, many-flowered. Calyxes warted, hairy. Petals crenulated, imbricate at the base. (*Don's Mill.*, i. p. 311.) A native of the south of Europe, producing its yellow flowers in June and July. It is an upright-growing shrub, and was introduced in 1820. It is a very distinct species, a free grower, and requires no protection. Plants were in the Chelsea Botanic Garden in 1828.

♣ 55 *H. LEPTOPHY'LLUM* Dunal. The slender-leaved Helianthemum, or Sun Rose.

Identification. Dunal ined., Dec. Prod., 1. p. 279.; Don's Mill., 1. p. 311.

Synonymes. *Cistus angustifolius* Lag.; *C. stoechadifolius* Hort.

Engraving. Swt. Cist., t. 50.

Spec. Char., &c. Stem suffruticose, woody, rather procumbent, branched. Branches ascendent, rather tomentose, greyish. Leaves narrow, oblong-linear, tapering into the short footstalks, with revolute margins; under surface covered with short cinereous tomentum; upper surface smoothish, green. Stipules awl-shaped, pilose, scarcely longer than the footstalks. Racemes long. Calyxes covered with long hairs. (*Don's Mill.*, i. p. 311.) A trailing plant, from Spain, in 1818, producing yellow flowers in June and July. It is a vigorous-growing plant, very distinct in its appearance, and very hardy. The petals of the flowers are large, and of a fine dark yellow. Plants were in the Chelsea Botanic Garden in 1826.

▮ 56. *H. ACUMINA'TUM* Pers. The acuminate Helianthemum, or Sun Rose.

Identification. Pers. Eneh., 2. p. 79.; Don's Mill., 1. p. 311.

Synonymes. *Cistus serpyllifolius* Balb. ined.; *Cistus acuminatus* Fiv. Fragm., 13. t. 14. f. 1.

Spec. Char., &c. Branches erect, pilose at the base and at the apex, middle naked. Leaves on long footstalks, oblong, with revolute margins, green on both surfaces, pilose; under surface rather tomentose. Stipules smoothish, linear, longer than the footstalks of the leaves. Racemes rather hairy, few-flowered, loose. Calyxes smooth, shining, transparent. (*Don's Mill.*, i. p. 311.) A native of Nice, and cultivated in British gardens in 1820; it grows to 1 ft. in height, and produces its yellow flowers in June and July.

♣ 57. *H. SERPYLLIFO'LIUM* Mill. The Wild-Thyme-leaved Helianthemum, or Sun Rose.

Identification. Mill. Dict. No. 8.; Swt. Cist., t. 60.; Don's Mill., 1. p. 311.

Synonymes. *Cistus serpyllifolius* Lin. Sp., 743.

Engraving. Swt. Cist., t. 60.

Spec. Char., &c. Stem suffruticose. Branches ascendent, glabrous at the base, and pilose at the apex. Leaves oblong-elliptical, with revolute margins; under surface hoary-tomentose; upper surface intensely green, shining, at first rather pilose, afterwards almost smooth. Stipules and bracteas green, ciliated. Calyxes canescent, with inconspicuous down, and with the nerves sparingly pilose. (*Don's Mill.*, i. p. 311.) A trailing shrub, with large yellow flowers, the petals of which are distinct. It is found on the Alps of Styria and Austria, as well as on the mountains of Spain. It was introduced into our gardens in 1731, and continues in flower from May to September. It was found wild in Somersetshire, by Mr. Sweet and some others, in 1826, in which year there were plants of it in Mr. Colvill's Nursery, Chelsea.

♣ 58. *H. VULGA'RE* Gart. The common Helianthemum, or Sun Rose.

Identification. Gart. Fruct., 1. p. 371. t. 76.; Don's Mill., 1. p. 311.

Synonymes. *Cistus Helianthemum* Lin. Sp., 1. p. 744., *Fl. Dan.*, t. 101., *Smill's Engl. Bot.*, 1321., *Curt. Fl. Lond.*, fasc. 5. t. 36.

Varieties. There is a very handsome double-flowered variety, with pale yellow flowers; and another, called Lee's new double yellow, with dark yellow flowers; both of which are in general cultivation in the nurseries. De Candolle also notices two forms of the species: one with tomentose pubescent branches, and stipules scarcely longer than the footstalks of the leaves; and another with branches glabrous at the base, but pubescent upwards, and the stipules twice or thrice the length of the petioles of the leaves.

Spec. Char., &c. Stem suffruticose, procumbent, branched, branches elongated.

Leaves scarcely revolute at the margins; under surface cinerously hoary, upper surface green, pilose, somewhat ciliated; lower leaves somewhat orbicular, middle ones ovate-elliptical, upper ones oblong. Stipules oblong-linear, ciliated, longer than the footstalks of the leaves. Racemes loose. Pedicels and calyxes pilose. (*Don's Mill.*, i. p. 311.) The stamens, if touched during sunshine, spread slowly, and lie down upon the petals. (*Smith.*) Native of dry and hilly pastures throughout Europe, and common in Britain. It was first recorded under the name of *Helianthemum vulgare* by Ray, who says that the flowers are very seldom white. Sir J. E. Smith says, "It is a variable species, but less so than authors make it." (*Eng. Fl.*, iii. p. 26.) It is a trailing shrub, flowering from May till September. The double-flowered varieties ought to be in every collection.

2. 59. *H. SURREJANUM Mill.* The Surrey Helianthemum, or *Sun Rose*.

Identification. Mill. Diet., No. 15; Swt. Cist., t. 28.; Don's Mill., i. p. 311.

Synonymic. *Cistus surrejanus Lin. Sp.*, 743., *Smith's Eng. Bot.*

Engravings. Dill. Elth., 177. t. 145. f. 174.; Swt. Cist., t. 28.; Eng. Bot., t. 2207.

Spec. Char., &c. Stem suffruticose, procumbent. Leaves ovate-oblong, rather pilose. Racemes many-flowered, terminal. Petals narrow, lanceolate, jagged. (*Don's Mill.*, i. p. 311.) Found wild in Surrey, near Croydon.

It is a procumbent shrub, with yellow flowers, the petals of which are distinct, and the calyxes pilose. It flowers from July to October. Plants were in the garden of Mrs. Dickson, of Croydon, in Surrey, in 1826.

2. 60. *H. OVATUM Dunal.* The ovate-leaved Helianthemum, or *Sun Rose*.

Identification. Dunal ined., Dec. Prod., 1. p. 280.; Don's Mill., i. p. 311.

Synonymic. *Cistus ovatus Viv. Frag.*, 1. p. 6.

Engraving. Viv. Frag., 1. t. 8. f. 2.

Spec. Char., &c. Stem suffruticose, procumbent, much branched. Branches villous. Leaves elliptic-lanceolate, tapering into the footstalks, bluntish, silky-villous on both surfaces, ciliated. Stipules somewhat longer than the footstalks of the leaves, villously ciliated. Peduncles 1-3-flowered, terminal. Calyxes rather villous. (*Don's Mill.*, i. p. 311.) A trailing plant, from the mountains between Viterbo and Ronciglione, and in the Alps about Geneva. Introduced in 1818, and producing its yellow flowers from June to August.

2. 61. *H. GRANDIFLORUM Dec.* The large-flowered Helianthemum, or *Sun Rose*.

Identification. Dec. Fl. Fr., 4. p. 821.; Don's Mill., i. p. 311.

Synonymic. *Cistus grandiflorus Scop. Carn.*, ed. 2. No. 648. t. 25.

Engravings. Swt. Cist., t. 69.; Scop. Carn., ed. 2. t. 25.; and our fig. 73.

Spec. Char., &c. Stem suffruticose, ascending. Branches hairy. Upper leaves flattish, oblong, rather pilose; upper surface green, under surface sometimes pale cinereous. Stipules ciliated, rather longer than the footstalks of the leaves. Flowers large. Calyxes rather hairy. (*Don's Mill.*, i. p. 311.) A native of the Pyrenees, from which it was introduced in 1800. It bears a close resemblance to *H. vulgare*, but is larger in all its parts; and its flowers, which appear from June to August, are considerably larger, and of a paler colour. It is quite as hardy as the indigenous sorts. Plants of it were in the Fulham Nursery in 1828.

73



2. 62. *H. OBSCURUM Pers.* The obscure Helianthemum, or *Sun Rose*.

Identification. Pers. Syn., 2. p. 79.; Don's Mill., i. p. 311.

Synonymic. *H. obscurum a Dec. Flor. Fr.*, 6 p. 624.

Spec. Char., &c. Stem suffruticose, ascending, much branched. Branches hairy. Leaves elliptical, hairy on both surfaces, greenish; upper ones elliptic. Stipules ciliated, longer than the footstalks. Racemes long. Calyxes hairy. (*Don's Mill.*, i. p. 311.) A native of Europe, in woods and dry wastes, and introduced into British gardens in 1816. It is hardly procumbent, forming a small roundish bush, about 1 ft. in height, and producing its pale yellow flowers, which are rather small, from May to August.

2. 63. *H. TAU'RICUM* Fisch. The Taurian Helianthemum, or Sun Rose.

Identification. Fisch. MSS. Swt. Cist., t. 105.; Don's Mill., 1. p. 312.

Engraving. Swt. Cist., 105.

Spec. Char., &c. Stem suffruticose, much branched, procumbent. Branches procumbent, beset with long hairs. Leaves oblong-lanceolate, with rather revolute margins, pilose on both surfaces, green above and paler beneath. Stipules lanceolate-linear, ciliated, longer than the petiole. Flowers large. Calyx shining, rather hairy. Petals imbricate. (*Don's Mill.*, i. p. 312.) A very ornamental species, from Tauria, with large pale yellow flowers, which appear from May to October. Sweet says, "It has, most probably, been confused with *H. grandiflorum* by many authors; but, when the plants are seen growing together, no two plants need appear more distinct: the present spreading flat on the ground, and extending its branches round to a great distance, and these being only slightly suffrutescent at the base; whereas *H. grandiflorum* grows upright, or has its branches spreading and ascendent, forming a neat little bushy shrub." Plants of this species were in the Fulham Nursery in 1829.

2. 64. *H. LU'CIDUM* Horn. The shining-leaved Helianthemum, or Sun Rose.

Identification. Horn. Cat. Hort. Hafn., p. 498.

Spec. Char., &c. Stem suffruticose, procumbent. Leaves stipulate, ovate, green, glossy, with revolute margins. (*Don's Mill.*, i. p. 312.) The native country of this sort is uncertain; but it is said to have been in cultivation in British gardens since 1826. The flowers are yellow, and produced in May and June.

2. 65. *H. NUMMULA'RIUM* Mill. The Money-wort-leaved Helianthemum, or Sun Rose.

Identification. Mill. Dict., No. 11.; Swt. Cist., t. 80.; Don's Mill., 1. p. 312.

Synonymes. *Cistus nummularius* α *Lin. Sp.*, 743., not of Desf. and Cav.; *H. obscurum*, β *nummularium* *Dec. Flor. Fr.*, 6. p. 624.; *H. angustifolium* of many botanic gardens.

Engraving. Swt. Cist., t. 80.

Spec. Char., &c. Stem suffruticose. Branches procumbent, hairy. Lower leaves orbicular, upper ones oblong-linear, hairy; under surface greenish cinereous. Stipules linear-oblong, twice the length of the footstalks of the leaves. Racemes and calyxes hairy. (*Don's Mill.*, i. p. 312.) A native of the south of France and of Italy, and strongly resembling *H. vulgare*; from which, according to Sweet, it is readily distinguished by its broader and flatter leaves. It has been in cultivation since 1752, producing its bright yellow flowers from June to August.

2. 66. *H. ANGUSTIFO'LIUM* Pers. The narrow-leaved Helianthemum, or Sun Rose.

Identification. Pers. Ench., 2. p. 79.; Don's Mill., 1. p. 312.

Synonyme. *Cistus angustifolius* *Jacq. Vind.*, 3. t. 53.

Engraving. *Jacq. Vind.*, 3. t. 53.

Spec. Char., &c. Stem suffruticose, diffuse. Branches rather tomentose, cinereous. Leaves on short footstalks; upper ones linear-oblong, with revolute margins, acutish; under surface clothed with canescent tomentum, upper surface rather hispid. Stipules pilose, longer than the footstalks. Racemes loose. Calyxes pubescent, rather hairy; hairs deciduous. (*Don's Mill.*, i. p. 312.) In cultivation since 1800; and, according to De Candolle, probably the same as *H. nummularium*.

2. 67. *H. HI'RТУM* Pers. The hairy Helianthemum, or Sun Rose.

Identification. Pers. Syn., 2. p. 79.; Swt. Cist., t. 109.; Dec. Prod., 1. p. 281.; Don's Mill., 1. p. 312.

Synonymes. *Cistus hirtus* *Lin. Sp.*, 744., *Smith's Fl. Græc.*, exclusive of synonyme of Barrelier, *Cav. Icon.*, 2. p. 37.

Engravings. Swt. Cist., t. 109.; *Smith's Fl. Græc.*, t. 501.; *Cav. Icon.*, 2. t. 146.

Spec. Char., &c. Stem suffruticose, branched. Branches ascendent, numerous, tomentosely hairy, cinereous. Leaves ovate or oblong, with revolute margins; under surface canescent, upper surface greenish cinereous. Stipules narrow, rather longer than the footstalks of the leaves. Calyxes densely covered with white hairs. Petals obovate, imbricate. (*Don's Mill.*, i. p. 312.) A native of Spain and of the south of France, and in cultivation in British gardens since 1759. It is a very distinct sort, readily distinguished by its large deep-yellow flowers, which appear in June and July, and by its very hairy leaves. Plants of it were in the Chelsea Botanic Garden in 1829.

Varieties. De Candolle mentions three forms of this species: *H. h. ba'licum*, *H. h. alvæum*, and *H. h. teretifolium*.

2. 68. *H. LAGA'SCÆ* Dunal. La Gasca's Helianthemum, or *Sun Rose*.

Identification. Dunal ined., and Dec. Prod., 1. p. 281.; Don's Mill., 1. p. 312.

Synonymy. *H. hirtum* var. *Lag.* in *Litt.*

Spec. Char., &c. Branches ascending, tomentosely hairy, hoary. Leaves linear, obtuse, with very revolute margins, almost terete, rather hairy, greenish. Stipules flat, scarcely pilose, twice the length of the footstalks of the leaves. Pedicels hispid, whitish. Calyxes shining, furrowed. Nerves ciliated with white hairs. (*Don's Mill.*, 1. p. 312.) A native of Spain, introduced into England in 1826, and so like the foregoing species as to be often considered a variety of it.

B. Petals white, rose-coloured, red, pale sulphur-coloured, or variegated with these Colours.

2. 69. *H. VIOLA'CEUM* Pers. The Violet-coloured-calyxed Helianthemum, or *Sun Rose*.

Identification. Pers. Syn., 2. p. 78.; Don's Mill., 1. p. 312.

Synonymy. *Cistus violaceus* Cav. *Icon.*, 2. p. 38.

Engraving. Cav. *Icon.*, 2. t. 147.

Spec. Char., &c. Stem erect or ascending, much branched. Branches opposite; branchlets slender, tomentosely hairy, hoary. Leaves small, almost linear, obtuse, with revolute margins, somewhat tomentose on both surfaces; under surface caescent. Stipules minute, pilose. Racemes few-flowered, loose. Calyxes smooth, violaceous, nervedly furrowed. (*Don's Mill.*, 1. p. 312.) A native of Spain, and in cultivation since 1826. The flowers are white, slightly tinged with violet, and they appear in June and July.

2. 70. *H. RACEMOSUM* Dunal. The racemose-flowered Helianthemum, or *Sun Rose*.

Identification. Dunal ined., and Dec. Prod., 1. p. 281.; Don's Mill., 1. p. 312.

Synonymy. *Cistus racemosus* Lin. *Mant.*, 76.?, *Lam. Dict.*, 2. p. 25., *Fakt. Symb.*, 1. p. 39., *Willd.*

Sp., 2. p. 1208.

Engraving. Swt. *Cist.*, t. 82.

Spec. Char., &c. Stem shrubby, branched. Branches erect, terete, hoary, tomentose. Leaves on short footstalks, narrow-linear, or linear-lanceolate, with revolute margins; under surface hoary, upper surface greenish, shining. Stipules awl-shaped, longer than the footstalks of the leaves. Pedicels hoary. Calyxes nervously furrowed, brownish violet. (*Don's Mill.*, 1. p. 312.) A native of Spain, Barbary, and Teneriffe, and in cultivation in the Bristol Nursery in 1828. It is a very beautiful species, with large white flowers, having imbricate crenulate petals, and the sepals marked with red or violet. It is readily distinguished, Sweet observes, from all the others of its section, by its upright growth, glossy leaves, and the red veins of its sepals. Its flowers are produced nearly the whole summer, and it requires very little protection during the winter.

2. 71. *H. FARINOSUM* Swt. The mealy-leaved Helianthemum, or *Sun Rose*.

Identification. Swt. *Cist.*, p. 18.; Don's Mill., 1. p. 312.

Synonymy. *H. racemodum* β *farinosum* Dec. *Prod.*, 1. p. 281.

Spec. Char., &c. Stem shrubby, erect, branched, tomentosely hoary. Leaves on short footstalks, linear, or lanceolate-linear, with revolute margins, hoary and powdered on both surfaces. Stipules awl-shaped, longer than the footstalks of the leaves. Calyx powdery, as well as beset with very short hairs. (*Don's Mill.*, 1. p. 312.) A native of Spain, with white flowers in June and July. It is said to have been cultivated in 1820. De Candolle considers it a variety of *H. racemodum*.

2. 72. *H. STRICTUM* Pers. The straight-branched Helianthemum, or *Sun Rose*.

Identification. Pers. *Ench.*, 2. p. 79.; Don's Mill., 1. p. 312.

Synonymy. *Cistus strictus*, Cav. *Icon.*, 3. p. 32.

Engraving. Cav. *Icon.*, 3. t. 263. f. 2.

Spec. Char., &c. Stem suffruticose, erect, branched. Branches straight, hoary-tomentose. Leaves almost sessile, very narrow, linear awl-shaped, with revolute margins, caescent. Stipules linear, setaceous. Calyxes pilose, nervously-stricted, yellowish, smoothish. (*Don's Mill.*, 1. p. 312.) A native of Spain, and in cultivation since 1820, producing white flowers in June and July.

2. 73. *H. PILOSUM* Pers. The pilose Helianthemum, or *Sun Rose*.

Identification. Pers. *Ench.*, 2. p. 79.; Dec. *Prod.*, 1. p. 282.; Don's Mill., 1. p. 312.

Synonymy. *Cistus pilosus* Lin. *Sp.*, 74. α .?; *Chamaecistus*, iv., *Clus. Hist.*, 1. p. 74.

Engraving. Swt. *Cist.*, t. 49.

Varieties. De Candolle has two forms of this species: one with linear leaves, hoary; and the other with linear leaves, oblong and shining.

Spec. Char., &c. Stem suffruticose, branched. Branches erectish. Leaves linear or linear-oblong, hoary on both surfaces, and bristly at the apex. Stipules awl-shaped. Calyxes rather pilose, nervously striated. (*Don's Mill.*, i. p. 312.) A native of Spain and the south of France, and in cultivation since 1731; producing its white flowers from May to July. According to Sweet, it is nearly related to *H. lineare*, and also to *H. apenninum*: but it differs from the former in having whiter leaves, and being more hairy; and from both, by its imbricate paper-white petals. Plants of it were in the Fulham Nursery in 1826.

2. 74. *H. LINEARE Pers.* The linear-leaved Helianthemum, or Sun Rose.

Identification. Pers. Ench., 2. p. 78.; Don's Mill., 1. p. 312.

Synonymes. *Cistus linearis Cav. Icon.*, 3. p. 8.; *Cistus pilosus Dec. Fl. Fr.*, 5. p. 823?

Engraving. Swt. Cist., t. 48.

Spec. Char., &c. Stem suffruticose. Branches elongated, ascendent, rather hoary, tomentose. Leaves linear, greenish hoary, with revolute margins. Stipules linear awl-shaped. Racemes loose, twiggly, few-flowered. Calyxes striated, glabrous, with the nerves somewhat violaceous. Sepals acute. (*Don's Mill.*, i. p. 312.) A native of Spain and the south of France; in cultivation since 1817; and producing white flowers from June to August. It requires a little protection during winter.

2. 75. *H. VIRGATUM Pers.* The twiggly Helianthemum, or Sun Rose.

Identification. Pers. Ench., p. 79.; Don's Mill., 1. p. 313.

Varieties. De Candolle mentions two forms of this species: *H. v. albiflorum*, in which the petals are white; and *H. v. roseum*, in which they are pale rose-coloured.

Spec. Char., &c. Stem suffruticose, with twiggly, hoary, ascending or erect branches. Leaves linear, hoary on the under surface, with revolute margins. Stipules linear awl-shaped. Calyxes hoary, powdery, pubescent. (*Don's Mill.*, i. p. 313.) A native of Barbary, and in cultivation since 1818. The flowers are of a pale rose colour, large in proportion to the leaves, and the petals have yellow spots at the base. The flowers appear from May to August. This is one of the most beautiful species of the section to which it belongs: it requires a little protection during winter, but grows and flowers freely during summer, and also ripens seeds. Plants of it were in the Epsom Nursery in 1828.

2. 76. *H. APENNINUM Dec.* The Apennine Helianthemum, or Sun Rose.

Identification. Dec. Fl. Fr., 4. p. 824.; Don's Mill., 1. p. 313.

Synonymes. *Cistus apenninus Lin. Sp.*, 744.?, *Dill. Eth.*, 170.; *Cistus hispidus β Lam. Dict.*, 2. p. 26.

Engraving. Swt. Cist., t. 62.

Varieties. De Candolle records two forms of this species; one with leaves broad and flattish, and other with leaves linear and narrow.

Spec. Char., &c. Stem suffruticose, branched. Branches spreading, hoary tomentose. Leaves stalked, oblong linear, with the margins scarcely revolute; under surface tomentose, upper surface glaucescent, but at length becoming smooth. Stipules awl-shaped, longer than the footstalks of the leaves. Calyxes covered with very short hairs, striated, cinerously glaucous, bluntish. (*Don's Mill.*, i. p. 313.) A native of Spain, France, Italy, and Germany, on dry hills and places exposed to the sun; and in cultivation in British gardens since 1731. The flowers are white, and the petals are distinct, and marked with yellow at their base. The plant is as hardy as the common species. It was in the Chelsea Botanic Garden in 1828.

2. 77. *H. HISPIDUM Dunal.* The hispid-herbaged Helianthemum, or Sun Rose.

Identification. Dunal ined., and Dec. Prod., 1. p. 282.; Don's Mill., 1. p. 313.

Synonymes. *Cistus hispidus Lam.*, 2. p. 26.; *Brot. Fl. Lus.*, 2. p. 271.; *Cistus pilosus β Gouan. Fl. Monsp.*, p. 265.; *H. majoranæfolium β Dec. Fl. Fr. Suppl.*, p. 625.

Spec. Char., &c. Stem suffruticose, branched. Branches ascending, hoary-tomentose. Leaves stalked, oblong, bluntish, somewhat mucronate, with revolute margins; under surface hoary, upper surface roughish, greenish-glaucous. Calyxes covered with long hairs. (*Don's Mill.*, i. p. 313.) Native of the south of France, and in cultivation since 1816; producing its white flowers from May to July.

2 78. *H. PULVERULENTUM* Dec. The powdered-leaved Helianthemum, or *Sun Rose*.

Identification. Dec. Fl. Fr., 4. p. 823; Don's Mill., 1. p. 313.

Synonymes. *Cistus pulverulentus* Pour. Act. Toul., 3. p. 311; *Cistus polifolius* Lam. Dict., 2. p. 26, but not of Lin.

Engraving. Swt. Cist., 1. 29.

Spec. Char., &c. Stem suffruticose, much branched, prostrate. Branches hoary-tomentose. Leaves oblong linear, with revolute margins, obtuse; under surface hoary, upper surface glaucous. Stipules subulate, ciliated, longer than the footstalks of the leaves. Calyxes hoary, minutely tomentose, pubescent. (*Don's Mill.*, i. p. 313.) A native of France, on sterile hills; and, according to Sweet, in cultivation in the Chelsea Botanic Garden in 1826. It has white flowers, about an inch broad, which appear in May and June; and, though they are not so showy as some of the other species, yet, as Sweet observes, they make a pleasing variety. It is nearly related to *H. apenninum*.

2 79. *H. MACRANTHUM* Swt. The large-flowered Helianthemum, or *Sun Rose*.

Identification. Swt. Cist., t. 103; Don's Mill., 1. p. 313.

Engraving. Swt. Cist., t. 103.

Spec. Char. &c. Stem suffruticose. Branches procumbent, rather tomentose. Leaves flat, ovate oblong, acutish; smooth above, and densely tomentose beneath, pale cinereous. Stipules rather pilose, about equal to, or longer than, the petioles. Calyx striated, pilose. Petals distinct. (*Don's Mill.*, i. p. 313.) The native country of this species is uncertain; but, according to Sweet, it was in cultivation in the Hammersmith Nursery in 1828. The leaves are large and flat, and the flowers larger than any other in the section. The petals are of a cream-coloured white, distinct, spreading, and very slender at the base, where they are marked with yellow. The plant is as hardy as the common species. It was cultivated in the Hammersmith Nursery in 1828.

Variety.

2 *H. m. 2 multiplex* Swt. Cist., t. 104, and our fig. 74., is a beautiful plant, not only on account of its fine double flowers, but of its habit of growth. It ought to be in every Cistacatum. Plants of it were in the Hammersmith Nursery in 1828.



2 80. *H. RHODANTHUM* Dunal. The red-flowered Helianthemum, or *Sun Rose*.

Identification. Dunal, med., and Dec. Prod., 1. p. 282; Don's Mill., p. 313.

Synonymes. *Cistus roseus* Jacq. Hort. Vin., 3. p. 65? *Cistus angustifolius*, formerly in the Royal Botanic Garden at Paris; *Cistus piluliferus* Thib. incd.

Engraving. Swt. Cist., t. 7.

Varieties. De Candolle records three forms of this species: *H. r. oblongifolium*, *H. r. subhirsutum*, and *H. r. carneum*.

Spec. Char., &c. Stem suffruticose, procumbent. Branches rather tomentose, and hoary. Leaves oblong, with revolute margins; under surface hoary-tomentose, upper surface greenish glaucous. Stipules awl-shaped, pilose, and bristly at the tip. Calyxes covered with short white tomentum. (*Don's Mill.*, i. p. 313.) A native of Spain, introduced in 1800, and, according to Sweet, in the Chelsea Botanic Garden in 1825. It is a very showy kind, with flowers of a bright red, inclining to crimson, which it produces in abundance, being quite hardy and of the easiest culture. Sweet says, "Our drawing was taken from a fine plant, growing luxuriantly, with many other handsome species, on the rockwork of the Chelsea Botanic Garden, in June, 1825. Nothing could make a more brilliant appearance than the different habits of growth and beautiful flowers of various colours with which the plants on this rockwork were decked every day for about two months. *H. rhodanthum*," he adds, "also makes a handsome appearance when grown in pots."

We may observe, here, that the rockwork in the Chelsea Botanic Garden forms a sort of truncated cone, flattened on the sides, terminating in a small basin for water plants; and it suggests the idea that a helianthemum wall, constructed somewhat on the same plan as the strawberry wall of Mr. Byers (*Gard. Mag.*, vol. v. p. 438.), would be a very suitable manner of growing the species.

♣ 81. *H. CANESCENS* Swt. The canescent-leaved Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 51.; Don's Mill., 1. p. 313.
Engraving. Swt. Cist., t. 51.

Spec. Char., &c. Stem suffruticose, branched, diffuse. Branches ascending, rather tomentose, canescent. Leaves flat, or hardly revolute at the margins; under surface tomentose hoary, upper surface greenish glaucous. Lower leaves ovate oblong, obtuse; upper ones lanceolate, acute. Stipules linear, ciliated, somewhat longer than the footstalks. Calyxes smoothish, but with the nerves pubescent. Petals imbricated. (*Don's Mill.*, i. p. 313.) A splendid plant, with reddish crimson flowers; the petals imbricated, and having a small orange spot at the base of each. Its native country is uncertain, but it was in cultivation in the Fulham Nursery in 1826. Sweet considers it as having the darkest-coloured, if not the handsomest, flowers of the genus. The flowers are also, he says, very large for the size of the plant. It is nearly related to *H. rhodanthum*, but is readily distinguished from it by its canescent leaves, and stronger habit of growth.

♣ 82. *H. CONFUSUM* Swt. The confused Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 91.; Don's Mill., 1. p. 313.
Synonymc. *H. polifolium* Dec. *Prod.*, 1. p. 283.
Engraving. Swt. Cist., t. 91.

Spec. Char., &c. Stem suffruticose. Branches procumbent, smoothish, rather tomentose at the apex. Leaves oblong, ovate, bluntish, rather flat; under surface tomentose, hoary; upper surface glabrous, green. Stipules and bracteas linear, green, ciliated. Calyxes striated, smoothish, rather shining. Petals imbricate. (*Don's Mill.*, i. p. 313.) A native of France and Spain, and, according to Sweet, cultivated in British nurseries in 1829. Its flowers are white, and of a delicate texture. This sort, Sweet observes, is generally found, in the nurseries, under the name of *H. polifolium*; but he thinks that it has nothing to do with the English species of that name. For some reason of this kind, we suppose, it has been called the *H. confusum*.

♣ 83. *H. LANCEOLATUM* Swt. The lanceolate-leaved Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 100.; Don's Mill., 1. p. 313.
Engraving. Swt. Cist., t. 100.

Spec. Char., &c. Stem suffruticose, much branched, procumbent. Branches ascending, smoothish, hoary-tomentose at the apex. Leaves lanceolate, acute, with somewhat revolute margins; green and smoothish above, but hoary-tomentose beneath. Stipules awl-shaped, linear, longer than the footstalks of the leaves. Sepals smoothish, or rather pilose. Petals imbricate. (*Don's Mill.*, i. p. 313.) A hybrid, found in gardens in 1818, and producing white flowers, marked with yellow, from May to August. "Confused with *H. polifolium* by some botanists," but readily distinguished by its sharp-pointed leaves, which are of a glossy green on the upper side, and by its broad imbricated petals. Plants of it were in the Chelsea Botanic Garden in 1829.

♣ 84. *H. POLIFOLIUM* Pers. The Polium-leaved Helianthemum, or Sun Rose.

Identification. Pers. *Ench.*, 2. p. 80.; Don's Mill., 1. p. 313.
Synonymc. *Cistus polifolius* Lin. *Sp.*, 745., *Smith's Engl. Bot.*, 1322., *Dill. Elth.*, 175. t. 145. f. 172.
Engravings. *Smith's Engl. Bot.*, t. 1322.; *Dill. Elth.*, 175. t. 145. f. 172.

Spec. Char., &c. Stem suffruticose, branched. Branches procumbent, densely tomentose. Leaves oblong-linear, with revolute margins, hoary-tomentose on both surfaces. Stipules narrow, linear, obtuse, longer than the petioles, and are, as well as the bractæas, tomentose and ciliated. Petals distinct, crenulated. (*Don's Mill.*, i. p. 313.) A native of England, producing its white flowers, marked with yellow, from May to August. It is found on stony hills near the sea side, particularly on Brent Downs, Somersetshire; also at Babicome, near Newton Abbot, and on Tor Hill, near Torquay, Devonshire.

85. *H. MUTABILE Pers.* The changeable-coloured-flowered Helianthemum, or *Sun Rose*.

Identification. Swt. Cist., 106; Pers. Ench., 2. p. 79.; Dec. Prod. I. p. 283.; Don's Mill., i. p. 318.
Synonymæ. *Cistus mutabilis Jacq. Icon. rar.*, i. t. 99.; *Misc.*, 2. p. 340.

Engravings. Swt. Cist., t. 106.; Jacq. Icon. rar., i. t. 99.

Varieties. There are three forms of this very handsome species; one with white flowers, another with smaller flowers of a rose red, and the third with double rose-coloured flowers.

Spec. Char., &c. Stem suffruticose. Branches procumbent, rather tomentose. Leaves flat, ovate-oblong, acutish; upper surface glabrous, under surface tomentose, pale-cinereous. Stipules rather pilose, generally equal in length with the footstalks of the leaves, or longer. Calyxes striated, smoothish. Petals imbricate. (*Don's Mill.*, i. p. 313.) A native of Spain, and in cultivation in British gardens in 1829. The flowers are produced from June to August, in great abundance; and seeds are ripened afterwards in fine seasons. The plants are quite hardy. They were cultivated for sale in the Clapton Nursery in 1829.

86. *H. VARIEGATUM Swt.* The variegated Helianthemum, or *Sun Rose*.

Identification. Swt. Cist., t. 33.; Don's Mill., i. p. 314.

Engraving. Swt. Cist., t. 38.

Spec. Char., &c. Stem suffruticose, procumbent. Branches tomentose, rather hoary, diffusely procumbent. Leaves lanceolate, acute, flattish; under surface hoary-tomentose, upper surface green, rather scabrous. Stipules linear, ciliated, longer than the petioles. Calyxes covered with short violaceous tomentum. Petals imbricate, undulated. (*Don's Mill.*, i. p. 314.) A hybrid between *H. rhodanthum* and *H. lineare*, found by Mr. Sweet in the Chelsea Botanic Garden in 1827. "When in full bloom, it makes a very pleasing appearance, from the diversity of colours in its flowers; some being nearly all red, others variegated with dark and light red and white, and some altogether white: it also continues to bloom, if the weather prove favourable, from May to November." It is as hardy as the indigenous species.

87. *H. VERSICOLOR Swt.* The various-coloured Helianthemum, or *Sun Rose*.

Identification. Swt. Cist., t. 26.; Barrel. Icon., 440.; Don's Mill., i. p. 314.

Engravings. Barrel. Icon., 440.; Swt. Cist., t. 26.

Spec. Char., &c. Stem shrubby, erect. Branches ascending, rather hoary from stellate down. Leaves oblong, flat, or concave above; under surface hoary-tomentose, upper surface green, glabrous. Stipules oblong-linear, ciliated, bristly at the top, somewhat longer than the leaves. Calyxes covered with short tomentum. Petals imbricate. (*Don's Mill.*, i. p. 314.) A native of the south of Europe, and readily distinguished from *H. variegatum* by its upright habit of growth, and its lanceolate and acutely pointed leaves. The flowers are very variable in colour, scarcely two on the plant being alike; some are of a bright red, others nearly yellow, some copper-coloured, and others with a mixture of all these colours, and of the different shades between them; so that the plant, when in flower, has a curious variegated appearance. (*Sweet.*)

2. 88. *H. SULPHU`REUM Willd.* The sulphur-colour-flowered Helianthemum, or Sun Rose.

Identification. Willd. Enum. Suppl., 39; Swt. Cist., t. 37; Don's Mill, i. p. 314.
Engraving. Swt. Cist., t. 37.

Spec. Char., &c. Stems branched, procumbent. Leaves lanceolate, flat; upper surface green, under surface paler, but beset with stellate pubescence on both surfaces. Racemes terminal, few-flowered. (*Don's Mill*, i. p. 314.) A native of Spain, and cultivated in British gardens in 1795. A very distinct variety, when in flower, during June and July. It is tolerably hardy, but requires a little protection during very severe frosts. Plants were in the Fulham Nursery in 1826.

2. 89. *H. STRAMI`NEUM Swt.* The straw-colour-flowered Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 93; Don's Mill, i. p. 314.
Engraving. Swt. Cist., t. 93.

Spec. Char., &c. Stems branched, elongated, procumbent, tomentosely pubescent at the apex. Leaves flat, or with the margins scarcely revolute; green above and pilose, hoary-tomentose beneath; lower ones roundish ovate, obtuse; upper ones oblong-lanceolate, acutish. Stipules lanceolate, acute, ciliated, twice the length of the petioles. Racemes many-flowered. Calyx striated, smoothish. Petals obovate, spreading, distinct. (*Don's Mill*, i. p. 314.) Found in gardens, and, probably, a hybrid. Its straw-coloured flowers are produced from May to August. Plants were in the Clapton Nursery in 1829.

Variety.

2. *H. s. 2 multiplex* (*Swt. Cist.*, t. 94.) has double straw-coloured flowers, with the petals orange-coloured at the base. It is a very beautiful variety, tolerably hardy. Plants of it were in the Hammer-smith Nursery in 1829.

2. 90. *H. DIVERSIFO`LIUM Swt.* The various-leaved Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 95; Don's Mill, i. p. 314.
Engraving. Swt. Cist., t. 95.

Variety.

2. *H. d. 2 multiplex.*—Flowers large, double, and of deep purplish red, mixed with lighter-coloured ones.

Spec. Char., &c. Stem suffruticose, ascending, branched. Branches rather tomentose, erectly ascending. Leaves stalked, green, hairy above, hoary-tomentose beneath; lower ones oval or oblong, obtuse, flat; upper ones linear-lanceolate, ciliated, three to four times longer than the petiole. Sepals pilose. Petals crenulated, distinct. (*Don's Mill*, i. p. 314.) It is not known of what country it is a native; but it is, or has been, cultivated in England, as Sweet's drawing was taken from a plant in the Hammersmith Nursery in 1829. It is nearly allied to *H. polifolium*, and has a very brilliant appearance from May to June, from the dark rich red of its flowers, each of the petals of which has a deep copper-coloured mark at its base. The plant is a very desirable one for rockwork. When grown in pots, it requires a light sandy soil.

2. 91. *H. ERIOSE`PALON Swt.* The woolly-sepaled Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 76; Don's Mill, i. p. 314.
Engraving. Swt. Cist., t. 76.

Spec. Char., &c. Stems branched, procumbent, rather tomentose, hoary at the apex. Leaves lanceolate, acute, with somewhat revolute margins, green on both surfaces, and beset with starry hairs. Stipules linear, acute, ciliate, twice as long as the footstalks of the leaves. Racemes terminal, many-flowered. Calyxes clothed with woolly hairs. Petals obovate, crenulated,

distinct at the base. (*Don's Mill.*, i. p. 314.) A hybrid, found in Colvill's Nursery, Chelsea, in 1828. It is nearly related to *H. sulphureum*, but differs from it in habit, and in having a woolly calyx. The leaves are narrow, and undulated in the margins.

2 92. *H. ROSEUM* Dec. The Rose-colour-flowered Helianthemum, or Sun Rose.

Identification. Dec. Fl. Fr., 4. p. 822; *Don's Mill.*, i. p. 314.
Synonymic. *Cistus roseus* *All. Ped.*, 2. p. 105. t. 45. f. 4., but not of Jacq.
Engraving. Swt. *Cist.*, t. 55.

Variety.
 2 *H. r. 2 multiplex.* (*Swt. Cist.*, 86.)—A very pretty double variety: it is well suited to rock-work, on which it grows vigorously, and produces large flowers. It is quite hardy, and continues in bloom the whole summer.

Spec. Char., &c. Stem suffruticose, rather procumbent, somewhat tomentose. Leaves ovate-lanceolate, tomentose on the under surface, green above, hairy. Stipules lanceolate-linear, ciliated. Pedicels and calyxes pilosely hairy. (*Don's Mill.*, i. p. 314.) A native of the south of Europe, and very nearly allied to *H. vulgare*. The petals are rose-coloured, and imbricate at the base. It flowers in June and July, and was introduced in 1815. It is a very pretty plant, from the delicate colour of its flowers; and it is very suitable for rockwork, as it will continue in bloom for several months. It requires protection during frosts. It was cultivated in the Fulham Nursery in 1827.

2 93. *H. FÆTIDUM* Pers. The fetid Helianthemum, or Sun Rose.

Identification. Pers. *Syn.*, 2. p. 79.; *Don's Mill.*, i. p. 314.
Synonymic. *Cistus fœtidus* *Jacq. Icon. rar.*, 1. p. 98., *Misc.*, 2. p. 341.

Spec. Char., &c. Stem suffruticose, procumbent, pilosely hairy. Leaves oblong, green on both surfaces, hairy, roughish. Stipules hairy, linear, longer than the footstalks of the leaves. Pedicels and calyx rather hairy. (*Don's Mill.*, i. p. 314.) A hybrid, resembling *H. vulgare*, but differing from it in having white flowers. The whole plant is said to smell like *Bryonia*. It produces its flowers from May to July, and was in cultivation in 1800.

2 94. *H. HYSSOPIFOLIUM* Tenore. The Hyssop-leaved Helianthemum, or Sun Rose.

Identification. Tenor. *Syn. Fl. Neap.*, p. 48.; *Don's Mill.*, i. p. 314.

Spec. Char. Stem suffruticose, ascending. Branches hairy-tomentose. Lower leaves oval, upper ones oblong-lanceolate, green on both surfaces, flat, hairy. Calyxes hairy. Petals imbricate. (*Don's Mill.*, i. p. 314.) Varying considerably by culture.

Varieties.

2 *H. h. 1 crocætum* (*Swt. Cist.*, t. 92.) has flowers saffron-coloured, with more or less, of a ferruginous tint, and may represent the species. Plants of it were in the Clapton Nursery in 1828.

2 *H. h. 2 cupreum* (*Swt. Cist.*, t. 58.) has flowers of a reddish copper-colour. Plants of it were in the Fulham Nursery in 1827.

2 *H. h. 3 multiplex* (*Swt. Cist.*, t. 72., and our fig. 75.) has double flowers, of a reddish copper-colour. Plants of this variety were in the Fulham Nursery in 1828.

Description, &c. All the three forms of this species are splendid plants; they are hardy, of luxuriant growth, flowering freely, and of the easiest culture, either in pots or on banks of light sandy soil, covered with flints or stones. The flowers of the copper-coloured variety, and also the leaves, are larger than those of the two other kinds. The double-flowered variety appears to be of a more upright habit of growth, and not quite so robust as the others. Sweet says that he is "acquainted with two other very distinct varieties; one with flowers of a lighter colour, and the other having double flowers."



2. 95. *H. CU'PREUM* Swt. The copper-colour-flowered Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 66.; Don's Mill., 1. p. 314.

Engraving. Swt. Cist., t. 66.

Spec. Char., &c. Stem suffruticose, procumbent. Branches ascending, rather tomentose; adult ones glabrous. Leaves oblong-lanceolate, channeled; upper surface green, hairy; under surface hoary-tomentose. Stipules lanceolate, acute, ciliated, bristly at the apex, twice as long as the footstalks of the leaves. Calyxes tomentosely pilose. Petals imbricated. (*Don's Mill.*, i. p. 314.) A hybrid, found by Mr. Sweet in the Hammersmith Nursery in 1827. The petals are of a dark copper colour, with a darker mark at the base of each. The flowers appear from May to August, and the plant requires very little protection.

2. 96. *H. VENU'STUM* Swt. The handsome Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 10.; Don's Mill., 1. p. 314.

Engraving. Swt. Cist., t. 10.

Spec. Char., &c. Stem suffruticose, ascending, branched. Branches glabrous, warted, somewhat tomentose at the apex. Leaves oblong-lanceolate, acute, flat, or hardly revolute in the margins, but denticulately scabrous; under surface hoary-tomentose, upper surface green, shining. Stipules lanceolate, hairy, ciliated, twice as long as the footstalks of the leaves. Inner sepals membranous, with hairy warted nerves. Petals imbricated. (*Don's Mill.*, i. p. 314.) The native country of this beautiful plant is not known, though it is now very common in collections about London. It is very suitable for rockwork, as it is quite hardy, and continues in flower during the whole summer. In some collections, Sweet tells us, it is considered only as a variety of *H. vulgare*; but he says that it is more nearly allied to *H. rhodanthum*; from which, however, it may be easily distinguished "by its warted stalks and calyxes, and its smooth shining stems." It was in Colvill's Nursery in 1825.

2. 97. *H. MI'LLERI* Swt. Miller's Helianthemum, or Sun Rose.

Identification. Swt. Cist., t. 101.; Don's Mill., 1. p. 315.

Engraving. Swt. Cist., t. 101.

Spec. Char. Stem suffruticose, procumbent. Branches hairy-tomentose. Leaves oblong, bluntish, flat, green on both surfaces, hairy. Stipules falcate, longer than the petioles. Calyxes hairy. Petals imbricate. (*Don's Mill.*, i. p. 315.) A hybrid, with saffron-coloured flowers, with a dark mark at the base of each petal. It is nearly related to *H. hyssopifolium*, from which it differs in its dull green and rough leaves, which are much more hairy. It is also related to *H. nummularium*, and may, probably, be a hybrid between the two. It is one of the most ornamental sorts of the section to which it belongs, and is so hardy as to require no protection in winter. Plants of it were in the Bristol Nursery in 1829.

2. 98. *H. MAJORANÆFO'LIUM* Dec. The Marjoram-leaved Helianthemum, or Sun Rose.

Identification. Don's Mill., 1. p. 315.

Synonymes. *H. m.* var. α *Dec. Fl. Fr.*, 6. p. 225.; *Cistus majoranæfolius Gouan. Herb.*, p. 26. ?

Spec. Char. Suffruticose, erect, much branched. Branches hairy-tomentose. Leaves stalked, ovate-oblong, acutish, with revolute margins; under surface hoary-tomentose, upper surface greenish glaucous, tomentosely hairy. Stipules awl-shaped, bristly. Calyxes densely clothed with white hairs. (*Don's Mill.*, i. p. 315.) A native of the south of Europe, and introduced in 1818. It produces its yellowish-white flowers in May and June.

2. 99. *H. HIRSU'TUM* Dec. The hairy Helianthemum, or Sun Rose.

Identification. Dec. Prod. 1. p. 284.; Don's Mill., 1. p. 315.

Synonymes. *Cistus hirsutus Lapcyr. Abr.*, 303., but not of Lam.

Spec. Char. Suffruticose, stipulate, hairy. Leaves stalked; under surface hoary. Lower leaves rounded, upper ones lanceolate, acute. Flowers secund, in terminal racemes. (*Don's Mill.*, i. p. 315.) A native of the eastern Pyrenees, on rocks. It has large white flowers, and is procumbent; but it has not yet been introduced into British gardens.

GENUS III.

HUDSONIA L. THE HUDSONIA. *Lin. Syst. Polyándria Monogýnia.*

Identification. Lin. Mant., 11.; Nutt. Gen. Amer., 2. p. 4.; Dec. Prod., 1. 284.

Derivation. Named in honour of William Hudson, a London apothecary, the author of *Flora Anglica*, published in 1762.

Gen. Char. Sepals 5, equal. Petals 5. Stamens 15—30. Filaments filiform. Anthers small, opening lengthwise. Style straight, simple, equalling the stamens in length. Stigma simple. Capsule 1-celled, 3-valved, 1—3-seeded, oblong or obovate, coriaceous, smooth or pubescent. Seeds granulated. Embryo immersed in a horny albumen. (*Don's Mill.*, i. p. 315.) Small heath-like shrubs, natives of North America, with yellow flowers, almost sessile, solitary, or aggregate.

1. II. ERICOIDES L. The Heath-like Hudsonia.

Identification. Lin. Mant. 74.

Engravings. Willd. Hort. Ber. t. 15.; Swt. Cist., t. 36.; Don's Mill., 1. p. 315.; and our fig. 76.

Spec. Char., &c. Pubescent. Stems suffruticose, erect. Branches elongated. Leaves filiform, awl-shaped, rather imbricated. Peduncles solitary, rising laterally from the leafy bud. Calyx cylindrical, obtuse. Capsule pubescent, always 1-seeded. Valves oblong. (*Don's Mill.*, i. p. 315.) A heath-like shrub, native of New Jersey and Virginia, in pine woods, and introduced into England in 1805. It is a short, densely branched, suffruticose plant, rather scarce in British collections. Its flowers are yellow, small, solitary, and produced from May to July. The plant is rather more difficult to cultivate than those of the other genera of this order; but it thrives very well in sandy peat; its native habitat being similar to that of the common heath in England.



2. II. [? E.] NUTTALLI Swt. Nuttall's Hudsonia.

Identification. Swt. Cist., p. 19.; Don's Mill., 1. p. 315.

Synonymes. II. ericoides Nutt. Gen. Amer., 2. p. 4.

Spec. Char., &c. Equally pubescent. Stem erect, much branched. Leaves about 2 lines long, filiform, rather imbricate, but distinct from the stem. Pedicels lateral, crowded; when in fruit, from 5 to 8 lines long. Calyx cylindrical, obtuse, pubescent, with the segments oblique and convolute; the two smaller ones hardly visible when in fruit, but sufficiently distinct in the unexpanded flowers. Capsules cylindrical, oblong, externally pubescent, always 1-seeded. Valves oblong; the central suture obsolete. (*Don's Mill.*, i. p. 315.) An evergreen undershrub, closely resembling II. ericoides, and perhaps only a variety of it.

3. II. [? E.] TOMENTOSA Nutt. The tomentose-leaved Hudsonia.

Identification. Nutt. Gen. Amer., 2. p. 5.; Swt. Cist., t. 57.; Don's Mill. 1. p. 316.

Engraving. Swt. Cist., t. 57.

Spec. Char., &c. Tufed, and hoary-tomentose. Stems intricate, dense. Leaves minute, densely imbricated, ovate, acute. Flowers aggregate, almost sessile. Calyxes rather cylindrical, with obtuse partitions. Capsules 1-seeded. Valves ovate, smooth. (*Don's Mill.*, i. p. 316.) Found in New Jersey, Delaware, Maryland, &c., in the sea sand. It was first discovered by Mr. Nuttall, and described by him as a very distinct species. Mr. James McNab, in "An Account of some of the rarer Plants, observed during an Excursion in the United States and the Canadas in 1834," published in the *Edinburgh New Philosophical Journal*, No. 37., July, 1835, says

that he found this species in dry sandy barrens in New Jersey, and that he is much inclined to think it is only a variety of *H. ericoides*: they were both seen in abundance together; and many subvarieties were observed, which seemed to unite the two. This curious shrub covers large tracts of the dry white sandy plains of New Jersey, in large round tufts, where it resembles very much the common ling (*Calluna vulgaris*) of Scotland." Were all botanists, who, like Mr. McNab, are at once scientific observers and practical cultivators, to exercise their common sense, as he has done in this instance, we are persuaded that the number of alleged species, in every genus which now contains a great many, would soon be considerably reduced. The flowers of this kind of hudsonia are small and yellow, and appear from May to July. Like *H. Nuttallii*, this is somewhat difficult of culture, requiring a peat soil, a shady situation, and protection by glass or by snow during winter. All the species are readily increased by layers, or by cuttings of the ripened wood, planted in sand under a hand-glass.

App. I. *Other Species of Cistàcæa.*

Long as is the list of *Cistàcæa*, and especially of helianthemums, in this chapter, it might have been increased by the addition of various other species, or sorts, described by botanists. The utility, however, of such additions is very questionable. We have confined ourselves, as much as possible, to sorts that are, or were lately, in existence in the neighbourhood of London; and for this information we have taken as our guide the *Cistineæ* of Sweet, commenced in 1825, and completed, in one volume, in January, 1830. We think we may safely assert that several of the sorts described in that work are now no longer in existence; because, in consequence of their tenderness, and liability to be neglected, they are continually dying off during winter, and as continually being replaced by others raised from seeds, either imported or saved in this country. The chief use which we propose to ourselves, in giving so long a descriptive list, is, to show the numerous and beautiful forms assumed by this family of plants, in order to promote their more extensive cultivation. Supposing a cultivator about to form a collection of *Cistàcæa*, we should attach much less importance to his being able to procure all the sorts described by Mr. Sweet, than to his obtaining all the sorts easily procurable, whatever names they might pass under, and cross-fecundating them, so as to produce new forms. There can be no doubt whatever that the sorts of both the genera *Cistus* and *Helianthemum* might, by cross-fecundation, be increased *ad infinitum*; and, considering their very great beauty as border and rockwork shrubs, we think they merit the attention of cultivators at least as much as many florist's flowers.

CHAP. XII.



OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE ORDER POLYGALA'CEÆ.

THERE is only one perfectly hardy ligneous plant belonging to this order, and it is of such humble growth, that, for all practical purposes, it may be considered as a herbaceous plant, rather than as a shrub. We have introduced the order, however, chiefly for the purpose of recommending gardeners to try some of the beautiful Cape species of *Polygala* against conservative walls.

■ 1. *POLY'GALA CHAMÆBU'XUS* L. The Dwarf Box Polygala, or Box-leaved Milkwort. (Jacq. Aust., t. 233.; Sims, Bot. Mag., t. 316.; and our fig. 77.)

Described as having fruticose, branched, procumbent stems, with oblong-lanceolate mucronated leaves; and the racemes 1—2-flowered; and the keel of the flower crested. It forms a little evergreen tuft, the leaves being like those of the dwarf box; and the yellowish flowers, which are slightly tipped with purple, resembling at a distance those of the order Leguminosæ. It is a native of mountainous woods in many parts of Europe, particularly in those of Germany and Switzerland. In rocky situations, it seldom exceeds 6 in. in height; but in heath soil, or in sandy loam enriched with leaf mould, it will grow to the height of a foot and upwards, flowering freely every year. This plant has been in cultivation in British gardens since 1658. Miller says that the seeds, which are with difficulty obtained from abroad, do not vegetate till they have been a whole year in the ground; unless they are sown soon after they are ripe, which is in August or September, in which case they will come up the following spring. It is readily propagated, however, by division of the plant, as it throws up suckers in abundance. This plant succeeds very well in most gardens, in a shady situation, and in peat soil kept rather moist. Intermixed with *Gaulthéria procumbens*, *Mitchella repens*, *Linnaea borealis*, and other dwarf-growing shrubs, *Polygala Chamæbuxus* will form a rich margin to American groups. The price, in the London nurseries, is 50s. a hundred; it being frequently sold in quantities for forming edgings to beds of peat-earth plants.



■ 2. The half hardy *Polygalas* are evergreen shrubs, natives of the Cape of Good Hope; and in Britain they are generally treated as greenhouse plants, though some of them have stood against a wall, with protection during winter: and so great is their beauty during summer, that, we think, whoever has a conservative wall ought to place some of them against it. Above twenty Cape species have been introduced; but the most common are, *P. oppositifolia* L. (*Bot. Reg.*, t. 636.), which is a native of the mountainous part of the Cape, and tolerably hardy; *P. oppositifolia major* (fig. 78.), called by some *P. grandiflora*, which is equally hardy, and is a fine variety; *P. latifolia* Ker, *P. myrtifolia*, *P. grandiflora* Lodd., *P. bracteolata* L., *P. speciosa* Bot. Mag., and *P. attenuata* Lodd., all fine plants, with bright purple flowers mixed with white, and some of them with red, and all procurable in the principal London nurseries. Like almost all other Cape shrubs, they grow best in heath soil, or in a mixture of sand and leaf mould; and, when they are cultivated against a wall in the open ground, great care should be taken not to let their stems be injured by damp in autumn, more especially at the surface of the ground; or, in technical language, at the collar. There are some trees and shrubs belonging to the order Polygalææ in the Himalaya; which, when introduced, will be worth trying against a conservative wall with the Cape species.



CHAP. XIII.

OF THE HARDY AND HALF-HARDY LIGNEOUS SPECIES OF THE ORDER PITTOSPORACEÆ.

Distinctive Characteristics. Thalamiflorous. (*H. B.*) Sepals 5, petals 5; both imbricate in aestivation. Stamens 5, distinct, alternate with the petals. Ovarium of several cells, with the placenta in the axis; cells or placenta 2 or 5 in number, and many-ovuled. Style 1. Stigmas as many as the placenta. Fruit capsular, or berryed. Seeds often covered with a glutinous or resinous pulp. (*Lindl. In. to N. S.*) The species contained in this order are all ligneous; and are either trees, or bushy or climbing shrubs, with terminal or axillary flowers, usually of a bell-shape, with a spreading border. They are natives of warm climates; but some species of *Pittosporum*, *Billardiæra*, and *Söllya*, are half-hardy, and suitable for a conservative wall.

GENUS I.



BILLARDIÆRA Sm. THE BILLARDIERA, or APPLE-BERRY. *Linn. Syst.* Pentândria Monogýnia.

Derivation. Named in honour of *Jean Jacques Julien La Billardière*, a celebrated French botanist, who visited Syria, and afterwards New Holland in D'Entrecasteaux's expedition. He was the author of *Nove Hollandicæ Plantarum Specimen*, and other works.

Gen. Char. Corolla tubularly bell-shaped. Anthers widely distant, opening lengthwise. Ovary 2-celled, many-ovuled. Pericarp soft, spongy, sub-baccate, the cells inflated. Seeds many, lying loose in the cells, not attended by pulp. Twining Australian shrubs, with entire or serrated leaves, and axillary flowers, sub-solitary, pendulous, and of a colour passing from green to yellow. (*Lindl.*, in *Bot. Reg.*, t. 1719.) The fruit, in most of the species, is of a bluish colour when ripe, and is eatable. (*Dec. Prod.*, and *Don's Mill.*)



79

1. *B. LONGIFLO'RA* Labill. The long-flowered Billardiera, or Apple-berry.

Identification. Labill. Nov. Holl., t. 89.; Don's Mill., l. p. 373.

Engravings. Labill. Nov. Holl., t. 89.; Bot. Mag., t. 1507.; and our fig. 79.

Spec. Char. Branches climbing, younger ones scarcely pubescent. Leaves lanceolate, entire. Pedicels 1-flowered, glabrous, one half shorter than the flower. Berries almost globose, torose, glabrous. (*Don's Mill.*, i. p. 373.)

An evergreen twiner, introduced from Van Diemen's Land in 1810; growing vigorously, and flowering and fruiting freely in conservatories; whence we are led to conclude that it will answer against a conservative wall. In a conservatory it is an interesting twiner, from its slender habit, abundant small leaves, and rather numerous dark blue fruits.

2. *B. OVA'LIS* Lindl. The oval-leaved Billardiera, or Apple-berry.

Identification. Lindl. in *Bot. Reg.*, t. 1719.

Engraving. Bot. Reg., t. 1719.

Spec. Char., &c. Of this species Dr. Lindley states that "it is nearly related to *B. longiflora*, from which it chiefly differs in its smaller and shorter flowers, and more oval obtuse leaves. Its flowers change from greenish yellow to dark purple, and appear in May. A native of Van Diemen's Land, whence it was introduced by Mr. Low of Clapton." (*Bot. Reg.*, t. 1719. Nov. 1834.) Dr. Lindley adds, "It is probable that *Billardiera ovalis* will be quite hardy enough to live in this country, trained to a west wall, if protected from wet in winter; at all events, a cold-pit would be an ample covering for it, and for all the other species. The beautiful *Sollya* grows with all its native luxuriance in such a situation." (*Ibid.*)



80

3. *B. MUTA'BILIS* H. K. The changeable-coloured-flowered Billardiera, or Apple-berry. (fig. 80.)

A native of New South Wales, and producing purplish flowers from June to September. It is not such a strong-growing plant as *B. longiflora*, and is probably more tender than that species; but, on account of the beauty of the flowers, and the large size of the fruit, it ought to be tried.

GENUS II.



SO'LLYA Lindl. THE SOLLYA. *Lin. Syst.* Pentándria Monogýnia.

Synonymy. Billardiera, in one species.

Derivation. Named in honour of Richard Horsman Solly, Esq., F.R.S., &c.; "whose general acquaintance with science, and, as far as botany is concerned, with vegetable physiology and anatomy, are such as to entitle him most fully to such a mark of respect." (*Lindl.*, in *Bot. Reg.*, t. 1466.)

Gen. Char. Calyx minute. Corolla spreadingly bell-shaped, petals rather unequal. Stamens opposite the petals. Anthers disposed into a cone, connate at the tip, and each opening by two pores at its tip. Ovary 2-celled, many-ovuled. Pericarp spindle-shaped, many-seeded. — Australian shrubs, twining in some degree. Leaves simple, alternate, remaining long on the plant, devoid of stipules. Flowers in cymes that are placed opposite the leaves, blue. (*Lindl.*, in *Bot. Reg.*, t. 1466. Jan. 1832.) Fruit 4-celled, the seeds enveloped by a soft pleasant pulp. (*D. Don*, in *Sw. Fl.-Gard.*, 2. ser. Aug. 1834.) Fruit, as examined in a half-ripe state, 2-celled; each cell occupied by two rows of seeds, set fast in a firm somewhat fleshy substance, which fills each cell, and which we presume to be what finally becomes the pulp that envelops the seeds. (*Lindl.*, in *Bot. Reg.*, t. 1719. Nov. 1834.)

1. *S. HETEROPHY'LLA* Lindl. The various-leaved Sollya.

Identification. Lindl., in *Bot. Reg.*, t. 1466. t. 1719.

Engravings. Bot. Reg., t. 1466.; Swt. Br. Fl.-Gar., 2. s., t. 232.; and our fig. 81.

Spec. Char., &c. Flowers constantly bright blue. Fruit including pulp that envelops the seeds. This is a very interesting plant, from its slender stems and branches, its fine full-green and abundant foliage, and its neat, simple-formed, pendulous flowers, with corollas of a beautiful bright blue. It is found wild on the south-western coast of New Holland, and was introduced in 1830. It has been since treated by some as a green-house plant, and by others as being nearly hardy. It grows in sand and loam, with a mixture of leaf mould, and is readily propagated by cuttings of the young wood in sand under a glass. Plants, in the London nurseries, cost 2s. each.



81

2. *S. ANGUSTIFOLIA* Lindl. The narrow-leaved Sollya.

Identification. Lindl., in *Bot. Reg.*, t. 1466.

Synonymes. *Billardiëra fusiformis* Lab. *Nov. Holl.*, *Don's Mill.*, 1. p. 373., and *Loudon's Hort. Brit.*, No. 5530.

Engraving. Labill. *Nov. Holl.*, t. 90.

Spec. Char., &c. Flowers cream-coloured, changing to bluish. Fruit dry, the pericarp villous, of the consistence of parchment. (*Lindl.*, in *Bot. Reg.*, t. 1466.; *Don's Mill.*, 1. p. 373., under *Billardiëra fusiformis* Lab.) Branches hardly climbing; younger ones rather villous. Leaves lanceolate, entire. Cymes few-flowered. Petals spreading. Native of Van Diemen's Land. A green-house climbing shrub. Flowers in May and August. Cultivated in Britain in 1823. (*Don's Mill.*, i. p. 373., under the name *Billardiëra fusiformis* Lab.) This species deserves trial against a conservative wall, along with the others.

GENUS III.

PITTO'SPORUM Banks. THE PITTOSPORUM. *Lin. Syst.* Pentándria Monogýnia.

Identification. Banks, in *Gært. Fr.* 1. p. 286. t. 59.; *Dec. Prod.*, 1. p. 346.; and *Don's Mill.*, 1. p. 373.

Derivation. From *pitta*, pitch, and *sporos*, a seed; in allusion to the seeds being covered with a sticky pulp.

Gen. Char. *Calyx* of 5 sepals. *Petals* 5, with the claws conniving into a connate tube. *Capsules* smooth or hairy; 2-5-valved, 1-celled, bearing a dissepiment in the middle of each valve. *Seeds* covered with a resinous pulp. (*Don's Mill.*, i. p. 373.)—Evergreen shrubs, or low trees, with entire permanent leaves, generally more or less lanceolate. The species known to be half-hardy are two, but all the rest may be equally so.

1. *P. TOBIRA* Ait. The Tobira Pittosporum.

Identification. Sims *Bot. Mag.*, 1396.; *Dec. Prod.*, 1. p. 346.; and *Don's Mill.*, 1. p. 373.


Synonymes. *Euonymus Tobira* Thunb., chap. 99.; *Pittosporum chinënsis* Donn's *H. Cantab.*, 48.

Tobira Japane, *Fr.*; *Chinesischer Klebsame*, *Ger.*

Engravings. Kæmpf. *Amœn.*, t. 797.; *Bot. Mag.*, t. 1396.; and our *fig.* 82.

Spec. Char., &c. Leaves obovate, obtuse, coriaceous, quite smooth.

Peduncles 1-flowered, pubescent, disposed in aggregate umbels. (*Don's Mill.*, i. p. 373.) An evergreen shrub, growing to the height of 12 ft. in its native country, and producing its cream-coloured flowers from March till August. It was introduced in 1804, and is generally treated as a green-house plant; but, planted in a warm situation against a wall, it endures the winters of the climate of London without any protection when they are mild; and with the ground covered with litter, and the branches screened by a mat, when they are severe. The plant will grow in any free soil, sufficiently drained, and is readily propagated by cuttings of the young wood in sand. The largest specimens of which we have received any accounts are in Ireland; one in the Cullenswood Nursery, near Dublin, 20 years planted, being 10 ft. high, as a standard in the open air. In the neighbourhood of London, there are various plants placed against the ends and fronts of green-houses, which have reached 5 ft. or 6 ft. in height in as many years; which are protected by a mat in very severe winters, and which, when not protected, sometimes die down to the ground, and spring up again the following season. Their glossy dark green leaves, and fragrant cream-coloured flowers, are very ornamental during summer. Price of plants, in the London nurseries, 2s. 6d. each.




2. *P. UNDLA'TUM*. The undulated-leaved Pittosporum.

Identification. *And. Bot. Rep.*; *Vent. Hort. Cels.*; *Ker Bot. Reg.*

Dec. Prod., 1. p. 346.; *Don's Mill.*, p. 373.

Engravings. *Vent. Hort. Cels.*, 1. t. 76.; *Bot. Rep.*, t. 293.; *Delanuy, Herb. Anat.*, t. 36.; *Schrad. Gen. III.*, t. 4.; *Bot. Reg.*, t. 16.; and our *fig.* 83.

Spec. Char., &c. Leaves oval-lanceolate, undulated, tapering at both ends, and, as well as the footstalks, glabrous. *Peduncles* terminal, aggregate, pubescent, branched, many-flowered. (*Don's Mill.*, i. p. 373.) An evergreen shrub, growing to the height of 10 ft., introduced in 1789, and producing its white flowers from May to June. As hardy as the preceding species. The finest plant in England is understood to be in the conservatory at Ashridge Park, which, when we saw it in 1829, was 12 ft. high. In various situations, it has lived against a wall, protected during winter and flowering freely in summer. Culture, uses, price, &c., as in the preceding species.



App. i. *Other Species of Pittosporum.*

P. revolutum Ait. (*Bot. Reg.*, 186., and *Bot. Cab.*, t. 506.), from Port Jackson; *P. tomentosum* Bonp. (*Sut. Fl. Aust.*, t. 33.), from New Holland; *P. filum* Rudge; *P. hirtum* Willd., from the Canary Islands; and various other species; are, in all probability, equally hardy with *P. Tobira*. *P. cricocarpum*, from the Himalaya, has already been noticed as probably half-hardy, p. 173.

App. I. *Other Pittosporaceæ probably half-hardy.*

Senecia nepalensis Dec. a shrub from Nepal, introduced in 1820, and treated as a green-house plant; and *Bursaria spinosa* Cav. figured in *Bot. Mag.*, 1767., a shrub from New Holland, producing a profusion of elegant little white blossoms; are probably as hardy as the species of *Pittosporum*, which have been tried against a conservative wall. *Cheiranthera linearis* which Dr. Lindley states (*Bot. Reg.*, t. 1719.), to be "one of the most beautiful plants in all the flora of New Holland," would be also, if a plant of it could be obtained, well worthy of a trial.

CHAP. XIV.

OF THE HARDY SUFFRUTICOSE PLANTS OF THE ORDER
CARYOPHYLLACEÆ.

This order is introduced chiefly for the sake of the tree carnation, one of the oldest inhabitants of British gardens, and one of the finest plants that can be placed against a conservative wall. There are various species and varieties of *Dianthus*, which, technically considered, are ligneous plants. Indeed, the common pink and carnation are shrubs, and that, too, evergreen; because they do not die down to an underground bud, at the end of the growing season, like, for example, *Ranunculus acris*.



84

1. *Dianthus Caryophyllus* var. *fruticosus* Hort., the shrubby *Clow Pink*, or Tree Carnation, in its wild state, is a native of the south of France, of the Alps of Switzerland; and, in England, it is found on old ruinous walls near towns, particularly on Rochester Castle, on the old walls of Norwich, and on ruins adjoining several other old English towns. It has been cultivated in gardens from time immemorial; and is highly valuable, no less for the brilliancy of its colours, than for the aromatic fragraney of its flowers. The tree variety is one which has been originated, in all probability, by training the plant against a wall, and thus keeping it continually in a growing state without permitting it to rest, and afterwards continuing this habit by propagating it by layers or cuttings. The flowers of the tree carnation are not so various and beautiful as those of the common dwarf carnation; but they are still objects of very great beauty, and are universally admired for their symmetry of form, rich colours, and grateful odour. Planted

against an east or west wall, in calcareous loam, and carefully trained, a plant will grow at the rate of a foot a year; and, if protected during very severe winters, it will attain the height of 6 ft. or 8 ft. In Scotland, in the neighbourhood of Edinburgh, in Dalroy Nursery, in 1800, a plant against the west end of a green-house covered, in five years, a space 8 ft. wide, and 6 ft. high, flowering beautifully every year.

2. *Other suffruticose evergreen Caryophyllacæ.* *Dianthus arbuseula* Bot. Reg., 1086., *D. arboreus* L. (*Bot. Cab.*, 459., and our fig. 84.), *D. fruticosus* L., and *D. suffruticosus* W., are all beautiful glaucous-leaved evergreens, which require a little protection during winter, and produce their fragrant pink flowers from June to August. *Silene fruticosus* L., *Arenaria verticillata* W., and *Drýpis spinosa* L. (*Bot. Mag.*, 2216., and our fig. 85.), are all beautiful little evergreens, from 1 ft. to 1½ ft. in height.



85

CHAP. XV.

OF THE HARDY AND HALF-HARDY SUFFRUTICOSE PLANTS OF THE ORDER LINA'CEÆ.

This order is included in our catalogue for the sake of the tree flax, *Linum arboreum* L. (*Bot. Mag.*, t. 234, and our fig. 86.) It is a native of Candia and Italy, on the mountains, and forms a neat little evergreen bush in dry soils, in warm situations in the neighbourhood of London, requiring little or no protection, except during the most severe winters. The largest plants which we have seen of it were in a sandy border in the garden at Nonsuch Park, in Surrey. They were about 2 ft. high, and 3 ft. broad, and they produced their fine large yellow flowers from May to September; in the same situation ripening seeds, from which, or from cuttings, they are readily propagated. *Linum tauricum* W., from Tauria, which grows to the height of 1½ ft., and *L. salsoloides* Lam. with pink flowers, are equally hardy with *L. arboreum*; and all three should be included in every complete arboretum and fruticetum. *L. suffruticosum*, from Spain, an old inhabitant of our green-houses, with pink flowers, is, in all probability, as hardy as the others.



CHAP. XVI.

OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE ORDER MALVA'CEÆ.

DISTINCTIVE Characteristics. Thalamiflorous. (*H. B.*) Calyx with a valvate aestivation, mostly with an involucre. Stamens with the filaments monadelphous, and the anthers 1-celled. Pubescence starry. (*Liud. Introd. to N. S.*) The hardy ligneous species of this order are few, but splendid; the *Hibiscus syriacus*, and its different varieties, being among the most ornamental of flowering shrubs. Chemically, all the species abound in a nutritive mucilage; and, medicinally, they are emollient. The fibrous threads of the inner bark may, in most of the species, when properly prepared, be manufactured into cordage or cloth. The genera containing hardy or half-hardy species are two: *Lavatera* and *Hibiscus*; the distinctive characters of which are:—

LAVA'TERA L. Carpels capsular, 1-seeded, disposed into a ring around the axis.

HIBI'SCUS L. Carpels joined into a 5-celled capsule.

GENUS I.

LAVA'TERA L. THE LAVATERA, or TREE MALLOW. *Lin. Syst.* Monadélphia Polyándria.

Identification. *Lin. Gen.*, n. 842; *Dec. Prod.*, l. 438; *Don's Mill.*, l. 468.

Synonymes. The Tree Mallow; *Lavatère, Fr.* and *Ger.*

Gen. Char., &c. *Calyx* 5-cleft, girded by a 3- or 5-cleft involucre; its leaflets being joined as far as the middle. *Carpels* capsular, 1-seeded, disposed into a ring around the axis, which is variously dilated above the fruit. (*Dec. Prod.*, i. p. 438.) The half-hardy ligneous species are *L. marítima*, *triloba*, *subovata*, and *africana*.

■ 1. *L. MARÍ'TIMA* Gouan. The sea-side-inhabiting *Lavatera*.

Identification. *Gouan. Ill.*, p. 46; *Dec. Prod.*, l. p. 429; *Don's Mill.*, l. p. 469.

Synonymes. *L. hispánica* *Mill. Dict.*, No. 9; *L. rotundifolia* *Lam.*

Engravings. *Gouan. Ill.*, t. 11. f. 2; *Cav. Diss.*, l. 32 fig. 4; and our fig. 57

Spec. Char., &c. Stem shrubby, downy. Leaves downy, roundish, bluntly angular, 5-lobed, crenated. Pedicels axillary, solitary. (*Don's Mill.*, i. p. 469.) A shrub, growing from 2 ft. to 3 ft. high, on the rocky shores of the south of France and Spain, producing its white flowers, the petals of which have purple claws, from April to June. It was cultivated in England in 1596, and generally treated as a frame plant; but, like the following species, it only requires a wall, and a little protection, during winter. Plants attained the height of 5 ft., trained against a wall with a south-east aspect, without any protection, in the Botanic Garden, Bury St. Edmunds, about the years 1825, 1826.

87



❖ 2. *L. TRI'LOBA L.* The three-lobed-leaved Lavater

Identification. Lin. Sp., 972; Dec. Prod., i. 439; Don's Mill., i. p. 46^a
Engravings. Cav. Diss., 2. t. 31, f. 1.; Bot. Mag., 2226.

Spec. Char., &c. Stem shrubby, downy. Leaves downy, rather cordate, and somewhat 3-lobed, round, crenated. Pedicels aggregate. Sepals acuminate. (*Don's Mill.*, i. p. 469.) A tomentose shrub, growing to the height of 4 ft. in the south of Spain, and producing large pale-purple flowers in June and July. It was introduced into England in 1759, and is generally treated as a frame plant; but it will grow, and flower beautifully, trained against a wall, and slightly protected during winter. It is easily propagated by cuttings of the young wood; and it also ripens seeds.

❖ 3. *L. SUBOVA'TA Dec.* The subovate-leaved Lavatera.

Identification. Dec. Prod., i. p. 439; Don's Mill., i. p. 469.

Spec. Char., &c. Stem suffruticose. Leaves rather downy, ovate, notched, somewhat 3-lobed, with the middle lobe longest. Pedicels 1 or 2, axillary, length of petioles; lobes of calyx acuminate. (*Don's Mill.*, i. p. 469.) A shrub, growing from 2 ft. to 4 ft. high, in fields about Mogadore, on the sea coast of Morocco, producing pale purple flowers in July. It has not yet been introduced into England, but would form a desirable addition to the half-hardy species.

❖ 4. *L. AFRICA'NA Cav.* The African Lavatera.

Identification. Cav. Diss., 5. p. 282; Dec. Prod., i. p. 348; Don's Mill., i. p. 468.

Synonyme. *L. hispida* var. *Willd.*

Engraving. Cav. Diss., 5. t. 139. f. 1.; Bot. Mag. t. 2541., as *L. hispida*

Spec. Char., &c. Stem shrubby, rather tomentose from flocky down. Leaves canescent, all bluntly 5-lobed. Pedicels twin, equal in length to the petioles. Involucel 3-parted, tomentose. (*Don's Mill.*, i. p. 468.) A shrub, found in the south of Spain and the north of Africa, where it grows from 4 ft. to 6 ft. high, producing its pale purple flowers in June and July. It was cultivated in England in 1820, or earlier. In the Bury St. Edmunds Botanic Garden there were plants of it, one or more of which attained the height of 8 ft. or 9 ft. in a warm sheltered border.

App. I. *Other Species of Lavatera likely to prove half-hardy.*

L. Pseudo-Olbia Poir., with pale purple flowers, introduced in 1817; *L. phœnicea* Vent., with scarlet flowers, introduced from the Canary Islands in 1816, and forming a tree 10 ft. high; *L. Olbia* L., with reddish purple flowers, a native of Provence, which has been in culture in England, as a frame-plant, since 1570, and attains the height of 6 ft. in its native country; with *L. unguiculata* and *L. hispida* Desf. and, perhaps, some other varieties or species, all highly beautiful; might be subjected to the same treatment. Indeed, there are few kinds of plants more ornamental, when trained against a wall, than the different species of *Lavatera* and *Hibiscus*: every one knows what a splendid appearance *Hibiscus Rhsa-sinensis* makes in stoves, when so trained.

GENUS II.



HIBISCUS L. THE HIBISCUS. *Lin. Syst.* Monadelphía Polyándria.

Identification. Lin. Gen., 546; Dec. Prod., i. p. 446; Don's Mill. i. p. 476.

Synonyms. *Ketmie, Fr.; Eibisch, Ger.*

Derivations. The word *hibiscos* is one of the names given by the Greeks to the mallow. The *Hibiscus* of Pliny appears to be an umbelliferous plant; while that of Virgil is a plant with plant branches, which was made into baskets. The word *Hibiscus* is supposed by some to be derived from *ibis*, a stork, which is said to feed on some of the species. *Ketmie (Fr.)* is derived from *Kétmia*, the name given to the genus by Tournefort. The German, *Eibisch*, is the German aboriginal word for the mallow.

Gen. Char., &c. *Calyx* encompassed by a many-leaved, rarely by a few-leaved, involucl, or one with its leaves connate *Petals* not auricled. *Stigmas* 5; carpels joined into a 5-celled 5-valved capsule, with a dissepiment in the middle of each valve on the inside. *Cells* many-seeded, rarely 1-seeded. (*Don's Mill.*, adapted.)—The only hardy ligneous species is *H. syriacus*.

❧ 1. *H. SYRIACUS* L. The Syrian Hibiscus, or *Althæa Frutex*.

Identification. Lin. Sp., 978; Dec. Prod., 1. p. 448; Don's Mill., 1. 478.

Synonymes. Ketznie des Jardins, Fr.; Syrischer Eibisch, Ger.

Derivation. It is called Althæa from the resemblance of its flowers to those of the *Althæa rosea*.

Engravings. Cav. Diss., 3. l. 69. t. 1; Bot. Mag., t. 83.; and our fig. 88.

Spec. Char., &c. Stem unarmed, arboreous. Leaves ovate, wedge-shaped, 3-lobed, toothed. Pedicels hardly longer than the leaves. Involucel 6—7-leaved. (*Don's Mill.*, i. p. 478.) A deciduous shrub, a native of Syria and Carniola, where it attains the height of 6 ft., and flowers in August and September. The flowers are large, single or double, purple, white, red, or variegated. It is one of our most ornamental hardy shrubs; and, having been in cultivation since 1596, the following varieties have proceeded from it:—

❧ II. s. 2 *foliis variegatis*, the variegated-leaved Syrian Hibiscus, or *Althæa Frutex*.

❧ II. s. 3 *fløre variegato*, the variegated-flowered Syrian Hibiscus, or *Althæa Frutex*.

❧ II. s. 4 *fløre purpureo*, the purple-flowered Syrian Hibiscus, or *Althæa Frutex*.

❧ II. s. 5 *fløre purpureo pleno*, the purple-double-flowered Syrian Hibiscus, or *Althæa Frutex*.

❧ II. s. 6 *fløre rubro*, the red-flowered Syrian Hibiscus, or *Althæa Frutex*.

❧ II. s. 7 *fløre albo*, the white-flowered Syrian Hibiscus, or *Althæa Frutex*.

❧ II. s. 8 *fløre albo pleno*, the white-double-flowered Syrian Hibiscus, or *Althæa Frutex*.




Description, History, &c. A deciduous shrub, from 6 ft. to 8 ft. in height, with numerous upright white-barked branches; their general character being rather fastigate than spreading. The leaves are variously lobed. The flowers are axillary, large, and bell-shaped. In English gardens, these flowers are produced from the middle of August to the end of September; and, when the season is dry and warm, they are succeeded by capsules containing ripe seeds. It is a native of Syria and Carniola, and was introduced into England previously to 1629; being mentioned by Parkinson, in his *Paradisus* of that date, as a new shrub, somewhat tender, requiring to be kept in a large pot or tub in the house, or in a warm cellar. In the neighbourhood of Paris, it has been known for upwards of two centuries; and it is found there, as well as about London, to be perfectly hardy. At Berlin and Vienna, in severe winters, it requires protection. In the time of Du Hamel, and of Miller, there were no double-flowered varieties; but these have since been procured from seeds. Double-flowered varieties are now common both in Europe and America. The only use to which the shrub is applied is as a garden ornament, of which it is one of the most conspicuous; and it is the more valuable, because it produces its flowers at a time of the year when few shrubs are in bloom. It forms beautiful garden hedges, more especially when the different sorts are planted in a harmonious order of succession, according to their colours; and when the plants are not clipped, but carefully pruned with the knife. In the colder parts of Britain, and in the north of Germany, few ornamental shrubs better deserve being planted against a wall. It will grow in almost any soil not too wet; but, like all the *Malvaceæ*, seems to prefer one sandy, deep, and rich rather than poor. An open airy situation, where it will ripen its wood, is essential. The single-flowered varieties are propagated by seed, which come up true to their respective colours; the double-flowered varieties are propagated by layers, by grafting on the common sorts, and sometimes by cuttings of the ripened wood, planted in sand in autumn, and covered with a hand-glass during the winter. Price, in the London nurseries: seedlings, 5s. a hundred; the different single-flowered varieties, 50s. a hundred; and the va-

riegated-leaved and double-flowered varieties, 1s. 6d. each: seeds are 6s. a pound. At Bollwyller, the price is 1 franc and 50 cents each plant, for the single-flowered varieties, and 2 francs 50 cents for each of the double-flowered varieties. At New York, the single-flowered varieties are 25 cents a plant, the white-double-flowered, and the purple-double-flowered, 37 cents a plant; and the other double-flowered varieties, 50 cents a plant: the seed is 56 cents a quart.

App. I. *Other ligneous Plants of the order Malva'ceæ, which will probably be found hardy or half-hardy.*

Sida pulchella Bonpl., *Abutilon pulchellum* Bot. Mag., t. 2573, and our fig. 89. An evergreen New Holland shrub, introduced in 1824, producing its clusters of beautiful white blossoms in the gullies about Sandy Bay, and at the foot of Mount Wellington, in the neighbourhood of Hobart Town, in the depth of winter; and, as might have been expected, it is found to stand the open air, in sheltered situations, in England. There is a plant against a wall in the Botanic Garden at Kew, which has stood there since 1822, without any protection whatever. There is a plant of it at Spring Grove, Middlesex, which forms a bush between 3 ft. and 4 ft. high, which has stood several winters without protection, and flowers freely every winter and spring. We have just (February, 1836) sent to ascertain how it has passed the late severe frosts, when the thermometer was at 10°; and learn, with satisfaction, that it has suffered little or no injury. The plant is easily increased by cuttings. There is another species, or, perhaps, a variety of this one, which has stood some winters, in a warm situation, at Redleaf, in Kent, where it flowers in January, February, and March. (See *Gard. Mag.*, xi. p. 208.)



CHAP. XVII.

OF THE HARDY AND HALF HARDY LIGNEOUS PLANTS OF THE ORDER STERCULIA'CEÆ.

This order is introduced chiefly for the sake of *Sterculia platanifolia* L. (*Cav. Dis.*, 5. t. 149, and our fig. 90.) It is a tree, a native of Japan and China, with fine large palmate leaves, smooth on both surfaces, and upright branches without visible buds; that is, with the buds concealed like those of the walnut, or the *Gymnocladus*. The flowers are small and green, or greenish yellow. The tree was introduced in 1757, and, at first, treated as a green-house plant; but it has since been found to be quite hardy in the neighbourhood of London, more especially when planted against a wall. There is a tree in the Chelsea Botanic Garden 12 ft. high, which has stood out many years with only a little litter thrown round it occasionally, to protect the roots.

S. lanceolata Cav. (*Bot. Reg.*, 1256.), from China; *S. diversifolia* G. Don, from New Holland; *S. tomentosa* Thunb., from Japan; *S. peltata* G. Don, from China; are all handsome deciduous trees, with very handsome foliage, growing to the height of 20 ft.; and are probably all equally hardy with *S. platanifolia*. Some of them can be purchased, in the nurseries, at from 5s. to 7s. each.



CHAP. XVIII.

OF THE HARDY LIGNEOUS PLANTS OF THE ORDER TILIA'CEÆ.

DISTINCTIVE Characteristics. Sepals 4 or 5, with a valvate æstivation, mostly without an involucre. Petals 4 or 5, or rarely not any. Stamens hypogynous, generally numerous, with filaments separate, and anthers 2-celled. Mostly glands between the petals and ovarium. Ovary and fruit single, of 4—10 carpels grown together; cells in the fruit, at least in some, not so many as the carpels. (*Lindley, Introd. to N. S.*) The species are chiefly trees and shrubs from warm climates. The only genus which is perfectly hardy is *Tilia*.

GENUS I.

*TILIA* L. THE LIME TREE. *Lin. Syst. Polyándria Monogýnia.*

Identification. *Lin. Gen.*, 660; *Dec. Prod.*, I. p. 512; *Don's Mill.*, I. p. 552.

Synonymes. *Lime Tree Gerard*; *Lind, Anglo-Sax.*; *Tilleul, Fr.*; *Linde, Ger. and Dutch*; *Tiglio, Ital.*; *Tilo, Span.*; *Lipa, Russ.*

Derivation. In London and *Wise's Retired Gardener*, the name of *Tilia* is derived from the Greek word *ptilon*, a feather, from the feathery appearance of the bracteas; but others derive it from the Greek word *tilai*, light bodies floating in the air like wool or feathers.

Gen. Char. *Calyx* 5-parted. *Petals* 5. *Stamens* numerous, free, or somewhat polyadelphous. *Ovary* globose, villous, 1-styled, 5-celled; cells 2-ovuled. *Nut* coriaceous, 1-celled 1—2-seeded, from abortion (*Don's Mill.*, i. p. 540.)—Timber trees, with mellifluous flowers, and a remarkable bractea attached to the peduncle of each of the cymes of flowers. The species are three, according to some; and more than twice that number, according to others. Our opinion is, that they may be all included under two, *T. europæa*, and *T. americana*.

♣ 1. *T. EUROPEA* L. The European, or common, Lime Tree.

Identification. *Linn. Sp.*, 733; *Don's Mill.*, I. p. 552; *Smith's Eng. Fl.*, iii. p. 16.

Synonymes. *T. intermedia Dec. Prod.*, I. 513; *T. vulgâris Hayne Dend.*; *T. europæa borealis Wahl.*

Engravings. *Eng. Bot.*, t. 610; *Ced. Fl. Dan.*, t. 553; and our plate in Vol. II.

Spec. Char. *Petals* without scales. *Leaves* cordate, acuminate, serrated, smooth, except a tuft of hair at the origin of the veins beneath, twice the length of the petioles. *Cymes* many-flowered. *Fruit* coriaceous, downy. (*Don's Mill.*, i. p. 552.) The extensive distribution and long cultivation of this tree in Europe have given rise to the following races, or varieties, described by De Candolle and others as species; from which high authority it may be considered presumption in us to differ; but we have not done so without due consideration, and after having examined the living plants of different ages, and in different situations, with the greatest care and attention.

♣ *T. e. 2 microphylla.* The small-leaved European Lime Tree.

Synonymes. *T. microphylla Vent.*, *Willd.*, *Dec.*, and *G. Don*; *T. e. var. 2 L.*; *T. uhni-folia Scop.*; *T. sylvêstris Desf.*; *T. parvifolia Ehrh.*, *Hayne Dend.*; *T. cordata Mill.*; *Tilleul à petites Feuilles, Fr.*; *kleinblättrige Linde, or Winterlinde, Ger.*

Engravings. *Willd. Holzart.*, t. 106; *Engl. Bot.*, t. 1705; and our plate in Vol. II.

Description, &c. *Petals* without scales. *Leaves* cordate, roundish, acuminate, sharply serrated, smooth above, glaucous, and bearded beneath on the axils of the veins, as well as in hairy blotches. *Fruit* rather globose, hardly ribbed, very thin, and brittle. *Native of* Europe, in sub-mountainous woods. In England, frequent in *Essen*

and Sussex. (*Don's Mill*, i. p. 552.) This variety appears to be the male linden tree of Gerard; the timber of which, he says, is much harder, more knotty, and more yellow, than the timber of the other sort; and not very different from the timber of the elm tree. This sort we conceive to have originated in inferior soils and situations; for example, in the rocky parts of the north of Sweden, and in the milly districts of the north of Germany. This variety, according to Steven (*Nouv. Mém. de la Soc. Imp. des Nat. de Moscou*, tome iii.), is found here and there in the woods of Tauria; also in Iberia, and on this side Caucasus. It varies, he says, in a wonderful manner, in the form of the fruit, in the sinus at the base of the leaves, and in the proportion of the disk of the leaves to the petioles, as well as in the number of flowers in a cyme, colouring of the twigs, &c.; whence he agrees with Sprengel in uniting *T. triflora*, *T. intermedia*, &c., which are usually separated by authors. We doubt much if this, or any other variety, is indigenous in Britain; but it exists in plantations, and is recognised as a distinct variety by practical men, the wood being preferred by pianoforte-makers. Sir James Edward Smith says, "This species being planted along with *T. europæa*, and *T. grandifolia*, in avenues or parks, will insure a longer succession of flowers than either of the others alone." This variety is distinguishable, at first sight, from all the others, by the smallness of its leaves, which are only about 2 in. broad, and sometimes scarcely longer than their slender footstalks. The flowers are also much smaller than in any of the other varieties; and they are very fragrant, having a scent like those of the honeysuckle. There appears to be a subvariety of this in the garden of the London Horticultural Society, under the name of *T. parvifolia glauca*.

‡ *T. c. 3 platyphýlla. The broad-leaved European Lime Tree.*

Synonyms. *T. platyphýlla Scop.*; *T. cordifolia Bess.*; *T. europæa Desf.*; *T. grandifolia Ehrh.* and *Smith*; broad-leaved downy Lime Tree; Tilleul à grandes Feuilles, or Tilleul de Hollande, *Fr.*
Engravings. Vent. Diss., p. 6. t. 1. f. 2.; Bull. Fr., t. 175.; Gärtn., 2. t. 113.; and our plate in Vol. II.

Description. Petals without scales. Leaves cordate, roundish, acuminate, sharply serrated, downy beneath, origin of their veins woolly. Branches hairy. Cymes 3-flowered. Fruit woody, downy, turbinate, with 5 prominent angles. (*Don's Mill*, i. p. 553.) This tree is of about the same size as *T. europæa*, from which it is readily distinguished by its larger and rougher leaves, and, also, by its rougher bark and hispid branches. *T. europæa*, *T. e. microphýlla*, and *T. e. platyphýlla*, may be seen together, in a young state, in the London Horticultural Society's Garden; and, full-grown, in the avenue at Syon House: in both places they are readily distinguishable from each other, and are perfectly distinct; not, we think, as species, but as races. According to Sir James Edward Smith, *T. platyphýlla* is the lime tree of the south of Europe, as *T. europæa* is of the north; and he is of opinion that they are unquestionably distinct species. Steven (in *Nouv. Mém. &c.*, tome iii. 1834.) says, that, though *T. platyphýlla* is cited in the *Flora Taurico-Caucasica*, as common in Tauria and Caucasus, he never found it there, or knew of its having been observed in those regions. There is a subvariety of this sort, called *T. e. platyphýlla minor*, with leaves somewhat smaller than those of *T. e. platyphýlla*, but the difference is so trifling that it is not worth while keeping it distinct.

‡ *T. c. 4 rubra. The red-twigged European Lime Tree.*

Synonyms. *T. triflora Puer.*, in *Horn. Cat.*, 2. p. 493.; *T. corinthiaca Bosc*; *T. corallina*

Hort. Kew.; *T. europæa* β *rùbra* *Sibtharp*; *T. europæa* γ *Smith's Flor. Brit.*, 571.; *T. grandifolia* β *Smith's Eng. Flora*, 3. 19.

Description. This variety is distinguished by the redness of its young branches, and it may be properly considered as a subvariety of all the above races or kinds. In Sweden, where lime woods extend over the low part of the country for many miles together, the common lime tree is met with, in some places, perhaps for a mile together, with the twigs bright red, yellow in others, and in others quite green. In the park at Shardeloes, near Amersham, in Buckinghamshire, may be seen large lime trees, all apparently of the commonest kind (*T. europæa*), some with yellow, others with red, and others with brown or green wood. In the garden of the London Horticultural Society there is one variety with small leaves and bright yellow wood; and another, with the large rough leaves of *T. e. platyphýlla*, and bright yellow wood. Our conclusion from these, and other facts that have come before us, is, that there is a red-twigged and yellow-twigged variety or subvariety of *T. europæa*, of *T. e. microphýlla*, and of *T. e. platyphýlla*; and that *T. rùbra* *Dec.* can be nothing more than a variety of *T. e. platyphýlla*.

‡ *T. e. 5 laciniàta.* *The cut-leaved European Lime Tree.*

Synonyms. *T. platyphýlla laciniàta* *Hort.*
Engraving. Our plate in Vol. II.

Description. The leaves are smaller than those of the common species, and deeply and irregularly cut and twisted, scarcely two on the tree being alike. This variety is seldom seen of a large size; as might be expected from the diminished power of the leaves, in consequence of their diminished surface. We have never heard of its attaining a greater height than 30 ft.

‡ *T. e. 6 àurea.* *The golden-twigged European Lime Tree.*—Differing from the species in the yellowness of its twigs; and, apparently, not so vigorous in its growth as any of the other varieties, except *T. e. laciniàta*. (See our plate in Vol. II.)

‡ *T. e. 7 p. àurea.* *The golden-twigged broad-leaved European Lime Tree.*—This differs from the common broad-leaved lime in no other respect than in the yellow colour of its twigs. It is, in winter, a very distinct and very handsome variety, and may be procured in some of the London nurseries. There is a small tree in the London Horticultural Society's Garden.

‡ *T. e. 8 dasýstyła.* *The hairy-styled European Lime Tree.* *T. dasýstyła* *Steven.*—This is described as having petals without scales; leaves smooth, somewhat hairy at the base beneath; axils of veins bearded; style tomentose. It is found on the south-west coast of Tauria, at the base of the mountain Castel Dagle, where there is one tree near the public road. Steven considers it as satisfactorily distinct in the form of its fruit, and especially in the hairiness of its style. To us it appears that this variety bears the same relation to the species that *Crataëgus Oxyacantha eriocárpa* does to the species.

Other Varieties. There is a variety with variegated leaves, but it is such a ragged ill-looking plant that we deem it altogether unworthy of culture. There are some names of varieties in nurserymen's catalogues, which we have not thought worth a detailed notice; the slightest deviation being often eagerly seized on for the sake of producing something new. In the *Bollwyller Catalogue* for 1833, we have *T. aspleniúfolia nõva*, which, we presume, is a subvariety of *T. europæa laciniàta*; and M. Baumann informs us that they have lately discovered a new variety of *T. e. àurea* in a forest in their neighbourhood. In the Botanic Garden of Antwerp, there is a plant

named *T. europæa rubicaulis*, which is said to be quite different from *T. europæa rubra*. There can be no doubt that where several of the varieties are growing together, and ripen seeds, these seeds will produce different new sorts, as the result of cross-fecundation. In a work published in 1750 at Leyden, entitled *Les Agrémens de la Campagne, &c.*, the author recommends continuing all the different sorts of the lime by layers: because, says he, those which are raised from seed come up of different species; and almost all hybrids, such as the poplar-leaved lime, or the birch-leaved lime, which never arrive at the size of large trees, or become finely furnished with leaves. Those which come up with red bark, he says, grow very rapidly for a while, as do the yellow-barked varieties, but neither do they ever form large trees. The only seedlings that should be planted, with a view to this end, are such as have green leaves and shoots. (p. 207.)

Geography of T. europæa and its Varieties. *T. europæa* appears to be confined to the middle and north of Europe. The variety *T. e. platyphýlla* is found on the Alps of Switzerland, and the north of Italy; and also in Spain, Portugal, and Greece. *T. europæa* and *T. microphýlla* appear to be indigenous chiefly in the north of Germany, in Russia, and in Sweden. We have already (p. 24.) expressed our doubts as to the genus *Tilia* being indigenous in Britain; though, as Sir J. E. Smith has observed, all the varieties (species with him) are naturalised, if not all originally indigenous. Ray seems to have thought that *T. e. microphýlla* was, or might be, indigenous; but he was of a different opinion with respect to the broad-leaved variety. He says, speaking of the latter kind, "I think that Turner and Gerard err in saying that this kind grows plentifully in Essex; for, although I am an inhabitant of Essex, I have never seen the *Tilia fœmina vulgãris platyphýllos* [which, according to Smith, is a synonyme of *T. europæa* (*Eng. Flora*)] growing spontaneously there, or elsewhere in England. What we frequently find with us, in woods and hedges," he says, "is the *Tilia minõre folio*" [which, according to Smith, is a synonyme of *T. parvifolia* (*Eng. Flora*)]. "This last species," Ray continues, "is called in Lincolnshire, by the rustics, bast; because ropes are made from its bark. It flowers later than the other, and ripens its seeds more perfectly." Sir J. E. Smith gives as a native habitat of *T. europæa*, "woods and hedges upon grassy declivities:" of *T. e. platyphýlla*, "Whitstable, Surrey; and near Dorking; on the banks of the Mole, near Boxhill; and a few other places in Surrey, Norfolk, and Oxfordshire." According to Watson, *T. europæa* is common all over Britain; and in the south-western, north-eastern, and north-western counties of Ireland: *T. e. platyphýlla* is found in the north-eastern parts of England, and in the southern counties of Scotland; and *T. e. microphýlla* is found in the south-eastern and north-eastern counties of England, and north-western counties of Scotland. Mr. Edwin Lees, Hon. Sec. of the Natural History Society of Worcester, informs us that at Shawley, eight miles north-west of Worcester, there is a wood, remote from any old dwelling or public road, of above 500 acres in extent, the greater part of the undergrowths of which is composed of *T. e. microphýlla*. He also states that, in the same part of the county, there are some trees estimated to be upwards of 300 years old. So extensive a tract in Britain covered with the lime tree, we had before never heard of, and the circumstance has considerably diminished our doubts as to the tree being truly indigenous. In the *Nouveau Du Hamel*, *T. europæa* is said to be found wild in Denmark, Sweden, Bohemia, and throughout Europe generally. Pallas states that it is found through the whole of Russia, and great part of Siberia. *T. e. platyphýlla* is said to inhabit Sweden, and most parts of Europe, as far south as the alpine regions of Spain.

History. The common lime tree appears to have been known to the Greeks and Romans. The tree, according to Theophrastus, is of both sexes, which are totally different as to form; probably referring to the small-leaved

and large-leaved varieties. The leaves, he says, are sweet, and used as fodder for most kinds of cattle. The tree was highly esteemed by the Romans for its shade; and, according to Pliny, for the numerous uses to which its wood might be applied. In modern times, the lime tree was one of the first to attract the notice of writers on plants; and, accordingly, it occupies a considerable space in the works of L'Obel, Gerard, Ray, and the various dendrological authors previously to the time of Linnæus, who describes only two species, *T. europæa* and *T. americana*; but M. Ventenat, in 1798, describes three European species and three American ones. De Candolle has described ten species. Evelyn, speaking of the lime tree, says, "It is a shameful negligence that we are no better provided with nurseries for a tree so choice, and so universally acceptable. We send, commonly, for this tree into Flanders and Holland, while our woods do in some places spontaneously produce them." The lime tree has long been a favourite tree for avenues and public walks; it is planted in the streets of some of the principal towns of France, Holland, and Germany; and it forms avenues to country seats, both on the Continent and in Great Britain. "The French," Du Hamel says, "growing tired of the horsechestnut for avenues, adopted the lime for that purpose, in the time of Louis XIV.; and, accordingly, the approaches to the residences of the French, as well as English, gentry of that date are bordered with lime trees"; and Fénélon, Sir J. E. Smith observes, "in conformity to this taste, decorates, with 'flowery lime trees,' his enchanted Isle of Calypso." The lime trees in St. James's Park are said to have been planted at the suggestion of Evelyn; probably with a view to the improvement of the air, and to avert, in part, the evils pointed out in his *Fumifugium*. The Dutch plant the lime in towns, along their widest streets, and by the sides of their canals; and the whole country is perfumed by their flowers during the months of July and August. In Miller's time, the tree began to be little esteemed, on account of its coming into leaf late in the spring, and beginning to decay early in autumn; more especially when planted in a dry soil. Since the modern style of laying out grounds has rendered straight avenues unfashionable, the lime tree has not been nearly so much planted as formerly; and its chief use at present, both in Britain and on the Continent, is for planting public walks and promenades.

Properties and Uses. The wood of the lime tree is of pale yellow or white, close-grained, soft, light, and smooth, and not attacked by insects. It is used by pianoforte-makers for sounding-boards, and by cabinet-makers for a variety of purposes. It is turned into domestic utensils of various kinds; carved into toys, and turned into small boxes for the apothecaries. The most elegant use to which it is applied is for carving, for which it is superior to every other wood. Many of the fine carvings in Windsor Castle, Trinity College Library at Cambridge, and in the Duke of Devonshire's mansion at Chatsworth, are of this wood. It is supposed by some, that the blocks employed by Holbein for wood-engravings were of this tree. The wood is said to make excellent charcoal for gunpowder; even better than alder, and nearly as good as hazel. Baskets and cradles were formerly made from the twigs; and shoemakers and glovers are said to prefer planks of lime tree for cutting the finer kinds of leather upon. The leaves of the lime tree, in common with those of the elm and the poplar, were used, both in a dried and in a green state, for feeding cattle, by the Romans; and they are still collected for the same purpose in Sweden, Norway, Carniola, and Switzerland; though in Sweden, Linnæus says, they communicate a bad flavour to the milk of cows. One of the most important uses of the lime tree, in the north of Europe, is that of supplying material for forming ropes and mats; the latter of which enter extensively into European commerce. The Russian peasants weave the bark of the young shoots for the upper parts of their shoes, the outer bark serves for the soles; and they also make of it, tied together with strips of the inner bark, baskets and boxes for domestic purposes. The outer bark of old trees supplies them, like that of the birch, with tiles for covering their cottages.

Ropes are still made from the bark of the tree in Cornwall, and in some parts of Devonshire; as appears by the *Agricultural Reports* of those counties; and this, according to Ray, was formerly the case in Lincolnshire. The manufacture of mats from the inner bark of the lime tree, however, is now chiefly confined to Russia, and some parts of Sweden. Trees of from 6 in. to 1 ft. in diameter are selected in the woods; and in the beginning of summer, when, from the expansion produced by the ascending sap, the bark parts freely from the wood, it is stripped from the trees in lengths of from 6 ft. to 8 ft. These are afterwards steeped in water, till the bark separates freely into layers; it is then taken out and separated into ribands or strands, which are hung up in the shade, generally in the wood where the tree grew from which they were taken; and, in the course of the summer, they are manufactured into the mats so much in use by gardeners and upholsterers, and for covering packages generally. The fishermen of Sweden make nets for catching fish of the fibres of the inner bark, separated, by maceration, so as to form a kind of flax; and the shepherds of Carniola weave a coarse cloth of it, which serves them for their ordinary clothing. The trees from which the bark is taken are cut down during the same summer, collected into open places in the woods, cut into short lengths, and burned in heaps, so as to form charcoal. The sap of the lime tree, drawn off in spring, and evaporated, affords a considerable quantity of sugar; and Adanson suggested the idea of employing it for this purpose in France, along with the sap of the birch and the maple. The honey produced by the flowers is considered superior to all other kinds for its delicacy, selling at three or four times the price of common honey; and it is used exclusively in medicine, and for making some particular kinds of *liqueurs*, more especially Rosoglia. This lime tree honey is only to be procured at the little town of Kowno, on the river Niemen, in Lithuania, which is surrounded by an extensive forest of limes. An account of this forest, of the mode of managing the bees in it, and of disposing of the honey, &c., was given to Sir John Sinclair by the botanist Hove, and will be found printed as an appendix to the *Husbandry of Scotland*. The Jews of Poland produce a close imitation of this honey, by bleaching the common sort in the open air during frosty weather. (See Bright's *Travels in Hungary*.) The fruit of the lime tree had long been thought of little use, till Missa, a physician of the faculty of Paris, by triturating it, mixed with some of its flowers, succeeded in procuring a butter, perfectly resembling chocolate; having the same taste, and giving the same paste, as the cocoa. This was in the time of Frederick the Great; who, feeling a greater interest in the discovery than the French, who were in possession of plantations of the cocoa in their colonies, engaged the chemist Marcgraf to prove the observations of Missa, which he did entirely to the satisfaction of Frederick; but, unfortunately, it was found that the lime tree chocolate did not keep. On this Ventenat remarks, that, if the subject had been pursued a little further, and the fruits of some of the American species of limes taken, the success would probably have been complete. In landscape-gardening the principal use of the lime is as a detached tree on a lawn, or in scenery which is decidedly gardenesque; because, from the symmetrical and regular form of the tree, it is unfitted for grouping with other trees in the picturesque manner. London and Wise recommended the lime tree, as preferable to the elm, for sheltering gardens or orchards; because the roots do not, like those of the elm, spread out and impoverish all around them. In the *Retired Gardener*, the chief use of the tree is said to be for bowers, or covered ways 18 ft. or 20 ft. high: the lime being trained to a shelter roof. Evelyn commends the lime for its "unparalleled beauty" for walks; "because" he says, "it will grow in almost all grounds, lasts long, soon heals its wounds when pruned, affects uprightness, stoutly resists a storm, and seldom becomes hollow." Scattered trees of it harmonise well with immense masses of Grecian or Roman architecture; but it is less suitable for the narrow perpendicular forms of the Gothic. For architectural

gardening it is well adapted, from the patience with which it bears the knife or the shears. In some of the public gardens of recreation on the Continent, and especially in those in the neighbourhood of Paris and Amsterdam, there are very imposing colonnades, arcades, walls, pyramids, and other architectural-looking masses, formed of this tree.

Soil and Situation. A deep and rather light soil is recommended for the lime tree by Du Hamel (*Traité des Arbres*); but the largest trees are generally found in a good loamy soil. In Lithuania, where the tree is more abundant, and of a larger size, than it is either in Russia or Poland, the soil, as we particularly remarked about Kowno, when in that country in 1813, is rather a clayey loam than a sandy one. This agrees with an observation of Du Hamel, in another of his works (*Exploitation des Bois*), that the lime tree gets to a prodigious size in an argillaceous soil inclining somewhat to sand, and rather moist. In dry situations, the tree never attains a large size, and it loses its leaves earlier than any other tree. Being a tree of the plains, rather than of the mountains, it does not appear suitable for exposed surfaces: but it requires a pure air rather than otherwise; for, though it is found in towns on the Continent, and sparingly so in Britain, the smoke of mineral coal seems more injurious to it than it is to the platanus, the elm, or some other trees.

Propagation and Culture. It is seldom propagated otherwise than by layers, which are made, in the nurseries, in autumn and winter, and which become rooted, so as to admit of being taken off, in a year. The tree, in Britain at least, appears seldom to ripen its seeds; but Evelyn states that he received many of these from Holland, and that plants may be raised from them; though, he says, with better success from suckers. Du Hamel says that the lime tree may be raised from seeds, which ought to be sown immediately after being gathered; because, if they are preserved dry till the following spring, they will often not come up till the second year. If, however, the seeds are mixed with sand, or with soil, not too dry, and kept in that state till the following spring, they will generally come up the first year. Owing to the slowness of the growth of plants raised from seeds, Du Hamel states, the French gardeners, when they want a supply of young lime trees, cut over an old one close by the surface of the ground, which soon sends up a great number of shoots: among these they throw in a quantity of soil, which they allow to remain one, or two, or three years; after which they find the shoots well rooted, and of a sufficient height and strength to be planted at once where they are finally to remain. This mode is still practised in France and Belgium, both with the lime and the elm. (See *Agréments de la Campagne*, liv. ii.) We have seen the plants, or shoots, 15 ft. or 20 ft. high, with very few roots when they were first taken off: but all the branches being cut off close to the stems, and the stems shortened to 6 ft. or 7 ft., and the roots also pruned, they are planted, and seldom fail to grow; all the young shoots produced the first season after planting being removed, except one to serve as a leader. The lime tree bears transplanting when of a considerable size; but, when it is grown in the nurseries for this purpose, it ought always to be taken up and replanted every two or three years. A tree which has stood some years without being removed should always have the roots cut round, at 3 ft. or 4 ft. from the stem, a year before removal, for the purpose of stunting the growth, both of the head and roots, and of forming smaller roots and fibres. Evelyn mentions some very large lime trees which the prince elector took out of his forests at Heidelberg, to a steep hill "exceedingly exposed to the heat of the sun, and that in the heat of summer. They grow behind that strong tower on the south-west and most torrid part of the eminence, being a dry, reddish, barren earth; yet do they prosper rarely well: but the heads were cut off, and the pits into which they were transplanted were (by the industry and direction of Monsieur De Son, a Frenchman, and an admirable mechanic, who himself related it to me) filled with a composition of earth and cow-dung,

which was exceedingly beaten, and so diluted with water, that it became almost a liquid pap. It was into this that he plunged the roots, covering the surface with the turf: a singular example of removing great trees at such a season, and therefore taken notice of here expressly." This operation was probably performed before midsummer, when the trees, not having spent their vital or growing force for the season, might still send out shoots and fibrous roots, which would preserve them alive till the following year, when they would probably grow freely. If it had not been intended that they should grow a little the first year, the puddle formed with so much care would have been unnecessary.

Statistics. We have received the dimensions and age of some hundreds of lime trees, with notices of the soil and situation in which they grow, in different parts of Britain and the continent of Europe: from which we shall select but a very few examples, the tree being sufficiently well known.

Tilia europæa in the Environs of London. The oldest tree that we know of is at Fulham Palace. The head of the tree has suffered great injury from time and the weather; and is not remarkable either for its height or breadth; but the trunk is between 7 ft. and 8 ft. in diameter. At Kenwood there are trees 90 years planted, which are 90 ft. high. At Syon there are trees of *T. europæa*, of *T. e. microphylla*, and *T. e. platyphyllo*, which are supposed to be about 80 years planted, and are 75 ft. high.

Tilia europæa South of London. In Kent, at Cobham Hall, there is a tree 97 ft. high, and above 9 ft. in diameter, which contains 136 ft. of timber. At Knowle there is an immense lime tree, the dimensions of which have not been sent us; but when we saw it, in 1820, it covered, as we estimated at the time, nearly a quarter of an acre of ground. The lower branches, which extended to a great length, had rested with their extremities on the soil, rooted into it, and sent up a circle of young trees, which surrounded the old or central one. The outer branches of this outer row of trees had, in their turn, stretched out, rested on the ground, and thrown up a second circle of trees, which, at the time we saw them, were from 20 ft. to 30 ft. high. The tree stands on a lawn in an ancient geometrical garden; and must be at least two centuries old; the soil is a deep sandy loam. At Ashstead, in Surrey, there are trees 95 ft. high, with trunks from 7 ft. to 8 ft. in diameter: the soil is an adhesive loam on chalk.

Tilia europæa North of London. In Berkshire, at Ditton Park, there is a tree which is known to be upwards of 200 years old; and, though it is only 80 ft. high, yet the diameter of the trunk, at 1 ft. from the ground, is 22 ft. 10 in.; it grows on strong loam on gravel, and is supplied with abundance of water, from this gravel being on a level with the Thames. In Hertfordshire, at Moor Park, there are several magnificent lime trees, one of which has been beautifully portrayed by Mr. Strutt; nineteen large branches, 6 ft. or 8 ft. in girth, strike out horizontally from 67 ft. to 70 ft. in length, and these support three or four upright limbs; the tree is in full vigour, and its branches droop down and rest on the ground; the trunk girths 23 ft. 3 in.; and the head is 122 ft. in diameter; it is nearly 100 ft. high; and contains, by actual measurement, 875 ft. of timber. In Norfolk, at Merton Park, there is a tree 75 ft. high, which, at 1 ft. from the ground, is about 6 ft. in diameter. In Somersetshire, at Brockleby Hall, are three lime trees with trunks from 15 ft. to 17 ft. in circumference, and 60 ft. high. In Staffordshire, at Enville, are some of the finest trees of *T. europæa* and *T. e. microphylla* in England; they are nearly 100 ft. high, and they are completely feathered to the ground. In Warwickshire, at Crompton, a tree, between 60 ft. and 70 ft. high, has a trunk measuring, at 4 ft. from the ground, 15 ft. in girth; in 9 ft. to 12 ft. high the trunk divides into six upright branches, each from 50 ft. to 60 ft. high. In Worcestershire, between Horford and Ombersley, on the edge of a small pool, there is a tree of *T. e. microphylla* estimated at upwards of 300 years of age; it is 70 ft. high, and the trunk, at 8 ft. from the ground, 30 ft. in circumference. There are several other trees of the same variety in the neighbourhood estimated to be as old, and nearly as large; at Croomie, *T. e. platyphyllo*, 40 years planted, is 70 ft. high, with a trunk 2½ ft. in diameter at 1 ft. from the ground, and the diameter of the space covered by its branches 70 ft., the soil is a strong red loam. In Yorkshire, at Bolton, are two lime trees of large size, supposed to be *T. e. microphylla*, noticed by Whitaker, in his *History of Craven*, which still exist. In the Park at Thrickleby Hall, there is a lime tree with a branch like a teapot handle, noticed by Sir J. E. Smith, in his correspondence.

Tilia europæa in Scotland. At Hopeton House, *T. e. rubra*, 100 years planted, is 70 ft. high, with a trunk 3½ ft. in diameter at 1 ft. from the ground, and the diameter of the space covered by the branches 33 ft.: it stands in an avenue among several others of about the same height. At Gordon Castle, in Aberdeenshire, there are trees above 80 ft. high, and with trunks above 5 ft. in diameter at 1 ft. from the ground. In Ayrshire, at Rosenearth Castle, there are several trees with trunks 3½ ft. in diameter, and from 75 ft. to 80 ft. high. In Perthshire, at Taymouth Castle, there is a noble avenue of lime trees, which has been 100 years planted, and the trees are 82 ft. high and upwards; and 6 ft., or more, in diameter, at a foot from the ground. In the Perth Nursery there is a tree of the *T. e. platyphyllo* minor, 60 years planted, 66 ft. high; the diameter of the trunk, at 1 ft. from the ground, 23 in.; and that of the space covered by its branches 42 ft. In general, when the soil is properly prepared, the tree grows at the average rate of 2 ft. a year, for the first 12 or 15 years, in the milder parts of Scotland.

Tilia europæa in Ireland. In the park at Charleville Forest, county of Meath, there is a tree 110 ft. high, with a trunk, at 1 ft. from the ground, 5½ ft. in diameter: it grows in brown loam resting on a limestone gravel, in an open situation. In the plantations on the same estate, the tree attains the height of from 25 ft. to 30 ft. in 10 years. At Florence Court there is a tree, 38 years planted, 46 ft. high, with a trunk 4 ft. in diameter at 1 ft. from the ground, and the diameter of the space covered by the branches 46 ft.; the soil a retentive loam. At Moira, near Beltast, *T. e. platyphyllo* minor has attained the height of 85 ft., with a trunk 4 ft. in diameter at 1 ft. from the ground; the branches covering a space of 60 ft. in diameter.

Tilia europæa in Foreign Countries. In France, in the Paris Garden, *T. e. platyphyllo*, 120 years planted, is 75 ft. high, and the space covered by its branches is 87 ft. in circumference; at Mereville, *T. e. microphylla*, 60 years planted, is 60 ft. high; its trunk, at 1 ft. from the ground, is 5 ft. in diameter; and the diameter of the space covered by its branches is 40 ft.; in the botanic garden at Toulon, *T. europæa*, 40 years planted, is 50 ft. high, with a trunk 4 ft. in diameter; in the public walks at Nantes, *T. europæa*, 70 years planted, is 80 ft. high, with a trunk 4½ ft. in diameter. In Belgium and Holland this species and its varieties abound: the largest are in "the wood" at the

Hague, some of which are between 70 ft. and 80 ft. in height, with trunks between 3 ft. and 4 ft. in diameter. In the neighbourhood of Ghent and of Brussels, the tree is seldom to be found above 60 ft. high; and in the native forests, where it is indigenous, not often so much. In Austria, in the park at Schönbrunn, there is a lime tree, 70 years planted, which is 75 ft. high; at Laxenburg, 60 years planted and 55 ft. high; and many others of similar heights, or higher, are to be found. In Wirtenberg, at Neustadt an der Linde, is a tree, from which the town takes its name, of unknown age, and great size; the trunk girths 54 ft., and rises 15 ft. high before the branches begin; and the whole height of the tree is about 100 ft. The branches extend to nearly 100 ft. on each side of the trunk, and they are supported by 108 pillars, some of which are of wood, and some of stone; there is a place of entertainment formed in the head of the tree, which is ascended to by a flight of steps. In the hollows of the branches, earth has been placed, and gooseberry bushes planted, which bear fruit which is sold to visitors. The avenue of lime trees in Berlin (*Der Linden Strasse*) is celebrated. In Denmark, *T. europæa* and *T. e. microphylla* attain the height of from 60 ft. to 70 ft. in the royal gardens in the neighbourhood of Copenhagen. In Sweden, in the botanic garden at Lund, there is a lime tree which is 60 ft. high, with a trunk 3 ft. 3 in. in diameter. In Switzerland, according to Cox, and to M. Alphonse De Candolle, p. 169, there are some very large lime trees. One, near Morges, has a trunk 24 ft. 4 in. in circumference; another, near the great church at Berne, which was planted before the year 1410, is 36 ft. in girth; and a third, near Morat, which is, probably, one of those referred to by M. De Candolle, is not less than 90 ft. high, and of the same girth as the last. In p. 162. some other remarkable lime trees are mentioned. Mr. Strutt, the most celebrated artist in dendrography which this country has ever produced, and who is now (January, 1836) in Switzerland, has, we believe, taken sketches of all these trees, which he will, in all probability, engrave and publish on his return to England.

Commercial Statistics. The common lime is propagated for sale in all the European nurseries, and in some of those of North America. The price varies according to the size of the plants. In London, plants from layers, 3 ft. to 4 ft. high, cost 20s. a hundred; from 5 ft. to 6 ft. high, 30s. a hundred; and from 7 ft. to 10 ft. high, 2s. 6d. each. At Bollwyler, plants of the common lime are 1 franc each; of the common yellow-twigg'd variety 2 francs each; and of the cut-leaved variety, which, we believe, was originally brought from that nursery, 5 francs each. In New York, ?

Æ 2. *T. (EUR.) ALBA* Waldst. & Kit. The white-leaved European Lime Tree.

Identification. Waldst. and Kit. Pl. Hung.; Wats. Dendr. Brit.; Hort.; Kew., 2. p. 230.; Hayne Dend., p. 113.; Don's Mill., 1. p. 553.

Synonymes. *T. americana* Du Roi; *T. argentea* Desf., Dec. Cat. Hort. Monsp., and Dec. Prod., 1. p. 513.; *T. rotundifolia* Vent. and N. Du Ham.; *T. tomentosa* Moench.

Engravings. Waldst. and Kit. Pl. Hung., 1. t. 3.; Vent. Diss., t. 4.; N. Du Ham., t. 52.; Wats. Dend., t. 71.; and our plate in Vol. II.

Spec. Char. Petals each with a scale at the base inside. Leaves cordate, somewhat acuminate, and rather unequal at the base, serrated, clothed with white down beneath, but smooth above, 4 times longer than the petioles. Fruit ovate, with 5 obscure ribs. (*Don's Mill.*, i. p. 553.) Fruit evidently ribbed. (*Steren*, in *Nouv. Mém. de la Soc. Imp. des Naturalistes de Moscou*, tome iii. p. 103.) A native of Hungary; with yellowish and very fragrant flowers, produced from June to August. Introduced in 1767.

Description. Our own opinion is, that this is nothing more than a very distinct race of the common lime; notwithstanding the circumstance of its having scales to its petals, as noticed by Watson in his *Dendrologia*, which no one of the other varieties of *T. europæa* is said to possess. Even allowing this structure to be permanent in the Hungarian lime, the tree bears such a general resemblance to *T. europæa* in all its main features, that it seems to us impossible to doubt the identity of their origin. We are strengthened in this opinion by the circumstance of its being found only in isolated stations in the Hungarian forests. We have, however, placed this lime by itself, rather than among the other varieties; because, from the whiteness of its foliage, it is far more obviously distinct than *T. e. platyphylla* or *T. e. microphylla*. The tree is at once distinguishable from all the other species and varieties by this white appearance, even at a considerable distance, and by the strikingly snowy hue of its leaves when they are ruffled by the wind. Its wood and shoots resemble those of the common lime; but it does not attain the same height as that tree. At High Clere, where a number of plants of this species are sprinkled along the approach road, its line of direction may be traced at some miles' distance, through the apparently dense forest, by their white tops appearing at intervals among the other trees.

Geography, History, &c. The white lime was discovered by Kitaibel in the woods of Hungary, where it is rare; it was also seen by Olivier near Constantinople. It was sent to Gordon, at Mile End, in the year 1767; whence it

passed into other nurseries, and has since been rather extensively cultivated, though not so much so as it deserves from its very striking appearance. Twelve years after it was introduced into England, we are informed in the *Nouveau Du Hamel*, printed in the time of the French Republic, that "the citizens Thouin and Cels received some plants from Kew, and propagated them with success; the former in the garden of the Museum, and the other in his nursery at Arcueil. Some of these plants grew with such vigour, that, in the 6th and 7th years of the Republic, citizen Thouin sowed their seeds, and raised several young plants from them." The tree alluded to in the Paris Garden is now according to the Return Paper sent to us by Professor Mirbel, 55 ft. high, and its branches cover a space of 63 ft. in diameter. It is propagated in the same manner as the common species, and requires a similar soil and situation. When first introduced, it was propagated by grafting on *T. e. platyphýlla*; and this is still the practice in some nurseries, especially in Continental ones.

Statistics. There is a good specimen of the white lime in the Kew arboretum, and a fine tree at Walton upon Thames, 60 ft. high. There are also a great many at High Clere, in Berkshire, some of which, in 36 years, have attained the height of 60 ft., with diameters of from 1 ft. 6 in. to 2 ft., on a retentive shallow soil on chalk. There are some good specimens at Deepdene, also on chalk. One at Croome, in Worcestershire, only 30 years planted, on a loamy soil, has attained the height of 50 ft., with a trunk 2 ft. in diameter. One in the Glasnevin Botanic Garden, 35 years planted, is 35 ft. high; and, in the principal botanic gardens on the Continent, there are trees of from 30 ft. to 50 ft. in height. Plants, in the London nurseries, cost 2s. each; at Bollwyller, 1 franc 50 cents; at New York, ?.

✠ *T. (?e.) a. petiolàris* Dec., the long-petioled-leaved Lime Tree, described by De Candolle from dried specimens without flower or fruit, is, probably, only a variety of *T. álba*. He has placed it in one of two sections in which the species have the petals each with a scale at its base, inside. It is said to have leaves cordate, acuminate, twice the length of the petioles, serrated, smooth above, but white beneath from close-pressed down. It is said to be cultivated in the gardens of Odessa, but has not yet been introduced into England. Steven has stated (*Nouv. Mem. de la Soc. Imp. des Nat. de Moscou*, tome iii. p. 104.), that *T. petiolàris* Dec. certainly belongs to *T. argentea* [*T. e. álba*]; for the length of the petioles varies often upon the same branch, not only in this, but in all species of *Tilia*; wherefore Sprengel has judiciously omitted it from his *Systema*.

✠ 3. *T. AMERICA`NA* L. The American Lime Tree.

Identification. Lin. Sp., 733.; Hort. Kew.; Willd. Spec.

Synonymes. *T. glàbra*, Vent.; *T. caroliniana* Wangenh.; *T. canadensis* Michaux; *T. glàbra* Dec., Hayne's Dendr., and Don's Mill.; the smooth-leaved, or black, Lime Tree, and Bass Wood, Amer. Engravings. Vent. Diss., t. 2.; Wats. Dendr. Brit., t. 134.; and our plate in Vol. II.

Spec. Char. Petals each with a scale at the base, inside. Leaves profoundly cordate, abruptly acuminate, sharply serrated, somewhat coriaceous, smooth. Petals truncate and crenate at the apex, equal in length to the style. Fruit ovate, somewhat ribbed. (*Don's Mill.*, i. p. 553.) Canada. Yellowish-white flowers. July and August. 1752. On a general view of the trees, the most obvious external differential characteristics of the European and American limes appear to us to be, that the former have regularly cordate, and the latter obliquely cordate, leaves. The other American limes we consider to be nothing more than varieties of this species.

Description. The American lime, in its native country, attains the height of 80 ft. or upwards, with a straight uniform trunk, having an ample finely tufted summit. In England, there are some specimens of from 50 ft. to 60 ft. in height: in summer, these are readily distinguished from the European limes by the largeness of the leaves, which are heart-shaped, acutely pointed, deep green and glabrous on their upper sides, and pale green beneath. Some of the leaves have a tendency to be slightly pubescent; but they are generally smooth and shining. The flowers, which are large, appear, in Canada, in June and July; but, in England, not till the end of July, or the beginning of August, when those of the common sort are decaying. In winter, this species is readily recognised by the robust appearance of the trunk and branches, and by the dark-brown colour of the bark on the young shoots. This circumstance alone is a very marked distinction; and has, no doubt, procured for the species the name of the black line tree. The largest tree, that we know of, of this

species is at White Knights, near Reading : it is about 60 ft. high ; and, at a distance, the tufting of the masses has a very singular appearance.

Geography, History, &c. This species is found in Canada, and in the northern parts of the United States : it is less common towards the south ; and, in Virginia, the Carolinas, and Georgia, it is found only on the Alleghany Mountains. It is abundant in Tennessee, on the borders of Lake Erie and Lake Ontario, on loose, deep, fertile soil. The wood of this tree is white and tender ; and, in the northern states of America, where the tulip tree does not attain a timber size, it is used for the panels of carriage bodies, and the seats of chairs. On the Ohio, the images affixed to the prows of vessels are made of this wood ; and the inner bark is formed into ropes, as that of the *T. europæa* is in the north of Europe. This lime was cultivated by Miller in 1752, but has not been very extensively distributed. There are trees of it of 10 years' growth in the London Horticultural Society's Garden ; and it is propagated, generally by grafting, in some of the British and Continental nurseries. Price, in London, 2s. 6d. a plant ; at Bollwyller, 1 franc 50 cents ; and at New York, 50 cents.

† *T. a. 2 laxiflora.* *The loose-cymed-flowered American Lime Tree.*

Synonyme. *T. laxiflora Michx., Pursh, Dec., Hayne's Dendr., and Don's Mill.*

Description, Geography, &c. Petals each with a scale at the base, inside. Leaves cordate, gradually acuminate, serrated, membranaceous, smooth. Cymes loose. Petals emarginate, shorter than the style. Fruit globose. (*Don's Mill.*, i. p. 553., adapted.) A most distinct species, according to Pursh. G. Don observes that it is generally confounded with *T. glabra* ; which, if the trees in the London Horticultural Society's Garden be correctly named, is not to be wondered at : for their general resemblance is so great, that we have no doubt of their being essentially the same species, notwithstanding the comparatively loose cymes of the flowers, which, however, we have never seen ; no plants of this variety, that we are aware of, having yet flowered in Britain. This variety is said to be found from Maryland to Georgia, near the sea-coast, where it grows to the height of 50 ft., and produces its yellowish-white sweet-scented flowers from May to July. It was introduced into England in 1820, and is but sparingly cultivated. There is a plant 6 ft. or 8 ft. high in the London Horticultural Society's Garden.

† *T. a. 3 pubescens.* *The pubescent-leaved American Lime Tree.*

Synonyms. *T. pubescens Hort. Kew, Willd., Dec., Don's Mill, Vent., Michaux, and Hayne's Dendr. ; T. caroliniana Mill. ; T. americana Walt. Engravings.* Vent. Diss., p. 10. t. 3. ; Wats. Dendr., t. 135., and our plate in Vol II.

Description. Petals each with a scale at the base inside. Leaves truncate at the base, somewhat cordate and oblique, denticulately serrated, pubescent beneath. Petals emarginate, shorter than the style. Fruit globose, even. (*Don's Mill.*, i. p. 553.) This variety is of much less vigorous growth than the preceding ; the leaves are much smaller, and the branches more slender. The leaves are most pubescent after their first expansion : as they increase in size, a part of the pubescence falls off, and the hairs which remain form little starry tufts. The colour of the bark is dark, and shows that it is more nearly allied to *T. americana* than to *T. a. laxiflora*. It is a native of the southern parts of the United States and the Floridas, where it grows on the borders of rivers and large marshes, where the soil is cool and rich, and not subject to inundation. It is the only variety found in the maritime parts of Carolina and Georgia. Michaux found it principally in the neighbourhood of Charleston, growing to the height of 40 ft. or 50 ft., and having the general appearance of the common American species. Its leaves, he says, differ widely in size, according to the exposure in which they grow : in dry and open places they are only 2 in. in diameter ; but in cool and shaded

situations they are twice that size. The flowers, which resemble those of the common American species, appear in June, and they vary in the same proportion as the leaves. Seeds of this variety were brought to England by Catesby in 1726; but it does not appear to have been much cultivated. There is a tree of it in the arboretum at Kew, one in the London Horticultural Society's Garden, and one at Messrs. Loddiges's; and it may be found in a few of the principal nurseries. In New York, the price is 50 cents a plant.

‡ *T. a. 4 pubescens leptophýlla.* *The thin-leaved pubescent American Lime Tree.*

Synonymes. *T. pubescens leptophýlla Vent.*; *T. mississippensis Desf.*

Description. This variety is described by Ventenat as having very thin leaves, with only a few fine serratures. De Candolle doubts whether it may not prove a distinct species. There is a plant bearing this name in the garden of the London Horticultural Society, which closely resembles *T. a. pubescens*; and, if this be correctly named, we should have no doubt of its being nothing more than a variety of that race.

‡ *T. a. 5. heterophýlla.* *The various-leaved American Lime Tree.*

Synonymes. *T. heterophýlla Vent., Dec.*; *T. álba Michx.*; the White Lime.

Engravings. *Vent. Diss., t. 5.*; *Michx. Arb., 3. t. 2.*; and our plate in Vol. II.

Description. Petals each with a scale at the base inside. Leaves ovate, downy beneath, sometimes cordate at the base, sometimes obliquely or equally truncate. Fruit globose, with 5 ribs. (*Don's Mill., i. p. 553.*) According to Michaux, this tree rarely exceeds the height of 40 ft. in its native habitats; and, according to the *Nouveau Du Hamel*, it does not exceed the height of 20 ft. in France, though it has been introduced into that country upwards of 70 years. The young branches of this variety are covered with a smooth silver-grey bark; by which, and by their thickness, rough surface, and the large size of their buds, the tree is easily recognised in winter. The leaves are larger than those of any other variety, either American or European; obliquely heart-shaped and pointed like those of all the other American varieties; of a dark green on the upper surface, and whitish beneath; with small reddish tufts of hairs at the intersections of the principal nerves. The flowers appear, in America, in June; and, as well as the floral leaf, are larger than those of any other lime tree. The petals are larger and whiter, and have an agreeable odour. The seeds are round, or, rather, oval, and downy. There is a tree of this variety in the London Horticultural Society's Garden, which, if correctly named, will prove it, as we think, to be only a variety of *T. americana*, more nearly approaching *T. a. laxiflora* than *T. a. pubescens*.

Geography, History, &c. *T. a. heterophýlla*, or the white lime, as it is called in America, is abundant in Maryland, Delaware, and the western states. It does not grow, like the common species (*T. americana*), in elevated places, nor amidst the trees of the forests, but is almost always found on the banks of rivers. It is particularly observed on those of the Susquehanna, the Ohio, and the streams which flow into them; but it rarely exceeds 40 ft. in height, with a trunk of from 12 in. to 18 in. in diameter. The wood is white and tender, and is seldom applied to any use in the arts. It is remarkable, that, although this tree was known in France in the time of Du Hamel, in 1755, it should not have been introduced into England till 1811. We are not aware of any plants of it, except those in the garden of the London Horticultural Society, which have not been planted above 8 or 10 years. Like all the other American varieties, it may be considered as highly ornamental, and well deserving a place in collections, where the climate is not much more severe than that of London, or where, if the cold is greater in winter, the heat is proportionately greater in summer, and is sufficient to give such a degree

of maturity to the young wood as will enable it to endure the winter.

App. i. *Doubtful Varieties of Tilia europæa and americana.*

In the garden of the London Horticultural Society there are some names attached to young plants of lime trees, which will not be found in the foregoing enumeration as distinct. The reason is, that we have not been able to satisfy ourselves that they were sufficiently distinct from the species and varieties which we have enumerated to be worth recording. Among these names are, *T. platyphyllo vera*, *T. p. minor*, *T. præcox*, *T. vitifolia*, *T. laxiflora microphyllo*, and *T. pubescens rugosa*.

App. I. *Other Species belonging to the Order Tiliaceæ.*

The genus *Grèwia* L. (*Dec. Prod.*, i. p. 508., and *Don's Mill.*, i. p. 547.) consists of a great number of species, mostly natives of tropical climates; but, as several of them inhabit Himalaya and the islands in the Indian Ocean, it is probable that some species might succeed in the neighbourhood of London, against a wall. In the Himalaya, Mr. Boyle states that the inner bark of *Grèwia oppositifolia* is used for the same purposes as that of the lime tree is in Europe; that the leaves of *G. didyma* and other species are given as fodder to cattle, and are dried and stacked up for winter use. The wood is used, on account of its lightness, for making boats. Some of the species of *Grèwia* yield pleasant acid berries, much used for making sherbet. The species cannot be considered as likely to prove very ornamental in our gardens, but they will enrich them by increasing the variety. The species which might be tried are the following: *G. oppositifolia Roxb.*, a Nepal shrub of 6 ft. with purple flowers; *G. biloba G. Don*, a shrub, native of China; *G. occidentalis L.*, a shrub, native of the Cape of Good Hope, which grows about the height of 10 ft., has leaves like those of the small elm, purple flowers, and has been cultivated in British green-houses since 1690; *G. populifolia Vahl*, a shrub with leaves like those of *Pópulus tremula*, found in Egypt; *G. pàmila Ham.*, found in Nepal; *G. velutina Vahl*, found in Arabia; *G. echinulata Delile*, found in the north of Africa. The only one of these species which is at present in British gardens is *G. occidentalis Bot. Mag.*, t. 422., which well deserves a trial against a conservative wall.

CHAP. XIX.



OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE ORDER
TERNSTRÖMIA[^]CEÆ.

DISTINCTIVE Characteristics. Thalamiflorous. (*H. B.*) Calyx with an imbricate æstivation. Stamens with filaments monadelphous or polyadelphous, and anthers 2-celled to 4-celled. Leaves alternate. (*Lindley's Introd. to N. S.*; and *Don's Mill.*) The species which endure the open air, belonging to this order, are included in two tribes, *Gordonicæ*, and *Canellicæ*.

Sect. I. *Hardy and half-hardy ligneous Plants belonging to the Tribe Gordonicæ.*

Common Character. Sepals 5, free, or joined together at the base. Petals usually connected at the base. Stamens numerous, with filiform filaments, connected at the base. Anther oval, vane-like. Styles 5, distinct, or connected only at the base, or usually united to the tip. Carpels 5, more or less united, 1—2-seeded. Seeds few. Albumen none. Embryo straight. Radicle oblong. Cotyledons leafy, folded lengthwise. Plumule inconspicuous. Trees and shrubs of America; a few of Asia. Leaves alternate, usually deciduous, oval or oblong, feather-nerved, and without stipules. (*Dec. Prod.*, i. p. 527.) The genera are three; and their differential characters are as follow:—

MALACHODE'NDRON Cav. Calyx attended by a single bractea. Petals 5—6, with the limb finely notched. Ovary marked with 5 furrows. Styles 5, free, separate to the base. Stigmas capitate. Carpels capsular, 5, connected, 1-seeded. (*Don's Mill.*, i. p. 564.)

STUA'RTIA. Sepals 5, connected almost to the middle, bibracteate. Petals 5. Style crowned by a 5-lobed stigma. Capsule woody, 5-celled, 5-valved. Cells 1—2-seeded. Seeds wingless. (*Don's Mill.*, i. p. 564.)

GORDO'NIA. Sepals 5, coriaceous. Petals 5, adhering to the tube of the stamens, and connected together at the base. Style crowned by 5 stigmas. Capsule 5-celled, 5-valved; cells 2—5-seeded. Seeds ending in a leafy wing. (*Don's Mill.*, i. p. 564.)

GENUS I.



[**MALACHODE'NDRON** Cav. THE MALACHODENDRON. *Lin. Syst.*
Monadélphia Polyándria.

Identification. Cav. Diss., 5. p. 502.; Juss. Gen., 275.; Mitch. Gen., 16. p. 38.; Dec. Prod., 1. p. 528.; and Don's Mill., 1. p. 572.

Synonymes. Stua'rtia L'Hérit.; Stewa'rtia L.

Derivation. From *malakos*, soft, and *dendron*, a tree; in allusion, perhaps, to the quality of the timber: or, possibly, from the flowers resembling those of the mallow, the Greek name for which is *malachē*.

Gen. Char., &c. Calyx 5-cleft, furnished with two bracteas at the base. Petals 5, with a crenulate limb. Ovary 5-furrowed. Styles 5, unconnected. Stigmas capitate. Carpels 5, capsular, connected, 1-seeded. Seeds unknown. (*Don's Mill.*, i. p. 572.) A deciduous low tree, with large white flowers.

‡ I. MALACHODE'NDRON OVA'TUM Cav. The ovate-leaved Malachodendron.

Identification. Cav., 1. c.; Lindl. Bot. Reg.; Don's Mill., 1. p. 272.

Synonymes. Stua'rtia pentagfonia L'Hérit.; Stewa'rtia Malachodéndron Mill.; Stewartia à cinq

Styles, Fr.

Engravings. Smith's Exot. Bot., t. 101.; Michx. t. 58.; Bot. Reg., t. 1104.; and our fig. 91.

Spec. Char., &c. Leaves ovate, acuminate. Flowers axillary, solitary, almost sessile. Petals waved, cut, of a pale cream-colour. (*Don's Mill.*, i. p. 572.) A deciduous tree, attaining, in its native country, the height of 20 ft.; but, in England, generally seen as a bush, and seldom above 10 ft. or 12 ft. high. It is a native of Virginia, Carolina, and Georgia, on mountains; and it was introduced into England in 1795, producing its large white flowers in August and September. It has been, since the period of its introduction, in frequent cultivation among other American or peat-earth plants. The largest specimens which we have seen in England are at Dropmore and White Knights: the former are between 10 ft. and 12 ft. high; and form wide-spreading bushes, flowering freely every year. Their flowers are very large (2½ in. or more across), and slightly fragrant.* There are a great many trees nearly equally large at White Knights, which flower magnificently every year, and make a fine appearance during the months of August and September, when they are in full bloom. The plant would have a much better effect if trained up with a single stem, so as to form a small tree. For this purpose, after a plant has been two or three years established, it may be advisable to cut it down to the ground; and, from the shoots that it will throw up, to select one, and train it as the stem of the future tree. The soil in which it is generally grown is a mixture of loam and peat, in which the latter prevails; but, in the Mile End Nursery, it shoots vigorously, and flowers freely, in deep sandy loam. The situation



should be sheltered; and shaded rather than otherwise. The usual mode of propagation is by layers; and the stools are sometimes protected, during winter, by mats. Plants, in the London nurseries, cost 5s. each; at Bollwyller, 15 francs; and at New York, 50 cents.

GENUS II.



STUARTIA Cav. THE STUARTIA. *Lin. Syst. Monadélphia Pentádrin.*

Identification. Cav. Diss., 5. p. 393.; Dec. Prod., 1. p. 528., as *Stewartia*; Don's Mill., 1. p. 573.

Derivation. Named in honour of John Stewart, Marquess of Bute, the patron of Sir John Hill, and a distinguished promoter of botanical science.

General Character, &c. *Calyx* permanent, 5-cleft, rarely 5-parted, furnished with two bracteas at the base. *Petals* 5. *Ovary* roundish. *Style* 1, filiform, crowned by a capitate 5-lobed stigma. *Capsule* woody, 5-celled, 5-valved; cells 1—2-seeded. *Seeds* wingless, ovate, even. (*Don's Mill.*, 1. p. 573.)—A deciduous shrub, or low tree, with large white flowers.

1. STUARTIA VIRGINICA Cav. The Virginian Stuartia.

Identification. Dec. Prod., 1. p. 528.; Don's Mill., 1. p. 573.

Synonymes. *Stewartia Malachodéndron* *Lin. Sp.*, 982.; *Stuártia marilándica* *Bot. Rep.*; *Stewartia à un Style*, *Fr.*; eingriffliche (one-styled) *Stuartie*, *Ger.*

Engravings. Lam. Ill., t. 593.; Bot. Rep., t. 397.; and our fig. 92.

Spec. Char., &c. Flowers large, white, with purple filaments and blue anthers, usually in pairs. Leaves ovate, acute.

Petals entire. (*Don's Mill.*, 1. p. 573.) A deciduous shrub; from 6 ft. to 8 ft. high in Virginia, its native country, and attaining nearly the same height in British gardens. It is found in swamps in the lower parts both of Virginia and Carolina; and was introduced into England in 1742. The general appearance of the plant is the same as that of the preceding genus; but it forms a smaller bush, and the foliage has a redder hue. The flowers are of the same size, white, with purple filaments and blue anthers. This plant is not so extensively cultivated as the other, from its being of somewhat slower growth; but its beauty, and the circumstance of its flowering from July to September, when but few trees or shrubs are in blossom, render it desirable for every collection. It thrives best in a peat soil, kept moist; but it will also grow in deep moist sand. In this, as in similar cases, care should be taken that no rampant plant be placed near it, the roots of which might penetrate into the mass of peat or sand, and, from their greater vigour, soon occupy it, and destroy, or greatly injure, those of the *Stuártia*. The propagation of this plant is the same as that of *Malachodéndron*; from which it is separated on account of a technical difference in the capsules, somewhat similar to that by which *Thea* is separated from *Caméllia*. There are large plants of this species in the Mile End Nursery. The price is the same as that of *Malachodéndron*.



GENUS III.



GORDONIA Ellis. THE GORDONIA. *Lin. Syst. Monadélphia Polyándria.*

Identification. Ellis, in Phil. Trans., 1770.; Cav. Diss., 597.; Dec. Prod., 1. p. 528.; Don's Mill., 1. p. 573.

Derivation. Named in honour of *Alexander Gordon*, a celebrated nurseryman at Mile End, near London, who lived in the time of Philip Miller.

Gen. Char. Calyx of 5 rounded coriaceous sepals. Petals 5, somewhat adnate to the urceolus of the stamens. Style crowned by a peltate 5-lobed stigma. Capsules 5-celled, 5-valved; cells 2—4-seeded. Seeds ending in a leafy wing fixed to the central column, filiform. (*Don's Mill.*, i. p. 573.)—There are only two hardy species, both sub-evergreen.

† 1. GORDONIA LASIANTHUS L. The woolly-flowered Gordonia, or Loblolly Bay.

Identification. Lin. Mant., 570.; Dec. Prod., i. p. 528.; Don's Mill., i. p. 573.

Synonymics. *Hypéricum Lasianthus* Lin. Sp., 1101., *Catesb. Carol.*, i. t. 44., *Pluk. Amalth.*, t. 352.; *Gordonia à Feuilles glabres*, and *Alcée de la Floride*, Fr.; *langsfiehlige Gordonie*, Ger.

Engravings. Cav. Diss., 6. t. 171.; Sims, Bot. Mag., t. 668.; *Catesb. Carol.*, i. t. 44.; *Pluk. Amalth.*, t. 352.; and our fig. 93.

Spec. Char., &c. Pedicels axillary, usually shorter than the leaves. Leaves oblong, coriaceous, smooth, serrated. Calyx silky. Capsules conoid, acuminate. (*Don's Mill.*, i. p. 573.) A tree growing, in its native country, to the height of 50 ft. or 60 ft., with a diameter of 18 in. or 20 in.; and a straight trunk of from 25 ft. to 30 ft. "The small divergency of its branches near the trunk gives it a regularly pyramidal form; but, as they ascend, they spread more loosely, like those of other trees of the forest. The bark is very smooth, while the tree is less than 6 in. in diameter: on old trees it is thick, and deeply furrowed. The leaves are evergreen, from 3 in. to 6 in. long, alternate, oval-acuminate, slightly toothed, and smooth and shining on the upper surface. The flowers are more than 1 in. broad, white, and sweet-scented: they begin to appear about the middle of July, and continue blooming in succession during two or three months. This tree possesses the agreeable singularity of bearing flowers when it is only 3 ft. or 4 ft. high. The fruit is an oval capsule, divided into five compartments, each of which contains small, black, winged seeds." (*Sylva Americana*, p. 164, 165.) In England, the *Gordonia Lasianthus* is seldom seen otherwise than as a sub-evergreen bush, of 5 ft. or 6 ft. in height; but it flowers beautifully, even at that size. It sometimes, however, reaches the height of 10 ft. or 12 ft.

Geography, History, &c. The loblolly bay has a comparatively limited range in North America, being confined to the swamps near the sea coast, from the Floridas to Lower Louisiana. "In the pine-barrens, tracts of 50 or 100 acres are met with, at intervals, which, being lower than the adjacent ground, are kept constantly moist by the waters collected in them after the great rains. These spots are entirely covered with the loblolly bay, and are called bay swamps. Although the layer of vegetable mould is only 3 in. or 4 in. thick, and reposes upon a bed of barren sand, the vegetation of these trees is surprisingly luxuriant." (*Sylva Amer.*, p. 164.) This plant seems to have been first recorded by Catesby; and it was soon afterwards described by Ellis, in the *Philosophical Transactions*; and figured there, as well as in Catesby's *Carolina*. It was introduced into England, about 1768, by Benjamin Bewick, Esq.; but it has never been very successfully cultivated, apparently from neglecting to imitate its natural habitat, a swampy soil in a low sheltered situation. The largest plants in the neighbourhood of London are at Purser's Cross, and are not above 10 ft. high.

Properties and Uses. The wood of this tree, in America, is considered of little use; but its bark is of great value for tanning, for which purpose it is employed throughout the maritime parts of the southern states, and of the Floridas. A bark fit for the purpose of tanning is more valuable, in America, than might at first sight be imagined; because, though they have many sorts of oak, there are very few the bark of which contains a sufficient quantity of tannin to be worth employing by the tanner. Hence the Americans import

93



the bark and acorn cups of *Quercus Ægilops* from Spain; and these they mix with the bark of the loblolly bay. Michaux remarks that the bark may be taken off this tree during three months; which shows that in it the sap is abundant, and in vigorous motion, during a much longer period than it is in the oak. In the northern parts of America, such as New York and Philadelphia, the loblolly bay is planted in gardens, along with the magnolia, as an ornamental tree, requiring there, as it does in Paris, some slight protection during winter.

Soil, Situation, &c. This most beautifully flowering tree well deserves to have a suitable soil prepared for it, and to be treated with more care after it is planted than it appears to have hitherto received in England. The soil ought to be peat, or leaf-mould and sand; and it should be so circumstanced as always to be kept moist. For this purpose a considerable mass of soil ought to be brought together, and placed in an excavation, on a retentive substratum, in a low situation. During summer, water ought to be supplied from below, rather than from the surface, in order that the degree of moisture be kept as steady as possible; which it never can be when the surface is alternately moistened by the watering-pot, and dried by the sun. A steady moisture may be produced by laying in the bottom of the foundation either one or two brick drains across it, or a substratum of broken stones or gravel, to which water can be supplied through a shaft or tube communicating with the surface. Where both species of *Gordonia* are to be grown along with some other American trees and shrubs, such as *Magnolia glauca*, &c., which require similar treatment, the expense of this preparation would be well worth incurring, in order to insure the successful growth of the plants. In British nurseries, the *Gordonia* is generally propagated by layers, but sometimes seeds are imported. These require to be raised on peat soil, kept moist, and shaded; and for this purpose a covering of *Sphagnum* is thought desirable, as the seeds which drop from the plants in their native habitats, according to Michaux, only germinate successfully in this moss.

Statistics. There are specimens of this tree of 8 ft. or 10 ft. in height at Purser's Cross, at White Knights, in some of the London nurseries, and at a few other places in England; but scarcely any in Scotland, from the tree being rather tender; and but few in Ireland, because there the summers seldom admit of the wood being ripened. In the neighbourhood of Paris, there are some plants in the nurseries which have attained the height of 8 ft. or 10 ft.; and there was formerly a large one at Malmaison. Price, in the London nurseries, 5s.; at Bollwyller, ?; and at New York, 1 dollar.

Æ 2. *G. PUBESCENS Ph.* The pubescent *Gordonia*.

Identification. Pursh, *Flor. Amer.*, 2. p. 451; Dec. Prod., 1. p. 528; Don's Mill., 1. p. 573.

Synonyms. *Lacathea florida* Sal. Par. Lond., t. 56; *Franklinia americana* Marsh.; the Frank-linia, Amer.; behaarte Gordonie, Ger.

Engravings. Sal. Flor. Lond., t. 56; Michx., t. 59; and our fig. 94.

94

Spec. Char., &c. Flowers almost sessile. Leaves obovate-lanceolate, pubescent beneath, somewhat serrated, membranaceous. Petals and sepals rather silky on the outside. (*Don's Mill.*, i. p. 573.) A deciduous tree, from 20 ft. to 30 ft. high, producing large, white, fragrant flowers, with yellow anthers, in September and October. 1774.

Variety. — De Candolle indicates two forms: — *G. p. velutina*, *G. pubescens* L'Hér., figured in *Car. Diss.*, 6. t. 162., with oblong leaves velvety beneath, which may be considered the species; and *G. p. subglabra*, *G. Franklini* L'Hér., and *Franklinia Alatamaha* Marsh., with leaves smoothish beneath.

Description, &c. This species is much smaller than the preceding one; in its native country forming a deciduous tree, rarely exceeding 30 ft. in height, with a trunk 6 in. or 8 in. in diameter. The bark of the trunk is smooth, and presents a ridged surface, somewhat like that of the common hornbeam. The flowers are more than 1 in. in diameter, white, and of an agreeable odour. In Carolina they appear about the beginning of July; and a month later near Philadelphia. They open in succession during two or three months, and



begin to appear when the tree is only 3 ft. or 4 ft. high. In the neighbourhood of London the tree seldom comes into flower before September; and it continues flowering till its flower buds are destroyed by frost. It is rather hardier than the preceding species.

Geography, History, &c. This species is found only on the banks of the Alatamaha river in Georgia; where it was discovered, in 1770, by John Bartram, who gave it the name of *Franklinia*, in honour of the celebrated Dr. Franklin. Its native soil is sandy wastes, where there is peat, and where there is abundance of moisture great part of the year. This tree was introduced into England, in 1774, by Mr. William Malcolm. It is considered somewhat hardier than the preceding species, and has been more generally cultivated. The soil, situation, &c., may be considered, in all respects, the same as for *Gordonia Lasiánthus*. There are plants from 6 ft. to 8 ft. high in the Mile End Nursery, and of a larger size at Purser's Cross and at Syon; there are, also, some very fine bushes, or low trees, of it at White Knights, which flower freely every year. In the *Nouveau Du Hamel* it is stated to be cultivated in the "*Jardin Impérial des Plantes*," in those of Malmaison, and of the Trianon, and in Cels's Nursery. There is, or was a few years ago, a tree of considerable size in the garden at Trianon; and there is one in Bartram's Botanic Garden, Philadelphia (now Carr's Nursery), 50 ft. high. (See *Gard. Mag.*, vol. viii. p. 272.) Price, in the London nurseries, 3s. 6d. a plant; at Bollwyller, 4 francs; and at New York, 40 cents, and the seeds 2 dollars a quart.

Sect. II. *Hardy and half-hardy ligneous Plants belonging to the Tribe Camelliææ.*

Common Character. Calyx of 5—9 sepals. Petals 5—7—9, alternating with the sepals when they are the same in number: sometimes they are connected at the base. Stamens numerous, usually monadelphous, but, in some, separated into many bundles at the base. Capsule 3—5-celled, 3—5-valved, valves sometimes with dissepiments in the middle, and sometimes so much bent in at the margins as to form dissepiments. Seeds large, few, fixed to the margins of the central placenta. Smooth evergreen trees or shrubs, inhabitants of the colder parts of Asia, China, Japan, &c. Flowers axillary, very showy, red, white, or striped. (*Don's Mill.*, i. p. 574.) The half-hardy genera are two, *Camellia* and *Thsea*; which are thus contradistinguished:—

CAME[^]LLIA. Stamens polyadelphous or monadelphous at the base. Valve of capsule bearing a dissepiment in the middle of each.

THEA. Stamens almost unconnected to the very base. Dissepiments of capsule formed from the inflexed margins of the valves.

GENUS I.



CAME[^]LLIA L. THE CAMELLIA. *Lin. Syst.* Monadélphia Polyándria.

Identification. *Lin. Gen.*, No. 848.; *Dec. Prod.*, I. p. 529.; *Don's Mill.*, I. p. 574.

Synonyms. The Japan Rose; Camellier, Rose du Japon, et de la Chine, *Fr.*; Camellie, *Ger.*

Derivation. Named in honour of *George Joseph Camellus*, or *Kamel*, a Moravian Jesuit, and traveller in Asia.

Gen. Char., &c. *Calyx* imbricate, surrounded by accessory bractæ or sepals. *Stamens* monadelphous. *Anthers* elliptical, 2-celled, bursting lengthwise; capsule furrowed, with a dissepiment in the middle of each valve, separating from the free triquetrous axis when ripe. *Cells* 1—2-seeded. Elegant evergreen trees or shrubs, with coriaceous, dark green, shining leaves and large flowers, resembling the rose, of various hues. (*Don's Mill.*, i. p. 574.)—The species are evergreen low trees or shrubs, from China, all of which will bear the open air in the neighbourhood of London,

with some protection during winter; and a few of them, when once established, with no protection whatever. They are all readily propagated by layers, which, for the commoner sorts, are made from stools planted in cold-pits. They are also propagated by cuttings of the ripe wood, planted in autumn, and kept in a cool frame through the winter, being put into heat when they begin to grow in the spring. Grafting, inarching, and budding are employed for propagating the sorts that are comparatively rare. Various new sorts have been raised from seeds ripened in this country.

An excellent work has been published on the *Camelliæ*, by Messrs. Chandler and Booth, entitled *Illustrations and Descriptions of the Camelliæ*; in which many of the finest varieties are figured, and scientifically described. Another work, on the same genus, is in course of publication by the Messrs. Baumann, at Bollwyller.

■ 1. *C. JAPONICA* L. The Japanese Camellia, or the red single-flowered Camellia.

Identification. Lin. Sp., 982; Dec. Prod., l. p. 529; Don's Mill., l. p. 574.

Engravings. Cav. Diss., 6. t. 160; Jacq. Icon. rar., 3. t. 553; Duh. Ed. nov., t. 71; Bot. Mag., t. 42; Chandl. Ill., t. 1; Andr. Bot. Rep., t. 25; Lod. Bot. Cab., t. 329. and 455; Lois. Herb. Amat., t. 43, 44, 45, and 46; and our fig. 94.

Spec. Char., &c. Leaves ovate, acuminate, acutely serrated. Flowers axillary, sessile, usually solitary. Ovary smooth. (*Don's Mill.*, i. p. 574.) An evergreen shrub or low tree, perfectly hardy in the neighbourhood of London, as a standard in the free soil, when once thoroughly established.

Geography, History, &c. The *Camellia japonica* is indigenous both in China and Japan, and probably in other parts of Asia not yet explored by botanists. In Japan, it forms a lofty tree in high esteem with the Japanese for the elegance of its large flowers, which there exhibit a great variety of colours, and are produced from October to April. The trees are universally planted in the Japanese gardens; and, according to Thunberg, there are there several double-flowered varieties; and, among others, a double purple.

The *Camellia* was introduced into England by Lord Petre, about or before 1739. The first plants brought over were killed by being kept in a stove; but it was afterwards reimported, and kept in a conservatory. The plant began to come into general estimation in England about the beginning of the present century; and it has since been more extensively propagated than any other genus of green-house plants, unless we except *Pelargonium* and *Erica*. Within the last 15 or 20 years plants of this species have been tried in the open air, some against walls, and others as bushes; and, provided they are protected for a few years after planting, till the roots become firmly established in the soil, they seem to be nearly as hardy, even in the climate of London, as the common laurel. Sweet, who, it will be allowed, is a competent judge, says, "the *Camellia* is not generally so much cultivated as it deserves to be, though it is very hardy, standing out our severest winters when planted out against a wall, or in any sheltered situation, without protection." In Devonshire camellias form immense evergreen bushes without any protection whatever, and have even ripened seeds from which young plants have been raised. There is a single red camellia at Bicton, about $2\frac{1}{2}$ miles from the sea, and 83 ft. above its level, which is 9 ft. 6 in. high, with the head covering a space 12 ft. 6 in. in diameter. The soil and subsoil are chiefly sand, and the situation not particularly well sheltered. It has been planted out for 16 years, and not protected for the last 12 years. During the last five years, it has endured a temperature of 10° Fah., without sustaining the slightest injury. At Bicton there are a number of other varieties of *C. japonica* planted out as bushes. In the Vauxhall Nursery 8 sorts have stood out against a north-west wall for 8 years, and flowered freely every year without the slightest protection;



95

and in the Goldworth arboretum there are 30 or 40 sorts, species and varieties, which have stood out as bushes for several years, also without any protection, and in an elevated, open, unsheltered situation. We are not aware of the *Camellia japonica* having been tried as a standard in the open air in France or Germany; but at Naples, and more especially at Caserta, it has attained the height of 20 ft. in a very few years. (See *Gard. Mag.*, vol. xi. p. 151.) The price of the single-flowered variety, in the London nurseries, is 1s. 6d. a plant; at Bollwyller, 3 francs; and at New York, ?.

Varieties. A great many varieties have been raised in the neighbourhood of London, chiefly in the nursery of Messrs. Chandler and Son, and in the garden of the Messrs. Loddiges. Some of these may be considered as rather tender, but the greater part of them would answer against a north-west or north-east wall, if protected. The following is an enumeration of the principal varieties known in British gardens.

A. *Chinese Varieties in general Cultivation.*

- *C. j. 2 variegata* Bot. Rep. *The variegated-flowered Japanese Camellia.*—Figured in Lodd. *Bot. Cab.*, t. 329.; Chandl. *Ill.*, t. 6.; and *Bot. Rep.*, t. 91. It has flowers of a fine dark red, irregularly blotched with white. This is one of the hardiest of the varieties, and has stood out in several places for eight or ten years as an evergreen bush; flowering freely every spring, though sometimes having the flowers injured by frost. It has stood in the Vauxhall Nursery, without protection, for eight years. There are stools of it in the open ground in the Leyton Nursery, where it is propagated for sale in the same manner as the *Laúrus nóbilis*, and other hardy evergreen shrubs. There are stools of it in the Vauxhall Nursery, in cold-pits, from which plants are raised, and sold as hardy evergreen shrubs in the same manner as at Leyton. It was imported from China, by Captain Connor, for the late John Slater, Esq., in 1792. Price, in London, 3s. 6d. a plant; and at Bollwyller, 5 francs.
- *C. j. 3 incarnata* Bot. Reg. *The flesh-colour-flowered Japanese Camellia*, Lady Hume's Camellia, or Blush Camellia.—Figured in *Bot. Reg.*, t. 112.; and Chandl. *Ill.*, t. 7. This is generally considered the next hardiest variety to *C. j. variegata*. The flowers are of a fine delicate, and yet glowing, blush colour, becoming richer as they expand; the leaves are narrower and more acuminate than those of the preceding variety, and the tree has a looser and more slender habit of growth. Imported in 1806, for the late Lady Amelia Hume, of Wormeleybury, Herts. Price, in London, 3s. 6d. a plant; and at Bollwyller, 4 francs.
- *C. j. 4 álba pléna* Bot. Rep. *The white-double-flowered Japanese Camellia.*—Figured in Chandl. *Ill.*, t. 11.; Lodd. *Bot. Cab.*, t. 269. The flowers are of a pure white, from 3 in. to 4 in. in diameter. Plants of this variety, between 6 ft. and 8 ft. in height, have stood out as bushes in the Mile End Nursery, at Messrs. Loddiges's, in the Vauxhall Nursery, and at Purser's Cross, for several years. "One of the most elegant varieties in cultivation; brought to England, in 1792, by the same gentleman who introduced the double-striped; viz. John Slater of the India House, according to Messrs. Chandler and Booth; but Thomas Slater, according to Mr. Main, who went out as collector for Gilbert Slater ([*Gard. Mag.*], vol. ii. p. 423.) in 1791." (*Gard. Mag.*, vol. vi. p. 471.) Price, in London, 3s. 6d.; at Bollwyller, 4 francs. Beautiful imitations of the flowers of this variety have been formed in wax.
- *C. j. 5 fimbriata* Lodd. *The fringed-petaled white-double-flowered Japanese Camellia.*—Figured in Chandl. *Ill.*, t. 15.; and Lodd. *Bot. Cab.*, t. 1103. In cultivation since 1816, and a very beautiful variety. "Mr. Colvill, of the King's Road Nursery, has the merit of being

the first who brought it into notice, and no collection ought to be without it." (*Chandl. Ill.*, t. 15.) Price, in London, 5s.; and at Bollwyller, 15 francs.

j. 6 rubra plèna Bot. Rep. *The red-double-flowered Japanese Camellia*, Old red, and Greville's red.—Figured in *Bot. Rep.*, t. 199.; and in *Chandl. Ill.*, t. 18. The flowers are of a crimson-red colour, and resemble those of a double *Hibiscus Rosa sinensis*. Imported in 1794, by Sir Robert Preston, of Valleyfield, in Perthshire, and Woodfield, in Essex. Of a free and robust habit, and growing very erect; flowers but sparingly produced before the plant gets old; and hence this is not so much cultivated as some other varieties. The flowers are about 3 in. or 3½ in. in diameter, and open at the same time as those of the waratah and àtro-rubens. Price, in London, 3s. 6d. a plant; and at Bollwyller, 5 francs.

- *C. j. 7 anemoneflora*. *The Anemone-flowered Japanese Camellia*, Waratah Camellia, Blush Waratah Camellia.—Figured in *Bot. Mag.*, t. 1654.; *Chandl. Ill.*, t. 8.; and our *fig. 96*. Named Waratah from the resemblance of the flower to that of the *Telœpa speciosissima*, or waratah plant. This is one of the most singular, as well as the most beautiful, varieties: the flowers resemble those of a double poppy anemone (*Anemone coronaria*), having the exterior petals of the usual form, and the centre ones narrow and numerous; they are 3 in. or 4 in. in diameter, and of a deep and brilliant scarlet colour. Price, in London, 5s. each; and at Bollwyller, 10 francs.



96

- *C. j. 8 crassinervis* Lodd. *The thick-nerved-leaved Japanese Camellia*, Kent's Camellia, Kent's hexangular.—Figured in *Chandl. Ill.*, t. 39.; and Lodd. *Bot. Cab.*, t. 1475. Resembles the waratah, but differs in the outer petals being paler and more cupped, and in the leaves being thinner and rounder. Introduced in 1820. Price, in London, 7s. 6d.; and at Bollwyller, 15 francs.
- *C. j. 9 myrtifolia* Bot. Mag. *The Myrtle-leaved Japanese Camellia*.—Figured in *Bot. Mag.*, 1670.; and *Chandl. Ill.*, t. 14. The leaves are rather smaller than in most of the other varieties, and the flowers large in proportion to them, being about 3 in. in diameter. The plant is somewhat slender in growth, but flowers freely. Supposed to have been imported in 1808, for the Kew Garden. Price, in London, 5s.; and at Bollwyller, 6 francs.
- *C. j. 10 involuta* Bot. Reg. *The involute-petaled Japanese Camellia*, Lady Long's Camellia.—Figured in *Bot. Reg.*, t. 633. In general appearance resembling *C. j. myrtifolia*, but more erect and of stronger growth, and having petals involute, instead of spreading. Mr. Sweet deemed it identical with *C. myrtifolia*, as appears by his *Hort. Brit.*, p. 73. Introduced in 1820. In London, 7s.; and at Bollwyller, 25 francs.
- *C. j. 11 variabilis*. *The variable-coloured-flowered Japanese Camellia*.—More than four different-coloured flowers are produced upon the same plant; namely, red, white, and blush varieties of the peony-flowered and the pomponc.
- *C. j. 12 Pompônia* Bot. Reg. *The Pomponc Japanese Camellia*, the Kew Blush Camellia.—Figured in *Bot. Reg.*, t. 22.; and *Chandl. Ill.*, t. 9. The name appears to be derived from some fancied resemblance of the flowers to the French head-dress called a pomponc. The petals are delicate in their texture, and, when fully expanded, the flowers are just 4 in. across. The colour of the petals is pure white, except for about a third of their length nearest the base, which is deeply tinged with red, of which there is a small stripe up the centre. Introduced in 1810. This variety is very hardy; plants

of it have stood out for eight winters in the Vauxhall Nursery. Price, in London, 3s. 6d.; and at Bollwyller, 4 francs. *C. j. pæoniæflora rosea*, figured in *Chandl. Ill.*, t. 19.; *C. j. p. pallida*, and *C. j. p. alba*; the red peony-flowered, introduced in 1810; the blush pæony-flowered, introduced in 1820; and the white pæony-flowered, introduced in 1810, may be considered as subvarieties of the pompone. The price of these subvarieties is somewhat higher than that of *C. j. Pomponia*.

- ✱ *C. j. 13 semiduplex* Bot. Rep. *The semidouble-flowered Japanese Camellia*.—Figured in *Bot. Rep.*, t. 559. The flowers consist of from 6 to 12 large roundish petals, in a single or double series, with a column of stamens in the centre; they are of a rich rose colour. Introduced in 1808. Middlemist's red camellia so closely resembles this sort as hardly to be distinguishable from it.
- ✱ *C. j. 14 atro-rubens* Bot. Cab. *The dark red-flowered Japanese Camellia*, Loddiges's red Camellia.—Figured in *Lodd. Bot. Cab.*, t. 170.; and *Chandl. Ill.*, t. 25. This is a very beautiful variety, and a vigorously growing one. It is always among the latest in coming into blossom, the flowers opening at about the same time as those of the waratah camellia. The plants have an erect fastigate habit, and the flowers are of a deep scarlet, and are very showy. They are not so large as in some of the other varieties; but they have a marked appearance from the middle of the flower being filled with small petals, which project so as to overtop the others. Imported by Messrs. Loddiges, from China, in 1809. Price, in London, 5s.; and at Bollwyller, 5 francs.
- ✱ *C. j. 15 Welbankii* *Chandl. Ill.*, t. 27.; *Welbank's white-flowered Japanese Camellia*. *C. j. luteo-âlbicans* *Bot. Reg.*, t. 708.; *C. j. flavescens*; and white Moutan Camellia.—Figured in *Bot. Reg.*, t. 708. The flowers, which have been compared to those of *Gardènia florida*, are of a yellowish-white colour, rather delicate, and they do not open very freely. The flowers are from 3 in. to 3½ in. in diameter. In Messrs. Chandler and Booth's account of this variety, published Feb., 1831, it is characterised as very different from any other white-flowered kind, and as being of robust habit, and remarkable for the convexity of its foliage. Introduced by Captains Welbank and Rawes in 1820. Price, in London, 5s.; and at Bollwyller, 10 francs.
- ✱ *C. j. 16 rosea* Don's Mill. *The rosy-flowered Japanese Camellia*.—The flower has the appearance of a small moss rose; it measures about 2½ in. in diameter, approaching in form that of the myrtle-leaved. Introduced in 1821. Price, in London, 3s. 6d. each; and at Bollwyller, 5 francs.
- ✱ *C. j. 17 speciosa* Hort. Trans. *The showy Japanese Camellia*, Rawes's variegated Waratah Camellia. *Camellia Rawesiana* Hort.—Figured in *Chandl. Ill.*, t. 32. An extremely handsome variety, with flowers of a deeper red than either those of *C. atro-rubens* or the waratah. They open very regularly, and, when expanded, are usually 4 in. in diameter. Nearly the whole of the petals have a little white stripe at their base, and some of them are variegated; all of them are disposed in the same manner as those of the waratah. It was imported by Captain Richard Rawes in 1824, who presented it to his relation, T. C. Palmer, Esq., Bromley, Kent. Price, in London, 10s. 6d.; and at Bollwyller, 50 francs.
- ✱ *C. j. 18 carnea* Bot. Reg. *The flesh-coloured-flowered Japanese Camellia*, Middlemist's red Camellia, rose-coloured Camellia.—Figured in *Bot. Reg.*, t. 22. The flowers are similar in colour to those of the semidouble red (No. 13.), but larger. The centre petals are short and vary in form; generally they are roundish and a little twisted, as well as marked with dark-coloured veins, and all of them have, more or less, a small white-coloured stripe down their centre. The stamens are generally all changed into petals, and the whole flower resembles a full-blown rose. Introduced in 1808.
- ✱ *C. j. 19 imbricata* Hort. Trans. *The imbricated-petaled Japanese Camellia*, crimson Shell Camellia.—Figured in *Chandl. Ill.*, t. 22.; and *Bot. Reg.*, t. 1398. "Without doubt, the best variety that has

been brought from China. The flowers are upwards of $3\frac{1}{2}$ in. in diameter, and are very regular in form, the petals being arranged one above another, and gradually diminishing in size towards the centre, exactly in the manner of the double white. The colour is of a fine crimson red, and remarkably showy. When the flowers first begin to open they are concave, but, as they expand, they become quite flat. The outer petals are nearly round. The centre petals are rather pointed, and rise upright." (*Don's Mill.*, i. p. 576.) Introduced in 1824. Price, in London, 7s. 6d. each.

- *C. j.* 20 *Parksii Hort. Trans.* Parks's Japanese Camellia, Parks's striped-Rose Camellia. — The flowers are of a bright rose colour, 4 in. in diameter, irregularly striped or blotched with white, and they are slightly odoriferous, like the flowers of the myrtle-leaved variety. In this and other respects, it differs from *C. j.* variegata (No. 2.). Introduced in 1824. Price, in London, 10s. 6d. each.
- *C. j.* 21 *Sabiniæna Hort. Trans.* Sabine's Japanese Camellia, Sabine's white Camellia. — The flowers are of a pure white, 3 in. across, and they resemble in form those of the pomponne. Introduced in 1824.

B. *Chinese and other Foreign Varieties not in general Cultivation, but in all probability as hardly as the others.*

- *C. j.* 22 *canadissima* is noticed, in *G. M.*, vol. xi. p. 78. and 190., as one that assimilates to *C. j.* *Wellbänkii* and as brought directly from Japan by Dr. Siebold, and called by some *C. Sieboldii*. It is deemed by some a species. It is in Mr. Knight's collection.
- *C. j.* 23 *Donklæri*. — It is said that this is a very fine variety. It was raised on the Continent, and is named after the head gardener at the botanic garden at Louvain. (*Gard. Mag.*, vol. xi. p. 85.) Mr. Knight possesses it in a living state. (p. 191.)
- *C. j.* 24 *francofurtensis*. — Flowers dark and light red, quite as large as those of *C. reticulata*. Raised from seeds of *C. argentea*, by M. J. Rinz, jun., nurseryman, Frankfort on the Maine; who deems it the finest variety that has ever been seen in Germany. It flowered for the first time in 1834. Mr. Low, at Clapton, has a plant of it. (*Gard. Mag.*, vol. xi. p. 265. 543.)
- *C. j.* 25 *kybrida* Makoy. — M. Jacob Makoy sent us, previously to March, 1835, a dried specimen of this kind of camellia, which he informed us was a hybrid from *C. japonica* var. *insignis* and *C. euryoides*. By the specimen, it assimilates in habit to *euryoides*: the leaves are ovate, acuminate, serrate, and slightly puberulous; the sprig and buds densely puberulous; the flower 1 in. across, perhaps more; and the petals 5, orbicular, and centred by the cluster of stamens. The petals, in a dried state, were of a buff colour; they might be, when living, white, tinted with red: nothing was stated of their colour when living. M. Makoy deemed the hybrid a fine variety. (*Gard. Mag.*, vol. xi. p. 143.)

Other names of Foreign Varieties of *C. japonica*. In *Gard. Mag.*, vol. xi., varieties of camellia, by the following names, are mentioned as extant on the Continent, which, it seems, are not yet so in Britain: *C. argentea*, in p. 265.; *C. Gunnêli*, in p. 543.; *C. Pronayana*, in p. 544.; and *C. violæcea superba*, in p. 544.

The *semidouble white* was purchased in 1822, on the Continent, by Mr. Palmer, and is considered a distinct variety; but it has not yet flowered in England. The *rose-coloured waratah* was introduced, by the London Horticultural Society, from China, in 1824; but it has not yet flowered. *C. j.* *hexangulâris*, the hexangular-flowered Japanese Camellia, is a very singular variety, only known by the Chinese drawings in the possession of the London Horticultural Society, it not having been yet introduced. There are, doubtless, other varieties in China; but, from our increased intercourse with that country, in consequence of the trade being thrown open, there can be no doubt that they will all, sooner or later, find their way into Britain.

C. *Varieties of Camellia japonica originated in Britain.*

The varieties of the common camellia originated in Britain are exceedingly numerous. The first seeds ripened were those of *C. j.* *anemoneflora*, about the year 1818, in the Count de Vandes's garden at Bayswater; and, subsequently, a great number of varieties have been raised by Messrs. Lodiges; Messrs. Chandler, of the Vauxhall Road Nursery; Mr. Press, gardener to Edward Gray, Esq., at Haringay, Hornsey; and various other nurserymen and gardeners. In Sweet's *Hortus Britannicus*, 2d edit., published in 1830, sixty-five sorts of camellias are enumerated; of which upwards of fifty are varieties of *C. japonica*. The following selection of these is taken from the *Illustrations* of Booth and Chandler, already mentioned; from the *Gardener's Magazine*, and from *Don's Miller*.

a. *Varieties raised in Britain that are figured and described in Chandler and Booth's Illustrations of the Camellieæ.*

- *C. j.* 26 *corallina* Chandl. Ill. *The coral-coloured-flowered J. C.* — Figured in Chandl. Ill., t. 10., and Chandler's *Camellieæ Britannicæ*, t. 5. A fine variety; its habit, like that of the waratah. Originated in 1819. Raised from seed by Messrs. Chandler of the Vauxhall Nursery. Price, in London, 7s. 6d.; and at Bollwyller, 10 francs.

- *C. j.* 27 *extima* Chandl. III. *The choice J. C.*, Chandler's choice-flowering Camellia.—Figured in Chandler's *Ill.*, t. 12; described there; whence the following traits are drawn. Of free growth, resembling, in some respects, the waratah. Flowers of a deep rose colour, very double, and 4 in. across. It also resembles *C. j.* *imbricata*; but the foliage of the two is totally distinct. Price, in London, 7s. 6d.
- *C. j.* 28 *Wiltoni* Chandl. *Ill.* *Lady Wilton's J. C.*—Figured in Chandl. *Ill.*, t. 13., and described there; where it is stated that it is a desirable well-marked variety, and distinct from either the double-striped, Parks's rose-striped, or the *C. j.* *Elphinstoni*. Flowers 2 in. or 2½ in. across, similar to those of the double-striped. Raised from seed by Mr. Knight of the Exotic Nursery, about 1814, probably from the single red, impregnated with the double-striped. Price, in London, 7s. 6d.; and at Bollwyller, 50 francs.
- *C. j.* 29 *Chandleri* Chandl. *Ill.*, *synon.* "versicolor *Bot. Reg.*, t. 887." (*Swt.*, in his *Hort. Brit.* p. 74.) *Chandler's J. C.*, Chandler's striped waratah *C.*—Figured in Chandl. *Ill.*, t. 10., Chandl. *Cam. Brit.*, t. 1 and 2., and *Bot. Reg.*, t. 887. In Chandl. *Ill.* it is stated of it, that it approaches nearer to the waratah than to any other variety, and is one of the very best which has been raised; and that it was originated by Mr. Chandler, in the Vauxhall Nursery, in 1819, from the seeds of the waratah, crossed with the double-striped. Price, in London, 7s. 6d.; and at Bollwyller, 20 francs.
- *C. j.* 30 *flore albo* Chandl. *Ill.* *The white single-flowered J. C.*—Figured in Chandl. *Ill.*, t. 17., and *Bot. Reg.*, t. 353. In Chandl. *Ill.* it is stated that it is more robust than almost any other variety. It abounds in flowers, and these usually open earlier than those of most other varieties; they are nearly 3 in. across, white, and not infrequently striped or spotted with red. It seeds freely, and some fine varieties, with double flowers, of different colours, have been raised from it. It was raised from seed of the double-striped, by Messrs. Rolleston of the Tooting Nursery, in about 1814.
- *C. j.* 31 *althæiflora* Chandl. *Ill.* *The Hollyhock-flowered J. C.*—Figured in Chandl. *Ill.*, t. 20., and Chandl. *Cam. Brit.*, t. 4.—Flowers red. It is stated that it resembles the waratah in some respects, and is well deserving of a place in every collection. Raised by Mr. Chandler, from seed of the waratah, in 1819. Price, in London, 7s. 6d.; and at Bollwyller, 15 francs.
- *C. j.* 32 *anemoneflora alba* Chandl. *Ill.* *The Anemone-flowered white-flowered J. C.*, the white Waratah *C.*—Figured in Chandl. *Ill.*, t. 21., and Chandl. *Cam. Brit.*, t. 8. This, from the account of it in Chandl. *Ill.*, is a most elegant variety, with blossoms of a delicate white colour, a little striped; occasionally spotted with pale red, from 3½ in. to 4 in. over. It was raised from the pomphone. Price, in London, 7s. 6d.; and at Bollwyller, 15 francs.
- *C. j.* 33 *Woodsii* Chandl. *Woods's J. C.*—Figured in Chandl. *Ill.*, t. 25. A seedling, raised by Mr. Chandler, who named it in compliment to Mr. Woods of Camberwell Grove, a great admirer of camellias. Its flowers are very handsome, symmetrically formed, and nearly 4 in. across. Their colour is pale red, similar to those of the common Provence rose, which, at a distance, they resemble, except in being larger, although not so double. (*Chandl. Ill.*, adapted.) Price, in London, 7s. 6d.; and at Bollwyller 50 francs.
- *C. j.* 34 *punctata* Chandl. *Ill.* *The dotted-flowered J. C.*, Gray's invincible *C.*—Figured in Chandl. *Ill.*, t. 24, whence the following particulars are derived. The expanded blossoms are from 3 in. to 4 in. over; of a very delicate bluish colour, almost white; striped, and slightly spotted with pale rose, in the manner of a rose-flake carnation. An extremely fine variety. It was raised, in 1824, by Mr. Press, gardener to E. Gray, Esq., from a seed of the semidouble red, the flowers of which had been fecundated with the pollen of the single white.
- *C. j.* 35 *elegans* Chandl. *Ill.* *Chandler's elegant J. C.*—Figured in Chandl. *Ill.*, t. 26. Of free growth; the flowers of a very delicate rose colour, and from 3½ in. to 4 in. across; in form between those of the waratah and those of the pæony-flowered, but, in other respects, distinct from those of both varieties. Raised from seed of the waratah, by Mr. Chandler, about 1822.
- *C. j.* 36 *florida* Chandl. *Ill.* *The flowery J. C.*, the cluster-flowering *C.*—Figured in Chandl. *Ill.*, t. 28., and in Chandl. *Cam. Brit.*, t. 7. The flowers are not large, but very beautiful; 3 in. in diameter; of a deep rose colour, intermediate between the deep red of the waratah and the bright rose of the pæony-flowered. They are produced in great abundance at the extremity of the branches, and open pretty early in the season. This variety was produced in 1819, from seed of the waratah, from a flower that had been fertilised with the pollen of the pæony-flowered. Price, in London, 7s. 6d.; and at Bollwyller, 15 francs.
- *C. j.* 37 *rosea* Chandl. *Ill.* *Middlemist's rosy-coloured-flowered J. C.*, Middlemist's red *C.*—Figured in Chandl. *Ill.*, t. 29. The flowers open late, and are of a rich rose colour; more than semidouble, and 3 in. or more across. The stamens are sometimes perfect, but oftener transformed into small petals; so that the flower altogether resembles that of a full-blown rose.
- *C. j.* 38 *eclipsis* Chandl. *Ill.* *Press's eclipse J. C.*—Figured in Chandl. *Ill.*, t. 50. The following is abridged from the description of it in Chandl. *Ill.* The flowers are handsome and well-formed, the petals being numerous, and neatly imbricated; the ground colour white, which is striped and feathered with pale red, in the manner of a flake carnation. One of the very fine varieties originated by Mr. Press, and noticed in *Gard. Mag.*, vol. ii. p. 358. Both *C. j.* *eclipsis* and *C. j.* *punctata* were raised from seeds contained in one capsule. Price, in London, 7s. 6d.; and at Bollwyller, 100 francs.
- *C. j.* 39 *insignis* Chandl. *Ill.*, *syn. C. j.* *dianthiflora Hort. Brit.*, p. 592. *The remarkable J. C.*, Chandler's splendid *C.*, the Carnation Waratah *C.*—Figured in Chandl. *Ill.*, t. 31., and Chandl. *Cam. Brit.*, t. 6. It is stated that this is a favourite variety with most cultivators, and that there are few collections in which it does not hold a conspicuous place. Its flowers are large and red. Price, in London, 7s. 6d.; and at Bollwyller, 20 francs.
- *C. j.* 40 *alba semi-duplex* Chandl. *Ill.*, t. 33. *Palmer's white-semidouble-flowered J. C.*—Figured in Chandl. *Ill.*, t. 33., and described there; whence it appears that the flowers average more than 4 in. in breadth, and consist of 8 or more petals, disposed in 2 or more rows; the largest round, and about 1½ in. across; the others roundish, oblong, and a little smaller; all are extremely delicate in texture, like those of Welbank's white. Price, in London, 10s. 6d.
- *C. j.* 41 *concinna* Chandl. *Ill.* *The neat J. C.*—Figured in Chandl. *Ill.*, t. 51. The flowers open

well, are very regularly formed, and of a fine rose colour; they exceed 3 in. in diameter, and are little inferior in appearance to those of *C. j. eximia*, or of *C. j. imbricata*; the petals being nearly as numerous, and arranged with equal symmetry. Raised from seeds of the warratah, by Mr. Chandler, in 1819. It possesses much beauty; but its flowers are less brilliant than those of some others: it is not so well known as it should be. (*Ill.*, t. 34.) Price, in London, 10s. 6d.

- *C. j. 42 splendens*, Chandl. *Ill.*, synon. *C. j. cœcinea Hort. Brit.*, p. 593. Allnutt's *splendens J. C.* — Figured in Chandl. *Ill.*, t. 35. A much admired and most desirable variety. The flowers are of a brilliant red, 3 in. broad, very showy, and produced abundantly on both young plants and old ones. The petals are all deeply veined, and, though less numerous than in some varieties, are so arranged in the centre as to form flowers to all appearance perfectly double. The petals, also, are so remarkable for their roundness as to give the flowers a peculiar character; by which the variety may be readily distinguished. The plant, in habit, is similar to the single red *C. japonica L.*, but is stronger and more bushy; the branches are upright and twiggy. Price, in London, 5s.
- *C. j. 43 Rōsa sinensis* Lodd. *Bot. Cab. The Chinese Rose (? Hibiscus flowered J. C.* — Figured in Lodd. *Bot. Cab.*, t. 1175, and Chandl. *Ill.*, t. 36. This is a bold-flowering, freely blooming, first-rate variety. Its pale purplish red flowers are tolerably full of petals, extremely handsome, 4 in. across, and bearing considerable resemblance to those of *C. j. elegans*. It was raised by Mr. Chandler about 1819. Price, in London, 7s. 6d.; and at Bollwyller, 15 francs.
- *C. j. 44 Bissin* Chandl. *Ill.*, t. 37. *Ross's J. C.* — Figured in Chandl. *Ill.*, t. 37., and described in that work. This is a desirable variety; it is briefly described in *Gard. Mag.*, vol. i. p. 211., under the name of Ross's *Camellia gloriosa*. The flowers are often 1 in. in diameter; in form they resemble those of *C. j. elegans*; but in colour they are of a much darker and deeper red.
- *C. j. 45 Aiton's* Chandl. *Cum. Brit.* *Aiton's J. C.* Aiton's large single red *C.* — Figured in Chandl. *Ill.*, t. 38., and in Chandl. *Cum. Brit.*, t. 3. It is a very striking variety, and a most valuable one to the cultivator of camellias, on account of its producing seeds more freely than any other kind whatever. This and four others were raised at the Vauxhall Nursery, from seeds contained in one capsule of the pomponne camellia, and sown in Nov. 1819. It was named in compliment to Mr. Aiton, the king's gardener. (*Ill.*) Price, in London, 7s. 6d.; and at Bollwyller, 50 francs.
- *C. j. 46 epsomensis* Chandl. *Ill.* *The Epsom J. C.*, Young's semidouble red *C.* — Figured in Chandl. *Ill.*, t. 41. Raised by Mr. Young of Epsom, previously to 1824. It is robust in habit; its flowers are much like those of the semidouble red, but of a deeper colour, and with more petals. It is prone to vary in the number of petals; when they are numerous, some of the stamens are transformed into small, roundish, spatulate, striped, red petals, all of which, as well as the large outer ones, are tolerably evenly arranged over one another, and distinctly marked with dark-coloured veins. The usual colour of the flowers is a deep red, approaching to scarlet, and their width about 3 in. (*Ill.*)

b. Varieties raised in Britain that are figured and described in British Works, exclusive of those figured and described in Chandler and Booth's Illustrations of the Camellieæ.

- *C. j. 47 Colvill's* Sweet. *Br. Fl.-Gard.*, 2 ser. *Colvill's J. C.* — Figured in *Sweet. Br. Fl.-Gard.*, 2 s. t. 2., and described there; whence it appears that its petals are striped like those of the carnation, and that, when Mr. Sweet wrote the description, published in June, 1829, he deemed it to be the finest and most beautiful variety that he had seen; this was previous to the flowering of *C. j. Sweetiana*. G. Don has described *C. j. Colvillia*, as "an elegant hybrid, with the petals regularly disposed, blotched with white on a red ground." *Don's Mill.*, i. p. 376. Price, in London, 10s. 6d.
- *C. j. 48 Sweetiana* Sweet. *Br. Fl.-Gard.*, 2 ser. *Sweet's J. C.*, Sweet's painted-flowered *C.* — Figured in *Sweet. Br. Fl.-Gard.*, 2 s. t. 133., and in March, 1832. The flower bears a strong resemblance to a beautiful variegated rose: it is generally very double, more spreading than that of many varieties, and elegantly marked and variegated with white, blush, and deep rosy red. It was the finest variety Mr. Sweet had seen, and one of numerous hybrid varieties that he had fertilised and raised from seeds, several years before (perhaps about 1824, in the nursery of Mr. Colvill. This was the offspring of the double-striped, fertilised by the pomponne; the foliage resembles most that of the latter, but the leaves are larger; and the plant, if not seen in flower, might be mistaken for a strong-growing single-flowered one. Sweet's camellia assimilates with Gray's *invincibile*; but its flower bud is larger, and its flower larger, and of a deeper colour, than those of that variety.
- *C. j. 49 Knight's* Lodd. *Bot. Cab. Knight's J. C.* — Figured in Lodd. *Bot. Cab.*, t. 1163. It is stated in Chandl. *Ill.*, t. 31., that this approximates closely to *C. j. insignis*, and that it was raised by Mr. Knight of the King's Road, Chelsea.

c. Varieties raised in Britain, of which some Description has been published.

- *C. j. 50 Rōsa mundi* Don's Mill. *The Rose of the World J. C.* — Flowers white and crimson. (*Don's Mill.*, i. p. 376.) Price, in London, 5s.; at Bollwyller, 50 francs.
- *C. j. 51 Pressii* Don's Mill. *Pressii's J. C.* — Flowers single red. (*Don's Mill.*, i. p. 376.)
- *C. j. 52 rubra-punctata* Don's Mill. *The red-spotted-flowered J. C.* — Flowers single, white, spotted with red. (*Don's Mill.*, i. p. 376.)
- *C. j. 53 Elphinstonia* Miss Elphinstone's *J. C.* — Flowers red. Raised by Mr. Knight. (*Sweet's II. B.*, p. 71.) Assimilates to *C. j. Wiltoni*. Chandl. *Ill.*, t. 13.; Price, in London, 7s. 6d.
- *C. j. 54 single-striped and dotted* Burnard, in *Gard. Mag.*, vol. ii. p. 338., has described its flowers as having a clear white ground, with pink stripes, and dotted all over with small dots: they are large and beautiful; and the variety was raised by Mr. Press, along with the varieties *punctata*, *Rōsa mundi*, *Pressii*, and *elopsis*, from seeds saved from a plant of the semidouble red, the flowers of which had been fecundated with pollen of the single white

- *C. j. 55 aeneæfolia* Loudon's H. B. *The Aucuba-leaved J. C.*— Splendid; its flowers red, and produced from February to May. Flowers of it were exhibited at a meeting of the London Horticultural Society, on March 3. 1835, from the Society's Garden. (*G. M.*, vol. xi. p. 216.)
- *C. j. 56 expansa* Loudon's H. B. *The expanded-flowered J. C.*— Described to be splendid; its flowers red, and produced from February to May. *C. j. Susanna*, *C. j. Märtha*, and *C. j. Wadieana* were raised from seeds of this variety. (*G. M.*, vol. xi. p. 294.)
- *C. j. 57 Susanna Gard. Mag.*, vol. xi. p. 294. Miss *Susanna Thomson's J. C.*— The flower assimilates to that of *C. j. Sweetiana*, and some have thought that it equals or surpasses it in merit. The petals have a white ground, with pink stripes, in the manner of those of the carnation, but fewer and fainter: the centre of the flower is pretty well filled with petals. The flowers are produced in plenty. It was raised in Thomson's Nursery, Mile End, from seed of the variety termed *C. expansa*, produced in 1827, and sown in 1828.
- *C. j. 58 Märtha Gard. Mag.*, vol. xi. p. 294. *Martha Poole's J. C.*— Its flower assimilates to that of *C. j. Colvilli*. The mode of its formation is somewhat that of the waratah. The petals have a blush ground and pink stripes; the centre of the flower is filled with petals. The flowers are produced in plenty. The foliage is fine. Named after Mrs. Poole, formerly *Martha Thomson*.
- *C. j. 59 Wadieana*.—Petals of a dead white colour. The flower large; formed differently from either the flowers of *C. j. alba plena*, or *C. j. fimbriata*: the petals are larger and less compactly disposed; though the centre is filled. The flowers stand long on the plant. The leaves are large and healthy. It was raised from the same stock of seeds as *C. j. Susanna*. Named after Mr. Wadie, propagator in Thomson's Nursery, Mile End.

d. *Names of Varieties of Camellia japonica that are mentioned in Gard. Mag., vol. xi., but without any Description of them being given.*

C. Cliveana, *conspicua*, *deebra*, *princeps*, *rotundifolia*, *Palmieri*, *Reevésii*, *longifolia*, are mentioned in p. 215. In p. 216, *C. Dorsètii*, *péndula*. In p. 326, *C. Allnütta alba*, and *superba*. Eight hybrid camellias, raised in the gardens of W. F. Campbell, Esq., M. P., Woodhall, Lanarkshire, the names of which are not given, are mentioned in *Gard. Mag.*, vol. xi. p. 295.

D. *Varieties of Camellia japonica included in the foregoing Lists, but placed here in the Order of their Hardiness in the Vauxhall Nursery, with the Retail Prices of Messrs. Chandler in 1836, for Plants of the smallest Size, affixed to each.*

a. *Varieties of C. japonica planted out against a North-west Wall, and which have grown and flowered well without any Protection, for Eight Years.*

<i>Camellia japonica</i> , or	<i>variegata</i> , 3s. 6d.	<i>præoniciflora</i> , 3s. 6d.
single red, 3s. 6d.	<i>røsea</i> , 3s. 6d.	<i>incarnata</i> , 3s. 6d.
<i>alba plena</i> , 3s. 6d.	<i>Pomponia</i> , 3s. 6d.	<i>variabilis</i> , 3s. 6d.

b. *Varieties of C. japonica considered the finest and most desirable, of all of which there are in the Vauxhall Nursery Stools planted in Cold-pits, and protected in severe Weather with only the Lights. The Names in the List are placed according to the Hardiness of the Sorts, and their Vigour of Growth.*

<i>rubra plena</i> , 3s. 6d.	<i>dianthiflora</i> , 5s.	<i>myrtifolia</i> , 5s.
<i>atro-rubens</i> , 5s.	<i>alba semi-duplex</i> ,	<i>fimbriata</i> , 5s.
<i>Welbankiana</i> , 5s.	10s. 6d.	<i>Elphinstoniaana</i> , 7s. 6d.
<i>corallina</i> , 7s. 6d.	<i>Sabini</i> , 10s. 6d.	<i>Parksii</i> , 10s. 6d.
<i>imbricata</i> , 7s. 6d.	<i>Aitonii</i> , 7s. 6d.	<i>compacta</i> , 7s. 6d.
<i>Chandleri</i> , 7s. 6d.	<i>Røsa sinensis</i> , 7s. 6d.	<i>Rössii</i> , 7s. 6d.
<i>Colvilli</i> , 10s. 6d.	<i>anemoneflora</i> , 5s.	<i>delicatissima</i> , 21s.
<i>élegans</i> , 10s. 6d.	<i>Røsa mundi</i> , 5s.	<i>Gilesii</i> , 31s. 6d.
<i>eximia</i> , 7s. 6d.	<i>eclypsis</i> , 7s. 6d.	<i>triumpfans</i> , 42s.
<i>speciosa</i> , 10s. 6d.	<i>punctata</i> , 7s. 6d.	<i>althææflora</i> , 7s. 6d.
<i>anemoneflora alba</i> ,	<i>spléndens</i> , 5s.	<i>spofforthiana</i> , 21s.
7s. 6d.	<i>Wiltoni</i> , 7s. 6d.	<i>crassinervis</i> , 7s. 6d.
<i>florida</i> , 7s. 6d.	<i>concinna</i> , 10s. 6d.	<i>Le Blanc's red</i> , or <i>røsea</i> ,
<i>insignis</i> , 7s. 6d.	<i>Woodsii</i> , 7s. 6d.	10s. 6d.

■ 2. *C. RETICULATA* Lindl. *The reticulated-leaved Camellia, or Captain Rawes's Camellia.*

Identification. Lindl. Bot. Reg., t. 1078.; Hook. Bot. Mag., t. 2784.; Don's Mill., I. p. 576. *Engravings.* Lindl. Bot. Reg., 1078.; Hook. Bot. Mag., t. 2784.; Chandl. Ill., t. 4.

Spec. Char., &c. Leaves oblong, acuminate, serrated, flat, reticulated. Flowers axillary, solitary. Calyx 5-sepaled, coloured. Ovary silky. (*Don's*

Mill., i. p. 576.) Introduced from China in 1824, and still scarce and high-priced. The plant is of vigorous growth, and appears as hardy as any of the other species and varieties. It is generally propagated by inarching on the common species. It appears to flower rather later than *C. japonica*; and, when it becomes more frequent, it will probably, on that account, be found well adapted for the open air, or a conservative wall. Messrs. Chandler and Booth express themselves of opinion that "it will be found to be hardier than the *C. japonica*, and that at no distant period, perhaps, it may ornament our shrubberies." There are stools of it in a cold-pit, in the Vauxhall Nursery, where plants cost a guinea and a half each.

■ 3. *C. MALIFLORA* Lindl. The Apple-blossom-flowered Camellia.

Identification. Lindl. Bot. Reg., 1078, in a note; Don's Mill., I. p. 576.

Synonymes. *C. Sasánqua* of Bot. Mag., t. 2080., Bot. Reg., t. 547., and Bot. Cab., t. 1134.; *C. Sasánqua rosea* Hort.; Palmer's double *Sasanqua*.

Engravings. Bot. Reg., t. 1078.; Chaudl. Ill., t. 2.; and, as *C. Sasánqua*, Bot. Mag., t. 2080.; Bot. Reg., t. 547.; Bot. Cab., 1134.; and our fig. 97.

Spec. Char., &c. Leaves obovate, convex, bluntly serrated. Flowers terminal and axillary, usually solitary.

Branches and petioles pubescent. Ovary smooth. (*Don's Mill.*, i. p. 576.)

Introduced from China, in 1816, by Captain Richard Rawes, into the garden of T. C. Palmer, Esq., at Bromley, in Kent, where it flowered in 1818, and was afterwards published in the *Botanical Magazine* as a variety of *C. Sasánqua*.

An elegant shrub, with a slender fastigate habit of growth, and a very free flowerer. It seldom exceeds 8 ft. in height; but, in the flowering season, its numerous semidouble apple-blossom-like flowers are extremely beautiful.

The leaves are obovate, and thinner, and of a smaller size, than those of *C. japonica*, or any of its varieties. It is propagated by inarching on the single red, or by layers.

Though this species will stand against a wall with very little protection, yet, to grow it properly, and to flower it in the best manner, it requires a little more heat than *C. japonica* and its varieties. In the Vauxhall Nursery there are stools of it in cold-pits. Price of plants, in London, 5s. each.



97

■ 4. *C. SASÁNQUA* Thun. *Sasanqua*, or *Lady Banks's*, Camellia.

Identification. Thun. Fl. Jap., p. 273. t. 30.; Don's Mill., I. p. 576.

Synonymes. *Sasánqua* Kempt. Amén., 853.; Cha-wha, *Chinese*, Staunt. Icon. Chin., 2. p. 466.

Engravings. Thun. Fl. Jap., t. 30.; Kempt. Amén., t. 853.; Chaudl. Ill., t. 5.; and our fig. 98.

Spec. Char., &c. Leaves ovate-oblong, serrated.

Flowers terminal and axillary, solitary. Branches and ovary villous. (*Don's Mill.*, i. p. 576.)

Introduced by Captain Wellbank, of the East India Company's service, in 1811. It forms a loose straggling bush when left unsupported, seldom rising higher than 3 ft. or 4 ft.; but, when tied to a stake, attaining the height of 6 ft. or 8 ft. It produces its white flowers in November and December, which very much resemble those of the tea tree.

It is extensively cultivated in China, for the same object as *C. oleifera*; that is, for crushing the seeds for oil, and using the leaves for adulterating tea. In China, it is said to grow on the debris of rocks and stones: here it succeeds best in moderately strong, rich, sandy soil, and is readily increased by inarching or grafting on *C. japonica*.

It is said to grow on the debris of rocks and stones: here it succeeds best in moderately strong, rich, sandy soil, and is readily increased by inarching or grafting on *C. japonica*.

It is said to grow on the debris of rocks and stones: here it succeeds best in moderately strong, rich, sandy soil, and is readily increased by inarching or grafting on *C. japonica*.

It is said to grow on the debris of rocks and stones: here it succeeds best in moderately strong, rich, sandy soil, and is readily increased by inarching or grafting on *C. japonica*.

It is said to grow on the debris of rocks and stones: here it succeeds best in moderately strong, rich, sandy soil, and is readily increased by inarching or grafting on *C. japonica*.

It is said to grow on the debris of rocks and stones: here it succeeds best in moderately strong, rich, sandy soil, and is readily increased by inarching or grafting on *C. japonica*.

It is said to grow on the debris of rocks and stones: here it succeeds best in moderately strong, rich, sandy soil, and is readily increased by inarching or grafting on *C. japonica*.

It is said to grow on the debris of rocks and stones: here it succeeds best in moderately strong, rich, sandy soil, and is readily increased by inarching or grafting on *C. japonica*.



98

■ 5. *C. KISSI* Wall. The Kissi Camellia.

Identification. Wall. Asiat. Res., 13. p. 429.; Don's Mill., I. p. 576.

Synonyme. *C. Keina* Hamil. MSS. in D. Don's Prod. Fl. Nep., p. 924.

Engravings. Wall. Pl. Asiat. Rar., t. 256., and our fig. 99.

Spec. Char., &c. Leaves elliptical, serrulated, bluntly acuminate. Flowers sessile, axillary, generally solitary, and somewhat terminal, usually 4-petaled, and with 3 distinct, furrowed, woolly styles, which are about equal in length to the stamens. Native of Nepal, at Narainhetty; where it flowers in September, and where it is called *kengua* by the inhabitants. This species is very like *C. Sasánqua*. The flowers are white and fragrant. It is called in the Newar language, *kissi*, or *kissi-swa*. The leaves have a very strong, but transient, smell of tea; but an infusion possesses only a very slight degree of flavour, owing, perhaps, as Mr. Gardner justly observes, to the defective manner of gathering and drying them. It has also been ascertained by Mr. Gardner that the Nepalese extract an oil from the seed of the *Kissi* by pressure, which is much valued by them as a medicine. (*Don's Mill.*, i. p. 676., adapted.) Introduced into the garden of the Horticultural Society in 1823; but, not being a very showy, species, it has not been extensively propagated. There are stools of it in a cold-pit in the Vauxhall Nursery. Plants, in London, cost 10s. 6d. each, and at Bollwyller, 30 francs.



■ 6. *C. OLEIFERA* Abel. The oil-bearing Camellia.

Identification. Abel's Chin., p. 174; Don's Mill., i. p. 577.

Engravings. Lodd. Bot. Cab., 1065.; Ker. Bot. Reg., 492.; Chaudl. Ill., t. 3.; and our fig. 100.

Spec. Char., &c. Leaves elliptic-oblong, acute, serrated, coriaceous, shining. Flowers solitary. Calyxes silky, deciduous. Petals 5—6, 2-lobed. (*Don's Mill.*, i. p. 577.) A native of Cochin-China, where it is cultivated, and forms a small tree 10 ft. high. This bears a close resemblance to the two preceding species: the flowers are very numerous, white, and fragrant. The Chinese call it "the oil-bearing tea plant," as it very closely resembles tea. Dr. Abel sometimes found it of the magnitude of a moderately sized cherry tree, and never less than the size of a shrub 6 ft. or 8 ft. high. At a distance, these plants looked as if they had been lightly clothed with snow; but nearer they exhibited one immense garden of white roses. This species is said to have been originally brought to the country by Lord Macartney; but it was afterwards lost till 1820, when it was reintroduced by Captain Nisbett. It is readily distinguished from *C. Sasánqua*, as it is of a much more robust habit, and larger in every respect, with thicker leaves, having moderately large serratures, and being sharp at the point. (*Gard. Mag.*, vol. vi. p. 290.) The Chinese extract an oil from the seed, which is in very general use in the domestic economy of China. The seeds are white, and are reduced to a coarse powder, which is afterwards chewed or boiled in bags, and then pressed, when the oil is produced. The seeds of all the different species of camellia are said to be used in China for the same purpose. (*Abel.*)



■ 7. *C. EURYOIDES* Lindl. The Eurya-like Camellia.

Identification. Lindl. Bot. Reg., t. 983.; Don's Mill., i. p. 577.

Synonyme. *Thèa euryoides* Booth, in *Hort. Trans.*

Engravings. Lodd. Bot. Cab., 1493.; Lindl. Bot. Reg., t. 983.; and our fig. 101.

Spec. Char., &c. Leaves ovate-lanceolate, acuminate, serrated, silicate beneath. Branches hairy. Peduncles lateral, 1-flowered, scaly. (*Don's Mill.*, i. p. 557.) Native of China, whence it was introduced, in 1824, in rather a singular manner. The grafted part of a camellia, brought from China, in 1822, by Mr. John Potts, having perished, the stock sent up young shoots, and

proved to be this species, which had been before unknown to botanists. It forms a diffuse bushy plant, with hairy branches, obovate, acuminate, serrated leaves, and small neat white flowers, never expanding fully, but in size resembling those of a *Thea*. It is inferior in showiness to any of the previously known camellias; but must be considered a subject of much interest to the cultivator, from its being one of the stocks employed by the Chinese in propagating the ornamental species of the genus. (*Hort. Trans.*)

This species is not much cultivated, because, though beautiful as an evergreen shrub, it is less so than the other sorts. It deserves a place, however, on the conservative wall, wherever the collection is extensive.

Soil, Situation, and general Management. The camellias will grow in any free soil; but a loam inclining to sand, enriched with leaf-mould or thoroughly rotten dung, seems to suit them best. Most of the species and varieties seem to prefer a situation somewhat shaded; which, as we have already observed, seems to be generally the case with evergreen plants having broad shining leaves. For this reason, an east or west wall, or even a north wall, inclining to the east or west, will be found preferable to a south wall for the more tender sorts; and for those which are to be treated as bushes, a situation in the shrubbery or arboretum, where they will be shaded by tall trees during the hottest part of the day, is desirable.

Where there is ample space and a decided taste for the genus, a camellia garden, containing all the species and varieties, might be formed in the following manner. Choose a situation, either level, or having a west or south-east aspect; and enclose a circle, a quarter of an acre in extent, with a 9-inch brick wall, built hollow, and having holes about the size of the end of a brick at 3 ft or 4 ft. apart, immediately under the coping, for the purpose of receiving rafters for supporting a temporary roof of thatched hurdles or of boards. Against both sides of this wall all the more tender sorts of camellia might be planted; and the interior of the area might be devoted to the hardier sorts, to the green tea, and to other half-hardy and evergreen shrubs from China and Japan, such as *Illicium*, *Magnolia fuscata*, &c. Where a number of conservative gardens are to be placed together, the walls may be arranged in the form of pentagons, by which means, as in the cells of the honey-bee, no space would be lost. The form of the garden being either circular or pentagonal, and the walls being hollow, the latter need not be more than a brick in width, and they may be carried to the height of 10 ft., which will be sufficient. Being hollow, they might all be heated by steam from one boiler, a small steam pipe being conducted along their foundations.



GENUS II.



THE A L. THE TEA TREE. *Lin. Syst.* Monadelphica Polyandria.

Identification. *Lin. Gen.*, No. 668; *Dec. Prod.*, 1. p. 530; *Don's Mill.*, 1. p. 577.

Derivation. Altered from *cha*, the Chinese name for tea.

Gen. Char., &c. *Calyx* of 5 sepals. *Petals* 5—9, disposed in two or three rows, covering at the base. *Stamens* almost unconnected to the very base. *Anthers* roundish. *Style* trifid at the apex. *Capsules* of 3 almost distinct carpels, 3-seeded; the dissepiments are formed from the edges of the valves being bent inwards. Beautiful evergreen shrubs from China. (*Don's Mill.*, 1. p. 577.) The species, in general appearance, closely resemble

camellias, with which genus they are united by various botanists. The leaves are large, shining, laurel-like, and the flowers white, axillary, pediceled, and sweet-scented. The culture may be considered the same as that of the camellia, but some of the species are less hardy.

❖ 1. *T. VIRIDIS* L. The common, or green Tea.

Identification. Lin. Sp., 755.; Don's Mill., l. p. 577.

Synonymes. *T. Bohæa stricta* Ait. Hort. Kew., ed. 2. vol. 3. p. 303.; *T. sinensis* Sims, Bot. Mag., t. 998.; *S. chinensis* var. *a. viridis* Dec. Prod., l. p. 530.; *Camellia viridis* Link, Enum., 2. p. 73.; *Thèa cantonensis* Lour. Coch., p. 539.

Engravings. Lodd. Bot. Cab., t. 227.; Woodv. Med. Bot. Suppl., 116. t. 256.; Black. Herb., t. 351.; Letts. Mon., t. 1.; and our fig. 102.

Spec. Char., &c. Leaves elliptic-oblong, serrated, 3 times longer than broad. Flowers of 5 sepals and 5—7 petals, axillary, solitary, erect. Fruit nodding, dehiscent. (*Don's Mill.*, i. p. 577.) An evergreen shrub, with light green laurel-like leaves, and large white fragrant flowers, which are produced from September till December. Introduced from China in 1768. Height from 6 ft. to 8 ft.



102

❖ 2. *T. BOHÆA* L. The Bohea, or Black, Tea.

Identification. Lin. Sp., 743.; Don's Mill., l. p. 577.

Synonymes. *T. chinensis* β *Bohæa* Sims, Bot. Mag., t. 998.; Dec. Prod., l. p. 530.

Engravings. Lodd. Bot. Cab., 226.; Lois. Herb. Amer., t. 255.; Kæmpf. Amœn., t. 606.; Sims. Bot., t. 998.; and our fig. 103.

Spec. Char., &c. Leaves elliptical-oblong, obtuse, crenated, twice as long as broad. Flowers of 5 sepals, and 5 petals, axillary, twin or ternary. (*Don's Mill.*, i. p. 577.) An evergreen shrub, with dark green leaves, much smaller than those of the preceding species; and white flowers, also smaller, but fragrant. Introduced from China in 1780, and generally treated as a frame or green-house plant.

History, Uses, &c. The genus *Thèa* (forming the *Thèacææ* of Mirbel, and included in the *Camelliææ* of Jussieu and De Candolle) is almost exclusively confined to China, Japan, and some of the neighbouring islands; but, as the species are plants which have been cultivated for an unknown length of time, it is difficult to ascertain their native country. Of late, the *Thèa viridis* has been discovered in Upper Assam through an extent of country of one month's march, and within the East India Company's territories, from Sadiya and Beesa to the Chinese frontier of the province of Yunnan, where the shrub is cultivated for the sake of its leaf. This discovery was made in 1826, by Mr. David Scott; and an account of it is given in the *Journal of the Asiatic Society of India* for January 1835, and in the *Gardener's Magazine*, vol. xi. p. 429. It appears that the inhabitants of these countries are in the habit of boiling the stalks and leaves, and then squeezing them into a ball, which they dry in the sun, and then retain for use.



103

Much has been written on the plant which produces the tea of commerce. Dr. Lettsom, who wrote a pamphlet on the subject in 1772, asserts that all the different kinds of tea brought to Europe are the produce of *Thèa viridis*, and that the whole difference in the qualities of teas depends, not on the species of plant, but on the soil and climate in which the plants are grown, the different ages and periods at which the leaves are gathered, and the different modes of preparing and drying them. A green tea plant, he asserts, planted in the bohea tea country, will produce bohea tea; and a plant from the bohea tea country, planted in the green tea country, will produce green tea. Among all the different opinions that have been advanced on the subject, this appears to us by far the most plausible. It is said, however, that *Thèa Bohæa* is cultivated in the southern provinces, as the *Thèa viridis* is in

the north; and that hyson, and all its numerous varieties, are made from the latter; and bohea, and all its varieties, from the former: but, comparing the price of green and black tea in this country, and the quantity of the latter which is consumed in proportion to that of the former, it is difficult to believe that the black teas are all made from the leaves of *Thèa Bohèa*, which is a very distinct species, constitutionally much more tender than *Thèa vridis*, and of comparatively slow and diminutive growth. From Kæmpfer, Thunberg, and Siebold we learn that the tea plant is extensively cultivated in Japan, the various islands of which extend from 30° to 41° of north latitude; from which, in considering its extensive culture in China, and the great extent of territory where it is found indigenous, it is easy to conceive that its varieties may be as numerous as those of the grape vine or the apple are in the south of Europe. Mr. Reeves has disputed, in *Gard. Mag.*, vol. ix. p. 713, 714., the correctness of some of Mr. Main's statements. Whoever wishes to pursue this subject at greater length, and to consult a digest of all that has been hitherto written on it, may peruse Royle's *Illustrations*, from p. 108. to p. 113.

The *Thèa vridis* is sufficiently hardy to stand the air in the neighbourhood of London, with little or no protection. There are bushes of it from 6 ft. to 8 ft. high, and 20 ft. or more in circumference, in the Mile End Nursery, which, in severe winters, have only a little pea-haulm or a mat thrown over them. There are, also, large plants at Syon, at Purser's Cross, at Vere's Villa, at Brompton, at Upton, near Ham, at Lady Tankerville's, Walton on Thames (40 years old, and 8 ft. high), &c., which, we believe, have never received any protection whatever. The plants at Mile End are in a deep sandy loam, and they are remarkable for sending down their strong, thick, black roots in a perpendicular direction to a great depth. There is a plant at Farnham Castle, Surrey, almost as large as those in the Mile End Nursery; and there are plants at White Knights, and at various other places, which leave no doubt of the hardness of this species. Indeed, it thrives far better in the open air than in pots; probably owing to its constitutional habit of throwing down its roots perpendicularly to a great depth. The only conservatory in which we have seen it in a prosperous state is that at Cashio-bury, in Hertfordshire, where the glass roof is taken off during the greater part of the year. The plant not only flowers freely in the open air, but sometimes, as at Farnham Castle, it ripens seed. It is easily propagated by layers; and its general treatment, both in the nursery and in the pleasure-ground, as a half-hardy shrub, may be considered the same as that of the camellia; with this difference, that, being more of a bush than that plant, and loving the shade still more, it does not seem to answer so well against a wall. In the warmest parts of Devonshire, and the south of Ireland, it might be grown as an article of field culture for its leaves; and, if our prejudice in favour of the Chinese mode of preparation could be got over, and the leaves could be slightly fermented, and dried in the same manner as the best meadow hay is about London, being afterwards compressed into cakes to keep for use, the principal nations of Europe might easily grow their own tea, instead of importing it from China, if such a measure were found necessary, or thought desirable. The culture of the plant for commercial purposes has been tried at Rio Janeiro, at Algiers, and, we believe, in Australia. There is much less difficulty in growing the plants, than in preparing the leaves in the Chinese manner; and, as this is principally performed by manual labour, it can only be done profitably where the population is extremely numerous, and the means of living proportionably as cheap as in China or India. At some future time an imitation of this process will, probably, be effected by means of steam.

The black tea (*Thèa Bohèa*) is a much more delicate plant, and is very seldom to be seen in England in the open air in a thriving state. It will neither thrive in pots, nor do well in a conservatory, unless it is quite close to the glass. The best situation for it seems to be a pit, where it may be covered

with glass during winter, and exposed to the air during summer. In a conservative arboretum, it may be preserved by placing litter, fern, or spruce branches round the roots, and covering the top with a case of wickerwork, which, in climates colder than that of London, may be thatched. There are stools of it in the open ground in the Kensington Nursery, and in some others; but they are protected with mats in winter.

* *Other Species and Varieties of Thèa.*

■ 3. *T. COCHINCHINE[']NSIS* Lour., the Cochin-China Tea,

is said to have narrower leaves than the other species, and to have 1-seeded fruit opening at the apex. It is a shrub, growing 8 ft. high, and the leaves are used by the inhabitants of Cochin-China medicinally, as a sudorific and refrigerant. It has not yet been introduced into Britain, and is, in all probability, only a variety of the green tea.

■ 4. *T. OLEO[']SA* Lour., the oily Tea,

has lanceolate leaves, and peduncles 3-flowered, and axillary. An oil is said to be obtained from the seed of this shrub, which has not yet been introduced into Britain.

A species of tea grown in the province of Canton, with a pale-coloured leaf, which is occasionally mixed with Congou tea, is mentioned by Mr. Reeves (*Royle's Illust.*, p. 111.); and this, with the numerous other sorts which are, doubtless, in cultivation in China and Japan, may be expected in Britain at some future period.

Sect. III. *Anticipated Ternströmiæcæ.*

We have already mentioned that there are, undoubtedly, many varieties of *Caméllia japonica* in China and Japan which have not yet found their way to Britain; and there can be no doubt that the varieties of the green tea are still more numerous. In p. 173., it appears that *Eurya acuminata*, which belongs to this order, is likely to be hardy. This shrub, Mr. Royle observes, attains the height of about 8 ft. or 10 ft., and is common in the Himalaya at an elevation of 6500 ft. The leaves are thick, coriaceous, finely serrated, smooth, and the young ones hairy. The flowers, owing to the early fall of the leaves from the lower parts of the branches, appear to be lateral; but they actually rise out of the axils of the fallen leaves. The plant is a handsome evergreen bush, well meriting introduction. (*Royle's Illust.*, p. 128.)

CHAP. XX.

OF THE HALF-HARDY LIGNEOUS PLANTS OF THE ORDER AURANTIA[^]CEÆ.

We introduce this order for the sake of noticing two genera, the species of which may, with care, be cultivated against flued walls without the protection of glass. These genera are *Citrus* and *Limonia*. The species of the first are trees universally known and admired, natives of India; and those of the second Himalayan shrubs, growing at considerable elevations in Nepal.

Orange trees, when first introduced into England in 1595, were grown against a wall at Bedington, in Surrey, and flowered and fruited there for many years; till, as Evelyn informs us (see *E. of Gard.*, ed. 1835, p. 967.), they were neglected. With a little care, and without the expense of glass, there can be no doubt that all the *Citrus* family might be grown against a hot wall in the climate of London, in as great, or greater perfection, than they are now to be seen in those formal artificial contrivances, tubs and boxes kept under glass, and which are far more expensive than hollow walls to be heated by steam or smoke flues, and protected by thatched hurdles, or reed or straw mats. Those who have seen the walls covered with orange and lemon trees at Woodhall in West Lothian, at Coombe Royal in Devonshire, and at M. Fion's in Paris, will not wonder at our great anxiety to encourage the culture of this plant in the conservative manner. We have also seen fine orange trees in the neigh-



bourhood of Paris, grown as standards in the open air, but enclosed during winter with double glass cases, which are removed in May, and replaced in September; the outer case being covered with straw mats, for weeks together, in severe weather. When the wood of the orange tree is ripened, and the sap is set to rest by withholding water, it may be kept in the dark, for two or three months together, without injury, provided the air be kept dry. The hardiest kind of orange appears to be the Seville, or bitter, orange. (fig. 104.) Mr. Spence, who passed some winters in Florence, states *Gard. Mag.*, vol. vii. p. 298. that the bitter orange is by far the most hardy variety grown in Italy; and that, in the neighbourhood of Florence, where the cold is so great that skating is sometimes practised occasionally during four months of the year, and where, in the winter of 1829-30, the thermometer was repeatedly down to 26° and 21° Fahr. at 8 a. m., there are standards in the open air, in sheltered situations, but not planted near a wall, with stems from 4 m. to 6 in. in diameter, which pass the most severe winters with no other injury than having the points of the last year's shoots turned a little yellow. The bitter orange is in general use among the Italians, as a stock for grafting all the other sorts on; another proof that it is one of the hardiest and most vigorous-growing sorts. When a trial is to be made in England, we would recommend commencing with this variety; and protecting the ground by litter, and both sides of the wall by mats or boards, for several years, till the plants were fully established. The simple protection of straw mats, placed over the trees, might, probably, be found enough afterwards, at least in favourable situations. The walls should always be fluted, or hollow, in order to afford the opportunity of lighting a fire occasionally. The north side of the wall might be kept warm by ivy; or, if it were an east and west wall, it might be clothed with half-hardy trees on both sides, and protected accordingly.

As examples of different species of the genus *Citrus* which have attained large dimensions in the open air in England, we cite the following from different places in Devonshire. At Salcombe, near Kingsbridge, in the garden of the Rev. Mr. Henshaw, are two walls covered with fine healthy orange trees: one wall has a broad coping, and glass shapes for placing before the trees in severe winters; the other has reed hurdles for the same purpose, and, though this sometimes excludes the light for three months, it is considered a better protection than the glass. The kinds are, the common, Seville, and blood-red orange, and the citron, lemon, lime, and shaddock. At Dartmouth, in the garden of Mr. Strong, is a very large citron tree. At Luscombe, the seat of C. Hoare, Esq., there are some remarkable large trees of citrons, shaddocks, and limes, which grow so vigorously, that they make shoots of from 6 ft. to 7 ft. in length in one year. They were raised from cuttings on the spot; and, the third year after being planted out, they produced a number of fine fruit, some of the lemons measuring from 15 in. to 18 in. in circumference, and weighing 14 oz. (*See Gard. Mag.*, vol. ii. p. 59.)

At Kitley, the seat of E. P. Bastard, Esq., there is a remarkably hardy variety of the shaddock, raised on the spot, and known as the Kitley shaddock, which is much harder than most other sorts of *Citrus*. At Coombe Royal, the seat of John Luscombe, Esq., are a number of lime, citron, orange, and lemon trees, of which a description will be found in the *Gard. Mag.*, vol. x. p. 36. The trees are planted against a south wall, which is divided by buttresses, forming recesses in which the trees are trained in the manner of common fruit trees. Each recess is 2½ ft. wide, and about the same in height; and is protected, during the winter, by a frame of wood, which is wholly removed in the summer, and partially by day, at other seasons; the trees only requiring to be guarded from severe frost. A more interesting sight cannot well be imagined by the lovers of horticulture, than that which is presented by these trees on a sunny day in winter, when the open frames furnish a display of the richest foliage, and of fruit rivalling the produce of foreign countries. The soil, which is seldom manured, is light and rich, on a slaty substratum, and seems admirably adapted to trees of the citron tribe. The situation in which they are grown is in a sheltered valley, protected from wind. It is necessary to add, that the lime is grown under glass in winter; but it is believed that the tree would thrive equally well under wood, and be more secure from frost and storms; a fine tree, in full bearing, having been destroyed, a few years since, by the accidental breaking of a pane or two of glass in winter. The fruit is produced abundantly by the citron, oranges, and lemons; by the lime more sparingly, from the tree being young, and in a vigorous growing state; and, when gathered at a proper period, the fruit is of a fine flavour, and full of juice. Citrons are every year ripened measuring from 15 in. to 18 in. in circumference; lemons, 11 in. by 9 in.; and oranges, 10 in. by 9 in.; weighing from 5 oz. to 9 oz. The leaves are generally 11 in. long, and 6 in. or 7 in. wide. (*See Gard. Mag.*, vol. x. p. 36.)

The genus *Linnæa* (from *lymoun*, the Arabic name of the citron, Latinised) is not very well known; but it includes thirteen or fourteen species, all of which bear more or less affinity to the genus *Citrus*. The species likely to prove hardy or half-hardy are the following, *L. Lauræola* Dec., a native of Nepal, with leaves like those of *Daphne Lauræola*, and white flowers, and which is found on the top of cold and lofty mountains, where it is, for some months in the year, buried under the snow. The leaves are highly fragrant; and are, like others of a similar kind, Mr. Royle informs us, employed in the religious ceremonies of the inhabitants of the hills. There can be no doubt of this species of *Linnæa* being perfectly hardy. — *L. citrifolia* Willd. is a native of China, and has white flowers and red berries, with simple or trifoliate leaves. — *L. parviflora* Bot. Mag., t. 2416, has impari-pinnate leaves, and white flowers. It is a native of China, and grows to the height of 6 ft. — *L. ambigua* Dec. is said to be cultivated in East Florida; but very little is known of this and the preceding species. — *L. australis* is a native of New Holland, where it forms a tree 25 ft. high; and there is a plant of it at Kew, introduced in 1830. There are four other species of this genus in our stoves and green-houses; but *L. Lauræola* remains to be introduced by Mr. Royle, or some other patriotic individual.

CHAP. XXI.

OF THE HARDY AND HALF-HARDY LIGNIOUS PLANTS OF THE ORDER
HYPERICACEÆ, AND TRIBE *HYPERICÆÆ*.

Distinctive Characteristics. Thalamiferous. (H. B.) Sepals 4 or 5, unequal, with an imbricate aestivation. Stamens, in nearly all, numerous, and in 3

or more parcels. Fruit, a capsule or berry of many valves and many cells; the edges of the valves curved inwards. Seeds attached to a placenta in the axis, or on the inner edge of the dissepiments. Leaves dotted; in most, opposite and entire. Flowers, in most, yellow. Sap yellow, resinous. (*Lindley, Introd. to N. S.*)

Description, &c. The hardy ligneous plants belonging to this order are all shrubs or undershrubs sub-evergreen or deciduous; with dotted leaves, smooth, oblong, or lanceolate; and yellow flowers. They are natives of Europe, North America, or Asia. Some few of Africa, but more of Australia. Medicinally, they are bitter and slightly astringent: the soft parts of many species contain a fragrant oil, and others secrete a yellow juice: from a Mexican species the gum Squitum Gulla of commerce is produced. In gardening, these shrubs must be considered more as flowering shrubs for dry borders, than as woody plants of permanent duration in the arboretum or shrubbery. There are, however, one or two exceptions. All the species throw up abundance of side suckers, and are readily propagated by division of the plant, or by cuttings; and some of them ripen seeds. They will grow in any soil, not too stiff, or too much charged with moisture.

HYPE`RICUM L. Capsule membranous. Stamens numerous, disposed in 3 to 5 bundles at the base.

ANDROSÆ`MUM Chois. Capsule baccate, 1-celled. Calyx 5-parted. Stamens numerous, monadelphous at the base.

GENUS I.



HYPE`RICUM L. THE ST. JOHN'S WORT. *Lin. Syst.* Polyadélphia Polyándria.

Identification. *Lin. Gen.*, 392; *Juss.*, 255; *Dec. Prod.*, 1. p. 542; *Don's Mill.*, 1. p. 601.

Synonymes. Fuga Dæmonum; Mille Pertuis, *Fr.*; Johannis Kraut, *Ger.*

Derivations. The name of *Hypéricum* is as old as the time of Dioscorides; but its origin and meaning are uncertain. Some derive it from the Greek words *hyper*, under, and *eilôn*, an image; and suppose it to signify that the upper part of the flower represents a figure. Others state that *hyper* signifies through, and that the name alludes to the pellucid dots in the leaves, which form small lenses, through which, when held up to the light, images might be seen. *Donnegan*, in his *Lexicon*, conjectures the word *Hypéricum* to be taken from *hyper*, for, in the place of, &c., and *ereikê*, heath; from the plant bearing some resemblance to the heath. By others, the word is supposed to be composed from *hyper*, through, on the other side, and *eilô*, to resemble; from the pellucid dots resembling holes or pores; and the French name of the plant, Mille Pertuis, a thousand pores, is evidently derived from the same source. The English name, St. John's Wort, and the German one, Johannis Kraut, are taken from the country people formerly, both in England and Germany, being in the habit of gathering this plant on St. John's day, to use it to protect themselves from evil spirits. This plant, with some others, was employed to make what was called John's fire, which was supposed to be a security, for those who kindled it, against witchcraft and all attacks of demons. For this reason, also, the *Hypéricum* received the name of Fuga Dæmonum.

Gen. Char. &c. Capsules membranous. Stamens numerous, free or joined at the bases into 3 or 5 bundles. Petals 5. Sepals 5, more or less connected at the base, unequal, rarely equal. Styles 3 to 5, rarely connate in one, manent. Capsule 1- or many-celled, many-seeded, 3—5-valved. Integument of seed double. Albumen none. Embryo with the radicle situated at the umbilicus, and with semicylindrical cotyledons. (*Don's Mill.*, i. p. 601.)—Low sub-evergreen shrubs; with yellow flowers, and oppositely placed sessile or subsessile leaves, usually full of pellucid dots on their disks, and some dark ones on their edges, lodging an essential oil. They are chiefly natives of Europe, and vary in height from 1 ft. to 5 ft. They are all considered medicinal, being powerfully astringent, and were formerly in great request by herbalists and other empirical practitioners. Gerard gives a receipt for making a balsam of them in his *Herbal*; which, he says, is "a most pretious remedie for deep wounds, and those that are thorow the body; for the sinnes that are prickt, or any wound made with a venomed weapon." (*Johnson's Gerard*, p. 541.)

§ i. *Ascyreia* Chois. Prod. Hyp.

Derivation. From *a*, not, and *skuros*, hard; that is to say, plants soft to the touch.

Secl. Char. Sepals connected at the base, and unequal. Stamens numerous. Styles 3 to 5. Flowers terminal, large, few, subcorymbose. (*Don's Mill.*, i. p. 601.)

A. *Styles commonly 3.*§ 1. *H. ELA'TUM* Ait. The tall St. John's Wort.

Identification. Ait. Hort. Kew., ed. 2. vol. 3. p. 104.; Don's Mill., i. p. 601.

Engravings. Juss. Ann. du Mus., 3. p. 162. t. 17; Wats. Dend. Brit., t. 85.

Spec. Char. &c. Younger stems reddish. Leaves ovate-oblong, acute, dilated at the base, somewhat emarginate, with the margins somewhat revolute. Flowers corymbose. Peduncles bibracteate. Sepals ovate-oblong. (*Don's Mill.*, i. p. 601.) Height 5 ft. A sub-evergreen shrub, native of North America, with yellow flowers in July and August. Introduced in 1762. Propagated by layers or division, and of the easiest culture in common garden soil.

§ 2. *H. HIRCI'NUM* L. The Goat-scented St. John's Wort.

Identification. Lin. Sp., 1103.; Don's Mill., i. p. 602.

Synonymes. Trægium *Clus.*; Androsœmum fœtidum *Bauh.*, *Park.*, and *Ray.*; Mille Pertuis à Odeur de Bouc, *Fr.*

Engravings. Schkuhr. Handb. 3. t. 213. f. 3.; Wats. Dend. Brit., t. 87.; and our fig. 105.

Spec. Char. &c. Branches winged. Leaves somewhat emarginate at the base, dilated, sessile, acute at the apex, ovate-lanceolate, with glandular margins. Peduncles bibracteate. Stamens exceeding the corolla in length. Seeds 2, appendiculated. (*Don's Mill.*, i. p. 602.) A deciduous undershrub, from the shores of the Mediterranean in 1640, and producing its yellow flowers from July to September. Height, from 3 ft. to 4 ft. The leaves of this species, when bruised, have a very disagreeable smell, resembling that of a goat, whence its name. Plants, in London, cost 6d. each; at Bollwyller, 50 cents.

105



Varieties. *H. h.* 2 *obtusifolium* Dec. has blunter leaves than the original species, and is found on the mountains of Corsica, on humid rocks. *H. h.* 3 *minus* Wats. is a smaller plant than the other, figured in *Dendrologia Britannica*, t. 87.

§ 3 *H. GRANDIFLO'RUM* Chois. The large-flowered St. John's Wort.

Identification. Chois. Prod. Hyp., p. 38. t. 3.; Don's Mill., i. p. 602.

Synonymes. *H. canariense* *Willd.*, not of Lin.

Engravings. Chois. Prod. Hyp., t. 3.

Spec. Char., &c. Stem round, reddish. Leaves ovate-oblong, cordate, somewhat clasping, acute at the apex, netted with pellucid veins. Flowers corymbose. Peduncles bibracteate. Calyx acutish, reflexed upon the peduncle after flowering, much shorter than the corolla. (*Don's Mill.*, i. p. 602.) A half-hardy evergreen shrub, from Tenerife in 1818, producing its fine large yellow flowers in July and August. Height 3 ft. It is commonly treated as a green-house plant; but, considering its native country, there can be no doubt that it would stand very well against a conservative wall.

§ 4. *H. FOLIO'SUM* Ait. The leafy St. John's Wort.

Identification. Ait. Hort. Kew., ed. 1. vol. 3. p. 104.; Don's Mill., i. p. 602.

Synonymes. Shining St. John's Wort.

Spec. Char., &c. Branches winged. Leaves sessile, oval-oblong, rather acute, finely perforated. Calyx lanceolate, caducous. (*Don's Mill.*, i. p. 602.) A deciduous undershrub, introduced from the Azores in 1778, and producing its yellow flowers in August. Height 2 ft. It is commonly treated as a green-house plant; but, in a dry sheltered situation, it requires very little protection.

§ 5. *H. FLORIB'NDUM* Ait. The abundant-flowered St. John's Wort.

Identification. Ait. Hort. Kew., ed. 1. vol. 3. p. 104.; Don's Mill., i. p. 602.

Synonymes. *H. frutescens* *Comm.* Hort. Amst., p. 137.; many-flowered St. John's Wort.

Engravings. Comm. Hort. Amst., t. 68.

Spec. Char., &c. Stem round. Leaves sessile, lanceolate, numerous, without dots. Peduncles dilated, and somewhat compressed towards the apex. Calyx obtuse. Corolla and stamens marcescent. (*Don's Mill.*, i. p. 602.) A deciduous undershrub, from the Canary Islands in 1779, producing its yellow flowers in August. Height 3 ft. It is usually treated as a green-house plant; but it is capable of resisting the winters of the climate of London, in a warm situation, with very little protection.

¶ 6. *H. OLYMPICUM* L. The Olympian St. John's Wort.

Identification. Lin. Sp., 1102.; Sm. Ex. Bot., 2. p. 71.; Dec. Prod., 1. 545.

Synonymes. *H. montis olympi* *Wied. Itin.*, Ray; *H. orientale* *Abre majs Tourne.*

Engravings. Sm. Exot. Bot., 2. t. 96.; Bot. Mag., t. 1867.; and our fig. 106.

Spec. Char., &c. Stem round. Leaves elliptical-lanceolate, rather acute, full of pellucid dots. Calyx ovate, acute. Peduncles bibracteate. Corolla and stamens withering. (*Don's Mill.*, i. p. 602.) An interesting little shrub, with glaucous sessile leaves, native of Mount Olympus and China, introduced in 1706, and producing its yellow flowers from July to September. It grows to the height of from 1 ft. to 2 ft., and requires protection during winter. All the half-hardy species of *Hypericum* might be grown on a conical piece of rock-work, a sort of miniature Mount Olympus, in a warm sheltered part of the pleasure-ground or arboretum. The protection required in winter might be given with complete effect, and at very little expense, by resting a number of poles on the protruding points of the larger rocks or stones, and on these placing thatched hurdles, or even, in warm districts, a few spruce fir branches.



106

¶ 7. *H. CANARIENSE* L. The Canary Island St. John's Wort.

Identification. Lin. Syst. p. 575.; Dec. Prod., 1. p. 544.; Don's Mill., 1. p. 602.

Engraving. Lodd. Bot. Cab., 953.

Varieties. De Candolle notices two: *H. c. triphyllum*, and *H. c. salicifolium*.

Spec. Char., &c. Stem bluntly quadrangular. Branches compressed. Leaves ovate-lanceolate, acute. Calyx ovate, obtuse. Styles 3—4, diverging. (*Don's Mill.*, i. p. 602.) A neat little shrub, a native of the Canary Islands, introduced in 1689, and producing its yellow flowers from July to September. Height 4 ft. It is commonly kept in green-houses or frames; but, like most of the other plants from the Canary Islands and similar climates, it will endure a London winter in the open air against a wall, with the protection of litter or leaves over the ground, and a couple of mats over the top.

B. Styles commonly 5.

¶ 8. *H. CHINENSE* L. The Chinese St. John's Wort.

Identification. Lin. Amœn., 8. p. 523.; Dec. Prod., 1. p. 545.; and Don's Mill., 1. p. 602.

Synonymes. *H. monogynum* *Mill. Illustr.*, 151.; *H. aureum* *Lour.*

Engraving. Mill. Illustr., 151. f. 2.

Spec. Char., &c. Stem round. Leaves elliptical, obtuse, with a few black dots. Peduncles bibracteate. Calyx oblong, obtuse, beset with black dots. Styles collected together. (*Don's Mill.*, i. p. 602.) A sub-evergreen shrub, a native of the East Indies and the Cape of Good Hope; introduced in 1753, and producing its yellow flowers from March to September. Height 3 ft. It is marked in the *Catalogues* as a green-house plant: it would, in all probability, endure the open air, with protection, during winter. It stood at Biel, in East Lothian, in 1825, in an exposed situation. A species bearing this name has stood against the wall in the garden of the London Horticultural Society, for four years, with very little protection. There is a species, named *H. monogynum* L., not of Miller, which is figured in *Bot. Mag.*, t. 334., which appears to be different from this one. It is a native of Japan and China, and grows to the height of 3 ft. In Nepal, a species nearly allied to this (*H. cernuum* *Rox.*, *H. speciosum* *Wall.*) is met with on hills at 3000 ft. of elevation.

¶ 9. *H. CORDIFOLIUM* Choix. The heart-leaved St. John's Wort.

Identification. Dec. Prod., 1. p. 545.; Don's Mill., 1. p. 602.

Synonymes. *H. bracteatum*, and *H. lungatum* *Ham.* MS. in D. Don. Prod., p. 317.

Spec. Char., &c. Stem round. Leaves elliptical, acute, coriaceous, smooth, somewhat stem-clasping, without dots; flower-bearing branches leafy below, crowded. Bractæas ovate-ovate, acute. Sepals ovate, mucronate, without dots. Petals oblong, unequally sided, obliquely mucronulate. Stamens short. Styles unconnected, scarcely longer than the corolla. (*Don's Mill.*, i. p. 602.) A sub-evergreen shrub from Nepal, in 1825, producing its yellow flowers from April to October. Height 2 ft. It is commonly kept in a frame, but would stand our winters, in a warm situation, with a very little protection.

¶ 10. *H. PATULUM* Thun. The spreading St. John's Wort.

Identification. Thun. Jap., p. 295. t. 17.; Don's Mill., 1. p. 603.

Engraving. Thun. Jap., t. 17.

Spec. Char., &c. Stem round, purplish. Leaves ovate-lanceolate, acute, tapering to the base with revolute margins, without dots. Flowers corymbose. Styles recurved at the apex, scarcely longer than the stamens. Peduncles bibracteate. Sepals sub-orbicular, very obtuse. (*Don's Mill.*, i. p. 603.) An evergreen shrub, a native of Japan and Nepal, introduced in 1823, and producing its yellow flowers from June to August. Height 6 ft.

11. *H. KALMI'NUM* Lam. Kalm's St. John's Wort.

Identification. Lam. Diet., t. p. 148.; Don's Mill., t. p. 603.

Synonymes. *H. Bartramianum* Mill.; Virginia St. John's Wort.

Spec. Char. &c. Branches tetragonal. Leaves linear-lanceolate. Flowers 3 to 7, in a terminal corymb. Sepals lanceolate, bluntish. (*Don's Mill.*, i. p. 603.) A sub-evergreen undershrub; a native of North America, in Pennsylvania and Virginia; introduced into England in 1759, and producing its yellow flowers in June and July. Height 3 ft. It was found by Mr. James M'Nab in great abundance in the neighbourhood of the Falls of Niagara, in dry places; and a variety of it (*H. K. elongatum*) was found in moist places in New Jersey. This variety has flowers somewhat smaller than those of the species, but they are equally rich in colour. (*Ed. N. Phil. Journ.*, vol. xix. p. 38.) This species is ornamental, forming a very neat compact bush, and is in very general cultivation. American seeds may be procured, in London, at 6*d.* a packet, and plants at 9*d.* each; at Bollwyller, 1 franc; and at New York, 25 cents.

12. *H. URAL'UM* Ham. The Urala St. John's Wort.

Identification. D. Don Prod. Nep., p. 218.; Don's Mill., t. p. 607.

Derivation. From its name, *Urala sive*, in the Newar language.

Engravings. Bot. Mag., t. 2375.; and our fig. 107.]

Spec. Char. &c. Branches compressed, 2-edged. Leaves elliptical, mucronulate, smooth, shining. Flowers terminal, somewhat corymbose. Sepals oval, very blunt. Petals orbicular. Styles shorter than the stamens. (*Don's Mill.*, i. p. 603.) An undershrub, growing 2 ft. high, introduced from Nepal, where it is found on the tops of mountains, in 1823; and producing its yellow flowers from July to September. In mild situations, and on a dry soil, it may safely be left through the winter without any protection; but this should not be the case where the situation is cold, and the soil tenacious or humid.



107

13. *H. CALYCT'NUM* L. The large-calyxed St. John's Wort.

Identification. Lin. Mant., 106.; Willd., 3. p. 1442.; Hook. Scot., 221.; Dec. Prod., t. p. 546.; Smith's Eng. Fl., 3. p. 323.; and Don's Mill., t. p. 603.

Synonymes. *Androsæmum constantinopolitanum flore máximo Wheeler's Journey*, 205.; the large-flowered St. John's Wort; the large-flowering Tutsan; the terrestrial Sun; Aaron's Beard; Mille Pertuis à grandes Fleurs, Fr.; grossblumiger Johannis Krant, Ger.

Derivations. This species was called *Androsæmum* by the old writers on botany, on account of the tinge of red in different places on the stems, and the redness of the anthers, which were supposed to give it the appearance of being spotted with blood. It was called *Constantinopolitanum* from its having been found near that city, in 1676, by Sir George Wheeler, Bart. The large size of its flowers is remarkable, and has given rise to most of its other names. The name of the Terrestrial Sun is very appropriate to the large golden flowers, with their long ray-like stamens, lying glittering on their bed of dark green shining leaves, which spread over the surface of the ground. The number and length of the stamens are, doubtless, also the origin of the name of Aaron's Beard.

Engravings. Eng. Bot., v. 29. t. 2017.; Bot. Mag., t. 146.; Jacq. Frag., 10. t. 6. f. 4.

Spec. Char. Stem tetragonal, dwarf. Leaves ovate, coriaceous, broad, full of pellucid dots. Flowers large, terminal, solitary. Sepals large, obovate, spreading; capsule nodding. (*Don's Mill.*, i. p. 603.) A beautiful little evergreen, with shining dark green leaves, and bright golden flowers 2 in. or 3 in. in diameter, and having innumerable reddish tremulous anthers. Height from 1 ft. to 18 in. *H. calycinum* is a native of bushy places in the west of Ireland and Scotland. It is extremely valuable for covering banks, rock-work, or the surface of the ground in old shrubberies or picturesque woods, especially for the latter purpose, as it thrives perfectly well under the drip and shade of trees. The root creeps, and a small plant will soon extend itself in every direction, especially if the soil be light, so as to cover a great many square yards in a very short space of time. It is an excellent shelter for game. Plants may be had, in the London nurseries, at 6*d.* each.

14. *H. BALEARIC'UM* L. The Majorca St. John's Wort.

Identification. Lin. Sp., 1101.; Don's Mill., t. p. 603.

Engraving. Curt. Bot. Mag., t. 137.

Spec. Char. Stem quadrangular, warted. Leaves ovate, obtuse, rather stem-clasping. An evergreen shrub, with small warted leaves; native of the Island of Majorca; introduced in the year 1714; and producing its yellow flowers from March to September. Height 2 ft. It requires some protection during winter, stood the winter of 1825, in a sheltered situation, at Biel, in East Lothian

§ ii. *Perforària* Chois.

Identification. Chois. Prod. Hyp., p. 44., Dec. Prod., 1. p. 546; Don's Mill., 1. p. 603.

Derivation. From *perforatus*, perforated; because the leaves are full of pellucid dots, which gives them the appearance of being perforated.

Sect. Char. Calyx of 5 equal sepals, toothed in some with glandular teeth, but entire in others, connected at the base. Stamens numerous, free or disposed in 5 sets. Styles commonly 3. Herbs or undershrubs. Flowers axillary, or in terminal paniced corymbs. Leaves rarely linear. (*Don's Mill.*, 1. p. 603.) Undershrubs, from 1 ft. to 3 ft. in height.

A. *Sepals entire.*■ 15. *H. PROLI'FICUM* L. The prolific St. John's Wort.

Identification. Lin. Mant., 106.; Don's Mill., 1. p. 605.

Synonymes. *H. folièsum* Jacq., Hort. Schönbr., 3. p. 27.; *H. Kalmianum* Du Roi, Harbk., 1. p. 310.

Engravings. Wats. Dend. Brit., t. 88.; Jacq. Hort. Schönbr., t. 200.

Spec. Char., &c. Stem round. Branches angular. Leaves linear-lanceolate, with revolute edges, full of pellucid dots. Corymbs few-flowered. Sepals ovate-lanceolate, stamens very numerous. Styles usually connected together. (*Don's Mill.*, 1. p. 605.) A sub-evergreen shrub, from Virginia and Canada, introduced in 1758, and producing its yellow flowers from June till August. Height 4 ft. Frequent in gardens, and forming a dense leafy bush, covered with flowers great part of the summer, and with seed-pods in the autumn. American seeds, in London, 6d. a packet; and plants, in London, 9d. each; and at Bollwyller, 50 cents each.

■ 16. *H. HETEROPHY'LLUM* Vent. The various-leaved St. John's Wort.

Identification. Vent. Hort. Cels., t. 68.; Don's Mill., 1. p. 607.

Engraving. Vent. Hort. Cels., t. 68.

Spec. Char., &c. Stem suffruticose, round. Leaves linear-lanceolate, full of pellucid dots; axillary ones crowded, imbricate, very short, blunt. Sepals acute, somewhat unequal. (*Don's Mill.*, 1. p. 607.) A low sub-evergreen undershrub, from Persia, in 1712, and producing its yellow flowers in July and August. Height 2 ft. It requires some protection during winter.

■ 17. *H. EGYPTI'ACUM* L. The Egyptian St. John's Wort.

Identification. Lin. Sp., 1103.; Don's Mill., 1. p. 607.

Engravings. Lin. Amœn., 8. t. 8. f. 3.; Ker Bot. Reg., t. 195.

Spec. Char., &c. Stem round. Leaves small, ovate, crowded, without dots. Flowers few, almost sessile. Sepals lanceolate, acute. Styles small, diverging. (*Don's Mill.*, 1. p. 607.) A sub-evergreen undershrub, with glaucous leaves and small flowers; introduced from Egypt in 1787, and producing its yellow flowers in June and July. Height 2 ft. It requires protection during winter.

■ 18. *H. GALIÖI'DES* Lam. The Galium-like-leaved St. John's Wort.

Identification. Lam. Diet., 4. p. 160.; Don's Mill., 1. p. 609.

Spec. Char. Stem suffruticose, round, straight. Leaves linear-lanceolate, tapering to the base, broadest at the apex, acute, with revolute dotted margins. Sepals linear, acute, reflexed after flowering. Styles at first connected, but at length free. Capsules conical, very acute. (*Don's Mill.*, 1. p. 609.) A sub-evergreen undershrub, native of North America, from New Jersey to Carolina, in sandy moist places near rivulets; producing its yellow flowers from July to September. Height 2 ft.

■ 19. *H. AXILLA'RE* Lam. The axillary-flowered St. John's Wort.

Identification. Lam. Diet., 4. p. 160.; Don's Mill., 1. p. 609.

Synonymes. *H. fasciculatum* Willd. Spec., 3. p. 1452., exclusive of the synonymes of Michx., Pursh, Fl. Amer. Sept., 2. p. 576.; *H. Coris* Walt. Fl. Carol., 190.

Spec. Char., &c. Stem shrubby, round, diffuse. Leaves lanceolate-linear, narrowed at the base, with revolute margins. Sepals rather unequal. Styles, at first joined, but afterwards free. (*Don's Mill.*, 1. p. 609.) A sub-evergreen undershrub, native of the pine woods of Georgia and Florida; producing its yellow flowers in July. Height 2 ft.

B. *Sepals toothed, usually with the Teeth glandular.*n. 20. *II. GLANDULOSUM* Ait. The glandular St. John's Wort.

Identification. Ait. Hort. Kew., ed. 1. vol. 3. p. 107.; Don's Mill., 1. p. 609.

Spec. Char., &c. Stem shrubby, round, erect, branched. Leaves elliptical-lanceolate, acute, with glandular margins, and pellucid dots. Calyx lanceolate, acute. (*Don's Mill.*, i. p. 609.) A sub-evergreen undershrub, native of Madeira and Teneriffe, introduced in 1777; producing its pale yellow flowers, the petals of which are full of brown dots, from May to August. Height 2 ft. It requires protection during winter.

n. 21. *II. SERPYLLIFOLIUM* Lam. The Wild-Thyme-leaved St. John's Wort.

Identification. Lam. Dict., 4. p. 176.; Don's Mill., 1. p. 610.

Engraving. Mor. Hist., 2. p. 469. sect. 5. t. 6. f. 2.

Spec. Char., &c. Stem suffruticose, round. Leaves ovate, obtuse, on very short petioles, with revolute margins. Calyx ovate, obtuse, fringed. (*Don's Mill.*, i. p. 610.) A neat little bush, which has been in cultivation as a half-hardy shrub since 1688. It produces its yellow flowers in July and August, and grows to the height of 1½ ft. It is well adapted for culture in pots; or on the warmest part of rockwork.

n. 22. *II. EMPETRIFOLIUM* Willd. The Empetrum-leaved St. John's Wort.

Identification. Willd. Spec., 3. p. 1452.; Don's Mill., 1. p. 610.

Engravings. Dend. Brit., t. 141.; and our fig. 108.

Spec. Char., &c. Stems suffruticose, round, with subulate branchlets.

Leaves linear, ternary, with revolute margins. Calyx small, obtuse.

Petals without glands. (*Don's Mill.*, i. p. 610.) A neat little shrub, a

native of the south of Europe, particularly near the Mediterranean; and

introduced in 1820, and producing its yellow flowers from May to

August. Height 2 ft. This is one of the neatest species of the genus,

and it well deserves a place on the hypericum mount, suggested

under *H. olympicum*, p. 599., because it is not altogether hardy. As

it is a slow-growing plant, and small in all its parts, it should not be

placed immediately adjoining any of the rapid-growing, broad-leaved,

or bulky species, unless required in the way of contrast. It would

suit very well to accompany *H. baleáricum*, *H. cricoides*, and *H.*

Coris, which are also half-hardy species.

n. 23. *II. CORIS* L. The Coris-leaved St. John's Wort.

Identification. Lin. Spec., 1107.; Don's Mill., 1. p. 610.

Spec. Char., &c. Stem shrubby, erect, round. Leaves in whorls, linear, with revolute margins. Calyx

linear, bluntish. (*Don's Mill.*, i. p. 610.) A small shrub, of the habit of the last, but a native of the

Levant, whence it was introduced in 1640. It produces its yellow flowers from May to September.

Height from 1½ ft. to 2 ft. This species stood out, in a sheltered situation at Biel, in the winter of

1825. The plant *Coris*, which it is said to resemble, is the *Coris monspeliensis* H., a herbaceous

biennial, one of the Primulaceæ.

n. 24. *II. ERICOIDES* L. The Heath-like St. John's Wort.

Identification. Lin. Spec., 1104.; Don's Mill., 1. p. 611.

Engravings. Cav. Icon., 2. p. 20. t. 122.; Pluk. Phyt., t. 93. f. 5.

Spec. Char., &c. Stem suffruticose, round, twisted, small. Leaves linear, acute, much crowded,

dotted, glaucous, small. Sepals acute, hardly glandular. (*Don's Mill.*, i. p. 611.) A neat little

heath-like shrub, a native of Spain, Portugal, and the Levant; introduced in 1821, and producing

its yellow flowers from June to September. It requires protection during winter.

§ iii. *Bràthyæ* Chois.

Identification. Chois. Prod., p. 58.; Dec. Prod., 1. p. 553.

Derivation. From *bràthys*, the Greek name of the savin tree (which is derived from *brazō*, to

overheat); in allusion to the habit of the shrubs, which resembles that of the savin tree, or juniper.

Sept. Char. Calyx of 5 entire equal sepals, usually very like the leaves.

Stamens numerous, disposed in bundles. Styles 3 to 4. Subshrub, with

axillary solitary flowers, and imbricate, whorled, or crowded leaves, which

are usually linear-awl-shaped. (*Don's Mill.*, i. p. 611.) Sub-evergreen under-

shrubs.

A. *Styles 3, with simple Stigmas.*n. 25. *II. FASCICULATUM* Lam. The fascicled-leaved St. John's Wort.

Identification. Lam. Dict., 4. p. 160., but not of Lapeyr.; Don's Mill., 1. p. 611.

Synonymes. *H. aspalathoides* Pursh, *Fl. Amer. Sept.*, 2. p. 376.

Spec. Char., &c. Stem round, compressed at the top. Branches erect. Leaves dense, without dots,

channelled, with somewhat revolute margins. Sepals equal, erect. Styles joined. (*Don's Mill.*, i.

p. 611.) An undershrub 1 ft. in height, a native of Carolina; introduced in 1811, and producing its yellow flowers in July and August. Somewhat tender.

App. i. *Other Species of Hypéricum.*

The only truly hardy shrubby species of *Hypéricum* are, *H. elátum*, *H. hircínium*, *H. calycinum*, *H. Kalmiánium*, and *H. prolificum*. The other hardy species are of such low growth, that they may be considered, for all practical purposes, as herbaceous plants. The same may be said of a number of the half-hardy species. The number of these might be increased partly by the addition of *H. repens* and *H. linearifólium*, from the south of Europe; by several species from North America, which will be found noticed in p. 179.; and by a few from Africa. *H. oblongifólium*, in the list, p. 173., appears to have been lost; and there are, probably, some other Himalayan species which will prove half-hardy. *H. japonicum* Dec. (*Royle* t. 24. f. 2.) is a plant enjoying a very extended distribution, being found in situations where the snow covers the soil for nearly six months in the year, along the Himalayas, and on the Neelgherries. It is also found in Japan. (*Royle's Illust.*, p. 131.)

GENUS II.



ANDROSÆMUM *Chois.* THE ANDROSÆMUM, or TUTSAN. *Lin. Syst.* Polyadélphia Polyándria.

Identification. *Chois.* Prod. Hyp., 37.; Dec. Prod., 1. p. 543.; Don's Mill., 1. p. 601.

Synonymes. *Hypéricum* L.; *Androsème*, *Fr.*; *Johanniskraut*, *Ger.*

Derivation. From *anēr*, *andros*, a man, and *haima*, blood; the capsules, when crushed between the fingers, giving out a blood-coloured juice. Tutsan is a corruption of *toute saine*, all heal; and it was applied to the plant formerly from its supposed vulnerary properties.

Gen. Char., &c. *Capsule* baccate; usually 1-celled. *Calyx* 5-parted, with unequal lobes. *Petals* 5. *Styles* 3. *Stamens* numerous, disposed in 3 sets. (*Don's Mill.*, i. p. 601.)—An evergreen suffruticose plant, with sessile leaves, and terminal stalked flowers.

1. A. OFFICINALE *Allioni.* The officinal *Androsæmum*, or common *Tutsan*.

Identification. *All. Ped.*, No. 1440.; Dec. Prod., 1. p. 543.; Don's Mill., 1. p. 601.

Synonymes. *Clymenon* *Italdrum L'Obel*; *Hypéricum Androsæmum* *Lin., Willd., Smith, and Hooker*; *Park Leaves* (because it is frequently found wild in parks); *Androsème officinale, Fr.*; *breit-blättriges* (broad-leaved) *Johanniskraut, Ger.*

Engravings. *Blackw.*, t. 94.; *Eng. Bot.*, t. 1225.; and our *fig. 109.*

Spec. Char., &c. Leaves ovate, and somewhat heart-shaped, sessile, widely spreading. Flower an inch wide. A native of moist shady lanes, thickets, and woods in England, in the western part of Scotland, and not unfrequent in woods in Ireland. It was formerly common in the woods about Hampstead and Highgate, till these were grubbed up, and the land where they grew subjected to cultivation. It is also a native of Italy, Greece, and Cacausus. It forms a dense bush, with many stems, attaining the height of 3 ft. and upwards, and producing its large yellow flowers from July to September. The fruit is an ovate capsule, assuming the appearance of a berry: it is, at first, yellowish green, then red or brownish purple; and, lastly, almost black when ripe. The juice of the capsules, and also that of the leaves, is claret-coloured. The latter, when bruised, have an aromatic scent, and were formerly applied to fresh wounds; and hence the French name of *la toute saine*. In gardening, the plant is valuable as growing under the drip of trees, and thriving and flowering freely in almost any soil or situation. Plants, in the London nurseries, may be obtained at 9d. each; and at Bollwyller for 50 cents.

109



CHAP. XXII.

OF THE HARDY LIGNEOUS PLANTS BELONGING TO THE ORDER
ACERACEÆ.

DISTINCTIVE Characteristics. Flowers either unisexual or bisexual. Calyx and corolla equal in the number of their parts, with an imbricated aestivation; the corolla sometimes absent. Petals without appendages. Stamens inserted upon a disk, which arises from below the pistillum, not agreeing in number with the divisions of the calyx and corolla. Pistillum 2-lobed, each lobe having a wing at its back. Style 1. Stigmas 2. Fruit formed of two samaræ, or keys, each containing 1 cell and 1 erect seed. Embryo curved, with leafy shriveled cotyledons and no albumen. Trees or shrubs, almost all deciduous, with opposite leaves, without stipules. (*Pen. Cyc.*) Calyx 4—9 lobes, mostly 5. Stamens mostly 8. Flowers in axillary corymbs. Sap sugary. (*Lindl. Introd. to N. S.*) Cotyledons, in the germination of the seed, produced above ground. The samaræ in *Acer Pseudo-Plátanus* are very rarely 3.

Description. The species are chiefly low and middle-sized deciduous trees, generally with lobed, but, in one or two cases, with entire or pinnated leaves. They are natives of Europe, North America, and the north of India. They are all highly ornamental; some of them valuable for their timber; and from *Acer saccharinum*, and other species, sugar is extracted. In point of magnitude, the species of the *Aceraceæ* may be arranged in three classes: those of the largest size, having large leaves, and the trunks of a timber size, fit for various purposes in architecture, such as the *A. Pseudo-Plátanus*, *A. eriocarpum*, &c.; those of the second size, with small leaves, the timber of which is chiefly used by cabinet-makers, turners, &c., and the trees as copsewood, such as *Acer campêtre*; and those of the third size, with small leaves, which are solely employed for ornamental planting, such as *Acer monspessulanum*, *A. creticum*, &c.

Geography and History. “The maples,” Michaux observes, “form extensive forests in the northern parts of North America: these forests appear, with those of the beech, to succeed the spruce fir, the larch, and the pine, and to precede the chestnut and the oak; at least, this is the case between 43° and 46° of N. lat., the region assigned by nature to the true sugar maple.” Seven species of *Acer*, and one of *Negundo*, are described by Michaux. In the *Pen. Cyc.*, under the article *Acer*, understood to be by Dr. Lindley, 34 species are enumerated or described; and, in *Dou’s Miller* 39 species, of which 26 are in cultivation in British gardens.

Most of the American species are already introduced into Britain; but there are some in the mountainous regions of India, and probably in Japan and China, which are likely to prove hardy in Britain, which are not yet introduced, the names of some of which will be found in p. 173. and p. 176., and in the concluding section of this chapter.

Properties and Uses. The wood of the *Aceraceæ* is moderately hard, compact, and more or less veined: it is useful in various departments of architecture, and is particularly valuable as fuel. Sugar is one of the constituent parts of the sap of all the acers and negundos, though that article is chiefly obtained from two species, which are natives of America.

Soil and Situation. The *Aceraceæ* prefer a free, deep, loamy soil, rich rather than sterile, and neither wet nor very dry. The situation that suits them best is one that is sheltered, and shady rather than exposed. They are seldom found on the north sides of lofty mountains, or on mountains at all, except among other trees; but in the plains they are found by themselves. Though the species only attain perfection in favourable soils and situations, they will spring up and live in any soil or situation whatever.

Propagation and Culture. The *Aceraceæ* are chiefly propagated from seeds; but some sorts are increased by layers, cuttings of the shoots or roots, or by budding or grafting. The seeds of most of the species ripen in October, and

they are gathered by hand, or by shaking the tree, when the keys begin to turn brown. The maturity of the seed may be proved by opening the key, and observing if the cotyledons are green, succulent, and fresh; if the green colour of the cotyledons is wanting, the seeds are good for nothing. The seeds of all the species may either be sown in autumn, after they are gathered, or in spring: and the latter method is preferable where moles abound, as they are very fond of the seeds. Sown in spring, they come up in five or six weeks afterwards, with the exception of those of the *A. campêtre*, which never come up till the second or third year. The seeds should not be covered with more than from a quarter to half an inch of soil. The surface of the ground in which they are sown may be advantageously shaded with leaves, fronds of firs, heath, or straw.

The genera which compose this order are three, *Acer*, *Negundo*, and *Dobinea*; and the species in cultivation in Britain are of the two former genera, which are thus contradistinguished in *Don's Mill.*, i. p. 647.

ACER L. Flowers polygamous. Calyx 5-lobed. Stamens 7—9, rarely 5. Leaves simple, usually lobed.

NEGUNDO Mœnch. Flowers diœcious. Calyx unequally 4—5-toothed. Anthers 4—5, linear, sessile. Leaves pinnate.

GENUS I.



ACER L. THE MAPLE. *Lin. Syst.* Polygãmia Monœc'ia.

Identification. *Lin. Gen.*, No. 1115; *Mœnch. Meth.*, 334; *Dec. Prod.*, l. p. 593; *Don's Mill.*, 1. p. 648.

Synonymes. Erable, *Fr.*; Ahorn, *Ger.*; Acero, *Ital.*; and Arce, *Spanish.*

Derivation. From *acer*, hard or sharp, derived from *ac*, Celtic, a point. The name is supposed to be applied to this genus because the wood of some species is extremely hard, and was formerly much sought after for the purpose of making pikes and lances.

Gen. Char., &c. Sexes hermaphrodite, or monœciously polygamous. Flowers with a calyx and corolla. *Calyx* divided into 5 parts, or some number between 4 and 9. *Petals* the same in number. *Stamens* 8, or some number between 5 and 12. *Anthers* 2-lobed. *Carpels* 2, very rarely 3, each a samara; that is, a fruit which is called, in England, vernacularly, a key.—Leaves lobed and toothed, or, rarely, neither lobed nor toothed. Flowers generally yellow, with more or less of green blended with the yellow; red in *A. rubrum*: not individually conspicuous, but interesting in the kinds that flower at leafing time, from their number, from the rarity of flowers generally at that season, and from the enlivening effect of the numerous bees, and other insects, that attend them. The tips of the wings of the samaræ of several of the species are of a light red, in England, at the end of summer, and in autumn. The species are middle-sized, or low deciduous trees, natives of Europe, North America, and, some, of the Himalayas. They are, in general, quite hardy in Britain, and most of them ripen seeds in this country, by which they are readily propagated. They are among the most ornamental trees of artificial plantations, on account of the great beauty and variety of their foliage, which changes to a fine scarlet, or rich yellow, in autumn. The larger-growing species are often many years before they come into flower, and, after they do so, they sometimes flower several years before they mature seeds; probably from having the flowers of only one sex. In general it may be observed that there is great uncertainty, in the different species of *Acer*, with regard to sex.

A. *Leaves simple.*

‡ 1. *A. OBLO'NGUM* Wall. The oblong-leaved Maple.

Identification. Wall. in *Litt.*; *Dec. Prod.*, l. p. 593; *Don's Mill.* l. p. 648.

Synonymes. *A. laurifolium* D. Don, *Prod. Fl. Ncp.*, p. 249; *A. Buximp'ata* Hamilt. *Engraving.* Our *fig.* 113. in p. 433.

Spec. Char., &c. Leaves oblong-lanceolate, acuminate, quite entire, coriaceous, smooth, rounded at the base. Racemes compound; wings of fruit parallel, smooth, separated. (*Don's Mill.*, i. p. 648.) Native of Nepal, where it flowers in February; introduced in 1824. Height 20 ft. This species is rather tender, and somewhat difficult to keep in the open ground. We are not aware of any living plant of it being in the neighbourhood of London; but there is one in the arboretum of John Thomas Brooks, Esq., at Flitwick House, Bedfordshire, which is frequently killed down to the ground during winter, but always shoots up vigorously the following spring. The leaves and general appearance of the shoots resemble those of a euca-lyptus; hence its character among maples is so very extraordinary, that to the botanist it must be a species of very great interest.

‡ 2. *A. TATARICUM* L. The Tartarian Maple.

Identification. Lam. Sp., 1495; Dec. Prod., 1. p. 593; Hayne Dend., p. 509; Don's Mill., 1. p. 648. *Synonymes.* Érabie de Tartarie, Fr.; Tartarische Ahorn, Ger.; Zarza-modon, or Locust Tree, Russ. *Engravings.* Pall. Fl. Rus., t. 3; Tratt. Arch., 1. No. 13; Wat. Dend. Brit., t. 160; our fig. 114. in p. 134; and the plate of this species in our Second Volume.

Spec. Char. Leaves cordate, undivided, serrated, with obsolete lobes. Racemes compound, crowded, erect; wings of fruit parallel, young ones puberulous. (*Don's Mill.*, i. p. 648.) A low tree, native of Tartary, introduced in 1759; flowering in May and June, and growing to the height of from 20 ft. to 30 ft.

Description, &c. The branches are numerous, and disposed into a compact head, densely covered with leaves, which are distinguished by a peculiarly veiny appearance, and lively green. The flowers are of a pale greenish yellow, sometimes slightly tinged with red, as are the fruit, or keys, before their maturity. When raised from seed, the plant will come into flower in 5 or 6 years; and, in good soil, it will attain the height of 15 ft. in 10 years.

Geography, History, &c. This species is common throughout all the south of European Russia; but it is not found on the Ural Mountains, or on Caucasus. Near the Wolga and its tributary streams, it forms a hemispherical tree, about 20 ft. in height, and the same in diameter. In New Russia, it attains the size of *Acer campêtre*. (*Pall.*) According to some, this species will thrive in a moister soil than most others. The wood is hard; and, being of whitish colour veined with brown, it may be used for cabinet-work. In ornamental plantations, this species is valuable on account of the early expansion of its leaves, which appear before those of almost every other kind of *Acer*. Pallas informs us, that the Calmucks, after depriving the keys of their wings, boil them in water, and afterwards use them for food, mixed up with milk and butter. In Britain, it is planted solely as an ornamental tree or bush.

Statistics. The largest specimen of it in the neighbourhood of London is at Syon, where it has attained the height of 25 ft. In Devonshire, at Endsleigh Cottage, 18 years planted, it is 40 ft. high. In Sussex, at West Dean, 15 years planted, it is 19 ft. high. In Staffordshire, at Trentham, 16 ft. high, with a head 20 ft. in diameter. In Worcestershire, at Croome, 30 years planted, and 30 ft. high. In Scotland, in the Perth Nursery, 14 years planted and 10 ft. high. Price, in London, 1s.; and at Bollwyller, 1 franc each.

B. *Leaves 3-lobed, or trifid; rarely 5-lobed.*

‡ 3. *A. SPICATUM* L. The spiked-flowered Maple.

Identification. Lam. Dict., 2. p. 381; Dec. Prod., 1. p. 593; Don's Mill., 1. p. 648. *Synonymes.* *A. montanum* *Al. Hort. Kew.*, 3. p. 17; *A. pennsylvanicum* *Du Roi, Hartek.*, t. 2; *A. parviflorum* *Ehrh.*; Mountain Maple, Érabie de Montagne, Fr.; Berg Ahorn, Ger. *Engravings.* Tratt. Arch., No. 13; our fig. 115. in p. 135; and the plate of this species in our Second Volume.

Spec. Char., &c. Leaves cordate, 3- or slightly 5-lobed, acuminate, pubescent beneath, unequally and coarsely serrated. Racemes compound, erect. Petals linear. Fruit smooth, with the wings rather diverging. (*Don's*

Mill., i. p. 648.) Flowers polygamous. A deciduous tree, a native of the mountains of Canada, and of the Alleghany Mountains; producing its very small greenish flowers in April and May, and attaining in its native country, according to Michaux, the height of 6 ft. or 8 ft. Introduced in 1750, by Archibald Duke of Argyle, and about as common in ornamental plantations in England as *A'cer tataricum*. In British gardens, it forms a low tree, 8 ft. or 10 ft. high, very ornamental in autumn, from its small keys, which are fixed upon slender pendulous spikes, and have their membranous wings, beautifully tinged with red when ripe. Michaux states that this species, grafted upon the sycamore, is, like the *A'cer striatum*, augmented to twice its natural dimensions; a fact which we have never had an opportunity of seeing verified.

Statistics. At Syon, 25 ft. high. In Worcestershire, at Croome, 30 years planted and 40 ft. high, the trunk 15 in. in diameter, and the diameter of the head 20 ft. In Scotland, at Edinburgh, in the Caledonian Horticultural Society's Garden, 9 years planted and 30 ft. high. Price, in London, 1s. 6d. a plant; at Bollywyler, 1 franc 50 cents; at New York, 25 cents, and seeds 1 dollar per quart.

‡ 4. *A. STRIA'TUM* L. The striped-barked Maple.

Identification. Lam. Dict., 2. p. 381; Dec. Prod., 1. p. 593; Don's Mill., 1 p. 648.

Synonymes. *A. pennsylvanicum* *Lin. Sp.*, 1496; *A. canadense* *Marsh.*, and *Dub. Arb.*, 1. t. 12.; Snake-barked Maple, Moose Wood, striped Maple; *E'rable jaspé*, *Fr.*; gestreifter Ahorn, *Ger.*

Engravings. Mill. t. 7.; *Trat. Arch.*, No. 11.; *Mich. Fel. Arb.*, 2. t. 17.; our *fig.* 116. in p. 436, 437.; and the plate of this species in our Second Volume.

Spec. Char. Leaves cordate, 3-lobed, acuminate, finely and acutely serrated. Racemes pendulous, simple. Petals oval. Fruit smooth, with the wings rather diverging. (*Don's Mill.*, i. p. 643.) A tree readily distinguished by the striped bark of the young shoots, growing in its native country to the height of 10 ft. or 12 ft., but to that of 20 ft. or upwards in a state of cultivation. It produces its flowers in May and June, and sometimes ripens seeds.

Description, &c. The trunk and branches are covered with a smooth green bark, longitudinally marked with black and white stripes, by which the tree is readily distinguished at all seasons of the year. In America, it is one of the first trees that announces the approach of spring. Its buds and leaves, when beginning to unfold, are rose-coloured. The leaves are of a thick texture, and finely serrated. The flowers are greenish, and are grouped on long peduncles. The fruit is remarkable for a cavity on one side of the capsules. It is a native of North America, in Nova Scotia, and from Canada to Carolina. It makes its first appearance in about latitude 47°, and is particularly abundant in Nova Scotia, the State of Maine, and New Hampshire. In approaching the Hudson, it becomes more rare; and, beyond this boundary, it is confined to the mountainous tracts of the Alleghanies, in which it is found in cold shaded exposures, along the whole range to its termination in Georgia. In many of the forests of Maine and New Hampshire, *A. striatum* constitutes a great part of the undergrowth, seldom exceeding 10 ft. in height; but, where it is not shaded by other trees, it attains the height of 20 ft. or upwards. The wood is white and fine-grained, and used by cabinet-makers as a substitute for holly. Cattle, in Nova Scotia, are fed with the leaves, both in a green and dried state; and in spring, when the buds begin to swell, both horses and cattle are turned into the woods to browse on the young shoots, which they consume with avidity. (*Michaux.*) From the great beauty of its bark, this tree deserves a place in every collection. It is propagated by seeds, which are received from America, or by grafting on *A. Pseudo-Platanus*.

Statistics. The largest specimen which we know of within a short distance of London, is at Mr. Needham's villa, near Maidenhead, where it has attained the height of 16 ft. 6 in. in 20 years. Near Reading, at White Knights, a tree 25 years planted is 21 ft. high; in Surrey, at Farnham Castle, 35 years planted, it

is 16 ft. high; in Yorkshire, at Ripley Castle, 11 years planted, it is 15 ft. high. In Scotland, in the garden of the Caledonian Horticultural Society, 9 years planted, it is 8 ft. high. In Ireland, at Oriel Temple, 35 years planted, it is 27 ft. high. Price of plants, in the London Nurseries, 1s. 6d. each; at Bollyweller, 1 franc 50 cents; and at New York, 25 cents a plant, and 1 dollar and 50 cents for a quart of seed.

C. *Leaves 5-lobed.*

Υ 5. *A. MACROPHYLLUM Pursh.* The long, or large, leaved Maple.

Identification. Pursh, Fl. Amer. Sept., 1. p. 267.; Dec. Prod., 1 p. 594.; Don's Mill., 1 p. 649.

Engravings. Hook. Fl. Bor. Amer., 1. t. 38.; our figs. 117. in p. 438, 439., and 118. in p. 440, 441.; and the plate of this species in our Second Volume.

Spec. Char., &c. Leaves digitately 5-lobed, with roundish recesses. Lobes somewhat 3-lobed, repandly toothed, pubescent beneath, racemes compound, erect. Stamens 9, with hairy filaments. Ovaries very hairy. (*Don's Mill.*, i. p. 648.) The leaves vary much in size, and also in the manner in which they are lobed. Those of the dried specimens sent home by Mr. Douglas, are cut nearly to the base, so as almost to merit the appellation of palmate, while those of young plants in the London Horticultural Society's Garden, and at Messrs. Loddiges's, are not more deeply cut than those of *A. platanoides*, as may be seen by our figs. in p. 440, 441., taken from leaves gathered in these gardens.

Description, &c. A tree of the largest size, a native of the north-west coast of North America, and introduced into England in 1812; where, however, it has not yet flowered. In its native country it is found exclusively in woody mountainous regions along the sea coast, between 40° and 50° N. lat., and on the great rapids of the Columbia. This noble tree, Dr. Hooker observes, was unquestionably discovered by Mr. Menzies, the first naturalist who visited the coast where it grows. Mr. Douglas, who subsequently found it, and sent dried specimens and seeds to the London Horticultural Society, observes, "It is one of the most graceful of trees in the country it inhabits, varying from 40 ft. to 90 ft in height, and from 6 ft to 16 ft. in the circumference of its trunk. The branches are widely spreading, the bark rough and brown, the wood soft, but beautifully veined. It contains, perhaps, as much sap as any species, except *A. saccharinum*; but the sap is not used for making sugar by the natives. The flowers are yellow, and very fragrant, appearing in April and May. Mr. Douglas prophetically adds, "It will, at some future time, constitute one of our most ornamental forest trees in England." (*Hooker's Fl. Bor. Amer.*, vol. i. p. 112.) Specimens of the timber, which were sent home by Mr. Douglas, exhibit a grain scarcely inferior in beauty to the finest satin wood. The largest specimen of the tree is in the garden of the London Horticultural Society; where, in 1835, it had attained the height of 25 ft. It is propagated by layers in the garden of the Society; and at Messrs. Loddiges's, and the annual shoots from them are often from 6 ft. to 10 ft. in length; so that there can be no doubt of the tree being one of very rapid growth. This magnificent species cannot be too warmly recommended to the attention of planters, as it is perfectly hardy, and well suited for general cultivation, both in useful and ornamental plantations, throughout Europe. Plants, in London, cost 2s. 6d. each; and when the tree has once ripened seeds in Britain, plants will be much cheaper.

Υ 6. *A. PLATANOIDES L.* The Platamis-like, or Norway, Maple.

Identification. Lin. Sp., 1496.; Dec. Prod., i. p. 649.; Don's Mill., 1 p. 649.

Synonymes. Erable plane, or Erable de Norvège, Fr.; spitz Ahorn, or spitz-blättriger Ahorn, Ger.

Engravings. Dub. Arb., 1. t. 10. f. 1.; Tratt. Arch., 1. t. 4.; Mill. Ic., t. 8. f. 1.; and our fig. 119. in p. 112., 443.

Spec. Char., &c. Leaves cordate, smooth, 5-lobed. Lobes acuminate, with a few coarse acute teeth. Corymbs stalked, erectish, and, as well as the fruit, smooth; fruit with divaricated wings. (*Don's Mill.*, i. p. 649.) A middle-

sized deciduous tree, a native of Europe, from Norway to Switzerland, and also of North America, but not of Britain, flowering in May and June. Introduced in 1683. Height from 40 ft. to 70 ft.

Description. A handsome tree of the first rank; in general appearance, at a distance, like the common sycamore; but, on a nearer approach, the leaves are found of a smoother and finer texture. The roots extend considerably both downwards and laterally. The trunk is somewhat shorter than that of the sycamore, seldom exceeding 60 ft. or 70 ft. in height. The bark is green on the young shoots, but it afterwards becomes of a reddish brown, dotted with white points: that of the trunk is brown, and rather cracked. The buds are large and red in autumn, becoming of a still darker red in the course of the winter: those on the points of the shoots are always the largest. The leaves are thin, green on both sides, and shining. When the petiole is broken an acrid milky sap issues from it, which coagulates with the air. The leaves are about 5 in. long, and nearly the same in width. The petioles are longer than the leaves. About the end of October, the leaves become either of a clear or a yellowish red, and then drop off. The flowers appear just before the leaves, near the end of April: they form a short raceme, somewhat corymbose. The fruits, or keys, have their wings yellow. They ripen in September and October; and it is not till the tree has attained the age of nearly 40 years that it produces fertile seeds, though it will flower many years before that period. The rate of growth of this species is considerable. In France, a plant has been known to attain the height of 12 ft. in three years from the seed. In England, when once established, it produces shoots from 18 in. to 3 ft. long every year, till it attains the height of 20 ft. or 30 ft.; which, in favourable situations, it does in 10 years.

Varieties.

‡ A. p. 2 *Lobèlii*. *Lobel's Platanus-like Maple*.

Synonyms. *A. Lobèlii* Tenore; *A. platanoides* Don's Mill., 1. p. 649.

Engraving. Our fig. 120. in p. 444.

Description. The leaves are very slightly heart-shaped, irregularly toothed, 5-lobed, with the lobes more or less abruptly pointed. The bark of the young wood striped, somewhat in the manner of that of *A. striatum*; by which circumstance the plant, in a young state, is readily distinguished from *A. platanoides*. A large tree, native of the kingdom of Naples, and found on mountains. The general appearance is said to be that of *A. platanoides*, of which it seems to us to be only a variety. We have seen small plants of this sort in the arboretum of Messrs. Loddiges, and in one or two of the nurseries. These were imported from Messrs. Booth, nurserymen, Hamburg. They appear to be grafted on *A. platanoides*.

‡ A. p. 3 *pubescens* Hayne. *The downy-leaved Platanus-like Maple*.—Leaves downy on the under side. This variety appears to be found in Germany; but we have not seen it in England.

‡ A. p. 4 *variegatum* Hort., *albo variegatum* Hayne. *The silvery variegated-leaved Platanus-like Maple*.—According to the figure in Schmidt's *Baumzucht*, the foliage of this variety is beautifully marked, and very handsome; but we have never seen it in Britain in a state to warrant us in recommending it for cultivation. From several specimens which we have seen, we consider it as decidedly inferior in beauty to the variegated sycamore.

‡ A. p. 5 *aurco variegatum*, the golden variegated-leaved *Platanus-like Maple*, is described in books, but we have never seen a plant of it.

‡ A. p. 6 *laciniatum* Dec. *The cut-leaved Platanus-like Maple*. (fig 121. in p. 445.)—A very distinct variety, with the leaves deeply and variously cut. It is frequently produced from seed, being found by nurserymen among seedlings of the species. In 1835, there were above 100 of them, in two beds of one year's seedlings, in the Goldworth Nursery. *A. p. crispum* Lanth seems to be nothing more than

a synonyme of this variety; which, in the nurseries, is sometimes called the eagle's claw, or hawk's foot, maple.

Geography. A native of Europe, from the west coast of Norway to Switzerland, and from France to the eastern boundary of European Russia. Pallas says, it does not occur beyond the Ural Mountains, or in Siberia, but that it is common through all the woods of Russia. We observed it in 1814, in all the woods bordering the public road from Wilna to Mittan, and from Moscow to Galicia. Next to the birch and the trembling poplar, it seemed to us the most abundant tree in the Russian woods. In the north, according to Pallas, it forms a stunted bush; but in the Ukraine it is a lofty tree.

History. This species is recorded as having been first cultivated in Britain in the Edinburgh Botanic Garden, by Mr. James Sutherland. It has since been very generally propagated in Britain, and is now to be found in most ornamental plantations made since the days of Miller. The largest trees in the neighbourhood of London are at Purser's Cross and Syon, where they have attained the height of nearly 60 ft.

Properties and Uses. The wood, in its young state, is white; but at a more advanced age it becomes grey. In a dry state, it weighs 43 lb. 4 oz. per cubic foot. It is easily worked, takes a fine polish, and absorbs and retains all kinds of colours. By drying it only loses a 24th part of its weight. It is used for all the various purposes of the wood of the common sycamore. From the sap, sugar has been made in Norway, Sweden, and in Lithuania. The German foresters have found that this sap is produced in less abundance than that of the sugar maple, or of the common sycamore; but that it contains more saccharine matter than the sap of the latter species. Some chemists have found that, after boring a hole at the base of the trunk, 35 quarts of sap have been produced in 8 days; and that 95 lb. of this sap have, by evaporation, given 4 lb. of syrup; and that from 80 lb. to 100 lb. of this syrup have given from 4 lb. to 6 lb. of crystallised sugar. After a great quantity of this sap has been drawn off, it begins to get thicker, muddy, and yellow in appearance, bitter in taste, and not productive of syrup.

Soil and Situation. To attain a considerable size, the tree ought to be planted in a free, deep, rich soil, not surcharged with moisture; and the situation ought to be low rather than high. It thrives remarkably well on the sea shore on the Baltic, and along the west coast of Norway, and the west coast of Scotland.

Propagation and Culture. After the tree has attained a considerable size and age, it produces abundance of fertile seeds in England. It does so at Purser's Cross, at Syon, and various other places. The varieties are propagated by grafting or layering. The seeds, as soon as they are gathered, should be either immediately sown, or mixed with sand or earth, and kept moderately dry till spring. In either case they come up the first year.

Accidents and Diseases. The leaves of this species, in common with those of *A. Pseudo-Platanus*, and perhaps most of the other species of *Acer*, are subject to what is commonly called the honey dew, which, from its clamminess in the neighbourhood of the smoke of mineral coal, is apt to attract and retain the particles of soot which are continually floating in the air. In consequence of insects resorting to these leaves, they are frequently blackened with their excrements. In some parts of France this honey dew is called manna. M. Tschoudi says that the manna is produced by the extravasated sap; and that the bees are so fond of it, that it would be worth while to plant the tree in the neighbourhood of places where hives are kept. According to others, the bitterness of the matter of the leaves prevents them from being attacked by insects.

Statistics. In the neighbourhood of London, the largest tree of *A. platanoides* is at Kew, where, in 70 years, it has attained the height of 76 ft.; at Syon, it has attained the height of 64 ft., with a trunk 24 ft. in diameter, and the diameter of the head 64 ft.; at Kenwood there is a tree, 35 years planted, 47 ft. high. In Surrey, at Bagshot Park, a tree, 14 years planted, has attained the height of 25 ft., with a trunk 16 in. in diameter. In Sussex, at West Dean, a tree of the cut-leaved variety, 9 years planted, has attained the height of 26 ft. In Lancashire, at Latham House, a tree, 40 years planted, is 45 ft. high. In Staffordshire, at Teddesley, a tree, 14 years planted, is 29 ft. high. In

Worcestershire, at Croome, a tree, 35 years planted, is 40 ft. high. In Yorkshire, at Grimstone, a tree, 13 years planted, is 30 ft. high. In Scotland, in Haddingtonshire, at Tynningham, there is a tree 42 ft. high. In Clackmannanshire, in the garden of the Dollar Institution, one 7 years planted is 14 ft. high. In Perthshire, at Taymouth, one 50 years planted is 50 ft. high; the diameter of the head 51 feet. According to Dr. Walker, this tree has attained a large size in the Island of Bute, at Bargally, and at various other places on the sea coast of Scotland. In Ireland, in King's county, at Charleville Forest, a tree, 60 years planted, is 78 ft. high, with a trunk 3 ft. 8 in. at 1 ft. from the ground. In France, in the neighbourhood of Paris, the tree attains the height of 60 ft. In Germany, in Hannover, at Schwöbber, it has attained the height of 80 ft. In Saxony, at Wörlitz, 40 ft. In the neighbourhood of Vienna, from 50 ft. to 60 ft. In Russia, where the tree is very common, it often exceeds the height of 40 ft., south of Kiow; but north of Moscow it is seldom above 30 ft. In Sweden, on the north-west coast, exposed to the sea breeze, it grows to the height of between 30 ft. and 40 ft.; as it does about Lund, and at different places on both shores of the Baltic.

Commercial Statistics. This tree is very generally propagated in European nurseries. In London, plants, 1 ft. high, cost 30s. a 1000; and 3 ft. high, 50s.; at Bollwyller, 20 cents each, or 40 francs a 1000; at New York, ?.

Æ 7. *A. SACCHARINUM L.* The Sugar Maple.

Identification. Lin. Sp., 1496; Hayne, Dend., p. 214; Dec. Profl., 1. p. 595.; Don's Mill., 1. p. 650.

Synonymes. Rock Maple, Hard Maple, Bird's-eye Maple, Amer.

Engravings. Michx. Fl. Arb., 2. t. 15.; Tratt. Arch., 1. No. 3.; our fig. 122. in p. 446, 447.; and the plate of this species in our Second Volume.

Variety. *A. s. 2 nigrum*; the *A. nigrum* of Michaux, De Candolle, and G. Don; the black Sugar Tree, or Rock Maple, figured in Michx. Arb., 2. t. 16. has the leaves resembling those of *Acer saccharinum*, but much darker. According to Michaux, the leaves are 5 in. or 6 in. long, and "exhibit, in every respect, nearly the same conformation as those of the true sugar maple. "They differ from it," he says, "chiefly in being of a darker green, and of a thicker texture; and in being somewhat more bluntly lobed. The tree is indiscriminately mixed with the common sugar maple through extensive ranges of country in New Hampshire, Vermont, and Connecticut; but is readily distinguished from it by the smaller size which it attains, and the darker colour of its leaves." The soil in which it flourishes best is a rich, strong, sandy loam; and there it usually grows to the height of 40 ft. or 50 ft. Trees of this alleged species were introduced into England in 1812; and there are plants bearing the name in the garden of the London Horticultural Society, and in the garden of the Messrs. Loddiges, and to us they have always appeared to be merely varieties of *A. saccharinum*, differing in nothing but in having the foliage somewhat darker. Plants, in the London nurseries, cost 1s. 6d., and seeds 4s. per ounce; at Bollwyller, 1 franc 50 cents a plant; and at New York, 50 cents a plant, and seeds 1 dollar per quart.

Spec. Char., &c. Leaves cordate, smooth, glaucous beneath, palmately 5-lobed; lobes acuminate, serrately toothed. Corymbs drooping, on short peduncles. Pedicels pilose. Fruit smooth with the wings diverging. (*Don's Mill.*, i. p. 650.) A deciduous tree, from North America, growing, in England, to the height of 40 ft. or 50 ft., and flowering in April and May. Introduced in 1735.

Description. In America, the sugar maple sometimes reaches the height of 70 ft. or 80 ft., with a proportionate diameter; but it does not commonly exceed 50 ft. or 60 ft., with a diameter of 12 in. or 18 in. Well-grown thriving trees are beautiful in their appearance, and easily distinguished by the whiteness of their bark. The leaves are about 5 in. broad; but they vary in length according to the age and vigour of the tree. They are opposite, attached by long petioles, palmated or unequally divided into 5 lobes, entire at the edges, of a bright green above, and glaucous or whitish underneath. In autumn, they turn reddish with the first frosts. Except in the colour of the under surface, they nearly resemble the leaves of the Norway maple. The flowers are small, yellowish, and suspended by slender drooping peduncles. The seed is contained in two capsules united at the base, and terminating in a membranous wing. It is ripe near New York in the beginning of October, though the capsules attain their full size six weeks earlier. Externally, they appear equally perfect; but Michaux informs us that he constantly found one of them empty; and the fruit is matured only once in two or three years. (*Michaux*, p. 225.) The wood, when cut, is white; but, after being wrought and exposed some time to the light, it takes a rosy tinge. Its grain is fine and close, and, when polished, it has a silky lustre. It is very strong, and sufficiently heavy, but wants the property of durability, for which the chestnut and the oak are so highly esteemed. When exposed to moisture it soon decays; and for this reason it is neglected in civil and naval architecture. (*Michaux*, p. 225, 226.) The buds of this species, like those of *Acer Pseudo-Platanus*, of which it may be considered the American representative, have a fine ruddy tint early in spring, before they begin to expand.

Geography. According to the elder Michaux's researches, the sugar maple begins a little north of Lake St. John, in Canada, near 48° of N. lat. which, in the rigour of its winter, corresponds to 68° of Europe. It is nowhere more abundant than between 46° and 43° of N. lat.; which space comprises Canada, New Brunswick, Nova Scotia, the states of Vermont and New Hampshire, and the district of Maine: in these regions, it enters largely into the composition of the forests, with which they are still covered. Farther south, it is common only in Genessee in the state of New York, and in the upper parts of Pennsylvania. It is estimated by Dr. Rush, that, in the northern parts of these two states, there are 10,000,000 of acres which produce these trees in the proportion of thirty to an acre. In the lower parts of Virginia, of the Carolinas, and of Georgia, and likewise in the Mississippi territory, this tree is unknown, or very rare. It is rapidly disappearing from the forests about New York and Philadelphia, where it is no longer tapped for sugar, but is felled for fuel and for other purposes.

The sugar maple covers a greater extent of the American soil than any other species of this genus. It flourishes most in mountainous places, where the soil, though fertile, is cold and humid. Besides the parts already mentioned, it is found along the whole chain of the Alleghanies to its termination in Georgia, and on the steep and shady banks of the rivers which rise in these mountains. (*Michaux*, 225.)

Properties and Uses. In America, in Vermont, New Hampshire, the district of Maine, and farther north, where the oak is not plentiful, the timber of the sugar maple is substituted for it, in preference to that of the beech, the birch, or the elm. When perfectly seasoned, which requires two or three years, it is used by wheelwrights for axle trees and spokes, and for similar purposes. It is also employed, as well as the red-flowered maple, in the manufacture of Windsor chairs. In the country, where the houses are wholly of wood, sugar maple timber is used for the framework; and in the district of Maine it is preferred to the beech for the keels of vessels, as it furnishes longer pieces: with the beech and the yellow pine it forms, also, the lower frame of vessels, which is always in the water. The wood exhibits two accidental forms in the arrangement of the fibre, of which cabinet-makers take advantage for making beautiful articles of furniture. The first consists in undulations like those of the curled maple (*A. rubrum*, see p. 426.) the second, which takes place only in old trees that are still sound, and which appears to arise from an inflexion of the fibre from the circumference towards the centre, produces spots of half a line in diameter, sometimes contiguous, and sometimes several lines apart. The more numerous the spots, the more beautiful and the more esteemed is the wood. This variety is called bird's-eye maple. Like the curled maple, it is used for inlaying mahogany. Bedsteads are made of it, and portable writing-desks, which are elegant and highly prized. To obtain the finest effect, the log should be sawn in a direction as nearly as possible parallel to the concentric circles. When cut at the proper season, the sugar maple forms excellent fuel. It is exported from the district of Maine, for the consumption of Boston, and is equally esteemed for that purpose with the hickory.

The ashes of the sugar maple are rich in the alkaline principle, and it may be confidently asserted, that they furnish four fifths of the potash exported to Europe from Boston and New York. In the forges of Vermont and the district of the Maine, the charcoal of this wood is preferred to any other, and it is said to be one fifth heavier than that made from the same species in the middle and southern states; a fact which sufficiently evinces that this maple acquires its characteristic properties in perfection only in northern climates.

The wood of the sugar maple is easily distinguished from that of the red-flowered maple, which it resembles in appearance, by its weight and hardness. There is, besides, a very simple and certain test: a few drops of sulphate of iron being poured on samples of the different species, the sugar

maple turns greenish, and the white maple and the red-flowered maple change to a deep blue.

The extraction of sugar from the maple is a valuable resource in a new country abounding in forests of that tree, and without much foreign commerce; but it is evident that this mode of obtaining sugar is only destined for a certain stage in the progress of society, and must, in America, very shortly give way to the sugar of commerce, produced by the cane; for this reason, we shall give but a short account of the process of manufacturing maple sugar, and this chiefly as a matter of historical interest, rather than of practical utility. In America, wherever there are canals and railroads, the making of maple sugar must soon become an unprofitable occupation.

The process of making maple sugar is commonly begun in February, or in the beginning of March, while the cold continues intense, and the ground is still covered with snow. The sap begins to be in motion at this season, two months before the general revival of vegetation. In a central situation, lying convenient to the trees from which the sap is to be drawn, a shed is constructed, called a sugar camp, which is destined to shelter the boilers, and the persons who tend them, from the weather. An auger $\frac{3}{4}$ of an inch in diameter; small troughs to receive the sap; tubes of elder or sumach, 8 in. or 10 in. long, corresponding in size to the auger, and laid open for a part of their length; buckets for emptying the troughs and conveying the sap to the camp; boilers of 15 or 18 gallons' capacity; moulds to receive the syrup when reduced to a proper consistency for being formed into cakes; and, lastly, axes to cut and split the fuel, are the principal utensils employed in the operation. The trees are perforated in an obliquely ascending direction, 18 in. or 20 in. from the ground, with two holes 4 in. or 5 in. apart. Care should be taken that the augers do not enter more than half an inch within the wood, as experience has shown that the most abundant flow of sap takes place at this depth. It is also recommended to insert the tubes on the south side of the tree. The troughs, which contain 2 or 3 gallons, are made, in the northern states, of the white pine, of white or black oak, or of maple; but on the Ohio, the mulberry, which is very abundant, is preferred. The chestnut, the black walnut, and the butternut should be rejected, as they impart to the liquid the colouring matter and bitter principle with which they are impregnated. A trough is placed on the ground, at the foot of each tree; and the sap is every day collected and temporarily poured into casks, from which it is drawn out to fill the boilers. The evaporation is kept up by a brisk fire; and the scum is carefully taken off during this part of the process. Fresh sap is added from time to time; and the heat is maintained till the liquid is reduced to a syrup; after which it is left to cool, and then strained through a blanket or other woollen stuff, to separate it from the remaining impurities, when it is poured into the moulds. The boilers are only half filled; and a steady heat is kept up till the liquor is reduced to the proper consistency for being poured into the moulds. The evaporation is known to have proceeded far enough, when, upon rubbing a drop of the syrup between the fingers, it is perceived to be granular. If it is in danger of boiling over, a bit of lard or of butter is thrown into it, which instantly calms the ebullition. When refined, this sugar equals in beauty the finest consumed in Europe; but it is used only in the districts where it is made, and there only in the country places, as, from prejudice or taste, Michaux observes, imported sugar is used in all the small towns, and in the inns.

The sap continues to flow for six weeks; after which it becomes less abundant, less rich in saccharine matter, and sometimes even incapable of crystallisation. In this case it is consumed in the state of molasses, or exposed for three or four days to the sun; when it is converted into vinegar by the acetous fermentation: a kind of beer is also made of it. The amount of sugar produced by each tree in a year varies from different causes. A cold and dry winter renders the trees more productive than a changeable and humid season. It is observed, that, when a frosty night is followed by a dry and

brilliant day, the sap flows abundantly; and two or three gallons of sap are sometimes yielded by a single tree in twenty-four hours. The yearly product varies from 2 lb. to 4 lb. of sugar each tree. Trees which grow in low and moist places afford a greater quantity of sap than those which occupy rising grounds; but it is less rich in the saccharine principle. That of insulated trees, left standing in the middle of fields, or by the side of fences, is the best. It is also remarked, that, in districts which have been cleared of other trees, and even of the less vigorous sugar maples, the product of the remainder is proportionably greater.

Wild and domestic animals are inordinately fond of maple juice, and break through their enclosures to sate themselves with it. (*Michaux*, p. 236.) In Europe, it is not likely that the extraction of sugar from any species of maple will ever be tried otherwise than as a matter of curiosity. Count Wingersky is said to have planted a great many trees of *A. saccharinum* on his estates in Moravia, and to have drawn off the sap from them at the age of 25 years, in order to make sugar. He succeeded in procuring a very good sugar; but, in consequence of drawing sap from the trees every year, they became sickly, and soon afterwards died.

Soil and Situation, Propagation, &c. The same soil may be recommended as for *A. platanoides*; but, as the species is considerably more tender, it requires a more sheltered situation. In British nurseries, it is always raised from American seeds.

Statistics. The largest tree in the neighbourhood of London is at Purser's Cross, where it has attained the height of 45 ft. In Berkshire, at 11th Clere, 6 years planted, it is 24 ft. high; at White Knights, 25 years planted, 21 ft. high; in Herefordshire, at East Ior Castle, 14 years planted, 30 ft. high; in Cumberland, at Ponsonby Hall, 20 years planted, and 24 ft. high; in Cheshire, at Kinnel Park, 20 years planted, and 24 ft. high; in Staffordshire, at Trentham, 56 years planted, and 35 ft. high; in Yorkshire, at Cannon Hall, 42 ft. high; in Hertfordshire, at Cheshunt, 6 years planted, and 18 ft. high; in Edinburghshire, at Dalhousie Castle, 7 years planted, and 9 ft. high; in Banffshire, at Gordon Castle, 33 ft. high. In France, in the Botanic Garden at Toulon, 36 ft. high. In Saxony, at Worlitz, 60 years planted, and 50 ft. high. In Austria, at Kopenzel, near Vienna, 6 years planted, and 14 ft. high. In Bavaria, in the Botanic Garden at Munich, 20 years planted, and 8 ft. high. In Cassel, at Wilhelmshoe, 24 years planted, and 30 ft. high. Some of the largest sugar maples in America, according to Mr. Douglas, are on Goat Island, at the Falls of Niagara.

Commercial Statistics. In the London nurseries, plants cost 2s. each, and seeds 2s. per ounce; at Bollwyller, plants are 1 franc 50 cents each; and at New York, 15 cents a plant, and seeds 2 dollars 25 cents per pound.

Y 8. *A. PSEUDO-PLATANUS L.* The Mock Plane Tree, the *Sycamore*, or *Great Maple*.

Identification. Lin. Sp., 1169; Don's Mill., i. p. 648.

Synonyms. Plane Tree, *Scotch*; *Erbule Sycamore*, *Erbale blanc de Montagne*, *fausse Platane*, or *grand Erable*, *Fr.*; *Ehrenbaum*, *weisser Ahorn*, or *cincine Ahorn*, *Ger.*

Engravings. Dub. Arb., t. 1. 35; Traut. Arb., l. No. 2; Willd. Ab., t. 213; Krause, t. 121; our fig. 123, in p. 183, 449; and the plate of this species in our second Volume.

Spec. Char. &c. Leaves cordate, smooth, with 5 acuminate, unequally toothed lobes. Racemes pendulous, rather compound, with the rachis, as well as the filaments of stamens, hairy. Fruit smooth, with the wings rather diverging. (*Don's Mill.*, i. p. 648.) A deciduous tree, native of Europe, flowering in May and June. Height from 30 ft. to 60 ft. Sexes mostly hermaphrodite.

Varieties.

Y A. P. 2 *flavovariegata.* The *yellow variegated Sycamore*, or *Costorphine Plane* with leaves variegated with yellow. — The original tree stands near an old pigeon-house in the grounds of Sir Thomas Dick Lauder, Bart., in the parish of Costorphine, near Edinburgh. Seeds of this variety, sown, have produced plants with green leaves; but in some of the plants the footstalks of the leaves were of a yellowish green colour, and this colour was partaken of by the leaf; in the other plants the petioles were strongly tinged with red, and the leaves were of a darker green than those of the first-mentioned plants.

Y A. P. 3 *albo variegata* Hayne. The *white variegated-leaved Sycamore*. —

Leaves blotched with white. This variety is much more common than the other. Tschoudi says of it, that it is one of the finest trees that can be seen; and that, in the beginning of summer, it is delightful to stand under it, and look through the leaves to the sun. At a short distance, he adds, the leaves are as beautiful as flowers. In Britain, however, like the leaves of most other variegated deciduous trees, they soon become ragged, and lose, in autumn, by dying off of a dirty colour and diseased appearance, what they have gained by their whiteness and transparency in spring. Of all the variegated varieties of *Acer*, however, it must be acknowledged that this variety is to be considered the most ornamental.

‡ A. *P. 4 purpurea* Hort. *The purple-leaved Sycamore*.—The leaves are of a fine purple underneath. This variety was originated in Saunders's Nursery, Jersey, about 1828, and is now to be met with in all the principal nurseries. The tree has a very fine appearance when the leaves are slightly ruffled by the wind, alternately appearing clothed in purple and in pale green. In spring, when the leaves first expand, the purple bloom is not obvious; but when they become matured it is very distinct.

‡ A. *P. 5 subobtusata* Dec. Prod., i. p. 594. *The half-obtuse-leaved Sycamore*.—Lobes of leaves blunter; fruit and wings larger. *A. opulifolium* Thuil. *Fl. Par.*, 538. *A. ritifolium* Opiz.

‡ A. *P. 6 laciniata* Loud. Hort. Brit., p. 412. *The cut-leaved Sycamore*.—Lobes of leaves jagged. (*Schm. Arb.*, i. 5.; *Don's Mill.*, i. p. 648.)


Other Varieties. In the garden of the London Horticultural Society there is a variety called *Hodgkins's Seedling*, with yellow blotched leaves; and another, called *Leslie's Seedling*. In Hayne's *Dendrologische Flora* there are, also, the following varieties: *A. P. stenoptera*, *A. P. macroptera*, and *A. P. microptera*, which differ in the proportions of the wings of the keys, and do not appear worth farther notice.

Description. A large handsome tree, of quick growth, with a smooth ash-coloured bark, and round spreading branches. Leaves on long footstalks, 4 in. or 5 in. broad, palmate, with 5 acute, variously serrated lobes; the middle one largest, pale or glaucous beneath. Flowers green, the size of a currant blossom, disposed into axillary, pendulous, compound clusters. Capsules 2 or 3, with broad spreading wings. (*Smith's Eng. Flora*, ii. p. 230., with adaptation.) The fruits of this species are botanically interesting, from the readiness with which the funiculus may be traced in its passage through the base of the samara to its union with the seed; and from the neat and copious lining of soft and glossy down, with which the interior of the cell of the samara is coated, as if for a commodious lodging for the seed, till wind shall have acted on the wing of the samara, and disseminated it, and the moisture of the earth whereon it falls shall have excited the seed it contains to germinate. In this species, the cotyledons are circinate folded, and incumbent on the radicle. The cotyledons, but, perhaps, after germination, and the primordial leaves (those first produced on germination), are, when chewed, bitter. Professor Henslow



has found, by "a careful search among the numerous young plants of this tree which every where spring up in its neighbourhood, many in which the cotyledons were either three or four. In some instances, where there were only two, as usual, one of them was more or less cloven down the middle (*fig.* 110. *a*); and these served to illustrate, in a marked manner, the way in which others had become possessed of more than their ordinary number. For, in these cases, either two of the cotyledons were not, at first, so large as the third, when there were three

only (*b*); or else, when four were present, they were all proportionally smaller than in those plants which bore two (*fig. 111. c*). This shows that the multiplication of the cotyledons, in some plants, may be the result merely of a



subdivision in the two which belong to them in their normal condition, and that it may not have originated in any supernumerary development of these organs themselves. Their comparative inequality, however, soon ceases as the plant develops itself. In one instance, I have

remarked a cohesion taking place between the two cotyledons nearly throughout their whole length (*fig. 110. d*); and then the young plant had strangely assumed the form of a monocotyledon. Sometimes the superfluous division was continued to the primordial leaves, of which there were one large, and two that were smaller (*fig 111. c*): but I have never observed the anomaly extend beyond them; the next in succession, and all after them, being developed in pairs, in the usual way. The above figures are selected from among several varieties which I possess of this anomalous germination of the sycamore." (*Mag. Nat. Hist.*, vol. v. p. 346, 347.) The growth of the sycamore is very rapid compared with that of other trees, particularly when it is in a deep, free, rich soil, and in a mild climate. It arrives at its full growth in 50 or 60 years; but it requires to be 80 or 100 years old before its wood arrives at perfection. In marshy soil, or in dry sand, and even on chalk, the tree never attains any size. It produces fertile seeds at the age of 20 years, but flowers several years sooner; sometimes even perfecting its seeds sooner also. The longevity of the tree is from 140 to 200 years, though it has been known of a much greater age. M. Hartig has felled sycamore trees 200 years old, and upwards of 100 ft. in height, the timber of which was perfectly sound.

Geography. Found in various parts of Europe, particularly in Switzerland, Germany, Austria, and Italy, in wooded mountainous situations. In England, it is found in hedges and about houses, but not truly wild, according to Smith; though others consider it indigenous. Gerard, in 1597, says it is a stranger in England, only found growing in the walks and places of pleasure of noblemen, where it is planted for the sake of its shadow. Parkinson observes, "It is cherished in our land only in orchards or elsewhere, for shade and walks." Ray speaks of it as very common in courtyards, churchyards, avenues, and about noblemen's houses; but says it began, in his time, not to be much in request, because of the great litter occasioned in gardens and walks by the falling leaves. Martyn, in his edition of *Miller's Dictionary*, says that, if it were truly indigenous, the country would have been full of it; since the tree comes up with such wonderful facility from the seed. For the same reason, Dr. Walker supposes it to have been one of the very earliest of foreign trees introduced into Scotland. Sir T. D. Lauder says, "It is a favourite Scotch tree, having been much planted about old aristocratic residences in Scotland; and, if the doubt of its being a native of Britain be true, which, however, we cannot believe, then it is probable that the long intimacy which subsisted between France and Scotland may be the cause of its being so prevalent in the latter country." (*Lauder's Gilpin*, i. p. 121.) In Switzerland, the tree is found from 2000 ft. to 3000 ft. above the level of the sea, reaching up the mountain s to the point where *Facchinum Fitis idæa* commences; provided, however, that the soil be dry and of a good quality. In such situations it suffers much less from frost and snow than many other trees.

History. The first record of the tree, as in cultivation in Britain, is in

Turner's *Herbal*, in 1551 : it is mentioned by all subsequent British authors as of doubtful indigeneness. From the facility with which it is propagated, the hardness and vigorous growth of the tree, its various uses, especially, as Dr. Walker observes, for forming domestic utensils, and also the beauty of its buds in spring, and of its foliage in early summer, it has been very generally planted.

Properties and Uses. The wood, when the tree is young, is white ; but, as it gets older, the wood becomes a little yellow, and often brown, especially towards the heart. It is compact and firm, without being very hard ; of a fine grain, sometimes veined, susceptible of a high polish, and easily worked, either on the bench, or in the turning-lathe. It does not warp, and is not likely to be attacked by worms. It weighs per cubic foot, newly cut, 64 lb. ; half dry, 56 lb. ; dry, 48 lb. It loses, in drying, about a twelfth part of its bulk.

In France and Germany, it is much sought after by wheelwrights, cabinet-makers, turners, sculptors in wood, manufacturers of musical instruments, and especially of violins, and makers of toys and other small wares. The roots, which are often agreeably veined, and the stools or stumps where the plant has been long treated as a bush, and cut periodically for coppice-wood, is eagerly sought after for curious cabinet-works and for inlaying. The wood is used for pestles, for tables, rollers, spoons, plates, and other household articles ; it is also used for gun-stocks, and in every kind of structure, whether under water or in the air. According to M. Hartig, the principal German writer on timber trees and their uses, the wood of the common sycamore is the most valuable of all woods as fuel, both for the quantity of heat which it gives out, and the time that it continues burning : it surpasses the beech, in these respects, in the proportion of 1757 to 1540. Converted into charcoal, it is superior to the beech in the proportion of 1647 to 1600. The leaves, gathered green and dried, form an excellent forage for sheep during the winter. The sap has been drawn from the trees in Germany, and various experiments made with it. At first, it is as clear as water, and sweet ; but, after it has run from the tree for some time, and begins to run slowly, it takes a whitish colour, and becomes sweeter and of a thicker consistence ; though this thick sap is found to contain less sugar than that which comes off first, and is quite clear. From a tree 18 in. in diameter, from which the sap was allowed to flow for five days, 36 quarts were obtained. The proportion of sugar produced by the sap varies. Sometimes an ounce of sugar from a quart of liquor has been obtained ; but, generally, not so much. The variations depend on the age of the tree, the vigour of its growth, the nature of the soil, the temperature of the season, and a number of other circumstances of which little is known. In Scotland, Sir Thomas Dick Lauder informs us, incisions were made in the trunk of a sycamore tree of 45 years' growth, at 5 ft. from the ground, in the beginning of March, 1816. " A colourless and transparent sap flowed freely, so as in two or three hours to fill a bottle capable of containing 1 lb. of water. Three bottles and a half were collected, weighing, in all, 3 lb. 4 oz. The sap was evaporated by the heat of a fire, and gave 214 grains of a product in colour resembling raw sugar, and sweet in taste, with a peculiar flavour. After being kept fifteen months, this sugar was slightly moist on the surface. The quantity of sap employed in the evaporation was 24,960 grains, from which 214 grains of sugar were obtained : therefore, 116 parts of sap yielded one part of sugar. The experiment was made at Cannon Park, in Stirlingshire, on the 7th and 8th of March, 1816. (*Lauder's Gilpin*, i. p. 124.) Dr. Walker states that the sap is made into wine in the Western Highlands of Scotland.

In Britain, the uses to which the *A. Pseudo-Platanus* is applied are much less various than in France and Germany. The species is a very umbrageous one, from its numerous branches, and numerous and large leaves ; and hence it is eligible in all cases where trees are wanted to afford dense shade : it may be on this account that it is sometimes seen bounding the homesteads of a farm, and on the sunny side of the dairy in the farm-yard. It is used in joinery and turnery, and cabinet-making ; by musical instrument makers ; for cider-

presses; and, sometimes, for gun-stocks. Formerly, when wooden dishes and spoons were more used than they are at present, it was much in demand, especially in Scotland, by the manufacturers of these articles.

As underwood, the sycamore shoots freely from the stool to the age of 80 or 100 years. As a timber tree, it is most advantageously cut down at the age of 80 years, or from that age to 100. In Germany, the tree which is commonly planted along with it, in plantations made with a view to profit, is the beech. As an ornamental tree, it produces the best effect; either singly, in groups of two or three, placed sufficiently near to form a whole, but not so as to touch each other; and in rows or avenues. The varieties with variegated leaves are very ornamental in the beginning of summer; but their leaves are almost always more or less imperfect, especially on the edges, and fall off much sooner in the autumn than those of the species. The leaves of the purple variety are not liable to the same objection as those of the variegated sorts.

In Scotland, children amuse themselves by cutting openings in the bark, and sipping the sap that flows from its wounds (*Mag. Nat. Hist.*); and they also play with the large buds which are found on the points of the shoots, which they call cocks, and the small side-buds, which they call hens. In England, children suck the wings of the growing keys, for the sake of obtaining the sweet exudation that is upon them.

Poetical and historical Allusions. The sycamore, in the language of flowers, signifies curiosity, because it was supposed to be "the tree on which Zacchæus climbed to see Christ pass on his way to Jerusalem, when the people strewed leaves and branches of palm and other trees in his way, exclaiming, 'Hosanna to the Son of David!'" (*Syl. Flor.*, p. 221.) The tree called the sycamore in the Bible, however, was not the *Acer Pseudo-Platanus*, but the *Ficus Sycómorus*; though the supposition that the first was the sycamore of the Scriptures induced many religious persons, in the fourteenth and fifteenth centuries, to plant it near their houses and in their gardens. Evelyn mentions this practice, and condemns it; as the sycamore, from the frequency of honey-dew on its leaves, is a very unwholesome and unsightly tree. It is mentioned by Chancer; and Cowper says,—

——— "Nor unnoticed pass
The sycamore, capricious in attire;
Now green, now tawny; and ere autumn yet
Has changed the woods, in scarlet honours bright."

Soil and Situation. The common sycamore will grow in any soil not saturated with moisture; but it seems to prefer one that is dry and free, rather than one stiff or moist. It will grow in exposed situations, and especially on the sea coast, and maintain its erect position against the sea breeze better than most other trees. It is in use for this purpose in Scotland, and also for planting round farm-houses and cottages on bleak hills. In such situations, an instance can hardly be found of the head of the tree leaning more to one side than another. Even when the wind blows strongly in one direction for nine months in the year, this tree maintains its perpendicularity and symmetrical form.

Propagation and Culture. This species is invariably propagated by seed; and the variegated-leaved and other varieties by layers, or by budding or grafting. It will also propagate freely by cuttings of the roots. The seeds may either be sown immediately after they are gathered, or they may be kept in sand till the following spring. If the seeds are kept dry, and unmixed with sand or earth till spring, they seldom come up the same year, and sometimes lose their vegetative properties altogether.

Accidents and Diseases. The leaves are attacked by various insects, and the young shoots eaten by goats, hares, cattle, horses, and mules. In a suitable soil, the tree is attacked by few diseases; but at great elevations, on unsuitable soils, and especially on such as are wet, the superabundance of sap produces ha morrhage, and, according to M. Werneck, dropsy. In both these cases, the

roots soon become spongy and rotten, and the plant becomes a prey to lichens and fungi, and finally dies. The cause of the disease being the humidity of the soil, it can only be prevented by planting the tree on soil sufficiently dry, either naturally or by drainage. Two parasitic species of *Fungus* are found upon the leaves: *Xyloma acérinum Pers.*, described and figured in the *Encyclopædia of Plants*, No. 16490., and *Erineum acérinum Pers.*, described and figured in the *Encyclopædia of Plants*, No. 16593. A sweet clammy matter exudes from the foliage, and is fed upon by insects, whose excrements tend to discolour it; which shows, in some seasons, considerable discolouration and want of cleanness and freshness, that may be referred in part to this cause, and may be in part referable to particles of dust and other matter floating in the atmosphere, and falling on the clammy surface of the foliage.

Statistics. There are a great many fine specimens of this tree in different parts of Europe; and, as it is a well-known species, we shall only select a few, as in other cases, partly to show the rate of growth, and partly to show the magnitude attained relatively to time.

A. Pseudo-Plátanus in the Environs of London. At Kew, there is a tree, 100 years planted, and 74 ft. high, the trunk 5½ ft. in diameter, and the diameter of the head 60 ft.; at Mount Grove, Hampstead, *A. P.* álbo variegáta, 69 ft. high, the trunk 2 ft. 10 in. in diameter, and the head 47 ft. in diameter; the soil a sandy loam, and the situation exposed.

A. Pseudo-Plátanus South of London. In Hampshire, in Wilkins's Nursery, Isle of Wight, 10 years planted, and 25 ft. high; at Alresford, 81 years planted, 70 ft. high, the trunk, at 1 ft. from the ground, 9 ft. in diameter; in Devonshire, at Endsleigh Cottage, 22 years planted, and 40 ft. high; in Dorsetshire, at Bridehead House, 77 years old, and 40 ft. high; in Kent, at Cobham Hall, 98 ft. high; the diameter of the trunk, at 1 ft. from the ground, 5 ft. 4 in.; the contents of the tree in timber, 450 ft.; in Somersetshire, at Broekley Hall, 90 ft. high, and the diameter of the trunk 2 ft. 10 in.

A. Pseudo-Plátanus North of London. In Berkshire, at Bear Wood, 15 years planted, 30 ft. high; in Worcestershire, at Hadzor House, 10 years planted, and 28 feet high; at Hagley, 9 years planted, and 17 ft. high; in Lancashire, at Lancaster, in the Friends' Burying Ground, several fine trees about a century old, between 60 ft. and 70 ft. high, with trunks from 2½ ft. to 3 ft. in diameter, and heads 60 ft. to 70 ft. in diameter; at Latham House, 40 years planted, and 45 ft. high; at Aldcliff Hall, 50 ft. high, diameter of the trunk 40 ft. and of the head 60 ft.; in Northumberland, at Woolstington, 55 ft. high, the diameter of the trunk 8 ft. 4 in., and of the head 50 ft.; in Suffolk, at Finborough Hall, 70 years planted, and 70 ft. high; at Livermore, *A. P.* variegáta, 13 years planted, 21 ft. high; in Yorkshire, at Hackness, 70 years planted, and 74 ft. high; at Grimston, 12 years planted, and 40 ft. high; in Warwickshire, at Coombe Abbey, *A. P.* variegáta, 60 years planted, and 53 ft. high; in Rutlandshire, at Belvoir Castle, the species 18 years planted, and 24 ft. high; in Hertfordshire, at Cheshunt, *A. P.* variegáta, 10 years planted, 28 ft. high.

A. Pseudo-Plátanus in Scotland. In Mid-Lothian, at Hopeton House, 50 ft. high, the trunk 4 ft. 10 in. in diameter, and the diameter of the space covered by the branches 90 ft.; at Moredun Park are two trees 50 ft. high and 70 ft. high, the trunks 3 ft. 3 in. and 4 ft. 6 in. in diameter, and the heads from 50 ft. to 60 ft. in diameter; at Calder House, one standing on the pleasure-ground on the road from the house to the church, measured, on the 4th of October, 1799, 17 ft. 7 in. in girth, and at the ground 20 ft. 3 in. Its trunk was 12 ft. high, with a clear bole, after which it divided into five great arms. Its branches extended in diameter about 60 ft. This tree is known to have been planted before the Reformation; and it is supposed to be not less than 300 years old; yet it has the appearance of being perfectly sound. It was the tree to which, long ago, the iron jugs (a species of pillory) were fastened. The tree came gradually to grow over them; and they have now been completely enclosed in its trunk for a considerable time. At the place where they are enclosed, there is a great protuberance, on the south side of the tree, at the height of between 4 ft. and 5 ft." (*Lauder's Gilpin*, i. 272.) At Preston Hall, there is a tree, 19 ft. 3 in. in circumference, at 1 ft. from the ground; at Nidry Marisshall, one which, at 3 ft. from the ground, measures 19 ft. 4 in. In Renfrewshire, there is one 65 ft. high, diameter of the trunk 5 ft. 2 in., and of the head 80 ft.; at Bishopton, one 60 ft. high, the trunk 6½ ft. in diameter, and the contents in timber 720 ft., figured by Strutt in his *Sylva Britannica*; in East Lothian, at Tynningham, 52 ft. high, diameter of the trunk 3 ft. 4 in., and of the head 96 ft.; in Berwickshire, at Yester, at 1 ft. from the ground, 19 ft. in circumference; at Newbattle Abbey, many large trees, planted before the Reformation; one planted before the year 1530; at Nisbet, 65 ft. high in 1795 (*Lauder's Gilpin*, vol. i. p. 272.); in Ayrshire, at Dornholm, 70 ft. high; at Blair, 100 years planted, 70 ft. high; at Cassilis, the diameter of the head 84 ft., and of the trunk 5 ft.; in Clackmannanshire, in the garden of the Dollar Institution, 12 years planted, and 28 ft. high; in Perthshire, in Messrs. Dickson and Turnbull's Nursery, Perth, 24 years planted, 57 ft. high, the diameter of the trunk 28 in., and of the head 30 ft.; at Taymouth, 200 years planted, and 100 ft. high, the diameter of the trunk 6 ft., and of the head 40 ft.; in Sutherlandshire, at Dunrobin Castle, 63 ft. high, diameter of the trunk 2 ft. 5 in., and of the head 27 ft.; another tree there, 180 years planted, 68 ft. high, and the trunk 3½ ft. in diameter; in Stirlingshire, at Airthrey Castle, 60 ft. high, the diameter of the space covered by the head 66 ft.; at Callendar Park, 70 ft. high, the diameter of the trunk 5 ft., and of the head 66 ft.; at Sauchie, 89 ft. high, the diameter of the trunk 3 ft. and of the head 50 ft.

A. Pseudo-Plátanus in Ireland. In Dublin, at the Glasnevin Botanic Garden, 25 years planted, and 24 ft. high; at Cypress Grove, 45 ft. high; in Connaught, at Makree Castle, 77 ft. high, diameter of the trunk 4 ft., and of the head 68 ft.; in Galway, at Coole, 40 ft. high, the diameter of the trunk 14 in., and of the head 38 ft.; in Down, at Castle Ward, 134 years planted, and 64 ft. high, the diameter of the trunk 3 ft., and of the head 42 ft.

A. Pseudo-Plátanus in Foreign Countries. In France, in the Botanic Garden at Toulon, 48 years planted, and 90 ft. high; in the neighbourhood of Nantes, 60 ft. high. In Hanover, at Schwöbber, 80 ft. high; in Saxony, at Wörlitz, 65 years planted, 50 ft. high; in Austria, in the garden of the University of Vienna, 30 years planted, and 40 ft. high; at Laxenburg, 50 years planted, and 35 ft. high. In Prussia, in the Pfauen Insel, at Potsdam, 40 years planted, and 45 ft. high. In Bavaria, at Munich, 26 years planted, and 15 ft. high. In Switzerland, at Friburg, is a tree supposed to be about 500 years old, the trunk is 26½ ft. in circumference at 1 ft. from the ground. In Sweden, at Lund, 14 years planted, and 38 ft. high.

Commercial Statistics. Price of plants, in the London nurseries, seedling,

10s. a 1000, plants 6 ft. high 2s. each, the variegated varieties 2s. 6d. each, the purple-leaved 3s. each. At Bollwyller, 80 cents a plant, and the variegated varieties 1 franc 20 cents a plant; at New York, ?.

‡ 9. *A. OBTUSUM* *Kit.* The obtuse-lobed-leaved Maple.

Identification. Kit. in Willd. Spec., 4. p. 918.; Dec. Prod., 1. p. 594.; Don's Mill., 1. p. 949.

Synonymes. *A. neapolitanum* Tenore; *A. hybridum*, in the Loud. Hort. Soc. Gard. in 1834; the Neapolitan Maple.

Engravings. Tratt. Arch., 1. No. 14.; our fig. 124., in p. 450, 451.; and the plate of this species in our Second Volume.

Spec. Char., &c. Leaves cordate, roundish, 5-lobed; lobes bluntish (or pointed), repandly toothed, velvety beneath. Corymbs pendulous. Pedicels hairy. Fruit rather hairy, with the wings somewhat diverging. (*Don's Mill.*, i. p. 649.) The flowers are pendulous, pale, and few in the panicle. A large tree, with the general habit of *A. Pseudo-Platanus*, but apparently of more vigorous growth; a native of Hungary, Croatia, and many parts of Italy; and introduced into England in 1825. "On all the hills and lower mountains of the kingdom of Naples, in Camaldoni, Castellamare, and the Abruzzi, it is found abundantly, growing, usually, to the height of 40 ft. It is extremely striking, with its reddish purple branches, in the wood of Lucania, between Rotonda and Rubia; and, in the Basilicate, and Calabria, it is said, by Tenore, to acquire colossal dimensions. It is certainly very singular that so fine a tree as this, occupying so large a tract of country frequently visited by English tourists, should be almost unknown in this country; and yet, although it is perfectly hardy, and very easily multiplied, it is scarcely ever met with in any but botanical collections." (*Pen. Cyc.*, vol. i. p. 77.) There is a noble specimen of this tree in the garden of the London Horticultural Society at Chiswick; which, though only 10 or 12 years planted, in 1835 had attained the height of 26 ft., with a trunk $5\frac{1}{2}$ in. in diameter, as shown in our plate in Vol. II.

‡ 10. *A. BARBATUM* *Michx.* The bearded-calyved Maple.

Identification. Michx. Fl. Bor. Amer., 1. p. 252.; Pursh, Fl. Amer. Sept., 1. p. 267.; Dec. Prod., 1. p. 595.; Don's Mill., 1. p. 649.

Synonymes. *A. carolinianum* *Bolt.*; *A. trilobatum*, in the London Hort. Soc. Gard. in 1835.

Engravings. A leaf is shown in *Pen. Cyc.*, vol. 1. p. 76.; and several in our fig. 125. in p. 452.

Spec. Char., &c. Leaves heart-shaped, 3-lobed, nearly equally serrated, nearly smooth beneath. Clusters sessile. The stalks of the female flowers simple, of the male flowers branched. Calyx bearded internally. Keys smooth, diverging but little. (*Pen. Cyc.*) This species, according to Pursh, inhabits North America, between New Jersey and Carolina, in deep pine and cedar swamps. It was found on the west side of the Rocky Mountains, about the sources of the Columbia, by Douglas; but Dr. Hooker says the specimens sent home by him are too young to enable him to form an opinion as to the correctness of the name. There are two plants of this species, under the name of *A. trilobatum*, in the garden of the London Horticultural Society, where they form low trees, or bushes, about 10 ft. high. In its native country, the tree is said to grow to the height of 20 ft. The plants named *A. barbatum*, in the London Horticultural Society's Garden, and in Messrs. Loddiges's arboretum, and in some of the nurseries in 1835, seem to be *A. platanoides*. The leaves, and the general appearance of the plant, are those of *A. Pseudo-Platanus*, diminished, in all its parts, to one third of its usual size. This species was introduced in 1812; and plants of it, under the name of *A. trilobatum*, may be obtained in some of the nurseries.

D. *Leaves 5, rarely 7-lobed.*

‡ 11. *A. OPALUS* *Dec.* The Opal, or Italian, Maple.

Identification. Ait Hort. Kew., 3. p. 136.; Dec. Prod., 1. p. 594.; Don's Mill., 1. p. 649.

Synonymes. *A. Opalus* *Lin.*, *Mill.*, and other authors; *A. rotundifolium* *Lam. Dict.*, 3. p. 582.; *A. italicum* *Lamb. Ac.*, No. 8.; *A. villosum* *Pres.*; l'Érable Opale, Érable à Feuilles rondes, or l'Érable d'Italie *P.*

Derivation. The specific appellation of *O'palus* has been given to this species, probably from the thick opal-like aspect of the leaves.

Engravings. Baudril. *Traité*, &c., vol. 5, p. 13. ; our *fig.* 126. in p. 453 ; and the plate of this species in our Second Volume.

Spec. Char., &c. Leaves more or less heart-shaped, roundish, 5-lobed, smooth beneath ; the lobes generally obtuse, and coarsely serrated. Flowers in drooping corymbs. Keys smooth. (*Pen. Cyc.*) A tree, a native of Corsica ; from which country it was brought to Paris by M. Richard, and thence to England, in 1752. It is described by Baudillart as a branchy tufted tree, covered with smooth leaves, somewhat coriaceous, roundish, indented, with five blunt lobes, deep green on the upper surface, and somewhat glaucous underneath, with long red petioles. Its flowers are whitish, in short racemes ; and the small fruits, or keys, which succeed them, are almost round. It found in forests and on mountains in Corsica, and in Italy ; where, from the denseness of its shade, it is sometimes planted by road sides, and in gardens near houses. The red colour of the petioles, of the leaves, of the fruits, and even the red tinge of the leaves themselves, more especially in autumn, give it rather a morbid appearance. It pushes later in the spring than most of the other species. The wood is veined, and very close : in Italy, it is used for gun-stocks ; and the roots, especially of those trees which have been often cut down, are very much sought after on account of their hardness, and their curious knots and blotches, which render them suitable for making snuff-boxes, and for inlaid work.

Variety. *A. coriaceum*, in the arboretum of the Messrs. Loddiges, seems to be a variety of this species ; but *A. opulifolium*, No. 14, as described by Baudrillart, seems quite distinct from it.

Statistics. There is a plant of this species in the garden of the London Horticultural Society, which answers perfectly to M. Baudrillart's description. The largest tree in the neighbourhood of London, bearing the name of *A. O'palus*, is at Fulham Palace ; where, in 25 years, it has attained the height of 35 ft. ; in Sussex, at Langham Park, 9 years planted, it is 20 ft. high ; in Staffordshire, at Trentham, 12 years planted, it is 7 ft. high ; in Yorkshire, at Grimston, 12 years planted, 24 ft. high ; in Scotland, in the garden of the Caledonian Horticultural Society, Edinburgh, 8 years planted, and 12 ft. high ; at Hopeton House, 18 years planted, and 18 ft. high ; in Argyllshire, at Toward Castle, there is a tree, which is said to be considered *A'cer O'palus* by Dr. Hooker, which is no less than 50 ft. high, and girths 4 ft. at 1 ft. from the ground. In France, in the Jardin des Plantes, 30 years planted, and 34 ft. high ; in the Botanic Garden at Toulon, 40 years planted, and 50 ft. high. In Belgium, in the Botanic Garden at Ghent, 15 ft. high. In Hanover, at Schwöbber, 80 ft. high (most probably some other species, ? *A. obtusatum*) ; in Saxony, at Wörlitz, 25 ft. high. In Austria, in Rosenthal's Nursery, at Vienna, 16 years planted, and 12 ft. high.

Commercial Statistics. Price, in London, from 1s. to 1s. 6d. a plant ; and at Bollwyller, 1 franc.

§ 12. *A. OPULIFOLIUM* Will. The Guelder-Rose-leaved Maple.

Identification. Will. Dauph., 4, p. 802. ; Don's Mill., 1, p. 649.

Synonymes. *A. hispanicum* Pomr. *Act. Toul.*, 3, p. 305. ; *A. vcrnum* Reyp. ; *A. montanum* C. Bannlin, Pin., 431. ; *E'rable durct*, or *E'rable à Feuilles d'Obier*, and *Ayart* in Dauphiné, Fr. ; *Schneeboll-blättriger Ahorn*, Ger.

Engraving. *Traité* Arch., 1, No. 13. ; and the plate of this species, in our Second Volume.

Spec. Char., &c. Leaves cordate, roundish, 5-lobed. Lobes obtuse, bluntly and coarsely toothed. Corymbs almost sessile. Ovaries and fruit smooth, with wings rather diverging. (*Don's Mill.*, i, p. 649.) According to Dr. Lindley, in the *Pen. Cyc.*, this kind is the same as *A. O'palus* ; but, according to Baudrillart, it is quite distinct. Its height, the latter says, is from 20 ft. to 25 ft. or 30 ft. It grows naturally in the French Alps, and on the Pyrenees. Its bark is grey ; its leaves have 5 lobes, somewhat rounded, a little toothed, and greener above than below. It flowers are in drooping racemes ; its fruits are swelled out, and their wings spreading ; so much so as to form almost a straight line, like those of *A. platanoides*. It is common on the rocks of Mount Jura ; and is considered preferable to all the other maples for its wood, which is hard and compact, without sap-wood, not easily split, and so homogeneous in its texture, that it is almost impossible to distinguish in it the annual layers. It takes the finest polish ; it is white, lightly shaded with lemon-colour, sometimes exhibiting flashes or shades of red, but not red veins. Completely dried, it weighs 52 lb. 11 oz. the cubic foot. In Bugly it is used by wheelwrights, and makes excellent naves to wheels.

"M. Maratray, inspector of forests in the department of Mount Jura, sent, in 1807, to the Administration of Forests in Paris, a specimen of this wood, which justified all that had been said of the fineness and homogeneity of its grain. He also sent seeds, part of which came up the first year, and the remainder the year following. The plants have made tolerable progress; but it remains to be known, whether, in fertile soil, the tree will preserve the valuable quality of its wood." (*Traité, &c.*, i. p. 51.) The plants have been distributed among the different government gardens of France; and particularly those of Versailles, under the direction of M. Bose. At Bollwyller, there are plants of this species which, it is said, can be furnished of considerable size; and they are designated in the *Catalogue*, "*Acer opulifolium*, non *A. Opalus*;" price 1 franc 50 cents. There was a tree in the garden of the London Horticultural Society, in 1835, which differed somewhat from *A. Opalus*, and seemed to us intermediate between that species and *A. barbatum*, or, as it was then marked, in 1835, *A. trilobatum*; but, whether it was the *A. opulifolium* of Villars and Baudrillart, we are unable to say.

‡ 13. *A. CIRGINATUM* Pursh. The round-leaved Maple.

Identification. Pursh *Fl. Amer. Sept.*, l. p. 267.; Dec. Prod., l. p. 535.; Don's Mill., l. p. 651.
Engraving. Hook. Amer., t. 39.; and our fig. 112., and fig. 127. in p. 454.

Spec. Char., &c. Leaves orbicular, rather cordate at the base, 7-lobed, smooth on both surfaces; lobes acutely toothed; nerves and veins hairy at their origins. (*Don's Mill.*, i. p. 651.) A tree from 20 ft. to 40 ft. high.

112



Branches slender, pendulous, and crooked; often taking root, in the manner of those of many species of *Picea*. Bark smooth, green when young, white when fully grown. Leaf the length of the finger, upon rather a short foot-stalk, membranaceous, heart-shaped, with 7—9-lobes, and 7—9-nerves, smooth above, except hairs in the axils of the nerves; downy beneath, and in the axils of the nerves woolly: lobes ovate, acute, and acutely serrated; the sinuses acute; the nerves radiate from the tip of the petiole, and one extends to the tip of each lobe. Flowers (produced in April and May) of a middling size, in nodding corymbs, that are on long peduncles. (*Hook. Fl. Bor.*

Amer.) This is a very marked and beautiful species; distinguishable, at sight, by the regular form of its leaves, and pale reddish green colour.

Geography. On the great rapids of Columbia River. (*Lewis*, in *Pursh's Fl. Am. Sept.*) Common along the north-west coast of North America, between lat. 43° and 49°. (*Douglas*; *D. Scouler.*) *A. circinatum*, like *A. macrophyllum*, is exclusively confined to the woody mountainous country that skirts the shores; and there, among the pine forests, it forms almost impenetrable thickets. (*Douglas*, in *Hook. Fl. Bor. Amer.*, vol. i. p. 112.)

Properties and Uses. The wood is fine, white, and close-grained, very tough, and susceptible of a good polish. From the slender branches the native tribes make the hoops of their scoop-nets, which they employ for taking the salmon at the rapids, and in the contracted parts of the river.

Statistics. There is a plant of this species, in the London Horticultural Society's Garden, about 2 ft. high; and one at Messrs. Loddiges's about the same height. In Berkshire, at High Clere, there is a tree which has blossomed and ripened seeds.

‡ 14. *A. PALMATUM* Thunb. The palmate-leaved Maple.

Identification. Thunb. *Fl. Jap.*, p. 161.; Dec. Prod., l. p. 595.; Don's Mill., l. p. 650.
Engraving. Tratt. Arch., l. No. 17.; and our fig. 128. in p. 455.

Spec. Char., &c. Leaves smooth, palmately divided into 5—7-lobes beyond the middle; lobes acuminate, oblong, serrated. Umbels 5—7-flowered. (*Don's Mill.*, i. p. 650.) A native of Japan, and introduced in 1832.

Branches and corolla purple. Fruit woolly. There are plants of this species in the garden of the London Horticultural Society, the leaves of which, as will be seen by our *fig.* 128., are strikingly distinct. The plants appear to be rather tender, and we would recommend them to be tried, in the first instance, against a wall.

‡ 15. *A. ERIOCÁRPUM Michx.* The hairy-fruited, or white, Maple.

Identification. Michx. *Fl. Amer. Bor.*, 2. p. 213; Don's Mill., 1. p. 650.

Synonymes. *A. dasycárpum Willd. Spec.*, 4. p. 985; *A. tomentosum Hort. Par.*; *A. glaucum*

Marsh.; *A. virginianum Duh.*; *A. rubrum Wagenh.*; white, or soft, Maple, *United States*; Sir

Charles Wager's Maple; *E'rablé à Fruits cotonneux, or E'rablé blanc, Fr.*; *rauhler Ahorn, Ger.*

Engraving. Desf. *Ann. Mus.*, 7. t. 25.; *Tratt. Arch.*, 1. No. 8.; our *fig.* 129. in p. 456.; and the plate of this species in our Second Volume.

Spec. Char., &c. Leaves truncate at the base, smooth and glaucous beneath, palmately 5-lobed, with blunt recesses, and unequally and deeply toothed lobes. Flowers conglomerate, on short pedicels, apetalous, pentandrous. Ovaries downy. (*Don's Mill.*, i. p. 650.) A large tree, with pale greenish yellow seeds, and flowers tinged with pale pink. They are produced in April and May; and seeds are ripened by midsummer, from which plants may be raised the same year. Introduced by Sir Charles Wager, in 1725.

Description. The trunk of the white maple is low, and divides itself into a great number of limbs, so divergent, that Michaux says they form a head more spacious, in proportion to the size of the trunk, than that of any other tree with which he is acquainted. The tree blooms early in the spring: its flowers are small and sessile, with a downy ovary. The fruit is larger than that of any other species which grows east of the Mississippi. It consists of two capsules joined at the base, each of which encloses one roundish seed, and is terminated by a large, membranous, falciform wing. In Pennsylvania, it is ripe about the 1st of May; and a month earlier on the Savannah river, and in Georgia. At this period the leaves, which have attained half their size, are very downy underneath: a month later, when fully grown, they are perfectly smooth. They are opposite, and supported by long petioles; they are divided by deep sinuses into 4 lobes, are toothed on the edges, of a bright green on the upper surface, and of a beautiful white beneath. The foliage, however, is scattered, and leaves an open thoroughfare to the sunbeams. "The young leaves, and young germs, are very downy; but the old leaves, and perfect fruit, are glabrous." (*Hook, Fl. Amer.*, p. 114.) The wood of this maple is very white, and of a fine grain; but it is softer and lighter than that of the other species in the United States, and, from its want of strength and durability, is little used. (*Michaux*, p. 215.) In the United States, as well as in England, this species is often confounded with *Acer rubrum*, which, in the leaves, it nearly resembles; but it differs in its inflated woolly fruit, expressed in the terms *eriocárpum* and *dasycárpum*, and in its flowers, which are produced in small compact axillary groups, and are almost, or quite, sessile; while those of *A. rubrum* are produced in axillary groups on peduncles of irregular length (the shortest being about 1 in., and the longest about 2 in.), and are succeeded by smooth compressed fruits.

Geography. *A. eriocárpum*, in the Atlantic parts of the United States, commences on the banks of Sandy River, in the district of Maine; and those of the Connecticut, near Windsor, in Vermont, are its most northern points. But, like many other trees, it is pinched by the rigorous winters of this latitude, and never reaches the size which it attains a few degrees farther south. It is found on the banks of all the rivers which flow from the mountains to the ocean; though it is less common along the streams which water the southern parts of the Carolinas and of Georgia. In no part of the United States is it more multiplied than in the western country; and nowhere is its vegetation more luxuriant than on the banks of the Ohio, and of the great rivers which empty themselves into it. There sometimes alone, and sometimes mingled with the willow, which is found along all these waters, it contributes singularly, by its magnificent foliage, to the embellishment of the scene. The brilliant white of the leaves beneath forms a striking contrast

with the bright green above; and the alternate reflection of the two surfaces in the water heightens the beauty of this wonderful moving mirror, and aids in forming an enchanting picture; "which," says "Michaux, during my long excursions in a canoe in these regions of solitude and silence, I contemplated with unwearied admiration. Beginning at Pittsburg, and even some miles above the junction of the Alleghany and Monongahela rivers, white maples, with trunks 12 ft. or 15 ft. in circumference, are continually to be met with at short distances." (*Michx.* p. 213, 214.)

History. This tree was introduced by Sir Charles Wager, in 1725, and has since been in general cultivation. It was at first supposed to be a variety of *A. rubrum*; and, as such, is mentioned in Martyn's *Miller*. In the first edition of *Du Hamel*, the two sorts are described as distinct varieties; and their specific distinctions were afterwards pointed out by Desfontaines, in the *Ann. du Mus. d'Hist. Nat. de Paris*, vol. vii. p. 412.; the principal distinction consisting in the fruit of *A. eriocarpum* being woolly, and that of *A. rubrum* being smooth. Both species are now in very general cultivation, throughout Europe, as ornamental trees.

Properties and Uses. In America, wooden bowls are sometimes made of the wood of this species, when that of the tulip tree cannot be procured. At Pittsburg, and in the neighbouring towns, it is used, in cabinet-making, instead of holly, and for inlaying furniture made of mahogany, cherry tree, and walnut; though it is not exactly suitable for this purpose, as it soon changes colour. The hatters of Pittsburg prefer the charcoal of this wood to every other for heating their boilers, as it affords a more uniform and durable heat than any other. Sugar is made from this species in districts where the tree abounds, but the produce is not above half that obtained from the sap of the sugar maple. The sap is in motion very early in this species, beginning to ascend about the 15th of January; so that, when sugar is made from it, the work of extracting it is sooner completed. The cellular integument rapidly produces a black precipitate with sulphate of iron. (*Michaux.*) In Europe, this tree is chiefly planted for ornamental purposes, for which it is admirably adapted by the rapidity of its growth, the graceful divergent direction of its branches, the beauty of its leaves, and the profusion of its early flowers. In mild seasons, such as the present spring of 1836, these flowers begin to burst from their buds in the first week in January; and they are often fully expanded by the end of February or beginning of March.

Soil, Situation, Propagation, &c. *A. eriocarpum* requires a deep free soil, and more moisture than most of the other species. Though it will not grow in swamps, yet it attains its greatest dimensions on the alluvial banks of rivers which are occasionally inundated. It ripens its seeds, both in America and Britain, by midsummer, or earlier; and, if these are immediately sown, they come up, and produce plants which are 8 in. or 10 in. high, by the succeeding autumn.

Statistics. The largest tree in the neighbourhood of London is at Kew, where, in 25 years, it has attained the height of 50 ft.; in Staffordshire, at Trentham, it is also 50 ft. high. Price of plants, in London, 1s. 6d. each; at Bollwyller, 1 franc; at New York, 25 cents, and seeds 1 dollar per quart.

Y 16. *A. RUBRUM* L. The red-flowering, or scarlet, Maple.

Identification. Lin. Spec. 1496.; Hayne Dend., p. 213.; Dec. Prod., l. p. 505.; Don's Mill., l. p. 650.

Synonymes. *A. virginianum* *Herm. Par.*, t. 1., *Mill. Ill.*, t. 8. f. 4., *Trew. Sel.*, t. 85, 86.; *A. coccineum* *Ait.*; soft Maple, Swamp Maple, red Maple; *E'table rouge, Fr.*; rother Aborn, *Ger.*

Engravings. *Michx. Arb.*, 2. t. 14.; *Desf. Ann. Mus.*, 7. p. 413. t. 25.; *Tratt. Arch.*, l. No. 9.; *Schmidt. Arb.*, l. t. 6.; *Krause*, t. 119.; our *fig.* 150. in p. 457.; and the plate of the tree in our Second Volume.

Spec. Char., &c. Leaves cordate at the base, glaucous beneath, deeply and unequally toothed, palmately 5-lobed, with acute recesses. Flowers conglomerate, 5-petaled, pentandrous. Ovaries smooth. (*Don's Mill.*, i. p. 650.) A tree of the middle size, from North America in 1656, remarkable for its red flowers, which are produced in April and May.

Varieties.

Y *A. r.* 2 *coccineum* and Y *A. r.* 3 *intermedium* are varieties of this species,

differing so very little from it as hardly to be worth cultivation as distinct. The leaves of *A. r. coccineum* are somewhat redder in spring, when they expand, than those of *A. rubrum*.

Description. In America, the red-flowering maple, Michaux observes, is the earliest tree, the bloom of which announces the return of spring; it is in flower, near New York, from the 10th to the 15th of April. The blossoms, which are of a beautiful purplish or deep red, unfold more than a fortnight before the leaves. They are aggregate, and situated at the extremity of the branches. The fruit is suspended by long flexible peduncles, and is of the same hue with the flowers; though it varies in size and in the intensity of its colouring, according to the exposure and dampness of the soil. The keys and the seeds are one half smaller than those of *A. eriocarpum*, and they ripen sooner. The leaves are also smaller than those of that species; but, in most other respects, they resemble them. They are glaucous and whitish underneath; and are palmated, or divided into 3 or 4 acuminate lobes, irregularly toothed. The extremities of this tree, which are formed by numerous twigs united at the base, have a remarkable appearance when garnished with flowers and seeds of a deep red, before vegetation has begun generally to revive. Before the red-flowering maple exceeds 25 ft. or 30 ft. in height, and 7 in. or 8 in. in diameter, its bark is perfectly smooth, and marked with white blotches, by which it is easily distinguished. Afterwards, the trunk, like that of the white oak (*Quercus alba*) and sweet gum (*Liquidambar Styraciflua*), becomes brown and chapped. In this tree, as in others which grow in wet places, the sap-wood bears a large proportion to the heart-wood, if, indeed, the name of heart-wood can properly be given to the irregular column, star-like in its horizontal section, which occupies the centre of large trunks, and which has points, from 1 in. to 3 in. in length, projecting into the sap-wood. (*Michaux*, p. 218, 219.) The wood weighs, when dry, 44 lb. per cubic foot. The tree only attains its full size in rich soils. Somewhat large forests of it in America, when their summits are covered with flowers, are said to present a very singular and grand appearance.

Geography. In America, towards the north, the red-flowering maple appears first, scantily, about Malebaye, in Canada, in 48° north latitude; but, in proceeding southward, it soon becomes more common, and abounds in Florida and Lower Louisiana. Of all the trees which flourish in grounds which are occasionally overflowed, this species is most multiplied in the middle and southern states. It occupies, in great part, the borders of the creeks, and abounds in all the swamps, which are often inundated, and always miry. In these situations it is accompanied by the black gum (*Nyssa sylvatica*), sweet gum (*Liquidambar Styraciflua*), shellbark hickory (*Carya squamosa*), swamp white oak (*Quercus Prinus discolor*), black ash (*Fraxinus sambucifolia*), and white ash (*Fraxinus quadrangulata*). To these are added, in the Carolinas and Georgia, the small magnolia, or swamp bay (*Magnolia glauca*), the water oak (*Quercus aquatica*), loblolly bay (*Gordonia Lasianthus*), tupelo (*Nyssa aquatica*), and the red bay (*Laurus carolinensis*). It is a remarkable fact, that, west of the mountains, between Brownsville and Pittsburg, the red-flowering maple is seen growing on elevated ground, with the oaks and the walnuts; but, in such situations, it does not attain such ample dimensions as in Pennsylvania and New Jersey: in these states exist extensive marshes, called maple swamps, exclusively covered with it; where it is found 70 ft. high, and 3 ft. or 4 ft. in diameter. (*Michaux*, p. 217, 218.) Elliot observes that, in "descending the mouths of our large rivers, the red maple is the last tree found in the swamps, the tree diminishing in size as the soil becomes impregnated with salt, until it dwindles down to a shrub, and, mingling with the candleberry myrtle (*Myrica cerifera*) and *Baccharis halimifolia*, it finally disappears." (*Hook, Fl. Bor. Amer.*, i. p. 114.)

History. This tree was cultivated by Mr. John Tradescant, jun., in 1656. It was for a long time confounded with *A. eriocarpum*. It is described, in the first edition of *Du Hamel*, as *Acer floribus rubris*; and *A. eriocarpum*, as *Acer*

virginiænum flore majore. Miller, in 1741, says, the Virginian flowering maple (*A. rubrum*) was raised from seeds, which were brought from Virginia many years since, by Mr. John Tradescant, in his garden at South Lambeth, near Vauxhall; and, since, in the garden of the Bishop of London at Fulham, where it has flowered for several years, and produced ripe seeds, from which several plants have been raised. Upon a survey of the state of the garden at Fulham, in October 1793, the *Acer rubrum* planted by Bishop Compton, at 3 ft. from the ground, measured, in girth, 4 ft. 3 in., and its computed height was 40 ft. On visiting the same gardens again in 1809, the tree was gone. (*Lyson's Env. London*, 1810, 2d edit. vol. ii. p. 228.) Miller, in 1755, speaking of *A. eriocarpum*, says, this the gardeners distinguish by the title of Sir Charles Wager's maple; the other being called the scarlet-flowering maple; but, as there is no difference in [the form of] their flowers, seeds, or leaves, so they must be deemed but one species, as they are only accidental varieties arising from seed. In continuation, he says, speaking of the scarlet variety, that it flowers about the middle of April, and that Sir Charles Wager's maple flowers about the middle or latter end of March. In Martyn's *Miller*, published in 1807, the two are still considered as the same species. We have no doubt of their being in fact only varieties, or races, of the same species; but we have kept them distinct, as they are remarkably different, both in the colour of their flowers, and in their habit of growth, though not much so in their leaves. Michaux was the first to distinguish the two kinds as species, and the distinction (as before observed, p. 424.) was first pointed out in detail by Desfontaines. As *A. rubrum* is not a rapidly growing tree, like *A. eriocarpum*, it has not been so extensively cultivated as that species; nevertheless, it is to be found in most collections, and it is propagated in the principal European nurseries.

Properties and Uses. In America, the wood of the red-flowering maple is applicable to several uses. It is harder than that of the white maple, and of a finer and closer grain; hence it is easily wrought in the lathe, and acquires, by polishing, a glossy and silken surface. It is solid, and, for many purposes, it is preferred by workmen to other kinds of wood. It is principally employed for the seats of Windsor chairs: the pieces are prepared in the country; and so considerable is the demand, that boats laden with them frequently arrive at New York and Philadelphia, where an extensive manufactory is carried on for the consumption of the neighbouring towns, and for exportation to the southern states and to the West India Islands. The whole frame of japanned chairs is made of this wood, except the back, for which hickory (*Carya*) is chosen, on account of its superior strength and elasticity. The frame, the nave, and the spokes of spinning-wheels are made of the red maple. At Philadelphia, it is exclusively used for saddle-trees; and, in the country, it is preferred for yokes, shovels, and wooden dishes, which are brought to market by the country people, and purchased by the dealers in wooden ware. It sometimes happens that, in very old trees, the grain, instead of following a perpendicular direction, is undulated; and this variety bears the name of the curled maple. This singular arrangement is never found in young trees, nor in the branches of even such as exhibit it in the trunk: it is also less conspicuous in the centre than near the circumference. Trees having this character of wood are rare, and do not exist in the proportion of one to a hundred. The serpentine direction of the fibre, which renders them difficult to split and to work, produces, in the hands of a skilful mechanic, the most beautiful effects of light and shade. These effects are rendered more striking, if, after smoothing the surface of the wood with a double-ironed plane, it is rubbed with a little sulphuric acid, and afterwards anointed with linseed oil. On examining it attentively, the varying shades are found to be owing entirely to the inflection of the rays of light; which is more sensibly perceived in viewing it in different directions by candle-light. Before mahogany became generally fashionable in the United States, the best furniture in use was made of the red-flowering maple, and bedsteads are still made of it, which, in richness and lustre, exceed the finest mahogany.

At Boston, some cabinet-makers saw it into thin plates for inlaying mahogany; but the most constant use of the curled maple is for the stocks of fowling-pieces and rifles, which to elegance and lightness unite toughness and strength, the result of the twisted direction of the fibres. The cellular matter of the inner bark is of a dusky red. By boiling, it yields a purplish colour, which, on the addition of sulphate of iron, becomes dark blue approaching to black. It is used in the country, with a certain portion of alum in solution, for dyeing black. The wood of the red-flowering maple does not burn well, and is so little esteemed for fuel, that it is rarely brought for that purpose into the cities. It has but little strength, is liable to injury from insects, and ferments and speedily decays when exposed to the alternations of dryness and moisture. For these reasons, though it is now extensively used in America, its importance in the arts is not sufficient to entitle it to preservation; and Michaux supposes that, when artificial plantations become necessary in that country, the red maple will be altogether omitted. The French Canadians make sugar from the sap of this maple, which they call *plaine*; but, as in the preceding species, the product of a given measure is only half as great as that obtained from the sugar maple. (*Michaux.*) In Britain, and throughout Europe, the sole use of the red-flowering maple is as an ornamental tree; and, whether we regard the beauty of its flowers and opening leaves in early spring; of its red fruits in the beginning of summer, or its red foliage in autumn, it deserves to be considered one of the most ornamental of hardy trees. Unfortunately, in British nurseries, it is generally raised by layers; whereas, if it were grafted on *A. eriocarpum*, or raised from seed, as it is said to be on the Continent (for, whether it be considered as a race or a species, it will reproduce itself), the plants would be of much more vigorous growth.

Soil, Situation, Propagation, &c. Contrary to the general character of the maples, this species is said to thrive best in moist soil, which must, however, at the same time, be rich; and, for the tree to attain a large size, the situation ought to be sheltered. In Britain, it is chiefly propagated by layers; but, on the Continent, almost always by seeds, which ripen before midsummer, even sooner than those of *A. eriocarpum*, and, if sown immediately, come up the same season. The seeds, even when mixed with soil, do not keep well; and, in general, but a small proportion of those sent home from America vegetate in Europe.

Statistics. In the neighbourhood of London, the largest tree is at Kew, which, in 45 years, has attained the height of 40 ft.: one at Purser's Cross is 35 ft. high; and at Kenwood, in 38 years, the tree has attained the height of 34 ft. In Berkshire, at White Knights, 30 years planted, it is 18 ft. high; in Hertfordshire, at Cheshunt, 8 years planted, it is 22 ft. high; in Staffordshire, at Trentham, it is 50 ft. high; in Surrey, on an eminence in the arborctum at Milford, there is a tree 40 ft. high, which, in autumn, when the leaves have assumed their dark red colour, looks like a column of scarlet, and is seen from a great distance all round the country. In Surrey, at Farnham Castle, a tree, 45 years planted, is 50 ft. high. In Scotland, in the Glasgow Botanic Garden, there is a tree, 16 years planted, and 20 ft. high; and in the Perth Nursery, one, 14 years planted, which is also 20 ft. high. In Ireland, in Down, at Ballyleady, 60 years planted, it is 28 ft. high; in Kilkenny, at Woodstock, 60 years planted, and 50 ft. high; in Louth, at Oriel Temple, 40 years planted, and 44 ft. high. In France, in the Botanic Garden at Toulon, 45 years planted and 29 ft. high. In Saxony, at Wörlitz, 65 years planted, and 55 ft. high. In Austria, at Laxenburg, 30 years planted, and 25 ft. high. At Brück on the Leytha, 35 years planted, and 32 ft. high. In Prussia, at Sans Souci, 40 years planted, and 38 ft. high. In Bavaria, at Munich, 24 years planted, and 40 ft. high.

Commercial Statistics. Plants, in London, cost from 1s. to 1s. 6d. each, and seeds 4s. a quart; at Bollwyller, plants are 1 franc 50 cents each; at New York, plants are 15 cents each, and seeds 2 dollars 25 cents a pound.

‡ 17. *A. MONSPESSULANUM* L. The Montpellier Maple.

Identification. Lin. Spec., 1497.; Hayne Dend., p. 210.; Dec. Prod., l. p. 595.; Don's Mill., l. p. 649.
Synonymes. *A. trilobum* Moench.; *A. trifolium* Duh.; *A. trilobatum* Lam.; *E. rable* de Montpellier, Fr.; Französischer Ahorn, Ger.
Engravings. Pluk. Alm., t. 251. f. 3.; Tratt. Arch., l. No. 20.; Duh. Arb., l. t. 10. f. 8.; Schmidt Arb., l. t. 14.; and Krause, t. 101.; our fig. 131. in p. 458.; and the plate of this species in our Second Volume.

Spec. Char., &c. Leaves cordate, 3-lobed; lobes almost entire, and equal. Corymbs few-flowered, pendulous. Fruit smooth, with the wings hardly diverging. A low tree or shrub, found in the south of Europe, and introduced in 1739; flowering in May.

Description, &c. In its native countries, (the south of France and Italy,) this tree attains the height of 30 ft. or 40 ft.; though, in rocky exposed situations, it often forms only a large bush. The trunk is covered with a reddish brown bark. The leaves are chiefly 3-lobed, with an entire margin: they are dark green, and bear a general resemblance to those of *A. campêtre*, which are about the same size, but of a paler green, and 5-lobed. The leaves, in mild seasons, remain on during great part of the winter, more especially in France. The flowers are produced just before the leaves: they are pendulous, on peduncles 1 in. or more long, disposed in dichotomous panicles, or corymbs, one from almost every bud, and consist of from 6 to 10 flowers. The flowers are of a pale yellow colour, and form a great source of attraction to bees. The tree of this species in the London Horticultural Society's Garden has the branches rather ascending, so as to form somewhat a fasciculate head; but in old trees, the head is roundish and spreading.

Geography, History, &c. The tree abounds, in a wild state, in the south of France, and also in Spain and Italy, chiefly on rocky exposed situations. It is also much planted in the south of France as hedges, on account of the persistency of the leaves, which remain on during a great part of winter. The tree was introduced into England in 1739, and is to be met with in most botanic gardens. The wood, which is hard and heavy, is used in France by turners and cabinet-makers; but, in England, the tree may be considered as purely one of ornament; and, as such, it well deserves a place in every collection. It is easily propagated by seeds or by layers.

Statistics. The largest tree in the neighbourhood of London is at Kenwood, where, in 55 years, it has attained the height of 47 ft.; there are two very handsome trees at Ham House, each nearly 30 ft. high: at Kew, 30 years planted, it is 29 ft. high; in the Oxford Botanic Garden, 40 years planted, it is 25 ft. high; in Worcester-shire, at Croome, 30 years planted, 25 ft. high; in Staffordshire, at Trent-ham, 26 years planted, 27 ft. high. In Scotland, in the Perth Nursery, 14 years planted, and 14 ft. high. In France, in the Jardin des Plantes, 139 years planted, and 55 ft. high. In Saxony, at Wörlitz, 45 years planted, 39 ft. high; in Austria, at Vienna, in the garden of the University, 40 years planted, and 36 ft. high; and at Laxenburg, 35 years planted, and 30 ft. high. In Prussia, in the Berlin Botanic Garden, 18 ft. high. In Bavaria, in the Botanic Garden at Munich, 24 years planted, and 20 ft. high.

Commercial Statistics. Plants, in the London nurseries, are 1s. 6d. each; at Bollwyler, 1 franc 50 cents; at New York, ?.

Y 18. *A. CAMPESTRIS* L. The common, or Field, Maple.

Identification. Lin. Spec., 1497; Hayne, Dend., p. 211; Dec. Prod., l. p. 595; Don's Mill., l. p. 649.

Synonymes. Erable champêtre, Fr.; kleiner Ahorn, Feld Ahorn, Ger.

Engravings. Engl. Bot., t. 304; Fel. Dan., t. 1288; Reitter and Abel, Abbild., t. 25; Willd. Abbild., t. 213; our fig. 132, in p. 458; and the plate of *A. campêtre* var. *austriacum* in our Second Volume.

Spec. Char., &c. Leaves cordate, with 5-toothed lobes. Racemes erect.

Wings of fruit much divaricated. (*Don's Mill.*, i. p. 649.)

Varieties. There are four forms of this species.

Y *A. c.* 1 *hebecarpum* Dec. Prod., i. p. 594. *The downy-fruited Field Maple.*—Fruit clothed with velvety pubescence. *A. campêtre* Wallr. in *Litt. Tratt. Arch.*, i. No. 7; *A. molle* Opiz. (*Don's Mill.*, i. p. 649.)

Y *A. c.* 2 *foliis variegatis.* *The variegated-leaved Field Maple.*—Next to the handsomest of all the variegated-leaved maples; the leaves preserving, with their variegation, the appearance of health, and the blotches, and stripes of white, or whitish yellow, being distinctly marked.

Y *A. c.* 3 *collinum* Wallr. in *Litt. Dec. Prod.*, i. p. 594. *The hill-inhabiting Field Maple.*—Fruit smooth. Lobes of leaves obtuse. Flower smaller. *A. affine* and *A. macrocarpum* Opiz. Native of France. (*Don's Mill.*, i. p. 649.)

Y *A. c.* 4 *austriacum* Tratt. Arch., i. No. 6. *The Austrian Field Maple.*—Fruit smooth. Lobes of leaves somewhat acuminate. Flowers larger than those of the species. Native of Austria, Podolia, and Tauria. (*Don's Mill.*, i. p. 649.) This variety is larger in all its parts than the original species, and is of much freer growth; the main stem rises erect and straight, and sends out its branches regularly on

every side, so as to form a sort of cone, almost like a fir, as exhibited in the plate of this variety, in our Second Volume. A very handsome tree, from which our drawing was taken, exists in the garden of the London Horticultural Society, and, not far from it, a round-headed shrubby tree of the original species. A subvariety of this sort, with variegated leaves, is propagated in the Bollwyller Nursery.

Description. *A. c. hebecarpum* is the British form of this species; and it is thus characterised by Smith in his *Eng. Flora*, ii. p. 231. A rather small tree, with spreading branches; the bark corky, and full of fissures; that of the branches smooth. Leaves about 1½ in. wide, downy while young, as are their footstalks, obtusely 5-lobed, here and there notched, sometimes quite entire. Flowers green, in clusters that terminate the young shoots, hairy, erect, short, and somewhat corymbose. Anthers hairy between the lobes. Capsules downy, spreading horizontally, with smooth, oblong, reddish wings. The character of the flowers, in being produced upon the young shoots, ascribed to the British field maple by Smith, is one which it possesses, and one in which it differs markedly from certain exotic kinds; as *A. eriocarpum*, *A. rubrum*, and *A. monspessulanum*; the flowers of which species are produced from buds distinct from those out of which the shoots are developed. In Britain, it seldom attains the height of more than 20 ft., except in a state of cultivation.

Geography. This species is found throughout the middle states of Europe, and in the north of Asia. According to Pallas, it is found in New Russia, and about Caucasus. In Britain, it is common in hedges and thickets, in the middle and south of England; but is rare in the northern counties and in Scotland. It is not a native of Ireland, and, perhaps, not of Scotland.

Properties and Uses. The wood weighs 61 lb. 9 oz. a cubic foot, in a green state, and 51 lb. 15 oz. when perfectly dry. It makes excellent fuel, and the very best charcoal. It is compact, of a fine grain, sometimes beautifully veined, and takes a high polish. It was celebrated among the ancient Romans for tables. In France, it is much sought after by turners, cabinet-makers, and the manufacturers of domestic utensils. The wood of the roots is frequently knotted; and, when that is the case, it is used for the manufacture of snuff-boxes, pipes, and other fanciful productions. The young shoots, being tough and flexible, are employed by the coachmen, in some parts of France, instead of whips. The tree is much used in France for forming hedges, and for filling up gaps in old fences. It is also employed in topiary works, in geometrical gardens, being found to bear the shears better than most other trees. The leaves and young shoots are gathered green, and dried for winter provender for cattle. The sap yields more sugar, in proportion to the quantity taken, than that of the sycamore; but the tree does not bleed freely. In Britain, the tree is seldom planted for any other purpose than that of ornament, in which it is effective by adding to the variety of a collection, rather than to its positive beauty. The variegated variety is showy, and, if a mule could be procured with red flowers, by cross-fecundation with *A. rubrum*, we should then have a singularly handsome little tree. Even a red tinge added to the autumnal foliage would be valuable. For the purpose of cross-fecundation, a plant of *A. campêtre* would require to be forced forward in a green-house, or a plant of *A. rubrum* retarded in an ice-house, as the two species flower at different periods.

Soil, Situation, &c. A dry soil suits this species best, and an open situation; but, to attain a timber-like size, it requires a deep free soil, and a situation sheltered by other trees. So circumstanced, it attains the height of 30 ft. or 40 ft., both in France and England, as may be seen in Eastwell Park, in Kent, and at Caversham Park, near Reading. In the nurseries, plants of this species are raised from seeds, most of which often remain eighteen months in the ground before they come up, though a few come up the first spring. The varieties are propagated by layers.

Casualties. The mistletoe is sometimes found growing on this species.

Statistics. In the environs of London, at Kew, *A. campêtre*. 50 years planted, is 26 ft. high; in

Dorsetshire, at Melbury Park, 100 years planted, and 38 ft. high, the trunk 2 ft. 9 in. in diameter, and the head 37 ft. in diameter, growing in stony clay; in Essex, at Braystock, 80 years planted, and 50 ft. high; in Surrey, at Farnham Castle, 50 years planted, 30 ft. high; in Suffolk, at Finborough Hall, 70 years planted, and 40 ft. high. In Denbighshire, at Llanbede, 20 years planted, 34 ft. high. In Scotland, near Edinburgh, at Hopeton House, 16 ft. high; in Argyllshire, at Hafton, 36 years planted, and 36 ft. high; in Clackmannanshire, in the garden of the Dollar Institution, 12 years planted, and 20 ft. high; in Forfarshire, at Airlie Castle, 10 years planted, and 14 ft. high; in Stirlingshire, at Blairlogie, 302 years old, and 55 ft. high, the diameter of the trunk 4 ft., and of the head 45 ft., the soil, a light loam on dry gravel, and the situation exposed. In France, in the Botanic Garden at Toulon, 18 years planted, and 45 ft. high. In Saxony, at Wörlitz, 65 years planted, and 40 ft. high; in Austria, in the University Botanic Garden at Vienna, 20 years planted, and 25 ft. high; at Laxenburg, 60 years planted, and 45 ft. high; at Kopenzel, 45 years planted, and 30 ft. high; at Hildersdorf, 40 years planted, and 21 ft. high. In Prussia, at Sans Souci, 40 years planted, and 55 ft. high. In Bavaria, at Munich, 24 years planted, and 20 ft. high.

Commercial Statistics. Plants, in London, seedlings 10s. a 1000, of a larger size 20s. a 1000; the variegated-leaved variety 2s. 6d. a plant. At Bollwyller, the broad-leaved variegated subvariety, 1 franc 50 cents a plant; at New York, ?.

Y 19. *A. CRÉTICUM* L. The Cretan Maple.

Identification. Lin. Spec., 1497.; Dec. Prod., l. p. 594.; Don's Mill, l. p. 649.

Synonymes. *A. heterophyllum* Willd. *Eu.*; *A. sempervirens* L. *Mant.*; *E'rabre de Crète*, *Fr.*; *Cresischer Ahorn*, *Ger.*

Engravings. Tratt. Arch., l. No. 19.; Duh. Arb., l. p. 28. t. 10. f. 9.; Alp. Exot., 9. t. 8.; Pöczek Orient., 197. t. 85.; Schmidt Arb., t. 15.; Krause, t. 120.; our fig. 132. in p. 459.; and the plate of this species in our Second Volume.

Spec. Char., &c. Leaves permanent, cuneated at the base, acutely 3-lobed at the top. Lobes entire, or toothleted; lateral ones shortest. Corymbs few-flowered, erect. Fruit smooth, with the wings hardly diverging. (*Don's Mill*, i. p. 649.) A diminutive slow-growing sub-evergreen tree; native of Candia, and of other islands in the Grecian Archipelago. Introduced in 1752.

Description, &c. This species is seldom seen in British collections, and then more frequently as a shrub than as a tree; but it is readily known from all the others, and from *A. monspessulanum*, to which it makes the nearest approach, by its being nearly evergreen; and by the great variety of the forms of its leaves; and, according to the specific character, by the flowers being erect; those of *A. monspessulanum* being pendulous. In young plants, the leaves are seldom much lobed; and this, we suppose, has given rise to the species or variety known in gardens as *A. heterophyllum*, which appears to be only *A. créticum* in a young state. We are only aware of three plants of this species which have assumed the character of trees in the neighbourhood of London; viz. that at Syon, figured in our Second Volume, which is 28 ft. high, flowers freely, and produces seeds almost every year; one in the Chelsea Botanic Garden, about 8 ft. high, which has stood there since the time of Miller; and a third, of equal age, which was in the Mile End Nursery, and which, in 1834, was 10 ft. high. This last tree has since been sold, and removed to the garden of the Rev. T. Williams, at Hendon, Middlesex. This species is generally propagated by layers; though it might, probably, be grafted on the Montpellier maple. Where a miniature arboretum is formed in a small garden, this species may be considered valuable, as exemplifying the order *Aceræceæ*, in a space not larger than what would be required for a herbaceous plant.

Statistics. The only specimens worth recording in Britain are those already referred to at Syon, Chelsea, and Hendon. The plants in the London Horticultural Society's Garden, and at Messrs. Loddiges's, are not above a foot and a half high. In France, in the Jardin des Plantes, a tree 130 years planted is 31 ft. high. In Saxony, at Wörlitz, one 55 years planted is 40 ft. high.

Commercial Statistics. Plants, in the London nurseries, cost 5s. each; at Bollwyller, where it is considered as synonymous with *A. heterophyllum*, 2 francs each; at New York, ?.

App. i. *Doubtful Species of Acer.*

We have not been able to satisfy ourselves respecting the distinctness of *A. Opalus* and *A. opulifolium*; and we are very much inclined to think that the sort which we have figured as *A. barbätum* is a European species, and, consequently, not the *A. barbätum* of Michaux. To us, it appears that the *A.*

barbatum, of which a leaf is figured in the *Penny Cyclopædia*, and several in our fig. 125. p. 452., may possibly be only a variety of *A. Pseudo-Plátanus*, diminished in all its parts. *A. coriáceum*, which we consider as a synonyme of *A. Opalus*, and which, in the *Penny Cycl.*, is considered a synonyme of *A. créticum*, may be a more distinct sort than we think it is, from having seen only small plants of it. These small plants have leaves as nearly as possible of the same form as those of *A. Opalus*, exhibited in fig. 126. p. 453.; and, therefore, the *A. coriáceum* mentioned in the *Penny Cycl.* as a synonyme of *A. créticum*, must refer to a different plant from the specimen of *A. coriáceum* in Loddiges's arboretum. *A. Lobèli Tenore*, of which there is a tree, 20 ft. high, at Croome, in Worcestershire, certainly appears very distinct from *A. platanoides* (of which we have set it down as a variety) in the foliage of the plants about 2 ft high in the London nurseries; but not so in the specimens which we have received from Croome. *A. nigrum*, which we consider as a variety of *A. saccharinum*, ought to be proved by raising plants from seeds, which can be procured from America at the same price as those of *A. saccharinum*. It would not surprise us if *A. platanoides* and *A. saccharinum* were ultimately to turn out to be races of one and the same species. Under the name of *A. híbridum*, it is probable that there is more than one variety or species in cultivation. The *A. híbridum* of Bose, with coriaceous leaves, profoundly trifid, seems to be different from the *A. híbridum* of Baudrillart, which he describes as intermediate between *A. monspessulanum* and *A. tatáricum*. In *Don's Miller*, there is *A. obtusifolium Flor. Græc.*, t. 361., a native of Crete; and there is also *A. obtusatum Kit.*, a native of Hungary: these may possibly be names applicable to one and the same species. *A. lobatum Don's Miller*, i. p. 651., a native of Siberia, and said to have been introduced in 1820, is there designated an "extremely doubtful" species.

App. ii. *Anticipated Species of Acer.*

All the species of this family are so interesting and ornamental, that it is desirable to procure additions to those already introduced into Britain, from every quarter of the globe, and by every resource which art supplies. It appears, from *Don's Miller* and *Royle's Illustrations*, that there are one or two European species, one from N. America, several from the Himalayas, and some from Japan, all likely to prove hardy, which are at present wanting in British collections. We shall give the names of these, in the hope of directing to them the attention of travellers, collectors, and patriotic amateurs.

☞ *A. ibéricum* Bieb., a tree 20 ft. high, native of Iberia, with greenish yellow flowers, and shining 3-lobed leaves, is described in Bieberstein's *Flora Turico-Caucasica*, vol. ii. p. 447.; in *Dec. Prod.*, i. p. 594.; and in *Don's Mill.*, i. p. 649.

☞ *A. obtusifolium* Sibth. and Smith, a native of Crete, a tree 15 ft. high, with 3-lobed crenated leaves, is described in the *Flora Græca*, and in *Don's Mill.*; in the latter work it is mentioned as being of doubtful introduction.

☞ *A. parvifolium* Tausch is described as having 5-lobed leaves, and nodding corymbs of flowers. It is a native of the south of Europe, growing to the height of 20 ft., by some confounded with *A. créticum*, and by others with *A. monspessulanum*.

☞ *A. glabrum* Torrey has smooth roundish 5-lobed leaves, and is found in North America, on the Rocky Mountains. Very little appears to be known of this species.

☞ *A. laevigatum* Wall. has oblong, acuminate, smooth, shining, leaves, and white flowers. It is a native of Nepal, on high mountains, where it forms a tree 40 ft. high. It is described and figured by Dr. Wallich in his *Plant. Asiat. rar.*, 2. p. 3. t. 104.

☞ *A. acuminatum* Wall., a tree of Nepal, with leaves varying from 3-lobed to 5-lobed, is described in *Don's Prod. Fl. Nep.*, p. 249.

☞ *A. cultratum* Wall., a tree of Himalaya, with cordate, 7-lobed, acuminate leaves, is described by Wallich in his *Planta Asiaticæ rariores*, vol. ii. p. 4.

☞ *A. caudatum* Wall., a tree from the highest regions of Nepal, with long pointed leaves, is described in the work last quoted, vol. ii. p. 4., and in *Don's Mill.*, i. p. 648.

☞ *A. villosum* Wall., a tree 50 ft. high, native of the high alps of the Himalaya, near to perpetual snow, has cordate 5-lobed leaves, and fragrant flowers. It is described in Dr. Wallich's work, and in *Don's Mill.*, as before quoted.

☞ *A. sterculiaceum* Wall., is a tree 50 ft. high, with a trunk 3 ft. in diameter. The leaves are very large, with long petioles, and the flowers are white. It is found in Nepal, on Mount Shiopore.

☞ *A. dissectum* Thunb. is a tree of Japan, with leaves 9—10-parted, and oblong acuminate lobes, with a red corolla. It is described by Thunberg in his *Flora Japonica*, p. 160.; in *Dec. Prod.*, i. p. 955.; and *Don's Mill.*, i. p. 650.

Y *A. japonicum* Thun., with roundish palmate leaves, is a Japan tree, 20 ft. high, with the bark of the branches, and the corollas purple, and the fruit woolly. It is described in Thunberg's *Flora Japonica*, p. 161.; in *Dec. Prod.*, i. p. 595.; and in Don's *Mill.*, i. p. 650.

Y *A. septenlobum* Thun., with smooth, acuminate, 7-lobed leaves, is a Japan tree, 40 ft. high, described by Thunberg, and, after him, by De Candolle and G. Don, as before quoted.

Y *A. pictum* Thun., with smooth, palmate, 7-lobed leaves, is a Japan tree, 30 ft. high, with ash-coloured branches, and leaves variegated with white. Described as above.

Y *A. trifidum* Thun., with undivided and trifid entire leaves, and twigs smooth and purplish, is a Japan tree, 20 ft. high, described in the works quoted.

Y *A. truncatum*, described in Bunge's *List of Plants of the North of China*, noticed in p. 176.


New Sorts from Cross-Fecundation. *A. obtusatum* has very much the appearance of a hybrid between *A. Pseudo-Platanus* and *A. O'palus*; but, whether this be the case or not, there seems no reason for doubting that cross-fecundation might be effected in this genus, as well as it is in various others. The objects ought to be, to get more colour into the flowers, and more red into the leaves. *A. monspessulanum*, with flowers as red as those of *A. rubrum*, would be a fine variety. Possibly some of the species might be fecundated by some species of the genus *Negundo*, which would lead to great changes in the leaves. As a number of the species of *Acer* do not flower and perfect seeds, till they become considerable trees, it would be desirable, when experiments are to be made by cross-fecundation, and the person wishing to try these experiments possesses only young trees, to graft them with scions from trees which already flower and fruit. In collecting species for this purpose, care must be taken, either to procure plants producing hermaphrodite flowers, or plants having male and female flowers on the same tree; unless, indeed, advantage is proposed to be taken of the circumstance of the tree being only of one sex, to fecundate its flowers, if female, or pistilliferous, with the male flowers of another species; or, if stamiferous, to fecundate with them the female flowers of some other kind. We are strongly inclined to believe that some of what are now considered aboriginal species of *Acer* are only the result of cross-fecundation, produced by accident; and hence we anticipate a number of new forms, when the attention of cultivators is powerfully directed to this object. *Negundo fraxinifolium* will, in all probability, fecundate, and be fecundated by, acers of different kinds; and this alone would produce something which would amply repay the curious cultivator. The tree produces flowers at an early age; and, as there are in almost every part of the country abundance of trees of *Acer campéstris*, and *A. Pseudo-Platanus*, which flower every year, we would recommend a trial to be made between these sorts without delay. Success may not attend the first trial, but the object ought to be persisted in till some result be obtained. *A. créticum*, *A. monspessulanum*, and *A. campéstre* will, doubtless, fertilise together; because, in foliage, in mode of growth, in time of flowering, and in the form of the fruit, they are very much alike; and something might be done with them with very little trouble.

Additional Sorts from accidental Forms of Growth. The eagle's claw maple is a well known and very curious variety. Whenever any appearance of the same kind is observed in any of the other species, it ought to be continued by grafting. By carefully looking out for sports from the average forms, we shall probably, at some future day, have weeping maples, as well as the weeping ash; fastigate maples, like the Lombardy poplar; and purple-leaved kinds, like the purple-leaved beech or common hazel. Though scarcely any of the variegated maples now known can be considered as very handsome, yet a new and beautiful variety of them may one day be procured.

App. iii. *Half-hardy Species of Acer.*

A. palmatum Thun., described p. 422., is, in all probability, only half-hardy; or, at all events, it is safe to treat this, and the other Japan species, and also those from Nepal, when once they are procured, as half-hardy, when in their young state. *A. oblongum*, described in p. 415. according to Mr. Boyle, descends to the lowest level of all the species found in Nepal, and is, therefore, probably the tenderest of those from that part of the world.


A'cer oblongum. The oblong-leaved Maple.
Leaves of the natural size.



113

Acer tataricum. The Tartarian, or entire-leaved, Maple.
Leaves and fruit of the natural size.

114




Acer spicatum. The spike-flowered, or mountain, Maple.
Leaves and fruit of natural size.




115

Acer striatum. The striped-
Leaves of

116



bark, or *Pennsylvanicum*, Maple.
the natural size.




Acer macrophyllum. The
Part of leaf, and



5-lobed Maple.
of the natural size.

117



Acer macrophyllum. The
Smaller leaves, also of the natural size, to

118



ge-leaved Maple.
ow how much they vary on the same tree.

118




Acer platanoides. The Platanus-
Leaves and fruit of

119




like, or *Norway*, Maple.
The natural size.

119




Acer Lobëlii. L'Obel's Maple.
Leaves of the natural size.

120



Acer platanoides laciniatum.
The cut-leaved Platanus-like, or Eagle's claw, Maple.
Leaves of natural size.



Acer saccharinum.

The leaves and fruit

122



the Sugar Maple.
the natural size.

122



Acer Pseudo-Platanus. The
Leaves and fruit

123



False Plane, or *Sycamore*, Maple.

of the natural size.



123

Acer obtusatum. The obtuse-
Leaves of the

124



lobed-leaved, or Neapolitan, Maple.

natural size.

124



Acer barbatum. The bearded-calyxed Maple.
Leaves and fruit of the natural size.


125



A`cer Opalus. The Opal, or Italian, Maple.

Leaves of the natural size.


126



Acer circinatum. The round-leaved Maple.

Leaves of the natural size.


125



Acer palmatum. The palmate-leaved Maple.


Leaves of the natural size.

128



Acer eriocarpum. The woolly-fruited Maple.

Leaves and fruit of the natural size.




A`cer ru`brum. The red-flowered Maple.
Leaves and fruit of the natural size.

130



Acer monspessulanum, and *A. campêtre*. The Montpellier Maple,
and the *common*, or field, Maple.


Leaves and fruit of the natural size.



Acer creticum. The Cretan, or various-leaved, Maple.

Leaves and fruit of the natural size.

132



Leaves and fruit from an old tree at Syon.



Leaves from a young tree in the garden of the London Hort. Soc.



GENUS II.



NEGUNDO Mœnch. THE NEGUNDO, or BOX ELDER. *Lin. Syst.*
Dioecia Pentandria.

Identification. Mœnch. Meth., 334.; Dec. Prod., 1. p. 596.; Don's Mill., 1. p. 647. and 651.

Synonymes. *Acer* Lin.; *Negundium* Rafinesque.

Derivations. This genus was constituted from *Acer Negundo* L.; but the meaning of the latter word is unknown. Probably, it may be merely the Illinois name of Giguères (from *giguer*, to romp, alluding to the tremulous and playful motion of the long pinnated leaves) Latinised.

Gen. Char. Sexes dioecious. Flowers without a corolla. Calyx with 4—5 unequal teeth. Male flowers upon thread-shaped pedicels, and disposed in fascicles: anthers 4—5 linear, sessile. Female flowers disposed in racemes. Leaves impari-pinnately divided. (*Dec. Prod.*, i. p. 596.)

♂ 1. *N. FRAXINIFOLIUM* Nutt. The Ash-leaved Negundo.

Identification. Nutt. Gen. Amer., 1. p. 253.; Dec. Prod., 1. p. 596.; Don's Mill., 1. p. 651.

Synonymes. *Acer Negundo* L.; *Mich. Arb.*, *Tratt. Arch.*; *N. aceroides* Mœnch; *Negundium americanum* Rafin.; the Ash-leaved Maple, the Black Ash; *Erable à Feuilles de Frêne*, Fr.; *Erable à Giguères*, Illinois; *Eschenblättriger Ahorn*, Ger.

Engravings. Mich. Arb., 2. t. 16.; *Tratt. Arch.*, 1. No. 10.; Schmidt, Arb., 1. t. 12.; Wats. Dend., t. 172.; E. of Pl., No. 14294.; and the plate of this species in Vol. II. It is the female plant that is represented in these figures, except, perhaps, that of *Tratt. Arch.*, which we have not seen.

Spec. Char. &c. Leaves of from 3 to 5 leaflets, the opposite ones coarsely and sparingly toothed, the odd one oftener three-lobed than simple. (*Dec. Prod.*, i. p. 596.) A tree from North America, in 1688, growing to the height of 50 ft. or 60 ft., and flowering in April.

Variety.

♂ *N. f. 2 crispum* G. Don. *The curled-leaved Ash-leaved Negundo, or Box Elder.*—Figured in our Second Volume. The plant of this variety in the arboretum of the London Horticultural Society is of the male sex: the inflorescence consists of pendulous panicles of flowers, that are green, with some redness from the colour of the anthers; and each is placed upon a slender peduncle of about 1 in. long.

Description. As far as is known, only the female plant of the species has, as yet, been introduced into England. The flowers are produced, profusely, about the middle of April, and appear with the leaves: they are green, small, and in slender pendulous racemes. The racemes of flowers are inconspicuous, so that the flowering of the tree may occur without being noticed, unless the tree be carefully watched at its flowering season. The racemes of fruits that succeed to the flowers, increasing gradually to the length of 6 in. or 7 in., become obvious as the season advances, and appear conspicuously among the foliage. The leaves are opposite, and are from 6 in. to 15 in. long, according to the vigour of the tree, and the moisture of the soil in which it grows. Each leaf is composed of two pairs of leaflets, with an odd one. The leaflets are petioled, oval-acuminate, and sharply toothed: towards autumn, the common petiole becomes of a deep red. The tree grows rapidly, arriving at maturity in 15 or 20 years; and in a suitable soil and situation, attains the height of 40 ft. or 50 ft.: it branches at a small distance from the ground, and forms a loose and wide-spreading head. The bark of its trunk is brown; and there is a disagreeable odour in the inner bark: that of the young branches is of a smooth rush-like appearance, interrupted by very few buds, and of a most beautiful pea-green, like the shoots of the common jessamine (*Jasminum officinale*), but on a larger scale. The proportion of the sap-wood to the heart-wood is large, except in very old trees: in these the heart-wood is variegated with rose-coloured and bluish veins.

Geography, &c. This tree is a native of the United States, and of Canada: in the latter country, it is abundant about the Red River, in N. lat. 54°, which is its most northern limit. It is seldom found in the northern states of the Union, or in the maritime parts of the southern states. It commences on the banks of the Delaware, in the neighbourhood of Philadelphia, and increases towards the Alleghany Mountains, to the west of which it is most abundant; and, instead of being confined, as in the upper parts of Virginia and of the Carolinas, to the river sides, it grows in the woods, with the locust (*Robinia*), wild cherry (*Cerasus virginiana*), and coffee tree (*Gymnocladus*). But in the bottoms which skirt the rivers, where the soil is deep, fertile, constantly moist, and often inundated, this tree is most abundant, and attains the largest size. Even here, however, the tree seldom exceeds the height of 50 ft., with a trunk 20 in. in diameter; and trees of these dimensions, Michaux adds, are found only in Tennessee, and in the back parts of Georgia, which lie far to the south. In Kentucky, it seldom attains above half this height.

History. This species was cultivated by Bishop Compton in 1688; and it was introduced into France by Admiral Gallisonière, in the time of Du Hamel. As already observed, only the female plant of the species is in British gardens, though the male plant exists in the form of the variety *N. f. crispum*. Both male and female plants appear to be in cultivation in France; because, according to Baudrillart, the Administration of Forests at Paris received a quantity of seeds from the neighbourhood of Lyons, from which a number of young plants were raised, and distributed through the national forests. The old tree of this species, in the garden at Fulham, planted by Bishop Compton in 1688, measured, in 1793, 6 ft. 4 in. in girth at 3 ft. from the ground, and was computed to be 45 ft high. In 1809, it measured 7 ft. 1½ in. in girth; and in 1835, as appears in p. 43., the dimensions are scarcely varied. The tree, from the rapidity of its growth, its showy appearance, the fine green of its shoots, its large pinnate leaves (which move with the slightest breeze) and spreading head, and its faculty of growing in almost any kind of soil, is a general favourite, and is to be found in most collections.

Properties, Uses, &c. The wood of this tree has a fine even grain, and is saffron-coloured, slightly mixed with violet, but it is rather tender; the proportion of the soft wood to the heart-wood is so great, that it has never been used for any purpose in America except for that of fuel: in Europe, it is used for cabinet-making, particularly for inlaying; it works well, and is elastic and sonorous. According to some, sugar has been extracted from the sap; but Michaux denies this to be the case. He suggests that, from its rapid growth, after being cut down to the ground, it might form a valuable underwood, to be cut every 3 or 4 years, for fuel, charcoal, or other purposes. It has been tried in this way in France; but, unless the soil be kept constantly humid, the stool is found to decay in a very few years. Baudrillart considers it the least useful of all the tree maples. In England, it is solely to be considered as an ornamental tree; and, as we have before observed, it is one of considerable show; and, being also one of rapid growth, it well merits the attention of planters, in situations where immediate effect is an object.

Soil, Situation, &c. The same soil and situation which suit the *Acer* will answer for the *Negundo*. The plant grows with amazing rapidity, when the soil is deep and somewhat moist; but, as it is not a long-lived tree, it should not be placed in situations where the permanent effect of wood is of importance. When raised from seed, it should always, if possible, be sown in the autumn, as soon as practicable after gathering, as it keeps with difficulty till the spring, even when mixed with sand.

Statistics. In the environs of London, the oldest tree is that already mentioned at the Bishop of London's; and the largest one, at Kenwood, 35 years planted, and 47 ft. high. There are trees at Syon, at Kew, and at Purser's Cross, above 40 ft. high; in Buckinghamshire, at Temple House, 40 years planted, and 40 ft high; in Yorkshire, at Grimston, 10 years planted, and 30 ft. high. In Wales, in Pembroke-shire, at Golden Grove, 30 years planted, and 25 ft. high. In Scotland, in the Glasgow Botanic Garden, 14 years planted, and 14 ft. high. In Ireland, in Galway, at Coole, 39 ft. high. In France, Michaux informs us, a row of these trees was planted in the Jardin des Plantes, in the Rue

de Buffon, which gave an excellent idea of their appearance in their native forests; the highest these trees now remaining, estimated at upwards of 60 years of age, are 51 ft. in height, with heads 54 ft. in diameter. At Toulon, in the Botanic Garden, a tree 20 years planted, is 50 ft. high. In Saxony, at Wörlitz, a tree, 65 years planted, is 40 ft. high. In Austria, in Held's Nursery, at Vienna, 20 years planted, and 25 ft. high; at Brück, on the Leytha, 48 years planted, and 80 ft. high, with a head 48 ft. in diameter, close by water. In Prussia, at Sans Souci, 45 years planted, 24 ft. high. In Bavaria, at Munich, 24 years planted, and 30 ft. high. In Hanover, in the Botanic Garden at Göttingen, 20 years planted, and 40 ft. high.

Commercial Statistics. In London, plants of the species are 1s. each, of the variety *N. f. crispum* 1s. 6d. each, and seeds of the species 4s. a quart. At Bollwyller, plants of the smallest size of the species are 50 cents each, and larger plants 2 francs. At New York, plants are 25 cents each, and seeds 1 dollar a quart.

App. i. *Anticipated Species of Negúndo.*

‡ *N. mexicanum* Dec. is a native of Mexico, with trifoliate leaves, which grows to the height of 40 ft. It has been described under the name of *A. ternatum* Moc.; and also by De Candolle, who expresses a doubt as to whether it may not be a variety of *N. fraxinifolium*. Seeds or plants of it ought, if possible, to be obtained from Mexico.

‡ *N. cochinchinense* Dec. is a doubtful species, probably belonging to Sapindaceæ rather than Aceraceæ. It is a tree 25 ft. high, in the woods of Cochin-China, and would form a very desirable addition to the British arboretum. A variety of *N. fraxinifolium* (the leaves having their two lower and their terminate leaflets ternate) is mentioned by Dr. Hooker, as having been received by him from Dr. Schwinitz in the state of dried specimens. (*Fl. Bor. Amer.*, i. 114.)

App. I. *Other Aceraceæ.*

• *Dobinea vulgaris* Hamilt. (altered from the Nepalese name), a shrub 6 ft. high, a native of Nepal, with elliptical, oblong, acutely serrated leaves, is, in all probability, hardy, and would add to the variety of British collections. There no other species of this genus, and no other genus belonging to the order Aceraceæ has been yet discovered and recorded, either hardy or tender.

CHAP. XXIII.

OF THE HARDY TREES AND SHRUBS OF THE ORDER ÆSCULACEÆ.

Identification. Lindl. Key.

Synonymes. Castaneæ Link; Hippocastaneæ Dec.

Distinctive Character. Calyx campanulate, 5-lobed. Ovary roundish, trigonal. Seeds large and globose; albumen wanting. Embryo curved, inverted, with fleshy, thick, gibbous cotyledons not produced above ground in germination. Plumule large, 2-leaved. Deciduous trees, with opposite digitate leaves. The fruit large and bitter, sternutatory, abounding in potash and starch. The bark astringent and tonic; and the fruit containing æsculin, a febrifuge. The genera are two, which are thus contradistinguished:—

ÆSCULUS L. Capsule echinated.

PAVIA Boeh. Capsule smooth.

GENUS I.



ÆSCULUS L. THE HORSECHESTNUT. *Lin. Syst.* Heptándria Monogýnia.

Identification. *Lin. Gen.*, No. 462; *Hayne Dend.*, p. 43; *Dec. Prod.*, 1. p. 597; *Don's Mill.*, 1. p. 532.

Synonymes. Hippocastanum *Tourn.*; Marronier d'Inde, *Fr.*; Rosskastanie, *Ger.*

Derivation. The word Æsculus, derived from *cæsa*, nourishment, is applied by Pliny to a species of oak, which had an eatable acorn. The word Hippocastanum, from *hippos*, a horse, and *castanea*.

a chestnut, is said by some to have been given to this tree ironically, the nuts, though they have the appearance of sweet chestnuts, being only fit for horses; and by others, because, in Turkey, it is said the nuts are used for curing horses of pulmonary diseases. According to Evelyn, they are also given in England to horses that are broken-winded, and to other cattle that have coughs and colds.

Description. All the species (except one, which is a shrub) are deciduous trees, with deeply cut leaves, and showy flowers; and they are distinguished from the genus *Pavia* by the roughness of their fruit, and by the comparative roughness, also, of their leaves. To us it appears doubtful if the roughness of the fruit be a sufficient generic distinction, since it varies much in different individuals, and since, in some of the sorts, which have apparently been originated between *Æsculus* and *Pavia*, the fruit is as smooth, or nearly as much so, as in the proper *pavias*. It is highly probable that the two genera consist, in fact, of only two, or at most three, distinct species: however, all the different sorts in cultivation are so truly ornamental, that they may very conveniently be kept distinct, as races or botanical species. The common horsechestnut is invariably propagated by the nuts, which are sown when newly gathered, or in the following spring; and in either case they will come up the succeeding summer. All the other sorts, as being varieties of the species, are propagated by budding or grafting.

Æ 1. *Æ. HIPPOCÆSTANUM L.* The common Horsechestnut.

Identification. Lin. Sp., 488.; Willd. Baum., p. 14.; Hayne Dend., p. 43.; Dec. Prod., l. p. 597.; and Don's Mill., l. p. 652.

Synonymes. *Hippocæstanum vulgare Tourn.*; *Marronnier d'Inde, Fr.*; *gemeine Rosskastanie, Ger.*
Engravings. Woodv. Med. Bot., t. 128.; Plenck Icon., t. 293.; Willd. Abbild., t. 40.; and the plate of this species in our Second Volume.

Spec. Char., &c. Leaflets 7, obovately cuneated, acute, and toothed. A tree, growing to the height of 50 ft. or 60 ft., in cultivation since 1629, and flowering in May.

Varieties.

Æ *Æ. H. 2 flore pleno.* The double-flowered Horsechestnut. — This variety is recorded in nurserymen's catalogues, but is not common.

Æ *Æ. H. 3 variegata.* The variegated-leaved Horsechestnut. — The leaves are blotched with yellow, or yellowish white, but they have a ragged and unhealthy appearance, and are by no means ornamental.

Description. A tree of the largest size, with an erect trunk, and a pyramidal head. The leaves are large, of a deep green colour, and singularly interesting and beautiful when they are first developed. When folded in the bud, they are covered with pubescence, which falls off as the leaves become expanded, sooner or later, according to the dryness or moistness of the season. The growth, both of the tree and of the leaves, is very rapid; both shoots and leaves being sometimes perfected in three weeks from the time of foliageation; in which time, says Miller, I have measured shoots 1½ ft. long, with their leaves fully expanded. The flowers appear a short time after the leaves, and are white, variegated with red and yellow: they expand in May, and the fruit ripens about the end of September or the beginning of October. It is allowed to fall from the tree; but, if wanted for seed, must be soon afterwards gathered up, and either sown, or mixed with earth; because, if exposed to the atmosphere, it will lose its germinating faculties in a month. (*Bau-drillart.*) The buds are covered with a gummy substance, which protects their downy interior from the wet. Miss Kent observes that "we cannot have a better specimen of the early formation of plants in the bud, than in that of the horsechestnut." A celebrated German naturalist detached from this tree, in the winter season, a flower bud not larger than a pea, and first took off the external covering, which he found consisted of seventeen scales. Having removed these scales, and the down which formed the internal covering of the bud, he discovered four branch leaves surrounding a spike of flowers, the latter of which were so distinctly visible, that, with the aid of a microscope, he not only counted 68 flowers, but could discern the pollen of the stamens, and perceive that some was opaque, and some transparent. This experiment

may be tried by any one, as the flowers may be perceived with a common magnifying glass; but, as detaching the scales requires care, it would be advisable for an unpractised student to gather the bud in early spring, when the sun is just beginning to melt away the gum with which the scales are sealed together. (*Mag. Nat. Hist.*, vol. iii. p. 135.)

Geography. "The native country of the common horsechestnut," Mr. Royle observes, "is yet unknown, though stated, in some works, to be the north of India." He adds that he has "never met with it, though often visiting the lofty mountains of Northern India, where, if anywhere, it was likely to be found, and where the nearly allied Indian *pavia* is so abundant." (*Illustr.*, p. 135.) As the genus *Pavia* is a native both of India and America, and as *Æsculus ohioënsis*, which is nothing but a variety of the common horsechestnut, is a native of North America, it seems to us highly probable that the genus *Æsculus* will ultimately be found to belong to both continents.

History. The horsechestnut passed from Asia to Europe about the middle of the sixteenth century. The tree was first described by Matthioli, and afterwards by Clusius in his *Rariorum Plantarum Historia, &c.* He there says, that, in 1581, it was still considered as a botanical rarity; but that in 1588 there was a plant of the species at Vienna, that had been brought there twelve years before, but which had not then flowered. In France, it was first raised from seed procured from the Levant, in the year 1615, by one Bachelier whose flower-gardens at Paris were then celebrated. The largest tree of the kind in France, and which was considered as the parent stock from which all the others have been propagated, formerly existed in the gardens of the Temple. (*Beckmann's Hist.*, vol. i. p. 317.) A tree of this species was planted in the Jardin des Plantes, in 1650, which was the second plant introduced into France: it died in 1767; and a section of its trunk is still preserved in the Museum of Natural History. According to M. Jaume Sainte-Hilaire (see p. 147.), and his account appears to us the most probable, the horsechestnut passed from the mountains of Thibet to England in 1550, and was afterwards taken to Vienna by Clusius, and thence to Paris by Bachelier. The earliest notice which we have of the horsechestnut being in England is in Gerard's *Herbal*, where, in 1579, he speaks of it as a rare foreign tree. In Johnson's edition of the same work, in 1633, the horsechestnut is said to be growing in Mr. Tradescant's garden at South Lambeth. Parkinson, in 1629, says, "Our Christian world had first a knowledge of it from Constantinople." The same author placed the horsechestnut in his orchard, as a fruit tree, between the walnut and the mulberries. How little it was then known may be inferred from his saying, not only that it is of a greater and more pleasant aspect for the fair leaves, but also of as good use for the fruit, which is of a sweet taste, roasted and eaten, as the ordinary sort. Houghton (1700) mentions some horsechestnut trees at Sir William Ashhurst's at Highgate, and especially in the Bishop of London's garden at Fulham. Those now standing at Chelsea College were then very young. There was also a very fine one in the Post-house Garden, in Old Street, and another not far from the Ice-house, under the shadow of the Observatory in Greenwich Park. (*Mart. Mill.*) In Germany, as we have seen in p. 147., the horsechestnut, after having been planted at Vienna, soon found its way to Baden, where it was planted about the end of the sixteenth century, and where some of the trees are still in existence. The tree is now generally cultivated in the middle states of Europe, and also in North America.

Properties and Uses. The wood weighs, when newly cut, 60 lb. 4 oz. per cubic foot; and, when dry, 35 lb. 7 oz.; losing, by drying, a sixteenth part of its bulk. According to other experiments, the wood, green, weighs 62 lb. 3 oz.; half dry, 46 lb. 2 oz.; and quite dry, 37 lb. 3 oz. It is soft, and unfit for use where great strength, and durability in the open air, are required; nevertheless, there are many purposes for which it is applicable when sawn up into boards; such as for flooring, linings to carts, packing-cases, &c. In France, sabots are made from it; and it is said to be used by carvers, turners,

&c. Bouteher says, that it is suitable for water-pipes that are to be kept constantly under ground; and it is also recommended for this purpose by Du Hamel. The charcoal of the horsechestnut may be used in the manufacture of gunpowder; and the ashes of every part of the plant, more especially of the fruit, afford potash in considerable quantity. The bark, which is very bitter, is employed for tanning, and also for dyeing yellow; and it has been used medicinally as a substitute for Jesuits' bark. In Turkey, the nuts are ground, and mixed with horse food, especially when the horses are broken-winded: and, in their natural state, they are eaten by goats, sheep, and deer. Ruddy says the nuts are an excellent food for deer and hogs: they also possess a detergent quality, and serve, in some degree, as a soap. They are used in Ireland to whiten flaxen cloth, and for this purpose are rasped into water, in which they are allowed to macerate for some time. (*Nat. Hist. of the County of Dublin*, vol. i. p. 107.) The nuts, when ground into flour, and mixed in the proportion of one third with the flour of wheat, are said to add to the strength of bookbinder's paste. (*Mech. Mag.*, vol. viii. p. 223.) M. Vergaud has proposed to change the starch which can be extracted from the nut into sugar, and afterwards to employ it in distillation. (*Gard. Mag.*, vol. i. p. 318.) Parmentier, in his *Nouveau Dictionnaire d'Histoire Naturelle*, after noticing most of the uses to which this tree and its nuts have been applied, says, it appears, after all, that no use applicable to every-day purposes has yet been discovered respecting it. (See *Baudrillart, Traité, &c.*, tom. ii. p. 364.) In Britain, the horsechestnut can only be considered as an ornamental tree, and, as such, is well known, and needs no eulogium. It produces a splendid effect when in flower, either singly, in avenues, or on the margins of plantations. The finest avenue of these trees in England is that at Bushy Park. Gilpin objects to the horsechestnut, as being lumpish in its form; but in saying this he evidently judged of the tree merely with reference to picturesque beauty, to which it has few pretensions till it becomes very old: whereas, in point of floral beauty, it and its several varieties, or races, are unequalled by any tree of equal size which will endure our climate. A very remarkable vegetable principle, called æsculine, which is found in this genus, and which is said to possess alkaline properties, and to act as a febrifuge, is mentioned in Stephenson and Churchill's *Medical Botany*, vol. ii. Miss Kent states that the fruit is not only sometimes ground and given to horses, but that deer are very fond of it, and that it is given to sheep, in a raw state, or when made into a paste, after maceration in lime water, to fatten them. Soap and starch have also been procured from these nuts. (*Mag. Nat. Hist.*, iii. p. 135.)

Poetical and legendary Allusions. The horsechestnut, where allowed to attain its proper shape on a lawn, is certainly a most magnificent tree. Some authors have compared it to an immense lustre or chandelier, its long racemes of flowers tapering up from its drooping foliage like lights. A horsechestnut tree, in full flower, has been called by Daines Barrington a giant's nosegay; and in the *Mag. Nat. Hist.*, vol. iv. p. 238., an eloquent description of this tree has been given by Mr. Dovaston, who compares its racemes of flowers to those of a gigantic hyacinth. Miss Kent, in the same work, vol. iii. p. 135., calls it a Brobdnagian lupine. In Paris, the magnificent trees in the garden at the Luxembourg have been celebrated by Castel.

“ Là de marronniers les hautes avenues
S'arrondissent en voûte, et nous cachent les nues.”

The manner in which it scatters its flowers on the grass, and the comparative uselessness of its fruit and timber, make it an excellent emblem of ostentation.

Soil, Situation, &c. The horsechestnut requires a deep, free, loamy soil, and will neither attain an ample size, nor flower freely, except in a situation rather sheltered than exposed. In England, it seldom suffers by the frost in spring; and the severest of our winters do not injure either its young shoots or its buds, which are covered with a resinous gluten. The species is always

propagated by the nut, sown in autumn or spring, and covered with from 2 in. to 4 in. of soil. The cotyledons do not rise to the surface, as in the oak, the beech, and some other trees. Some nurserymen cause the nuts to germinate, before sowing them, in order to have an opportunity of pinching off the extremity of the radicle; by which means the plants are prevented from forming a taproot; or, at least, if a taproot is formed, it is of a much weaker description than it otherwise would be, and the number of lateral fibres is increased; all which is favourable for transplanting. When the tree is intended to attain the largest size, in the shortest time, the nut ought to be sown where the tree is finally to remain; because the use of the taproot is mainly to descend deep into the soil, to procure a supply of water, which, in dry soils and seasons, can never be obtained in sufficient quantities by the lateral roots, which extend themselves near the surface in search of nourishment and air.

Statistics. Æ. Hippocastanum in the Environs of London. At Syon, 80 ft. high, the diameter of the trunk 3 ft. 8 in., and of the head 48 ft.; at Enfield, 100 ft. high (see our plate in Vol. II); at Ham House, 60 ft. high; the diameter of the trunk, at 1 ft. from the ground, 6 ft. 4 in., and that of the head 48 ft.; a magnificent tree, the trunk dividing, at 8 ft. from the ground, in three immense branches, which are respectively 2 ft. 2 in., 2 ft. 3 in., and 1 ft. 11 in. in diameter, at 4 ft. from the point where they separate from the trunk; at York House, Twickenham, 100 years planted, and 70 ft. high.

Æ. Hippocastanum South of London. In Cornwall, at Caulen Penryn, 55 ft. high, the trunk 5 ft. in diameter, and the diameter of the head 86 ft. In Devonshire, in Veitch's Nursery, Exeter, 34 years planted, and 43 ft. high; at Luscombe, 34 years planted, and 44 ft. high. In Hampshire, in Wilkins's Nursery, Isle of Wight, 30 years planted, and 35 ft. high; in Alresford, 81 years planted, and 59 ft. high; at Southampton, 70 years planted, and 80 ft. high. In Somersetshire, at Nettiecombe, 60 years planted, and 60 ft. high. In Wiltshire, at Wardour Castle, 45 years planted, and 50 ft. high; and 10 years planted, and 22 ft. high.

Æ Hippocastanum North of London. In Herefordshire, at Eastnor Castle, 18 years planted, and 40 ft. high. In Gloucestershire, at Dodington, 60 ft. high, and the diameter of the head 80 ft. In Suffolk, at Finborough Hall, 80 years planted, and 100 ft. high; the diameter of the trunk, at 1 ft. from the ground, 5 ft., and of the head 30 ft. In Lincolnshire, at Noeton, 59 ft. high; said to be the largest in Britain. This is a most magnificent tree, with immense branches, extending over a space 305 ft. in circumference: the branches are so large as to require props; so that at a little distance it looks like an Indian banyan tree. In Oxfordshire, in a field adjoining the Botanic Garden at Oxford, 60 ft. high, the head 50 ft. in diameter; and a tree of the pendulous variety, also 60 ft. high. In Staffordshire, at Trentham, 60 ft. high. In Warwickshire, at Coombe Abbey, 100 years planted, 70 ft. high, the diameter of the trunk 7 ft. 3 in., and of the head 103 ft. In Derbyshire, at Keddleston, a tree, the head of which was broken to pieces by lightning soon after the family had been drinking tea under it: this head is 62 ft. in diameter, and the branches touch the ground all round: the trunk is 16 ft. in circumference. In Worcestershire, at Croome, 50 years planted, and 60 ft. high; in Yorkshire, at Grimston, 10 years planted, and 36 ft. high.

Æ. Hippocastanum in Scotland. In the environs of Edinburgh, at Hopeton House, 45 ft. high, the diameter of the trunk 4 ft. 3 in., and of the head 50 ft. In Renfrewshire, in the Glasgow Botanic Garden, 12 years planted, and 18 ft. high. In Banffshire, at Gordon Castle, 66 ft. high, the diameter of the trunk 1 ft. 7 in., and of the head 80 ft. In Forfarshire, at Cortachy Castle, 102 years planted, and 40 ft. high, the diameter of the head 43 ft. In Haddingtonshire, at Tynningham, 44 ft. high, the diameter of the head 45 ft.; at Fountain Hall, the trunk 8 ft. 2 in. in girth, at 3 ft. from the ground, and the diameter of the head 50 ft. In Kirkeudbrightshire, at St. Mary's Isle, 55 ft. high. In Fifeshire, 35 years planted, and 56 ft. high, the head 52 ft. in diameter. In Perthshire, in the Perth Nursery, 40 years planted, and 54 ft. high; at Taymouth, 50 ft. high. In Rosshire, at Brahan Castle, 60 ft. high, the diameter of the head 50 ft. In Clackmannanshire, at Tullibodie, 70 ft. high, the diameter of the head 40 ft. Sir Thomas Dick Lauder, speaking of horsechestnuts in Scotland, says, "A horsechestnut, at Halton, in the county of Edinburgh, measured 11 ft. 4 in. in girth, at 4 ft. from the ground. A horsechestnut, at Preston Hall, at 1 ft. from the ground, measures 14 ft. 3 in.; and at 3 ft. from the ground, 10 ft. 9 in. There are many trees near it of dimensions little inferior. The horsechestnuts on the lawn, which was formerly the garden of Dawick, the seat of Sir John Murray Nasmyth, Bart., a few miles above Peebles, in Tweeddale, are certainly the oldest and finest in Scotland; or, perhaps, we should say there are none equal to them in Britain. They stand 12 ft. apart from each other; but they support a mass of foliage that appears to be but one head, which takes a beautiful form, and covers an area of ground the diameter of which is 96 ft. The larger of the two is, in girth, immediately above the root, 16½ ft. The smaller tree is 12½ ft. in circumference at the base, and 10 ft. at 3 ft. high. These measurements are kindly communicated to us by Sir John Nasmyth himself; and, from what Dr. Walker has said of these two horsechestnuts, we have no doubt that they are from 180 to 190 years of age." (*Lauder's Gilpin*, i. p. 270, 271.)

Æ. Hippocastanum in Ireland. In the environs of Dublin, at Cypress Grove, 60 ft. high, the diameter of the trunk 3 ft., and of the head 75 ft. In Connaught, at Makree Castle, 71 ft. high, the diameter of the trunk 3 ft. 3 in., and of the head 46 ft. In Down, at Ballyleady, 60 years planted, and 40 ft. high, the diameter of the trunk 14 in., and of the head 30 ft. In Fermanagh, at Florence Court, 70 years planted, and 48 ft. high, the diameter of the head 80 ft.

Æ. Hippocastanum in Foreign Countries. In France, in Paris, there are some very fine specimens in the gardens of the Luxembourg; though the original trees, mentioned in p. 147, are dead. At Nerrières, near Nantes, there is a tree 100 years planted, which is 70 ft. high. In Saxony, at Wörlitz, 65 years planted, and 40 ft. high. In Austria, at Vienna, at Laxenburg, 50 years planted, and 35 ft. high; at Kopenzel, 40 years planted, and 30 ft. high. In Prussia, at Sans Souci, 60 ft. high. In Bavaria, at Munich, 24 years planted, and 40 ft. high.

Commercial Statistics. Plants, in the London nurseries, are, seedlings 7s. a thousand; transplanted plants, from 2 ft. to 5 ft. high, 5s. a hundred; and the variegated variety 2s. 6d. a plant; at Bollwyller, seedlings, 2 years transplanted,

are 15 francs a hundred; the variegated-leaved variety, and a variety with fern-like leaves (*Æ. aspleniifolia*), 3 francs each: at New York, ?.

‡ 2. *Æ.* (H.) OHIOE'NSIS Michx. The Ohio *Æsculus*, or *Horsechestnut*.

Identification. Mich. Arb., 3. p. 242.; Dec. Prod., 1. p. 597.; Don's Mill., 1. p. 652.

Distinct. Char., &c. The fruit is said to be of about half the size of that of the common horsechestnut. Judging from the two trees in the garden of the London Horticultural Society, one of which flowered in 1835, we should say that this was nothing more than a variety of the common species, and far inferior to it in point of beauty.

‡ 3. *Æ.* (H.) RUBICU'NDA Lois. The reddish-flowered *Æsculus*, or *Horsechestnut*.

Identification. Loiseleur Herb. Amat.; Dec. Prod., 1. p. 597.; Don's Mill., 1. p. 652.; Marronnier rubicund, Fr.; scharlachrothe Rosskastanie, Ger.

Synonyms. *Æ. carnea Hort.*, and *Lindl. Bot. Reg.*; *Æ. rosea Hort.*; *Æ. coccinea Hort.*; *Æ. Hippocastanum* var. *rubicundum Schubert*; Whitley's fine scarlet.

Engravings. Herb. Amat., t. 367.; Hayne, Abbild., t. 22.; Bot. Reg., t. 1056., as *Æ. carnea*; Wats. Dendr., t. 121., as *Æ. carnea*; and the plate in our Second Volume.

Distinct. Char., &c. Petals 4, with the claws shorter than the calyx. The flowers are scarlet, and very ornamental; the leaves of a deeper green than those of any other sort. It is doubtful whether this tree be a native of North America, or originated in British gardens. It passes under different names in different nurseries, as will be seen by our list of synonymes, and may be considered as differing little, if at all, from *Æ. carnea Lindl.* It is distinguished from *Pavia rubra* by its larger and rougher leaves, and from *Æ. Hippocastanum* by the leaves being fuller and more uneven on the surface, and of a deeper green. The tree is also smaller, and of much less vigorous growth; but, as it has only been in cultivation since 1820, sufficient time has not elapsed to know its ultimate size. It is, without doubt, the most ornamental sort of the genus.

Statistics. In the environs of London, at Kenwood, 8 years planted, and 12 ft. high; in Devonshire, at Endsleigh Cottage, 18 years planted, and 30 ft. high; in Staffordshire, at Arley Hall, 27 ft. high, diameter of the trunk 10 in., and of the head 24 ft.; in Suffolk, at Ampton Hall, 12 years planted, and 12 ft. high; in Surrey, at Farnham Castle, 20 years planted, and 10 ft. high.

Commercial Statistics. Plants of this sort, in the London nurseries, cost from 1s. 6d. to 3s. 6d. each; at Bollwyller, 2 francs; at New York, ?.

‡ 4. *Æ.* (H.) GLA'BRA Willd. The smooth-leaved *Æsculus*, or *Horsechestnut*.

Identification. Willd. Enum., p. 405.; Hayne Dend., p. 44.; Dec. Prod., 1. p. 597.; Don's Mill., 1. p. 652.

Engravings. Hayne Abbild., t. 34.; and our fig. 133.



Distinct. Char., &c. Claws of the petals of about the length of the calyx. Leaflets of a pale green, very smooth. Flowers of a greenish yellow. A low tree, a native of North America, introduced in 1822, and flower-

ing, with the other sorts, in June. This sort is very distinct; but whether it is a species, or not, appears to us doubtful. The whole plant is comparatively glabrous, and even the fruit

partakes of that quality. The tree is of less vigorous growth than *Æ. rubicunda*; and the shoots take a more upright direction. It appears to lose its leaves sooner than most of the other sorts. There is a tree of it in the London Horticultural Society's Garden. Plants, in the London nurseries, cost 3s. 6d. each.

¶ 5. *Æ. (H.) PALLIDA* Willd. The pale-flowered *Æsculus*, or *Horsechestnut*.

Identification. Willd. Enum., p. 406; Hayne Dend., p. 44; Dec. Prod., 1. p. 597; Don's Mill., 1. p. 652.

Synonyme. Gelbliche Rosskastanie, *Ger.*

Engravings. Hayne Abbild., t. 25.; and our fig. 134.



Distinct. Char., &c. Petals with the claws shorter than the calyx. Stamens twice as long as the corolla. Flowers greenish yellow or whitish. A native of North America, in the forests of Kentucky, introduced in 1812. This sort so closely resembles *Æ. glabra* as to leave no doubt in our mind of its being essentially the same. It is of somewhat more robust growth, and the leaves are, perhaps, not quite so smooth. The tree in the Horticultural Society's Garden, in 1834, measured 12 ft. in height, after having been 8 years planted; the diameter of the trunk was 3½ in., and of the head 7 ft.

¶ 6. *Æ. (H.) LYONII* Hort. Lyon's *Æsculus*.

Plants of this species, or variety, are in the garden of the London Horticultural Society; but they are so small, that it is difficult to say what they will ultimately prove to be.

App. i. Other Sorts or Varieties of *Æsculus*.

In consequence of this genus ripening its seeds freely, and admitting of cross-fecundation with the genus *Pavia*, several varieties have, within these few years, been raised by British cultivators; and, indeed, there seems no limit to the number which may be raised by these means. In the Fulham Nursery are, *Whitley's fine scarlet*, which seems little, if at all, different from *Æ. rubicunda*; *Æ. americana*, which also differs little from *Æ. rubicunda*; and several other varieties, which will be noticed in our appendix to the genus *Pavia*. (See *Gard. Mag.*, vol. xi. p. 218.) In the garden of the London Horticultural Society are the following names:—*Æ. H. incisum*, *Æ. H. præcox*, *Æ. H. tortuosum*, and *Æ. H. nigrum*; but the plants to which they are applied are all quite small and young. It may be worth while to remark, that purchasers of the different varieties should always take care

to have worked plants; and indeed they should, if practicable, endeavour to see the tree from which the plants have been worked, when that tree is in flower. For this reason we think that nursery-men, who propagate varieties of these and other select flowering trees, ought always to keep specimen plants in their own grounds, from which to take scions for propagation. At all events, such varieties ought never to be raised from seed; because, though there can be no doubt but that the progeny would bear a general resemblance to the parent, yet the particular feature for which the variety was cultivated might be wanting. For example, the flowers of the seedlings may come earlier or later, larger or smaller, than those of the parent. Early and late varieties of all showy-flowered trees, and trees which retain their leaves late in the season, are also desirable; and in this respect the common horsechestnut varies exceedingly, as any one may observe, by walking along the avenue of horsechestnuts in Bushy Park in spring and autumn. There is one variety of the common horsechestnut, exemplified in a tree in the garden of the Tuilleries, which we should like much to see introduced into Britain. This tree is easily distinguished, even in summer, from all the others in the same garden, by the profusion of flowers with which it is covered, and also by the earliness of their appearance, and that of the leaves. The tree was first mentioned to us by our correspondent, Mr. Blaikie, some years since, as flowering always a fortnight earlier than the others; and another friend has supplied us with a fact respecting the individual tree at the Tuilleries, which is of some historical interest. On Napoleon's entry into Paris, on the 20th March, 1815, after his return from Elba, this tree furnished to him and his friends foliage for their personal decoration, being the only tree then in leaf in the garden of the Tuilleries. (See *Gard. Mag.*, vol. xi. p. 557.)

GENUS II.



PAVIA Boerh. THE PAVIA, or SMOOTH-FRUITED HORSECHESTNUT TREE.
Lin. Syst. Heptándria Monogýnia.

Identification. Boerh. Lugd., t. 260.; Dec. Prod., l. p. 598.; Don's Mill., l. p. 652.

Synonymæ. Pavier, Fr.

Derivation. In honour of Peter Paw, a Dutch botanist, once Professor of Botany at Leyden.

Gen. Char. Middle-sized deciduous trees or shrubs, distinguishable from the horsechestnuts by the smoothness of their fruit, and the comparative smallness of their flowers, which have their petals erect and narrower. The leaves, also, are generally smaller, and smoother. There are probably only three, or possibly only two, aboriginal species.

‡ 1. P. RU'BRA Lam. The red-flowered Pavia.

Identification. Lam. Illust.; Dec. Prod., l. p. 598.; Don's Mill., l. p. 653.

Synonymes. Æsculus PAVIA Lin.; Æ. PAVIA var. *æ rubra* Hayne *Dend.*, p. 44.; PAVIA parviflora Hort.; small Buckeye, Amer.; Marronier Pavie, or Pavie à Fleurs rouges, Fr.; rothe Rosskastanie, Ger.


Engravings. Lam. Illust., t. 273.; Hayne Abbild., t. 21.; Wats. *Dend.*, t. 120.; Krause, t. 55.; and our plate of the tree in Vol. II.

Spec. Char., &c. Fruit smooth. Corolla of 4 petals, that are longer than the stamens. Leaflets 5, elliptic-oblong, tapered to both ends, and smooth, as is the petiole; axils of the nerves hairy on the under surface of the leaf. (*Dec. Prod.*, l. p. 598.) A slender-growing tree, from the mountains of Virginia and Carolina; and said, also, to be a native of Brazil and Japan. Introduced in 1711, and producing flowers, which are of a brownish scarlet colour, in May and June. Height, from 10 ft. to 20 ft.; or, in some cases, to 30 ft. The tree in the garden of the London Horticultural Society was, in 1834, 10 ft. high, after having been planted 8 years.

Description, &c. In its native country, the P. rubra varies in magnitude from a low rambling shrub to a tree of 20 ft. or more in height. In England, it is in cultivation in various forms: as a tree, in which character it has, at Syon (see our plate in Vol. II.), attained the height of 26 ft.; as a pendulous tree, of 12 ft. or 14 ft. in height; and as a trailing shrub, under the name of P. humilis in the London Horticultural Society's Garden, and in the arboretum of Messrs. Loddiges. In addition to these forms, there are several others which are enumerated below.

Varieties.

‡ P. r. 2 arguta G. Don. The sharp-toothed-leaved red-flowered Pavia.— Figured in the *Botanical Register*, t. 993., and in our fig. 135. Introduced into the garden of the London Horticultural Society from the nursery of M. Catros of Bordeaux, under the name of Æsculus PAVIA parviflora. It is a handsome small tree, with dark brownish red flowers,



135

differing little from those of *P. rubra*. The tree in the garden of the London Horticultural Society, which, in 1834, was marked *Æsculus Pavia parviflora*, was then 15 ft. high, after having been 10 years planted.

- † *P. r. 3 sublaciniata* Wats. *The slightly cut-leaved red-flowered Pavia*.—Figured in *Wats. Dend.*, t. 120. Leaflets acutely serrated: in other respects it differs little from the species. In 1823, plants of it were in the Fulham Nursery, whence it was figured by Watson. The plants in the same nursery named *Æsculus Pavia serrata* (see *Gard. Mag.*, vol. xi. p. 248.) appear to be the same sort.
- ‡ *P. r. 4 humilis*. *P. humilis* G. Don. in *H. B.*, and in his *Mill.*; and *Æsculus humilis* Lodd. *The dwarf red-flowered Pavia*.—Figured in the *Botanical Register*, t. 1018. A diminutive, weak, straggling form of the species, probably obtained from some sport, and which, on its own root, is only a recumbent bush, from 2 ft. to 3 ft. in height; but which, when grafted on the common horsechestnut, forms the very beautiful pendulous low tree noticed below. A plant of *P. humilis*, in the garden of the London Horticultural Society, was, in 1834, 3 ft. high, after having been planted 7 years.
- § *P. r. 5 humilis pendula*. *The pendulous-branched dwarf red-flowered Pavia*.—Figured in our Second Volume. This is not properly a variety, but only a variation in form, produced by changing the position of the plant by grafting. There is a very handsome low tree of it in the arboretum at Messrs. Loddiges's, which continues flowering and fruiting almost the whole summer. We consider this one of the most beautiful and interesting forms of *Pavia*, and would recommend horsechestnut trees of 20 or 30 years' growth to be grafted all over with it at the points of the shoots, care being taken afterwards, once or twice in every year, to rub off all the buds from the stock as

soon as they appear, so that the entire force of the plant may be directed to the nourishment of the scions. Plants of *P. r. h. pendula*, in the London nurseries, are 5*s.* each: but, as they are rather scarce, the readiest way of obtaining pendulous trees is, to procure plants of *P. r. humilis*, which can be had for 2*s.* each, and common horse-chestnut trees 12 ft. high, which can be had for 2*s.* 6*d.* each; and to graft the shoots of the former on the tips of those of the latter.

Statistics. In the environs of London, at Syon, there is a tree of *P. rubra* 26 ft. high (as exhibited in the plate in our Second Volume); at Ham House, Essex, 21 ft. high, the diameter of the head, 32 ft.; in Hampshire, at Southampton, 12 years planted, and 20 ft. high; in Surrey, at Bagshot Park, 12 years planted, and 14 ft. high; in Wiltshire, at Wardour Castle, 20 years planted, and 30 ft. high; in Suffolk, at Finborough Hall, 6 years planted, and 10 ft. high. In France, at Paris, in the Jardin des Plantes, 23 years planted, and 30 ft. high; in the Botanic Garden at Toulon, 40 years planted, and 12 ft. high. In Hanover, at Schwöbber, 40 ft. high. In Saxony, at Wörlitz, 25 years planted, and 30 ft. high.

Commercial Statistics. Plants, in the London nurseries, are 1*s.* 6*d.* to 2*s.* 6*d.*; at Bollwyler, 80 cents; at New York, 25 cents, and nuts 40 cents a quart.

‡ 2. *P. FLAVA* Dec. The yellow-flowered Pavia.

Identification. Dec. Prod., i. p. 598.; Don's Mill, i. p. 653.

Synonymes. *Æsculus flava* *Ait.*, *Hayne*; *Æ. lutea* *Wangh.*; *Pavia lutea* *Poir.*; the large Buckeye, big Buckeye, *Amer.*; the yellow Pavia.

Engravings. Wangh. in Act. Nat. Scrut. Berl., 8. t. 6.; Hayne Abbild., t. 23.; Krause, t. 44.; Wats. Dend. Brit., t. 163.; and our plate in Vol. II.

Distinct. Char., &c. Petioles pubescent, flattish towards the tip. Leaflets 5—7, pubescent beneath, and above upon the nerves. (*Dec. Prod.*, i. p. 598.) This species differs from *P. rubra*, in being a much stronger plant, assuming the character of a tree of the second rate, and attaining the height of 30 ft. or 40 ft., or upwards, in England. The leaves are pubescent, and much paler than those of *P. rubra*, and the flowers are yellow. The tree seems to partake both of the character of *Æsculus* and *Pavia*. It is a native of Carolina and Virginia, in mountainous woods, and was introduced into England in 1764. In its native country, on the declivities of mountains, where the soil is loose, deep, and fertile, this tree attains the height of 60 ft. or 70 ft., with a trunk 3 ft. or 4 ft. in diameter. The largest tree in England is at Syon, and is 40 ft. high, as exhibited in the plate in our Second Volume. The tree in the garden of the London Horticultural Society was, in 1834, 12 ft. high, having been planted 8 years. This species is not quite so free a flowerer as *P. rubra*, and it is one of the first of the genus to drop its leaves in autumn: they generally commence falling about the middle of August, and the tree is frequently naked by the 1st of September. Like all the *Æsculacææ*, to thrive, it requires a deep rich soil. It is commonly propagated by buds, because the colour of the flower is found to vary much in plants raised from seed. A nurseryman, writing on this subject in the *Gardener's Magazine* (vol. xi. p. 249.), observes that there are two varieties of *P. flava* in cultivation in the English nurseries: one, an inferiorly flowering variety, generally raised from seed; and the other variety producing a larger flower, and of brighter colour, only to be propagated by budding or grafting. There are two fine grafted trees of this variety in the Fulham Nursery, with trunks 5 ft. and 6 ft. in circumference, and nearly 30 ft. in height.

Statistics. In the environs of London, at Syon, there is a tree 40 ft. high, already mentioned; in the Fulham Nursery, two, nearly 30 ft. high; at Kenwood, one, 40 years planted, which is 37 ft. high; at Ham House, Essex, one, planted by Dr. Fothergill, which is 28 ft. high; in the arboretum at Kew, the tree is 30 ft. high; at York House, Twickenham, 40 years planted, and 30 ft. high. In the Isle of Jersey, in Saunders's Nursery, 10 years planted, and 12 ft. high. In Lancashire, at Latham House, 14 years planted, and 25 ft. high; in Monmouthshire, at Dowlais House, 20 years planted, and 15 ft. high; in Oxfordshire, in the Oxford Botanic Garden, 40 years planted, and 34 ft. high. In Scotland, in Perthshire, in the Perth Nursery, 20 years planted, and 15 ft. high; in Stirlingshire, in Calendar Park, 10 years planted, and 14 ft. high. In Ireland, in the environs of Dublin, at Castle-town, 35 ft. high; in the Glasnevin Botanic Garden, 30 years planted, and 30 ft. high; in Louth, at Oriel Temple, 40 years planted, and 31 ft. high. In France, at Paris, in the Jardin des Plantes, 55 years planted, and 44 ft. high; in the Botanic Garden at Toulon, 40 years planted, and 18 ft. high. In Hanover, at Schwöbber, 40 ft. high. In Saxony, at Wörlitz, 20 ft. high. In Austria, at Kopenzel, 12 years planted, and 18 ft. high; at Brück on the Leytha, 24 ft. high.

Commercial Statistics. Plants, in London, cost from 1s. 6d. to 2s. 6d. each; at Bollwyller, 80 cents; and at New York, 25 cents, and nuts 40 cents a quart.

§ 3. *P. DISCOLOR* Swt. in *H. B.* The two-coloured-flowered Pavia.

Identification. Swt. Hort. Brit., p. 83.; Don's Mill, 1. p. 653.

Synonymy. *Æsculus discolor Ph. and Bot. Reg.*

Engraving. Bot. Reg., t. 310.

Distinct. Char., &c. The whole plant, including the young wood, is covered with pubescence. The flowers are large, showy, continuing a long time expanding, and numerous, though they are but sparingly succeeded by fruit. When the plant is raised from seed, it is remarkable for its thick, fleshy, carrot-like roots, which, in free soil, penetrate perpendicularly to the depth of 8 ft. or 10 ft., as has been found to be the case in the Hammersmith Nursery. This sort was introduced from North America (where it is found principally in the western territory of Georgia), in 1812, by Mr. Lyon. Unless when grafted on *Æ. Hippocástanum*, it is seldom seen above 4 ft. or 5 ft. in height; but it is a very free flowerer, and, considered as a shrub, one of the most ornamental in May that can be planted. The plant in the garden of the London Hort. Soc. was, in 1834, 4 ft. high, after having been 3 years planted. There are varieties of it in the nurseries under different names; one of these, raised by Messrs. Rivers at Sawbridgeworth, bears the name of *P. cárnea pubescens*. There is a handsome specimen of this bush in the Hammersmith Nursery; it is about 5 ft. high, and 6 ft. or 8 ft. in diameter, and is profusely covered with bloom every year. A large tree of the common horsechestnut, covered with grafts of this species, as recommended in the case of *P. hùmilis*, would form a noble object. Scions taken from flowering trees, and grafted on *P. hùmilis*, would form beautiful miniature trees for pots. Scions from flowering trees, grafted on stocks of this species, flower the second year, and form the most beautiful flowering shrubs for small gardens that can well be recommended. The same may truly be said of *P. hùmilis*, and *P. macrostàchya*, and yet none of these species are ever to be met with in the suburban gardens of the metropolis. Plants, in the London nurseries, cost 2s. 6d. each.

‡ 4. *P. HYBRIDA* Dec. The hybrid Pavia.

Identification. Dec. Prod., 1. p. 598.; Don's Mill, 1. p. 653.

Synonymy. *Æsculus hybrida Dec. Hort. Monsp.*, 1813, p. 75.

Distinct. Char., &c. Leaves clothed beneath with velvety pubescence, petioles smooth; flowers variegated with yellow, white, and purple. The tree in the London Horticultural Society's Garden was, in 1834, 6 ft. high, after having been 5 years planted. The leaves and flowers bear some resemblance to those of *P. discolor*; but the flowers of *P. hybrida* are more sparingly produced. This sort is not in general cultivation; though, like every other kind of *Pavia* and *Æsculus*, it well deserves to be so.


‡ 5. *P. NEGLECTA* G. Don. The neglected Pavia.

Identification. Loud. Hort. Brit., p. 143.; Don's Mill, p. 653.; Swt. Hort. Brit., p. 83.

Synonymy. *Æsculus neglecta Lindl. in Bot. Reg.*

Engravings. Bot. Reg., t. 1009.; and our fig. 136.

Distinct. Char., &c. Leaves with rufous down on the veins on the upper side, smooth beneath; rather plicate. Flowers pale yellow, veined with red. This is a tree resembling the preceding sort, and, like it, is apparently a hybrid between *P. rubra* and *P. discolor*. It was purchased by the London Horticultural Society from M. Catros of Bordeaux, under the name of *Æ. ohioënsis*. In the *Botanical Register*, it is said to be most nearly related to *Æ. (Pavia) flava*, but to differ from it in the flowers appearing 10 days earlier, and in the leaflets being more glabrous, with rufous down on the veins on the upper side, and with hairs in the axils of the veins on the under surface. There is a tree of this sort in the garden of the London Horticultural Society, which, in 1834, was 12 ft. high, after having been 8 years planted; but it has scarcely yet found its way into the nurseries.



‡ 6. *P. MACROCARPA Hort.* The long-fruited Pavia.

Synonymes. *Æsculus Pavia macrocarpa Lodd. Cat.*, 1830; *Pavia macrocarpa* in the Hort. Soc. Gard.

Engraving. Our plate in Vol. II.

Distinct. Char. &c. This tree is upwards of 20 ft. high; and that in the garden of the London Horticultural Society, was, in 1834, 12 ft. high, after having been 8 years planted. This sort appears to us to be intermediate between some variety of *Æsculus Hippocastanum* and *Pavia rubra*. The leaves are large, smooth on the upper surface, and shining. The flowers are nearly as large as those of the common horsechestnut, but with the petals less spreading, and of a pale red colour mixed with yellow. The branches are spreading and loose; and the whole tree has an open graceful appearance, quite different from that compactness of form and rigidity of branches which belong to most of the tree species and varieties both of *Æsculus* and *Pavia*. This sort can scarcely be said to be in cultivation in the nurseries, notwithstanding its claims to a place in every collection of ornamental trees.

‡ 7. *P. MACROSTACHYA Lois.* The long-racemed Pavia.


Identification. Lois. Herb. Amat.; Dec. Prod., 1. p. 598.; Don's Mill., 1. p. 652.

Synonymes. *Æsculus parviflora Walt.*; *Æ. macrostachya Mx. and Hayne*; *Pavia alba Poir.*; *Pavia edulis Poit. Arb. Fr.*, t. 88.; *Pavier à longs Epis, Pavier nain, Fr.*; *langährige Rosskastanie, Ger.*

Engravings. Lois. Herb. Amat., t. 212.; Jacq. Ecl., t. 9.; Poit. Arb. Fr., t. 88.; Hayne Abbild., t. 26.; Colla Hort. Rip., t. 19.; and our fig. 137.

Spec. Char., &c. Stamens much longer than the corolla; racemes very long. Root stoloniferous. Flowers white. (*Dec. Prod.*, i. p. 598.) A shrub, with loose racemes of white flowers, with long projecting stamens, which give the spike a fine fringed appearance. A native of North America, on the banks of rivers, more particularly in Georgia, near the little town of St. Augustin;

introduced in 1820; flowering in June, July, and August. The shoots are slender, spreading, and rooting at the joints where they happen to rest on the soil, with ascendent extremities. The tree comes into flower about a month or six weeks later than the other *Æsculaceæ*, and continues flowering, in the case of large plants on moist soil, for three months or longer, forming one of the greatest floral ornaments of the shrubbery, at a season when very few trees or shrubs are in flower. The fruit, which is small, seldom ripens in England; but in America it is said to be eaten, boiled or roasted: and M. Poiteau, accordingly, has included this species of *Pavia* in his list of fruit trees. (See *Bon Jard.*, 1835, p. 775.) When plants are to be raised from the nuts, he says they ought to be sown immediately; as, if kept exposed to the air, they shrink, and soon lose their vegetative power. The flowers are agreeably fragrant, and, as before observed, very ornamental; as are the spreading leaves, supported on long slender petioles; which, from their graceful disposition, combined with the feathery lightness of the racemes of flowers, give the whole plant an air of elegance quite different from that of any other species of dwarf pavia.



137

Statistics. The largest plant in the environs of London is at Syon; but it is not more than 12 ft. high. The plant of this species in the garden of the London Horticultural Society, 7 years planted, was, in 1834, 5 ft. high. In Berkshire, at White Knights, there are a great many plants which flower profusely the whole season, and among them is one, 25 years planted, which is 15 ft. high; in Lancashire, at Latham House, one, 12 years planted, 10 ft. high; in Suffolk, in the Bury Botanic Garden, 7 years planted, and 6 ft. high; and, in Surrey, at Farnham Castle, several plants, 10 ft. high.

Commercial Statistics. Plants, in London, are 1s. 6d. each; at Bollwyller, 2 francs; and at New York, 25 cents, and nuts 50 cents per quart.

App. i. Other Varieties of *Pavia*.

In the Fulham Nursery are plants belonging to *Pavia*, or intermediate between *Pavia* and *Æsculus*, with the names, *Pavia serrata* and *P. erecta*; and in the garden of the London Horticultural Society are plants marked *Æsculus Pavia flava* var. In different nurseries, there are different names for the same variety; and, as almost all the sorts seed freely, and hybridise as freely, both with *Æsculus*, and with each other, new varieties may be expected in abundance. All the species and varieties are so truly beautiful, that this is not to be regretted, more especially if they are kept distinct, and so described and named as to indicate what they are, and to enable purchasers to be certain of obtaining them. It is almost unnecessary to observe, that all the most valuable varieties are best perpetuated by budding or grafting, and that, with regard to the pavia, as well as to the *æsculus* (see p. 469), collectors ought always to see that the plants they purchase have been worked.

CHAP. XXIV.



OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE ORDER SAPINDACEÆ.

Of this order there is only one hardy ligneous plant in the country, namely, *Kolrenteria paniculata* Laxm.; and the half-hardy species, which chiefly belong to the genus *Dodonæa*, are not much cultivated even in green-houses.

GENUS I.



KÖLREUTE'RIA Laxm. THE KOLREUTERIA. *Lin. Syst. Octándria*
Monogýnia.

Identification. Laxm. Acad. Petr. 16. p. 561.; L'Hérit. Sert., 18. t. 19.; Willd. Spec. Pl. 330.; Dec. Prod., 1. p. 616.; Don's Mill., 1. p. 672.

Synonyme. Sapindus sp. *Lin. Fil.*

Derivation. In honour of *John Theophilus Kölreuter*, once Professor of Natural History at Carlsruhe, and celebrated for his researches on the pollen of plants.

Gen. Char., &c. *Calyx* of 5 sepals. *Petals* 4, each with 2 scales at the base. *Capsule* 3-celled, inflated. *Seeds* ovate-globose, the seed-coat penetrating into the seed, and occupying in the place of an axis the centre of the embryo, which is spirally convoluted. *Leaves* impari-pinnate, of many pairs of leaflets that are ovate, and coarsely toothed. *Flowers*, yellow, in panicles. (*Dec., Prod., i. p. 616.*)—A deciduous tree of the middle size.

‡ 1. K. PANICULA'TA Laxm. The paniced-*flowering* Kolreuteria.

Identification. N. D. Ham., t. 36.; Dec. Prod., 1. p. 616.; Hayne Dend. p. 45.; Don's Mill., 1. p. 672.

Synonymes. Sapindus chinénsis *Lin. Fil. Supp.*, p. 221.; K. paullinióides L'Hérit. Sert.; Savonnier paniculé, *Fr.*; rispentragede Kölreuteric, *Ger.*

Engravings. L'Hérit. Sert., 18. t. 19.; N. Du Ham., 1. t. 36.; Bot. Reg., t. 320.; and the plate of the tree in our Second Volume.

Description, History, &c. A tree of the middle size, with a loose irregular head, polygamous; that is, sometimes hermaphrodite, and sometimes unisexual: a native of China, and introduced in 1763. It was first cultivated at Croome, in Worcestershire, by the Earl of Coventry; and, being highly ornamental, both from its large compound leaves and fine loose terminal spikes of yellow flowers, it is to be found in most collections. Considering that it is a native of China, it is very hardy; the hermaphrodite plants not unfrequently ripening seeds in the neighbourhood of London. It has not only a very fine appearance when in flower, but also in autumn, when the tree is covered with its large bladdery capsules, and the leaves change to a deep yellow, which they do before they fall off. It was introduced into France in 1789, and is perfectly hardy in the neighbourhood of Paris, and also in the south of Germany. It is of the easiest culture in any common soil, and is readily propagated either by seeds or cuttings of the root or branches. In the London nurseries, it is generally propagated by seed. Though there are trees of this species of considerable size, both in Britain and on the Continent, we have never heard anything of the quality of its wood; which, from the prevalence of a yellow colour in its foliage and flowers, may probably be of a fine colour, and yield a yellow dye. The tree ought to be in every collection, on account of the beauty of its leaves, flowers, and fruit. In a young state, it is sometimes seen with a ragged head, owing to the young shoots dying back, after wet summers and cold autumns; but, as it gets older, it makes shorter shoots, and these have more time to ripen. Accordingly, old trees have generally much handsomer heads than young ones. The general contour of these heads is hemispherical, as may be seen by the fine old specimens at Kew, in the Fulham Nursery, and in the Jardin des Plantes at Paris.

Statistics. In the environs of London, the largest tree is at Ham House, where it is 42 ft. high, and the diameter of the trunk, at 1 ft. from the ground, is 16 in.; at Kew, it is 30 ft. high; in the Fulham Nursery, 25 ft.; at Fulham Palace, 17 years planted, it is 20 ft. high; at Syon, 20 ft. high; in Berkshire, at White Knights, 25 years planted, and 23 ft. high; in Hertfordshire, at Cheshunt, 6 years planted, and 11 ft. high; in Staffordshire, at Alton Towers, 10 years planted, and 10 ft. high; in Yorkshire, in the Hull Botanic Garden, 12 years planted, and 8 ft. high. In Scotland, in Lawson's Nursery, Edinburgh, 4 years planted, and 4 ft. high; in Sutherlandshire, at Dunrobin Castle, 20 ft. high. In Ireland, in the environs of Dublin, at Castletown, 15 ft. high; in the Glasnevin Botanic Garden, 20 years planted, and 12 ft. high; at Terenure, 10 years planted, and 6 ft. high; at Oriel Temple, 25 years planted, and 30 ft. high. In France, in the Jardin des Plantes, 30 years planted, and 25 ft.

high; at Se'caux, 10 years planted, and 20 ft. high; in the Toulon Botanic Garden, 20 years planted, and 12 ft. high; at Nerrières, near Nantes, 20 years planted, and 15 ft. high. In Austria, in the University Botanic Garden at Vienna, 25 years planted, and 25 ft. high.

Commercial Statistics. Plants, in London, cost from 1s. to 2s. 6d. each; at Bollwyller, 1 franc each; at New York, ?.

App. I. *Half-hardy ligneous Species of Sapindæcæ.*

Dopoxæa, a genus of plants named in honour of *Rambrot Dodoens*, author of *Historia Plantarum*, who died in 1585, consists of nearly 30 species of green-house plants, which are chiefly natives of new Holland, though some of them are from the East and West Indies and South America. They are all shrubs, with exstipulate, simple, or pinnate leaves, and small greenish yellow flowers. They are not showy, but they are interesting to the botanist, as illustrating this order, and also on account of the ramified venation of their leaves.

♣ *D. viscosa* L. is a native of the Caribbee Islands, where it is a shrub growing to the height of 6 ft. It has been in the country since 1690, and is occasionally to be met with in green-houses. It is highly probable that it would stand our winters against a wall, with sufficient protection.

♣ *D. alternata* Cuming, is a native of New Holland, and has been in cultivation since 1824.

♣ *D. jamaicensis* Dec., *D. angustifolia* Siez., *D. viscosa* Cav., is a native of the colder parts of Jamaica, where it grows to the height of 6 ft.; and, being very sour and bitterish in all its parts, it is known there by the name of switch sorrel. It has been in our green-houses since 1810.

♣ *D. salicifolia* Dec., *D. angustifolia* Lam., is in cultivation in French gardens under the name of *bois de reinette*, and has been in our green-houses since 1820. The leaves are very narrow, and they are sweet-scented. It is supposed to be a native of New Holland.

♣ *D. laurina* Sieb., *D. trilobata* Bot. Rep. t. 231., *D. cuneata* Smith, and *D. asplenifolia* Rudge, are all natives of New Holland, occasionally to be met with in green-houses. They are generally cultivated in loam and peat, or in any light soil; and, when they are tried against a conservative wall, care should be taken that they are not overpowered at the root, or at the top, by other plants.

CHAP. XXV.

OF THE HALF-HARDY LIGNEOUS PLANTS OF THE ORDER MELIACEÆ.

♣ 1. *Melia Azedarach* L., the bead tree, or Indian lilac, (fig. 138.) is an old inhabitant of British green-houses, and well known to all those who have travelled in Italy. The word *Melia* is derived from *melita*, the Greek name for the manna ash (from *melis*, honey); from a fancied resemblance between the leaves and those of the ash; and *Azedarach* from an Arabic word signifying a poisonous plant; the berries of the melia being formerly supposed to be poisonous. Its foliage and its spikes of flowers are large, the plant being remarkably showy when fully developed; and it is by no means tender. It grows in its native country, Syria, to the height of 40 ft.; and there are trees of nearly that height in the neighbourhood of Naples. It is planted as an ornamental tree in Spain, Portugal, Italy, and the south of France. There are trees of it in the public walks at Montpellier, at Toulon, and in various cities in Italy. In the southern states of North America, more particularly in Carolina, it is planted near houses, and known there by the name of the pride of India; a name also given to the *Lagerstræmia indica*. In Greece, and along the shores of the Grecian Archipelago and the Mediterranean, the *Melia Azedarach* is always planted in the area of monasteries for the sake of the nuts, contained in its fruits, which are made into rosaries by the monks; and hence its name of the bead tree. The fruit, which is of the size of a cherry, but more cylindrical, and of a pale yellow colour when ripe, was said by the Arabian physician Avicenna to be poisonous; and the pulp was mixed with grease, for the purpose of killing rats and dogs. According to Royle, however, the fruit can only be considered poisonous when used in large doses. It is used in Java as a vermifuge. The nuts, which are of a brown colour, are bored, and, as already stated, strung as beads in Catholic countries. In Britain, the tree frequently flowers in green-houses, and sometimes ripens seeds; it has been tried in the open air, both as a standard and against a wall. It has stood through several winters, in the open air, at Biel, in East Lothian; and at Bungay, in Suffolk, a plant, which had been 9 years planted against a wall, was, in 1831, $9\frac{1}{2}$ ft. high, the trunk 9 in. in diameter, and had branches extending 18 ft. on each side of the trunk. One, raised from seed in 1828, which has stood ever since against a wall in our garden at Bayswater, protected by a glass case during winter, flowered in 1855. In the warmest parts of Devonshire and Cornwall, it might be treated as a standard tree. Plants are generally raised from seeds; and they may be procured in the London nurseries at 2s. each; at Bollwyller, for 1 franc and 50 cents; and at New York, for 25 cents a plant, and 1 dollar a quart of seeds.

♣ 2. *M. sempervirens* Swz., the evergreen *Melia*, or Bead Tree, known in the West Indies by the name of the Indian lilac, is said to be a tree growing to the height of about 25 ft. It has been in our green-houses since 1636; and is by some considered as only a variety of *M. Azedarach*.



¶ 3. *M. australis* Swt. is a native of New Holland, introduced in 1810, and said to grow, in its native country, to the height of 20 ft.

¶ 4. *M. japonica* G. Don is a green-house species, growing 30 ft. high, which has not yet been introduced; and *M. Buckayun Royle* is a species of which little seems to be known.

All the species of *Melia*, being deciduous trees, without visible buds, are peculiarly eligible for growing against a conservative wall; because, by the application of heat artificially, and by preventing (which can be done by thatching the ground) the rain from falling on the soil under the trees at the end of summer, the wood may be ripened to such a degree as to enable it to stand our winters with very little or no protection.

CHAP. XXVI.

OF THE HARDY LIGNEOUS PLANTS OF THE ORDER VITACEÆ.

THALAMIFLOROUS. (*H. B.*) Calyx small. Petals 4 or 5, inserted on the outside of a disk surrounding the ovarium; in æstivation, turned inwards at the edge in a valvate manner. Stamens equal in number to the petals, inserted upon the disk that surrounds the ovarium; filaments distinct or slightly cohering at the base. Anthers versatile. Ovarium 2-celled. Fruit a pulpy berry. Seeds 4 or 5, fewer by abortion; embryo erect; albumen hard. Climbing shrubs, with tumid separable joints. Leaves with stipules. Properties, acidity and sugar. (*Lindl. Introd. to N. S., and Key.*) The species are trailing and climbing shrubs, and they include the grape vine, which may be considered as the type of the order. "The genus *Vitis* is found in the equinoctial parts of the Old and New Worlds, extending into both the temperate zones; as, southwards, to the Cape of Good Hope and New Holland; and, northwards, to Japan and North America, as well as from the plains of India to the defiles of Caucasus." (*Royle, Illustr.*, p. 144.) The genera which contain hardy species are two, which are thus distinguished:—

VITIS. Calyx 5-toothed. Style wanting. Berry, 2-celled, 4-seeded.

AMPELOPSIS. Calyx nearly entire. Petals 5. Stamens 5. Style 1, crowned by a capitate stigma.

CYSSUS. Calyx nearly entire. Petals 4. Stamens 4. Ovary 4-celled.

GENUS I.



VITIS L. THE GRAPE VINE. *Lin. Syst.* Pentándria Monogýnia.

Identification. *Lin. Gen.*, 284; *Dec. Prod.*, 1. p. 633.; *Don's Mill.*, 1. p. 695.

Synonymes. *Giud. Celtic*; *Vid. Span*; *Vigne, Fr.*; *Wein, Ger.*

Gen. Char. *Flowers* hermaphrodite, diœcious or trœcious. *Calyx* commonly 5-toothed. *Petals* 5, cohering at the top, separating at the base, and deciduous. *Stamens* 5.—Climbing shrubs, deciduous, with leaves simple, lobed, or serrated, sometimes compound, and small greenish yellow flowers in thyrsoid racemes. (*Dec. Prod.*, i. p. 633.) The species are deciduous climbers, one of which has long been celebrated in the Old World as the grape vine; and all the others are natives of North America. The varieties of the first species have been described at length by Du Hamel in France, Don Roxas de Clemati in Spain, and Siecker in Germany; and the species and varieties of North America by Rafinesque.

§ 1. *V. VINÍFERA* L. The wine-bearing Vine.

Identification. *Lin. Spec.*, 293; *Dec. Prod.*, 1. p. 675.; *Don's Mill.*, 1. p. 695.

Synonymes. *Vigne, Fr.*; gemeiner Weinstock, *Ger.*

Engravings. *Dub. Arb. Fr.*, 2. t. 16.; *Jacq. Ic.*, 1. p. 53.; and our *fig.* 139.

Spec. Char., &c. Leaves lobed, toothed, sinuated, or serrated, naked or downy. (*Dec. Prodrom.*, i. p. 633.) A deciduous climber, in cultivation from the remotest period of history, in the warmest parts of the temperate zones of the Old World, and of which there are innumerable varieties.

139



Geography and History. The grape vine is generally considered to be a native of Persia; and Dr. Sickler, in the first volume of his *Geschichte der Obst-cultur*, has given an interesting account of its migration to Egypt, Greece, and Sicily. From Sicily, which is generally considered to be one of the oldest seats of civilisation in the western hemisphere, the vine is said to have found its way into Italy, Spain, and France. It is supposed to have been cultivated in the latter country in the time of Antoninus, and to have been introduced into Britain by the Romans, but during what reign is uncertain. There were vineyards, however, in England, according to the venerable Bede, in the year, A. D. 280. The vine has been for ages in a wild state, in the woods and hedges of Provence, Languedoc, and Guienne, in France, where it differs from the cultivated plant, in having smaller and more cottony leaves, and very small fruit, rather austere than sweet. These wild vines, which were called by the ancients *labrusca*, are still known, in the south of France, by the names of *lambrusco*, and *lambresquiero*. (*N. Du Ham.*) The history of the vine as a fruit shrub, and all that relates to its varieties and their propagation and culture, will be found given at length in our *Encyclopædia of Gardening*; and we shall here only notice those varieties which we think deserving of introduction, as ornamental and fragrant-flowered standard climbers, for training against a prop in the free ground, in a British arboretum; or to be trained against a wall, in the arboretums of colder countries. Plants, in the European nurseries, are procurable at 1s. or 1 franc each; and at New York, for 37½ cents each.

♣ *V. v. 2 foliis incanuis.* *The hoary-leaved Grape Vine.* Miller's Grape, or Miller's black Cluster Grape.—Leaves almost entire, small, woolly, and whitish. Fruit round, small, in compact bunches, black. This variety is selected on account of the whiteness of its leaves.

♣ *V. v. 3 foliis rubescentibus.* *The rubescent-leaved Grape Vine.* The Claret Grape; Tenturier, Fr. (*N. Du Ham.*, var. 75., not *Clairette Du Ham.*, var. 12.)—The leaves are larger than those of the preceding variety, and more lobed and notched: in the autumn, before they die off, they change to a deep claret colour, in which state they are highly ornamental.

♣ *V. v. 4 apiifolia* Hort. *The Parsley-leaved Grape Vine.* Crotal, Fr. (*fig. 140.*)—The leaves are beautifully lacinated, middle-sized, and the fruit black. This variety is by some considered as a species, and, as such, is known as *V. laciniosa* L. It forms a very handsome climbing shrub, which has been in cultivation for its fruit since 1648.



140

§ 2. *V. LABRUSCA* L. The wild Vine, or Fox Grape.

Identification. Lin. Spec., 293; Dec. Prod., l. p. 634; Don's Mill, l. p. 711.

Synonymes. *V. taurina* Walt.; filziger Wein., *Ger.*

Engravings. Plum. Icon., t. 259. fig. 1.; Jacq. Schœn., t. 426.; and our fig. 141.

Spec. Char., &c. Sexes diœcious or polygamous.

Leaves heart-shaped, rather 3-lobed, acutely toothed beneath, and the peduncles tomentose and rather rusty. (*Dec. Prod.*, i. p. 634.) A tendriled climber, growing to about the same height as the common vine, but with much larger leaves (which are scarcely lobed, and downy, especially when young), and berries which are large and black, with a rough acid flavour, but are, nevertheless, eatable in a wild state, and much improved by cultivation. The whole plant has a disagreeable foxy smell, whence the name. "The fruit is, according to Professor Bigelow, large, purple, and pleasantly tasted; while Torrey remarks that it has a strong disagreeable flavour in a wild state, but that, when cultivated, it is as pleasant as any of the varieties of *V. vinifera*." (*Hook. Flor. Bor. Amer.*, p. 115.) There are two varieties growing in the vineyards of North America; one with white berries, and the other with red ones. From both of the varieties, and from the species, an excellent wine is made; which, when kept for five or six years, resembles Moselle. In America, the varieties have been much improved by culture; and, according to Rafinesque (*Med. Fl.*, i. p. 121.), greatly increased in number by culture, with a view to the production of wine. In Britain, the plant can only be considered as ornamental; and, from the largeness of its foliage and fruit, it forms a very distinct species of *Vitis*. A plant of the red-fruited variety, in the garden of the London Horticultural Society, placed against a wall with a west aspect, ripens fruit every year, which we have tasted, and found by no means disagreeable. We have also had some bottles of the wine sent us from America, which was not inferior to the weaker sorts of Rhenish wines. Possibly this plant might deserve cultivation on the Continent, with a view to the mixing of the fruit with that of the varieties of the grape vine, in making wine; since austere varieties of apple and pear, mixed with sugary varieties, are found to make the best kinds of cider and perry. Plants of this species, in the London nurseries, are 1*s.* 6*d.* each; at Bollwyller, 1 franc; and at New York, the species and its varieties are 37½ cents each.




§ 3. *V. ÆSTIVALIS* Michx. The Summer Vine, or Grape Vine.

Identification. Michx. Fl. Bor. Amer., 2. p. 230; Dec. Prod., l. p. 634; Don's Mill, l. p. 711.

Synonymes. *V. Labrusca* Walt. *F. Car.*, 242; *V. vulpina* Willd. *Spec.*, p. 1181.; and *E. of Pl.*, No. 2860.

Engravings. Jac. Hort. Schœn., t. 425.; *E. of Pl.*, 2860.; and our fig. 142.

Spec. Char., &c. Sexes diœcious, or polygamous. Leaves broadly heart-shaped, with from 3 to 5 lobes; the under surface of the young ones invested with a cottony down; of the adult ones, smooth. Racemes fertile, oblong. Berries small (*Dec. Prod.*, l. p. 634.) A native of North America, and abounding there in woods and wastes, from Virginia to Carolina. The berries are small, of a dark blue colour, finely covered with bloom, not disagreeable to the taste, and made into a very tolerable wine by the inhabitants. It was introduced into England in 1656, but is not very common in collections.



§ 4. *V. SINUATA* G. Don. The scallop-leaved Vine, or Summer Grape Vine.

Identification. Don's Mill, l. p. 711.

Synonymes. *V. æstivalis* var. *sinuata* Ph. *Flor. Amer.*, Sept., l. p. 169; *Dec. Prod.*, l. 634. "Probably the *V. labruscoides* of Muhl. *Cat.*, 27." (*G. Don*)

Spec. Char., &c. Sexes diœcious or polygamous. Leaves sinuately palmate, coarsely toothed, with rhomboid recesses; young ones covered beneath with cobwebby rusty down, adult ones smooth. (*Don's Mill.*, i. p. 711.) Found in woods from Virginia to Carolina, along with the two preceding species, of the last of which, notwithstanding Mr. G. Don's opinion, we think this only a variety. The berries are dark blue, agreeable to eat, and a very good wine is made from them. Introduced into England in 1656, but not much cultivated.

♣ 5. *V. CORDIFOLIA* Michx. The heart-shaped-leaved Vine, or Chicken Grape.

Identification. Michx. Fl. Bor. Amer., 2. p. 231; Dec. Prod., 1. p. 654; Don's Mill. 1. p. 711.

Synonymes. *V. incisa* Jacq. Schœn., t. 427; *V. vulpina* Lin Spec., p. 293, *Watt. Flor. Car.*, 243.

Engravings. Jacq. Schœn., t. 247; E. of Pl., 286; and our fig. 143.

Spec. Char., &c. Sexes diœcious or polygamous. Leaves heart-shaped, acuminate, toothed in the mode of incisions, smooth on both surfaces. Racemes loosely many-flowered. Berries small, greenish, ripened late. (*Dec. Prod.*, i. p. 634, 635.) Found wild from Canada to Florida, on the edges of rivers and in woods, where it is called the winter grape, probably from the late ripening of the fruit; and chicken grape, perhaps from the very small size of the berries. Dr. Torrey considers this to be the true *V. vulpina* of Linnæus, on account of its glabrous leaves. (*Hook.*) Introduced in 1806. Plants of it (but whether male or female, we are uncertain) are in the arboretum of Messrs. Loddiges, price 1s. 6d. each.

143



♣ 6. *V. RIPAÏRIA* Michx. The river-side, or sweet-scented, Vine.

Identification. Michx. Flor. Bor. Amer., 2. p. 821; Dec. Prod., 1. p. 635; Don's Mill. 1. p. 711

Synonymes. *V. odoratissima* Donn Hort. Cant., and *Lodd. Cat.*; Vigne de Bataures, Amer.

Engravings. Bot. Mag., t. 2429; E. of P., 2862; and our fig. 144.

Spec. Char., &c. Leaves heart-shaped, shallowly 3-cleft, toothed in the mode of incisions and unequally. Footstalk, and the margin of the nerves, pubescent. (*Dec. Prod.*, i. p. 635.) A native of North America, from Pennsylvania to Carolina, on the gravelly shores of islands and banks of rivers. "Extending to the south end of Lake Winipeg, in lat. 52°." (*Richardson*, in *Hook.*) Dr. Hooker observes that some of his specimens of this plant have the leaves so slightly lobed, that he scarcely knows how they are to be distinguished from *V. vulpina* (our No. 3.). Female plants are very seldom found north of the Potowmac river, though the male extends very far beyond it. The flowers have an exquisitely sweet smell, somewhat resembling that of mignonette. The female plant is in the arboretum of Messrs. Loddiges, where its shoots extend to the length of 20 ft.

144



♣ 7. *V. ROTUNDFOLIA* Michx. The round-leaved Vine, or Bullet Grape.

Identification. Michx. Flor. Bor. Amer., 2. p. 231; Dec. Prod., 1. p. 635; Don's Mill., 1. p. 711.

Spec. Char., &c. Sexes diœcious or triœcious. Leaves between heart-shaped and kidney-shaped, toothed in rather an equal manner, shining on both surfaces. Racemes composed of several little heads of flowers. Berries of a deep blue colour. (*Dec. Prod.*, i. p. 635.) Found in North America, from Virginia to Florida, on river sides, and on islands. The berries are as large as those of the common muscadine grape, by which name it is sometimes called: they are agreeable to eat. Introduced in 1806, but not often met with in collections. There is a plant in the garden of the London Horticultural Society.

♣ 8. *V. CARIBEA* Dcc. The Caribbean Vine.

Identification. Dec. Prod., 1. p. 634; Don's Mill., 1. p. 711.

Synonymes. *V. indica* Swz. Obs., 95., *Potr. Diet.*, 8. p. 607.

Engraving. Sloane Hist., 2. p. 104. t. 210. fig. 4.


Spec. Char., &c. Sexes diœcious or triœcious. Leaves heart-shaped, acuminate, toothed with acute and rather projecting teeth; rather glabrous above,

beneath, and the peduncles, tomentose. (*Dec. Prod.*, i. p. 634.) Flowers small and white. Berries small, brownish green, watery and acid, but eatable. This plant produces a great quantity of clusters of small black grapes, of an austere taste; but they would, doubtless, make a good red wine. When it grows luxuriantly, as it generally does on the higher woody lands of Jamaica, it is so full of juice, that a piece of a shoot, about 3 ft. long, will yield near a pint of clear tasteless water, which has saved the lives of many persons who have wandered long in the woods without any other refreshment of a liquid sort. For this reason, the plant is called, in Jamaica, water withe. According to Sloane, the fruit is red or deep purple, the size of currants, and agreeably acid, as well as astringent. (*Don's Mill.*, i. p. 711.) Introduced in 1800, but seldom to be met with. Not in the London Horticultural Society's Garden, nor in the collection of Messrs. Loddiges.

App. i. Other hardy or half-hardy Species of *Vitis*.

Dr. Wallich has enumerated upwards of fifty species of *Vitis*, natives of India. Of these *V. Wittichii* Dec., a native of Nepal, was introduced in 1822, and treated as a stove climber; but it will probably be found half-hardy. *V. glabrata* Roth is a native of the East Indies, introduced in 1819. It resembles in foliage *V. vulpina*, and is considered half-hardy. *V. indica* L. (*fig. 145.*) was introduced in 1692, and is generally treated as a stove climber; but, being deciduous, if its wood could be ripened in sufficient time, it might stand our winters against a conservative wall. The same may be said of *V. cæ'sia* Hort. Trans., a native of Sierra Leone, introduced in 1822. There are numerous other Indian and some Japan species, which remain to be introduced. There are also some species natives of South America, described by Humboldt, of which very little is known; but four of them, which are described in *Don's Miller*, are considered to be hardy.

Of North American species and varieties no fewer than 130 have been described by Professor Rafinesque in his *Medical Flora*, already quoted, and in his *Monograph of American Vines*. (See *Gard. Mag.*, vol. viii. p. 248.)



GENUS II.



AMPELO'PSIS Michx. THE AMPELOPSIS. *Lin. Syst.* Pentándria Monogýnia.

Identification. Michx. Fl. Bor. Amer., 1. p. 159; *Dec. Prod.*, 1. p. 632; *Don's Mill.*, 1. p. 694.

Synonymes. *Vitis* sp., and *Cissus* sp.

Derivation. *Ampelos*, a vine, and *opsis*, resemblance; similarity in the habits of the species.

Gen. Char., &c. *Calyx* almost entire. *Petals* 5, falling off separately. *Stigma* capitate. *Ovary* not immersed in a disk, including 2—4 ovules. (*Kunth, Nov. Gen. Am.*, 5. p. 222, quoted in *Dec. Prod.*, i. p. 632.)—A genus intermediate between *Cissus* and *Vitis*. (*Dec. Prod.*, i. p. 632.) The species are found in North America, in the north of Africa, in China, and in the Himalaya. They are all climbing shrubs, mostly deciduous, of the easiest propagation and culture: some of them, as the *A. hederacea*, are very ornamental.

§ 1. A. CORDATA Michx. The cordate-leaved Ampelopsis.

Identification. Michx. Bor. Amer., 1. p. 159; *Dec. Prod.*, 1. p. 633; *Don's Mill.*, 1. p. 694.

Synonymes. *Cissus Ampelopsis Pers. Syn.*, 1. p. 142; *Vitis indivisa Willd. Baum.*, 538.

Spec. Char., &c. Leaves heart-shaped, acute, toothed, indistinctly 3-lobed; the nerves villous beneath. Racemes doubly bifid. (*Dec. Prod.*, i. p. 633.) Found in North America, from Pennsylvania to Carolina, among hedges,

and by the sides of rivers. The flowers are reddish, and produced in May and June; and the berries are of a pale red colour. Introduced in 1803, but rare in British collections.

‡ 2. *A. HEDERA'CEA Michx.* The Ivy-like Ampelopsis, or Five-leaved Ivy.

Identification. Michx. Fl. Bor. Amer., 1. p. 160.; Dec. Prod., 1. p. 633.; Don's Mill., 1. p. 694.

Synonymes. *Hedera quinquefolia Linn. Spec.*, 292.; *Fitis quinquefolia Lam. Ill.*, No. 2815.; *Cissus hederacea Ph. Fl. Amer. Sept.*, 1. p. 170.; *Cissus quinquefolia Hort. Par.*; *Fitis hederacea Willd. Spec.*, p. 1182.; *Ampelopsis quinquefolia Hook. Fl. Bor. Amer.*, 1. 114.; *Vigne Vierge, Fr.*; *Jungfern Reben, or wilder, Wein, Ger.*

Engravings. Cornut. Canad., t. 100.; E. of Pl., 2868.; and our fig. 146.

Spec. Char. &c. Leaves digitate, of from 3 to 5 leaflets, that are stalked, oblong, toothed with mucronated teeth. Racemes dichotomously corymbose. (*Dec. Prod.*, i. p. 633.) A vigorous-growing climber, or trailer, rooting at the joints; a native of North America, from Pennsylvania to Carolina, in woods on the Alleghany Mountains. It was introduced into England in 1629; and, from its rapid growth, and the beauty of its foliage (especially in autumn, when it changes to a deep rich red), it soon became popular all over Europe. It grows freely in the smoke of cities; and in London, and more especially in Paris, it may be found reaching to the tops of houses from 50 ft. to 60 ft. in height. In fine seasons, it produces flowers, which are of a greenish purple colour, succeeded by corymbs of small black fruit. Plants, in the London nurseries, cost 1s. each, and seeds 1s. a packet; at Bollwyller, plants are 1 franc each; at New York, 15 cents.



‡ 3. *A. (H.) HIRSU'TA Donn (Hort. Cant.)*. The hairy-leaved Ampelopsis.

Identification. Donn Hort. Cant.; Dec. Prod., 1. p. 633.; Don's Mill., 1. p. 694.

Synonymes. *Cissus hederacea* var. *hirsuta Ph. Fl. Amer. Sept.*, 1. p. 170.

Spec. Char., &c. Leaves pubescent on both surfaces. A native of the Alleghany Mountains, introduced in 1806, and, in our opinion, likely to be only a variety of *A. hederacea*.

‡ 4. *A. BIPINNA'TA Michx.* The bipinnate-leaved Ampelopsis.

Identification. Michx. Fl. Bor. Amer., 1. p. 160.; Dec. Prod., 1. p. 633.; Don's Mill., 1. p. 694.

Synonymes. *Fitis arborea Willd. Spec.*, 1. p. 1183.; *Cissus stans Pers. Syn.* 1. p. 183., *Ph. Fl. Amer. Sept.*, 1. p. 170.

Engraving. Pluk. Mant., p. 412. fig. 2.

Spec. Char., &c. Leaves bipinnate, smooth; leaflets cut in a lobed manner. Racemes pedunculate, almost doubly bifid. Berries globose and cream-coloured. (*Dec. Prod.*, i. p. 633.) A deciduous shrub, with slender stems, but scarcely a climber; a native of Virginia and Carolina, in shady woods. It was introduced in 1700; and, being much admired for the beauty of its foliage, is not uncommon in collections. Plants, in the London nurseries, cost 1s. 6d. each; at Bollwyller, ?; at New York, 50 cents.

App. i. *Anticipated hardy Species of Ampelopsis.*

‡ *A. bôtrya* Dec. is a native of the eastern coast of Africa, with cordate leaves, reddish flowers, and berries black and eatable. It is described by Loureiro, but has not yet been introduced.

‡ *A. heterophylla* Blume, *Fitis javanica Spreng.*, a native of Java, has palmate leaves, and is considered as likely to endure our winters in the open air.

‡ *A. capreolata* G. Don, *Fitis capreolata D. Don*, is a native of Nepal, and resembles *A. hederacea* in every particular, except that it is one half smaller. Mr. Royle has given a figure of this plant (*Illus.*, i. 26.), and observes that he considers it the same as the *Fitis hederacea* of Dr. Wallich. The genera *Fitis*, *Ampelopsis*, and *Cissus* are so mixed together in the older botanical works, that there may probably be some of the above names that belong to *Fitis* or *Cissus*, and some names under *Fitis* and *Cissus* which belong to *Ampelopsis*.

GENUS III.

CISSUS L. THE CISSUS. *Lin. Syst. Tetrándria Monogýnia.*

Identification. *Lin. Gen.*, No. 147.; *Dec. Prod.*, 1. p. 627.; *Don's Mill.*, 1. p. 689.

Derivation. *Kissos* is the Greek name of the ivy, which these plants in some manner resemble.

Gen. Char., &c. *Calyx* almost entire. *Petals* 4, falling off separately. *Ovary* 4-celled. *Berry* 1—4-seeded. (*Dec. Prod.*, i. p. 630.) Climbing plants, chiefly ligneous, with simple, trifoliolate, or palmate leaves, and cymes or corymbs of small flowers, greenish, yellow, and sometimes purplish. Above 70 ligneous species are described in *Don's Miller*, a few of which are green-house plants, and already introduced into British gardens.

1. *C. orientális* Lam., figured in *Lam. Ill.*, t. 84. fig. 2., is a native of the Levant, and, according to *Sweet's Hortus Britannicus*, was introduced in 1818. It is a green-house climber; and, being considered tolerably hardy, it might be tried against a conservative wall.

2. *C. quinata* Ait., a native of the Cape of Good Hope, introduced in 1790, has palmate leaves, and is treated as a green-house plant.

3. *C. antarctica* Vent. Choix, t. 21., and our fig. 147, is a native of New Holland, whence it was introduced in 1790, and is commonly called the kangaroo vine. It has large, cordate, serrated, smoothish leaves, and but seldom, if ever, flowers in our green-houses. It is, probably, as hardy as other New Holland shrubs.


147



148



149



4. *C. capensis* Willd. is a native of the Cape of Good Hope, introduced in 1792.

C. vitigena, (fig. 148), *C. quinquefolia* (fig. 149), and, probably, other species now kept in our green-houses, and some even in our stoves, might, probably, prove half-hardy, if judiciously treated.

CHAP. XXVII.



OF THE HALF-HARDY LIGNEOUS PLANTS OF THE ORDER
GERANIA'CEÆ.

We introduce this order chiefly for the sake of recommending a trial of some of the hardier varieties of the common pelargonium; the roots of which, at least, will live through the winter at the bottom of a wall, if the soil be kept quite dry during that season, and covered with straw. The

150



151



152



following sorts may, perhaps, be chosen for a trial, in preference to some others: — *P. Barringtonii*, *cucullatum* (fig. 150.), *macranthon*, *megalanthon*, *calamistratum*, *quercifolium*, *peltatum*, *zonale* (fig. 151.), *Bentinckianum*, *inquinans* (fig. 152.)

CHAP. XXVIII.

OF THE HALF-HARDY LIGNEOUS PLANTS OF THE ORDER
ZYGOPHYLLACEÆ.

154 *Meliánthus major* L. (*Bot. Reg.*, t. 45, and our fig. 153.) is a well-known suffruticose green-house plant, admired for the beauty of its glaucous leaves, and the sharp and remarkable notching of its leaflets. It has stood out in many situations in the neighbourhood of London, at the base of a wall, protected only during the most severe weather. It has even flowered in the open air in some situations. In the Oxford Botanic Garden, this plant has stood out for many years at the base of a south wall, and also at the base of an east wall, and has flowered in both situations, the last time in 1834. In Devonshire, at Kingsbridge, in the garden of the Moults, it has stood out for many years as a bush; and, in November, 1855, was between 10 ft. and 12 ft. high.



153 *Zygophyllum sessilifolium* L. (figured in *Bot. Mag.*, t. 2184, and our fig. 154.), a native of the Cape of Good Hope, and *Z. Morgsana* L., also from the Cape, both green-house plants, which have been upwards of a century in this country, and which are interesting for their peculiar foliage and yellow flowers, may, with other ligneous Cape species, deserve a trial in favourable situations against a conservative wall.

CHAP. XXIX.

OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE
ORDER RUTAÆÆ.

IN this order there are two genera, *Ruta* and *Aplophýllum*, which contain some undershrubs, hardy or half-hardy. They are thus contradistinguished:—

RUTA L. Calyx 4-parted. Stamens 8. Styles 4, connected. Ovary almost stalked. Capsule 4-lobed, 4-celled. (*Don's Mill.*, i. p. 775.) Leaves compoundly divided.

APLOPHYLLUM Andr. *Juss.* Calyx 5-parted. Stamens 10. Styles 5, connected. Capsule 5-lobed, 5-celled. (*Don's Mill.*, i. p. 775.) Leaves undivided.

GENUS I.



RUTA L. THE RUE. *Lin. Syst.* Octándria Monogýnia.

Identification. Tourn. Inst., t. 133; *Lin. Gen.*, 523; *Dec. Prod.*, l. p. 769; *Don's Mill.*, l. p. 778.
Synonyms. Rue, Fr; Raute, Ger

Derivation. According to De Theis, incapable of explanation; but the same in all the most ancient languages; namely, *ruz* in Runic; *rude, rata, ruta, or rutu*, in Anglo-Saxon; *rutiza* in Slavonian; *ruta* in Italian and Latin; *ruda* in Spanish; *rutē* in Greek; said to be from *ruō*, to flow, in allusion to some expelling qualities of the plants.

■ 1. *R. GRAVE^OLENS L.* The heavy-scented, or common, Rue.

Identification. Lin. Spec., 548.; Dec. Prod., 1. p. 710.; Don's Mill., 1. p. 778.

Synonymes. *R. hortiensis* Mill. Dict., No. 1.; Rue, Fr.; Gartenraute, Ger.

Engravings. Du Ham. Arb., 2. t. 61.; Woodv. Med. Bot., t. 37.; Pluk. Icon., t. 332.; E. of Pl., 5886.; and our fig. 155.

Spec. Char., &c. Leaves supra-decompound; the lobes oblong, the terminal one obovate. Petals entire or toothed. (*Dec. Prod.*, i. p. 710.) A beautiful evergreen undershrub, native of the south of Europe, in sterile places, and cultivated in British gardens from time immemorial. It flowers from June to September, and ripens seeds.



Geography, History, &c. The rue may be found in a wild state in the south of France, in Spain, and in the north of Italy. We have gathered it, growing along with *Psoralea bituminosa*, on the rocks about Nice, and along the coast near Genoa. The rue was first recorded by Turner in 1562; but, from its reputed medicinal qualities, and its use in religious ceremonies, it was probably introduced into England by the monks, many centuries before.

Properties and Uses. "The rue and its allies," Professor Barnet observes, "are bitter stimulating plants, with a strong but rather unpleasant smell, and a hot bitter taste. *R. graveolens* is, indeed, so acrid, that the bruised leaves will excoriate the lips and nostrils, and inflame the skin, if applied as a cataplasm. Rue was much esteemed in ancient medicine: Hippocrates commends it: for many ages it was considered a preventive of contagion, and called the herb of grace; and, in later times, Boerhaave observes that the greatest commendations he can bestow upon it fall short of its merits. 'What medicine,' says he, 'can be more efficacious for promoting perspiration for the cure of hysteric passion, and of epilepsies, and for expelling poison?' But, notwithstanding all these praises, which are truly questionable, rue is now seldom employed, except in the form of tea, by village doctresses." (*Burnet's Outlines of Botany*, vol. 2. p. 882.) Notwithstanding these observations of Professor Burnet, the medicinal properties of the rue have been spoken of in terms of respect by Lewis, Woodville, Thomson, and other authors; and the following is a summary of their observations. The internal use of the rue is unsafe in large quantities; but eaten with bread and butter, as it very commonly is in France and Germany, a considerable dose may be taken without injury. In Britain, it is given to children as a vermifuge; and, mixed with butter, to poultry, as a remedy for the roup, &c. It is also given to dogs as a cure for the distemper, and is considered by the country people generally as an excellent "cleanser of the blood." By distillation with water, an essential oil is obtained from it; and by infusion in alcohol, a tincture which is warm, pungent, acrid, and penetrating. A conserve, made by beating the fresh leaves with fine sugar, is the most commodious form for using the herb in substance: the extract is given in doses of from 10 to 15 grains. To labouring men, if used with discretion, it might prove a condiment to their food, in the same way as garlic does in France, Spain, and Italy, and onions in most parts of Europe.

Poetical and legendary Allusions. Rue, as it is observed in Martyn's *Miller*, was anciently named herb grace, or the herb of grace; and it is to this day called ave grace in Sussex, in allusion, doubtless, to *Ave Maria, gratia plena*; and it is remarkable that Mary, in Hebrew, means bitter. Warburton says that rue had its name, "herb of grace," from its having been used in exorcisms. Ophelia, in Shakspeare's *Hamlet*, says to the Queen, "There's rue for you, and here's some for me; we may call it herb of grace o' Sundays." Herb of grace

was, indeed, the common name for rue in Shakspeare's time; and Greene, in his *Quip for an upstart Courtier*, has this passage:—"Some of them smiled, and said rue was called herb-grace, which, though they scorned in their youth, they might wear in their age, and that it was never too late to say *miserere*." The gardener in *Richard II.* says of the Queen,—

"Here did she drop a tear; here in this place,
I'll set a bank of rue, sour herb of grace:
Rue, even for ruth, here shortly shall be seen,
In the remembrance of a weeping queen."

Perdita, in *The Winter's Tale*, says,—

"Reverend sirs,
For you there 's rosemary and rue; these keep
Scenting and savour all the winter long;
Grace and remembrance be to you both."

They are both evergreens, retaining their appearance and taste during the whole year, and, therefore, are proper emblems of remembrance and grace. Rue seems to have been used formerly in nosegays; for the Clown, in *All's Well that Ends Well*, having said of the Countess, "She was the sweet-marjoram of the salad, or rather the herb of grace," Lafew replies, "They are not salad herbs, you knave, they are nose herbs;" upon which the Clown, in character, remarks, "I am no great Nebuchadnezzar, Sir, I have not much skill in grass;" thus punning upon the name of grace, as the gardener did upon the other name of rue. (*Don's Mill*, i. p. 779.) "Among the ancients, rue was used in several superstitious practices: 'You are not yet at the parsley, nor even at the rue,' was a common saying with the Greeks to those persons who, having projected an enterprise, had not begun to put it in execution. In ancient times, gardens were edged with borders of parsley and rue; and those persons who had not passed these borders were not accounted to have entered a garden: thence the proverb originated." (*Reid's Historical and Literary Botany*, p. 153.)

Physiological Phenomenon. "Linnæus having observed that the rue moved one of its stamens every day to the pistil, Sir James Smith examined the *Ruta angustifolia*, and found many of the stamens in the position which he describes, holding their anthers over the stigma; while those which had not come to the stigma were lying back upon the petals, as well as those which had already performed their office, and had returned to their original situation. Trying with a quill to stimulate the stamens, he found them all quite void of irritability: they are strong, stout, conical bodies, and cannot, without breaking, be forced out of the position in which they happen to be. The same phenomenon has been observed in several other flowers; but it is nowhere more striking, or more easily examined, than in the species of rue." (*Don's Mill*, i. p. 779.)

The Rue as a hardy Shrub. Though the rue is seldom seen in British gardens otherwise than as an herb of 1 ft. or 1½ ft. in height, yet when planted in dry, deep, calcareous soil, and suffered to grow without being cut over, it forms a singularly handsome evergreen shrub, attaining the height of 6 ft., or even 8 ft., in as many years. The manner in which the leaves are cut, their glaucous hue, the profusion of fine dark yellow flowers, which are produced for several months in succession, and often throughout the whole winter, justify us in strongly recommending the rue for cultivation as an ornamental plant. It will not succeed, however, if mixed with other trees and shrubs of rampant growth, nor attain a large size, unless in a sheltered situation, and in a soil that is deep, free, and calcareous. It forms beautiful evergreen separation hedges for cottage gardens; and some fine hedges of this sort, and also large single plants, may be seen in the bottoms of old chalk-pits on the south bank of the Thames, about Gravesend, in Kent. The plant is propagated in the easiest manner, by seeds or cuttings, and requires no other pruning during its whole existence than cutting off the withered flower-stalks. It appears to be a shrub of very great durability. In point of ultimate magnitude, rate of growth, soil, situation, and culture, the rosemary, the lavender, the sage, the hyssop, the thyme, and the more hardy *teucriums* may be considered as suitable associates for the rue.

App. i. *Half-hardy Species of Rûta.*

The following species of *Rûta* are generally kept in the frame or green-house; but there is little doubt that they would live in very dry soil or in lime rubbish, at the base of a wall, with some protection during severe weather. *R. pinnata* L. (*Bot. Reg.*, t. 307.), a native of the Canary Islands, where it grows to the height of 6 ft.; *R. bracteosa* Dec., a shrub 2 ft. high, a native of Sicily; *R. angustifolia* Pers. (fig. 156.), a native of the south of France, which was considered by Linnæus and others as a variety of the common rue; *R. macrophylla* Sol., from the north of Africa, where it grows 3 ft. high; *R. montana* Cus., from the south of Europe, also growing 3 ft. high; *R. divaricata* Tenore, from the south of Italy; *R. corsica* Dec., from Corsica; *R. albiflora* Hook., from Nepal, which was introduced in 1823, and which is found in the Himalayas, at elevations of from 500 ft. to 800 ft.; together with some other species from Nepal, from the south of Europe, and from the north of Africa; might all, we think, be tried at the base of a conservative wall, with every prospect of success. Perhaps half or more of the sorts above enumerated are only varieties of *Rûta graveolens*, but the shrub is so truly beautiful in the form and colour of its foliage, in its neat and compact shape, and its numerous flowers, that every variety is well worth cultivating.



156

GENUS II.



APLOPHY'LLUM Andr. Juss. THE APLOPHYLLUM, or SIMPLE-LEAVED RUE. Lin. Syst. Decândria Monogynia.

This genus, which forms a section of *Rûta* in De Candolle's *Prodromus*, was instituted by A. Jussieu in *Mém. Mus.*, 12. p. 464., and is adopted by G. Don. It contains two or three species of small under-shrubs, which are hardy, but which are more frequently treated as herbaceous than as ligneous plants.

- 1. *A. unifolium* G. Don., *Rûta unifolia* L., (*Bot. Rep.*, 565., and our fig. 157.) has entire oblong-lanceolate leaves, and yellow flowers in corymbs. It is a native of Spain, near Valencia, and also of Greece. It was introduced in 1752, grows to about 1 ft. in height, and flowers from July to September.
- 2. *A. suaveolens* G. Don., *Rûta suaveolens* Dec., has spatulately lanceolate glaucous leaves, and yellow flowers in corymbs, smelling like those of *Prîmula officinâlis*. It is a native of Tauria, where it forms a shrub about 2 ft. high; and was introduced in 1800. It flowers from June to September.
- 3. *A. fruticosum* G. Don, *Rûta fruticulosa* Lab., is a native of the country about Damascus. It grows about 1 ft. high, but has not yet been introduced into Britain.



157

CHAP. XXX.

OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE ORDER XANTHOXYLA'CEÆ.

THE genera belonging to this order which contain hardy species are three, *Xanthoxylum*, *Ptelea*, and *Ailantus*, which are thus distinguished in Don's *Mill.*, i. p. 777.

XANTHOXYLUM L., and *H. et Kth.* Flowers bisexual. Calyx 3—5-parted, with an equal number of petals and stamens. Carpels 1—5, 2-valved. Leaves simple, ternate, abruptly and impari-pinnate.

PTELEA L. Flowers bisexual. Calyx 4—5-parted. Petals 4—5. Stamens 4—5. Fruit compressed, 2—3-celled; cells 1-seeded, turgid in the centre, each cell extended into an orbicular reticulated wing. Leaves of 3 leaflets, rarely of 5 leaflets.

AILANTUS Desf. Flowers polygamous. Calyx 5-cleft. Petals 5. Stamens 10, unequal. Styles 3—5, arising from the notches of the ovaries. Carpels 3—5, membranous, 1-celled, 1-seeded. Leaves abruptly or impari-pinnate.

GENUS I.



XANTHOXYLUM L., and *H. B. et Kth.* THE XANTHOXYLUM, or
TOOTHACHE TREE. *Lin. Syst. Diœcia Tri-Pentândria.*

Identification. *Lin. Gen.*, No. 150. and 1109.; *Dec. Prod.*, 1. p. 725.; *Don's Mill.*, 1. p. 801.

Synonymes. *Zanthoxylum* (it is thus spelled in many botanical works); *Kampânna* *Ratin.*; *Clavaler*, *Fr.*; *Zahwehholz*, *Ger.*

Derivation. From *xanthos*, yellow, and *xylon*, wood; from the yellowness of the wood, more especially of the roots. The French name means club tree, and the German name, the toothache tree.

‡ 1. **X. FRAXINEUM** Willd. The Ash-leaved *Xanthoxylum*, or common *Toothache Tree*.

Identification. Willd. *Sp.*, 4 p. 757.; *Dec. Prod.*, 1. p. 726.; *Don's Mill.*, 1. p. 802.

Synonymes. *Zanthoxylum ramidorum* *Michx. Fl. B. A.*, 2. p. 25.; *Z. Clava Hérculis* var. *Lin. Sp.*, 1455., *Lam. Dict.*, 2. p. 38.; *Z. americanum* *Mill. Dict.*, No. 2.; *Z. caribæum* *Gart. Fruct.*, but not of *Lam.*; *Clavaler à Feuilles de Frêne*, *Fr.*; *Eichen-blättriges Zahwehholz*, *Ger.*

Engravings. *Du Ham. Arb.*, 1. t. 97.; *Catesb. Carol.*, 1. t. 26.; *E. of Pl.*, 1886.; our *fig. 158.*; and the plate of this species in Vol. II.

Spec. Char., &c. Leaves pinnate, of 4 to 5 pairs of leaflets, and an odd one; the leaflets ovate, obscurely sawed, equal at the base; the petiole round and devoid of prickles; prickles in the situation of stipules. Flowers in axillary umbels, without petals: the sexes diœcious. (*Dec. Prod.*, i. p. 726, 727.) A low deciduous tree, a native of North America, from Canada to Virginia and Kentucky, in woods near rivers; cultivated in England since 1740, and flowering in March and April. In its native country, this tree is seldom seen higher than from 12 ft. to 15 ft.; but its stem is decidedly that of a tree rising to the height of from 3 ft. or 4 ft. without side shoots, and then branching out, and forming a regular head. The flowers are yellowish, with red anthers. The bark and capsules are of a hot acrid taste, and are used for relieving the pains of the toothache; whence the popular name. A tincture of the bark is also used for curing rheumatism. This tree is common in British collections, but is never seen of any great size. There is one at Syon, about 13 ft. high; in the Cambridge Botanic Garden, one about 10 ft.; and some at White Knights, from 10 ft. to 13 ft. high. In the Edinburgh Botanic Garden, 10 years planted, it is 6 ft. high. It is generally propagated by seeds or by cuttings of the roots. Plants, in the London nurseries, are 1s. 6d. each; at Bollywiler, 1 franc and 20 cents; at New York, 25 cents, and seeds 1 dollar a quart.

Variety.

‡ *X. f. 2 virginicum*, the *X. virginicum* of *Lodd. Cat.*, of which there is a plant in the garden of the London Horticultural Society, and several in the arborctum of Messrs. Loddiges, appears to us only a variety of *X. fraxineum*.

‡ 2. **X. TRICARPUM** Michx. The three-fruited *Xanthoxylum*, or *Toothache Tree*.

Identification. *Michx. Fl. Bor. Amer.*, 2. p. 355.; *Dec. Prod.*, 1. p. 726.; *Don's Mill.*, 1. p. 803.

Synonymes. *Fagâra fraxinifolia* *Lam. Ill.*, 1. t. 354.

Engraving. *Lam. Ill.*, 1. t. 354.

Spec. Char., &c. Leaves pinnate; the leaflets 5 to 5 pairs, and an odd one, all on short stalks, oblong-oval, acuminate, finely sawed, oblique at the base. Petioles and branches prickly. Panicles terminal. Petals 5. (*Dec. Prod.*, i. p. 725.) A deciduous shrub, a native of Carolina and Florida, introduced in 1846, and flowering in July. Height 6 ft.

158



‡ 3. X. MI'LE Willd. The smooth, or thornless, Xanthoxylum, or Toothache Tree.

Identification. Willd. Enum., 1013.; Dec. Prod., 1. p. 727.; Don's Mill., 1. p. 802.

Spec. Char., &c. Thornless. Leaves impari-pinnate, downy beneath. Flowers axillary. (*Dec. Prod.*, i. p. 727.) Introduced from North America in 1812, and said to be a tree growing to the same height as *X. fraxineum*, with flowers of the same colour, and produced in the same months. It may, possibly, be only a variety of the *X. fraxineum*, as *Gleditschia inermis* is only a variety of *G. triacanthos*.

App. i. Half-hardy Species.

The species of this genus are not very ornamental, otherwise, there are some others, which are natives of China and Japan, which might be tried against a conservative wall: see the enumeration of all the species that have been introduced into Britain, in our *Hortus Britannicus*.

GENUS II.



PTE'LEA L. THE PTELEA, or SHRUBBY TREFOIL. *Lin. Syst.* Monœ'cia Tetra-Pentândria.

Identification. Lin. Gen., No. 152.; Dec. Prod., 2. p. 82.; Don's Mill., 1. p. 806.

Synonymes. *Bellücia* Adans.; Orme de Samarie, *Fr.*; Lederblume, *Ger.*

‡ 1. P. TRIFOLIATA L. The three-leafleted-leaved Ptelea, or Shrubby Trefoil.

Identification. Lin. Sp., 173.; Willd. Sp. Pl., 1. 670.; Dec. Prod., 2. p. 82.; Don's Mill., 1. p. 806.

Synonymes. Orme de Samarie à trois Feuilles, *Fr.*; dreyblättrige Lederblume, *Ger.*

Engravings. Dill. Elth., t. 182.; Mill. Ic., t. 211.; N. Du Ham., t. 57.; Hayne Abbild., t. 74.; Schmidt Arb., 2. t. 76.; and the plate in our Second Volume.

Spec. Char., &c. Leaf of three leaflets that are ovate acute, the middle one much tapered towards its base. Flowers in corymbs, usually tetrandrous. (*Dec. Prod.*, ii. p. 82.) A shrub or low tree from North America, where it is found from New York to Carolina in shady moist hedges, and on the edges of woods among rocks. It was introduced in 1704, and produces its small greenish white flowers in corymbose clusters in June and July. These are succeeded by flattened winged capsules, somewhat resembling those of the elm; whence the French name of *orme*.

Varieties.

P. t. 2 *pentaphylla* Munchh. Hans., 3. p. 342., has 5 leaflets; but we have not seen a plant.

P. t. 3 *pubescens* Pursh has pubescent leaves, and is a native of Pennsylvania; but we are not aware of its being in cultivation in Britain.

Description, &c. When this plant is pruned up with a single stem, it forms a handsome low tree with a hemispherical head; but in British gardens it is more frequently found as a large shrub, with numerous stems proceeding from the same basal point. The species was originally sent to England by Banister, and plants of it were raised by Bishop Compton at Fulham; but they were lost, and the plant was reintroduced from Carolina by Catesby in 1724. Being hardy, and of easy culture in any common soil, the tree is not uncommon in collections; and it well deserves a place there, both on account of the beauty of the leaves, and of the fruits, and the handsome general form of the tree. It is easily propagated by cuttings (put in in the autumn, and covered with a hand-glass), or by seeds.

Statistics. At Purser's Cross, there is a tree 25 ft. high, with a trunk 14 in. in diameter; at Gloucester Lodge, Brompton, is a somewhat deformed specimen, 16 ft. high, with a trunk 8 in. in diameter at the ground; in Sloane Square is one upwards of 12 ft. high; in the Fulham Nursery, one 12 years planted, and 14 ft. high; in the Hammersmith Nursery, one 5 years planted, 10 ft. high; in Middlesex, near Shepperton, by the road side, a very handsome tree, about 25 ft. high, with the head 30 ft. in diameter; in Surrey, at Claremont, 15 ft. high; in Wiltshire, at Longleat, one 50 years planted, and 18 ft. high, the diameter of the trunk 14 in., and of the head 24 ft.; in Worcestershire, at Croome, 10 years planted, and 15 ft. high. In Scotland, in the Edinburgh Botanic Garden, 15 years planted, and 12 ft. high; in Banffshire, at Gordon Castle, 45 ft. high, the diameter of the trunk 15 in., and of the head 27 ft., in a loamy soil, and a sheltered situation. In Ireland, at Terenure, near Dublin, 15 years planted, and 8 ft. high. In France, at Paris, in the Jardin des Plantes, 60 years planted, and 57 ft. high, the diameter of the head 40 ft.; at Scéaux, 10 years planted, and 20 ft. high; in the Botanic Garden at Toulon, 10 years planted, and 10 ft. high; at Nerrières, near Nantes, 15 ft.

high. In Saxony, at Würthz, 45 years planted, and 25 ft. high; and the variety *P. t. pentaphylla*, 34 years planted, and 15 ft. high. In Austria, at Vienna, in the University Botanic Garden, 15 ft. high. In Bavaria, at Munich, in the Botanic Garden, 24 years planted, and 18 ft. high.

Commercial Statistics. Plants, in London, cost 1s. 6d. each, and seeds 1s. a packet; at Bollwyller, plants 15 francs a hundred; at New York, 25 cents each, and seeds 1 dollar a quart.

App. i. Other ligneous Species of *Plèca*, hardy and half-hardy.

♁ *P. monophylla* Lam. has simple ovate lanceolate leaves, and grows to the height of 4 ft. in Carolina; but, though hardy, it has not yet been introduced.

♁ *P. podocarpa* Moc. is a native of Mexico, where it forms a shrub from 6 ft. to 10 ft. in height; *P. podocarpa* Dec. is a Mexican shrub of the same size; and *P. ovata* Lour. is a simple-leaved species, a native of Cochin-China. These, if introduced, would probably prove hardy or half-hardy.

GENUS III.



AILANTUS Desf. THE AILANTO. *Lin. Syst. Polygàmia Monœ'cia.*

Identification. Desf. Act. Acad. Par., 1786, p. 263; Dec. Prod., 2 p. 88; Don's Mill., 1. p. 807.

Synonymes. *Rhûs Ehrh., Ellis, and Mench;* Verne du Japon, *Fr.*; Götterbaum, *Ger.*

Derivation. Ailanto is the name of *Ailantus glandulosa* Desf. in the Moluccas. It was long considered as a species of *Rhûs*, whence the French name; and the meaning of the aboriginal word being, it is said, Tree of Heaven, hence the German name, Götterbaum, Tree of the Gods.

♁ 1. *A. GLANDULO'SA* Desf. The glandulous-leaved Ailanto.

Identification. Desf. Act. Acad. Par., 1786, p. 263; Dec. Prod., 1. p. 89; Don's Mill., 1. p. 807.

Synonymes. *A. procera* Sal. Prod., p. 271; *Rhûs hypselodéndron Mench;* *R. caecodéndron Ehrh.;*

R. sinense Ellis; *Aylanthé glanduleux, Fr.;* drüsiger Götterbaum, *Ger.*

Engravings. L'Hérit. Stirp., t. 84; Wats. Dend. Brit., t. 108; N. Du Ham., 1. t. 35; our fig. 159; and the plate of the tree in our Second Volume.

Spec. Char., &c. Leaves impari-pinnate; the leaflets coarsely toothed at the base; the teeth glandulous on the under side. (*Dec. Prod., i. p. 89.*) A deciduous tree of the first rank, introduced from China in 1751, and growing to the height of 60 ft. or upwards. In some years, the tree is said to bear only male flowers; and L'Héritier states that only twice in 10 years it bore both male and female flowers at the same time in France. In his time, it had produced fruit in the Jardin des Plantes at Paris, and in the Botanic Garden at Leyden; but in both cases it was immature. It has since, however, produced perfect fruit, from which plants have been raised.

It has also ripened fruit at White Knights in England. The flowers, which appear in August, are in large, upright, rather compact panicles, of a whitish green colour, and exhale a disagreeable odour. The fruit resembles the keys of the ash, but is smaller. The leaves are from 3 ft. to 6 ft. in length; those produced by vigorous suckers, in favourable situations, attaining the latter dimensions. The tree grows with great rapidity for the first 10 or 12 years, producing shoots from 3 ft. to 6 ft. in length at first, and attaining the height of 15 ft. or 20 ft. in 5 or 6 years in favourable situations. Afterwards its growth is much slower. The wood is of a fine grain; it has a satiny aspect, and is hard, and well fitted for the purposes of cabinet-making. The tree has a noble appearance when clothed with leaves; and its gigantic boughs and shoots, and its straight, erect, thick

159



trunk, seem to justify its original appellation of tree of heaven. On the first approach of frost, the leaflets begin to drop, without having previously shown any great change of colour, displaying in this respect a striking difference from the leaves of most species of *Rhûs*, to which those of this tree bear a general resemblance.

Geography, History, &c. This species of *Allantus* is a native of the northern provinces of China, more particularly in the neighbourhood of Pekin. Seeds were sent to England, to the Royal Society of London, by the Jesuit missionary D'Incarville, in 1751; and they were sown by Miller in the Chelsea Botanic Garden, and by Philip Carteret Webb, at Busbridge, in Surrey, in the same year. As the tree produced suckers freely, it was soon generally propagated; and there are many fine specimens of it in different parts of the country. The original tree planted by Mr. Webb was cut down some years ago; but several others, which have sprung up from the roots left in the soil, were in existence about the same spot when we visited it in 1834. (See *Gard. Mag.*, vol. ix. p. 481.) The tree was introduced into France, in 1780, by Mr. Blaikie, and the oldest specimens are at St. Leu, and in the Jardin des Plantes. We have not heard of the timber having been applied to any useful purpose in Europe, because trees of a large size are not yet sufficiently numerous to admit of their being cut down for profitable application. In France and Italy, it is much valued as a tree for shading public walks, and is planted for that purpose along with the tulip tree, the horsechestnut, the platanus, and other large-leaved exotic trees. Its leaves are not liable to be attacked by insects, which is a very great recommendation, and they continue on the tree, and retain their green colour, till the first frosts in November; when the leaflets drop suddenly off, the petioles remaining on often a week or two longer. The tree grows in any soil, though one that is light and somewhat humid, and a sheltered situation, suit it best. In France, it is said to thrive on chalky soils, and attain a large size, where scarcely any other tree will grow. It is readily propagated by cuttings of the roots. It might probably be found a valuable tree to be treated as coppice, and cut down every third or fourth year for fuel.

Statistics. A. glandulosa in the Environs of London. The largest tree is at Syon; it is 70 ft. high, the diameter of the trunk 3 ft. 10 in., and of the head 40 ft.; the trunk forms an erect column of about 30 ft. before it branches, and the head is hemispherical. This tree flowers and fruits occasionally. At Kew there is a tree 60 ft. high; in the Fulham Nursery, one 50 ft. high; in the Mile End Nursery, one 36 ft. high, with a trunk 2 ft. in diameter; the leaves, even in the tree of that age and height, measuring 4 ft. 6 in. in length. At Fulham Palace there is a tree, 20 years planted, which is 25 ft. high. In the London nurseries, plants are frequently to be met with, of two or three years' growth, 12 ft. high.

A. glandulosa South of London. In Kent, at Cobham Hall, 20 years planted, and 36 ft. high, the diameter of the trunk 1 ft., and of the head 15 ft. In the Isle of Jersey, in Saunders's Nursery, 10 years planted, and 16 ft. high. In Sussex, at Langham Park, 9 years planted, and 12 ft. high; at Kidbrooke, 30 years planted, and 30 ft. high.

A. glandulosa North of London. In Bedfordshire, at Amptill Park, 3 years planted, and 12 ft. high. In Berkshire, at White Knights, there are several trees, 19 years planted, and from 27 ft. to 30 ft. high, the diameter of the trunks about 9 in., and of the heads about 30 ft.: these trees produce flowers every year, and fruit occasionally. In Buckinghamshire, at Temple House, 3 years planted, and 7 ft. high. In Cambridge, in the grounds of St. John's College, there are two trees, both near the river Cam, one of which is 40 ft. high, with a trunk 2 ft. 7 in. in diameter. In Warwickshire, at Combe Abbey, 10 years planted, and 12 ft. high; in the Handsworth Nursery, near Birmingham, 12 years planted, and 16 ft. high. In Worcestershire, at Croome, 45 years planted, and 60 ft. high, the diameter of the head 90 ft.

A. glandulosa in Scotland. In Berwickshire, at the Hirscl, 3 years planted, and 6 ft. high. In Perthshire, at Kinfauns Castle, 8 years planted, and 16 ft. high. In Stirlingshire, at Airthrie Castle, 10 years planted, and 25 ft. high. In Sutherlandshire, at Dunrobin Castle, 43 ft. high, the diameter of the trunk 1 ft. 6 in., and of the head 35 ft.

A. glandulosa in Ireland. At Dublin, in the Glasnevin Botanic Garden, 20 years planted, and 18 ft. high; at Terenure, 20 years planted, and 14 ft. high. In the Clonmel Nursery, 15 years planted, and 14 ft. high. In the Kilkenny Nursery, 35 years planted, and 21 ft. high.

A. glandulosa in Foreign Countries. In France, at Paris, in the Jardin des Plantes, 68 ft. high, with the head 44 ft. in diameter, flowering most years, and ripening seeds occasionally; at St. Leu, where it was planted on a large scale by Mr. Blaikie in 1791, it is 80 ft. high, with a trunk from 3 ft. to 3½ ft. in diameter; in the Botanic Garden at Toalon, 50 years planted, and 60 ft. high; at Nerrères, near Nantes, 40 years planted, it is 50 ft. high. At Geneva, at the entrance to the Botanic Garden, there is a tree, from 45 ft. to 50 ft. high, the trunk of which, in 1833, measured 7 ft. 3 in. in circumference at the surface of the ground; when in flower, the disagreeable odour which proceeds from it is felt at a distance of nearly a quarter of a mile (*cinq minutes de distance*); and its suckers occupy the ground for 40 ft. or 50 ft. around it in every direction. In Saxony, at Wörlitz, a tree, 20 years planted, is 25 ft. high. In Austria, at Vienna, in the University Botanic Garden,

40 years planted, it is 35 ft. high; in Rosenthal's Nursery, 20 years planted, and 30 ft. high; at Brück on the Leytha, 40 years planted, and 42 ft. high. In Prussia, at Sans Souci, 30 years planted, and 20 ft. high; at the Pfauen Insel, 8 years planted, and 22 ft. high. In Bavaria, at Munich, in the Botanic Garden, 20 years planted, and 20 ft. high. In Hanover, at Göttingen, in the University Botanic Garden, 10 years planted, and 30 ft. high. In Cassel, at Wilhelmshöhe, 60 ft. high. In Sweden, at Lund, in the Botanic Garden, 2 years planted, and 4 ft. high. In Italy, at Monza, 29 years planted, and 60 ft. high.

Commercial Statistics. Plants, in the London nurseries, are 1s. 6d. each; at Bollwyller, from 1 franc to 1 franc 50 cents; at New York, ?.

CHAP. XXXI.

OF THE HARDY AND HALF-HARDY LIGNEOUS PLANTS OF THE ORDER CORIACEÆ.

This order consists of only one genus, of which there is one species quite hardy, and one or two others, natives of New Zealand and Nepal, which are probably half-hardy.

GENUS I.



CORIARIA Niss. THE CORIARIA. *Lin. Syst. Diœcia Decândria.*

Identification. Niss. in Act. Par., 1711. t. 12.; Dec. Prod., 1. p. 739.; Don's Mill., 1. p. 818.

Synonymes. Redoul, Fr.; Gerberstrauch, Ger.

Derivation. From *corium*, a hide; *C. myrtifolia* being used both in tanning leather and in dyeing it black.

Gen. Char., &c. Flowers either hermaphrodite, monœcious, or diœcious.

Calyx 5-parted. *Petals* 5, sepaloid, smaller than the lobes of the calyx.

Stamens 10, hypogynous, 5 between the lobes of the calyx and the angles of the ovarium, 5 between the petals and the furrows of the ovarium. *Anthers* bursting by longitudinal slits. *Style* none. *Stigmas* 5, long, awl-shaped.

Carpels 5, surrounding a fleshy axis; when ripe, close together, but separate, not opening, 1-seeded, surrounded with glandular lobes. *Ovule* and seed pendulous. *Albumen* none. *Embryo* straight. — Branches square, opposite. Leaves opposite, simple, 3-ribbed. (*Lindley, Introd. to N. S., and Key.*)

♁ 1. *C. MYRTIFOLIA* L. THE Myrtle-leaved Coriaria.

Identification. Lin. Sp., 1467.; Dec. Prod., 1. p. 739.; Don's Mill., 1. p. 818.

Synonymes. Fustel des Corroyeurs, or Redoul à Feuilles de Myrte, Fr.; Myrtenblättriger Gerberstrauch, Ger.

Engravings. Lam. Ill., t. 822.; Du Ham., 1. t. 73.; Wats. Dend. Brit., t. 103.; and our fig. 160.

Spec. Char., &c. Leaves ovate-lanceolate, acute, three-nerved, on short foot-stalks, glabrous.

160

Flowers in rather upright racemes. (*Dec. Prod.,* 1. p. 739.)

A deciduous shrub, growing to the height of from 4 ft. to 6 ft., in the south of Europe and north of Africa, in hedges and waste places. It was introduced into England in 1629, and has since been frequent in collections, flowering from May to August. In its native country, it is said to be used for tanning, and for dyeing black; but whether it is cultivated for this purpose, or merely gathered where found wild, we have not been able to ascertain. In Britain, it is cultivated as an ornamental undershrub, chiefly remarkable for its myrtle-like leaves, and the handsome frond-like form of its branches.



According to Dumont, the leaves, and more especially the berries, are a deadly poison, both to man and animals. The leaves have been employed in France to adulterate senna leaves, and have produced fatal consequences. It is stated by Féc, that several soldiers of the French army in Catalonia became stupefied by eating the berries, and three of them died in consequence. The shoots of this plant very frequently die down to the ground; so that it is never to be seen, in Britain at least, with shoots of above 3 or 4 years' growth; but it sends up shoots from its roots freely every year; and these shoots are sometimes 3 ft. or 4 ft. in length.

☞ 2. *C. NEPALE'NSIS* Wall. *Pl. As. Rar.*, t. 289., The Nepal Coriaria, grows in Nepal at heights of from 5000 ft. to 7000 ft., and is applied to the same purposes as *C. myrtifolia*; but what is remarkable is, the berries are eaten by the inhabitants.

☞ 3. *C. MICROPHY'LLA* Poir., The small-leaved Coriaria, from Peru; *synon. C. sarmentosa* Forst., from New Zealand, introduced in 1823; and some other Mexican and Peruvian species not yet in the country, may, probably, be found half-hardy; because, as the great body of the plant is under ground, it may be protected by leaves or litter during winter; even if the top should die down every year, like that of a herbaceous plant, shoots may spring up again from the root every spring.

CHAP. XXXII.

OF THE HARDY LIGNEOUS PLANTS OF THE ORDER STAPHYLEA'CEÆ.

GENUS I.



STAPHYLE'NA L. THE STAPHYLEA, or BLADDER-NUT TREE. *Lin. Syst.* Pentándria Di-Trigýnia.

Identification. *Lin. Gen.* No. 374.; *Dec. Prod.*, 2. p. 2.; Don's Mill, 2. p. 2.

Synonymes. *Staphylodéndron* Tourn.; *Staphilier*, faux Pistachier, *Fr.*; *Pimpernuss*, *Ger.*

Derivation. Abridged from *Staphylodéndron*, its name before the days of Linnaeus, derived from *staphulé*, a bunch or cluster, and *déndron*, a tree; the flowers and fruits being disposed in clusters, and the plant being ligneous.

Gen. Char., &c. *Calyx* of 5 coloured sepals, connected at the base, in æstivation imbricate. *Petals* 5, in æstivation imbricate. *Stamens* 5, perigynous, alternate with the petals, and opposite the sepals. A large urceolate disk, or nectary, within the corolla. *Ovarium* 2- or 3-celled, superior. *Fruit* membranous. *Seeds* with a bony testa, and a large truncate hilum. *Leaves* opposite, pinnate, with both common and partial stipules. *Flowers* in terminal stalked racemes. (*Lindley, Introd. to N. S.*; from the character of the order.)

☞ 1. *S. TRIFO'LIA* L. The three-leafletted-leaved Staphylea, or Bladder-nut Tree.

Identification. *Lin. Sp.*, 386.; *Dec. Prod.*, 2. p. 2.; Don's Mill, 2. p. 2.

Synonymes. *Staphilier à Feuilles ternées*, *Fr.*; *Virginische Pimpernuss*, *Ger.*

Engravings. *Schmidt Baum.*, t. 81.; *N. Du Ham.*, vi. t. 12.; *Hayne Abbild.*, t. 36.; *Krauss*, t. 109.; *E. of Pl.*, No. 3823.; and our *fig.* 161. in flower, and *fig.* 162. in fruit.

Spec. Char., &c. The leaf of 3 leaflets, which are ovate, acuminate, regularly sawed, and, when young, pubescent; the style smooth; the capsule bladdery. (*Dec. Prod.*, ii. p. 2.) A deciduous shrub, a native of North America, and found from New York to Carolina, on rocks. It was introduced in 1640, and produces its whitish flowers in May and June.

161



It grows to the height of 6 ft. or 8 ft. Though this species was cultivated by the elder Tradescant, it has never become very common in British gardens. It is propagated either by seeds or cuttings. Plants, in London, cost 1s. 6d. each; at Bollwyler, 1 franc; and at New York, 25 cents.

162



■ 2. *S. PINNATA* L. The pinnated-leaved *Staphylea*, or *Bladder-nut Tree*.

Identification. Lin. Sp., 386.; Dec. Prod., 2. p. 3.; Don's Mill., 2. p. 3.

Synonymes. *Staphylodendron pinnatum* Ray; *Staphilier à Feuilles ailées*, Fr.; *gemeine Pimper-nuss*, Ger.

Engravings. Eng. Bot., t. 1560.; Hayne Abbild., t. 36.; E. of Pl., 5822.; and our fig. 163.

Spec. Char., &c. Leaves pinnate, of 5—7

oblong, perfectly glabrous, serrated leaflets; the flowers in racemes; the capsules membranaceous and bladderly. (*Dec. Prod.*, ii. p. 3.)

A deciduous shrub, with leaves somewhat like those of the ash or the elder; a native of Europe, in hedges and thickets; and generally considered indigenous to England; though, according to Ray, it was scarcely found in sufficient plenty to be deemed certainly wild. Smith describes it (*Eng. Flor.*, ii. p. 111.) as a smooth branching shrub, throwing up many side suckers. In gardens, it is to be found from 6 ft. to 12 ft. high, and exhibiting a much more luxuriant growth than the preceding species;

and forming a singular object, when in fruit, from its large bladderly capsules. Each of these capsules contains a hard smooth nut, which, in some parts of Europe, is strung as a bead by the Roman Catholics. Haller says that the kernels taste like those of the pistacia, and are eaten in Germany by children; and this appears to have been formerly the case in England; for Gerard says the kernels, though sweet at first, are succeeded by a nauseous taste, and, finally, they act as an emetic. The wood is hard, of a yellowish white, and close grained; but it is seldom found of a sufficient size to be applied to any useful purpose. The flowers contain a great deal of honey, and are very attractive to bees. In the London nurseries, the plant is generally cultivated by side suckers, by cuttings put in during the month of September, or by seeds, which are ripened in abundance. The seeds ought to be sown as soon as they are ripe; because, as they contain an oil, they very soon become rancid. They should not be covered with more than half an inch of soil. They will come up the following June, with two large, lance-shaped, seminal leaves; though sometimes they do not come up for two years. Price as in the preceding.





App. i. *Half-hardy Species of Staphyléa.*

■ *S. Bumalda* Dec. has leaves like *S. trifolia*, and white nodding flowers. It is a native of the mountains of Japan, and was introduced in 1812.

■ *S. heterophylla* Ruiz et Pav. has leaves like *S. pinnata*. It is a native of Peru, where it grows to the height of 12 ft.

END OF THE FIRST VOLUME.





UNIVERSITY OF ILLINOIS-URBANA



3 0112 051870233